Arc-disjoint in- and out-branchings rooted at the same vertex in compositions of digraphs

Gregory Gutin¹ and Yuefang Sun^{2*} ¹ Department of Computer Science Royal Holloway, University of London Egham, Surrey, TW20 0EX, UK g.gutin@rhul.ac.uk ² Department of Mathematics, Shaoxing University Zhejiang 312000, P. R. China, yuefangsun2013@163.com

Abstract

A digraph D = (V, A) has a good pair at a vertex r if D has a pair of arc-disjoint in- and out-branchings rooted at r. Let T be a digraph with t vertices u_1, \ldots, u_t and let H_1, \ldots, H_t be digraphs such that H_i has vertices u_{i,j_i} , $1 \le j_i \le n_i$. Then the composition $Q = T[H_1, \ldots, H_t]$ is a digraph with vertex set $\{u_{i,j_i} \mid 1 \le i \le t, 1 \le j_i \le n_i\}$ and arc set

 $A(Q) = \cup_{i=1}^{t} A(H_i) \cup \{ u_{ij_i} u_{pq_p} \mid u_i u_p \in A(T), 1 \le j_i \le n_i, 1 \le q_p \le n_p \}.$

When T is arbitrary, we obtain the following result: every strong digraph composition Q in which $n_i \ge 2$ for every $1 \le i \le t$, has a good pair at every vertex of Q. The condition of $n_i \ge 2$ in this result cannot be relaxed. When T is semicomplete, we characterize semicomplete compositions with a good pair, which generalizes the corresponding characterization by Bang-Jensen and Huang (J. Graph Theory, 1995) for quasi-transitive digraphs. As a result, we can decide in polynomial time whether a given semicomplete composition has a good pair rooted at a given vertex.

Keywords: branching; semicomplete digraph; digraph composition.

AMS subject classification (2010): 05C20, 05C70, 05C76, 05C85.

1 Introduction

We use a standard digraph terminology and notation as in [4, 5]. A digraph D = (V, A) is strongly connected (or strong) if there exists a path from x to y and a path from y to x in D for every pair of distinct vertices x, y of D. An out-tree (in-tree, respectively) rooted at a vertex r is an orientation of a tree such that the in-degree (out-degree, respectively) of every vertex but r equals one. An out-branching (in-branching, respectively) in a digraph

^{*}Corresponding author. This author was supported by NSFC No. 11401389.

D is a spanning subgraph of D which is out-tree (in-tree, respectively). It is well-known and easy to show [4,5] that a digraph has an out-branching (in-branching, respectively) rooted at r if and only if D has a unique initial strong connectivity component (terminal strong connectivity component, respectively) and r belongs to this component. Out-branchings and inbranchings when they exist can be found in linear-time using, say, depth-first search from the root.

Edmonds [11] characterized digraphs with k arc-disjoint out-branchings rooted at a specified vertex r. Furthermore, there exists a polynomial algorithm for finding k arc-disjoint out-branchings with a given root r if they exist [4]). However, it is NP-complete to decide whether a digraph D has a pair of arc-disjoint out-branching and in-branching rooted at r, which was proved by Thomassen, see [1]. Following [9] we will call such a pair a good pair rooted at r. Note that a good pair forms a strong spanning subgraph of D and thus if D has a good pair, then D is strong. The problem of the existence of a good pair was studied for tournaments and their generalizations, and characterizations (with proofs implying polynomial-time algorithms for finding such a pair) were obtained in [1] for tournaments, [7] for quasi-transitive digraphs and [9] for locally semicomplete digraphs. Also, Bang-Jensen and Huang [7] showed that if r is adjacent to every vertex of D (apart from itself) then D has a good pair rooted at r.

In this paper, we study the existence of good pairs for digraph compositions. Let T be a digraph with t vertices u_1, \ldots, u_t and let H_1, \ldots, H_t be digraphs such that H_i has vertices u_{i,j_i} , $1 \le j_i \le n_i$. Then the *composition* $Q = T[H_1, \ldots, H_t]$ is a digraph with vertex set $\{u_{i,j_i} \mid 1 \le i \le t, 1 \le j_i \le n_i\}$ and arc set

$$A(Q) = \bigcup_{i=1}^{t} A(H_i) \cup \{ u_{ij_i} u_{pq_p} \mid u_i u_p \in A(T), 1 \le j_i \le n_i, 1 \le q_p \le n_p \}.$$

Digraph compositions generalize some families of digraphs. In particular, semicomplete compositions, which are compositions where T is a semicomplete digraph, generalize strong quasi-transitive digraphs as every strong quasi-transitive digraph is a strong semicomplete composition in which H_i is either a one-vertex digraph or a non-strong quasi-transitive digraph. To see that strong compositions form a significant generalization of strong quasi-transitive digraphs, observe that the Hamiltonian cycle problem is polynomial-time solvable for quasi-transitive digraphs [13], but NP-complete for strong compositions (see, e.g., [6]). When H_i is the same digraph H for every $i \in [t], Q$ is the lexicographic product of T and H, see, e.g., [15]. While digraph compositions has been used since 1990s to study quasi-transitive digraphs and their generalizations, see, e.g., [3, 4, 12], the study of digraph decompositions in their own right was initiated only recently in [16].

In the next section, we obtain the following somewhat surprising result: every strong digraph composition Q in which $n_i \ge 2$ for every $i \in [t]$, has a good pair rooted at every vertex of Q. The condition of $n_i \ge 2$ in this result cannot be relaxed. Indeed, the characterization of quasi-transitive digraphs with a good pair [7] provides an infinite family of strong quasi-transitive digraphs which have no good pair rooted at some vertices. In Section 3, we characterize semicomplete compositions with a good pair generalizing the corresponding result in [7]. This allows us to decide in polynomial time whether a given semicomplete composition has a good pair rooted at a given vertex. In Section 4, we discuss some open problems and a recent related result.

Let p and q be integers. Then $[p..q] := \{p, p+1, \ldots, q\}$ if $p \leq q$ and \emptyset , otherwise. In particular, if p > 0, [p] will be a shorthand for [1..p].

2 Compositions of digraphs: T arbitrary

A digraph D = (V, A) has a strong arc decomposition if A has two disjoint sets A_1 and A_2 such that both (V, A_1) and (V, A_2) are strong. Sun et al. [16] obtained sufficient conditions for a digraph composition to have a strong arc decomposition. In particular, they proved the following:

Theorem 2.1 Let T be a digraph with t vertices $(t \ge 2)$ and let H_1, \ldots, H_t be digraphs. Then $Q = T[H_1, \ldots, H_t]$ has a strong arc decomposition if T has a Hamiltonian cycle and one of the following conditions holds:

- t is even and $n_i \ge 2$ for every $i = 1, \ldots, t$;
- t is odd, n_i ≥ 2 for every i = 1,...,t and at least two distinct subgraphs H_i have arcs;
- t is odd and $n_i \ge 3$ for every i = 1, ..., t apart from one i for which $n_i \ge 2$.

Lemma 2.2 Let $Q = T[H_1, \ldots, H_t]$, where $t \ge 2$. If T has a Hamiltonian cycle and H_1, \ldots, H_t are arbitrary digraphs, each with at least two vertices, then Q has a good pair at any root r.

Proof: For the case that t is even, by Theorem 2.1, Q has has a pair of arc-disjoint strong spanning subgraphs Q_1 and Q_2 . Observe that in Q_1 (Q_2 , respectively), we can find an out-branching (in-branching, respectively) at r (in polynomial time), as desired.

Now we assume that t is odd. Without loss of generality, let $u_{1,1}$ be the root. Let T'_1 be the path $u_{1,1}u_{2,1}\ldots u_{t,1}u_{1,2}u_{2,2}\ldots u_{t,2}$, and let T'_2 be the in-tree rooted at $u_{1,1}$ with arc set $\{u_{i,2}u_{i+1,1} \mid 1 \leq i \leq t-1\} \cup \{u_{i,1}u_{i+1,2} \mid 2 \leq i \leq t-1\} \cup \{u_{t,1}u_{1,1}, u_{t,2}u_{1,1}\}$. By definition, $V(T'_1) = V(T'_2) = \{u_{i,j} \mid 1 \leq i \leq t, 1 \leq j \leq 2\}$. For any vertex $u_{i,j}$ with $1 \leq i \leq t$ and $j \geq 3$, we add the arcs $u_{i-1,1}u_{i,j}$ and $u_{i,j}u_{i+1,1}$ to T'_1 and T'_2 , respectively. Note that here $u_{0,1} = u_{t,1}$ and $u_{t+1,1} = u_{1,1}$. Observe that the resulting two subgraphs form a pair of out-branching and in-branching rooted at $u_{1,1}$, which are arc-disjoint.

We will use the following decomposition of strong digraphs. An *ear* decomposition of a digraph D is a sequence $\mathcal{P} = (P_0, P_1, P_2, \dots, P_t)$, where P_0 is a cycle or a vertex and each P_i is a path, or a cycle with the following properties:

(a) P_i and P_j are arc-disjoint when $i \neq j$.

(b) For each $i \in [0..t]$, let D_i denote the digraph with vertices $\bigcup_{j=0}^{i} V(P_j)$ and arcs $\bigcup_{j=0}^{i} A(P_j)$. If P_i is a cycle, then it has precisely one vertex in common with $V(D_{i-1})$. Otherwise the end vertices of P_i are distinct vertices of $V(D_{i-1})$ and no other vertex of P_i belongs to $V(D_{i-1})$. (c) $\bigcup_{j=0}^{t} A(P_j) = A(D)$.

The following result is well-known, see, e.g., [4].

Theorem 2.3 Let D be a digraph with at least two vertices. Then D is strong if and only if it has an ear decomposition. Furthermore, if D is strong, every cycle can be used as a starting cycle P_0 for an ear decomposition of D, and there is a linear-time algorithm to find such an ear decomposition.

Lemma 2.4 Let $Q = T[\overline{K_2}, \ldots, \overline{K_2}]$, where $|V(T)| = t \ge 2$ and $\overline{K_2}$ is the digraph with two vertices and no arcs. If T is strong, then Q has a good pair at any root r.

Proof: Without loss of generality, let $r = u_{1,1}$. Since T is strong, u_1 belongs to some cycle C in T. By Theorem 2.3, T has an ear decomposition $\mathcal{P} = (P_0, P_1, P_2, \cdots, P_p)$, such that $P_0 = C$ is the starting cycle. Let G_i denote the subgraph of G with vertices $\bigcup_{i=0}^{i} V(P_i)$ and $\operatorname{arcs} \bigcup_{i=0}^{i} A(P_i)$.

We will prove the lemma by induction on $i \in \{0, 1, \ldots, p\}$. For the base step, by Lemma 2.2, the subgraph $P_0[\overline{K_2}, \ldots, \overline{K_2}]$ has a good pair rooted at u_1 . For the inductive step, assume that $G_i[\overline{K_2}, \ldots, \overline{K_2}]$ has a pair of arcdisjoint out-branching T'_1 and in-branching T'_2 rooted at r. Without loss of generality, let $P_{i+1} = u_s u_{s+1} \ldots u_\ell$. The following argument will be divided into two cases according to whether P_{i+1} is a cycle.

Case 1: P_{i+1} is a cycle. In this case $u_s = u_\ell \in V(G_i)$. By Lemma 2.2, in the subgraph $P_{i+1}[\overline{K_2}, \ldots, \overline{K_2}]$, there is a pair of arc-disjoint out-branching T_1'' and in-branching T_2'' rooted at $u_{s,1}$. Let $T_1 = T_1' \cup T_1''$ and $T_2 = T_2' \cup T_2''$. Observe that T_1 is an out-branching and T_2 is an in-branching rooted at r in $G_{i+1}[\overline{K_2}, \ldots, \overline{K_2}]$. Since $P_{i+1}[\overline{K_2}, \ldots, \overline{K_2}]$ and $G_i[\overline{K_2}, \ldots, \overline{K_2}]$ are arc-disjoint, T_1 and T_2 are arc-disjoint.

Case 2: P_{i+1} is a path. In this case, $u_s, u_\ell \in V(G_i)$ and $s \neq \ell$. We just consider the case that $\ell - s \geq 2$ since the remaining case is trivial (no need to change the current pair of out- and in-branchings). Let T_1 be the union of T'_1 and the two paths $u_{s,i}u_{s+1,i} \ldots u_{\ell-1,i}$ where $1 \leq i \leq 2$. Let T_2 be the union of T'_2 and the two paths $u_{s,i}u_{s+1,i} \ldots u_{\ell-1,i}$ where $1 \leq i \leq 2$. Let T_2 be the union of T'_2 and the two paths $u_{s,1}u_{s+1,2}u_{s+2,1}u_{s+3,2} \ldots u_{\ell,i}$ and $u_{s,2}u_{s+1,1}u_{s+2,2}u_{s+3,1} \ldots u_{\ell,j}$, where $\{i, j\} = \{1, 2\}$. Observe that T_1 is an out-branching and T_2 is an in-branching rooted at r in $G_{i+1}[\overline{K_2}, \ldots, \overline{K_2}]$, moreover, they are arc-disjoint.

Thus, by induction, we are done.

Theorem 2.5 Let $Q = T[H_1, \ldots, H_t]$, where $t \ge 2$. If T is strong and H_1, \ldots, H_t are arbitrary digraphs, each with at least two vertices, then Q has a good pair at any root r. Furthermore, this pair can be found in polynomial time.

Proof: Without loss of generality, let $r = u_{1,1}$. Let Q' be the subgraph of Q induced by the vertex set $\{u_{i,j} \mid 1 \leq i \leq t, 1 \leq j \leq 2\}$. In Q' delete arcs between vertices $u_{i,1}$ and $u_{i,2}$ for every $i \in [t]$. By Lemma 2.4, Q' contains a pair of arc-disjoint out-tree T'_1 and in-tree T'_2 rooted at r. By definition of out-tree, there is an arc $u_{p_i,q_i}u_{i,2}$ in T'_1 for every $i \in [t]$. For every $i \in [t]$ and $j \in [3..n_i]$, add $u_{p_i,q_i}u_{i,j}$ to T'_1 . This results in an out-branching T_1 . By definition of in-tree, there is an arc $u_{i,2}u_{a_i,b_i}$ in T'_2 for every $i \in [t]$. For every $i \in [t]$ and $j \in [3..n_i]$, add $u_{i,j}u_{a_i,b_i}$ to T'_2 . This results in an in-branching T_2 . Observe that T_1 and T_2 are arc-disjoint since T'_1 and T'_2 are arc-disjoint and the added arcs have heads and tails from $\{u_{i,j} \mid 1 \leq i \leq t, 3 \leq j \leq n_i\}$, respectively, in the arcs added to T'_1 and T'_2 , respectively. Note that the proofs of Theorem 2.1, Lemmas 2.2 and 2.4, and this theorem are constructive and can be converted into polynomial-time algorithms. This fact and the polynomial-time algorithm of Theorem 2.3 imply that T_1 and T_2 can be constructed in polynomial time.

3 Compositions of digraphs: *T* semicomplete

We use $N^{-}(v)$ ($N^{+}(v)$, respectively) to denote the set of all in-neighbours (out-neighbours, respectively) of a vertex v in a digraph D.

The next result was obtained by Bang-Jensen and Huang [7].

Theorem 3.1 Let D be a strong digraph and r a vertex of D such that $V(D) = \{r\} \cup N^{-}(r) \cup N^{+}(r)$. There is a polynomial-time algorithm to decide whether D has a good pair at r.

For a path $P = x_1 x_2 \dots x_p$ and $1 \leq i \leq j \leq p$, let $P[x_i, x_j] := x_i x_{i+1} \dots x_j$. We now prove the following result on semicomplete compositions which generalizes a similar result for quasi-transitive digraphs by Bang-Jensen and Huang [7].

Theorem 3.2 A strong semicomplete composition Q has a good pair rooted at r if and only if $Q' = Q[\{r\} \cup N^-(r) \cup N^+(r)]$ has a good pair rooted at r.

Proof: Let $Q = T[H_1, \ldots, H_t]$ and $A = V(Q) \setminus V(Q')$. Without loss of generality, assume that $r \in V(H_1)$. By definitions of a semicomplete composition and Q', we have $A = V(H_1) \setminus \{r\}$.

Assume that Q' has a good pair rooted at r, an out-branching B'_r^+ and an in-branching B'_r^- . Starting with B'_r^+ , we can construct an out-branching B'_r in Q as follows. Let v be a vertex such that $vr \in B'_r^-$. Then add the arc vu to B'_r^+ for each $u \in A$. Similarly, starting with B'_r^- , we could construct an in-branching B_r^- in Q as follows: for each $u \in A$, add the arc uv' to B'_r^- , where $rv' \in B'_r^+$. Observe that B'_r^+ and B'_r^- are arc-disjoint, as desired.

Now we prove the other direction. Assume that Q has a good pair, an out-branching B_r^+ and an in-branching B_r^- , rooted at r. If $B_r^+[V(Q')]$ and $B_r^-[V(Q')]$ are branchings, then we are done. Otherwise, we will obtain an in-branching (out-branching, respectively) from B_r^- (B_r^+ , respectively) using the following procedure.

Choose a maximal path P of B_r^- to r, which contains a vertex $w \in A$, and assume that w is furthest from r among vertices in $A \cap V(P)$. If w is the first vertex of P, then delete it. Otherwise, the previous vertex u on Phas an arc ur to r (the arc ur exists since $A \subseteq V(H_1)$), and we replace P in B_r^- by two paths: one is P[p, u]r, where p is the first vertex of P, and the other is P[w, r].

Note that the in-degree $d^-(w)$ of w has decreased by one. Thus, after $d^-(w)$ such replacements the in-degree of w becomes equal to zero, i.e., w is the first vertex on its maximal path Q to r and therefore w will be deleted when we consider Q. This means that after a finite number of replacements, we will delete all vertices of A in B_r^- and obtain an in-branching B'_r^- of Q' rooted at r. Similarly, we can construct an out-branching B'_r^+ of Q'. Note that to build B'_r^- we add only arcs to r and to build B'_r^+ we add only arcs from r. This fact and the fact that B_r^- and B_r^+ are arc-disjoint, imply that B'_r^- and B'_r^+ are arc-disjoint, too.

By Theorems 3.1 and 3.2, we immediately have the following:

Theorem 3.3 Given a semicomplete composition and a vertex r, we can decide in polynomial time whether D has a good pair rooted at r.

4 Open Problems and Related Results

Theorem 3.2 provides a characterization for the following problem for semicomplete compositions: given a digraph D and a vertex $r \in V(D)$ decide whether D has a good pair rooted at r. The theorem generalizes a similar characterization by Bang-Jensen and Huang [7] for quasi-transitive digraphs. Strong semicomplete compositions is not the only class of digraphs generalizing strong quasi-transitive digraphs. Other such classes have been studied such as k-quasi-transitive digraphs [12] and it would be interesting to see whether a characterization for the problem (or, at least non-trivial sufficient conditions) on k-quasi-transitive digraphs can be obtained. As we mentioned above, Bang-Jensen and Huang [9] obtained a characterization for the problem on locally semicomplete digraphs. It would be interesting to see whether a characterization for the problem on in-locally semicomplete digraphs [2, 4] can be obtained.

An out-branching and in-branching B_r^+ and B_r^- are called *k*-distinct if B_r^+ has at least k arcs not present in B_r^- . The problem of deciding whether a digraph D has a k-distinct pair of out- and in-branchings is NP-complete since it generalizes the good pair problem (k = |V(D)| - 1). Bang-Jensen and Yeo [10] asked whether the k-distinct problem is fixed-parameter tractable when parameterized by k, i.e., whether there is an $O(f(k)|V(D)|^{O(1)})$ -time algorithm for solving the problem, where f(k) is an arbitrary computable function in k only. Gutin, Reidl and Wahlström [14] answered this open question in affirmative by designing an $O(2^{O(k \log^2 k)}|V(D)|^{O(1)})$ -time algorithm for solving the problem.

References

- J. Bang-Jensen, Edge-disjoint in- and out-branchings in tournaments and related path problems, J. Combin. Theory Ser. B, 51(1), 1991, 1–23.
- [2] J. Bang-Jensen, Locally semicomplete digraphs and generalizations, in *Classes of Directed Graphs* (J. Bang-Jensen and G. Gutin, eds.), Springer, 2018.
- [3] J. Bang-Jensen and G. Gutin, Paths and cycles in extended and decomposable digraphs. Discrete Math. 164 (1997) 5–19.
- [4] J. Bang-Jensen and G. Gutin, Digraphs: Theory, Algorithms and Applications, 2nd Edition, Springer, London, 2009.
- [5] J. Bang-Jensen and G. Gutin, Basic terminology, notation and results, in *Classes of Directed Graphs* (J. Bang-Jensen and G. Gutin, eds.), Springer, 2018.
- [6] J. Bang-Jensen, G. Gutin and A. Yeo, Arc-disjoint strong spanning subgraphs of semicomplete compositions, arXiv:1903.12225v1 [cs.DM].
- [7] J. Bang-Jensen and J. Huang, Quasi-transitive digraphs, J. Graph Theory, 20(2), 1995, 141–161.
- [8] J. Bang-Jensen and J. Huang, Decomposing locally semicomplete digraphs into strong spanning subgraphs, J. Combin. Theory Ser. B, 102, 2012, 701–714.
- [9] J. Bang-Jensen and J. Huang, Arc-disjoint in- and out-branchings rooted at the same vertex in locally semicomplete digraphs, J. Graph Theory, 77, 2014, 278–298.
- [10] J. Bang-Jensen and A. Yeo, The minimum spanning strong subdigraph problem is fixed parameter tractable. Discrete Appl. Math.,156:2924– 2929, 2008.
- [11] J. Edmonds, Edge-disjoint branchings, in *Combinatorial Algorithms* (B. Rustin ed.), Academic Press, 1973, 91–96.
- [12] H. Galeana-Sánchez and C. Hernández-Cruz, Quasi-transitive digraphs and their extensions, in *Classes of Directed Graphs* (J. Bang-Jensen and G. Gutin, eds.), Springer, 2018.
- [13] G. Gutin, Polynomial algorithms for finding Hamiltonian paths and cycles in quasi-transitive digraphs. Australasian J. Combin. 10, 1994, 231–236.
- [14] G. Gutin, F. Reidl and M. Wahlström, k-Distinct In- and Out-Branchings in Digraphs. J. Comput. Syst. Sci. 95 (2018) 86–97.
- [15] R.H. Hammack, Digraphs Products, in Classes of Directed Graphs (J. Bang-Jensen and G. Gutin, eds.), Springer, 2018.

[16] Y. Sun, G. Gutin, J. Ai, Arc-disjoint strong spanning subgraphs in compositions and products of digraphs, Discrete Math. 342(8), 2019, 2297–2305.