FINITE DISTRIBUTIVE NEARLATTICES

LUCIANO J. GONZÁLEZ AND ISMAEL CALOMINO

ABSTRACT. Our main goal is to develop a representation for finite distributive nearlattices through certain ordered structures. This representation generalizes the well-known representation given by Birkhoff for finite distributive lattices through finite posets. We also study finite distributive nearlattices through the concepts of dual atoms, boolean elements, complemented elements and irreducible elements. We prove that the sets of boolean elements and complemented elements form semi-boolean algebras. We show that the set of boolean elements of a finite distributive lattice is a boolean lattice.

1. Introduction and preliminaries

It is well known that the class of distributive lattices have many and important applications, especially in logic and computer science. Thus, it is also important to study some natural generalizations of distributive lattices which may result interesting. An important class of distributive lattices for computer science is the class of finite distributive lattices. This paper deals with the concept of *finite distributive nearlattice*, which is a natural and nice generalization of finite distributive lattice.

Distributive nearlattices were study from different points of view: algebraic, topological and logical [15, 21, 14, 2, 7, 11, 19, 8, 12, 6, 9, 10, 17, 18].

In Section 2 we consider the notions of dual atom and boolean element. We prove that the boolean elements form a semi-boolean algebra ([1]). In Section 3 we show that the semi-boolean algebra of boolean elements of a finite distributive nearlattice **A** is a homomorphic image of **A**. We generalize the concept of complement elements from distributive lattice to distributive nearlattice and, we show that the set of complement elements of a distributive nearlattice is a semi-boolean algebra. Section 4 is concerned with the free distributive lattice extension of a distributive nearlattice ([11]). Finally, in Section 5 we develop a discrete representation for the class of finite distributive nearlattice, which is a generalization of the well-known representation for finite distributive lattice given by Birkhoff [4].

We assume that the reader is familiar with the theory of ordered sets and lattices. Our main references for Order and Lattice theory are [16, 20, 3]. Now we introduce the notational conventions that we use throughout the paper.

Let $P = \langle P, \leq \rangle$ be a poset. A subset $U \subseteq P$ is said to be an *upset* when for all $a, b \in P$, if $a \leq b$ and $a \in U$, then $b \in U$. Dually we have the notion of *downset*. Given an element $a \in P$, the *principal upset* generated by a is $[a) := \{x \in P : a \leq x\}$. Dually we have the *principal downset* (a]. Given a subset Q of P and $a \in P$, $[a)_Q := \{x \in Q : a \leq x\}$.

²⁰¹⁰ Mathematics Subject Classification. Primary 06D75; Secondary 06A12; 06D05. Key words and phrases. Nearlattice; Lattice; Representation theorem; Irreducible elements.

Definition 1.1 ([15]). A distributive nearlattice is a join-semilattice $\mathbf{A} = \langle A, \vee, 1 \rangle$ with top element 1 such that for all $a \in A$, the principal upset [a] is a bounded distributive lattice (with respect to the induced order).

Let $\mathbf{A} = \langle A, \vee, 1 \rangle$ be a distributive nearlattice. For each $a \in A$, we denote the meet in [a) by \wedge_a . It should be noted that for all $x,y \in A$, the meet $x \wedge y$ exists in A if and only if x,y have a common lower bound in A. Thus, for all $x,y \in [a)$, the meet of x,y in [a) coincides with their meet in A, that is, $x \wedge_a y = x \wedge y$. This should be kept in mind since we will use it without mention. A subset $F \subseteq A$ is called *filter* if (i) $1 \in F$; (ii) F is an upset; and (iii) if $a,b \in F$ and $a \wedge b$ exists in A, then $a \wedge b \in F$. Let us denote by $\mathrm{Fi}(\mathbf{A})$ the collection of all filters of \mathbf{A} . It is straightforward that $\mathrm{Fi}(\mathbf{A})$ is a closure system. For $X \subseteq A$, we denote by $\mathrm{Fig}_A(X)$ the filter generated by X. Thus, we have that $\langle \mathrm{Fi}(\mathbf{A}), \cap, \vee, \{1\}, A \rangle$ is a distributive lattice where $F \vee G = \mathrm{Fig}_A(F \cup G)$, for all $F, G \in \mathrm{Fi}(\mathbf{A})$.

A non-empty subset $I \subseteq A$ is said to be an *ideal* when (i) I is a downset; (ii) if $a, b \in I$, then $a \lor b \in I$. A proper ideal I of A is called *prime* when for all $a, b \in A$, if $a \land b$ exists and $a \land b \in I$, then $a \in I$ or $b \in I$.

Theorem 1.2 ([21]). Let **A** be a distributive nearlattice. If I is an ideal and F is a filter of **A** such that $I \cap F = \emptyset$, then there exists a prime filter P of A such that $I \subseteq P$ and $P \cap F = \emptyset$.

Given two distributive nearlattices \mathbf{A}_1 and \mathbf{A}_2 , a map $f: A_1 \to A_2$ is said to be an *N-homomorphism* if f(1) = 1; for all $a, b \in A_1$, $f(a \lor b) = f(a) \lor f(b)$, and $f(a \land b) = f(a) \land f(b)$, whenever $a \land b$ exists in A_1 .

The reader is referred to [13, 15, 5] for further information about distributive nearlattices.

2. Dual atoms and boolean elements

Definition 2.1. Let **A** be a distributive nearlattice and $a \in A$. We will say that a is a dual atom if $a \neq 1$ and for each $x \in A$, $a \leq x \leq 1$ implies x = a or x = 1. We say that a is a boolean element if the principal filter [a] is a boolean lattice.

Let us denote by $\operatorname{At}_{d}(\mathbf{A})$ and $\operatorname{Bo}(\mathbf{A})$ the collections of all dual atoms of \mathbf{A} and of all boolean elements of \mathbf{A} , respectively. It follows that $\operatorname{At}_{d}(\mathbf{A}) \subseteq \operatorname{Bo}(\mathbf{A})$. If $a \in \operatorname{Bo}(\mathbf{A})$ and $x \in [a)$, we shall denote the complement of x in [a) by $\neg_a x$. The next properties will be used throughout the paper without mention.

Remark 2.2. Let $a \in At_d(\mathbf{A})$. Then, for all $x \in A$, we have that $x \le a$ or $x \lor a = 1$. Moreover, if \mathbf{A} is a finite distributive nearlattice, then for every x < 1, there exists $a \in At_d(\mathbf{A})$ such that $x \le a$.

Proposition 2.3. Let **A** be a distributive nearlattice. Let $a_1, \ldots, a_n \in \operatorname{At_d}(\mathbf{A})$ be such that $a_1 \wedge \cdots \wedge a_n$ exists in A. Then $a_1 \wedge \cdots \wedge a_n \in \operatorname{Bo}(\mathbf{A})$.

Proof. Let $a^* = a_1 \wedge \cdots \wedge a_n$. Since $\langle [a^*), \wedge_{a^*}, \vee, a^*, 1 \rangle$ is a bounded distributive lattice, it only remains to verify that each element of $[a^*)$ has a complement. For each a_i , let us denote the complement of an element $x \in [a_i)$ by $\neg_i x$. That is, for every $x \in [a_i)$, $\neg_i x \in [a_i)$ such that $x \wedge_{a_i} \neg_i x = a_i$ and $x \vee \neg_i x = 1$.

Let $x \in [a^*)$. Thus $x = (a_1 \vee x) \wedge \cdots \wedge (a_n \vee x)$. We define the element $\neg x := \neg_1(a_1 \vee x) \wedge \cdots \wedge \neg_n(a_n \vee x)$ (notice that the last meet exists because

 $a^* \le a_i \le \neg_i(a_i \lor x)$, for all i). It is clear that $\neg x \in [a^*)$. Now we show that $\neg x$ is the complement of x in $[a^*)$. On the one hand, we have

$$\neg x \land x = \bigwedge_{1 \le i \le n} \neg_i(a_i \lor x) \land \bigwedge_{1 \le i \le n} (a_i \lor x) = \bigwedge_{1 \le i \le n} [\neg_i(a_i \lor x) \land (a_i \lor x)]$$
$$= \bigwedge_{1 \le i \le n} a_i = a^*.$$

On the other hand, we have

$$\neg x \lor x = \left(\bigwedge_{1 \le i \le n} \neg_i(a_i \lor x) \right) \lor x = \bigwedge_{1 \le i \le n} \left(\neg_i(a_i \lor x) \lor x \right).$$

Since every a_i is a dual atom, it follows that

$$\neg_i(a_i \lor x) = \begin{cases} 1 & \text{if } x \le a_i \\ a_i & \text{if } x \nleq a_i \end{cases}$$

Hence, in any case, we obtain that $\neg_i(a_i \lor x) \lor x = 1$. Then $\neg x \lor x = 1$. We have proved that $\neg x$ is the complement of x in $[a^*)$. Therefore, $\langle [a^*), \wedge_{a^*}, \vee, a^*, 1 \rangle$ is a boolean lattice, and thus $a^* = a_1 \land \cdots \land a_n \in \operatorname{Bo}(\mathbf{A})$.

Let **A** be a distributive nearlattice and $a \in A$. We define the following set:

$$X_a := \{ x \in At_d(\mathbf{A}) : a \le x \}.$$

Thus, $X_a = \operatorname{At_d}(\mathbf{A}) \cap [a) = \operatorname{At_d}([a))$. If **A** is finite and $a \in \operatorname{Bo}(\mathbf{A})$, then since [a) is a finite boolean lattice it follows that

$$a = \bigwedge X_a$$
.

Lemma 2.4. Let **A** be a distributive nearlattice and $a, b \in A$. Then $X_{a \vee b} = X_a \cap X_b$. If $a \wedge b$ exists, then $X_{a \wedge b} = X_a \cup X_b$.

Proof. It is straightforward to show directly that $X_{a \vee b} = X_a \cap X_b$.

It is clear that $X_a \cup X_b \subseteq X_{a \wedge b}$. Let now $x \in X_{a \wedge b}$. Since $a \wedge b \leq x$, we have that $x = (a \vee x) \wedge (b \vee x)$. Given that $x \in \operatorname{At}_{\mathbf{d}}(\mathbf{A})$, it follows that $(a \leq x \text{ or } a \vee x = 1)$ and $(b \leq x \text{ or } b \vee x = 1)$. If $a \vee x = 1$ and $b \vee x = 1$, then x = 1, which is a contradiction. Hence, $a \leq x$ or $b \leq x$. That is, $x \in X_a \cup X_b$. Therefore $X_{a \wedge b} \subseteq X_a \cup X_b$.

Proposition 2.5. Let $\mathbf{A} = \langle A, \vee, 1 \rangle$ be a finite distributive nearlattice. Then $\langle \operatorname{Bo}(\mathbf{A}), \vee, 1 \rangle$ is a nearlattice subalgebra of \mathbf{A} .

Proof. We need to prove that Bo(**A**) is closed under \vee , and if $a, b \in \text{Bo}(\mathbf{A})$ are such that $a \wedge b$ exists in A, then $a \wedge b \in \text{Bo}(\mathbf{A})$.

Let $a, b \in \operatorname{Bo}(\mathbf{A})$. Notice that $a \vee b \leq \bigwedge X_{a \vee b}$, and since $a, b \in \operatorname{Bo}(\mathbf{A})$, it follows that $a = \bigwedge X_a$ and $b = \bigwedge X_b$. By Proposition 2.3, we have $\bigwedge X_{a \vee b} \in \operatorname{Bo}(\mathbf{A})$. Let us show that $a \vee b = \bigwedge X_{a \vee b}$. Suppose, towards a contradiction, that $a \vee b \neq \bigwedge X_{a \vee b}$. So $\bigwedge X_{a \vee b} \nleq a \vee b$. Then, there is a prime ideal P such that $a \vee b \in P$ and $\bigwedge X_{a \vee b} \notin P$. Thus, $a, b \in P$ and $P \cap X_{a \vee b} = \emptyset$. Since $\bigwedge X_a \in P$, $\bigwedge X_b \in P$ and P is prime, it follows that there is $x_a \in X_a$ such that $x_a \in P$ and there is $x_b \in X_b$ such that $x_b \in P$. Since x_a and x_b are dual atoms, we have

$$x_a \lor x_b = \begin{cases} 1 & \text{if } x_a \neq x_b \\ x_a = x_b & \text{if } x_a = x_b \end{cases}$$

Since P is proper and $x_a \vee x_b \in P$, we have that $x_a = x_b$. Let $x := x_a = x_b$. Thus $a \vee b \leq x$. Then $x \in X_{a \vee b} \cap P$, which is a contradiction. Hence $a \vee b = \bigwedge X_{a \vee b} \in Bo(\mathbf{A})$. Let now $a, b \in Bo(\mathbf{A})$ be such that $a \wedge b$ exists in A. By Lemma 2.4, we have

$$\bigwedge X_{a \wedge b} = \bigwedge (X_a \cup X_b) = \bigwedge X_a \wedge \bigwedge X_b = a \wedge b.$$

Then, it follows by Proposition 2.3 that $a \wedge b = \bigwedge X_{a \wedge b} \in Bo(\mathbf{A})$. Therefore, $Bo(\mathbf{A})$ is a nearlattice subalgebra of \mathbf{A} .

Proposition 2.6. Let **A** be a finite distributive nearlattice. Let $a \in Bo(\mathbf{A})$ and $b \in A$. If $a \leq b$, then $b \in Bo(\mathbf{A})$. That is, $[a]_{Bo(\mathbf{A})} = [a]_A$.

Proof. Let $a \in Bo(\mathbf{A})$ and $b \in A$ be such that $a \leq b$. Since a is a boolean element, we have $a = \bigwedge X_a$. We also have that $b \leq \bigwedge X_b$. Suppose that $\bigwedge X_b \nleq b$. Then, there is a prime ideal P such that $b \in P$ and $\bigwedge X_b \notin P$. Thus, $a \in P$ and $X_b \cap P = \emptyset$. Since $\bigwedge X_a \in P$ and P is prime, we obtain that there is $x \in X_a \cap P$. Then, $x \notin X_b$. Since $x \in At_d(\mathbf{A})$ and $b \nleq x$, it follows that $b \vee x = 1$. This is a contradiction because $b, x \in P$ and P is a proper ideal. Hence $b = \bigwedge X_b \in Bo(\mathbf{A})$.

A semi-boolean algebra ([1]) is a join-semilattice with a top element such that every principal upset is a Boolean lattice. In [1] Abbot show that the semi-boolean algebras are in a one-to-one correspondence with the implication algebras (also known as Tarski algebras).

Theorem 2.7. Let $\mathbf{A} = \langle A, \vee, 1 \rangle$ be a finite distributive nearlattice. Then, the nearlattice subalgebra $\langle \operatorname{Bo}(\mathbf{A}), \vee, 1 \rangle$ is a semi-boolean algebra.

Proof. Let $a \in Bo(\mathbf{A})$. By Proposition 2.6, we have $[a]_{Bo(\mathbf{A})} = [a]_A$, and since $Bo(\mathbf{A})$ is a nearlattice subalgebra of \mathbf{A} , it follows that $[a]_{Bo(\mathbf{A})}$ is a boolean lattice. Hence $Bo(\mathbf{A})$ is a semi-boolean algebra.

Proposition 2.8. Let A be a finite distributive nearlattice. The following are equivalent:

- (1) $\bigwedge \operatorname{At}_{d}(\mathbf{A})$ exists in A;
- (2) There is $a \in A$ such that $Bo(\mathbf{A}) = [a]$;
- (3) $Bo(\mathbf{A})$ is a boolean lattice.

Proof. (1) \Rightarrow (2) Let $a := \bigwedge \operatorname{At_d}(\mathbf{A})$. It follows by Proposition 2.3 that $a \in \operatorname{Bo}(\mathbf{A})$. Let $b \in \operatorname{Bo}(\mathbf{A})$. Then, $a = \bigwedge \operatorname{At_d}(\mathbf{A}) \le \bigwedge X_b = b$. Thus $b \in [a)$. Hence $\operatorname{Bo}(\mathbf{A}) \subseteq [a)$. Now, by Proposition 2.6, we have $[a) = [a]_{\operatorname{Bo}(\mathbf{A})} \subseteq \operatorname{Bo}(\mathbf{A})$. Therefore, $\operatorname{Bo}(\mathbf{A}) = [a)$.

- $(2) \Rightarrow (3)$ It follows straightforward by Theorem 2.7.
- $(3) \Rightarrow (1)$ Let $a \in A$ be the least element of Bo(**A**). Thus $[a) = [a]_{Bo(\mathbf{A})} = Bo(\mathbf{A})$. Since $At_d(\mathbf{A}) \subseteq Bo(\mathbf{A}) = [a)$, we obtain that $\bigwedge At_d(\mathbf{A})$ exists in A.

Corollary 2.9. If **L** is a finite distributive lattice, then $Bo(\mathbf{L})$ is a boolean lattice. Moreover, for each $a \in Bo(\mathbf{L})$ and $x \in [a)$, we have $\neg_a x = \neg x \lor a$, where $\neg x$ denotes the complement of x in $Bo(\mathbf{L})$.

Proof. From the previous proposition, it is clear that Bo(**L**) is a boolean lattice. Let $a \in \text{Bo}(\mathbf{L})$ and $x \in [a)$. Since [a) is a boolean lattice, there is unique $\neg_a x \in [a)$ such that $x \vee \neg_a x = 1$ and $x \wedge \neg_a x = a$. As $a \leq x$, by Proposition 2.6 we have $x \in \text{Bo}(\mathbf{L})$ and, there is $\neg x \in \text{Bo}(\mathbf{L})$ such that $x \vee \neg x = 1$ and $x \wedge \neg x = c$, where $c \in A$ is the least element of Bo(**L**). Consider the element $\neg x \vee a \in [a)$. So,

 $(\neg x \lor a) \lor x = (\neg x \lor x) \lor a = 1 \lor a = 1 \text{ and } (\neg x \lor a) \land x = (\neg x \land x) \lor (a \land x) = c \lor a = a.$ Hence, we have $\neg_a x = \neg x \lor a$.

3. Annihilators

Let $\mathbf{A} = \langle A, \vee, 1 \rangle$ be a distributive nearlattice. For every $a \in A$, let $a^{\top} = \{x \in A : a \vee x = 1\}$ (see [7, 14]). It follows that $a^{\top} \in \mathrm{Fi}(\mathbf{A})$, for all $a \in A$. We will say that an element $a \in A$ is dense if $a^{\top} = \{1\}$. Denote by $\mathrm{D}(A)$ the set of all dense elements of A.

Proposition 3.1. Let **A** be a finite distributive nearlattice. The following are equivalent:

- (1) $\bigwedge \operatorname{At}_{d}(\mathbf{A})$ exists in A;
- (2) There is $a \in A$ such that D(A) = (a];
- (3) D(A) is not empty.

Proof. (1) \Rightarrow (2) Let $a := \bigwedge \operatorname{At}_{\operatorname{d}}(\mathbf{A})$. Let $x \in a^{\top}$. So $x \vee a = 1$. Suppose that x < 1. Then, there is $b \in \operatorname{At}_{\operatorname{d}}(\mathbf{A})$ such that $x \le b$. Thus $b = b \vee a = 1$, which is a contradiction. Hence x = 1. It follows that $a^{\top} = \{1\}$, and thus $a \in \operatorname{D}(A)$. We obtain that $(a] \subseteq \operatorname{D}(A)$. Let now $b \in \operatorname{D}(A)$. So $b^{\top} = \{1\}$. Let $x \in \operatorname{At}_{\operatorname{d}}(\mathbf{A})$. Since $x \notin b^{\top}$, we have $b \vee x \ne 1$. Thus $b \le x$. Then $b \le \bigwedge \operatorname{At}_{\operatorname{d}}(\mathbf{A}) = a$. Hence $b \in (a]$. Therefore, $\operatorname{D}(A) \subseteq (a]$.

- $(2) \Rightarrow (3)$ It is immediate.
- $(3) \Rightarrow (1)$ Let us show that $\operatorname{At}_{d}(\mathbf{A})$ has a lower bound. As $\operatorname{D}(A) \neq \emptyset$, there is $a \in A$ such that $a^{\top} = \{1\}$. Let $x \in \operatorname{At}_{d}(\mathbf{A})$ and suppose that $a \nleq x$. Thus, $x \vee a = 1$. Then $x \in a^{\top} = \{1\}$, which is a contradiction because $x \in \operatorname{At}_{d}(\mathbf{A})$. Hence $a \leq x$, for all $x \in \operatorname{At}_{d}(\mathbf{A})$. Therefore, $\bigwedge \operatorname{At}_{d}(\mathbf{A})$ exists in A.

Proposition 3.2. Let **A** be a finite distributive nearlattice. The following properties are satisfied:

- (1) Let $a, b \in Bo(\mathbf{A})$. Then, $a \leq b$ if and only if $a^{\top} \subseteq b^{\top}$.
- (2) For each $a \in A$, we have $a^{\top} = (\bigwedge X_a)^{\top}$.

Proof. (1) It is easy to show that $a \leq b$ implies $a^{\top} \subseteq b^{\top}$. Conversely, assume that $a^{\top} \subseteq b^{\top}$. Let $x \in X_b$. Then $x \in \operatorname{At}_{d}(\mathbf{A})$ and $b \vee x = x \neq 1$, i.e., $x \notin b^{\top}$. So, $x \notin a^{\top}$. Since $a \vee x \neq 1$ and x is dual atom, we have $a \leq x$. Thus $x \in X_a$. Thus $X_b \subseteq X_a$. Therefore, $a = \bigwedge X_a \leq \bigwedge X_b = b$.

(2) Let $a \in A$ and $b := \bigwedge X_a$. By Proposition 2.3, $b \in \text{Bo}(\mathbf{A})$. Since $a \leq b$ we have $a^{\top} \subseteq b^{\top}$. Let $x \in b^{\top}$. Thus,

$$1 = b \lor x = \left(\bigwedge X_a \right) \lor x = \bigwedge \{ y \lor x \colon y \in X_a \}.$$

Then $y \lor x = 1$, for all $y \in X_a$. Suppose that $a \lor x < 1$. So, there exists $y \in \operatorname{At}_{d}(\mathbf{A})$ such that $a \lor x \le y < 1$. Thus $a \le y$ and $x \le y$. It follows that $y \in X_a$. Then $y = y \lor x = 1$, which is a contradiction. Hence, $a \lor x = 1$. Thus $x \in a^{\top}$. Then, we obtain that $b^{\top} \subseteq a^{\top}$. Therefore, $a^{\top} = b^{\top} = (\bigwedge X_a)^{\top}$.

Given a finite distributive nearlattice **A** we define the map $\pi_A \colon A \to \mathrm{Bo}(\mathbf{A})$ as follows: for every $a \in A$,

$$\pi_A(a) = \bigwedge X_a.$$

From Proposition 2.3, it follows that π_A is well defined.

Remark 3.3. Notice that $a \leq \pi_A(a)$, for all $a \in A$. Moreover, for each $a \in A$, $\pi_A(a)$ is the least boolean element b such that $a \leq b$. Thus, we also have that $\pi_A(a) = a$, for all $a \in Bo(\mathbf{A})$.

Proposition 3.4. Let **A** be a finite distributive nearlattice. Then, the map $\pi_A : A \to Bo(\mathbf{A})$ is an onto N-homomorphism.

Proof. Let $a, b \in A$. By definition and Lemma 2.4, we have $\pi_A(a \vee b) = \bigwedge X_{a \vee b} = \bigwedge (X_a \cap X_b)$. Since $X_a \cap X_b \subseteq X_a, X_b$, it follows that $\pi_A(a) \vee \pi_A(b) \leq \pi_A(a \vee b)$. By Remark 3.3, we have $a \leq \pi_A(a)$ and $b \leq \pi_A(b)$. Thus $a \vee b \leq \pi_A(a) \vee \pi_A(b)$. Then, since $\pi_A(a) \vee \pi_A(b)$ is a boolean element, it follows by Remark 3.3 that $\pi_A(a \vee b) \leq \pi_A(a) \vee \pi_A(b)$. Therefore, $\pi_A(a \vee b) = \pi_A(a) \vee \pi_A(b)$.

Now assume that $a \wedge b$ exists in **A**. Then,

$$\pi_A(a \wedge b) = \bigwedge X_{a \wedge b} = \bigwedge (X_a \cup X_b) = \bigwedge X_a \wedge \bigwedge X_b = \pi_A(a) \wedge \pi_A(b).$$

Finally, $\pi_A(1) = \bigwedge X_1 = \bigwedge \emptyset = 1$.

It is well known that if **L** is a bounded distributive lattice, then the subset $\operatorname{Comp}(\mathbf{L})$ of all complemented elements of L form a boolean algebra. Given that in a distributive nearlattice may not exists the least element, we generalize the concept of complemented element as follows. Recall that \veebar denotes the supremum in $\operatorname{Fi}(\mathbf{A})$ and $a^{\top} \in \operatorname{Fi}(\mathbf{A})$, for all $a \in A$.

Definition 3.5. Let **A** be a distributive nearlattice. An element $a \in A$ is said to be *complemented* if $[a) \veebar a^{\top} = A$.

We denote by $C(\mathbf{A})$ the set of all complemented elements of \mathbf{A} .

Proposition 3.6. If **A** is a bounded distributive lattice, then $Comp(\mathbf{A}) = \mathcal{C}(\mathbf{A})$.

Proof. Let $a \in \mathcal{C}(\mathbf{A})$. So $[a) \veebar a^\top = A$. Let 0 be the least element of A. Since $0 \in [a) \veebar a^\top$, there is $b \in a^\top$ such that $a \land b = 0$. Thus, we have $a \lor b = 1$ and $a \land b = 0$. Then, b is the complement of a in A. Hence $a \in \operatorname{Comp}(A)$.

Let now $a \in \text{Comp}(A)$. Let $a^* \in A$ be the complement of a. Thus $a \wedge a^* = 0$ and $a \vee a^* = 1$. Let $x \in A$. Then $(a \vee x) \wedge (a^* \vee x) = (a \wedge a^*) \vee x = x$. Since $x \vee a \in [a)$ and $a^* \vee x \in a^{\top}$, it follows that $x \in [a) \vee a^{\top}$. Hence $[a) \vee a^{\top} = A$. Therefore, $a \in \mathcal{C}(\mathbf{A})$.

Proposition 3.7. Let $\mathbf{A} = \langle A, \vee, 1 \rangle$ be a distributive nearlattice. Then $\langle \mathcal{C}(\mathbf{A}), \vee, 1 \rangle$ is a nearlattice subalgebra of \mathbf{A} .

Proof. Let $a, b \in \mathcal{C}(\mathbf{A})$. Thus $[a) \vee a^{\top} = [b) \vee b^{\top} = A$. Then,

$$[a \lor b) \veebar (a \lor b)^{\top} = ([a) \cap [b)) \veebar (a \lor b)^{\top} = ([a) \veebar (a \lor b)^{\top}) \cap ([b) \veebar (a \lor b)^{\top}).$$

Since $a \leq a \vee b$, we have $a^{\top} \subseteq (a \vee b)^{\top}$. Thus $A = [a) \vee a^{\top} \subseteq [a) \vee (a \vee b)^{\top}$. Then $A = [a) \vee (a \vee b)^{\top}$. Analogously, we have $A = [b) \vee (a \vee b)^{\top}$. It follows $[a \vee b) \vee (a \vee b)^{\top} = A$. Hence, $a \vee b \in \mathcal{C}(\mathbf{A})$.

Assume now that $a \wedge b$ exists in A. Then,

$$[a \wedge b) \veebar (a \wedge b)^{\top} = ([a) \veebar [b)) \veebar (a^{\top} \cap b^{\top})$$
$$= ([a) \veebar [b) \veebar a^{\top}) \cap ([a) \veebar [b) \veebar b^{\top}) = A.$$

Hence $a \wedge b \in \mathcal{C}(\mathbf{A})$. Therefore, $\langle \mathcal{C}(\mathbf{A}), \vee, 1 \rangle$ is a nearlattice subalgebra of \mathbf{A} .

Proposition 3.8. Let $\mathbf{A} = \langle A, \vee, 1 \rangle$ be a distributive nearlattice. Then $\langle \mathcal{C}(\mathbf{A}), \vee, 1 \rangle$ is a semi-boolean algebra.

Proof. By the previous proposition we know that $\langle \mathcal{C}(\mathbf{A}), \vee, 1 \rangle$ is a distributive nearlattice. Thus, $[a)_{\mathcal{C}(\mathbf{A})}$ is a bounded distributive lattice, for all $a \in \mathcal{C}(\mathbf{A})$. Let $a \in \mathcal{C}(\mathbf{A})$. Let us show that each element of the lattice $[a)_{\mathcal{C}(\mathbf{A})}$ has a complement. Let $b \in [a)_{\mathcal{C}(\mathbf{A})}$. So $a \leq b$ and $b \in \mathcal{C}(\mathbf{A})$. Then $[b) \veebar b^\top = A$. Given that $a \in [b) \veebar b^\top$, there exist $x \in [b)$ and $y \in b^\top$ such that $a = x \wedge y$. Thus $a \leq y$ and $y \vee b = 1$. Notice that $a \leq b, y$. Then $a \leq b \wedge y$. Since $b \leq x$, it follows that $b \wedge y \leq x \wedge y = a$. Hence $a = b \wedge y$. Now we show that $y \in \mathcal{C}(\mathbf{A})$. Since $a = b \wedge y$, we obtain that $[a) = [b) \veebar [y)$ and $a^\top = b^\top \cap y^\top$. Moreover, since $b \in y^\top$, it follows that $[b) \subseteq y^\top$. Then, we have

$$\begin{split} A &= [a) \veebar a^\top = ([b) \veebar [y)) \veebar \left(b^\top \cap y^\top\right) = \left([b) \veebar [y) \veebar b^\top\right) \cap \left([b) \veebar [y) \veebar y^\top\right) \\ &= [y) \veebar y^\top. \end{split}$$

Hence, we obtain that $y \in [a)_{\mathcal{C}(\mathbf{A})}$, $b \vee y = 1$ and $b \wedge y = a$. That is, y is the complement of b in $[a)_{\mathcal{C}(\mathbf{A})}$. Then $[a)_{\mathcal{C}(\mathbf{A})}$ is a boolean algebra. Therefore, $\mathcal{C}(\mathbf{A})$ is a semi-boolean algebra.

Proposition 3.9 ([6]). Let **A** be a distributive nearlattice. Then, **A** is a semi-boolean algebra if and only if $A = \mathcal{C}(\mathbf{A})$.

Proposition 3.10. Let **A** be a finite distributive nearlattice. Then, **A** is a semi-boolean algebra if and only if $Bo(\mathbf{A}) = \mathcal{C}(\mathbf{A})$.

Proof. Assume that **A** is a semi-boolean algebra. Then, $A = \operatorname{Bo}(\mathbf{A})$ and $A = \mathcal{C}(\mathbf{A})$. Hence $\operatorname{Bo}(\mathbf{A}) = \mathcal{C}(\mathbf{A})$. Conversely, assume that $\operatorname{Bo}(\mathbf{A}) = \mathcal{C}(\mathbf{A})$. Let $a \in A$. Then, by Proposition 3.2, $a^{\top} = b^{\top}$, where $b = \bigwedge X_a$. Thus, $b \in \operatorname{Bo}(\mathbf{A}) = \mathcal{C}(\mathbf{A})$ and $a \leq b$. Then $A = [b) \vee b^{\top} \subseteq [a) \vee b^{\top} = [a) \vee a^{\top}$. Thus $a \in \mathcal{C}(\mathbf{A})$. Hence $A = \mathcal{C}(\mathbf{A})$. Therefore, **A** is a semi-boolean algebra.

4. Connection with the free distributive lattice extension

Definition 4.1. Let **A** be a distributive nearlattice. A pair $\langle L(\mathbf{A}), e \rangle$, where $L(\mathbf{A})$ is a bounded distributive lattice and $e \colon A \to L(\mathbf{A})$ is an N-embedding, is said to be a *free distributive lattice extension* of **A** if e[A] is finitely meet-dense in $L(\mathbf{A})$ and the following universal property holds: for every bounded distributive lattice **M** and every N-homomorphism $h \colon A \to M$, there exists a unique lattice homomorphism $\widehat{h} \colon L(\mathbf{A}) \to M$ such that $h = \widehat{h} \circ e$.

Lemma 4.2. Let **A** be a distributive nearlattice and let $\langle L(\mathbf{A}), e \rangle$ be the free distributive lattice extension of **A**. The following properties are satisfied:

(1)
$$e[[a]] = [e(a)].$$

(2) $\operatorname{Fig}_{L(\mathbf{A})}(e[a^{\top}]) = e(a)^{\top}.$

Proof. (1) It is clear that $e[[a]] \subseteq [e(a))$. Conversely, if $u \in [e(a))$, then $e(a) \leq u$. Since e[A] is finitely meet-dense in $L(\mathbf{A})$, there exist $x_1, \ldots, x_n \in A$ such that $u = e(x_1) \wedge \ldots \wedge e(x_n)$. Then $e(a) \leq e(x_i)$, for all $i \in \{1, \ldots, n\}$. Since e is injective, we have $a \leq x_i, \forall i \in \{1, \ldots, n\}$. Then, there exists $y := x_1 \wedge \ldots \wedge x_n \in [a]$ such that e(y) = u. Hence $u \in e[[a]]$. Therefore, e[[a]] = [e(a)).

(2) We know that $e(a)^{\top}$ is a filter of $L(\mathbf{A})$, and it is clear that $e\left[a^{\top}\right] \subseteq e(a)^{\top}$. Then, we have $\mathrm{Fig}_{L(\mathbf{A})}\left(e\left[a^{\top}\right]\right) \subseteq e(a)^{\top}$. Let now $u \in e(a)^{\top}$. So $u \vee e(a) = 1_{L(\mathbf{A})}$.

Let $a_1, \ldots, a_n \in A$ be such that $u = e(a_1) \wedge \cdots \wedge e(a_n)$. Thus $1_{L(\mathbf{A})} = u \vee e(a) = (e(a_1) \vee e(a)) \wedge \cdots \wedge (e(a_n) \vee e(a))$. Then $1_{L(\mathbf{A})} = e(a_i \vee a)$ for all i. Hence $a_i \vee a = 1$ for all i. That is, $a_1, \ldots, a_n \in a^{\top}$. Thus, $e(a_1), \ldots, e(a_n) \in e\left[a^{\top}\right]$. Then $u = e(a_1) \wedge \cdots \wedge e(a_n) \in \operatorname{Fig}_{L(\mathbf{A})}\left(e\left[a^{\top}\right]\right)$. Hence, $\operatorname{Fig}_{L(\mathbf{A})}\left(e\left[a^{\top}\right]\right) = e(a)^{\top}$.

Proposition 4.3. Let **A** be a finite distributive nearlattice and $\langle L(\mathbf{A}), e \rangle$ its free distributive nearlattice extension. Then,

- (1) $e\left[\operatorname{At}_{d}(\mathbf{A})\right] = \operatorname{At}_{d}(L(\mathbf{A}));$
- (2) $e[Bo(\mathbf{A})] \subseteq Bo(L(\mathbf{A}));$
- (3) $e[\mathcal{C}(\mathbf{A})] \subseteq \text{Comp}(L(\mathbf{A})).$

Proof. (1) Let $a \in \operatorname{At}_{\operatorname{d}}(\mathbf{A})$. Let $u \in L(\mathbf{A})$ be such that $e(a) \leq u \leq 1_{L(\mathbf{A})}$ (recall that $1_{L(\mathbf{A})} = e(1)$). Thus, there are $a_1, \ldots, a_n \in A$ such that $u = e(a_1) \wedge \cdots \wedge e(a_n)$. Since $e(a) \leq u \leq e(a_i)$ for all i, we obtain that $a \leq a_i \leq 1$ for all i. Then, for every i, $a = a_i$ or $a_i = 1$. If $a_i = 1$ for all i, then $u = 1_{L(\mathbf{A})}$. Otherwise, there is $i \in \{1, \ldots, n\}$ such that $a_i = a$. Thus, $e(a) \leq u \leq e(a_i) = e(a)$. Then u = e(a). Hence $e(a) \in \operatorname{At}_{\operatorname{d}}(L(\mathbf{A}))$. Let now $u \in \operatorname{At}_{\operatorname{d}}(L(\mathbf{A}))$. There are $a_1, \ldots, a_n \in A$ such that $u = e(a_1) \wedge \cdots \wedge e(a_n)$. Since $u \neq 1_{L(\mathbf{A})}$, it follows that there is i, such that $a_i \neq 1$. Then, since $u \leq e(a_i) < 1_{L(\mathbf{A})}$, we have $u = e(a_i)$. Moreover, it is straightforward to show directly that $a_i \in \operatorname{At}_{\operatorname{d}}(\mathbf{A})$. Hence $u \in e[\operatorname{At}_{\operatorname{d}}(\mathbf{A})]$.

- (2) Let $a \in \operatorname{Bo}(\mathbf{A})$. It is clear that $\langle [e(a)), \wedge, \vee, e(a), 1_{L(\mathbf{A})} \rangle$ is a bounded distributive sublattice of $L(\mathbf{A})$. We need only to prove that every element of [e(a)) is complemented. Recall that [e(a)) = e[[a)]. Thus, let $e(x) \in [e(a))$ with $x \in [a)$. Since [a) is a boolean lattice, there is $\neg_a x \in [a)$ such that $x \vee \neg_a x = 1$ and $x \wedge_a \neg_a x = a$. Then $1_{L(\mathbf{A})} = e(x \vee \neg_a x) = e(x) \vee e(\neg_a x)$ and $e(a) = e(x \wedge_a \neg_a x) = e(x) \wedge e(\neg_a x)$. Thus, $e(\neg_a x)$ is the complement of e(x) in [e(a)), i.e., $e(\neg_a x) = \neg_{e(a)} e(x)$. Therefore, $e(a) \in \operatorname{Bo}(L(\mathbf{A}))$.
- (3) Let $a \in \mathcal{C}(\mathbf{A})$. So $[a) \veebar a^{\top} = A$. We need to prove that $e(a) \in \text{Comp}(L(\mathbf{A}))$. From Proposition 3.6, it is equivalent to show that $[e(a)) \veebar e(a)^{\top} = L(\mathbf{A})$ (here \veebar is the join of $\text{Fi}(L(\mathbf{A}))$). Since $[a) \veebar a^{\top} = A$, it follows that

$$\operatorname{Fig}_{L(\mathbf{A})}\left(e\left[[a) \veebar a^{\top}\right]\right) = \operatorname{Fig}_{L(\mathbf{A})}\left(e[A]\right) = L(\mathbf{A}).$$

By [11, Thm. 3.3], we know that the map $\Phi \colon \text{Fi}(\mathbf{A}) \to \text{Fi}(L(\mathbf{A}))$ defined by $\Phi(F) = \text{Fig}_{L(\mathbf{A})} (e[F])$ for every $F \in \text{Fi}(\mathbf{A})$, is a lattice isomorphism. Then,

$$\begin{split} \operatorname{Fig}_{L(\mathbf{A})} \left(e \left[[a) \veebar a^\top \right] \right) &= \Phi \left([a) \veebar a^\top \right) = \Phi \left([a) \right) \veebar \Phi \left(a^\top \right) \\ &= \operatorname{Fig}_{L(\mathbf{A})} \left(e \left[[a] \right] \right) \veebar \operatorname{Fig}_{L(\mathbf{A})} \left(e \left[a^\top \right] \right) \\ &= \left[e(a) \right) \veebar \operatorname{Fig}_{L(\mathbf{A})} \left(e \left[a^\top \right] \right). \end{split}$$

Hence, by Lemma 4.2, we have

$$L(\mathbf{A}) = \operatorname{Fig}_{L(\mathbf{A})} \left(e \left[[a] \veebar a^\top \right] \right) = [e(a)) \veebar e(a)^\top.$$

Therefore, we have shown that $e(a) \in \mathcal{C}(L(\mathbf{A}))$.

Remark 4.4. If **A** is a finite distributive nearlattice, then $L(\mathbf{A})$ is a finite distributive lattice. Hence, by Corolary 2.9, it follows that Bo $(L(\mathbf{A}))$ is a boolean lattice and for all $a \in \text{Bo}(\mathbf{A})$, $e(\neg_a x) = \neg e(x) \lor e(a)$ for all $x \in [a)$.

Theorem 4.5. Let **A** be a finite distributive nearlattice and $\langle L(\mathbf{A}), e \rangle$ its free distributive lattice extension. Then $\pi_{L(\mathbf{A})} \circ e = e \circ \pi_A$, i.e., the following diagram

commutes:

$$A \xrightarrow{\pi_A} \operatorname{Bo}(\mathbf{A})$$

$$\downarrow e$$

$$L(\mathbf{A}) \xrightarrow{\pi_{L(\mathbf{A})}} \operatorname{Bo}(L(\mathbf{A}))$$

Proof. Let $a \in A$. By Proposition 4.3, we obtain that

$$\pi_{L(\mathbf{A})}(e(a)) = \bigwedge \{ z \in \operatorname{At}_{d}(L(\mathbf{A})) : e(a) \leq z \}$$

$$= \bigwedge \{ e(x) : x \in \operatorname{At}_{d}(\mathbf{A}), e(a) \leq e(x) \}$$

$$= e \left[\bigwedge \{ x \in \operatorname{At}_{d}(\mathbf{A}) : a \leq x \} \right] = e \left(\pi_{\mathbf{A}}(a) \right). \quad \Box$$

5. A discrete representation

In this section we develop a representation for the class of finite distributive nearlattices through certain ordered structures. This representation is a nice generalization of that given by Birkhoff for finite distributive lattice through finite posets ([4]).

Definition 5.1. Let **A** be a distributive nearlattice. An element $a \in A$ is said to be *meet-irreducible* (or simply *irreducible*) if for all $x, y \in A$ such that $x \wedge y$ exists in A, $a = x \wedge y$ implies a = x or a = y.

Let us denote by $Irr(\mathbf{A})$ the set of all irreducible elements of \mathbf{A} . Notice that $At_d(\mathbf{A}) \subseteq Irr(\mathbf{A})$.

Proposition 5.2. Let **A** be a distributive nearlattice and $a \in A$. Then, a is irreducible if and only if for all $x, y \in A$ such that $x \wedge y$ exists, $x \wedge y \leq a$ implies $x \leq a$ or $y \leq a$.

Theorem 5.3. Let **A** be a finite distributive nearlattice. Then, for every $a \in A$,

$$a = \bigwedge \{x \in Irr(\mathbf{A}) : a \le x\}.$$

Proof. Let $S = \{a \in A : a \neq \bigwedge \{x \in \operatorname{Irr}(\mathbf{A}) : a \leq x\}\}$. We suppose by contradiction that $S \neq \emptyset$. Then, since A is finite, S has a maximal element m. Thus $m \notin \operatorname{Irr}(\mathbf{A})$. Then, there exist $a, b \in A$ such that $m = a \wedge b$, m < a and m < b. Since m is a maximal element of S, it follows that $a, b \notin S$. Thus $a = \bigwedge \{x \in \operatorname{Irr}(\mathbf{A}) : a \leq x\}$ and $b = \bigwedge \{\operatorname{Irr}(\mathbf{A}) : b \leq x\}$. Hence, by Proposition 5.2, we have

$$m = \left(\bigwedge \{ x \in \operatorname{Irr}(\mathbf{A}) : a \le x \} \right) \land \left(\bigwedge \{ x \in \operatorname{Irr}(\mathbf{A}) : b \le x \} \right)$$
$$= \bigwedge \{ x \in \operatorname{Irr}(\mathbf{A}) : m \le x \}.$$

Thus, we obtain that $m \notin S$, which is a contradiction. Therefore, $S = \emptyset$.

Proposition 5.4. Let **A** be a finite distributive nearlattice and $\langle L(\mathbf{A}), e \rangle$ its free distributive lattice extension. Then, $e[\operatorname{Irr}(\mathbf{A})] = \operatorname{Irr}(L(\mathbf{A}))$.

Proof. Let $x \in \text{Irr}(\mathbf{A})$. Let $u, v \in L(\mathbf{A})$ be such that $e(x) = u \wedge v$. There are $a_1, \ldots, a_n, b_1, \ldots, b_m \in A$ such that $u = e(a_1) \wedge \cdots \wedge e(a_n)$ and $v = e(b_1) \wedge \cdots \wedge e(b_m)$. Thus, we have that $e(x) \leq e(a_i), e(b_j)$, for all $i = 1, \ldots, n$ and $j = 1, \ldots, m$. Then $x \leq a_i, b_j$, for all i, j, which implies that there exists $a_1 \wedge \cdots \wedge a_n \wedge b_1 \wedge \cdots \wedge b_m$.

Then $e(x) = e(a_1 \wedge \cdots \wedge a_n \wedge b_1 \wedge \cdots \wedge b_m)$. So $x = a_1 \wedge \cdots \wedge a_n \wedge b_1 \wedge \cdots \wedge b_m$. Since x is irreducible, it follows that $x = a_1 \wedge \cdots \wedge a_n$ or $x = b_1 \wedge \cdots \wedge b_m$. Then e(x) = u or e(x) = v. Hence $e(x) \in \operatorname{Irr}(L(\mathbf{A}))$. Therefore $e[\operatorname{Irr}(\mathbf{A})] \subseteq \operatorname{Irr}(L(\mathbf{A}))$. Now it is straightforward to show the inclusion $\operatorname{Irr}(L(\mathbf{A})) \subseteq e[\operatorname{Irr}(\mathbf{A})]$.

From now on, given a poset $\langle X, \leq \rangle$, $\mathcal{D}(X)$ will denote the collection of all downsets of X and let us consider the bounded distributive lattice $\langle \mathcal{D}(X), \cap, \cup, \emptyset, X \rangle$.

Let **A** be a finite distributive nearlattice and $\langle L(\mathbf{A}), e \rangle$ its free distributive lattice extension. Let us consider $\operatorname{Irr}(\mathbf{A})$ ($\operatorname{Irr}(L(\mathbf{A}))$) as a sub-poset of A ($L(\mathbf{A})$), that is, $x \leq y$ if and only if $x \vee y = y$, for all $x, y \in \operatorname{Irr}(\mathbf{A})$. Since $L(\mathbf{A})$ is a finite distributive lattice, it follows that the map $\alpha \colon L(\mathbf{A}) \to \mathcal{D}(\operatorname{Irr}(L(\mathbf{A})))$ defined by $\alpha(u) = \{z \in \operatorname{Irr}(L(\mathbf{A})) : u \nleq z\}$ is an isomorphism. Hence, by Proposition 5.4, we have that the map $\widehat{\alpha} \colon L(\mathbf{A}) \to \mathcal{D}(\operatorname{Irr}(\mathbf{A}))$ given by $\widehat{\alpha}(u) = \{x \in \operatorname{Irr}(\mathbf{A}) : u \nleq e(x)\}$ is an isomorphism. Therefore, we have the following.

Proposition 5.5. For every finite distributive nearlattice \mathbf{A} , we have that $\langle \mathcal{D}(\operatorname{Irr}(\mathbf{A})), \widehat{e} \rangle$ is the free distributive lattice extension of \mathbf{A} , where $\widehat{e} \colon A \to \mathcal{D}(\operatorname{Irr}(\mathbf{A}))$ is given by $\widehat{e}(a) = \{\widehat{\alpha} \circ e \mid (a) = \{x \in \operatorname{Irr}(\mathbf{A}) : a \nleq x\}.$

Definition 5.6. A *DN-structure* is a pair $\langle X, \gamma \rangle$ such that X is a poset and $\gamma \colon \mathcal{D}(X) \to \{0, 1\}$ is a map satisfying the following:

- (S1) $\gamma(X) = 1$;
- (S2) $\gamma([x)^c) = 1$, for all $x \in X$;
- (S3) for all $U, V \in \mathcal{D}(X)$, $U \subseteq V$ implies $\gamma(U) \leq \gamma(V)$.

We say that a DN-structure $\langle X, \gamma \rangle$ is *finite* if the poset X is finite. Let $\langle X, \gamma \rangle$ be a DN-structure. We define

$$\mathbf{N}(X) := \{ U \in \mathcal{D}(X) : \gamma(U) = 1 \}.$$

Proposition 5.7. Let $\langle X, \gamma \rangle$ be a finite DN-structure. Then $\langle \mathbf{N}(X), \cup, X \rangle$ is a distributive nearlattice and $\langle X, \leq \rangle \cong \langle \operatorname{Irr}(\mathbf{N}(X)), \subseteq \rangle$.

Proof. First let us show that $\mathbf{N}(X)$ is closed under \cup . Let $U_1, U_2 \in \mathbf{N}(X)$. Thus $\gamma(U_1) = \gamma(U_2) = 1$. Then, since $U_1 \subseteq U_1 \cup U_2$, it follows by (S3) that $\gamma(U_1 \cup U_2) = 1$. Hence $U_1 \cup U_2 \in \mathbf{N}(X)$

Now let $U_1, U_2, V \in \mathbf{N}(X)$ be such that $V \subseteq U_1, U_2$. Thus $V \subseteq U_1 \cap U_2$. By (S3), we have $U_1 \cap U_2 \in \mathbf{N}(X)$. Then, $U_1 \cap U_2$ is the meet of U_1 and U_2 in $[V)_{\mathbf{N}(X)}$. Hence $\langle [V)_{\mathbf{N}(X)}, \cap, \cup, U, X \rangle$ is a bounded distributive lattice. Therefore, $\langle \mathbf{N}(X), \cup, X \rangle$ is a distributive nearlattice.

It is well-known that $\operatorname{Irr}(\mathcal{D}(X)) = \{[x)^c : x \in X\}$, and thus $\langle \operatorname{Irr}(\mathcal{D}(X)), \subseteq \rangle \cong \langle X, \leq \rangle$. Let us prove that $\operatorname{Irr}(\mathbf{N}(X)) = \operatorname{Irr}(\mathcal{D}(X))$. By (S2), we have $\operatorname{Irr}(\mathcal{D}(X)) \subseteq \operatorname{Irr}(\mathbf{N}(X))$. Let now $U \in \operatorname{Irr}(\mathbf{N}(X))$. Since U^c is a finite upset of X, we have that $U^c = [x_1) \cup \cdots \cup [x_n)$, for some $x_1, \ldots, x_n \in U^c$. Thus $U = [x_1)^c \cap \cdots \cap [x_n)^c$. Then, since $U \in \operatorname{Irr}(\mathbf{N}(X))$, we obtain that $U = [x_i)^c$, for some $i = 1, \ldots, n$. Hence $\operatorname{Irr}(\mathbf{N}(X)) \subseteq \operatorname{Irr}(\mathcal{D}(X))$. Therefore, we have that $\langle \operatorname{Irr}(\mathbf{N}(X)), \subseteq \rangle \cong \langle X, \leq \rangle$.

Let $\langle A, \vee, 1 \rangle$ be a finite distributive nearlattice. Let $\mathbf{S}(A) = \langle \operatorname{Irr}(\mathbf{A}), < \rangle$.

Proposition 5.8. Let $\langle A, \vee, 1 \rangle$ be a finite distributive nearlattice. Then, the pair $\langle \mathbf{S}(A), \gamma_A \rangle$ is a DN-structure, where $\gamma_A \colon \mathcal{D}(\mathbf{S}(A)) \to \{0, 1\}$ is the map defined by $\gamma_A(U) = 1$ if and only if $\bigwedge (\mathbf{S}(A) \setminus U)$ exists in A.

Proof. It is straightforward to show directly that the map γ_A satisfies conditions (S1)–(S3).

Given a finite distributive nearlattice \mathbf{A} , we have that $\langle \mathbf{N}(\mathbf{S}(A)), \cup, \mathbf{S}(A) \rangle$ is a finite distributive nearlattice, where $\mathbf{N}(\mathbf{S}(A)) = \{U \in \mathcal{D}(\mathbf{S}(A)) : \gamma_A(U) = 1\}$.

Theorem 5.9 (Discrete representation). Let $\mathbf{A} = \langle A, \vee, 1 \rangle$ be a finite distributive nearlattice. Then, $\mathbf{A} \cong \mathbf{N}(\mathbf{S}(A))$.

Proof. From Proposition 5.5, we have that the map $\widehat{e}: A \to \mathcal{D}(\mathbf{S}(A))$ is an N-embedding, where $\widehat{e}(a) = \{x \in \operatorname{Irr}(\mathbf{A}) : a \nleq x\}$, for every $a \in A$. Thus, $\mathbf{A} \cong \widehat{e}[A]$. Let us show that $\widehat{e}[A] = \mathbf{N}(\mathbf{S}(A))$. Then,

$$U \in \mathbf{N}(\mathbf{S}(A)) \iff U \in \mathcal{D}(\mathbf{S}(A)) \text{ and } \gamma_A(U) = 1$$

$$\iff U \in \mathcal{D}(\mathbf{S}(A)) \text{ and } \bigwedge \mathbf{S}(A) \setminus U \text{ exists in } A$$

$$\iff \text{there exists } a \in A \text{ such that } U = \{x \in \operatorname{Irr}(\mathbf{A}) : a \nleq x\}$$

$$\iff \text{there exists } a \in A \text{ such that } U = \widehat{e}(a)$$

$$\iff U \in \widehat{e}[A].$$

Hence, $\mathbf{A} \cong \widehat{e}[A] = \mathbf{N}(\mathbf{S}(A)).$

Remark 5.10. Let **A** be a finite distributive nearlattice. If **A** is in fact a lattice, then it follows that $\mathbf{N}(\mathbf{S}(\mathbf{A})) = \mathcal{D}(\mathbf{S}(\mathbf{A}))$. Thus $\mathbf{A} \cong \mathcal{D}(\mathbf{S}(\mathbf{A}))$. Moreover, it is clear that the finite posets are in one-to-one correspondence with the DN-structures $\langle X, \gamma \rangle$ such that $\gamma(U) = 1$, for all $U \in \mathcal{D}(X)$. Therefore, from the representation above established we can obtain the representation given by Birkhoff for finite distributive lattices.

ACKNOWLEDGMENTS

This work was partially supported by ANPCyT (Argentina) under the Grant PICT-2019-00882 and by CONICET (Argentina) under the Grant PIP 112-201501-00412. The first author was also partially supported by ANPCyT under the Grant PICT-2019-00674 and by Universidad Nacional de La Pampa under the Grant P.I. No 78M, Res. 523/19.

References

- [1] J. Abbott. Semi-boolean algebra. Matematički Vesnik, 4(19):177–198, 1967.
- [2] J. Araújo and M. Kinyon. Independent axiom systems for nearlattices. Czech. Math. J., 61(4):975-992, 2011.
- [3] R. Balbes and P. Dwinger. Distributive lattices. University of Missouri Press, 1974.
- [4] G. Birkhoff. Rings of sets. Duke Math. J., 3(3):443-454, 1937.
- [5] I. Calomino. Supremo álgebra distributivas: una generalización de las álgebra de Tarski. PhD thesis, Universidad Nacional del Sur, 2015.
- [6] I. Calomino. Note on α -filters in distributive near lattices. *Mathematica Bohemica*, 144(3):241–250, 2019.
- [7] I. Calomino and S. Celani. A note on annihilators in distributive nearlattices. Miskolc Math. Notes, 16(1):65-78, 2015.
- [8] I. Calomino, S. Celnai, and L. J. González. Quasi-modal operators on distributive nearlattices. Rev. Un. Mat. Argentina, In Press, 2019.
- [9] I. Calomino and L. J. González. Remarks on normal distributive nearlattices. Quaest. Math., pages 1–12, 2020.
- [10] S. Celani and I. Calomino. Stone style duality for distributive nearlattices. Algebra universalis, 71(2):127–153, 2014.

- [11] S. Celani and I. Calomino. On homomorphic images and the free distributive lattice extension of a distributive nearlattice. Rep. Math. Logic, 51:57–73, 2016.
- [12] S. Celani and I. Calomino. Distributive near lattices with a necessity modal operator. Math. Slovaca, 69:35–52, 2019.
- [13] I. Chajda, R. Halaš, and J. Kühr. Semilattice structures, volume 30 of Research and Exposition in Mathematics. Heldermann Verlag, Lemgo, 2007.
- [14] I. Chajda and M. Kolařík. Ideals, congruences and annihilators on nearlattices. Acta Univ. Palacki. Olomuc. Fac. rer. nat. Mathematica, 46(1):25–33, 2007.
- [15] I. Chajda and M. Kolařík. Nearlattices. Discrete Math., 308(21):4906-4913, 2008.
- [16] B. Davey and H. Priestley. Introduction to lattices and order. Cambridge University Press, 2002.
- [17] L. J. González. The logic of distributive nearlattices. Soft Computing, 22(9):2797–2807, 2018.
- [18] L. J. González. Selfextensional logics with a distributive nearlattice term. Arch. Math. Logic, 58:219–243, 2019.
- [19] L. J. González and I. Calomino. A completion for distributive nearlattices. Algebra Universalis, 80: 48, 2019.
- [20] G. Grätzer. Lattice theory: foundation. Springer Science & Business Media, 2011.
- [21] R. Halaš. Subdirectly irreducible distributive near lattices. *Miskolc Math. Notes*, 7:141–146, 2006.

Universidad Nacional de La Pampa. Facultad de Ciencias Exactas, y Naturales. Santa Rosa, Argentina.

 $Email\ address: {\tt lucianogonzalez@exactas.unlpam.edu.ar}$

CIC and Departamento de Matemática, Facultad de Ciencias Exactas, Universidad Nacional del Centro, Pinto 399, 7000 Tandil, Argentina.

Email address: calomino@exa.unicen.edu.ar