



**HAL**  
open science

## Packing and covering immersion-expansions of planar sub-cubic graphs

Archontia C. Giannopoulou, O-Joung Kwon, Jean-Florent Raymond,  
Dimitrios M. Thilikos

► **To cite this version:**

Archontia C. Giannopoulou, O-Joung Kwon, Jean-Florent Raymond, Dimitrios M. Thilikos. Packing and covering immersion-expansions of planar sub-cubic graphs. *European Journal of Combinatorics*, 2017, 65, pp.154-167. 10.1016/j.ejc.2017.05.009 . lirmm-01610005v1

**HAL Id: lirmm-01610005**

**<https://hal-lirmm.ccsd.cnrs.fr/lirmm-01610005v1>**

Submitted on 4 Oct 2017 (v1), last revised 12 Jun 2018 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Contents lists available at ScienceDirect

European Journal of Combinatorics

journal homepage: [www.elsevier.com/locate/ejc](http://www.elsevier.com/locate/ejc)

# Packing and covering immersion-expansions of planar sub-cubic graphs<sup>☆</sup>



Archontia C. Giannopoulou<sup>a</sup>, O-joung Kwon<sup>a</sup>,  
Jean-Florent Raymond<sup>b,c</sup>, Dimitrios M. Thilikos<sup>c,d</sup>

<sup>a</sup> Logic and Semantics, Technische Universität Berlin, Berlin, Germany

<sup>b</sup> Institute of Computer Science, University of Warsaw, Poland

<sup>c</sup> AIGCo project team, CNRS, LIRMM, Montpellier, France

<sup>d</sup> Department of Mathematics, National and Kapodistrian University of Athens, Greece

## ARTICLE INFO

### Article history:

Received 1 August 2016

Accepted 24 May 2017

Available online 24 June 2017

## ABSTRACT

A graph  $H$  is an immersion of a graph  $G$  if  $H$  can be obtained by some subgraph  $G'$  after lifting incident edges. We prove that there is a polynomial function  $f : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ , such that if  $H$  is a connected planar sub-cubic graph on  $h > 0$  edges,  $G$  is a graph, and  $k$  is a non-negative integer, then either  $G$  contains  $k$  vertex/edge-disjoint subgraphs, each containing  $H$  as an immersion, or  $G$  contains a set  $F$  of  $f(k, h)$  vertices/edges such that  $G \setminus F$  does not contain  $H$  as an immersion.

© 2017 Elsevier Ltd. All rights reserved.

## 1. Introduction

All graphs in this paper are finite, undirected, loopless, and may have multi-edges. Whenever we call a graph *simple* we also assume that it does not have multi-edges. Let  $\mathcal{C}$  be a class of graphs. A  $\mathcal{C}$ -vertex/edge cover of  $G$  is a set  $S$  of vertices/edges such that each subgraph of  $G$  that is isomorphic to a graph in  $\mathcal{C}$  contains some element of  $S$ . A  $\mathcal{C}$ -vertex/edge packing of  $G$  is a collection of vertex/edge-disjoint subgraphs of  $G$ , each isomorphic to some graph in  $\mathcal{C}$ .

<sup>☆</sup> An extended abstract of this paper appeared in the proceedings of WG 2016. Archontia C. Giannopoulou and O-joung Kwon have been supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (ERC consolidator grant DISTRUCT, agreement No 648527). Part of the research of Archontia C. Giannopoulou took place while she was a post-doc at the Warsaw Center of Mathematics and Computer Science. O-joung Kwon has been supported by ERC Starting Grant PARAMTIGHT (No. 280152). Jean-Florent Raymond has been supported by the (Polish) National Science Centre grant PRELUDIUM 2013/11/N/ST6/02706. Dimitrios M. Thilikos has been supported by the ANR project DEMOGRAPH (ANR-16-CE40-0028).

E-mail addresses: [archontia.giannopoulou@gmail.com](mailto:archontia.giannopoulou@gmail.com) (A.C. Giannopoulou), [ojoungkwon@gmail.com](mailto:ojoungkwon@gmail.com) (O.-j. Kwon), [jean-florent.raymond@mimuw.edu.pl](mailto:jean-florent.raymond@mimuw.edu.pl) (J.-F. Raymond), [sedthilk@thilikos.info](mailto:sedthilk@thilikos.info) (D.-M. Thilikos).

We say that a graph class  $\mathcal{C}$  has the *vertex/edge Erdős–Pósa property* (shortly *v/e-E &P property*) for some graph class  $\mathcal{G}$  if there is a function  $f : \mathbb{N} \rightarrow \mathbb{N}$ , called a *gap function*, such that, for every graph  $G$  in  $\mathcal{G}$  and every non-negative integer  $k$ , either  $G$  has a vertex/edge  $\mathcal{C}$ -packing of size  $k$  or  $G$  has a vertex/edge  $\mathcal{C}$ -cover of size  $f(k)$ . In the case where  $\mathcal{G}$  is the class of all graphs we simply say that  $\mathcal{C}$  has the v/e-E &P property. An interesting topic in Graph Theory, related to the notion of duality between graph parameters, is to detect instantiations of  $\mathcal{C}$  and  $\mathcal{G}$  such that  $\mathcal{C}$  has the v/e-E &P property for  $\mathcal{G}$  and, optimize the corresponding gap. Certainly, the first result of this type was the celebrated result of Erdős and Pósa in [11] who proved that the class of all cycles has the v-E &P property with gap function  $O(k \cdot \log k)$ . This result has triggered a lot of research on its possible extensions. One of the most general ones was given in [24] where it was proven that the class of graphs that are contractible to some graph  $H$  has the v-E &P property iff  $H$  is planar (see also [4,5,8] for improvements on the gap function).

Other instantiations of  $\mathcal{C}$  for which the v-E &P property has been proved concern odd cycles [18,21], long cycles [2], and graphs containing cliques as minors [9] (see also [14,16,23] for results on more general combinatorial structures).

As noticed in [8], cycles have the e-E &P property as well. Interestingly, only few more results exist for the cases where the e-E &P property is satisfied. It is known for instance that graphs contractible to  $\theta_r$  (i.e. the graph consisting of two vertices and an edge of multiplicity  $r$  between them) have the e-E &P property [3]. Moreover it was proven that odd cycles have the e-E &P property for planar graphs [19] and for 4-edge-connected graphs [18].

Given two graphs  $G$  and  $H$ , we say that  $H$  is an *immersion* of  $G$  if  $H$  can be obtained from some subgraph of  $G$  by lifting incident edges (see Section 2 for the definition of the lift operation). Given a graph  $H$ , we denote by  $\mathcal{I}(H)$  the set of all graphs that contain  $H$  as an immersion. Using this terminology, the edge variant of the original result of Erdős and Pósa in [11] implies that the class  $\mathcal{I}(\theta_2)$  has the v-E &P property (and, according to [8], the e-E &P property as well). A natural question is whether this can be extended for  $\mathcal{I}(H)$ , for other graphs  $H$ , different than  $\theta_2$ . This is the question that we consider in this paper. A distinct line of research is to identify the graph classes  $\mathcal{G}$  such that for every graph  $H$ ,  $\mathcal{I}(H)$  has the e-E &P property for  $\mathcal{G}$ . In this direction, it was recently proved in [20] that for every graph  $H$ ,  $\mathcal{I}(H)$  has the e-E &P property for 4-edge-connected graphs.

In this paper we show that if  $H$  is non-trivial (i.e., has at least one edge), connected, planar, and sub-cubic, i.e., each vertex is incident with at most 3 edges, then  $\mathcal{I}(H)$  has the v/e-E &P property (with polynomial gap in both cases). More concretely, our main result is the following.

**Theorem 1.** *Let  $k \in \mathbb{N}$ . If  $H$  is a connected planar sub-cubic graph and  $G$  is a graph without any  $\mathcal{I}(H)$ -vertex/edge packing of size greater than  $k$  then  $G$  has a  $\mathcal{I}(H)$ -vertex/edge cover of size bounded by a polynomial function of  $h$  and  $k$ , where  $h = |E(H)|$ .*

The main tools of our proof are the graph invariants of tree-cut width and tree-partition width, defined in [28] and [10] respectively (see Section 2 for the formal definitions). Our proof uses the fact that every graph of polynomially (on  $k$ ) big tree-cut width contains a wall of height  $k$  as an immersion (as proved in [28]). This permits us to consider only graphs of bounded tree-cut width and, by applying suitable reductions, we finally reduce the problem to graphs of bounded tree-partition width (Theorem 2). The result follows as we next prove that for every  $H$ , the class  $\mathcal{I}(H)$  has the e-E &P property for graphs of bounded tree-partition width (Theorem 3).

One might conjecture that the result in Theorem 1 is tight in the sense that both being planar and sub-cubic are necessary for  $H$  in order for  $\mathcal{I}(H)$  to have the e-E &P property. In this direction, in Section 7, we give counterexamples for the cases where  $H$  is planar but not sub-cubic and is sub-cubic but not planar.

## 2. Definitions and preliminary results

We use  $\mathbb{N}^+$  for the set of all positive integers and we set  $\mathbb{N} = \mathbb{N}^+ \cup \{0\}$ . Given a function  $f : A \rightarrow B$  and a set  $C \subseteq A$ , we denote by  $f|_C = \{(x, f(x)) \mid x \in C\}$ .

## Graphs

As already mentioned, we deal with loopless graphs where multi-edges are allowed. Given a graph  $G$ , we denote by  $V(G)$  its set of vertices and by  $E(G)$  its multiset of edges. The notation  $|E(G)|$  stands for the total number of edges, that is, counting multiplicities. We use the term *multi-edge* to refer to a 2-element set of adjacent vertices and the term *edge* to deal with one particular instantiation of the multi-edge connecting two vertices. The function  $\text{mult}_G$  maps a set of two vertices of  $G$  to the multiplicity of the edge connecting them, or zero if they are not adjacent. If  $\text{mult}_G(\{u, v\}) = k$  for some  $k \in \mathbb{N}^+$ , we denote by  $\{u, v\}_1, \dots, \{u, v\}_k$  the distinct edges connecting  $u$  and  $v$ . For the sake of clarity, we identify a multi-edge of multiplicity one and its edge and write  $\{u, v\}$  instead of  $\{u, v\}_1$  when  $\text{mult}_G(\{u, v\}) = 1$ .

For  $X \subseteq V(G)$ , we denote by  $G \setminus X$  the graph obtained by removing  $X$  and all edges incident to a vertex in  $X$ , and for  $x \in V(G)$ , we simply use  $G \setminus x$  to refer to  $G \setminus \{x\}$ . Similarly, for  $F \subseteq E(G)$ , we denote by  $G \setminus F$  the graph obtained by removing edges in  $F$ , and for  $e \in E(G)$ , we simply use  $G \setminus e$  to refer to  $G \setminus \{e\}$ .

We denote by  $\text{deg}_G(v)$  the *degree* of a vertex  $v$  in a graph  $G$ , that is, the number of vertices that are adjacent to  $v$ . The *multidegree* of  $v$ , that we write  $\text{mdeg}_G(v)$ , is the number of edges (i.e. counting multiplicities) incident with  $v$ . We drop the subscript when it is clear from the context. We also set  $\text{m}\delta(G) = \min_{v \in V(G)} \text{mdeg}_G(v)$ . We say that a graph  $G$  is *sub-cubic* whenever  $\text{mdeg}_G(v) \leq 3$  for every  $v \in V(G)$ . Two paths are *internally disjoint* if they do not share internal vertices.

## Immersion

Let  $H$  and  $G$  be graphs. We say that  $G$  contains  $H$  as an *immersion* if there is a pair of functions  $(\phi, \psi)$ , called an *H-immersion model*, such that  $\phi$  is an injection of  $V(H) \rightarrow V(G)$  and  $\psi$  sends  $\{u, v\}_i$  to a path of  $G$  between  $\phi(u)$  and  $\phi(v)$ , for every  $\{u, v\} \in E(H)$  and every  $i \in \{1, \dots, \text{mult}_H(\{u, v\})\}$ , in a way such that distinct edges are sent to edge-disjoint paths. This definition of an immersion correspond to what is sometimes called a weak immersion. Every vertex in the image of  $\phi$  is called a *branch vertex*. We will make use of the following easy observation.

**Observation 1.** *Let  $H$  and  $G$  be two graphs. If  $(\phi, \psi)$  is an  $H$ -immersion model in  $G$  then for every vertex  $x$  of  $H$ , we have  $\text{mdeg}_H(x) \leq \text{mdeg}_G(\phi(x))$ .*

An *H-immersion expansion*  $M$  in a graph  $G$  is a subgraph of  $G$  defined as follows:  $V(M) = \phi(V(H)) \cup \bigcup_{e \in E(H)} V(\psi(e))$  and  $E(M) = \bigcup_{e \in E(H)} E(\psi(e))$  for some  $H$ -immersion model  $(\phi, \psi)$  of  $G$ . We call the paths in  $\psi(E(H))$  *certifying paths* of the  $H$ -immersion expansion  $M$ .

We say that two edges are *incident* if they share some endpoint. A *lift* of two incident edges  $e_1 = \{x, y\}$  and  $e_2 = \{y, z\}$  of  $G$  is the operation that removes the edges  $e_1$  and  $e_2$  from the graph and then, if  $x \neq z$ , adds the edge  $\{x, z\}$  (or increases the multiplicity of  $\{x, z\}$  by 1 if this edge already exists). Notice that  $H$  is an immersion of  $G$  if and only if a graph isomorphic to  $H$  can be obtained from some subgraph of  $G$  after applying lifts of incident edges.<sup>1</sup>

The *dissolution* of a vertex of multidegree two of a graph is the operation that deletes this vertex and, if it had two neighbors, adds an edge joining them. The *subdivision* of an edge is the operation that adds a vertex of degree two adjacent to its endpoints and deletes this edge. We say that a graph  $G$  is a *subdivision* of a graph  $H$  if  $G$  can be obtained from  $H$  by repeatedly subdividing edges.

## Packings and coverings

An *H-cover* of  $G$  is a set  $C \subseteq E(G)$  such that  $G \setminus C$  does not contain  $H$  as an immersion. An *H-packing* in  $G$  is a collection of edge-disjoint  $H$ -immersion expansions in  $G$ . We denote by  $\text{pack}_H(G)$  the maximum size of an  $H$ -packing and by  $\text{cover}_H(G)$  the minimum size of an  $H$ -cover in  $G$ .

<sup>1</sup> While we mentioned this definition in the introduction, in the rest of the paper, we adopt the more technical definition of immersion in terms of immersion models as this will facilitate the presentation of the proofs.

Rooted trees

A *rooted tree* is a pair  $(T, s)$  where  $T$  is a tree and  $s \in V(T)$  is a vertex referred to as the *root*. Given a vertex  $x \in V(T)$ , the set of *descendants* of  $x$  in  $(T, s)$ , denoted by  $\text{desc}_{(T,s)}(x)$ , is the set containing each vertex  $w$  such that the unique path from  $w$  to  $s$  in  $T$  contains  $x$ . If  $y$  is a descendant of  $x$  and is adjacent to  $x$ , then it is a *child* of  $x$ . Two vertices of  $T$  are *siblings* if they are children of the same vertex. Given a rooted tree  $(T, s)$  and a vertex  $x \in V(G)$ , the *height* of  $x$  in  $(T, s)$  is the maximum distance between  $x$  and a vertex in  $\text{desc}_{(T,s)}(x)$ .

We now define two types of decompositions of graphs: tree-partitions (cf. [15,26]) and tree-cut decompositions (cf. [28]).

Tree-partitions

We introduce, especially for the needs of our proof, an extension of the parameter of tree-partition width defined in [15,26] to multigraphs, where we consider both the number of edges between the bags and the number of vertices in the bags. A *tree-partition* of a graph  $G$  is a pair  $\mathcal{D} = (T, \mathcal{X})$  where  $T$  is a tree and  $\mathcal{X} = \{X_t\}_{t \in V(T)}$  is a partition of  $V(G)$  such that either  $|V(T)| = 1$  or for every  $\{x, y\} \in E(G)$ , there exists an edge  $\{t, t'\} \in E(T)$  where  $\{x, y\} \subseteq X_t \cup X_{t'}$ . We call the elements of  $\mathcal{X}$  *bags* of  $\mathcal{D}$ . Given an edge  $f = \{t, t'\} \in E(T)$ , we define  $E_f$  as the set of edges with one endpoint in  $X_t$  and the other in  $X_{t'}$ . The *width* of  $\mathcal{D}$  is defined as  $\max\{|X_t|_{t \in V(T)} \cup \{|E_f|\}_{f \in E(T)}\}$ . The *tree-partition width* of  $G$  is the minimum width over all tree-partitions of  $G$  and will be denoted by  $\text{tpw}(G)$ . A *rooted tree-partition* of a graph  $G$  is a triple  $\mathcal{D} = ((T, s), \mathcal{X})$  where  $(T, s)$  is a rooted tree and  $(T, \mathcal{X})$  is a tree-partition of  $G$ .

Tree-cut decompositions

A *near-partition* of a set  $S$  is a family of pairwise disjoint subsets  $S_1, \dots, S_k \subseteq S$  (for some  $k \in \mathbb{N}$ ) such that  $\bigcup_{i=1}^k S_i = S$ . Observe that this definition allows some sets  $S_i$  of the family to be empty. A *tree-cut decomposition* of a graph  $G$  is a pair  $\mathcal{D} = (T, \mathcal{X})$  where  $T$  is a tree and  $\mathcal{X} = \{X_t\}_{t \in V(T)}$  is a near-partition of  $V(G)$ . As in the case of tree-partitions, we call the elements of  $\mathcal{X}$  *bags* of  $\mathcal{D}$ . A *rooted tree-cut decomposition* of a graph  $G$  is a triple  $\mathcal{D} = ((T, s), \mathcal{X})$  where  $(T, s)$  is a rooted tree and  $(T, \mathcal{X})$  is a tree-cut decomposition of  $G$ . Given that  $\mathcal{D} = ((T, s), \mathcal{X})$  is a rooted tree-partition or a rooted tree-cut decomposition of  $G$  and given  $t \in V(T)$ , we set  $G_t = G \left[ \bigcup_{u \in \text{desc}_{(T,s)}(t)} X_u \right]$ .

The *torso* of a tree-cut decomposition  $(T, \mathcal{X})$  at a node  $t$  is the graph obtained from  $G$  as follows. If  $V(T) = \{t\}$ , then the torso at  $t$  is  $G$ . Otherwise let  $T_1, \dots, T_\ell$  be the connected components of  $T \setminus t$ . The torso  $H_t$  at  $t$  is obtained from  $G$  by *consolidating* each vertex set  $\bigcup_{b \in V(T_i)} X_b$  into a single vertex  $z_i$ . The operation of consolidating a vertex set  $Z$  into  $z$  is to replace  $Z$  with  $z$  in  $G$ , and for each edge  $e$  between  $Z$  and  $v \in V(G) \setminus Z$ , adding an edge  $\{z, v\}$  in the new graph. Given a graph  $G$  and  $X \subseteq V(G)$ , let the *3-center* of  $(G, X)$  be the unique graph obtained from  $G$  by dissolving vertices in  $V(G) \setminus X$  of multidegree two and deleting vertices of multidegree at most 1. For each node  $t$  of  $T$ , we denote by  $H_t$  the *3-center* of  $(H_t, X_t)$ , where  $H_t$  is the torso of  $(T, \mathcal{X})$  at  $t$ .

Let  $\mathcal{D} = ((T, s), \mathcal{X})$  be a rooted tree-cut decomposition of  $G$ . The *adhesion* of a vertex  $t$  of  $T$ , that we will denote by  $\text{adh}_{\mathcal{D}}(t)$ , is the number of edges of  $G$  with exactly one endpoint in  $G_t$ . The *width* of a tree-cut decomposition  $(T, \mathcal{X})$  of  $G$  is  $\max_{t \in V(T)} \{\text{adh}_{\mathcal{D}}(t), |V(H_t)|\}$ . The *tree-cut width* of  $G$ , denoted by  $\text{tcw}(G)$ , is the minimum width over all tree-cut decompositions of  $G$ .

A vertex  $t \in V(T)$  is *thin* if  $\text{adh}_{\mathcal{D}}(t) \leq 2$ , and *bold* otherwise. We also say that  $\mathcal{D}$  is *nice* if for every thin vertex  $t \in V(T)$  we have  $N(V(G_t)) \cap \bigcup_{b \text{ is a sibling of } t} V(G_b) = \emptyset$ . In other words, there is no edge from a vertex of  $G_t$  to a vertex of  $G_b$ , for any sibling  $b$  of  $t$ , whenever  $t$  is thin. The notion of nice tree-cut decompositions has been introduced by Ganian et al. in [13]. Furthermore, they proved the following result.

**Proposition 1** ([13]). *Every rooted tree-cut decomposition can be transformed into a nice one without increasing the width.*

We say that an edge of  $G$  *crosses* the bag  $X_t$ , for some  $t \in V(T)$  if its endpoints belong to bags  $X_{t_1}$  and  $X_{t_2}$ , for some  $t_1, t_2 \in V(T)$  such that  $t$  belongs to the interior of the (unique) path of  $T$  connecting  $t_1$  to  $t_2$ .

### 3. From tree-cut decompositions to tree-partitions

The purpose of this section is to prove the following theorem. The graph  $H^+$  will be uniquely defined from  $H$  later on.

**Theorem 2.** *For every connected graph  $G$ , and every connected graph  $H$  with at least one edge, there is a graph  $G'$  and a graph  $H^+$  such that*

- $\text{tpw}(G') \leq (\text{tcw}(G) + 1)^2/2$ ,
- $\text{pack}_{H^+}(G') \leq \text{pack}_H(G)$ , and
- $\text{cover}_H(G) \leq \text{cover}_{H^+}(G')$ .

Theorem 2 will allow us in Section 4 to consider graphs of bounded tree-partition width instead of graphs of bounded tree-cut width. Before we proceed with the proof of Theorem 2, we need some definitions and a series of auxiliary results.

For every graph  $G$ , we define  $G^+$  as the graph obtained from  $G$  if, for every vertex  $v$ , we add two new vertices  $v'$  and  $v''$  and the edges  $\{v', v''\}$  (of multiplicity 2),  $\{v, v'\}$  and  $\{v, v''\}$  (both of multiplicity 1). Observe that for every connected graph  $G$  with at least two vertices, we have  $m\delta(G^+) \geq 3$ . We also define  $G^*$  as the graph obtained from  $G$  by adding, for every vertex  $v$ , the new vertices  $v'_1, \dots, v'_{\text{mdeg}(v)}$  and  $v''_1, \dots, v''_{\text{mdeg}(v)}$  and the edges  $\{v'_i, v''_i\}$  (of multiplicity 2),  $\{v, v'_i\}$ , and  $\{v, v''_i\}$  (both of multiplicity 1), for every  $i \in \{1, \dots, \text{mdeg}(v)\}$ . If  $v$  is a vertex of  $G$  then we denote by  $Z_{v,i}$  the subgraph  $G^*[\{v, v'_i, v''_i\}]$ , where  $i \in \{1, \dots, \text{mdeg}_G(v)\}$ .

Our first aim is to prove the following three lemmata.

**Lemma 1.** *Let  $G$  be a graph, let  $H$  be a connected graph with at least one edge and let  $G'$  be a subdivision of  $G^*$ . Then we have*

- $\text{pack}_{H^+}(G^*) = \text{pack}_{H^+}(G')$  and
- $\text{cover}_{H^+}(G^*) = \text{cover}_{H^+}(G')$ .

**Proof.** We denote by  $S$  the set of subdivision vertices added during the construction of  $G'$  from  $G^*$ . As  $G'$  is a subdivision of  $G^*$ , we have  $\text{pack}_{H^+}(G') \geq \text{pack}_{H^+}(G^*)$  and  $\text{cover}_{H^+}(G') \geq \text{cover}_{H^+}(G^*)$ .

As a consequence of Observation 1 and the fact that  $m\delta(H^+) \geq 3$ , if  $M$  is an  $H^+$ -immersion expansion in  $G'$  then no branch vertex of  $M$  belongs to  $S$ . Indeed, every vertex of  $S$  has multidegree 2 in  $G'$ . Therefore, by dissolving in  $M$  the vertices of  $S$  that belong to  $V(M)$ , we obtain an  $H^+$ -immersion expansion in  $G^*$ . It follows that  $\text{pack}_{H^+}(G^*) \geq \text{pack}_{H^+}(G')$ , hence  $\text{pack}_{H^+}(G^*) = \text{pack}_{H^+}(G')$ .

On the other hand, let  $X$  be an  $H^+$ -cover of  $G^*$  and let  $X'$  be a set of edges constructed by taking, for every  $e \in X$ , an edge of the path of  $G'$  connecting the endpoints of  $e$  that has been created by subdividing  $e$ . Assume that  $X'$  is not an  $H^+$ -cover of  $G'$ . According to the remark above, this implies that  $X$  is not an  $H^+$ -cover of  $G^*$ , a contradiction. Hence  $X'$  is an  $H^+$ -cover of  $G'$  and thus  $\text{cover}_{H^+}(G^*) = \text{cover}_{H^+}(G')$ .  $\square$

**Lemma 2.** *For every two graphs  $H$  and  $G$  such that  $H$  is connected and has at least one edge, we have  $\text{pack}_{H^+}(G^*) \leq \text{pack}_H(G)$ .*

**Proof.** In  $G^*$  (respectively  $H^+$ ), we say that a vertex is *original* if it belongs to  $V(G)$  (respectively  $V(H)$ ). Let  $(\phi, \psi)$  be an  $H^+$ -immersion model in  $G^*$ .

We first show that if  $u$  is an original vertex of  $H^+$ , then  $\phi(u)$  is an original vertex of  $G^*$ . By contradiction, let us assume that  $\phi(u)$  is not original, for some original vertex  $u$  of  $H^+$ . Then  $\phi(u) = v'_i$  or  $\phi(u) = v''_i$ , for some  $v \in V(G)$  and  $i \in \{1, \dots, \text{mdeg}_G(v)\}$ .

Observe that since  $H$  is connected and has at least one edge, every vertex of  $H^+$  has degree at least three: let  $x, y$ , and  $z$  be the endpoints of three multi-edges incident with  $u$  (notice that as the degree of  $H^+$  is at least three we can assume that  $x, y$ , and  $z$  are distinct). Then  $\psi(\{u, x\})$ ,  $\psi(\{u, y\})$ , and  $\psi(\{u, z\})$  are edge-disjoint paths connecting  $\phi(u)$  to three distinct vertices. This is not possible because there is an edge cut of size two,  $\{\{v, v'_i\}, \{v, v''_i\}\}$ , separating the two vertices  $v'_i$  and  $v''_i$  (among which is  $\phi(u)$ ) from the rest of the graph. Consequently, if  $u \in V(H^+)$  is original, then  $\phi(u)$  is original.

Let us now consider an edge  $\{u, v\} \in E(H)$ . By the above remark,  $\phi(u)$  and  $\phi(v)$  are original vertices of  $G^*$ . It is easy to see that  $\psi(\{u, v\})$  contains only original vertices of  $G^*$ . Indeed, if this path contained a non-original vertex  $w'$  or  $w''$  for some original vertex  $w$  of  $V(G^*)$ , it would use  $w$  twice in order to reach  $u$  and  $v$ , a contradiction to the fact that it is a path. Therefore, from the definition of  $H^+$ , the pair  $(\phi|_{V(H)}, \psi|_{E(H)})$  is an  $H$ -immersion model of  $G$ .

We proved that every  $H^+$ -immersion-expansion of  $G^*$  contains an  $H$ -immersion-expansion that belongs to the subgraph  $G$  of  $G^*$ . Consequently every  $H^+$ -packing of  $G^*$  contains an  $H$ -packing of the same size that belongs to  $G$ , and the desired inequality follows.  $\square$

**Lemma 3.** For every two graphs  $H$  and  $G$  such that  $H$  is connected and has at least one edge, we have  $\text{cover}_H(G) \leq \text{cover}_{H^+}(G^*)$ .

**Proof.** Similarly to the proof of Lemma 2, we say that an edge of  $G^*$  is *original* if it belongs to  $E(G)$ . Let  $X \subseteq E(G^*)$  be a minimum cover of  $H^+$ -immersion expansions in  $G^*$ .

*First case:* all the edges in  $X$  are original. In this case,  $X$  is an  $H$ -cover of  $G$  as well. Indeed, if  $G \setminus X$  contains an  $H$ -immersion expansion  $M$ , then  $G^* \setminus X$  contains  $M^*$  that, in turn, contains  $H^+$ . Hence in this case,  $\text{cover}_H(G) \leq \text{cover}_{H^+}(G^*)$ .

*Second case:* there is an edge  $e \in X$  that is not original. Let  $v$  be the original vertex of  $G^*$  such that  $e \in Z_{v,i}$  for some  $i \in \{1, \dots, \text{mdeg}_G(v)\}$ . Let us first show the following claim.

**Claim.** For every  $i \in \{1, \dots, \text{mdeg}_G(v)\}$ , there is an edge of  $Z_{v,i}$  that belongs to  $X$ .

**Proof of claim.** Looking for a contradiction, let us assume that we have  $E(Z_{v,i}) \cap X = \emptyset$ , for some  $i \in \{1, \dots, \text{mdeg}_G(v)\}$ . Clearly  $i \neq l$ . By minimality of  $X$ , the graph  $G^* \setminus (X \setminus \{e\})$  contains an  $H^+$ -immersion expansion  $M$  that uses  $e$ . Observe that  $M' = M \setminus E(Z_{v,i}) \cup E(Z_{v,l})$  contains an  $H^+$ -immersion expansion (since  $Z_{v,l}$  and  $Z_{v,i}$  are isomorphic). Hence,  $M'$  is a subgraph of  $G^* \setminus (X \setminus \{e\})$  that contains an  $H^+$ -immersion expansion. This is not possible as  $X$  is a cover, so we reach the contradiction we were looking for and the claim holds.  $\diamond$

We build a set  $Y$  as follows. For every edge  $f \in X$ , if  $f$  is original then we add it to  $Y$ . Otherwise, if  $v_f$  is the (original) vertex of  $G^*$  such that  $f \in E(Z_{v_f,i})$  for some  $i \in \{1, \dots, \text{mdeg}_G(v_f)\}$ , then we add to  $Y$  all edges of  $G$  that are incident to  $v_f$ .

The above claim ensures that when a non-original edge  $f$  of  $X$  is encountered, then  $X$  contains an edge in each of  $Z_{v_f,1}, \dots, Z_{v_f,\text{mdeg}_G(v_f)}$ . Therefore, the same set of edges, of size  $\text{mdeg}_G(v_f)$ , will be added to  $Y$  when encountering another edge from  $Z_{v_f,1}, \dots, Z_{v_f,\text{mdeg}_G(v_f)}$ . Consequently,  $|X| \geq |Y|$ .

Let us now show that  $Y$  is an  $H^+$ -cover of  $G^*$ . Suppose that there exists an  $H^+$ -immersion expansion  $M$  in  $G^* \setminus Y$ . Observe that since  $H$  is connected and has at least one edge,  $M$  does not belong to  $\bigcup_{i \in \{1, \dots, \text{mdeg}_G(u)\}} Z_{u,i}$ , for every original vertex  $u$  of  $G^*$ . Let

$$Z = \bigcup_{u \in V(G)} \bigcup_{i \in \{1, \dots, \text{mdeg}_G(u)\}} E(Z_{u,i}).$$

Then  $M$  is a subgraph of  $G \setminus (Y \cup Z)$ . As  $X \subseteq Y \cup Z$ , this contradicts the fact that  $X$  is a cover. Therefore,  $Y$  is an  $H^+$ -cover. Moreover all the edges in  $Y$  are original. As this situation is treated by the first case above, we are done.  $\square$

We are now ready to prove the main result of this section.

**Proof of Theorem 2.** Let  $k = \text{tcw}(G)$ . We examine the nontrivial case where  $G$  is not a tree, i.e.,  $\text{tcw}(G) \geq 2$ . Let us consider the graph  $G^*$ . We claim that  $\text{tcw}(G^*) = \text{tcw}(G)$ . Indeed, starting from an optimal tree-cut decomposition of  $G$ , we can, for every vertex  $v$  of  $G$  and for every  $i \in \{1, \dots, \text{mdeg}_G(v)\}$ , create a bag that is a child of  $v$  and contains  $\{v'_i, v''_i\}$ . According to the definition of  $G^*$ , this creates a tree-cut decomposition  $\mathcal{D} = ((T, s), \{X_t\}_{t \in V(T)})$  of  $G^*$ . Observe that for every vertex  $x$  that we introduced to the tree of the decomposition during this process,  $\text{adh}_{\mathcal{D}}(x) = 2$  and the corresponding bag has size two. This proves that  $\text{tcw}(G^*) \leq \max(\text{tcw}(G), 2) = \text{tcw}(G)$ . As  $G$  is a subgraph of  $G^*$ , we obtain  $\text{tcw}(G) \leq \text{tcw}(G^*)$  and the proof of the claim is complete.

According to Proposition 1, we can assume that  $G^*$  has a nice rooted tree-cut decomposition of width  $\leq k$ . For notational simplicity we again denote it by  $\mathcal{D} = ((T, s), \{X_t\}_{t \in V(T)})$  and, obviously, we can also assume that all leaves of  $T$  correspond to non-empty bags.

Our next step is to transform the rooted tree-cut decomposition  $\mathcal{D}$  into a rooted tree-partition  $\mathcal{D}' = ((T, s), \{X'_t\}_{t \in V(T)})$  of a subdivision  $G'$  of  $G^*$ . Notice that the only differences between the two decompositions are that, in a tree-cut decomposition, empty bags are allowed as well as edges connecting vertices of bags corresponding to non-adjacent vertices of  $T$ .

We proceed as follows: if  $X$  is a bag crossed by edges, we subdivide every edge crossing  $X$  and add the obtained subdivision vertex to  $X$ . By repeating this process we decrease at each step the number of bags crossed by edges, that eventually reaches zero. Let  $G'$  be the obtained graph and observe that  $G'$  is a subdivision of  $G$ . As  $G$  is connected, the obtained rooted tree-cut decomposition  $\mathcal{D}' = ((T, s), \{X'_t\}_{t \in V(T)})$  is a rooted tree partition of  $G'$ .

Notice that the adhesion of any bag of  $T$  in  $\mathcal{D}$  is the same as in  $\mathcal{D}'$ . However, the bags of  $\mathcal{D}'$  may grow during the construction of  $G'$ . Let  $t$  be a vertex of  $T$  and let  $\{t_1, \dots, t_m\}$  be the set of children of  $t$ . We claim that  $|X'_t| \leq (k + 1)^2/2$ .

Let  $E_t$  be the set of edges crossing  $X_t$  in  $G$ . Let  $H_t$  be the torso of  $\mathcal{D}$  at  $t$ , and let  $H'_t = H_t \setminus X_t$ . Observe that  $|E_t|$  is the same as the number of edges in  $H'_t$ . Let  $z_p$  be the vertex of  $H'_t$  corresponding to the parent of  $t$ , and similarly for each  $i \in \{1, \dots, m\}$  let  $z_i$  be the vertex of  $H'_t$  corresponding to the child  $t_i$  of  $t$ . Notice that if  $t_i$  is a thin child of  $t$ , then  $z_i$  can be adjacent to only  $z_p$  as  $\mathcal{D}$  is a nice rooted tree-cut decomposition. Thus the sum of the number of incident edges with  $z_i$  in  $H'_t$  for all thin children  $t_i$  of  $t$  is at most  $\text{adh}_{\mathcal{D}}(t) \leq k$ . On the other hand, if  $t_i$  is a bold child of  $t$ , then  $z_i$  is incident with at least 3 edges in  $H_t$  (none of which is a child of  $t$ ), and thus it is contained in the 3-center of  $(H_t, X_t)$ . Therefore, the number of all bold children of  $t$  is bounded by  $k - |X_t|$ . Since each vertex in  $H'_t$  is incident with at most  $k$  edges, the total number of edges in  $H'_t$  is at most  $(k - |X_t| + 1)k/2 + k$ . As  $|E(H'_t)| = |E_t| = |X'_t \setminus X_t|$ , it implies that  $|X'_t| \leq |X_t| + k \cdot (k - |X_t| + 2)/2 \leq \max\{2k, k(k + 2)/2\} \leq (k + 1)^2/2$ . We conclude that  $G'$  has a rooted tree-partition of width at most  $(\text{tcw}(G) + 1)^2/2$ .

Recall that  $G'$  is a subdivision of  $G^*$ . By the virtue of Lemma 3, Lemma 2, and Lemma 1, we obtain that  $\text{pack}_{H^+}(G') \leq \text{pack}_H(G)$  and  $\text{cover}_H(G) \leq \text{cover}_{H^+}(G')$ . Hence  $G'$  satisfies the desired properties.  $\square$

#### 4. Erdős–Pósa in graphs of bounded tree-partition width

Before we proceed, we require the following lemma and an easy observation.

**Lemma 4.** *Let  $G$  and  $H$  be two graphs such that  $H$  has no isolated vertices and let  $X \subseteq V(G)$ . Let  $\mathcal{C}$  be the collection of connected components of  $G \setminus X$ . If  $M$  is an  $H$ -immersion expansion of  $G$  then  $M$  contains vertices from at most  $(|X| + 1) \cdot |E(H)|$  graphs of  $\mathcal{C}$ .*

**Proof.** Let  $P$  be a certifying path of  $M$  connecting two branch vertices of  $M$ . Since  $P$  is a path, it cannot use twice the same vertex of  $X$ . Besides, as  $X$  is a separator,  $P$  must go through a vertex of  $X$  in order to go from one graph of  $\mathcal{C}$  to another one. Therefore,  $P$  contains vertices from at most  $|X| + 1$  graphs of  $\mathcal{C}$ . The desired bound follows as  $E(M)$  is partitioned into  $|E(H)|$  certifying paths.  $\square$

**Observation 2.** *Let  $G$  and  $H$  be graphs and let  $F \subseteq E(G)$ . Then it holds that  $\text{cover}_H(G) \leq \text{cover}_H(G \setminus F) + |F|$ .*

For a graph  $H$ , we define  $\omega_H : \mathbb{N} \rightarrow \mathbb{N}$  so that  $\omega_H(r) = \lceil r \cdot \frac{3r+1}{2} \cdot |E(H)| \rceil$ . The next theorem is an important ingredient of our results. It essentially states that  $\mathcal{I}(H)$  has the Erdős–Pósa property in graphs of bounded tree-partition width, for every connected graph  $H$ .

**Theorem 3.** *Let  $H$  be a connected graph with at least one edge. Then for every graph  $G$  it holds that  $\text{cover}_H(G) \leq \omega_H(\text{tpw}(G)) \cdot \text{pack}_H(G)$ .*

**Proof.** Let us show by induction on  $k$  that if  $\text{pack}_H(G) \leq k$  and  $\text{tpw}(G) \leq r$  then  $\text{cover}_H(G) \leq \omega_H(r) \cdot k$ . The case  $k = 0$  is trivial. Let us now assume that  $k \geq 1$  and that for every graph  $G$  of tree-partition width at most  $r$  and such that  $\text{pack}_H(G) = k - 1$ , we have  $\text{cover}_H(G) \leq \omega_H(r)(k - 1)$ . Let  $G$  be a

graph such that  $\mathbf{pack}_H(G) = k$  and  $\mathbf{tpw}(G) \leq r$ . Let also  $\mathcal{D} = ((T, s), \{X_t\}_{t \in V(T)})$  be an optimal rooted tree-partition of  $G$ . We say that a vertex  $t \in V(T)$  is *infected* if  $G_t$  contains an  $H$ -immersion expansion. Recall that the height of a vertex in a rooted tree is the maximum distance to a descendant. Let  $t$  be an infected vertex of  $T$  of minimum height.

**Claim.** *If some of the  $H$ -immersion expansions of  $G$  shares a vertex with  $G_{t'}$  for some child  $t'$  of  $t$ , then it also shares an edge with  $E_{\{t,t'\}}$ .*

**Proof of claim.** Let  $M$  be some  $H$ -immersion expansions of  $G$ . Notice that, by the choice of  $t$ ,  $M$  cannot be entirely inside  $G_{t'}$ . This fact, together with the connectivity of  $M$ , implies that  $E(M) \cap E_{\{t,t'\}} \neq \emptyset$ .  $\diamond$

Suppose that  $M$  is an  $H$ -immersion expansion of  $G_t$  and let  $U$  be the set of children of  $t$  corresponding to bags which share vertices with  $M$ . We define the multisets  $A = E(G[X_t]) \cap E(M)$ ,  $B = \bigcup_{t' \in U} E_{\{t,t'\}}$  and  $C = \bigcup_{t' \in U} E(G_{t'})$ . We also set  $D = A \cup B$ . By the definition of  $U$ , it follows that

$$E(M) \subseteq C \cup D. \tag{1}$$

Let us upper-bound the size of  $|D|$ . Applying Lemma 4 for  $G_t, H$ , and  $X_t$ , we have  $|U| \leq (r+1) \cdot |E(H)|$ , hence  $|B| \leq r(r+1) \cdot |E(H)|$ . Besides, every path of  $M$  connecting two branch vertices meets every vertex of  $X_t$  at most once (as it is a path), thus  $E(M)$  does not contain an edge of  $G[X_t]$  with a multiplicity larger than  $|E(H)|$ . It follows that  $|A| \leq \frac{r(r-1)}{2} \cdot |E(H)|$  and finally we obtain that  $|D| = |A| + |B| \leq r \cdot \frac{3r+1}{2} \cdot |E(H)| \leq \omega_H(r)$ .

Let  $G' = G \setminus D$ . We now show that  $\mathbf{pack}_H(G') \leq k - 1$ . Let us consider an  $H$ -immersion expansion  $M'$  in  $G'$ . As  $E(M') \subseteq E(G) \setminus D$ , it follows that

$$E(M') \cap D = \emptyset. \tag{2}$$

Recall that  $B \subseteq D$ , which together with (2) implies that  $E(M') \cap B = \emptyset$ . This fact, combined with the claim above, implies that

$$E(M') \cap C = \emptyset. \tag{3}$$

From (2) and (3), we obtain that  $E(M') \cap (C \cup D) = \emptyset$ , which, combined with (1), implies that  $E(M) \cap E(M') \neq \emptyset$ . Consequently, every maximum packing of  $H$ -immersion expansions in  $G'$  is edge-disjoint from  $M$ . If such a packing had size  $\geq k$ , it would form, together with  $M$ , a packing of size  $k+1$  in  $G$ , a contradiction. Thus  $\mathbf{pack}_H(G') \leq k - 1$ , as desired. By the induction hypothesis applied on  $G'$ ,  $\mathbf{cover}_H(G') \leq \omega_H(r) \cdot (k - 1)$  edges. Therefore, from Observation 2,  $\mathbf{cover}_H(G) \leq |D| + \mathbf{cover}_H(G') \leq |D| + \omega_H(r) \cdot (k - 1) \leq \omega_H(r) \cdot k$  edges as required.  $\square$

We set  $\sigma : \mathbb{N} \rightarrow \mathbb{N}$  where  $\sigma(r) = \lceil \frac{1}{8}(3(r+1)^4 + 2(r+1)^2) \rceil$ .

**Theorem 4.** *Let  $H$  be a connected graph with at least one edge,  $r \in \mathbb{N}$ , and  $G$  be a graph where  $\mathbf{tcw}(G) \leq r$ . Then  $\mathbf{cover}_H(G) \leq \sigma(r) \cdot (4 \cdot |V(H)| + |E(H)|) \cdot \mathbf{pack}_H(G)$ .*

**Proof.** Clearly, we can assume that  $G$  is connected, otherwise we work on each of its connected components separately. By Theorem 2, there is a graph  $G'$  where  $\mathbf{tpw}(G') \leq (r+1)^2/2$ ,  $\mathbf{pack}_{H^+}(G') \leq \mathbf{pack}_H(G)$  and  $\mathbf{cover}_H(G) \leq \mathbf{cover}_{H^+}(G')$ . The result follows as, from Theorem 3,  $\mathbf{cover}_{H^+}(G') \leq \omega_{H^+}((r+1)^2/2) \cdot \mathbf{pack}_{H^+}(G')$  and  $\omega_{H^+}((r+1)^2/2) = \sigma(r) \cdot |E(H^+)| \leq \sigma(r) \cdot (4 \cdot |V(H)| + |E(H)|)$ .  $\square$

### 5. Erdős-Pósa for immersions of sub-cubic planar graphs

#### Grids and walls

Let  $k$  and  $r$  be positive integers where  $k, r \geq 2$ . The  $(k \times r)$ -grid  $\Gamma_{k,r}$  is the graph with vertex set  $\{1, \dots, k\} \times \{1, \dots, r\}$  and edge set  $\{(i, j), (i', j')\}, |i - i'| + |j - j'| = 1$ . We denote by  $\Gamma_k$  the  $(k \times k)$ -grid. The  $k$ -wall  $W_k$  (also called wall of height  $k$ ) is the graph obtained from a  $((k+1) \times (2k+2))$ -grid with vertices  $(x, y), x \in \{1, \dots, k+1\}, y \in \{1, \dots, 2k+2\}$ , after the removal of the “vertical” edges  $\{(x, y), (x+1, y)\}$  for odd  $x+y$ , and then the removal of all vertices of degree 1.




Fig. 1. The graph  $\widehat{W}_5$ .

Let  $W_k$  be a wall. We denote by  $P_j^{(v)}$  the shortest path connecting vertices  $(1, 2j)$  and  $(k + 1, 2j)$ ,  $j \in \{1, \dots, k\}$  and call these paths the *vertical paths* of  $W_k$ , with the assumption that  $P_j^{(v)}$  contains only vertices  $(x, y)$  with  $y \in \{2j, 2j - 1\}$ . Note that these paths are vertex-disjoint. Similarly, for every  $i \in \{1, \dots, k + 1\}$  we denote by  $P_i^{(h)}$  the shortest path connecting vertices  $(i, 1)$  and  $(i, 2k + 2)$  (or  $(i, 2k + 1)$  if  $(i, 2k + 2)$  has been removed) and call these paths the *horizontal paths* of  $W_k$ . Let  $E = \{e \mid e \in E(P_j^{(v)}) \cap E(P_i^{(h)}), j \in \{1, 2, \dots, k\}, i \in \{1, 2, \dots, k + 1\}\}$ . We obtain  $\widehat{W}_k$  from  $W_k$  by adding a second copy of every edge in  $E$  (cf. Fig. 1).

*Strong immersions*

If we additionally require in the definition of the immersion containment that no branch vertex is an internal vertex of any certifying path, then the function  $(\phi, \psi)$  is an *H-strong-immersion model*. We then say that  $G$  contains  $H$  as a *strong immersion*, what we denote by  $H \leq_{\text{sim}} G$ .

*Topological minors*

If we additionally require in the definition of the immersion containment that certifying paths are pairwise internally disjoint, then  $(\phi, \psi)$  is an *H-topological model*. We then say that  $G$  contains  $H$  as a *topological minor*, what we denote by  $H \leq_{\text{tm}} G$ .

Observe that in a graph, every topological minor is a strong immersion of  $G$ , and every strong immersion is an immersion. The expansion of a strong immersion or topological model is defined as the one of an immersion model.

The next observation is a formal statement of what is depicted in Fig. 2:  $\widehat{W}_n$  contains  $\Gamma_n$  as a strong immersion. Branch vertices are depicted by white nodes and horizontal (respectively vertical) paths use the color green (respectively red).


**Observation 3.** Let  $k \geq 2$  be an integer. If we define  $\phi$  and  $\psi$  with domains  $V(\Gamma_n)$  and  $E(\Gamma_n)$ , respectively, as follows:

$$\begin{aligned} \phi((i, j)) &= (i, 2j - 1) \\ \psi(\{(i, j), (i, j + 1)\}) &= (i, 2j - 1)(i, 2j)(i, 2j + 1) - 1 \\ \psi(\{(i, j), (i + 1, j)\}) &= (i, 2j - 1)(i + 1, 2j - 1) \text{ for odd } i \\ \psi(\{(i, j), (i + 1, j)\}) &= (i, 2j - 1)(i, 2j)(i + 1, 2j)(i + 1, 2j - 1) \text{ for even } i, \end{aligned}$$

then  $(\phi, \psi)$  is a  $\Gamma_k$ -strong-immersion model in  $\widehat{W}_k$  (where we assume that  $\Gamma_k$  has vertex set  $\{1, \dots, k\}^2$ ).

We also need the following result.

**Lemma 5 ([17]).** Every simple planar sub-cubic graph of  $n$  vertices is a topological minor of the  $\lfloor \frac{n}{2} \rfloor$ -grid.



**Fig. 2.** Finding  $\Gamma_5$  as a strong immersion in  $\widehat{W}_5$ . (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

The next result is mentioned in [27] but its proof is not provided.

**Lemma 6.** Every planar sub-cubic graph on  $n$ -vertices is a topological minor, and hence also a strong immersion, of the wall  $W_n$ .

**Proof.** Let  $H$  be a graph on  $n$  vertices. The proof goes as follows: we first construct a topological expansion  $H'$  of  $H$  that is a simple graph. Then we prove that  $H'$  is a strong-immersion of  $\widehat{W}_n$  and obtain the following ordering:

$$H \leq_{\text{tm}} H' \leq_{\text{tm}} \Gamma_n \leq_{\text{sim}} \widehat{W}_n. \tag{4}$$

Finally, we construct a topological model of  $H'$  in  $\widehat{W}_n$ . The expansion of this model is simple, hence it will be a subgraph of  $W_n$ , as required.

Let  $H$  be a planar sub-cubic graph and let  $H'$  be the simple sub-cubic planar graph obtained from  $H$  by subdividing all but one edges of every multi-edge. Notice that the first inequality of Eq. (4) is satisfied. Let us count how many vertices are added during the construction of  $H'$ . As  $H$  is sub-cubic, among the edges incident to a given vertex, at most two are being subdivided. That way we count each subdivided edge twice (once for each of its endpoints), hence we get:


$$|V(H')| \leq 2|V(H)|.$$

According to Lemma 5,  $H'$  is a topological minor of  $\Gamma_n$ ; this gives the second inequality of the equation. Observation 3 gives the third inequality.

Let  $(\phi_1, \psi_1)$  be an  $H'$ -topological model in  $\Gamma_n$  and let  $(\phi_2, \psi_2)$  be the  $\Gamma_n$ -strong-immersion model in  $\widehat{W}_n$  given by Observation 3. These two models can be used to construct an  $H'$ -strong immersion model  $(\phi, \psi)$  in  $\widehat{W}_n$ , as the composition of  $(\phi_1, \psi_1)$  and  $(\phi_2, \psi_2)$ : for every  $v \in V(H')$ ,  $\phi(v) = \phi_2(\phi_1(v))$  and for every  $e \in E(H')$ ,  $\psi(e)$  is the concatenation of the paths obtained by applying  $\psi_2$  to the edges of the path  $\psi_1(e)$  (taken in the same order as they appear in this path). Observe that this model satisfies the following properties:

- the expansion of  $(\phi, \psi)$  is a subgraph of the expansion of  $(\phi_2, \psi_2)$ ; and
- the branch vertices of  $(\phi, \psi)$  are branch vertices of  $(\phi_2, \psi_2)$ .

We provide the following diagram to recall the roles of the different models we use (topological models are indicated by double arrows and strong immersion models by simple ones).






Fig. 3. Swapping branch vertices.

Let us show the following claim.

**Claim 1.** *Let  $e, f \in E(H')$ . If  $v$  is an internal vertex of both  $\psi(e)$  and  $\psi(f)$ , then these paths also share an endpoint, which is adjacent to  $v$ .*

If  $\psi(e)$  and  $\psi(f)$  share an internal vertex  $v$ , there are two edges  $a \in \psi_1(e)$  and  $b \in \psi_1(f)$  such that both  $\psi_2(a)$  and  $\psi_2(b)$  contain  $v$ . By definition of  $(\phi_2, \psi_2)$ , such a situation occurs only if  $a = \{(i, j), (i + 1, j)\}$  (for even  $i$ ) and  $b = \{(i, j), (i, j + 1)\}$  or  $b = \{(i + 1, j), (i + 1, j + 1)\}$ , for some even  $i \in \{1, \dots, n\}$  and some  $j \in \{1, \dots, n\}$  (see Fig. 2). Observe that in both cases  $a$  and  $b$  share an endpoint. As  $(\phi_1, \psi_1)$  is a topological minor model,  $\psi_1(e)$  and  $\psi_1(f)$  may meet on endpoints only. Therefore the common endpoint of  $a$  and  $b$  is an endpoint of both  $\psi_1(e)$  and  $\psi_1(f)$ , hence  $\psi(e)$  and  $\psi(f)$  have a common endpoint. This proves the first part of the claim. The second part is now clear from the definition of  $(\phi_2, \psi_2)$ , as we know that the paths  $\psi_1(e)$  and  $\psi_1(f)$  start from the same vertex, one with a “vertical” edge, the other with a “horizontal” edge (see Fig. 2).  $\diamond$

If  $(\phi, \psi)$ , which is a strong immersion model, is a topological model, then we can directly jump to the next step. Otherwise, according to Claim 1, there are two edges  $e = \{u, v\}, f = \{u, w\}$  of  $H$  and vertices  $x, y \in \widehat{W}_n$  such that  $\psi(e)$  and  $\psi(f)$  both start with  $x = \phi(u)$  followed by  $y$ . Hence  $\{x, y\}$  is a double edge of  $\widehat{W}_n$ . As  $(\phi, \psi)$  is a strong immersion model of a sub-cubic graph,  $x$  has degree at most three in the expansion of  $(\phi, \psi)$ . We can therefore modify  $(\phi, \psi)$  as follows: we set  $\phi(u) = x$  and we shorten  $\psi(\{u, v\})$  and  $\psi(\{u, w\})$  by removing the edge  $\{x, y\}$  from each of them. In the case where there is a third vertex  $t \in V(H) \setminus \{v, w\}$  adjacent to  $u$ , we also extend the path  $\psi(\{t, u\})$  by adding the edge  $\{x, y\}$ . See Fig. 3 for an example.

It is easy to see that by applying these changes we still get an  $H'$ -strong immersion model, with less crossings of certifying paths. By repeatedly applying these steps we eventually obtain an  $H'$ -topological model in  $\widehat{W}_n^+$ . Notice that its expansion is a simple graph, as  $H'$  is a simple graph. Therefore this expansion is also a subgraph of  $W_n$ . We proved that  $H'$  is a topological minor of  $W_n$ . It follows that the same holds for  $H$  and we are done.  $\square$

By combining [28] with the main result of [6] (see also [7]) we can readily obtain the following.

**Theorem 5.** *There is a function  $f : \mathbb{N}^+ \rightarrow \mathbb{N}$  such that the following holds: for every graph  $G$  and  $r \in \mathbb{N}^+$ , if  $\text{tcw}(G) \geq f(r)$  then  $W_r$  is an immersion of  $G$ . Moreover,  $f(r) = O(r^{29} \text{polylog}(r))$ .*

**Lemma 7.** *Let  $G$  be a graph and let  $H$  be an  $h$ -vertex graph that is connected, planar, and sub-cubic. Then  $\text{tcw}(G) = O(h^{29} \cdot (\text{pack}_{\mathcal{I}(H)}(G))^{14.5} \cdot (\text{polylog}(h) + \text{polylog}(\text{pack}_{\mathcal{I}(H)}(G))))$ .*

**Proof.** Let  $\text{pack}_H(G) \leq k$ . Let  $g(h, k) = f((h + 1) \cdot \lceil (k + 1)^{1/2} \rceil)$ , where  $f$  is the function of Theorem 5. Suppose that  $\text{tcw}(G) \geq g(h, k)$ . Then, from Theorem 5, we obtain that  $G$  contains the wall  $W$  of height  $(h + 1) \cdot \lceil (k + 1)^{1/2} \rceil$  as an immersion. Notice that  $W$  contains  $k + 1$  vertex-disjoint walls of height  $h$ . From Lemma 6, each one of these walls contains  $H$  as an immersion and thus an  $H$ -immersion expansion. Since, these walls are vertex-disjoint they are also edge-disjoint. Hence, we have found a packing of  $H$  of size  $k + 1 > k$ , a contradiction. Therefore,  $\text{tcw}(G) \leq g(h, k)$ . Notice now that, from Theorem 5,  $g(h, k) = O(h^{29} k^{14.5} (\text{polylog}(h) + \text{polylog}(k)))$  as required.  $\square$

The edge version of Theorem 1 follows as a corollary of Theorem 4 and Lemma 7.

## 6. The vertex case

Proving the vertex version of [Theorem 1](#) is a much easier task. For this, we follow the same methodology but use the graph parameter of treewidth instead of tree-cut width, and topological minors instead of immersions.

### Treewidth

We call a graph  $H$   $k$ -chordal if it does not contain any induced cycle of length at least 4 and every clique has at most  $k+1$  vertices. The treewidth of a graph  $G$  is the minimum  $k$  for which  $G$  is a subgraph of a  $k$ -chordal graph.

For the proof of the vertex case of [Theorem 1](#), we require the following two “vertex counterparts” of [Theorem 4](#) and [Lemma 7](#) respectively.

**Proposition 2.** *Let  $\mathcal{H}$  be a class of connected graphs and let  $t$  be a non-negative integer. Then  $\mathcal{H}$  has the  $v$ -E&P property for the graphs of treewidth at most  $t$  with a gap that is a polynomial function on  $t$ .*

**Lemma 8.** *Let  $G$  be a graph and let  $H$  be a connected planar graph on  $h$  vertices, without any  $\mathcal{I}(H)$ -vertex packing of size greater than  $k$ . Then  $\mathbf{tw}(G) = (h \cdot k)^{O(1)}$ .*

[Proposition 2](#) was proven by Thomassen in [\[27\]](#) (see also [\[4, 12\]](#)). For [Lemma 8](#), we need the fact that there is a polynomial function  $\lambda : \mathbb{N}^+ \rightarrow \mathbb{N}$  such that for every  $r \in \mathbb{N}^+$ , every graph with treewidth at least  $\lambda(r)$  contains  $W_r$  as a topological minor. The existence of such a function  $\lambda$  follows from the *grid exclusion theorem* of Robertson and Seymour in [\[24\]](#) (see also [\[8, 25\]](#)) and the polynomiality of  $\lambda$  was proved recently by Chekuri and Chuzhoy in [\[5\]](#) (see also [\[6, 7\]](#) for improvements). Then [Lemma 8](#) can be proved using the same arguments as in [Lemma 7](#), taking into account [Lemma 6](#).

The vertex version of [Theorem 1](#) follows from [Proposition 2](#) and [Lemma 8](#) if, in [Proposition 2](#), we set  $\mathcal{H} = \mathcal{I}(H)$  and  $t = (h \cdot k)^{O(1)}$ .

## 7. Discussion

Notice that in [Theorem 1](#) we demand that  $H$  is a connected graph. It is easy to extend this result if instead of  $H$  we consider some finite collection  $\mathcal{H}$  of connected graphs, at least one of which is planar sub-cubic, and where we define  $\mathcal{I}(\mathcal{H})$  as the class of all graphs containing some graph in  $\mathcal{H}$  as an immersion. Moreover, it is possible to drop the connectivity condition for the vertex variant using arguments from [\[24\]](#), to the price of a slight increase of the gap. However it remains open whether this can be done for the edge variant as well.

Naturally, the most challenging problem on the Erdős–Pósa properties of immersions is to characterize the graph classes:

$$\mathcal{H}^{v/e} = \{H \mid \mathcal{I}(H) \text{ has the } v/e - \text{E\&P property}\}.$$

In this paper we prove that both  $\mathcal{H}^v$  and  $\mathcal{H}^e$  contain all planar sub-cubic graphs. It is an interesting question whether  $\mathcal{H}^{v/e}$  are wider than this. Using arguments similar to [\[22, 24\]](#) it is possible to prove the following.

**Lemma 9.** *No graph of  $\mathcal{H}^v$  and  $\mathcal{H}^e$  is sub-cubic and not planar.*

Actually, the arguments of [\[22, 24\]](#) permit to exclude all non-planar graphs from  $\mathcal{H}^v$ . For the non-sub-cubic case, we can first observe that  $K_{1,4}$ , which is planar and non-sub-cubic belongs to both  $\mathcal{H}^v$  and  $\mathcal{H}^e$ . However, this is not the case for all planar and non-sub-cubic graphs as is indicated in the following observation.

**Observation 4.** *There is a graph  $H$  that is 3-connected, non-sub-cubic, planar, and does belong neither to  $\mathcal{H}^v$  nor to  $\mathcal{H}^e$ .*

**Proof.** Thomassen in [\[27\]](#) provided an example of a tree that does not belong in  $\mathcal{H}^v$  (the same graph does not belong in  $\mathcal{H}^e$  either). Inspired by the construction of [\[27\]](#), we consider first the graph  $H$




Fig. 4. A biconnected graph  $H$  for which  $\mathcal{I}(H)$  does not have the v/e-E&P property.




Fig. 5. The host graph  $G$ .

depicted in Fig. 4. To see that  $H \notin \mathcal{H}^v$  and  $H \notin \mathcal{H}^e$ , consider as host graph  $G$  the graph in Fig. 5. This graph consists of a main body that is a wall of height 3 and three triples of graphs attached at its upper, leftmost, and lower paths. Each of these triples consists of three copies of some of the 3-connected components of  $H$ . Notice that  $G$  does not contain more than one  $H$ -immersion expansion. However, in order to cover all  $H$ -immersion expansions of  $G$  one needs to remove at least 3 edges/vertices. By increasing the height of the wall of  $G$ , we may increase the minimum size of an  $\mathcal{I}(H)$ -vertex/edge cover while no  $\mathcal{I}(H)$ -vertex/edge packing of size greater than 1 will appear. It is easy to modify  $H$  so to make it 3-connected: just add a new vertex and make it adjacent with the tree vertices of degree 4. The resulting graph  $H'$  remains planar. The same arguments, applied to an easy modification of the host graph, can prove that  $H'$  is not a graph in  $\mathcal{H}^v$  or  $\mathcal{H}^e$ .  $\square$

Providing an exact characterization of  $\mathcal{H}^v$  and  $\mathcal{H}^e$  is an insisting open problem. A first step to deal with this problem could be the cases of  $\theta_4 =$   and the 4-wheel = . Especially for the 4-wheel, the structural results in [1] might be useful.

## References

- [1] Rémy Belmonte, Archontia Giannopoulou, Daniel Lokshtanov, Dimitrios M. Thilikos, The Structure of  $W_4$ -Immersion-Free Graphs. *ArXiv E-Prints*, February 2016.
- [2] Etienne Birmelé, Joh Adrian Bondy, Bruce A. Reed, The Erdős-Pósa property for long circuits, *Combinatorica* 27 (2) (2007) 135–145.
- [3] D. Chatzidimitriou, J.-F. Raymond, I. Sau, D.M. Thilikos, Minors in graphs of large  $\theta_r$ -girth, *European J. Combin.* 65 (2017) 106–121.
- [4] Chandra Chekuri, Julia Chuzhoy, Large-treewidth graph decompositions and applications, in: 45st Annual ACM Symposium on Theory of Computing, STOC, 2013, pp. 291–300.
- [5] Chandra Chekuri, Julia Chuzhoy, Polynomial bounds for the grid-minor theorem, *CoRR*, abs/13056577, 2013.
- [6] Julia Chuzhoy, Improved bounds for the excluded grid theorem, *CoRR*, abs/160202629, 2015.
- [7] Julia Chuzhoy, Excluded grid theorem: Improved and simplified, in: Proceedings of the Forty-Seventh Annual ACM on Symposium on Theory of Computing, STOC 2015, Portland, OR, USA, June 14–17, 2015, pp. 645–654.
- [8] Reinhard Diestel, Graph Theory, third ed., in: Graduate Texts in Mathematics, vol. 173, Springer-Verlag, Heidelberg, 2005.
- [9] Reinhard Diestel, Ken-Ichi Kawarabayashi, Paul Wollan, The Erdős-Pósa property for clique minors in highly connected graphs, *J. Combin. Theory Ser. B* 102 (2) (2012) 454–469.
- [10] Guoli Ding, Bogdan Oporowski, On tree-partitions of graphs, *Discrete Math.* 149 (1–3) (1996) 45–58.
- [11] Paul Erdős, Louis Pósa, On independent circuits contained in a graph, *Canad. J. Math.* 17 (1965) 347–352.
- [12] Fedor V. Fomin, Saket Saurabh, Dimitrios M. Thilikos, Strengthening erdős-Pósa property for minor-closed graph classes, *J. Graph Theory* 66 (3) (2011) 235–240.
- [13] Robert Ganian, Eun Jung Kim, Stefan Szeider, Algorithmic applications of tree-cut width, in: Giuseppe F. Italiano, Giovanni Pighizzini, Donald T. Sannella (Eds.), Mathematical Foundations of Computer Science 2015, in: Lecture Notes in Computer Science, vol. 9235, Springer, Berlin, Heidelberg, 2015, pp. 8–360.
- [14] Jim Geelen, Kasper Kabell, The Erdős-Pósa property for matroid circuits, *J. Combin. Theory Ser. B* 99 (2) (2009) 407–419.
- [15] Rudolf Halin, Tree-partitions of infinite graphs, *Discrete Math.* 97 (1–3) (1991) 203–217.
- [16] Naonori Kakimura, Ken-ichi Kawarabayashi, Fixed-parameter tractability for subset feedback set problems with parity constraints, *Theoret. Comput. Sci.* 576 (2015) 61–76.
- [17] Goos Kant, Drawing planar graphs using the canonical ordering, *Algorithmica* 16 (1) (1996) 4–32.
- [18] Ken-Ichi Kawarabayashi, Atsuhiko Nakamoto, The Erdős-Pósa property for vertex- and edge-disjoint odd cycles in graphs on orientable surfaces, *Discrete Math.* 307 (6) (2007) 764–768.
- [19] Daniel Král', Heinz-Jürgen Voss, Edge-disjoint odd cycles in planar graphs, *J. Combin. Theory Ser. B* 90 (1) (2004) 107–120.
- [20] Chun-Hung Liu, Packing and covering immersions in 4-edge-connected graphs. *CoRR*, abs/150500867, 2015.
- [21] Dieter Rautenbach, Bruce A. Reed, The Erdős-Pósa property for odd cycles in highly connected graphs, *Combinatorica* 21 (2) (2001) 267–278.
- [22] Jean-Florent Raymond, Ignasi Sau, Dimitrios M. Thilikos, An edge variant of the Erdős-Pósa property, *Discrete Appl. Math.* (2017) in press.
- [23] Bruce A. Reed, Neil Robertson, Paul D. Seymour, Robin Thomas, Packing directed circuits, *Combinatorica* 16 (4) (1996) 535–554.
- [24] Neil Robertson, Paul D. Seymour, Graph minors. V. Excluding a planar graph, *J. Combin. Theory Ser. B* 41 (2) (1986) 92–114.
- [25] Neil Robertson, Paul D. Seymour, Robin Thomas, Quickly excluding a planar graph, *J. Combin. Theory Ser. B* 62 (2) (1994) 323–348.
- [26] Detlef Seese, Tree-partite graphs and the complexity of algorithms, in: Proceedings of Fundamentals of Computation Theory, in: Lecture Notes in Computer Science, vol. 199, Springer, Berlin, Heidelberg, 1985, pp. 412–421.
- [27] Carsten Thomassen, On the presence of disjoint subgraphs of a specified type, *J. Graph Theory* 12 (1) (1988) 101–111.
- [28] Paul Wollan, The structure of graphs not admitting a fixed immersion, *J. Combin. Theory Ser. B* 110 (2015) 47–66.