Corners and simpliciality in oriented matroids and partial cubes

Kolja Knauer¹

Tilen Marc²

March 14, 2023

Abstract

Building on a recent characterization of tope graphs of Complexes of Oriented Matroids (COMs), we tackle and generalize several classical problems in Oriented Matroids (OMs), Lopsided Sets (aka ample set systems), and partial cubes via Metric Graph Theory.

Our first main result is that every element of an OM from a class introduced by Mandel is incident to a simplicial tope, i.e, such OMs contain no mutation-free elements. This allows us to refute a conjecture of Mandel from 1983, that would have implied the famous Las Vergnas' simplex conjecture. Further, we show that the mutation graph of uniform OMs of order at most 9 are connected, thus confirming a stronger conjecture of Cordovil-Las Vergnas in this setting.

The second main contribution is the introduction of corners of COMs as a natural generalization of corners in Lopsided Sets. Generalizing results of Bandelt and Chepoi, Tracy Hall, and Chalopin et al. we prove that realizable COMs, rank 2 COMs, as well as hypercellular graphs admit corner peelings. Using this, we confirm Las Vergnas' conjecture for antipodal partial cubes (a class much lager than OMs) of small rank or isometric dimension.

1 Introduction

The hypercube Q_n of dimension n is the graph whose vertex set is $\{+,-\}^n$ where two vertices are adjacent if they differ in exactly one coordinate. A graph G is called a partial cube if G is an isometric subgraph of a hypercube Q_n , i.e., $d_G(u,v)=d_{Q_n}(u,v)$ for all $u,v\in G$. In this case, the minimum such n is the isometric dimension of G.

An Oriented Matroid (OM) is a mathematical structure that abstracts the properties of hyperplane arrangements, directed graphs, vector arrangements and others. It has been studied extensively in combinatorial, as well as in topological and algebraic settings. For graph theorists it might be attractive to consider the *tope graph* of an OM. It can easily be shown that the tope graph is a partial cube and determines a simple OM uniquely up to isomorphism [6], hence no information is lost in this approach.

In this paper based on recent graph theoretical characterization [31], we analyse OMs and related structures through their tope graphs. In particular, in the first part of the paper this view leads us to new results concerning Las Vergnas simplex conjecture and mutations. While the approach to this section is graph theoretical, we try to present it in a way that a reader more familiar with the standard approach to OMs can understand it. In the second part of the paper we explore how these problems and ideas can be extended beyond OMs into two directions. First, we consider Complexes of Oriented Matroids (COMs), which are a recent generalization of OMs, who share the property that their tope graph is a partial cube and determines a simple COM uniquely up to isomorphism [4]. Second, we consider *antipodal* partial cubes, i.e., those isometric subgraphs G of Q_n such that for every $v \in G$ also its *antipode* -v, where all

¹Laboratoire d'Informatique et Systhèmes, Aix-Marseille Université, Marseille, France & Departament de Matemàtiques i Informàtica, Universitat de Barcelona (UB), Barcelona, Spain, kolja.knauer@lis-lab.fr

²Faculty of Mathematics and Physics, Ljubljana, Slovenia, Institute of Mathematics, Physics and Mechanics, Ljubljana, Slovenia & XLAB d.o.o., Ljubljana, Slovenia, tilen.marc@fmf.uni-lj.si

signs are reversed, is a vertex of G. Note that if G is isometrically embedded in Q_n with minimal n, the embedding is unique up to an isomorphism [42, Chapter 5], so an antipode is well defined even if the embedding is not given. Antipodal partial cubes are the tope graphs of so-called acycloids [20]. Both of these generalizations are natural since G is the tope graph of an OM if and only it is antipodal and the tope graph of a COM [31].

So, what is a tope graph? One of the standard ways to define an OM, a COM (or an acycloid) is as a pair $\mathcal{M} = (E, \mathcal{L})$ of a finite *ground set E* and a set of *covectors* $\mathcal{L} \subset \{+, -, 0\}^E$ with certain properties, see Section 2. One obtains the *tope graph*, by considering the subgraph induced in the hypercube by the *topes*, i.e., covectors without 0-entries. Considering an OM as its tope graph, we will denote it just as G. All OM notions have an equivalent description in terms of the tope graph, e.g., the order of E is the isometric dimension of G. The *rank* of an OM $\mathcal{M} = (E, \mathcal{L})$, classically defined as the length of the maximal chains in \mathcal{L} with respect to the product order induced by 0 < +, 0 < -, can be read of its tope graph G as well, see Section 2.

If G is an OM, then a vertex $v \in G$ is called *simplicial* if deg(v) = r(G). Simplicial vertices correspond to simplicial topes and it is a well known fact that the degree of each vertex (tope) must be at least r(G). The well-known simplex conjecture of Las Vergnas [34] in terms of tope graphs of OMs reads as follows:

Conjecture 1 (Las Vergnas). Every OM has a simplicial tope.

The conjecture is motivated by the fact that it holds for all OMs that are *realizable* (by hyperplanes in Euclidean space) [52]. The largest class known to satisfy Conjecture 1 was found in [37, Theorem 7]. We call that class *Mandel* here and consider it in depth in Section 4. Realizable OMs and OMs of rank at most 3 are *Euclidean* and the latter are Mandel, but the class is larger. Indeed, Mandel [37, Conjecture 8] even conjectured the following as a "wishful thinking statement", since by the above it would imply the conjecture of Las Vergnas.

Conjecture 2 (Mandel). Every OM is $Mandel^1$.

Let us now consider some strengthenings of Las Vergnas' conjecture. Recall that G is a partial cube, and hence its edges are partitioned into so-called Θ -classes corresponding to E, i.e., the coordinates of the hypercube G is embedded in. We say that such G is Θ -Las Vergnas, if every Θ -class of G contains an edge incident to a simplicial vertex. In the language of OMs this means that G has no mutation-free elements. It is known that rank 3 OMs are Θ -Las Vergnas [36]. This brings us to the first main result of this paper. In Theorem 4.5 we extend the class of Θ -Las Vergnas OMs significantly, by showing that Mandel OMs are Θ -Las Vergnas. On the other hand, OMs that are not Θ -Las Vergnas have been discovered [8,47,53]. See Figure 1 for an illustration of one of them. Together with Theorem 4.5 this disproves Mandel's conjecture (Corollary 4.6).

Result 1. Mandel OMs are Θ -Las Vergnas, hence Mandel's Conjecture fails.

The next is a strengthening of Las Vergnas' conjecture for uniform Oriented Matroids (UOMs), see Section 2 for a definition. A well-known operation that one can apply to a simplicial vertex of an UOM is a *mutation* - transforming one UOM to another one, see Section 3. One can consider the *mutation graph* $\mathcal{G}^{n,r}$, whose vertices are (reorientation classes of) UOMs of rank r and of order n. Two reorientation classes are connected if and only if there exists a mutation between them. The graph $\mathcal{G}^{n,r}$ is motivated by the topological representation of OMs. In particular, by Ringel's Homotopy Theorem [48, 49] it follows that $\mathcal{G}^{n,3}$ is connected. Moreover, the subgraph of $\mathcal{G}^{n,r}$ induced by the realizable UOMs is connected by [50]. Las Vergnas' conjecture would imply that $\mathcal{G}^{n,r}$ has minimum degree at least 1 (where loops can occur). A much stronger affirmation appears in [50]:

Conjecture 3 (Cordovil-Las Vergnas). For all r, n the graph $\mathcal{G}^{n,r}$ is connected.

¹Note that it was us and not Mandel who baptized this class of OMs, we merely want to combine eponymous homage and correct attribution.




Figure 1: A halfspace E_e^+ of a non-Mandel OM G, where E_e is the Θ-class witnessing that G is not Θ-Las Vergnas. The bold subgraph is an OM of rank 3 on the vertices incident with E_e in G.

In Section 3 we introduce two variations of $\mathscr{G}^{n,r}$ that seem natural in the graph theoretic language of OMs, denoted $\overline{\mathscr{G}}^{n,r}$ and $\underline{\mathscr{G}}^{n,r}$. In the graph $\underline{\mathscr{G}}^{n,r}$ the vertices are graph isomorphism classes, instead of the reorientation classes, while in $\overline{\mathscr{G}}^{n,r}$ the vertices are all the OMs of rank r and order n (over the same ground set). Indeed, it is easy to see that the connectivity of $\overline{\mathscr{G}}^{n,r}$ implies the connectivity of $\mathscr{G}^{n,r}$ and the connectivity of $\mathscr{G}^{n,r}$ implies that $\underline{\mathscr{G}}^{n,r}$ is connected. (Observation 3.1).

Our results with respect to Conjecture 3 include that $\overline{\mathscr{G}}^{n,3}$ is connected, which is a consequence of Ringel's Homotopy Theorem [48,49] and strengthens the fact that $\mathscr{G}^{n,3}$ is connected (Proposition 3.2). Moreover, we show that connectivity of $\underline{\mathscr{G}}^{n,r}$ implies connectivity of $\mathscr{G}^{n,r}$ (Proposition 3.3). Since $\underline{\mathscr{G}}^{n,r}$ is much smaller that $\mathscr{G}^{n,r}$, the fact that Conjecture 3 is closed under duality, see [7, Exercise 7.9], allows us to verify Conjecture 3 for all $n \leq 9$, computationally. See Table 1 for orders of the graphs $\underline{\mathscr{G}}^{n,r}$ and Figure 5 for a depiction of $\mathscr{G}^{8,4}$.

Result 2. The Cordovil-Las Vergnas Conjecture holds for UOMs of order at most 9.

In the last part of the paper (Sections 5, 6) we explore how the notion of simpliciality generalizes to classes of partial cubes containing OMs. First we consider Complexes of Oriented Matroids (COMs) [4]. This class can be seen as a common generalization of OMs as well as,e.g., distributive lattices, antimatroids, median graphs, CAT(0) cube complexes, linear extensions of posets, pseudoline arrangements, hyperplane arrangements, Pasch graphs, and Lopsided Sets (LOPs). COMs have been acknowledged already several times in their short existence, see [5, 16, 25, 27, 33, 40, 43, 46]. As their name suggest, COMs may be seen as complexes whose cells are OMs. In terms of the tope graph G of a COM its OM cells can be identified as its so-called antipodal subgraphs. If a COM is a LOP, then these cells are just subgraphs isomorphic to hypercubes, see Section 2 for the details.

We generalize the notion of simpliciality from OMs to COMs as follows: a vertex $v \in G$ is *simplicial* if it is contained in a unique maximal antipodal subgraph $A \subseteq G$ and deg(v) = r(A). In LOPs simplicial vertices are usually called *corners* and have important implications for results in computational

learning theory, see [9]. The notion of corners in COMs allows us to prove the existence of simplicial vertices in certain COMs. In Section 5.2 we show that realizable COMs (Proposition 5.5), COMs of rank 2 (Theorem 5.11), and hypercellular graphs (Theorem 5.14) admit *corner peelings*, which is a way of deconstructing a COM in a well-behaved way. This result generalizes the corresponding results on realizable LOPs [53], LOPs of rank 2 [9], and bipartite cellular graphs [3]. Furthermore, together with the examples from [8, 47, 53] this yields locally realizable COMs that are not realizable and refutes a conjecture of [4, Conjecture 2] (Remark 5.6).

Lastly, we consider the question of generalizing Conjecture 1 to antipodal partial cubes. First we consider the case where the rank is at most 3. In this case the distinction between general antipodal partial cubes and OMs is easy, since OMs of rank at most 3 are exactly the planar antipodal partial cubes [20]. Based on this, for rank 3 OMs Conjecture 1 can be deduced from Euler's Formula and the fact that partial cubes are triangle-free. We extend this result by showing that all (hence also the non-planar) antipodal partial cubes of rank at most 3 have simplicial vertices in Section 6.

Conjecture 1 has been verified for UOMs of rank 4 up to 12 elements [8]. In Section 6 we prove that even general antipodal partial cubes of isometric dimension up to 7 have simplicial vertices. It could be that every antipodal partial cube of rank r has minimum degree at most r.

Result 3. Realizable COMs, COMs of rank 2 and hypercellular graphs have corner peelings, thus in particular simplicial vertices. The minimum degree of antipodal partial cubes of rank 3 or of isometric dimension at most 7 is at most their rank, i.e., they satisfy the Las-Vergnas property.

2 Preliminaries

Based on a recent result from [31], we start by presenting OMs, Affine Oriented Matroids (AOMs) and Uniform Oriented Matroids (UOMs) solely as graphs, in particular as a subfamily of partial cubes. A reader familiar with these concepts can avoid this view and read the paper nevertheless. We begin with necessary standard definitions from Metric Graph Theory:

As stated above, a partial cube is an isometric subgraph of a hypercube, where the minimal dimension it embeds into is called its *isometric dimension*. The notion is well defined since an embedding of a partial cube is unique up to automorphisms of the hypercube [42, Chapter 5]. As mentioned above, we call a partial cube G of isometric dimension n antipodal if when embedded in Q_n for every vertex v of G also its antipode -v, i.e., the vertex with all coordinates flipped, is in G. A subgraph H of G is convex, if all the shortest paths in G connecting two vertices of H are also in H. A convex subgraph H of a partial cubes G is a partial cube as well, since every convex (or isometric) subgraph of an isometric subgraph of a hypercube is isometric in the hypercube as well. In fact the embedding of H is inherited from G. We say a subgraph H of G is antipodal, if it is a convex subgraph of G and an antipodal partial cube on its own (where antipodes in H are considered with respect to its embedding in a minimal hypercube). Finally, we call a subgraph H of a graph G gated if for every vertex x of G there exists a vertex v_x of H such that for every $u \in H$ there is a shortest path in G from x to u passing through v_x . Gated subgraphs are convex [23, Section 12.1]. As mentioned earlier, embedding a partial cube G in Q_n yields a partition of the edges of G into so-called Θ -classes corresponding to the dimensions of Q_n . Each Θ -class consists of the edges corresponding to a flip of a fixed coordinate. We denote by E_f a Θ -class of G where fcorresponds to a coordinate in $\{1,\ldots,n\}$. We write E_f^+ for the vertices having the coordinate f equal to + and analogously define E_f^- . The sets E_f^+ and E_f^- are called *halfspaces* of G and induce graphs that are partial cubes as well since they are already isometrically embedded in a hypercube. If $\{E_{f_1}^{s_1}, E_{f_2}^{s_2}, \dots E_{f_k}^{s_k}\}$ are halfspaces in a partial cube G, for f_i coordinate and $s_i \in \{+, -\}$, $1 \le i \le k$, the partial cube induced by the intersection $\bigcap_{i=1}^k E_{f_i}^{s_i}$ is sometimes called a *restriction* of G. Note that in the standard language of oriented matroids, Θ -classes correspond to *elements* of a simple OM, hence also isometric dimension is sometimes just called the number of elements.

Starting from a graph theoretical perspective we state a characterization – that serves as a definition for now – of (simple) OMs, AOMs, UOMs, LOPs, and COMs.

Theorem 2.1 ([13,15,31]). There is a one to one correspondence between the classes of (simple):

- (i) OMs and antipodal partial cubes whose antipodal subgraphs are gated [15, 31],
- (ii) AOMs and halfspaces of antipodal partial cubes whose antipodal subgraphs are gated [31],
- (iii) UOMs and antipodal partial cubes whose proper antipodal subgraphs are hypercubes [13, 31].
- (iv) LOPs and partial cubes whose antipodal subgraphs are hypercubes [35], hypercubes [13, 31].
- (v) COMs and partial cubes whose antipodal subgraphs are gated [31],

While the proof of the above theorem is rather complicated, the correspondence with the standard definition of the OMs, AOMs, UOMs through covectors $\mathcal{L} \subset \{+,-,0\}$ (see below) is simple. One direction we already described above, namely given \mathcal{L} the vertices of the tope graph are the elements without 0-entries (topes) inducing a subgraph in Q_E . Conversely, one can get the covectors from the tope graph by associating to its antipodal subgraphs sign-vectors, that encode their relative position to the halfspaces. This is explained in more detail below.

An operation that is well known in the study of partial cubes is a (partial cube) contraction π_f that for a coordinate f contracts all the edges in a Θ -class E_f . The family of partial cubes is closed under the operation of contraction as well as are the families of OMs, AOMs (in their graph representation) and the class of antipodal partial cubes, see [31]. The partial cube contraction operation corresponds to deletion in covector systems.

The $rank\ r(G)$ of a partial cube G is the largest r such that G can be transformed to Q_r by a sequence of partial cube contractions. The definition of rank in oriented matroid theory is equivalent, see [13,15]. Furthermore, notice that viewing the vertices of $G \subseteq Q_n$ as a set $\mathscr S$ of subsets of $\{1,\ldots,n\}$, r(G) coincides with the VC-dimension of $\mathscr S$, see [54]. The latter correspondence has led to some recent interest in partial cubes of bounded rank, see [12]. See Figure 2 for an antipodal partial cube of rank 3.




Figure 2: Contracting an antipodal partial cube to Q_3 .

The inverse of a contraction is an expansion. In fact one can look at expansions in partial cubes the following way: if H is a contraction of G, i.e. $H=\pi_e(G)$, then one can consider in H sets $H_1=\pi_e(E_e^+)$ and $H_2=\pi_e(E_e^-)$. Notice that they completely determine the expansion, since G can be seen as a graph on the disjoint union of H_1 and H_2 where edges between them correspond to $H_1\cap H_2$. By expansion of H we refer to the subgraphs H_1, H_2 and sometimes to G. If G and H are OMs, then we say that H_1, H_2 is an OM-expansion, also called single-element extension in OM theory (we define OM-expansion in terms of covectors as well, see below). More generally, if H is an antipodal partial cube, then an expansion H_1, H_2

of H is antipodal if the expanded graph G is again antipodal. It is well-known, see e.g [31,45], that G is antipodal if and only if $H_1 = -H_2$. Clearly, a OM-expansion is antipodal. An OM-expansion H_1, H_2 is in general position if each maximal proper antipodal subgraph is either completely in H_1 or completely in H_2 but not in $H_1 \cap H_2$. We reestablish the definition of the general position in terms of covectors below. Expansions in general position are central to the notion of Mandel OMs as well as to the definition of corners in COMs, i.e., in both Section 4 and Section 5.2. It is well-known, that every OM admits an expansion and indeed even an expansion in general position, see [51, Lemma 1.7] or [7, Proposition 7.2.2].

We introduced COMs and their subclasses as graphs. Nevertheless, for certain results it is convenient to define covectors and cocircuits of COMs which are standard ways to introduce OMs. The covectors are represented as a subset $\mathcal{L} \subseteq \{+,-,0\}^n$ and have to satisfy certain axioms in order to encode a COM, OM, AOM, LOP or UOM. If $X \in \mathcal{L}$ and $e \in [n]$ is a coordinate of X, we shall write $X_e \in \{+,-,0\}$ for the value of X in coordinate e. When considering tope graphs, one restricts usually to simple systems. Here, a system of sign-vectors \mathcal{L} is simple if it has no "redundant" elements, i.e., for each $e \in [n]$, $\{X_e : X \in \mathcal{L}\} = \{+,-,0\}$ and for each pair $e \neq f$ in [n], there exist $X, Y \in \mathcal{L}$ with $\{X_e X_f, Y_e Y_f\} = \{+,-\}$. We will assume simplicity without explicit mention. By the graph-theoretical representation of COMs given in [31], the covectors correspond to the antipodal subgraphs of a COM G. Indeed, in a partial cube every convex subgraph is an intersection of halfspaces, see e.g. [1], and one can assign to any convex subgraph H a unique sign-vector $X(H) \in \{0,+,-\}^n$ by setting for any coordinate $e \in \{1,\ldots,n\}$:

$$X(H)_e = \begin{cases} + & \text{if } X \subseteq E_e^+, \\ - & \text{if } X \subseteq E_e^-, \\ 0 & \text{otherwise.} \end{cases}$$

This correspondence yields a dictionary in which important concepts on both sides, graphs and sign-vectors translate to each other. We have spoken about isometric dimension and rank. Another instance of this is that if the $v \in G$ is a vertex of an antipodal partial cube, then for its antipode u we have X(u) = -X(v). Thus, we will often denote the antipodes of a set of vertices H just by -H. Another noteworthy instance is the relation of gates and composition. The *composition* of two sign vectors $X, Y \in \{0, +, -\}^n$ is defined as the sign-vector obtained by setting for any coordinate $e \in \{1, ..., n\}$:

$$(X \circ Y)_e = \begin{cases} X_e & \text{if } X_e \neq 0, \\ Y_e & \text{otherwise.} \end{cases}$$

Proposition 2.2. [31] If H is a gated subgraph of a partial cube G, then for a vertex v with gate u in H, we have $X(u) = X(H) \circ X(v)$.

A set of covectors forms a COM if it satisfies the following two axioms:

- (FS) $X \circ -Y \in \mathcal{L}$ for all $X, Y \in \mathcal{L}$.
- (SE) for each pair $X,Y\in\mathcal{L}$ and for each $e\in[n]$ such that $X_eY_e=-1$, there exists $Z\in\mathcal{L}$ such that $Z_e=0$ and $Z_f=(X\circ Y)_f$ for all $f\in[n]$ with $X_fY_f\neq-1$.

It is easy to see, check e.g. [4], that a set of covectors is an OM if and only if it is a COM satisfying additionally:

(Z) $0 \in \mathcal{L}$.

For similar axiomatizations for LOPs, AOMs, and UOMs we refer to [4,5]. We now reestablish some definitions stated in terms of graphs in terms of covectors. What was defined as a partial cube contraction on the tope graph of an OM, corresponds to so called *deletion* in OMs. In particular, if \mathcal{L} is a set of

covectors of an OM and $e \in [n]$ is an element of it, then $\mathcal{L}' = \mathcal{L} \setminus e = \{X \setminus e \mid X \in \mathcal{L}\}$ is a set of covectors of a smaller OM, that is said to the obtain by the deletion of e. On the contrary, of \mathcal{L} is said to be a single-element extension of \mathcal{L}' (recall that in the graph setting it is named an expansion). The extension is said to be in general position if the union of images of $\{X \in \mathcal{L} \mid X_e = +\}$ and $\{X \in \mathcal{L} \mid X_e = -\}$ under the deletion covers all the covectors of \mathcal{L}' . In fact, as we will see in Section 4, the image of $\{X \in \mathcal{L} \mid X_e = +\}$ under deletion is maximal with the property that it induces a COM in \mathcal{L}' . We will use this to define a corner of an OM. For G_1, G_2 being the images of $\{X \in \mathcal{L} \mid X_e = +\}$ and $\{X \in \mathcal{L} \mid X_e = -\}$ under deletion, we will sometimes also say that the expansion is along G_1 and G_2 .

We will content ourselves with the definition of OMs and COMs with covectors and for other classes use Theorem 2.1 as a definition. In order to formulate (SE) in terms of antipodal subgraphs, we give some more terminology about how subgraphs and Θ -classes can relate. We will say that a Θ -class E_f crosses a subgraph H of G if at least one of the edges in H is in E_f . Moreover, E_f separates subgraphs H,H' if $H\subseteq E_f^+$ and $H'\subseteq E_f^-$ or the other way around. We collect the coordinates separating H and H' in the set S(H,H') – usually called the separator. Let us now state the axiom of strong elimination for graphs:

- (SE) Let X, Y be antipodal subgraphs of G and E_e a Θ -class such that $X \subseteq E_e^+$ and $Y \subseteq E_e^-$, i.e., $e \in S(X,Y)$. There is an antipodal subgraph Z of G that is crossed by E_e and for all $f \neq e$ we have:
 - if $X,Y\subseteq E_f^+$ then $Z\subseteq E_f^+$, and if $X,Y\subseteq E_f^-$ then $Z\subseteq E_f^-$
 - if E_f crosses one of X,Y and the other is a subset of E_f^+ then $Z \subseteq E_f^+$, and if E_f crosses one of X,Y the other is a subset of E_f^- then $Z \subseteq E_f^-$
 - if E_f crosses both X and Y, then it crosses Z.

In some parts of the paper we will abuse a bit the distinction between covectors and antipodal subgraphs in the way that we have suggested in the above definitions.




Figure 3: A realizable COM and its tope graph. Its bounded faces, edges and vertices are the antipodal subgraphs.

Some COMs, OMs, AOMs, and LOPs can be particularly nicely represented by a geometric construction. Let $\{H_1,\ldots,H_n\}$ be a set of hyperplanes in an Euclidean space \mathbb{R}^d and C a full-dimensional open convex set in \mathbb{R}^d . The hyperplanes cut C in d-dimensional chambers and to every point in C one can associate a vector in $\{+,-,0\}^n$ that denotes its relative position to the hyperplanes (if they are given with positive and negative side). These are the covectors. One can form a graph G whose vertices are the chambers where two chambers are adjacent if and only if they are separated by exactly one hyperplane. Such graphs are always COMs. If $C = \mathbb{R}^d$ then the graph is an AOM, and if moreover all $\{H_1,\ldots,H_n\}$ cross the origin point, then the graph is an OM. If $\{H_1,\ldots,H_n\}$ are the coordinate hyperplanes of \mathbb{R}^d and C is arbitrary, then the graph is a LOP. If a COM, OM, AOM, or LOP G is isomorphic to a graph obtained

in this way, we call it *realizable*. See Figure 3 for a realizable COM and its tope graph. Realizable COMs embody many nice classes such as linear extension graphs of posets, see [4] and are equivalent to special convex neural codes, namely so-called stable hyperplane codes, see [27,33].

If G is an OM, then the set of antipodal subgraphs is usually ordered by reverse inclusion to yield the big face lattice $\mathscr{F}(G)$ whose minimum is G itself. This ordering in terms of covectors can be equivalently defined by setting $X \leq Y$ for covectors X, Y, as the product ordering of $\{+,-,0\}^n$ relative to the ordering $0 \leq +,0 \leq -$. If G is of rank r, then it follows from basic OM theory that the $\mathscr{F}(G)$ is atomistic and graded, where r+1 is the length of any maximal chain. Hence the latter can also be used to define the rank of G. The atoms, i.e. the covectors corresponding to the antipodal subgraphs of rank r-1, are called *cocircuits* in the standard theory of OMs.

Another graph associated to an OM G of rank r is its *cocircuit graph* of G, i.e., the graph G^* whose vertices are the antipodal subgraphs of G of rank r-1 (equivalently cocircuits) where two vertices are adjacent if their intersection in G is an antipodal subgraph of rank r-2 (i.e. two cocircuits X,Y are adjacent if there exists a covector Z with $X,Y \prec Z$). We denote the cocircuit graph of G by G^* , since it generalizes planar duality in rank 3, see [20], however in higher rank $(G^*)^*$ is not well-defined, because the cocircuit graph does not uniquely determine the tope graph, see [14]. There has been extensive research on cocircuit graphs [2, 17, 32, 41]. However their characterization and recognition remains open. Cocircuit graphs play a crucial role for the notion of Euclideaness and Mandel in Section 4.

In a general COM G, the poset $\mathscr{F}(G)$ remains an upper semilattice, since antipodal subgraphs are closed under intersection, but there is no minimal element. There are different possible notions of cocircuits that allow to axiomatize COMs, see [4]. We consider cocircuits in the setting of *pure* COMs G, i.e., all maximal antipodal subgraphs of G are of the same rank and G^* is connected. If G is a non-antipodal pure COM, then its cocircuits are just the maximal antipodal subgraphs. We will introduce the cocircuit graph of pure COMs in Section 5.2, where it will serve for proving the existence of corners in COMs of rank 2.

3 Mutation graphs of uniform oriented matroids

Let us define mutations of UOMs through their tope graph. If v is a simplicial vertex in a UOM G of rank r, then v is contained in a unique convex hypercube minus a vertex, let us denote it by Q_r^- . This is a well known assertion and it directly follows from Lemma 4.3. If one fills in the missing vertex of Q_r^- and instead removes v and does the same to the antipodes of the vertices in G, one obtains a new UOM G' of rank r. This operation is called a *mutation*. Hence, a mutation is an operation that transforms a UOM into another UOM. A simple analysis shows that the operation is reversible, i.e. the inverse operation is also a mutation, and that the rank of both UOMs is equal.

Thus, one can now consider a *mutation graph* whose vertices are UOMs embedded into Q_n , for some $n \in \mathbb{N}$ of fixed rank r and edges are corresponding to mutations. In fact, one can consider three mutation graphs corresponding to the different notions of equivalence of OMs, that will now be introduced.

As stated above, the isometric embedding of a partial cube into a hypercube of minimum dimension is unique up to automorphisms of the hypercube, see e.g. [42, Chapter 5]. Indeed, partial cubes $G_1, G_2 \subseteq Q_n$ are isomorphic as graphs if and only if there is $f \in \operatorname{Aut}(Q_n)$ such that $f(G_2) = G_1$. This leads to the fact that isomorphisms of simple OMs and their tope graphs correspond to each other. Since (unlabeled, non-embedded) isomorphic tope graphs are sometimes considered as equal, this allows to speak about isomorphism classes of OMs in a natural way. The situation is the same if one consider OMs in the standard definition since an isomorphism of an OM directly translates to an isomorphism of its tope graph, see e.g. [6,7].

A more refined notion of isomorphisms for partial cubes are reorientations. For partial cubes $G_1, G_2 \subseteq Q_n$ one says that G_1 is a *reorientation* of G_2 if there is $f \in \mathbb{Z}_2^n \subseteq \operatorname{Aut}(Q_n)$ such that $f(G_2) = G_1$, i.e., f only switches signs. This yields an equivalence relation whose classes are called *reorientation classes*.

We give the three resulting graphs:

- $\overline{\mathscr{G}}^{n,r}$ is the graph whose vertices are UOMs of rank r and isometric dimension n, embedded into Q_n . Two graphs are connected if and only if there exists a mutation between them.
- $\mathcal{G}^{n,r}$ is the graph whose vertices are reorientation classes of UOMs of rank r and isometric dimension n embedded into Q_n . Two reorientation classes are connected if and only if there exists a mutation between them.
- $\underline{\mathscr{G}}^{n,r}$ is the graph whose vertices are graph isomorphism classes of UOMs of rank r and isometric dimension n. Two classes are connected if and only if there exists a mutation between them.

The three different mutation graphs are related as follows:

Observation 3.1. Let $0 \le r \le n$. We have $\overline{\mathscr{G}}^{n,r}$ connected $\Longrightarrow \mathscr{G}^{n,r}$ connected $\Longrightarrow \underline{\mathscr{G}}^{n,r}$ connected.

Proof. If $\overline{\mathscr{G}}^{n,r}$ is connected, then also $\mathscr{G}^{n,r}$ is, since there is a weak homomorphism from the first to the second, mapping an OM to its equivalence class. Similarly, if $\mathscr{G}^{n,r}$ is connected then also $\underline{\mathscr{G}}^{n,r}$ is, since a reorientation can be seen as an isomorphism.

We start by analyzing the connectivity of mutation graphs for small rank. Since OMs of rank 1 or 2 are simply isomorphic to an edge or an even cycle, respectively, the first interesting case is when the rank of OMs is 3. Our first result on the topic, together with Observation 3.1 settles the connectivity of all three graphs in rank 3:

Proposition 3.2. For every n the graph $\overline{\mathscr{G}}^{n,3}$ is connected.

Proof. It is a well known fact, implied by the Topological Representation Theorem [18], that OMs of rank 3 can be represented as pseudo-circle arrangements on a 2-dimensional sphere. Further, this arrangement can be assumed to be centrally symmetric and then by restricting the sphere along an equator that does not contain any intersection of the pseudo-circles, one obtains a wiring diagram, see Figure 4, i.e., a set of pseudo-lines traversing from left to right pairwise intersecting exactly once. Note that different choices of equator yield different wiring diagrams, but the all represent the OM in the same way, and only differ in what line is the topmost entering on the left. Then, every pseudoline splits the plane into positive and negative half, and every point in the plane yields a covector, by associating to each pseudo-line a coordinate and putting a +, -, 0 in it, depending on the point being on positive, negative side or on the pseudo-line. This, way for each covector Y of the OM exactly one of Y, -Y is represented. Maximal cells correspond to topes where two are adjacent if the cells intersect on a line-segment. The OM being uniform corresponds to arrangement being *simple*, i.e., no three lines cross in a point. A *mutation* of the wiring-diagram means pulling a bounding line of a simplicial cell over its opposite corner. A special case is if the simplicial cell is crossed by the bounding vertical lines, see Figure 4. This operation corresponds to what we described as mutation in tope graphs.

We now use the proof of Ringel's Homotopy Theorem [48, 49] as shown in [7, Section 6.4]. Ringel shows that any two labeled simple wiring diagrams can be transformed into another by performing mutations.

Suppose now that we have two UOMs of rank three \mathscr{M} and \mathscr{M}' and want to transform one into the other by mutations. Both can be represented by wiring diagrams, as stated above. Then by Ringel's Theorem we can assume that \mathscr{M} can be transformed into a reorientation of \mathscr{M}' . So to prove that $\overline{\mathscr{G}}^{n,3}$ is connected we only need to consider the case where \mathscr{M} and \mathscr{M}' differ in the reorientation of one element e. Represent \mathscr{M} as a wiring-diagram \mathscr{A} , where e is the top-element on the left and construct a wiring diagram \mathscr{A}' as shown in Figure 4, with e being the bottom element on the left. Note that \mathscr{A}' represents \mathscr{M}' , and the mutations transforming \mathscr{A} into \mathscr{A}' by Ringel's Theorem push the line e without changing its orientation. Thus, we have obtained \mathscr{M}' from \mathscr{M} by mutations.

We show that two of the open questions about the connectivity of mutation graphs are equivalent.




Figure 4: How to construct the new pseudoline arrangement in the proof of Proposition 3.2. The arrow points to the positive side of *e*. The grey cell is the simplicial cell on which a mutation is applied.

$n \backslash r$	2	3	4	5	6	7	8	9	10
2	1	1	1	1	1	1	1	1	1
3		1	1	1	4	11	135	482	312356
4			1	1	1	11	2628	9276595	?
5				1	1	1	135	9276595	?
6					1	1	1	4382	?
7						1	1	1	312356
8							1	1	1
9								1	1
10									1

Table 1: Known orders of $\mathcal{G}^{n,r}$, retrieved from http://www.om.math.ethz.ch/.

Proposition 3.3. If $\mathcal{G}^{n,r}$ is connected, then $\mathcal{G}^{n,r}$ is connected.

Proof. Notice that the property of being a realizable OM is independent of reorientation or permuting the elements. If $\underline{\mathscr{G}}^{n,r}$ is connected, then there exists a sequence of mutations from any $[A] \in \underline{\mathscr{G}}^{n,r}$ to a realizable class $[B] \in \underline{\mathscr{G}}^{n,r}$. This sequence can then be lifted to a sequence of mutations from any $A' \in \mathscr{G}^{n,r}$ to a realizable $B' \in \mathscr{G}^{n,r}$. This proves that there exists a path from every $A' \in \mathscr{G}^{n,r}$ to a reorientation classes of realizable OMs. Since by [50] the induced subgraph of all realizable classes in $\mathscr{G}^{n,r}$ is connected, this proves that $\mathscr{G}^{n,r}$ is connected.

Proposition 3.3 allows to approach Conjecture 3 for small values of n and r from the computational perspective, since it allows computations on the smaller graph $\underline{\mathscr{G}}^{n,r}$. To provide an idea of the computational weight of this task, Table 1 shows the known orders of such graphs $\underline{\mathscr{G}}^{n,r}$ for small n and r. Moreover, Figure 5 displays the graph $\underline{\mathscr{G}}^{8,4}$.

We verified computationally that for all the parameters from Table 1 where the isomorphism classes of OMs are known, their mutation graph $\underline{\mathscr{G}}^{n,r}$ is connected. By Proposition 3.3 also the corresponding $\mathscr{G}^{n,r}$ are connected. This was possible by considering UOMs as (tope)graphs in which finding possible mutations is easy, since only degrees of vertices need to be checked. We calculated the isomorphism class of the mutated graphs using the software Bliss [28] that is designed for calculations of isomorphisms of graphs. The computationally most demanding task was the graph $\underline{\mathscr{G}}^{9,4}$ where efficient graph representation was needed.

Corollary 3.4. The graph $\mathcal{G}^{n,r}$ is connected for $n \leq 9$.

Testing graph isomorphism instead of OM-isomorphism was an essential ingredient in order to obtain Corollary 3.4. Checking connectivity of $\overline{\mathscr{G}}^{n,r}$ is far more demanding. We do not know anything about the connectivity of this graph beyond rank 3, see Proposition 3.2.




Figure 5: The graph $\underline{\mathscr{G}}^{8,4}$ where red vertices are isomorphism classes of realizable UOMs and blue ones are non-realizable

4 Mandel's OMs and Euclideanness

In this section we focus on Las Vergnas' and Mandel's Conjectures. The main result is that Mandel OMs are Θ -Las Vergnas and therefore not all OMs are Mandel. In order to make the result accessible to readers familiar with the OM terminology as well as to readers more familiar with graphs we try to argue in the proof along both views.

The concept of Euclideanness is based on the orientations of cocircuit graph introduced in Section 2. Furthermore, the property of being Mandel relies on the definition of extensions in general position and Euclideanness.

Let G be an OM of rank r. Recall the definition of cocircuit graph G^* from Section 2. Graph G^* has cocircuits of G for vertices, i.e. in graph theoretical language the antipodal subgraphs of G of rank r-1. Two cocircuits X,Y are adjacent if there exists a covector Z with $X,Y \prec Z$ (hence Z has rank r-2) in the big face lattice $\mathscr{F}(G)$, or equivalently if the intersection of the antipodal subgraphs corresponding to X and Y in G is an antipodal subgraph of rank r-2.

Consider now a maximal subset F of the Θ -classes of G such that there exist a covector Z of rank r-2 with $Z_e=0$, for all $e\in F$. In other words, there exists an antipodal subgraph of rank r-2 crossed by the Θ -classes in F. Let X_1,\ldots,X_n be a maximal path in G^* such that $(X_i)_e=0$ for all $e\in F$, i.e. all the corresponding antipodal subgraph are crossed by the Θ -classes in F. It follows from the topological representation of OMs [37], that X_1,\ldots,X_n induce a cycle in G^* . Moreover, $X_{n/2+i}$ is the antipode of X_i in G, for $1\leq i\leq n/2$. Furthermore, for each Θ -class $E_f\notin F$ there are exactly two antipodal cocircuits X_i and $X_{n/2+i}$ such that $(X_i)_f=(X_{n/2+i})_f=0$, meaning that E_f crosses exactly two pairwise antipodal subgraphs corresponding to X_i and $X_{n/2+i}$ on the cycle. The cocircuit graph G^* is the (edge-disjoint) union of such cycles.

Let now H be a halfspace of G, i.e. H is (the tope graph of) an AOM. The induced subsequence X_k, \ldots, X_ℓ of the above cycle is called a *line L* in H. This name comes from the fact that in the topological

representation the sequence corresponds to a pseudo-line, see [37]. Let now e be an element (or E_e be a Θ -class) of H. We say that e (or E_e) crosses a line E of E_e 0, i.e. the corresponding antipodal subgraph is crossed by E_e 1. Note that for a line E2 of an AOM, if there exists E3 with E4 with E5 and E6. Equivalently, exactly one or all of the antipodal subgraphs on the line are crossed by E6. This allows to define the orientation of line E4 with respect to E6 (or E6): If E6 is not crossed by E7 (equivalently E8) we leave its edges undirected. Otherwise, let E6 be the element of E7 that is crossed by it and assume that E8 and E9 is directed from E9 to E9. We orient all edges of the form E9 to E9. This is, the path E8 is directed from E9 to E9.

The edges of the cocircuit graph H^* of an AOM H are partitioned into lines and for every element e (or Θ -class E_e) we obtain a partial orientation of H^* by orienting every line with respect to E_e . Let us call this mixed graph the orientation of H^* with respect to E_e . Following Mandel [37, Theorem 6], an AOM is *Euclidean* if for every element e (or Θ -class E_e) the orientation of the cocircuit graph H^* with respect to e is *strictly acyclic*, i.e., any directed cycle (following undirected edges or directed edges in the respective orientation) consists of only undirected edges. In other words, any cycle that contains a directed edge contains one into each direction.

Following Fukuda [19], an OM is called *Euclidean* if all of its halfspaces are Euclidean AOMs. Since non-Euclidean AOMs exist, see [19,37], also non-Euclidean OMs exist. However, there is a larger class of OMs that inherits useful properties of Euclidean AOMs and that was introduced by Mandel [37].

We call an OM Mandel if it has an expansion in general position such that G_1 (and implicitly G_2 , since they are isomorphic with the antipodal map) are Euclidean AOMs. In terms of the covectors description, this can be stated as follows: if $\mathscr L$ is a set of covectors of an OM, and there exists $\mathscr L'$ a single-element extension in general position of it with respect to an element e, i.e. $\mathscr L' \setminus e = \mathscr L$, with the property that $\{X \in \mathscr L' \mid X = +\}$ induces an Euclidean AOM, then $\mathscr L$ is said to be Mandel. Mandel [37, Theorem 7] proved (and it is up to today the largest class known to have this property) that these OMs satisfy the conjecture of Las Vergnas:

Theorem 4.1 ([37]). *If an OM G is Mandel, then it has a simplicial vertex.*

In the following we go along the lines of Mandel's proof of Theorem 4.1 from [37] while refining it to such an extent, that we can disprove Conjecture 2. In particular, we use a stricter version of the induction in the proof to obtain Θ -Las Vergnas properties of Mandel OMs.

We shall repeatedly use the following fact, that is trivial with the covectors definition of OMs but is a bit more complicated to prove in the graph view. Nevertheless, it was also proved with graphs in [31, Lemma 6.2].

Observation 4.2. Let G, G' be OMs such that G' is an OM deletion (i.e. a partial cube contraction) of G, with respect to an element e (a Θ -class E_e). Then for every cocircuit X' of G' there exists a cocircuit X in G, such that $X_f = X_f'$ for all $f \neq e$. In graph theoretical view, there exists a maximal antipodal subgraph of G that contracts to the maximal antipodal subgraph corresponding to X'. The same holds for COMs.

The following is a characterization of simplicial vertices in an OM, that can be found in [37, Proposition 5, page 320] and [34, Proposition 1.4]. We provide a slightly different formulation together with a formulation in terms of the tope graph. Recall that in an OM G a vertex v is simplicial if the degree of v coincides with the rank of G. Also recall, that by definition G is Θ -Las Vergnas if for every Θ -class E_e of G, there exists a simplicial vertex incident with E_e . In terms of the covectors description, this means that for every element e there exists a simplicial tope E and a covector E with E and E such that E and E will say that E is incident with E.

Lemma 4.3. Let G be an OM of rank r. A tope $v \in G$ is simplicial if and only if there is a set of elements F of size r and r cocircuits, such that for each $e \in F$ there is a corresponding cocircuit X with $X_f = 0$ for $f \in F \setminus \{e\}$, $X_e \neq 0$ and $X \leq v$. In the graph interpretation, this is equivalent to $r \in G$ -classes F and r

antipodal subgraphs incident with v such that each is crossed by $F \setminus \{E_e\}$ but not by E_e , each for a different E_e . In this case, v is incident with elements (Θ -classes) in F.

As a last basic ingredient for the Theorem 4.5 we need the following. Let G be the tope graph of a simple OM and e one of its elements. Let v be a tope in the positive halfspace of G with respect to e. Since the lattice of covectors is atomistic and graded, there is a set of cocircuits such that their join is exactly v. In particular, for at least one of the latter cocircuits, say X, it must hold $X_e \neq 0$. This is:

Observation 4.4. Let e be an element (E_e be a Θ -class) of a simple OM G. There exists a cocircuit X (a maximal proper antipodal subgraph A) such that $X_e \neq 0$ (A is not crossed by E_e).

We are prepared to give the main theorem of the section. In the proof we will make use of the direct sum $G_1 \oplus G_2$ of OMs G_1 and G_2 . In the OM language this is defined as the OM with covectors $\{(X_1, X_2) \mid X_1 \in G_1, X_2 \in G_2\}$. Considering OMs as graphs, this is equivalent to the *Cartesian product* of graphs G_1, G_2 , being defined as the graph $G_1 \square G_2$ whose vertices are $V(G_1) \times V(G_2)$ where two vertices $(x_1, y_1), (x_2, y_2)$ adjacent if and only if $x_1 = x_2$ and y_1 is adjacent to y_2 in y_2 or $y_1 = y_2$ and y_2 is adjacent to y_2 in y_2 or y_2 and y_3 is adjacent to y_3 in y_4 or y_4 and y_5 in y_5 and y_6 in y_6 or equivalently the unique simple OM with ground set of size 1.

Theorem 4.5. Let G be a simple, Mandel OM of rank r and e one of its elements (E_e a Θ -class of G). Then there is a simplicial tope (a vertex of degree r) in G, incident with e (incident with E_e), i.e., G is Θ -Las Vergnas.

Proof. By the definition of Mandel OM, there exists a single-element extension (an expansion) of G in general position such that the two halfspaces in the obtained OM are Euclidean AOMs. Both halfspaces correspond to intersecting subsets of covectors of G, say G_1 and G_2 . Speaking of OMs as graphs this means that G is expanded along G_1 and G_2 . Let $E_e \in \mathcal{E}$ be a Θ -class of G. We prove the slightly stronger assertion (than asserted in the theorem) that there exists a simplicial tope V (i.e. of degree P) in $G_1 \setminus G_2$ incident with E_e , for any chosen G_1, G_2 , making G Mandel. We will proceed by induction on the size of G and distinguish two cases:

Case 1. *G* is not the direct sum (the Cartesian product) with factor K_2 , where the factor K_2 corresponds to element $e(\Theta$ -class $E_e)$.

By Observation 4.4 and the antipodality, G has at least two cocircuits X_1, X_2 (maximal antipodal subgraphs) with the property $(X_1)_e, (X_2)_e \neq 0$ (not crossed by E_e). Then at least one lies in G_1 . Without loss of generality assume that it is in E_e^- , otherwise reorient E_e . This is, there is a line in G_1 that has at least one cocircuit X_1 (an antipodal subgraph of rank r-1) with $(X_1)_e = -$ (in E_e^-).

Orient the lines of G_1 with respect to E_e . Note that every line of G_1 is a subline of a cycle in G^* that either has all its cocircuits X with $X_e = 0$ (i.e. maximal proper antipodal subgraphs crossed by E_e) or exactly two of them. Since the expansion according to G_1 and G_2 is in general position this implies that every line of G_1 is either crossed by E_e in exactly one cocircuit, or in all of them. Thus, all the lines are oriented, except the ones with all its cociruits X (antipodal subgraphs) having $X_e = 0$ (being crossed by E_e).

By the definition of Euclideaness, the orientation is strictly acyclic, hence we can find in $G_1 \cap E_e^-$ a cocircuit (i.e. an antipodal subgraph of rank r-1) X such that each of its out-neighbors Y in the cocircuit graph has $Y_e = 0$ (i.e. is intersected by E_e). Let X^0 be the set of all elements (Θ -classes E_f), such that $X_f = 0$ (that cross X). Let H be the OM-deletion (a partial cubes contraction) of G along all its elements (G-classes) besides G (G-classes) and the ones in G-classes) besides G-classes and the ones in G-classes are image of a covector in G-classes are image. Then by Observation 4.2, each covector of G-classes are image of a covector in G-classes are image. In graph theoretical language, this means that all the antipodal subgraphs of G-classes are image. This proves that G-classes are image. The cover image of G-classes are image. In G-classes are image, this means that all the antipodal subgraphs of G-classes are image. The cover image of G-classes are image. The cover image image image is G-classes are image. The cover image i

H with respect to E_e is strictly acyclic as well. This proves that H is Mandel by the expansion in general position according to H_1 and H_2 .

By definition, H is obtained by deleting elements f (contracting Θ -classes), for which $X_f \neq 0$ (not crossing X). It is not hard to see, that the sub-lattice of covectors Y with $X \leq Y$ has a maximal chain of length r-1 which is unaffected by the deletion. Thus, the rank of H is r since it properly contains X of rank r-1. In graph theoretical sense, this also follows from the gatedness of antipodal subgraphs in an OM implying that the structure of the antipodal subgraph corresponding to X is not affected by any of the contractions. We will denote by X' the cocircuit obtained from X by the deletion.

Let $\overline{X'}$ be the antipode of X' in H. Then $\overline{X'}$ is also a cocircuit of H and only e separates X' and $\overline{X'}$. As graphs they are disjoint antipodal subgraph connected by edges in E_e . Thus $H \cong H' \oplus K_2$, where H' consist of all Y with $X \leq Y$ in H. In the graph language, $H \cong A \square K_2$, where A is the antipodal subgraph corresponding to X'. Since we are in the Case 1, this gives that H is strictly smaller than G.

By the induction assumption, H has a tope v' of degree r in $H_1 \setminus H_2$, incident with e (with E_e). In fact, since $H \cong H' \oplus K_2$ (or $H \cong H' \Box K_2$) with the K_2 factor corresponding to e (E_e), the latter holds for all the topes in H. By Lemma 4.3, there is a set \mathscr{B} of r cocircuits and a set F of r Θ -classes, such that for each $E_g \in F$ there is a corresponding cocircuit Y with $Y_f = 0$ for $f \in F \setminus \{E_g\}$, $Y_g \neq 0$ and $Y \leq v'$. Note that $E_e \in F$. In other words, there are e antipodal subgraphs of rank e e 1 incident with e 1 incident with e 1 incident with e 2 with e 2 v', we have e 3. By assumption e 4 and since e 1 and since e 4 define an expansion in general position, all the members of e 3 are in e 4.2, there is a set e 0 of e 2 cocircuits of e 3 to 4 to 5 degree e 3. Moreover, since the members of e 3 are in e 4 contract to corresponding antipodal subgraphs of e 3. Moreover, since the members of e 3 are in e 4 are in e 6. Clearly, e 6.

Consider a tope $v \in G_1$ such that $X \leq v$ (v is an element of the antipodal subgraph corresponding to X) and the OM-deletion (partial cube contraction) maps v to v'. To prove that v has degree r in G, by Lemma 4.3, it suffices to prove that for each cocircuit Y in $\mathscr C$ it holds $Y \leq v$ (the antipodal subgraphs in $\mathscr C$ include v). Let $Y \in \mathscr C \setminus \{X\}$, and Y' the corresponding cocircuit (antipodal subgraph) in $\mathscr B$. Then Y' and X' are adjacent in H^* , hence X and Y lie on a line in G_1 . Since $Y'_e = 0$ also $Y_e = 0$, i.e. the corresponding antipodal subgraphs are crossed by E_e . Thus the line they are both on is oriented from X to Y. Since X was chosen to have all its out-neighbors crossed by E_e , Y must be adjacent to it. This implies that for each $E_f \notin X^0 \cup \{E_e\}$, $Y_f \neq -X_f$. In particular, since $X \leq v$ and by assumption $Y' \leq v'$, the former implies that $Y \leq v$. In the tope graphs this is seen as v being in the intersection of the antipodal subgraphs corresponding to X' and Y'.

We have found a tope ν that has degree r. Since e (or E_e) is in F, it is incident with it. By construction it lies in $G_1 \setminus G_2$.

Case 2. G is the direct sum (the Cartesian product) with factor K_2 , where the factor K_2 corresponds to element $e(\Theta$ -class $E_e)$.

If G has an element f (a Θ -class E_f) such that G is not the direct sum with factor K_2 corresponding to it, then by Case 1, G has a simplicial tope v (of degree r) in $G_1 - G_2$. Moreover, in this case all the vertices of G are incident with E_e , in particular also v is.

If all of the elements (Θ -classes) of G correspond to factors K_2 , then G is a hypercube and all its vertices are simplicial and incident to E_e .

OMs with a Θ -class not incident to a simplicial vertex have been found of different sizes [8, 47, 53]. We conclude:

Corollary 4.6. There exist OMs that are not Mandel. The smallest known such OM has 598 vertices (topes), isometric dimension 13 (elements) and is uniform of rank 4. See Figure 1.

5 Corners and corner peelings

In [9] the corners of lopsided sets (LOPs) have been defined as vertices of degree r that lie in a unique maximal hypercube subgraph of dimension r. This has a nice correspondence to the results from Section 4: If G is a LOP and also an AOM, i.e, a halfspace E_e^+ of an OM G', then G has a corner if and only if G' has a simplicial vertex incident to E_e . By the examples from [3,9,53] this proves that there are LOPs without corners. This was first observed in [9], where it is translated to an important counter example in computational learning theory.

In the present section we introduce corners and corner peelings for general COMs. The first subsection is concerned with the first definitions and results, and in particular contains a proof for existence of corner peelings of realizable COMs. The second subsection contains corner peelings for COMs of rank 2 and hypercellular graphs. Since the results in this section are more concerned with graphs, we use the graph theoretic language and use covectors only when needed.

5.1 First definitions and basic results

We will approach our general definition of corner of a COM, that generalizes corners on LOPs and has strong connections with simplicial vertices in OMs. The intuitive idea of a corner in a COM, is a set of vertices whose removal gives a new (maximal) COM. As a matter of fact it is convenient for us to first define this remaining object and moreover within an OM.

Recall the definition of an expansion in general position from Section 2. We will say that the subgraph T of an OM H is a *chunk* of H, if H admits an expansion in general position H_1, H_2 , such that $T = H_1$. We call the complement $C = H \setminus H_1$ a *corner* of H. In the case that H has rank 1, i.e. H is isomorphic to an edge K_2 , then a corner is simply a vertex of H.

This definition extends to COMs by setting C to be a *corner* of a COM G if C is contained in a unique maximal antipodal subgraph H and C is a corner of H. We need two more helpful observation:

Lemma 5.1. If G' is an isometric subgraph of a COM G such that the antipodal subgraphs of G' are antipodal subgraphs of G, then G' is a COM.

Proof. By Theorem 2.1 all antipodal subgraphs of G are gated, but since G' is an isometric subgraph and it has no new antipodal subgraph also the antipodal subgraphs of G' are gated. Thus, by Theorem 2.1 G' is a COM.

We are now ready to prove that chunks and corners as we defined them achieve what we wanted. This proof uses the correspondence between sign-vectors and convex subgraphs as introduced in Section 2.

Lemma 5.2. If C is a corner of a COM G, then the chunk $G \setminus C$ is an inclusion maximal proper isometric subgraph of G that is a COM.

Proof. Let us first consider the case where G = H is an OM. Let T, -T be an expansion of H in general position, i.e., -T is the set of antipodes of T. Since expansions in general position are OM-expansions, T is a halfspace of an OM. Thus, T is a COM — even an AOM. This proves that T is a sub-COM. Assume that it is not maximal and let $R \supseteq T$ be a COM contained in H. Let $X \subseteq R$ be an antipodal subgraph of H that is not completely in T, and is maximal with this property. Since the expansion is in general position, it holds $X \subseteq -T$ and $-X \in T \subseteq R$. Let $E_e \in S(X, -X)$, i.e., E_e separates X from -X. Such E_e clearly exists, since R is a proper sub-COM of H, thus $X \ne H$.

Considering R as a COM we apply (SE) to X, -X with respect to E_e in order to obtain $Z \subset R$ that is crossed by E_e . Note that X and -X are crossed by the same set of Θ -classes X^0 . By (SE) the set Z^0 of Θ -classes crossing Z strictly contains X^0 . Thus, if $S(Z,X) = \emptyset$, then Z is an antipodal subgraph containing X, i.e., X was not a maximal antipodal subgraph of R. Let otherwise $E_f \in S(Z,X)$. Apply (SE) to X,Z with respect to E_f in order to obtain $Z' \subset R$ which is crossed by E_f . Since $Z^0 \supseteq X^0$, we have $Z'^0 \supseteq X^0$ and furthermore $S(Z',X) \subseteq S(Z,X)$. Proceeding this way, we will eventually obtain an $\widetilde{Z} \subseteq R$

with $\widetilde{Z}^0 \supseteq X^0$ and $S(\widetilde{Z},X) = \emptyset$. Thus, $\widetilde{Z} \in R$ is an antipodal subgraph containing X. By the choice of X, \widetilde{Z} is not completely in T. This violates the assumption that X was maximal. Thus, R = T.

Now, let G be a COM that is not an OM and let H be the unique maximal antipodal subgraph of G containing G. By the above $T = H \setminus G$ is an isometric subgraph of H and a COM. Now, it follows that $G \setminus G$ is an isometric subgraph of G. Namely, since no vertex of G is adjacent to a vertex of $G \setminus H$ and G being an isometric subgraph of G, all shortest paths in G through G, can be replaced by shortest paths through G. Finally, Lemma 5.1 implies that $G \setminus G$ is a COM. Maximality follows from the first paragraph. G

Recall that simplicial vertices in LOPs are called corners. Before providing further central properties of corners in COMs, let us see that we indeed generalize corners of LOPs.

Proposition 5.3. A subset C of the vertices of a LOP G is a corner if and only if $C = \{v\}$ is contained in a unique maximal cube of G.

Proof. OMs that are LOPs are cubes, so let ν be a vertex of Q_n . The expansion with $H_1 = Q_n \setminus \{\nu\}$ and $H_2 = Q_n \setminus \{-\nu\}$ clearly is antipodal. Moreover, every proper antipodal subgraph of Q_n is contained in either H_1 or H_2 . Thus, this expansion is in general position. Consequently ν is a corner or Q_n . Since chunks are maximal sub-COMs, by Lemma 5.2, single vertices are precisely the corners of Q_n .

If now v is a corner of a LOP G, then by the definition of corners of COMs and the fact that in a LOP all proper antipodal subgraphs are cubes, v is contained in a unique maximal cube of G.

Conversely if ν is a vertex of a LOP contained in a unique cube, then this cube is also the unique maximal antipodal subgraph of the LOP, since in a LOP all proper antipodal subgraphs are cubes. Thus ν is a corner of the LOP.

Note that, as mentioned earlier, every OM admits an expansion in general position, see [51, Lemma 1.7] or [7, Proposition 7.2.2]. This yields directly from the definition:

Observation 5.4. Every OM has a corner.

Note however that COMs do not always have corners, e.g., with Proposition 5.3 one sees that the AOMs obtained from the UOMs with a mutation-free element have no corner.

Lemma 5.2 yields the following natural definition. A *corner peeling* in a COM G is an ordered partition C_1, \ldots, C_k of its vertices, such that C_i is a corner in $G - \{C_1, \ldots, C_{i-1}\}$. In the following we generalize a results from [53] for realizable LOPs.

Proposition 5.5. Every realizable COM has a corner peeling.

Proof. We show that a realizable COM G has a realizable chunk T. Represent G as a central hyperplane arrangement \mathcal{H} in an Euclidean space intersected with an open polyhedron P given by open halfspaces O. Without loss of generality we can assume that the supporting hyperplanes of the halfspaces in O are in general position with respect to the hyperplanes in \mathcal{H} . We shall call points in the Euclidean space, that can be obtained as intersection of subset of hyperplanes \mathcal{H} minimal dimensional cells. It follows from the correspondence between antipodal subgraphs and covectors of a COM [31, Theorem 4.9], that topes (chambers) surrounding minimal dimensional cells correspond to antipodal subgraphs of G.

Now, take some halfspace $O \in \mathcal{O}$ and push it into P until it contains the first minimal dimensional cell C of \mathcal{H} . The obtained realizable COM T is a chunk of G, because restricting the antipodal subgraph (an OM) corresponding to the cell C with respect to O is taking a chunk of C, while no other cells of G are affected and the resulting graph T is a COM.

In [4, Conjecture 2] it was conjectured that all *locally realizable* COMs, i.e., those whose antipodal subgraphs are realizable OMs, are realizable. Proposition 5.5 yields a disproof of this conjecture, since all antipodal subgraphs of a LOP are hypercubes, i.e., LOPs are locally realizable, but by the example in Figure 1 and others there are LOPs that do not have corner peelings. Thus, they cannot be realizable.

Remark 5.6. There are locally realizable COMs, that are not realizable.

Indeed, LOPs are even *zonotopally realizable*, i.e., one can choose realizations for all cells such that common intersections are isometric. It remains open whether every locally realizable COM is zonotopally realizable, see [4, Question 1].

5.2 Corners and corner peelings in further classes

In this section we consider the question of the existence of corners and corner peelings in various classes of graphs. By Proposition 5.3 simplicial vertices in LOPs are corners. Thus, Theorem 4.5 yields:

Corollary 5.7. *Every halfspace of a Mandel UOM has a corner.*

In the following we focus on COMs of rank 2 and hypercellular graphs. In both these proofs we use the zone graph $\zeta_f(G)$ of a partial cube G with respect to a Θ -class f, see [30]. In general the zone graph is not a partial cube, but indeed a characterization of COMs from [31] (generalizing a result of Handa for OMs [24]) allows the following definition in COMs. Let G be a COM, and F a subset of its Θ -classes. The zone graph $\zeta_F(G)$ is the graph obtained from G, whose vertices are the minimal antipodal subgraphs of G that are crossed by all the classes in F. It turns out that all such antipodal subgraphs have the same rank, say r. Two such antipodal subgraphs are connected in $\zeta_F(G)$ if they lie in a common antipodal subgraph of G of rank G of rank

Cocircuit graphs of COMs

In the following we generalize the concept of orientation of the cocircuit graph introduced in Section 4 from AOMs to general COMs.

Lemma 5.8. If G is a COM and a hypercube Q_r a minor of G, then there is an antipodal subgraph H of G that has a Q_r minor. In particular, the rank of a maximal antipodal subgraph of a COM G is the rank of G.

Proof. Since Q_r is antipodal, by Lemma 4.2, there exist an antipodal subgraph H of G that contracts to it. Then H is the desired subgraph.

We define the *cocircuit graph* of a non-antipodal rank r COM as the graph whose vertices are the rank r antipodal subgraphs and two vertices are adjacent if they intersect in a rank r-1 antipodal subgraph. By Lemma 5.8 the vertices of the cocircuit graph of a non-antipodal COM G correspond to the maximal antipodal subgraphs of G. The cocircuit graph of a COM can be fully disconnected hence we limit ourselves to COMs having all its maximal antipodal subgraphs of the same rank with G^* connected. We call them *pure* COMs. Note that AOMs are pure COMs.

Let G be a pure COM, $\{A_1, A_2\}$ be an edge in G^* and F be the set of Θ -classes crossing $A_1 \cap A_2$. We have seen in Section 4 that if G is an AOM, then the maximal proper antipodal subgraphs of G crossed by Θ -classes in F induce a path of G^* which we called a line. The following lemma is a generalization of the latter and of general interest with respect to cocircuit graphs of COMs, even if we will use it only in the case of rank 2.

Lemma 5.9. Let G be a COM that is not an OM, $\{A_1, A_2\}$ be an edge in G^* , and F be the set of Θ -classes crossing $A_1 \cap A_2$. Then the maximal proper antipodal subgraphs of G crossed by Θ -classes in F induce a subgraph of G^* isomorphic to the line graph of a tree.

Proof. Let G, A_1, A_2, F be as stated and r the rank of G. Consider the zone-graph $\zeta_F(G)$. Recall that its vertices are antipodal subgraphs of rank r-1 crossed by Θ -classes in F where two subgraphs are adjacent if they lie in a common rank r antipodal subgraph. Since $\zeta_F(G)$ is a COM, see e.g. [31] and has rank 1, we have that $\zeta_F(G)$ is a tree. By definition, the maximal proper antipodal subgraphs of G crossed by G-classes in G correspond to edges of G with two such edges connected if they share a vertex. Hence they form a subgraph of G that is isomorphic to the line graph of G.

Lemma 5.9 implies that G^* can be seen as the edge disjoint union of line graphs of trees. We can use this to orient edges of G^* . Similarly as in the settings of AOMs, we will call a *line* in G a maximal path $L = A_1, \ldots, A_n$ in the cocircuit graph G^* such that $A_{i-1} \cap A_i$ is the set of antipodes of $A_i \cap A_{i+1}$ with respect to A_i . Let now $E_e \in \mathscr{E}$ be a Θ -class of G. Similarly as before we say that E_e crosses a line E_e of E_e if there exists E_e or E_e is the orientation of that is crossed by E_e but E_e is the orientation of the path E_e in E_e in E_e or E_e is the orientation of the path E_e or E_e in E_e in E_e or E_e in E_e in E_e by orienting all the lines simultaneously. The orientation of each edge (if it is oriented) is well defined: If E_e is an edge in a line graph of a tree that is crossed by E_e in E_e and E_e in E_e is closer to E_e than E_e is, then E_e is oriented from E_e to E_e in E_e and E_e in E_e and E_e is closer to E_e than E_e is, then E_e is oriented from E_e to E_e in E_e and E_e in E_e in than E_e is oriented from E_e to E_e in E_e and E_e in E_e in than E_e in than E_e in that E_e is oriented from E_e to E_e in E_e and E_e in E_e in than E_e in that E_e is oriented from E_e to E_e in E_e and E_e in the oriented if E_e in the same distance to E_e in the oriented from E_e in the oriented if E_e in the oriented if E_e in the same distance to E_e is oriented from E_e in the oriented in this way we can orient the edges of E_e in the oriented if E_e in the oriented in E_e in E_e in the oriented in E_e in E_e in the oriented in E_e in the oriented in E_e in E_e in the oriented in E_e in




Figure 6: A pure rank 2 COM and its cocircuits graph oriented with respect to E_e .

COMs of rank 2

Mandel proved that every AOM of rank 2 is Euclidean, which by Corollary 5.7 implies that every rank 2 halfspace of a UOM has a corner. We generalize this result.

Let us first consider what corners in rank 2 COMs are. Up to isomorphism the only rank 2 OMs are even cycles. An expansion in general position of an even cycle $G = C_{2n}$ is given by $G_1, G_2 = -G_1$, where G_1 consists of an induced path on n+1 vertices. Hence a corner in a rank 2 COM consist of n-1 vertices inducing a path, included in a unique antipodal C_{2n} . For example, the COM in Figure 6 has 11 corners. Those contained in a square are single vertices and the ones contained in the C_6 are paths with two vertices.

Proposition 5.10. Let G be a pure COM of rank 2 and E_e a Θ -class of G. Then G has a corner in E_e^+ and in E_e^- .

Proof. Let $E_e \in \mathcal{E}$ be a Θ-class of a pure rank 2 COM G. We shall prove that G has a corner in E_e^+ , by symmetry it follows that it has one in E_e^- as well. Without loss of generality assume that there is no Θ-class E_f completely contained in E_e^+ , otherwise switch E_e^+ with E_f^+ or E_f^- depending on which is entirely in E_e^+ . Orient the edges of G^* with respect to E_e .

Since the rank of G is 2, the maximal antipodal subgraphs are even cycles and each line consist of sequence of cycles pairwise crossing in edges from E_f , for some E_f . We will say that the line *follows* E_f . The induced subgraph of G^* of all the maximal antipodal subgraphs crossed by E_f is the line graph of a tree, by Lemma 5.9. We will denote it by G_f^* . Let G_f^* be crossed by some E_g in A_i . Then this splits the

vertices of $G_f^* - \{A_i\}$ into the ones lying in E_g^+ and the ones lying in E_g^- . We denote these by $E_g^+(G_f^*)$ and $E_{\sigma}^{-}(G_{f}^{*})$, respectively.

We shall prove that there is no directed cycle in $G^* \cap E_{\rho}^+$ consisting of only directed edges. For the sake of contradiction assume that such a cycle exists and take one that is the union of as few parts of lines as possible. Let the cycle be a union of a part of L_1 , a part of L_2 ,..., and a part of L_n . Also denote by E_{e_1}, \ldots, E_{e_n} the respective Θ -classes followed by L_1, \ldots, L_n .

Since L_i and L_{i+1} intersect, L_i must be crossed by $E_{e_{i+1}}$ and L_{i+1} must be crossed by E_{e_i} . Without loss of generality assume that L_2 passes E_{e_1} from $E_{e_1}^-$ to $E_{e_1}^+$, L_3 passes E_{e_2} from $E_{e_2}^-$ to $E_{e_2}^+$, ..., and L_1 passes E_{e_n} from $E_{e_n}^-$ to $E_{e_n}^+$, otherwise reorient E_{e_1}, \ldots, E_{e_n} .

First we show that each L_i passes $E_{e_{i+1}}$ from $E_{e_{i+1}}^+$ to $E_{e_{i+1}}^-$. Assuming otherwise the intersection of L_{i-1} and L_i lies in $E_{e_{i+1}}^-$, while the intersection of L_{i+2} and L_{i+3} lies in $E_{e_{i+1}}^+$, by the assumption in the previous paragraph. Then one of the lines $L_{i+4}, L_{i+5}, \dots L_{i-1}$ must pass $E_{e_{i+1}}$ from $E_{e_{i+1}}^+$ to $E_{e_{i+1}}^-$, say L_j passes it. If this passing is in $E_{e_i}^+$, then the cycle is not minimal, since one could just replace the lines L_{i+1}, \ldots, L_i by the line following $E_{e_{i+1}}$ starting from the intersection of L_i and L_{i+1} to the crossing of $E_{e_{i+1}}$ and L_j . In fact such a directed line exists since by assumption the lines following $E_{e_{i+1}}$ pass E_i from E_j to E_i^+ thus the orientation of the shortest path from the intersection of L_i and L_{i+1} to the crossing of L_j

On the other hand, assume the passing is in $E_{e_i}^-$. By assumption, the intersection of L_{i+1} and L_{i+2} is in $E_{e_i}^+$. Hence one of the lines $L_{i+3}, L_{i+4}, \dots L_j$ must pass E_{e_i} , say L_l . In particular it must pass it in $E_{e_{i+1}}^+$, by the choice of L_j . But then again the cycle is not minimal, since one could just continue on the line following E_{e_i} starting from the intersection of L_{i-1} and L_i to the crossing of E_{e_i} and L_i . This cannot be.

Now we show that for each e_i, e_{i+1} it holds that $E_{e_{i+1}}^-(G_e^*) \subset E_{e_i}^-(G_e^*)$. Since L_{i+1} passes E_{e_i} from $E_{e_i}^$ to $E_{e_i}^+$ it follows that L_{i+1} passes E_e in $E_{e_i}^-$. Since G_e^* is the line graph of a tree, this implies that either $E_{e_{i+1}}^+(G_e^*) \subset E_{e_i}^-(G_e^*)$ or $E_{e_{i+1}}^-(G_e^*) \subset E_{e_i}^-(G_e^*)$. On the other hand, L_i passes $E_{e_{i+1}}$ from $E_{e_{i+1}}^+$ to $E_{e_{i+1}}^-$ hence L_i passes E_e in $E_{e_{i+1}}^+$. Again since G_e^* is the line graph of a tree, this implies that $E_{e_i}^+(G_e^*) \subset E_{e_{i+1}}^+(G_e^*)$ or $E_{e_i}^-(G_e^*) \subset E_{e_{i+1}}^+(G_e^*). \text{ Hence, } E_{e_{i+1}}^-(G_e^*) \subset E_{e_i}^-(G_e^*) \text{ and } E_{e_i}^+(G_e^*) \subset E_{e_{i+1}}^+(G_e^*).$ Inductively $E_{e_n}^-(G_e^*) \subset E_{e_{n-1}}^-(G_e^*) \subset \cdots \subset E_{e_1}^-(G_e^*) \subset E_{e_n}^-(G_e^*) - \text{contradiction. This proves that there is}$

no directed cycle in $G^* \cap E_e^+$ consisting of only directed edges.

We can now prove that G has a corner in E_e^+ . First, assume that $G^* \cap E_e^+$ is non-empty and let $A \in G^* \cap E_e^+$ be a maximal antipodal subgraph, i.e., an even cycle, that has no out-edges in G^* . By the choice of E_e , each line L that passes A is crossed by E_e . We now analyze how lines pass A. Let L_1, L_2 be lines passing A, following E_{f_1} , E_{f_2} , respectively. Since E_{f_2} crosses at most one antipodal subgraph of $G_{f_1}^*$, this implies that L_1 and L_2 simultaneously pass only A. In particular each antipodal subgraph of G_e^* is passed by at most one line passing A. Since G_e^* is the line graph of a tree, its every vertex is a cut vertex. Then each line L, passing A and $A_f \in G_e^*$, and following some E_f , splits $G_e^* - \{A_f\}$ into two connected components, $E_f^+(G_e^*)$ and $E_f^-(G_e^*)$. Thus we can inductively find L such that any other line passing A passes an antipodal subgraph in G_e^* in $E_f^+(G_e^*)$, reorienting E_f if necessary.

We now show that A includes a corner. Let A' be an antipodal subgraph on L that is a neighbor of A. Then $A \cap A'$ corresponds to an edge in E_f . Define the set C to include all the vertices of A in E_f^- besides the one vertex lying in $A \cap A'$. Then C is a corner of A, we will show that C is a corner of G. For the sake of contradiction assume that a vertex ν of C lies in a maximal antipodal subgraph A'' different from A.

We prove that we can choose A'' such that it shares an edge with A. Assuming otherwise, since Gis a pure COM, there is a path in G^* between A and A". This implies that there is a cycle C_k in G with subpath v'vv'', where $v' \in A \setminus A''$ and $v'' \in A'' \setminus A$. By [11, Lemma 13], the convex cycles span the cycle space in a partial cube. If A is one of the convex cycles spanning C_k , then there is a convex cycle incident with A in v and sharing an edge with A. If A is not used to span C_k , then a convex cycle incident with edge v'v must be used, thus again we have a convex cycle sharing v and an edge with A.

We hence assume that A and A" share an edge g. By definition of C, either $g \in E_f^-$, or $g \in E_f$ but

not in $A \cap A'$. The latter case implies that L can be extended with A'', which cannot be since A in G^* has no out-edges. Moreover, by the choice of L, all the other lines passing A pass E_f form E_f^+ to E_f^- . Then in the former case, some other line passing A can be extended, leading to a contradiction. This implies that G has a corner.

Finally, consider the option that $G^* \cap E_e^+$ is empty. Since G_e^* is the line graph of a tree, we can pick $A \in G_e^*$ that corresponds to a pendant edge in a tree, i.e. an edge with one endpoint being a leaf. Then it is easily seen that A has a corner in E_e^+ . This finishes the proof.

The following is a common generalization of corresponding results for cellular bipartite graphs [3] (being exactly rank 2 hypercellular graphs, which in turn are COMS [11]) and LOPs of rank 2 [9].

Theorem 5.11. Every rank 2 COM has a corner peeling.

Proof. Notice that a rank 2 COM is pure if and only if it is 2 connected. Consider the blocks of 2-connectedness of a rank 2 COM G. Then a block corresponding to a leaf in the tree structure of the block graph has 2 corners by Proposition 5.10. This implies that G has a corner. Proposition 5.2 together with the observation that G minus the corner has rank at most 2 yield a corner peeling.

Hypercellular graphs

Hypercellular graphs were introduced as a natural generalization of median graphs, i.e., skeleta of CAT(0) cube complexes in [11]. They are COMs with many nice properties one of them being that all their antipodal subgraphs are Cartesian products of even cycles and edges, called *cells*. See Figure 7 for an example.




Figure 7: A hypercellular graph.

More precisely, a partial cube G is *hypercellular* if all its antipodal subgraphs are cells and if three cells of rank k pairwise intersect in a cell of rank k-1 and altogether share a cell of rank k-2, then all three lie in a common cell, for all $2 \le k \le r(G)$. See Figure 8 for three rank 2 cells (cycles) pairwise intersecting in rank 1 cells (edges) and sharing a rank 0 cell (vertex) lying in a common rank 3 cell (prism). Since median graphs are realizable COMs, see [39], which is also conjectured for hypercellular graphs [11], they have corner peelings by Proposition 5.5. Here, we prove that hypercellular graphs have a corner peeling, which can be seen as a support for their realizability.

The following lemma determines the structure of corners in hypercellular graphs, since the corners of an edge K_2 and an even cycle C_{2n} are simply a vertex and a path P_{n-1} , respectively.

Lemma 5.12. Let $G = \Box_i A_i$ be the Cartesian product of even cycles and edges. Then the corners of G are precisely sets of the form $\Box_i D_i \subset G$, where D_i is a corner of A_i for every i.

Proof. Let $D = G_2 \setminus G_1$ be a corner of G, i.e. G_1, G_2 define an expansion in general position. Every subset of the form $A_1 \square \ldots \square A_{i-1} \square \{v\} \square A_{i+1} \square \ldots \square A_n$ is an antipodal subgraph, thus it is either completely in G_1 or in G_2 . We can use the latter to define an expansion in general position of A_i according to H_1, H_2 , by $v \in H_j$ if $A_1 \square \ldots \square A_{i-1} \square \{v\} \square A_{i+1} \square \ldots \square A_n \in G_j$, for $j \in \{1,2\}$. In fact each maximal antipodal subgraph A of A_i is either completely in H_1 or in H_2 (but not both) since $A_1 \square \ldots \square A_{i-1} \square A \square A_{i+1} \square \ldots \square A_n$




Figure 8: Any three cycles of a hypercellular graph that pairwise intersect in an edge and share a vertex lie in a common cell.

is a maximal antipodal of G either completely in G_1 or in G_2 (but not both). Moreover, since $G_1 = -G_2$ also $H_1 = -H_2$.

It remains to prove that H_1, H_2 are isometric. Let $v_1, v_2 \in H_1$. If there exists a unique shortest v_1, v_2 -path P in A_i , then pairs of vertices of $A_1 \square \ldots \square A_{i-1} \square \{v_1\} \square A_{i+1} \square \ldots \square A_n$ and $A_1 \square \ldots \square A_{i-1} \square \{v_2\} \square A_{i+1} \square \ldots \square A_n$ also have unique shortest paths. This implies that also $A_1 \square \ldots \square A_{i-1} \square P \square A_{i+1} \square \ldots \square A_n \in G_1$, since G_1 is isometric. Thus $P \in H_1$. The only option that v_1, v_2 do not have a unique v_1, v_2 -path is that A_i is a cycle of length 2k, for k > 2, and v_1, v_2 antipodal vertices in A_i . Then for each vertex u of A_i different form v_1, v_2 holds that $A_1 \square \ldots \square A_{i-1} \square \{u\} \square A_{i+1} \square \ldots \square A_n \in G_2$, thus $u \in H_2$. But then the neighbors of v_1 in A_i are in H_2 which by the previous case implies that $v_1 \in H_2$. Similarly $v_2 \in H_2$, which cannot be, since then $G_2 = G$. This proves that H_1, H_2 are isometric. In particular, if A_i is a cycle C_{2n} , then H_2, H_2 are paths P_{n+1} , and if A_i is an edge K_2 , then H_1, H_2 are vertices. Thus each $D_i = H_2 \setminus H_1$ is a corner.

By definition of the corners D_j , it holds $D = (G_2 \setminus G_1) \subset \square_i D_i$, i.e. $G \setminus \square_i D_i \subset G_1$. We prove that the equality holds. Let F be a set of Θ -classes that $\operatorname{cross} \square_i D_i$. Contracting all the Θ -classes in F gives a COM $\pi_F(G) = \square_i A_i'$ where $A_i' = \pi_F(A_i)$ is a 4-cycle if A_i is a cycle and $A_i' = A_i$ if A_i is an edge. Thus the rank of $\pi_F(G)$ is the same as the rank of G. Now if $(G_2 \setminus G_1) \subsetneq \square_i D_i$, it holds $\pi_F(G_1) = \pi_F(G)$, since $\pi_F(\square_i D_i)$ is a vertex. Defining the expansion of $\pi_F(G_1)$ according to $H_1 = \pi_F(G_1) = \pi_F(G)$, $H_2 = \pi_F(G_2) = \pi_F(G)$, gives a graph G that has a higher rank than $\pi_F(G)$ and G. But G can be obtain as a contraction of the graph G obtained by expanding G with respect to G and G. Since G, G define an expansion in general position G has the same rank as G. This is impossible.

We have proved that if G has a corner, then it is of the form $\Box_i D_i$. As mentioned, every OM has a corner. By symmetry, every set of vertices of the form $\Box_i D_i$ is a corner of G.

We shall need the following property about hypercellular graphs.

Lemma 5.13. Every zone graph $\zeta_f(G)$ of a hypercellular graph G is hypercellular.

Proof. Every zone graph of the Cartesian product of even cycles and edges is the Cartesian product of even cycles and edges, as it can easily be checked. Let $\zeta_f(G)$ be a zone graph of a hypercellular graph G. Then every cell of rank r in $\zeta_f(G)$ is an image of a cell of rank r+1 in G. Hence for every three rank r cells pairwise intersecting in rank r-1 cells and sharing a rank r-2 cell from $\zeta_f(G)$, there exist three rank r+1 cells pairwise intersecting in rank r cells and sharing a rank r-1 cell in G. Additionally the latter three cells lie in a common cell G. Then the image of G is a common cell of the three cells from G.

Let E_e be a Θ -class of a COM G. As usual, see e.g. [4, 11], we call the union of antipodal subgraphs crossed by E_e the *carrier* of E_e . The following is another generalization of the corresponding result for cellular graphs [3] and as mentioned above for median graphs.

Theorem 5.14. Every hypercellular graph G has a corner peeling.

Proof. We prove the assertion by induction on the size of G. The technical difficulty of the proof is that removing a corner in a hypercellular graph possibly produces a non-hypercellular graph. Hence we shall prove the above statement for the larger family \mathscr{F} of COMs defined by the following properties:

- (1) Every antipodal subgraph of $G \in \mathcal{F}$ is a cell.
- (2) Every carrier of $G \in \mathcal{F}$ is convex.
- (3) Every zone-graph of $G \in \mathcal{F}$ is in \mathcal{F} .

We first prove that hypercellular graph are a part of \mathscr{F} . By Lemma 5.13 only the first two properties must be checked. Now, (1) holds by definition of hypercellular graphs. Moreover, (2) follows from the fact that for any Θ -class E_e in a hypercellular graph the carrier of E_e is gated [11, Proposition 7], thus also convex.

We now prove that the graphs in \mathscr{F} have a corner peeling. Let $G \in \mathscr{F}$ and E_e an arbitrary Θ -class in G. Since the carrier of E_e is convex the so-called Convexity Lemma [26] implies that for any edge $g \in E_e^+$ with exactly one endpoint in the carrier its Θ -class E_g does not cross the carrier. Now if the union of cells crossed by E_e does not cover the whole E_e^+ , then for any edge g in E_e^+ with exactly one endpoint in the union, one of E_g^+ or E_g^- is completely in E_e^+ . Repeating this argument with E_g one can inductively find a Θ -class E_f with the property that the carrier of E_f completely covers E_f^+ , without loss of generality.

Let $\zeta_f(G)$ be the zone graph of G with respect to E_f , i.e., the edges of E_f are the vertices of $\zeta_f(G)$ and two such edges are connected if they lie in a common convex cycle. By (3) $\zeta_f(G)$ is in \mathscr{F} , thus by induction $\zeta_f(G)$ has a corner D_f . By definition there is a maximal antipodal subgraph A_f in $\zeta_f(G)$ such that the corner D_f is completely in A_f . Moreover, there exists a unique maximal antipodal subgraph A_f in G whose zone graph is A_f .

We lift the corner D_f from A_f to a corner D of A in the following way. If E_f in A corresponds to an edge factor K_2 , then A is simply $K_2 \square A_f$. In particular we can define $D = \{v\} \square D_f$ where v is a vertex of K_2 in E_f^+ . By Lemma 5.12, this is a corner of A. Since D_f lies only in the maximal antipodal graph A_f , D lies only in A.

Otherwise, assume E_f in A corresponds to a Θ -class of a factor C_{2k} (an even cycle). We can write $A = C_{2k} \square A'$. Then $A_f = K_2 \square A'$ with a corner $D_f = \{v\} \square D'$, where D' is a corner of A' by Lemma 5.12. We lift D_f to $D = P_{k-1} \square D'$. Here P_{k-1} is the path in C_{2k} consisting of the vertices in E_f^+ apart from the one lying on the edge not corresponding to v in the zone graph. As above since D_f lies only in the maximal antipodal graph A_f , D lies only in A.

We have proved that G has a corner D. To prove that it has a corner peeling it suffice to show that $G \setminus D$ is a graph in \mathscr{F} . Since removing a corner does not produce any new antipodal subgraph, all the antipodal subgraphs of $G \setminus D$ are cells, showing (1). The latter holds also for all the zone graphs of $G \setminus D$. To prove that (2) holds for $G \setminus D$ consider a Θ -class E_e of $G \setminus D$. By Lemma 5.2, $G \setminus D$ is an isometric subgraph of G, i.e. all the distances between vertices are the same in both graphs. Since the carrier of E_e in G is convex and removing a corner does not produce any new shortest path, the carrier of E_e is convex in $G \setminus D$. The same argument can be repeated in any zone graph of $G \setminus D$. This finishes the proof.

We have shown corner peelings for COMs of rank 2 and hypercellular graphs. A common generalization are Pasch graphs [10,11], which form a class of COMs [31] that exclude the examples from [3,9,53]:

Question 1. Does every Pasch graph have a corner peeling?

6 The minimum degree in antipodal partial cubes

Las Vergnas' conjecture can be seen as a statement about the minimum degree of an OM of given rank. Here we examine the relation of rank and minimum degree in general antipodal partial cubes.

6.1 Lower bounds

As stated in Section 1, if G is the tope graph of an OM, then $r(G) \le \delta(G)$, see [7, Exercise 4.4]. In general rank r antipodal partial cubes the minimum degree is not bounded from below by r. More precisely:

Proposition 6.1. For every $r \ge 4$ there is an antipodal partial cube of rank r and minimum degree 4. Moreover, there is an antipodal partial cube of rank 4 and minimum degree 3.

Proof. In [31] it is been shown that every partial cube G with n Θ -classes – thus embeddable in Q_n – is a convex subgraph of an antipodal partial cube A_G . Here, A_G is obtained by replacing in a Q_{n+3} one Q_n by G and its antipodal Q_n by -G. It is straight-forward to see that the minimum degree of A_G is $\delta(G) + 3$ and that the rank of A_G is at least n + 2. Indeed, for instance taking G as a path of length k > 1 we get $\delta(A_G) = 4$ and $r(A_G) = k + 2$.

Another construction is as follows. Take $Q_n^{--}(i)$, with $1 \le i < n$ and $n \ge 4$, to be the graph obtained from Q_n by removing a vertex ν , its antipode $-\nu$ and i neighbors of $-\nu$. Such a graph is affine and each antipode (in Q_n) of the removed neighbors of $-\nu$ is without the antipode in $Q_n^{--}(i)$, is of degree n-1 and of rank n-1. Then construct the antipodal graph taking two antipodal copies of it. Such graph will have minimum degree n-1 and rank n. For n=4 this gives the second part of the result.

On the other hand, it is shown in [44] that if an antipodal partial cube G has $\delta(G) \leq 2$, then $r(G) = \delta(G)$. This implies that if an antipodal partial cube G has $r(G) \leq 3$, then $r(G) \leq \delta(G)$. In relation to a question about cubic non-planar partial cubes we ask the following:

Question 2. Are there antipodal partial cubes with minimum degree 3 and arbitrary rank?

Indeed, since planar antipodal partial cubes are tope graphs of OMs of rank 3, see [20], any example for the above question has to be a non-planar antipodal partial cube of minimum degree 3. It has been wondered whether the only non-planar cubic partial cube is the (antipodal) Desargues graph [29], see the left of Figure 2. To our knowledge even the restriction to antipodal partial cubes remains open. For transitive cubic partial cubes it is known that the Desargues graphs is the only non-planar one, see [38]. On the other hand, it is open whether there are infinitely many non-planar partial cubes of minimum degree 3.

6.2 Upper bounds

Bounding the minimum degree in a partial cubes G from above by its rank is a generalization of Las Vergnas conjecture. As discussed in previous sections Las Vergnas conjecture is proved for OMs of rank at most 3. In fact tope graphs of OMs of rank 3 are even Θ -Las Vergnas, by Theorem 4.5 and the fact that they are Euclidean. We show that this property extends to general antipodal partial cubes of rank 3.

For this approach we introduce a couple of natural notions from [31]. A partial cube G is called *affine* if it is a halfspace E_e^+ of an antipodal partial cube. The *antipodes* A(G) of an affine partial cube are those $u \in G$ such that there is $-u \in G$ such that the *interval*

```
[u, -u] = \{v \in G \mid \text{there is a shortest path from } u \text{ to } -u \text{ through } v\}
```

coincides with G. The antipodes of G are exactly the vertices of E_e^+ incident to E_e when G is viewed as subgraph of G'. We need a auxiliary statement about the rank of affine partial cubes.

Lemma 6.2. If an affine partial cube G is a halfspace of an antipodal partial cube G' of rank r, then G has rank at most r-1.

Proof. Suppose there is a sequence of contractions from G to Q_k . Then the same sequence of contraction in G' yields a minor H with Q_k as a halfspace. Since H is antipodal, $H = Q_{k+1}$.

Lemma 6.3. Every affine partial cube of rank at most 2 has a vertex of degree at most 2 among its antipodes.

Proof. Since antipodal partial cubes of rank 2 are even cycles, it only remains to consider non-antipodal affine partial cubes. If G is an affine partial cube of rank 2, then by Lemma 6.2 it is a halfspace of an antipodal partial cube G' of rank 3. By [21, Theorem 7] since G' is not a cycle its minimum degree is at least 3. Hence if in G there are vertices of degree 2, then they are among its antipodes.

To show that G has vertices of degree 2, we use that by [12, Section 6.1], that every rank 2 partial cube can be augmented by adding vertices of degree at least 3 to a COM G'' of rank 2.

Finally, by Proposition 5.10 the graph G'' has a corner ν , which in particular is a vertex of degree at most 2. By the above ν must already have been present in G. This concludes the proof.

Proposition 6.4. Let G be an antipodal partial cube of rank 3 and E_e a Θ -class. There is a degree 3 vertex incident to E_e .

Proof. Suppose that the claim is false. Let G be a counterexample and E_e a Θ -class, such that all vertices incident to E_e have degree at least 4. Consider the contraction $G' = \pi_e(G)$ of G and let G'_2, G'_1 be the antipodal expansion of G' leading back to G. Since their preimage under π_e has degree at least 4, all vertices in $G'_1 \cap G'_2$ have degree at least 3. But $G'_1 \cap G'_2$ are the antipodes of the affine partial cube G'_1 . Moreover, G'_1 is of rank 2 by Lemma 6.2. Thus, we have a contradiction with Lemma 6.3.

While we have already used several times, that even OMs of rank 4 are not Θ -Las Vergnas, surprisingly enough Las Vergnas' conjecture could still hold for general antipodal partial cubes. We have verified it computationally up to isometric dimension 7. See Table 2 for the numbers.

n	2	3	4	5	6	7	8
antipodal	1	2	4	13	115	42257	?
OM	1	2	4	9	35	381	192449

Table 2: Numbers of antipodal partial cubes and OMs of low isometric dimension. The latter can also be retrieved from http://www.om.math.ethz.ch/.

Since already on isometric dimension 6 there are 13488837 partial cubes, instead of filtering those of isometric dimension 7 by antipodality, we filtered those of isometric dimension 6 by affinity. There are 268615 of them. We thus could create all antipodal partial cubes of dimension 7 and count them and verify Las Vergnas' conjecture also for this set. We extend the prolific Las Vergnas' conjecture to a much wider class.

Question 3. Does every antipodal partial cube of rank r have minimum degree at most r?

7 Conclusions and future work

We have shown that Mandel OMs have the Θ -Las Vergnas property, therefore disproving Mandel's conjecture. Finally, Las Vergnas' conjecture remains open and one of the most challenging open problems in OM theory. After computer experiments and a proof for rank 3, we dared to extend this question to general antipodal partial cubes, see Question 3. Another strengthening of Las Vergnas' conjecture is the conjecture of Cordovil-Las Vergnas. We have verified it by computer for small examples and it holds for low rank in general. However, here we suspect the existence of a counter example at least in the setting of $\overline{\mathscr{G}}^{n,r}$.

Our second main contribution is the introduction of corner peelings for COMs and the proof of their existence in the realizable, rank 2, and hypercellular cases. A class that is a common generalization of the latter two is the class \mathcal{S}_4 of Pasch graphs. Do these graphs admit corner peelings? See Question 1.

Let us close with two future directions of research that appear natural in the context of the objects discussed in this paper.

7.1 Shellability

In the context of an OM or AOM, a shelling is a special linear ordering of the vertices of its tope graph that yields a *recursive coatom ordering* of the full face lattice. See [7] for the definitions. It thus, is natural to compare corner peelings and shellings.

It is known that AOMs and OMs are shellable, see [7]. Shellability is defined for pure regular complexes thus the question of existence of shellings can be asked for all such COMs. However, the pure COM consisting of two 4-cycles joined in a vertex is not shellable.

Question 4. Which COMs are shellable?

A necessary condition might be r-connectedness, if a COM has rank r. Another useful hint might be the fact that an amalgamation procedure for COMs described in [4] is similar to the notion of constructibility, which is a weakening of shellability, see [22].

Corner peelings of LOPs are related to extendable shellability of the octahedron, see [9, 53]. While OMs have corners, AOMs do not always have corners, as shown by the example of Figure 1. Hence, shellability does not imply the existence of a corner or a corner peeling. However, there still might be a connection:

Question 5. If a shellable COM G has a corner peeling, can one find a shelling sequence that is a refinement of a corner peeling sequence of G?

7.2 Murty's conjecture

An important open problem in OMs is a generalization of the *Sylvester-Gallai Theorem*, i.e., for every set of points in the plane that does not lie on a single line there is a line, that contains only two points.

The corresponding conjecture in OMs can be found in Mandel's thesis [37], where it is attributed to Murty. In terms of OMs it reads:

Conjecture 4 (Murty). Every OM of rank r contains a convex subgraph that is the Cartesian product of an edge and an antipodal graph of rank r-2.

The realizable case of Murty's conjecture is shown by [52] and more generally holds for *Mandel* OMs [37]. Indeed, we suspect that along our strengthening of Mandel's theorem (Theorem 4.5) a Θ -version of Mandel's results can be proved:

Conjecture 5. Every Θ -class in a Mandel OM of rank r is incident to a vertex of an antipodal graph that is the Cartesian product of an edge and an antipodal graph of rank r-2.

On the other hand it would be interesting to find OMs, that do not have this strengthened property. Still, Murty's conjecture in general seems out of reach. We propose a reasonable weaker statement to attack:

Question 6. Does every OM of rank r contain a convex $Q_{\lceil \frac{r}{2} \rceil}$.

8 Acknowledgments

We thank the referees for helpful comments and Arnaldo Mandel for providing us with a copy of his PhD thesis. This work was supported by the Slovenian Research Agency (research core funding No. P1-0297 and projects J1-9109, N1-0095, J1-1693). Tilen Marc wishes to express his gratitude to Institut français de Slovénie for supporting a visit in Marseille that started the present research. Kolja Knauer was furthermore supported by the ANR project DISTANCIA: ANR-17-CE40-0015 and by the Spanish *Ministerio de Economía, Industria y Competitividad* through grant RYC-2017-22701.

References

- [1] M. Albenque and K. Knauer, Convexity in partial cubes: the hull number, Discrete Math., 339 (2016), pp. 866–876.
- [2] E. Babson, L. Finschi, and K. Fukuda, Cocircuit graphs and efficient orientation reconstruction in oriented matroids, Eur. J. Combin., 22 (2001), pp. 587–600.
- [3] H.-J. BANDELT AND V. CHEPOI, Cellular bipartite graphs., Eur. J. Comb., 17 (1996), pp. 121-134.
- [4] H.-J. BANDELT, V. CHEPOI, AND K. KNAUER, COMs: complexes of oriented matroids., J. Comb. Theory Ser. A, 156 (2018), pp. 195–237.
- [5] A. BAUM AND Y. ZHU, The axiomatization of affine oriented matroids reassessed., J. Geom., 109 (2018).
- [6] A. BJÖRNER, P. H. EDELMAN, AND G. M. ZIEGLER, *Hyperplane arrangements with a lattice of regions.*, Discrete Comput. Geom., 5 (1990), pp. 263–288.
- [7] A. BJÖRNER, M. LAS VERGNAS, B. STURMFELS, N. WHITE, AND G. M. ZIEGLER, *Oriented matroids*, vol. 46 of Encyclopedia of Mathematics and its Applications, Cambridge University Press, Cambridge, second ed., 1999.
- [8] J. Bokowski and H. Rohlfs, On a mutation problem for oriented matroids., Eur. J. Comb., 22 (2001), pp. 617-626.
- [9] J. CHALOPIN, V. CHEPOI, S. MORAN, AND M. K. WARMUTH, Unlabeled Sample Compression Schemes and Corner Peelings for Ample and Maximum Classes, in 46th International Colloquium on Automata, Languages, and Programming (ICALP 2019), vol. 132, 2019, pp. 34:1–34:15.
- [10] V. Chepoi, Separation of two convex sets in convexity structures., J. Geom., 50 (1994), pp. 30-51.
- [11] V. CHEPOI, K. KNAUER, AND T. MARC, Hypercellular graphs: Partial cubes without Q_3^- as partial cube minor, Discrete Math., 343 (2020), p. 111678.
- [12] V. Chepoi, K. Knauer, and M. Philibert, *Two-dimensional partial cubes*, Electron. J. Comb., 27 (2020), pp. research paper p3.29, 40.
- [13] V. Chepoi, K. Knauer, and M. Philibert, Ample completions of oriented matroids and complexes of uniform oriented matroids, SIAM J. Discrete Math., 36 (2022), pp. 509–535.
- [14] R. CORDOVIL, K. FUKUDA, AND A. G. DE OLIVEIRA, On the cocircuit graph of an oriented matroid, Discrete Comput. Geom., 24 (2000), pp. 257–266.
- [15] I. P. F. DA SILVA, Axioms for maximal vectors of an oriented matroid: a combinatorial characterization of the regions determined by an arrangement of pseudohyperplanes, Eur. J. Combin., 16 (1995), pp. 125–145.
- [16] G. DORPALEN-BARRY, J. S. KIM, AND V. REINER, Whitney Numbers for Poset Cones., Order, (2021).
- [17] S. Felsner, R. Gómez, K. Knauer, J. J. Montellano-Ballesteros, and R. Strausz, *Cubic time recognition of cocircuit graphs of uniform oriented matroids.*, Eur. J. Comb., 32 (2011), pp. 60–66.
- [18] J. FOLKMAN AND J. LAWRENCE, Oriented matroids, J. Combinatorial Theory Ser. B, 25 (1978), pp. 199–236.
- [19] K. Fukuda, Oriented matroid programming, 1982. PhD Thesis, University of Waterloo.
- [20] K. FUKUDA AND K. HANDA, Antipodal graphs and oriented matroids, Discrete Math., 111 (1993), pp. 245–256.
- [21] F. GLIVJAK, A. KOTZIG, AND J. PLESNÍK, Remark on the graphs with a central symmetry, Monatsh. Math., 74 (1970), pp. 302–307.
- [22] M. HACHIMORI, Combinatorics of constructible complexes, 2000. PhD thesis, University of Tokyo.
- [23] R. HAMMACK, W. IMRICH, AND S. KLAVŽAR, *Handbook of product graphs*, Discrete Mathematics and its Applications (Boca Raton), CRC Press, Boca Raton, FL, 2011.
- [24] K. HANDA, A characterization of oriented matroids in terms of topes, Eur. J. Combin., 11 (1990), pp. 41-45.
- [25] W. HOCHSTÄTTLER AND V. WELKER, The Varchenko determinant for oriented matroids., Math. Z., 293 (2019), pp. 1415–1430.
- [26] W. IMRICH AND S. KLAVŽAR, A convexity lemma and expansion procedures for bipartite graphs, Eur. J. Comb., 19 (1998), pp. 677–685.
- [27] V. ITSKOV, A. KUNIN, AND Z. ROSEN, *Hyperplane neural codes and the polar complex*, in Topological data analysis. Proceedings of the Abel symposium 2018, Geiranger, Norway, June 4–8, 2018, Cham: Springer, 2020, pp. 343–369.
- [28] T. JUNTTILA AND P. KASKI, Engineering an efficient canonical labeling tool for large and sparse graphs, in 2007 Proceedings of the Ninth Workshop on Algorithm Engineering and Experiments (ALENEX), SIAM, 2007, pp. 135–149.
- [29] S. KLAVZAR, Hunting for cubic partial cubes, Convexity in Discrete Structures, 5 (2008), pp. 87–95.
- [30] S. KLAVŽAR AND S. SHPECTOROV, Convex excess in partial cubes, J. Graph Theory, 69 (2012), pp. 356-369.
- [31] K. KNAUER AND T. MARC, On tope graphs of complexes of oriented matroids, Discrete Comput. Geom., (2019), pp. 377-417.

- [32] K. KNAUER, J. J. MONTELLANO-BALLESTEROS, AND R. STRAUSZ, A graph-theoretical axiomatization of oriented matroids, Eur. J. Combin., 35 (2014), pp. 388–391.
- [33] A. Kunin, C. Lienkaemper, and Z. Rosen, Oriented matroids and combinatorial neural codes, 2020.
- [34] M. LAS VERGNAS, Convexity in oriented matroids., J. Comb. Theory Ser. B, 29 (1980), pp. 231–243.
- [35] J. LAWRENCE, Lopsided sets and orthant-intersection by convex sets, Pacific J. Math., 104 (1983), pp. 155-173.
- [36] F. LEVI, Die Teilung der projektiven Ebene durch Gerade oder Pseudogerade, Ber. Math.-Phys. Kl. Sächs. Akad. Wiss, 78 (1926), pp. 256–267.
- [37] A. MANDEL, Topology of oriented matroids, 1982. PhD Thesis, University of Waterloo.
- [38] T. MARC, Classification of vertex-transitive cubic partial cubes., J. Graph Theory, 86 (2017), pp. 406-421.
- [39] T. MARC, Cycling in hypercubes, PhD thesis, Univ. of Ljubljana, Faculty of Mathematics and Physics, 2018.
- [40] S. MARGOLIS, F. V. SALIOLA, AND B. STEINBERG, Cell complexes, poset topology and the representation theory of algebras arising in algebraic combinatorics and discrete geometry, vol. 1345 of Mem. Am. Math. Soc., Providence, RI: American Mathematical Society (AMS), 2021.
- [41] J. J. Montellano-Ballesteros and R. Strausz, A characterization of cocircuit graphs of uniform oriented matroids, J. Combin. Theory Ser. B, 96 (2006), pp. 445–454.
- [42] S. OVCHINNIKOV, Graphs and cubes., Berlin: Springer, 2011.
- [43] A. PADROL AND E. PHILIPPE, Sweeps, polytopes, oriented matroids, and allowable graphs of permutations, 2021.
- [44] N. POLAT, On some properties of antipodal partial cubes, Discuss. Math., Graph Theory, (2018).
- [45] N. POLAT, On some characterizations of antipodal partial cubes., Discuss. Math., Graph Theory, 39 (2019), pp. 439-453.
- [46] H. RANDRIAMARO, The Varchenko determinant for apartments, Result. Math., 75 (2020), p. 17. Id/No 86.
- [47] J. RICHTER-GEBERT, Oriented matroids with few mutations., Discrete Comput. Geom., 10 (1993), pp. 251–269.
- [48] G. RINGEL, Teilungen der Ebenen durch Geraden oder topologische Geraden., Math. Z., 64 (1956), pp. 79–102.
- [49] G. RINGEL, Über Geraden in allgemeiner Lage., Elem. Math., 12 (1957), pp. 75-82.
- [50] J.-P. ROUDNEFF AND B. STURMFELS, Simplicial cells in arrangements and mutations of oriented matroids., Geom. Dedicata, 27 (1988), pp. 153–170.
- [51] F. Santos, Triangulations of oriented matroids., vol. 741, American Mathematical Society (AMS), 2002.
- [52] R. W. SHANNON, Simplicial cells in arrangements of hyperplanes., Geom. Dedicata, 8 (1979), pp. 179-187.
- [53] H. Tracy Hall, Counterexamples in Discrete Geometry, PhD Thesis, University Of California, Berkeley, 2004.
- [54] V. N. VAPNIK AND A. Y. CHERVONENKIS, On the uniform convergence of relative frequencies of events to their probabilities., in Measures of complexity, Springer, 2015, pp. 11–30.