Graph coloring with no large monochromatic components

NATHAN LINIAL*

School of Computer Science and Engineering Hebrew University Jerusalem, Israel

Or Sheffet[†]

School of Computer Science and Engineering Hebrew University Jerusalem, Israel

Jiří Matoušek

Department of Applied Mathematics and Institute of Theoretical Computer Science (ITI) Charles University, Malostranské nám. 25 118 00 Praha 1, Czech Republic

Gábor Tardos[‡]

School of Computing Science Simon Fraser University, Burnaby, BC, Canada and Rényi Institute, Budapest, Hungary

Abstract

For a graph G and an integer t we let $mcc_t(G)$ be the smallest m such that there exists a coloring of the vertices of G by t colors with no monochromatic connected subgraph having more than m vertices. Let \mathcal{F} be any nontrivial minor-closed family of graphs. We show that $mcc_2(G) = O(n^{2/3})$ for any n-vertex graph $G \in \mathcal{F}$. This bound is asymptotically optimal and it is attained for planar graphs. More generally, for every such \mathcal{F} , and every fixed t we show that $mcc_t(G) = O(n^{2/(t+1)})$. On the other hand we have examples of graphs G with no K_{t+3} minor and with $mcc_t(G) = \Omega(n^{2/(2t-1)})$.

It is also interesting to consider graphs of bounded degrees. Haxell, Szabó, and Tardos proved $mcc_2(G) \leq 20000$ for every graph G of maximum degree 5. We show that there are n-vertex 7-regular graphs G with $mcc_2(G) = \Omega(n)$, and more sharply, for every $\varepsilon > 0$ there exists $c_{\varepsilon} > 0$ and n-vertex graphs of maximum degree 7, average degree at most $6 + \varepsilon$ for all subgraphs, and with $mcc_2(G) \geq c_{\varepsilon}n$. For 6-regular graphs it is known only that the maximum order of magnitude of mcc_2 is between \sqrt{n} and n.

We also offer a Ramsey-theoretic perspective of the quantity $mcc_t(G)$.

1 Introduction

In the classical graph coloring problem we assign a color to each vertex so that no two vertices of the same color are adjacent. In other words, each monochromatic connected component must be a single vertex. In the problems that we study here, this requirement is relaxed and we only demand that monochromatic connected components should have small cardinality. Concretely, for a graph G and an integer t we define $mcc_t(G)$ as

^{*}Supported by the Israel Science Foundation.

[†]This paper is partially based on this author's undergraduate Amirim honours program project.

[‡]Supported by NSERC grant 611470 and the Hungarian Foundation for Scientific Research Grant (OTKA) Nos. T037846, T046234, AT048826 and NK62321.

the smallest integer m such that the vertices of G can be t-colored so that no monochromatic connected component has cardinality exceeding m. In particular, $mcc_t(G) = 1$ iff G can be properly t-colored.

Here are some technicalities before we survey earlier work on this subject. When we consider a graph G, n always denotes the number of vertices. For a set $S \subseteq V(G)$, we let G[S] denote the subgraph of G induced by S. We use the standard asymptotic language and conveniently ignore integrality issues in our computations. Some of the examples we consider are line graphs. We recall that the line graph of a graph H is defined as $L(H) = (E(H), \{\{e, e'\} : e \cap e' \neq \emptyset\})$. Clearly, $mcc_t(L(H))$ is the smallest m such that there is a t-coloring of the edges of H so that every monochromatic connected subgraph of H has $\leq m$ edges.

The earliest reference investigating the parameter mcc_t we are aware of is Kleinberg, Motwani, Raghavan, and Venkatasubramanian [KMR+97], where the question was motivated by a problem in computer science concerning dynamically evolving databases. Among others, the authors prove that mcc_3 is unbounded for planar graphs, and that there is a constant $\varepsilon > 0$ such that for all (sufficiently large) d, there are d-regular graphs G with $mcc_{\varepsilon\sqrt{d}}(G) = \Omega(n)$.

Apparently independently, the possibility of bounding $mcc_t(G)$ by a constant for graphs of bounded degree has been investigated by graph theorists. The results concern mainly the case t=2. It is easy to see that $mcc_2(G) \leq 2$ for any graph G of maximum degree 3. Alon, Ding, Oporowski, and Vertigan [ADO+03] proved that $mcc_2(G) \leq 57$ for every graph G of maximum degree 4. Haxell, Szabó, and Tardos [HST03] improved this to $mcc_2(G) \leq 6$ and proved that $mcc_2(G) \leq 20000$ for every graph G of maximum degree 5. On the other hand, Alon et al. [ADO+03] constructed 6-regular graphs G with $mcc_2(G)$ arbitrarily large. For graphs G of maximum degree 3 it was also shown in [BS05] that they admit two-coloring where one color induces an independent set, while the other color induces components of size at most 189. Earlier work on this subject [DOS+96], [JW96] mainly focused on more specific questions concerning line graphs of 3-regular graphs. These investigations culminated in [Tho99] showing that the edges of every 3-regular graph can be 2-colored so that each monochromatic component is a path of length at most 5.

We should also mention that there is a fairly rich literature that deals with the notion of t-vertex coloring where each monochromatic connected component has a small diameter, see e.g. [LS93]. This line of research originated in the field of distributed computing.

Here are the main results of the present paper:

Theorem 1.1 For every planar graph G we have $mcc_2(G) = O(n^{2/3})$. More generally, for every nontrivial minor-closed family of graphs \mathcal{F} there exists a constant $C = C_{\mathcal{F}}$ such that if G belongs to \mathcal{F} , then $mcc_2(G) \leq Cn^{2/3}$. This bound is tight even for planar graphs.

For every fixed integer $t \geq 2$ and every $G \in \mathcal{F}$ as above, there holds $mcc_t(G) = O(n^{2/(t+1)})$. On the other hand, for every t there exist graphs with no K_{t+3} minor and with $mcc_t(G) = \Omega(n^{2/(2t-1)})$.

If the tree-width of G is bounded by a constant (equivalently, if G excludes a fixed planar minor), then $mcc_t(G) = O(n^{1/t})$. This bound is asymptotically optimal for every fixed t.

Theorem 1.2 For every $\varepsilon > 0$ there exists a constant $c_{\varepsilon} > 0$ and arbitrarily large graphs G so that

- Every vertex in G has degree at most 7,
- Every subgraph of G has average degree at most $6 + \varepsilon$,
- $mcc_2(G) \ge c_{\varepsilon} n$.

As already mentioned, the questions we consider here have independently originated in computer science and in graph theory. Graph coloring is, of course, one of the most fascinating parts of graph theory. Due to its great significance and the famous open questions about it, many different variations on the basic theme are being investigated (see, e.g., [JT95]) and the present problems can be viewed as part of this ongoing research effort.

What is less obvious is the connection between the graph invariants we consider here and Ramsey Theory. The usual perception is that Ramsey-type theorems express the fact that large systems necessarily contain "highly regular islands". We suggest that many Ramsey-type results can be viewed as "sum theorems". Specifically, let \mathcal{G} be a class of graphs closed under taking subgraphs. Given a graph G, we ask for the smallest number of members in \mathcal{G} whose union is G. Thus when \mathcal{G} consists of all graphs without a k-clique we encounter the classical Ramsey problem. When it is the class of all graphs not containing a given subgraph we recover the so-called Graph Ramsey Theory. Finally if \mathcal{G} contains all graphs in which each connected component has small cardinality we arrive at our present problem. We believe that this perspective deserves further research. Needless to say, this concept extends beyond graphs and unions. Other mathematical objects and other appropriate operations can be considered. Given an object G and a class \mathcal{G} , one seeks the most economical way of expressing G as a "sum" of members in \mathcal{G} .

There are certain classes of graphs for which the study of coloring with small monochromatic connected components is particularly interesting. Let D_m^d denote the d-dimensional grid with all diagonals; that is, D_m^d is the graph with vertex set $\{1,2,\ldots,m\}^d$ where the two vertices u,v are adjacent if $||u-v||_{\infty}=\max_i|u_i-v_i|\leq 1$. The study of mcc for this graph and its relatives leads to very interesting problems which bring together combinatorics geometry and topology. The d-dimensional version of the well-known HEX lemma (see Gale [G79] and Linial and Saks [LS93]) implies that $mcc_d(D_m^d)\geq m$ (and a simple coloring shows an O(m) upper bound for every fixed d). More colors allow for constant size components, i.e., we have $mcc_{d+1}(D_m^d)=O(1)$ for every fixed d. For two colors, Matoušek and Přívětivý [MP07] proved $mcc_2(D_m^d)\geq m^{d-1}-d^2m^{d-2}$, which nearly matches the obvious upper bound (layer-by-layer 2-coloring) of m^{d-1} . The behavior of $mcc_t(D_m^d)$ for $3\leq t< d$ is still unknown and remains an intriguing open problem.

2 Excluded minors, separators, and coloring

In this section we prove all the upper bounds in Theorem 1.1.

A subset $C \subseteq V$ of the vertex set of a graph G = (V, E) is called a *separator* if no component of $G[V \setminus C]$ has more than $\frac{2}{3}|V|$ vertices (the choice of the constant $\frac{2}{3}$




Figure 1: The recursive algorithm for 2-coloring via separators.

is somewhat arbitrary). By the well-known planar separator theorem of Lipton and Tarjan [LT79], every planar graph G has a separator with at most $O(\sqrt{n})$ vertices. More generally, for every h-vertex graph H, every G containing no minor isomorphic to H has a separator with at most $h^{3/2}\sqrt{n}$ vertices [AST90]. We will also need that graphs of bounded tree width have bounded size separators. In particular, a graph of tree width w has a separator of size w+1 as stated, e.g., in Remark 2 of [R92]. (We haven't found the proof of this very simple fact anywhere, but a constant size separator follows from the combination of the analogous result on branch width in Lemma 3.1 of [RS95] and the connection between branch width and tree width as stated in [RS91].)

In view of these results, the upper bounds in Theorem 1.1 all follow from the following proposition.

Proposition 2.1 Let \mathcal{G} be a class of graphs closed under taking induced subgraphs such that every $G \in \mathcal{G}$ has a separator with at most Kn^{γ} vertices, where K and $\gamma \in [0,1)$ are constants depending only on \mathcal{G} . Then for every $G \in \mathcal{G}$ we have

$$mcc_2(G) = O(n^{1/(2-\gamma)}),$$

and more generally,

$$mcc_t(G) = O(n^{1/(t-(t-1)\gamma)}),$$

where the hidden constant of proportionality depends on K, γ , and (in the second case) on t.

Proof. First we deal with the special case t=2. Given an n-vertex G, let $n_0:=\lfloor n^{1/(2-\gamma)}\rfloor$ be a threshold parameter. We present a simple algorithm producing a 2-coloring of V(G). The algorithm maintains a list \mathcal{L} of induced subgraphs of G, which is initialized to $\mathcal{L}:=\{G\}$, and a set S of vertices, initialized to \emptyset . While \mathcal{L} contains at least one graph with more than n_0 vertices, we select one such graph $G_i \in \mathcal{L}$ arbitrarily, we remove it from \mathcal{L} , we find a separator C_i of G_i of size at most $K|V(G_i)|^{\gamma}$, we set $S:=S\cup C_i$, and we add all of the components of $G_i[V(G_i]\setminus C_i]$ to \mathcal{L} . The algorithm ends when \mathcal{L} contains only graphs of size at most n_0 ; at this moment, we color the vertices of S blue and all remaining vertices (i.e., the vertices of all graphs in \mathcal{L}) red and we finish. The algorithm is illustrated in Fig. 1.

By construction, no red component in this coloring has more than n_0 vertices, and it suffices to show that $|S| = O(n_0)$ at the end of the algorithm (S can form a single




Figure 2: The lower bound construction for planar graphs, drawn for k=3.

blue component at worst). We use the following charging scheme: Whenever we color a separator C_i in a graph G_i blue, we let each vertex of G_i pay $K|V(G_i)|^{\gamma-1}$ units. Since $|C_i| \leq K|V(G_i)|^{\gamma}$, the total paid by all vertices of G_i at this step is at least $|C_i|$. Now we consider an individual vertex $v \in V(G)$ and we bound the total charge paid by it throughout the whole algorithm. There may be several successive charges, since v first pays as a vertex of G, and then possibly as vertex of some of the G_i . Let x_j be the amount paid when v is charged the jth time, $j = 1, 2, \ldots, q$. We observe that $x_q \leq K n_0^{\gamma-1}$ (since only graphs G_i with at least n_0 vertices get partitioned), and that $x_{j-1} \leq (2/3)^{1-\gamma} x_j$ for all j, since the component of G_i containing v always has at most two-third of the vertices of G_i . Hence the x_j are bounded from above by a decreasing geometric series, and thus the total charge paid by v is $O(n_0^{\gamma-1})$. So $|S| \leq O(n n_0^{\gamma-1}) = O(n^{1/(2-\gamma)})$.

Next, we consider the case of t>2 colors. We proceed by induction on t, assuming that for every n-vertex graph $G\in\mathcal{G}$ we can construct a coloring with t-1 colors witnessing $mcc_{t-1}(G)=O(n^{1/(t-1-(t-2)\gamma)})$.

For the induction step, we consider a $G \in \mathcal{G}$ and we apply to it the algorithm above with the following modifications: This time we let the threshold be $n_0 := \lfloor n^{1/(t-(t-1)\gamma)} \rfloor$, and at the end, we color G[S] by t-1 colors using the inductive assumption, while the vertices not belonging to S get color t.

By the above analysis, we have $|S| = O(nn_0^{\gamma-1})$, and by induction, the monochromatic components in colors 1 through t-1 have size at most $O(|S|^{1/(t-1-(t-2)\gamma)}) = O(n^{1/(t-(t-1)\gamma)})$. The components in color t have size at most n_0 by construction. This finishes the proof of Proposition 2.1.

3 Lower bounds for planar graphs and for excluded minors

To prove that the bound in Theorem 1.1 for $mcc_2(G)$ is tight for planar graphs G, we construct planar graphs G with $mcc_2(G) = \Omega(n^{2/3})$. For every integer k we construct $G = G_k$ on $n = 2k^3 + 1$ vertices, as indicated in Fig. 2. This G is constructed from a k by k^2 grid Z, a path P of k^3 vertices, and an extra vertex x. For $1 \le i \le k^2$ we denote the ith column of Z by C_i , and we let v_i be the top vertex of C_i . We break the path P

into consecutive intervals I_1, \ldots, I_{k^2} of k vertices each and we connect the vertices of I_i with v_i . We let $R_i = C_i \cup I_i$ and call this a rib of G. We connect x with all vertices in P. Finally, we add diagonals to all quadrilateral faces of the planar graph constructed so far, so that it becomes a triangulated polygon, that is, a planar graph where all faces except possibly for one are triangles.

Our main tool is the following lemma about triangulated polygons (a very similar lemma appears in [MP07]). For a set S of vertices in a graph we denote by ∂S the set of vertices that are not in S but have a neighbor in S.

Lemma 3.1 Let G be a triangulated polygon and let $S \subseteq V(G)$ be such that G[S] is connected. Suppose that two vertices $u, v \in \partial S$ are not separated by S in G. Then there is a path between u and v that is entirely included in ∂S .

Proof. This is a simple consequence of the planar HEX lemma. Since G[S] is connected and $u, v \in \partial S$, there is an u-v path P_1 with all internal vertices in S. Since S doesn't separate u from v, there is another u-v path P_2 that avoids S. We consider subgraph of H consisting of the cycle $P_1 \cup P_2$ plus the part of G that triangulates the interior of this cycle (assuming that the single non-triangular face of G is the outer face). We add two new vertices z and t and we connect z to all vertices of P_1 and t to all vertices of P_2 . The resulting graph H is a triangulation of the cycle uzvt.

We color blue all vertices in ∂S including u and v, and we color red all other vertices of H including z and t. By the HEX lemma (as stated, e.g., in [MN98]) we have either a blue u-v path, which is what we want, or a red z-t path Q. We want to exclude the latter possibility. Let us imagine that we follow the red path Q from t to z and we watch the distance to S in H. Since t is adjacent only to P_2 , whose vertices are not in S, initially at t this distance is at least z. On the other hand, since the red vertices connected to z are inner vertices of P_1 and thus in S, the penultimate vertex in Q is in S. Consequently, there is a vertex in Q at distance z from z, but such a vertex was colored blue—a contradiction.

We need the following consequence of Lemma 3.1:

Corollary 3.2 Consider a red-blue vertex coloring of the graph $G = G_k$, where x is red. Let S be the connected component of x in the red subgraph. If there is a connected component of $G \setminus S$ containing at least r S-free ribs (i.e., ribs with no vertex in S), then G has a blue connected subgraph with at least r k vertices.

Proof. If a rib R_i is S-free, then the k vertices in the interval I_i are contained in ∂S and are therefore blue. For any two S-free ribs R_i , R_j contained in the same connected component of $G \setminus S$, we choose vertices $u \in I_i$ and $v \in I_j$ (arbitrarily). The previous lemma now shows that u and v are connected by a blue path.

We can now show that any two-coloring of G has a monochromatic connected subgraph of at least $k^2/2$ vertices. As in the corollary, we assume x red and we let S be the connected component of x in the red subgraph. We may assume $|S| \leq k^2/2$, for otherwise, we have a large red component. Hence there are at least $k^2/2$ S-free ribs. We want to show that at least k/2 of them are in the same connected component of $G \setminus S$; then we will be done by the corollary.

Since $|S| < k^2$, at least one of the k rows of the grid Z contains fewer than k vertices from S. It follows that the S-free ribs live in at most k connected components of $G \setminus S$. So there must be at least k/2 of them in the same connected component as claimed. \square

For $m \geq 1$ and a graph G let cone(mG) be a graph constructed by taking m isomorphic and pairwise disjoint copies of G and connecting all of their vertices to an additional new vertex, called the apex.

Lemma 3.3 Let $t \ge 1$, let G be a graph, and let $m = mcc_t(G)$. Then $mcc_{t+1}(cone(mG)) \ge m$.

Proof. Let us consider a coloring of the vertices of cone(mG) with t+1 colors, and let us assume that the apex has color t+1. Clearly, all vertices of color t+1 form a connected subgraph, so if there are at least m of them we have our large monochromatic connected subgraph. Otherwise, one of the copies of G lacks color t+1 and the claim follows.

Notice that as we pass from G to cone(mG), the number of vertices grows (approximately) m times, but other parameters grow slowly: the tree width grows by at most one, the size of the largest clique minor grows by one, and if G is outerplanar, then cone(mG) is planar.

To prove the statement of Theorem 1.1 about the existence of K_{t+3} minor free graphs with high mcc_t , we simply take the planar graph G_k constructed at the beginning of this section and we apply the above lemma t-2 times with $m=k^2/2$. Then mcc_t of the resulting graph is at least $k^2/2$, and the number of vertices is $O(k^{2t-1})$.

To prove the similar statement about constant tree-width graphs, we need a different base graph: let F_k be the "fan" consisting of a k-vertex path and an additional vertex adjacent to all vertices of this path. Clearly, F_k is an outerplanar graph of tree width 2. A straightforward computation shows that $mcc_2(F_k) = \Theta(\sqrt{k})$ (this also appears in [ADO+03]). Applying the above lemma to F_k t-2 times with $m = mcc_2(F_k)$, we obtain a graph of tree width at most t on $O(k^{t/2})$ vertices with $mcc_t = \Omega(\sqrt{k})$. This finishes the proof of Theorem 1.1.

The case t=3 in the just finished proof yields n-vertex planar graphs with mcc_3 at least $\Omega(n^{1/3})$. These very graphs were used in [KMR+97] to show that mcc_3 is not bounded for planar graphs.

4 Edge expansion and degree $6 + \varepsilon$

In this section we prove Theorem 1.2. The graphs we construct are line graphs G = L(H). So the property that we need is that in every 2-coloring of E(H) there are monochromatic connected components containing a positive fraction of the edges of H. To this end, it suffices to show that small subgraphs of H have small average degrees, as the next observation shows:

Lemma 4.1 Let H be a graph with average degree \overline{d} . Suppose that every subgraph on p or fewer vertices in H has average degree strictly smaller than \overline{d}/t . Then $mcc_t(L(H)) \geq p$.

Proof. If $F \subseteq E(H)$ is the largest color class in a t-coloring of E(H), then the graph (V(H), F) has average degree at least \overline{d}/t . Consequently, some connected component of (V(H), F) has average degree at least \overline{d}/t . Such a connected component must have more than p vertices, and thus at least p edges.

In the proof of Theorem 1.2, we will use a suitable random graph (with bounded vertex degrees) for H. The basic idea is that random graphs are typically very good expanders. This means that every sufficiently small set $S \subset V(H)$ has many neighbors, hence most of the edges incident to S leave S, and consequently, the average degree of H[S] is small. More precisely, it turns out that if |S| is sufficiently small and if H[S] is connected, then this subgraph is nearly a tree in the sense that the average degree in H[S] is just a little bigger than 2. Here sufficiently small means that $|S| < \beta |V(H)|$, where $\beta > 0$ is a suitable small constant depending on how close we want to get to average degree 2.

A result about this almost-tree behavior of small sets in random regular graphs appears explicitly in [HLW06]. In particular, Theorem 4.16, part (1) in [HLW06] tells us that for every $d \geq 3$ and every $\delta > 0$ there exists $\beta = \beta(d, \delta) > 0$ such that almost every d-regular graph H on m vertices has average degree of H[S] at most $2 + \delta$ for all S with at most βm vertices. This, together with Lemma 4.1, immediately yields the following weaker analogue of Theorem 1.2: There exist arbitrarily large 8-regular graphs G with $mcc_2(G) = \Omega(n)$. Indeed, we choose H as a 5-regular graph on m vertices (thus, G = L(H) is 8-regular) satisfying the conclusion of the statement quoted above with $\delta = 0.4$. Then for $\beta = \beta(5, 0.4)$, every $S \subseteq V(H)$ with at most βm vertices induces a subgraph average degree at most 2.4, and hence Lemma 4.1 with t = 2 and $\overline{d} = 5$ shows $mcc_2(L(H)) \geq \beta m$.

In order to lower the maximum degree of G to 7 and the maximum average degree to $6 + \varepsilon$, we will use a random H where a small fraction of vertices have degree 5, all others have degree 4, and no two degree-5 vertices are connected.

The random graph model. It is easier to deal with random bipartite graphs. Most of the literature in this area deals with regular random graphs, but we need a suitable mixture of vertex degrees, and so we prescribe the degree individually for each vertex. That is, we have two disjoint sets A and B of vertices, |A| + |B| = m, and for every $v \in A \cup B$ we specify a number $d(v) \in \{1, 2, ..., D\}$, where D is a constant. These degrees and the sizes of A and B are related by the condition d(A) = d(B), where we use the notation $d(S) = \sum_{v \in S} d(v)$. Hence $|A| = c_A m$, $|B| = c_B m$ for constants c_A, c_B . To generate the random graph, every vertex v starts with d(v) "half-edges", and then the half-edges of all vertices in A are matched at random to the half-edges of all vertices in B (this is a configuration model of generating random bipartite graphs). We note that the resulting B may have multiple edges, but the line graph C(B) we are interested in is still a simple graph.

The following lemma speaks about number of *vertices* adjacent to S; the number of edges is then obtained as a simple consequence.

 $^{^{1}}$ It is well known that for random regular graphs of fixed degree, the configuration model yields a simple graph with probability bounded away from 0, and consequently, any property that holds almost surely in the configuration model also holds for almost all simple regular graphs of the given degree; see, e.g., [JLR00]. By slightly modifying the proof of this fact, we could also get a similar result for our model with mixed degrees, and hence have H simple.

Lemma 4.2 Let D be a fixed integer. Then for every $\delta > 0$ there exists $\beta = \beta(D, \delta) > 0$ such that if H is generated according to the above model (with an arbitrary choice of the d(v)'s), then with probability 1 - o(1) (as $m \to \infty$), every $S \subseteq A$ with $|S| \le \beta m$ has at least $d(S) - (1 + \delta)|S|$ neighbors in B.

Proof. Let us write $w(S) = d(S) - (1 + \delta)|S|$. Since every S has at least |S|/D neighbors, it suffices to consider only the S with $w(S) \ge |S|/D$. The calculation is a variation of that in [HLW06].

For sets $S \subseteq A$, $|S| = s \le \beta m$, and $W \subseteq B$, $|W| = \lfloor w(S) \rfloor$, let $X_{S,W}$ be the event "all neighbors of S lie in W." If no $X_{S,W}$ occurs, then H satisfies the conclusion of the lemma. We have

$$\Pr[X_{S,W}] = \frac{d(W)(d(W) - 1) \cdots (d(W) - d(S) + 1)}{d(B)(d(B) - 1) \cdots (d(B) - d(S) + 1)} \le \left(\frac{d(W)}{d(B)}\right)^{d(S)} \le \left(\frac{D^2 w(S)}{m}\right)^{d(S)}.$$

Thus for S fixed, the probability of $X_{S,W}$ occurring for some w is at most

$$\binom{|B|}{w(S)} \left(\frac{D^2 w(S)}{m}\right)^{d(S)}.$$

Estimating the binomial coefficient as $\binom{x}{y} \leq (ex/y)^y$ and using $s/D \leq w(S) \leq Ds$, this can be bounded by

$$\left(\frac{e|B|}{w(S)}\right)^{w(S)} \left(\frac{D^2 w(S)}{m}\right)^{d(S)} \le \left(\frac{e|B|}{s}\right)^{w(S)} \left(\frac{D^3 s}{m}\right)^{d(S)} \le C_1^s \left(\frac{s}{m}\right)^{(1+\delta)s}$$

 $(C_1 \text{ a constant independent of } \beta)$. Then the probability of any $X_{S,W}$ occurring at all is bounded by

$$\sum_{1 < s < \beta m} \binom{|A|}{s} C_1^s \left(\frac{s}{m}\right)^{(1+\delta)s} \leq \sum_s C_2^s \left(\frac{s}{m}\right)^{\delta s},$$

where C_2 is another constant. The term for s=1 is $O(m^{-\delta})=o(1)$, and the ratio of consecutive terms is at most $C_2\beta^{\delta}$, which can be made smaller than $\frac{1}{2}$, say, by fixing β small enough. Then the entire sum is o(1) as claimed.

Corollary 4.3 In the setting of Lemma 4.2, the following holds almost surely: The average degree of the subgraph of H induced by any set of at most βm vertices is at most $2+2\delta$.

Proof. Let us consider the subgraph of H induced by $S \cup T$, $S \subseteq A$, $T \subseteq B$, $|S \cup T| \leq \beta m$. By Lemma 4.2, we may assume that S has at least w(S) neighbors. Hence at least w(S) - |T| neighbors of S do not lie in T, and each such neighbor "consumes" at least one edge among the d(S) edges incident to S. Thus the number of edges in $H[S \cup T]$ is at most $d(S) - w(S) + |T| = (1+\delta)|S| + |T|$, and the average degree is at most $2+2\delta$ (actually, at most $2+\delta$, if we use symmetry and assume $|S| \geq |T|$). \square

Proof of Theorem 1.2. We let $\rho = \rho(\varepsilon) > 0$ be a sufficiently small constant, and we set the parameters of our random graph model as follows: we let d(v) = 5 for some ρm

vertices in A, and all remaining vertices in $A \cup B$ have d(v) = 4. Clearly, the average degree of H is $4 + \Omega(\rho)$, and thus if we choose δ sufficiently small in terms of ρ and let $\beta = \beta(5, \delta)$, then Lemma 4.1 and Corollary 4.3 guarantee $mcc_2(L(H)) \geq \beta m$ almost surely.

The maximum degree of L(H) is 7; the degree-7 vertices of the line graph correspond to edges of H incident to the ρm vertices of degree 5. It remains to bound the maximum average degree of L(H).

To this end, we apply Corollary 4.3 once again, this time with $\delta = \frac{1}{2}$, say, and we let $\beta_0 = \beta(5, \frac{1}{2})$ be the corresponding parameter. We note that β_0 is independent of ρ , and hence we can assume that β_0/ρ is sufficiently large.

Now let $F \subseteq E(H)$ be an arbitrary subset of edges, and let U be the set of all vertices incident to edges of F. If $|U| \leq \beta_0 m$, then by Corollary 4.3 the graph K := (U, F) has average degree at most 2.5. The average degree of L(K) is

$$\overline{d}(L(K)) = \frac{1}{|F|} \sum_{\{u,v\} \in F} (\deg_K(u) + \deg_K(v) - 2) = \frac{1}{|F|} \sum_{u \in U} \deg_K(u) (\deg_K(u) - 1).$$

If we denote by x_i the fraction of vertices $u \in U$ with $\deg_K(u) = i$, i = 1, 2, ..., 5, then the x_i satisfy the constraints $\sum_{i=1}^5 x_i = 1$ and $\sum_{i=1}^5 ix_i \le 2.5$ (this reflects the bound on the average degree of K), and we have $\overline{d}(L(K)) = 2(2x_2 + 6x_3 + 12x_4 + 20x_5)/(x_1 + 2x_2 + 3x_3 + 4x_4 + 5x_5)$. One can use, e.g., linear programming to verify that the above constraints imply $\overline{d}(L(K)) \le 6$ as needed.

If, on the other hand, $|U| > \beta_0 m$, then since H has at most ρm vertices of degree 5, these vertices constitute at most ρ/β_0 fraction of U, and all other vertices have degree at most 4. Hence at most a small fraction of the vertices of L(K) can have degree 7, while others have degrees at most 6, and it follows that the average degree of L(K) can be pushed below $6 + \varepsilon$ by making ρ sufficiently small. Theorem 1.2 is proved.

5 The Hamming cube

An interesting example, where mcc_2 can be determined exactly, is the Hamming cube. Let Q_d denote the d-dimensional Hamming cube with vertex set $\{0,1\}^d$ and with two vectors $u, v \in \{0,1\}^d$ adjacent in Q_d if they differ in exactly one coordinate. Let $L(Q_d)$ be the line graph of Q_d .

Proposition 5.1 For every even d we have $mcc_2(L(Q_d)) = \frac{d}{4}2^{d/2}$.

Proof. The upper bound is witnessed by the following coloring: Color an edge $\{u,v\} \in E(Q_d)$ red if u and v differ in one of the first d/2 coordinates, and blue otherwise. Then the monochromatic components are (d/2)-dimensional subcubes.

For the lower bound, it suffices to show that whenever the edges of Q_d are colored red and blue, there exists a monochromatic connected subgraph with at least $\frac{d}{4}2^{d/2}$ edges. Let us assume that, e.g., blue is the majority color; that is, at least $\frac{d}{4}2^d$ edges are blue. Let B_1, B_2, \ldots, B_k be the connected components of the blue subgraph, and let m_i be the number of blue edges in B_i . Hence $\sum_{i=1}^k m_i \geq \frac{1}{2}|E(Q_d)| = \frac{d}{4}2^d$.

We recall the following formulation of the edge-isoperimetric inequality for the cube (E.g., [Bol86], Chapter 16): Every subgraph of Q_d on v vertices has at most $\frac{1}{2}v\log_2 v$

edges. Let $\beta_i > 0$ be the real number satisfying $\frac{1}{2}\beta_i \log_2 \beta_i = m_i$ (where $\beta_i = 1$ for $m_i = 0$). Thus, β_i is a lower bound for the number of vertices of B_i , and consequently, $\sum_{i=1}^k \beta_i \leq 2^d$. Assuming for contradiction that $m_i < \frac{d}{4}2^{d/2}$ for all i, we have $\beta_i < 2^{d/2}$ for all i, and thus

$$\sum_{i=1}^{k} m_i = \sum_{i=1}^{k} \frac{1}{2} \beta_i \log_2 \beta_i < \frac{d}{4} \sum_{i=1}^{k} \beta_i \le \frac{d}{4} \sum_{i=1}^{k} |V(B_i)| \le \frac{d}{4} 2^d.$$

But as was noted above, $\sum_{i=1}^k m_i \geq \frac{d}{4}2^d$, and this contradiction establishes the proposition.

The proof also shows that the monochromatic components in any extremal coloring have to be (d/2)-dimensional subcubes.

6 Open problems

There are quite a few interesting open questions suggested by the present paper. Here are some of them.

- 1. How large can $mcc_2(G)$ be for graphs of maximum degree 6? By the planar HEX lemma, the triangulated planar grid is an example with $mcc_2(G) = \Theta(\sqrt{n})$, but this is at present the best we know.
- 2. A special case of the previous question, which seems interesting in its own right, is when G = L(H) for some 4-regular H. The best lower bound we know is $\Omega(\log n)$, from a construction by Alon et al. [ADO+03], where H is a 4-regular graph of logarithmic girth.
- 3. The examples we know for planar graphs G with large $mcc_2(G)$ and $mcc_3(G)$ have at least one vertex of high degree. Can anything better be said if we assume that G has bounded degrees? More specifically, the following was asked in [KMR+97]: Is there a function f such that for every planar graph G of maximum degree Δ we have $mcc_3(G) \leq f(\Delta)$?
- 4. For two colors we cannot hope for constant monochromatic component size in bounded-degree planar graphs, as shown by a triangulated planar grid, but similar to the previous question, we can ask if there exists a function g such that every n-vertex planar graph G of maximum degree Δ satisfies $mcc_2(G) \leq g(\Delta)\sqrt{n}$.
- 5. There is still a gap between the best bounds we know for $mcc_t(G)$ for graphs from minor-closed families of graphs. Can this gap be closed?
- 6. A question suggested to us by Emo Welzl concerns the possible behavior of $mcc_t(G)$ when the chromatic number of G as well as its number of vertices are known.
- 7. The proof of Theorem 1.2 can be adapted to show that for any fixed t there exist n-vetrex graphs with maximum degree 4t-1 and with $mcc_t(G) = \Omega(n)$. Note that in [ADO+03] it was shown that $mcc_t(G)$ is not bounded by a constant even

for graphs of maximum degree 4t-2; however, that proof gives only a logarithmic lower bound for $mcc_t(G)$ (cf. our first open problem). From the other direction [HST03] show that $mcc_t(G)$ is bounded by a constant for all t and all graphs G with maximum degree at most 3t-1. The constant 3 here is not optimal, since the same paper shows that for some constant $\varepsilon > 0$ and all sufficiently large t, the value $mcc_t(G)$ is bounded by a constant for all graphs G of maximum degree at most $(3+\varepsilon)t$. It would be interesting to find the asymptotic behaviour of the maximal value of $mcc_t(G)$ for graphs G with maximum degree d in the intermediate range $(3+\epsilon)t < d < 4t-2$. In particular, it would be interesting to know if there exist t and d for which the above maximum is sublinear but not a constant.

8. There are several natural conjectures pertaining to mcc_t for triangulations of the d-dimensional grid graph. These questions suggest an interesting "combinatorial dimension theory" waiting to be discovered. More on this subject can be found in [MP07].

Acknowledgements

We would like to thank Maria Chudnovsky and Tibor Szabó for fruitful conversations and to Eyal Ackerman and Robin Thomas for important references.

References

- [ADO+03] N. Alon, G. Ding, B. Oporowski, D. Vertigan. Partitioning into graphs with only small components. *Journal of Combinatorial Theory* (Series B), 87:231–243, 2003.
- [AST90] N. Alon, P. Seymour and R. Thomas. A separator theorem for nonplanar graphs. J. Am. Math. Soc. 3,4:801–808, 1990.
- [Bol86] B. Bollobás. Set systems, hypergraphs, families of vectors and combinatorial probability. Cambridge University Press, Cambridge, 1986.
- [BL91] B. Bollobás and I. Leader. Edge-isoperimetric inequalities in the grid. Combinatorica 11(4):299–314, 1991.
- [BS05] R. Berke and T. Szabó. Relaxed two-coloring of cubic graphs. *DMTCS Proceedings of 2005 European Conference on Combinatorics, Graph Theory and Applications (EUROCOMB)*, 341–344, 2005.
- [DOS+96] G. Ding, B. Oporowski, D. Sanders, D. Vertigan, preprint, Louisiana State University, Baton Rouge, 1996.
- [G79] D. Gale. The game of hex and the Brouwer fixed-point theorem. Amer. Math. Monthly 86:818–827, 1979.
- [HST03] P. Haxell, T. Szabó, G. Tardos. Bounded size components—partitions and transversals. *Journal of Combinatorial Theory* (Series B) 88:281–297, 2003.

- [HLW06] S. Hoory, N. Linial, and A. Wigderson. Expander graphs and their applications. *Bull. Amer. Math. Soc.* 43,4:439–561,2006.
- [JLR00] S. Janson, T. Łuczak, A. Ruciński: *Topics in random graphs*, J. Wiley and Sons, New York, NY, 2000.
- [JT95] T. R. Jensen and B. Toft. *Graph coloring problems*. Wiley-Interscience Series in Discrete Mathematics and Optimization, John Wiley & Sons Inc., New York, 1995.
- [JW96] B. Jackson and N. Wormald. On the linear k-arboricity of cubic graphs. Discrete Math 162:293–297, 1996.
- [KMR+97] J. Kleinberg, R. Motwani, P. Raghavan, and S. Venkatasubramanian. Storage management for evolving databases, FOCS 1997, 353–362.
- [LS93] N. Linial and M. Saks. Low diameter graph decompositions. *Combinatorica* 13:441–454, 1993.
- [LT79] R. Lipton and R. Tarjan. A separator theorem for planar graphs. SIAM J. of Applied Mathematics 2:177–190, 1979.
- [MN98] J. Matoušek and J. Nešetřil. *Invitation to Discrete Mathematics*. Oxford University Press, Oxford 1998.
- [MP07] J. Matoušek and A. Přívětivý. Large monochromatic components in two-colored grids. Submitted, 2007.
- [R92] B. Reed. Finding approximate separators and computing tree width quickly. STOC 1992, Victoria B.C., 1992.
- [RS91] N. Robertson and P. Seymour. Graph Minors X. Obstructions to tree-decomposition. J. Combin. Theory Ser. B 52:153–190, 1991.
- [RS95] N. Robertson and P. Seymour. Graph Minors XIII. The Disjoint Path Problem. J. Combin. Theory Ser. B, 63:65–110, 1995.
- [Tho99] C. Thomassen. Two-colouring the edges of a cubic graph such that each monochromatic component is a path of length at most 5. *Journal of Combinatorial Theory* (Series B) 75:100–109, 1999.