### LICCI BINOMIAL EDGE IDEALS

#### VIVIANA ENE, GIANCARLO RINALDO, NAOKI TERAI

ABSTRACT. We give a complete characterization of graphs whose binomial edge ideal is licci. An important tool is a new general upper bound for the regularity of binomial edge ideals.

#### Introduction

Binomial edge ideals associated to simple graphs have been intensively studied in the last decade. Their algebraic and homological properties are intimately related to the combinatorics of the underlying graph. A lot of effort has been dedicated to study the Cohen-Macaulay property of these ideals. As in the case of classical edge ideals, an exhaustive classification of graphs whose binomial edge ideals are Cohen-Macaulay seems to be a hopeless task. There are successful attempts to characterize graphs with specific properties which have Cohen-Macaulay binomial edge ideals. For example, the Cohen-Macaulay property of binomial edge ideals is known for block graphs which include the trees [3] and for bipartite graphs [1]. We refer also to the papers [12, 20, 21, 22] for other classes of Cohen-Macaulay binomial edge ideals.

Let G be a simple graph (that is, undirected, with no loops, and no multiple edges) on the vertex set  $[n] := \{1, 2, ..., n\}$  and  $S = K[x_1, ..., x_n, y_1, ..., y_n]$  the polynomial ring in 2n variables. The binomial edge ideal  $J_G \subset S$  of G is generated by all the binomials of the form  $f_{ij} = x_i y_j - x_j y_i$  where  $\{i, j\}$  is an edge of G. In other words,

binomials of the form 
$$f_{ij} = x_i y_j - x_j y_i$$
 where  $\{i, j\}$  is an edge of  $G$ . In other words,  $J_G$  is generated by the 2-minors of the generic matrix  $X = \begin{pmatrix} x_1 & x_2 & \dots & x_n \\ y_1 & y_2 & \dots & y_n \end{pmatrix}$  which correspond to the edges of  $G$ .

In this paper, we study binomial edge ideals which are in the linkage class of a complete intersection. We call such ideals licci, in brief. Besides the Cohen-Macaulay property, they satisfy some extra conditions which make possible a full characterization of graphs whose binomial edge ideals are licci. Linkage theory has a rich history in commutative algebra and algebraic geometry. Peskine and Szpiro [19] in 1974 reduced general linkage to questions on ideals over commutative algebras and after then, a lot of work has been done to develop this theory in commutative algebra and algebraic geometry. If I, J are proper ideals in a local regular ring R, they are called directly linked and we write  $I \sim J$  if there exists a regular sequence

<sup>2010</sup> Mathematics Subject Classification. 14M06, 13C40, 13H10, 05E40,

Key words and phrases. licci ideals, binomial edge ideals, regularity.

The second author was supported by GNSAGA of INdAM (Italy). The third author was supported by the JSPS Grant-in Aid for Scientific Research (C) 18K03244.

 $\mathbf{z} = z_1, \dots, z_g$  in  $I \cap J$  such that  $J = (\mathbf{z}) : I$  and  $I = (\mathbf{z}) : J$ . One says that I and J belong to the same *linkage class* if there exists a sequence of direct links

$$I = I_0 \sim I_1 \sim \cdots \sim I_m = J.$$

If J is a complete intersection ideal, then I is said to be licci. The ideals in the same linkage class share several properties. For example, if I and J are linked, then I is Cohen-Macaulay if and only if J is Cohen-Macaulay. In particular, it follows that a licci ideal is Cohen-Macaulay.

The following natural question arises. May we give a full characterization of the graphs G with the property that the associated binomial edge ideal is licci?

In this paper, we give a complete answer to this question. In [10] a necessary condition for a Cohen-Macaulay homogeneous ideal in a polynomial ring to be licci is given. In the case of binomial edge ideals, this condition implies that if  $(J_G)_{\mathfrak{m}} \subset S_{\mathfrak{m}}$  (here  $\mathfrak{m}$  is the maximal graded ideal of the ring S) is licci, then  $\operatorname{reg}(S/J_G) \geq n-2$ . This condition turns to be also sufficient for Cohen-Macaulay binomial edge ideals as we are going to show in this paper.

The regularity of binomial edge ideals have been intensively studied in the last years. In [15] it was proved that the regularity of  $S/J_G$  is upper bounded by n-1 and it was conjectured that this upper bound is attained if and only if G is a path graph. This conjecture was later proved in [13]. Inspired by the paper [13], we prove a new upper bound for  $reg(S/J_G)$  which is stronger than n-1 and it plays an essential role in the characterization of the graphs G whose binomial edge ideal is licci.

The structure of the paper is as follows. In Section 1, we recall the basic results on licci and binomial edge ideals needed in the next sections. In Section 2, we prove that if G is a connected graph, then  $\operatorname{reg}(S/J_G) \leq n - \dim \Delta(G)$ , where  $\Delta(G)$  is the clique complex of G (Theorem 2.1). We believe that this new general upper bound for the regularity of binomial edge ideals will inspire new results on their resolution. In brief, in Theorem 2.1, we prove that for every clique  $W \subset [n]$  of the connected graph G, we have  $\operatorname{reg}(S/J_G) \leq n - |W| + 1$ . The proof is based on a double induction. First we make induction on n - |W| and, secondly, on a combinatorial invariant of G.

The characterization of graphs whose binomial edge ideal is licci is given in Section 3. In Theorem 3.5 we show that, for a connected graph G on n vertices, the following statements are equivalent:

- (i)  $(J_G)_{\mathfrak{m}} \subset S_{\mathfrak{m}}$  is licci.
- (ii)  $J_G$  is Cohen-Macaulay and  $n-2 \leq \operatorname{reg}(S/J_G) \leq n-1$ .
- (iii) G is a path graph or it is a triangle with possibly some paths attached to some of its vertices.

The most technical part in the proof is to show that there is no indecomposable graph G with  $n \geq 4$  vertices with  $\operatorname{reg}(S/J_G) = n-2$  and  $J_G$  Cohen-Macaulay. In order to make this part easier to understand, we proved some preparatory lemmas. We can reformulate the above statement by saying that the only indecomposable graphs G with  $J_G$  a Cohen-Macaulay ideal and  $\operatorname{reg}(S/J_G) = n-2$  are the path with

one edge and the triangle. Next we combine this fact with Lemma 3.2 which shows that for any decomposable graph G with  $reg(S/J_G) = n - 2$ , one of the components must be a path. In this way we derive the combinatorial characterization from Theorem 3.5 (iii).

A straightforward consequence of Theorem 3.5 is Corollary 3.6 which says that for a connected bipartite graph G, the ideal  $(J_G)_{\mathfrak{m}} \subset S_{\mathfrak{m}}$  is licci if and only if G is a path graph. The case when G is a disconnected graph is treated in Proposition 3.7.

In the last section of the paper, we show that for chordal graphs, in the equivalent statements of Theorem 3.5, we may replace the Cohen-Macaulay property with the unmixedness of the ideal  $J_G$  (Theorem 4.2). For the proof we use a theorem of Dirac which characterizes the chordal graphs in terms of their clique complex.

## 1. Preliminaries

We recall some notions and fundamental results needed in the later sections.

1.1. Licci ideals. Let R be a regular local ring and I, J proper ideals of R. Then I and J are called directly linked and we write  $I \sim J$  if there exists a regular sequence  $\mathbf{z} = z_1, \ldots, z_g$  in  $I \cap J$  such that  $J = (\mathbf{z}) : I$  and  $I = (\mathbf{z}) : J$ . One says that I is linked to J or that I and J belong to the same linkage class if there exists a sequence of direct links

$$I = I_0 \sim I_1 \sim \cdots \sim I_m = J.$$

If J is a complete intersection ideal, that is, it is generated by a regular sequence, then I is said to be in the linkage class of a complete intersection (*licci* in brief).

Several properties are preserved in the same linkage class. For example, if I is linked to J, then R/I is Cohen-Macaulay if and only if R/J is Cohen-Macaulay [19]. In particular, any licci ideal is Cohen-Macaulay. A necessary condition for a homogeneous ideal in a polynomial ring to be licci is given in [10].

**Theorem 1.1.** [10, Corollary 5.13] Let I be a Cohen-Macaulay homogeneous ideal in a standard graded polynomial ring  $S = K[x_1, \ldots, x_n]$  with the graded maximal ideal  $\mathfrak{m}$ . If  $I_{\mathfrak{m}} \subset R = S_{\mathfrak{m}}$  is licci, then

(1) 
$$\operatorname{reg}(S/I) \ge (\operatorname{height} I - 1)(\operatorname{indeg} I - 1)$$

where indeg I is the initial degree of the ideal I, that is, indeg  $I = \min\{i : I_i \neq 0\}$ .

Although, in general, inequality (1) is not a sufficient condition, if I is the edge ideal of a graph, then  $I_{\mathfrak{m}} \subset R = S_{\mathfrak{m}}$  is licci if and only if inequality (1) holds [14]. We will see a similar behavior in Section 3 for binomial edge ideals.

1.2. **Graphs and binomial edge ideals.** Let G be a simple graph on the vertex set V(G) := [n] with the edge set E(G) and  $S = K[x_1, \ldots, x_n, y_1, \ldots y_n]$  the polynomial ring in 2n variables over a field K. The binomial edge ideal of the graph G is generated by the binomials  $f_e := x_i y_j - x_j y_i$  with  $e = \{i, j\} \in E(G)$ . In other words,  $J_G$  is generated by the 2-minors of the matrix  $X = \begin{pmatrix} x_1 & x_2 & \ldots & x_n \\ y_1 & y_2 & \ldots & y_n \end{pmatrix}$  which correspond to the edges of G. For example, if G is the complete graph  $K_n$  on

n vertices, then  $J_G$  is the ideal  $I_2(X)$  generated by all the 2-minors of X, while if G is the path graph  $P_n$  on n vertices with edge set  $\{\{i, i+1\}: 1 \leq i \leq n-1\}$ , then  $J_G$  is the ideal of all adjacent maximal minors of X.

The binomial edge ideals were introduced independently in the papers [6] and [17]. In the last decade, these ideals have been studied by many authors. The interested reader may find a thorough introduction to this topic in the monograph [7]. Fundamental results regarding the minimal free resolutions of binomial edge ideals are surveyed in [24].

In this paper, we need to recall the primary decomposition of binomial edge ideals and some fundamental results on their regularity.

The minimal primary decomposition of a binomial edge ideal is strongly related to the combinatorics of the underlying graph; see [6] or [7, Chapter 7]. Let S be a (possibly empty) subset of [n] and let  $G_S$  be the restriction of G to the vertex subset  $[n] \setminus S$ . Let  $G_1, \ldots, G_{c(S)}$  be the connected components of this restriction and, for every  $1 \le i \le c(S)$ , let  $\tilde{G}_i$  be the complete graph on  $V(G_i)$ . Then, the ideal

$$P_{\mathcal{S}}(G) = (\{x_i, y_i : i \in \mathcal{S}\}) + J_{\tilde{G}_1} + \dots + J_{\tilde{G}_{c(\mathcal{S})}}$$

is a prime ideal in S which contains  $J_G$ , and by [6, Lemma 3.1] we have

(2) 
$$\operatorname{height}(P_{\mathcal{S}}(G)) = n - c(\mathcal{S}) + |\mathcal{S}|.$$

**Theorem 1.2.** [6] In the above notation, we have

$$J_G = \bigcap_{S \subset [n]} P_S(G).$$

In particular,  $J_G$  is a radical ideal and its minimal prime ideals are among  $P_S(G)$  with  $S \subset [n]$ . The following proposition characterizes the sets S for which the prime ideal  $P_S(G)$  is minimal.

**Proposition 1.3.** [6, Corollary 3.9]  $P_{\mathcal{S}}(G)$  is a minimal prime ideal of  $J_G$  if and only if either  $\mathcal{S} = \emptyset$  or  $\mathcal{S}$  is non-empty and for each  $i \in \mathcal{S}$ ,  $c(\mathcal{S} \setminus \{i\}) < c(\mathcal{S})$ .

In graph theoretical terminology, for a connected graph G,  $P_{\mathcal{S}}(G)$  is a minimal prime ideal of  $J_G$  if and only if  $\mathcal{S}$  is empty or  $\mathcal{S}$  is non-empty and is a *cut set* of G, that is, i is a cut vertex of the restriction  $G_{([n]\setminus\mathcal{S})\cup\{i\}}$  for every  $i\in\mathcal{S}$ . We recall that a vertex v of the graph H is a *cut vertex* of H if its removing breaks H into more connected components than H has. Let  $\mathcal{C}(G)$  be the set of all sets  $\mathcal{S} \subset [n]$  such that  $P_{\mathcal{S}}(G)$  is a minimal prime ideal of  $J_G$ . Equality (2) implies then the following.

**Corollary 1.4.** Let G be a connected graph on the vertex set [n]. Then  $J_G$  is unmixed if and only if for every  $S \in C(G)$ , c(S) = |S| + 1.

A general upper bound for the regularity of binomial edge ideals was first given in [15], namely,  $reg(S/J_G) \le n-1$ , and in the same paper it was conjectured that  $reg(S/J_G) = n-1$  if and only if G is a path graph. This conjecture was proved in [13].

**Theorem 1.5.** [13] Let G be a graph on n vertices which is not a path. Then  $reg(S/J_G) \le n-2$ .

For a chordal graph G, in [23, Theorem 3.5] it was shown that the number c(G) of maximal cliques of G is an upper bound for  $reg(S/J_G)$ .

Recall that a subset  $C \subset [n]$  is a *clique* of G if the induced subgraph of G on the vertex set G is a complete graph. The set of cliques of G forms a simplicial complex  $\Delta(G)$  called the *clique complex* of G. Its facets are the maximal cliques of G. By a famous theorem of Dirac ([2] or [5, Section 9.2]), a connected graph G is chordal if and only if either G is a complete graph or the facets of  $\Delta(G)$  can be ordered as  $F_1, \ldots, F_c$  such that, for all i > 1,  $F_i$  is a leaf of the simplicial complex generated by  $F_1, \ldots, F_i$ . A leaf of a simplicial complex G is a facet of G which has a branch, that is, a facet G such that for all facets G of G with G is the problem of G is a facet of G.

2. A NEW UPPER BOUND FOR THE REGULARITY OF BINOMIAL EDGE IDEALS

In this section, we give a new general upper bound for the regularity of  $S/J_G$ .

**Theorem 2.1.** Let G be a connected graph on [n]. Then  $reg(S/J_G) \leq n - \dim \Delta(G)$ .

When G is not connected, we derive the following upper bound for the regularity of  $S/J_G$ .

Corollary 2.2. Let G be a graph on n vertices with the connected components  $G_1, \ldots, G_c$ . Then

$$\operatorname{reg}(S/J_G) \le n - (\dim \Delta(G_1) + \dots + \dim \Delta(G_c)).$$

Let us make some short remarks before proving the above theorem. This new bound will be an essential tool in proving Theorem 3.5. Although for chordal graphs, this bound might be larger than the number of maximal cliques of G, it is a general bound which is better than n-1.

In what follows, we will need some notation and known results. If H is a graph and  $e \in E(H)$ , we denote by  $H \setminus e$  the subgraph of H obtained by removing the edge e from E(H) and if  $e_1, \ldots, e_m \in E(H)$ , we write  $H \setminus \{e_1, \ldots, e_m\}$  for the subgraph of H which is obtained by removing the edges  $e_1, \ldots, e_m$ . If  $e = \{i, j\}$  where i, j are vertices of H and  $e \notin E(G)$ , then  $H \cup e$  is the graph with the same vertex set as H and edge set  $E(H) \cup \{e\}$ , and  $H_e$  is the graph with  $V(H_e) = V(H)$  and  $E(H_e) = E(H) \cup \{\{k, \ell\} : k, \ell \in N(i) \text{ or } k, \ell \in N(j)\}$  where N(i) denotes the set of all neighbors of i in H.

The next proposition is a direct consequence of the behavior of the regularity with respect to short exact sequences; see [18, Corollary 18.7].

**Proposition 2.3.** [13, Proposition 2.1] Let H be a graph on n vertices and  $J_H \subset S$  its binomial edge ideal. Let  $e = \{i, j\}$  be an edge of H and  $f_e = x_i y_j - x_j y_i$ . Then, the following inequalities hold:

- (a)  $\operatorname{reg}(J_H) \leq \max\{\operatorname{reg}(J_{H \setminus e}), \operatorname{reg}(J_{H \setminus e}: f_e) + 1\};$
- (b)  $\operatorname{reg}(J_{H \setminus e}) \le \max\{\operatorname{reg}(J_H), \operatorname{reg}(J_{H \setminus e}: f_e) + 2\};$
- (c)  $reg(J_{H \setminus e} : f_e) + 2 \le max\{reg(J_{H \setminus e}), reg(J_H) + 1\}.$

In the settings of the above proposition, we have the following.

**Theorem 2.4.** [16, Theorem 3.7]

$$J_{H\backslash e}: f_e = J_{(H\backslash e)_e} + I_{H,e}$$

where  $I_{H,e}$  is the monomial ideal generated by the set

$$\{g_{\pi,t}|\pi:i,i_1,\ldots,i_s,j \text{ is a path between } i \text{ and } j \text{ and } 0 \leq t \leq s\}$$

and

$$g_{\pi,0} = x_{i_1} \cdots x_{i_s}, g_{\pi,t} = y_{i_1} \cdots y_{i_t} x_{i_{t+1}} \cdots x_{i_s} \text{ for } 1 \le t \le s.$$

Proof of Theorem 2.1. Clearly, the statement of the theorem follows if we show that for any clique  $W \subset [n]$ , we have

(3) 
$$\operatorname{reg}(S/J_G) \le n - |W| + 1$$
 or, equivalently,  $\operatorname{reg}(J_G) \le n - |W| + 2$ .

We prove this by induction on n - |W|. If n = |W|, then G is the complete graph on n vertices and it is well known that  $reg(S/J_G) = 1$ .

Let n-|W|>0. We proceed with the inductive step. For the remaining part of the proof, we need to define the following. For a vertex  $v\in V(G)$ , we set  $\alpha_G(v):=\binom{\deg v}{2}-|E(G_{N(v)})|$ . Here, we used the usual notation  $G_U$  for the restriction of G to the subset U of V(G). Obviously,  $\alpha_G(v)=0$  if and only if v is a simplicial vertex in G. Recall that a vertex of a graph is called *simplicial* if it belongs to exactly one maximal clique. In addition, for a subset  $W\subset V(G)$ , we define  $\alpha_G(W):=\min\{\alpha_G(v):v\in V(G)\setminus W\}$ . Further on, we proceed by induction on  $\alpha_G(W)$ .

**Step 1.** Let  $\alpha_G(W) = 0$ . Thus, there exists a simplicial vertex  $v \in V(G) \setminus W$ . Now we consider two cases, namely deg v = 1 and deg  $v \ge 2$ .

Case 1. Let deg(v) = 1 and  $e = \{v, w\} \in E(G)$ . By Proposition 2.3 (a), we have

$$reg(J_G) \le \max\{reg(J_{G \setminus e}), reg(J_{G \setminus e} : f_e) + 1\}.$$

Therefore, it is enough to show that

$$reg(J_{G \setminus e}) \le n - |W| + 2$$

and

(5) 
$$\operatorname{reg}(J_{G \setminus e} : f_e) \le n - |W| + 1.$$

Since  $\deg(v) = 1$ , the vertex v becomes isolated in the graph  $G \setminus e$ , thus  $\operatorname{reg}(J_{G \setminus e}) = \operatorname{reg}(J_{(G \setminus e) \setminus v})$ . So, for showing inequality (4), we simply apply the inductive hypothesis to the graph  $(G \setminus e) \setminus v$ . For showing inequality (5), we first apply Theorem 2.4 and get

$$reg(J_{G\backslash e}:f_e) = reg(J_{(G\backslash e)_e}),$$

since,  $I_{G,e} = (0)$  because the only path connecting v and w in G is the edge  $\{v, w\}$ . In the graph  $(G \setminus e)_e$ , v is an isolated vertex, thus,

$$reg(J_{(G\backslash e)_e}) = reg(J_{((G\backslash e)_e)\backslash v}).$$

Now we can apply again the inductive hypothesis for  $(G \setminus e)_e \setminus v$  and obtain

$$reg(J_{(G \setminus e)_e \setminus v}) \le (n-1) - |W| + 2 = n - |W| + 1.$$

Therefore, Case 1 is completed.

Case 2. Let v be a simplicial vertex of  $deg(v) = t \ge 2$ . Before discussing this case,

we prove the following.

**Claim**. Assume that there exists  $v \in V(G) \setminus W$  a simplicial vertex with  $\deg(v) \geq 2$ . Let e be an edge of G which contains v. Then

$$reg(J_{G \setminus e} : f_e) \le n - |W| + 1.$$

Proof of the Claim. Let  $\deg(v)=t$ , let  $N_G(v)=\{v_1,\ldots,v_t\}$  be the set of neighbors of v in G, and set  $e_i=\{v,v_i\}$  for  $1\leq i\leq t$ . We may assume that  $e=e_t$  and let us consider the monomial ideal  $I_{G,e}$  from Theorem 2.4. Since v is a simplicial vertex, for any  $1\leq i\leq t-1$ ,  $v_t,v_i,v$  is a path in G, thus  $x_{v_i},y_{v_i}\in I_{G,e}$  for all  $1\leq i\leq t-1$ . Moreover, every path from v to  $v_t$  must pass through some neighbor  $v_i$  with  $1\leq i\leq t-1$ . This implies that

$$I_{G,e} = (x_{v_i}, y_{v_i} : 1 \le i \le t - 1).$$

By Theorem 2.4, we get

$$J_{G \setminus e} : f_e = J_{(G \setminus e)_e} + (x_{v_i}, y_{v_i} : 1 \le i \le t - 1).$$

Set  $H := (G \setminus e)_e$ . Then

$$J_{G \setminus e} : f_e = J_{H_{[n] \setminus \{v_1, \dots, v_{t-1}\}}} + (x_{v_i}, y_{v_i} : 1 \le i \le t - 1),$$

because the binomial generators of  $H=J_{(G\backslash e)_e}$  corresponding to the edges which contain some  $v_i$  with  $1\leq i\leq t-1$  are contained in  $I_{G,e}$ . Since v becomes an isolated vertex in  $H_{[n]\backslash \{v_1,\ldots,v_{t-1}\}}$ , we get

$$J_{G \setminus e}: f_e = J_{H_{[n] \setminus \{v, v_1, \dots, v_{s-1}\}}} + (x_{v_i}, y_{v_i}: 1 \le i \le t-1),$$

which implies that

$$\operatorname{reg}(J_{G \setminus e} : f_e) = \operatorname{reg}(J_{H_{[n] \setminus \{v, v_1, \dots, v_{t-1}\}}}).$$

The graph  $H_{[n]\setminus\{v,v_1,\ldots,v_{t-1}\}}$  has n-t vertices and the clique  $W\setminus\{v,v_1,\ldots,v_{t-1}\}$ , thus we may apply the inductive hypothesis because

$$(n-t)-|W\setminus \{v,v_1,\ldots,v_{t-1}\}| \le n-t-|W|+t-1=n-|W|-1.$$

Therefore, we get

$$reg(J_{G \setminus e} : f_e) = reg(J_{H_{[n] \setminus \{v, v_1, \dots, v_{t-1}\}}}) \le (n-t) - |W \setminus \{v, v_1, \dots, v_{t-1}\}| + 2 \le n - |W| + 1,$$
 and the claim is proved.

We now go back to the discussion of Case 2. Let  $N_G(v) = \{v_1, \ldots, v_t\}$  be the set of the neighbors of v in G and  $e_i = \{v, v_i\}$  for  $1 \le i \le t$ . By Proposition 2.3 and the Claim, we have

$$\operatorname{reg}(J_G) \le \max\{\operatorname{reg}(J_{G \setminus e_1}), \operatorname{reg}(J_{G \setminus e_1}: f_{e_1}) + 1\} \le \max\{\operatorname{reg}(J_{G \setminus e_1}), n - |W| + 2\}.$$

Applying the same argument to  $G \setminus e_1$ , we obtain

$$reg(J_G) \le max\{reg(J_{G\setminus \{e_1,e_2\}}), n - |W| + 2\}.$$

After t-1 steps, we get

$$\operatorname{reg}(J_G) \le \max\{\operatorname{reg}(J_{G\setminus\{e_1,e_2,\dots,e_{t-1}\}}), n-|W|+2\}.$$

In the graph  $G \setminus \{e_1, e_2, \dots, e_{t-1}\}$ , we have  $\deg(v) = 1$ . Consequently, by Case 1, we derive that  $\operatorname{reg}(J_G) \leq n - |W| + 2$  which completes the proof of Step 1.

Now we proceed to prove the inductive step on  $\alpha_G(W)$ .

**Step 2.** Let  $\alpha_G(W) > 0$ . This implies that there exists a non-simplicial vertex  $v \in V(G) \setminus W$  such that  $\alpha_G(W) = \alpha_G(v)$ . Moreover, since v is not simplicial, there exist  $v_1, v_2 \in N_G(v)$  such that  $e = \{v_1, v_2\} \not\in E(G)$ . By Proposition 2.3 (b) where  $H = G \cup e$ , it follows

(6) 
$$\operatorname{reg}(J_G) \le \max\{\operatorname{reg}(J_{G \cup e}), \operatorname{reg}(J_G : f_e) + 2\}.$$

By the definition of  $\alpha_G(v)$ , we have  $\alpha_{G \cup e}(v) = \alpha_G(v) - 1$ , therefore  $\alpha_{G \cup e}(W) \le \alpha_G(W) - 1$ . By induction on  $\alpha_G(W)$ , we then derive that

$$reg(J_{G \cup e}) \le n - |W| + 2.$$

In order to complete this last step, by using (6), it is enough to show that

(7) 
$$\operatorname{reg}(J_G: f_e) + 2 \le n - |W| + 2.$$

By Theorem 2.4, we have

(8) 
$$J_G: f_e = J_{G_e} + I_{G \cup e,e}.$$

Since  $v_1, v, v_2$  is a path, the variables  $x_v, y_v$  belong to  $I_{G \cup e,e}$ . This implies that

$$I_{G \cup e,e} = (x_v, y_v) + I_{(G \setminus v) \cup e,e}.$$

By replacing  $I_{G \cup e,e}$  in equality (8), we can rewrite it as

$$J_G: f_e = J_{(G \setminus v)_e} + I_{(G \setminus v) \cup e, e} + (x_v, y_v).$$

This implies that

$$\operatorname{reg}(J_G: f_e) = \operatorname{reg}(J_{(G \setminus v)_e} + I_{(G \setminus v) \cup e, e}).$$

On the other hand, by Theorem 2.4 applied for  $G \setminus v$ , we get

$$\operatorname{reg}(J_{(G \setminus v)_e} + I_{(G \setminus v) \cup e,e}) = \operatorname{reg}(J_{G \setminus v} : f_e),$$

thus,

$$reg(J_G: f_e) = reg(J_{G \setminus v}: f_e).$$

Next, by Proposition 2.3, (c) we have

$$\operatorname{reg}(J_{G\setminus v}: f_e) + 2 \le \max\{\operatorname{reg}(J_{G\setminus v}), \operatorname{reg}(J_{(G\setminus v)\cup e}) + 1\}.$$

By the inductive hypothesis on n-|W|, we have

$$reg(J_{G\setminus v}) \le (n-1) - |W| + 2 = n - |W| + 1,$$

and

$$reg(J_{(G\setminus v)\cup e}) + 1 \le (n-1) - |W| + 3 = n - |W| + 2.$$

Consequently, we proved inequality (7) and this completes Step 2 and the whole proof of the theorem.

### 3. LICCI BINOMIAL EDGE IDEALS

As in the previous section, let G be a simple graph on the vertex set [n] and  $S = K[x_1, \ldots, x_n, y_1, \ldots, y_n]$  the polynomial ring over a field K. Let  $\mathfrak{m}$  be the maximal graded ideal of S and set  $R = S_{\mathfrak{m}}$ .

We recall the notion of decomposable graphs from [8].

**Definition 3.1.** A connected graph G is called decomposable if there exists two subgraphs  $G_1$  and  $G_2$  of G such that  $G = G_1 \cup G_2$  with  $V(G_1) \cap V(G_2) = \{v\}$  where v is a simplicial vertex in  $G_1$  and  $G_2$ . In this case we say that G is decomposable in the vertex v. Otherwise, the graph G is called indecomposable.

As it was proved in [8], if G is decomposable, then  $\operatorname{reg}(S/J_G) = \operatorname{reg} S_1/J_{G_1} + \operatorname{reg} S_2/J_{G_2}$  where  $S_i = K[\{x_j, y_j : j \in V(G_i)\}]$  for i = 1, 2. Moreover, by [20, Theorem 2.7],  $J_G$  is Cohen-Macaulay if and only if  $J_{G_1}$  and  $J_{G_2}$  are Cohen-Macaulay.

Before proving the main theorem of this section, we state some lemmas which are useful in what follows.

**Lemma 3.2.** Let G be a decomposable graph as  $G = G_1 \cup G_2$  with  $|V(G_i)| = n_i$  for i = 1, 2 and let  $S_i = K[\{x_j, y_j\} : j \in V(G_i)]$  for i = 1, 2. If  $reg(S/J_G) = n - 2$ , then  $reg(S_1/J_{G_1}) = n_1 - 2$  and  $G_2$  is a path or  $reg(S_2/J_{G_2}) = n_2 - 2$  and  $G_1$  is a path.

*Proof.* We have

$$n-2 = \operatorname{reg}(S/J_G) = \operatorname{reg}(S_1/J_{G_1}) + \operatorname{reg}(S_2/J_{G_2}) \le (n_1-1) + (n_2-1) = n-1.$$

This implies that  $reg(S_1/J_{G_1}) = n_1 - 2$  and  $reg(S_2/J_{G_2}) = n_2 - 1$ , which means that  $G_2$  is a path by Theorem 1.5, or  $reg(S_2/J_{G_2}) = n_2 - 2$  and  $reg(S_1/J_{G_1}) = n_1 - 1$ , that is,  $G_1$  is a path.

**Lemma 3.3.** Let G be a connected graph on the vertex set [n]. Suppose that G has a cut vertex v with  $\deg_G(v) \geq 4$ . Then  $\operatorname{reg}(S/J_G) \leq n-3$ .

*Proof.* Since v is a cut vertex of G, by [17, Lemma 4.8], we get

$$J_G = J_{G_v} \cap (J_{G \setminus v} + (x_v, y_v))$$

where  $G_v$  is the graph on  $V(G_v) = V(G)$  with the edge set

$$E(G_v) = E(G) \cup \{\{u, w\} : u, w \in N_G(v)\}.$$

Consequently, we have the following exact sequence

$$0 \to \frac{S}{J_G} \to \frac{S}{J_{G_v}} \bigoplus \frac{S}{J_{G \setminus v} + (x_v, y_v)} \to \frac{S}{J_{G_v \setminus v} + (x_v, y_v)} \to 0,$$

since  $J_{G_v} + (J_{G \setminus v} + (x_v, y_v)) = J_{G_v \setminus v} + (x_v, y_v)$ . From this exact sequence we obtain

(9) 
$$\operatorname{reg} \frac{S}{J_G} \leq \max \{ \operatorname{reg} \frac{S}{J_{G_v}}, \operatorname{reg} \frac{S}{J_{G \setminus v} + (x_v, y_v)}, \operatorname{reg} \frac{S}{J_{G_v \setminus v} + (x_v, y_v)} + 1 \}.$$

By our assumption, v has at least 4 neighbors in G. Therefore, in  $G_v$  we have a maximal clique with at least 5 vertices. By Theorem 2.1, we have  $\operatorname{reg}(S/J_{G_v}) \leq n-4$ .

The graph  $G \setminus v$  has n-1 vertices and at least two connected components, say  $G_1, \ldots, G_c$  with  $c \geq 2$ , because v is a cut vertex of G. Let  $S' = K[\{x_j, y_j\} : j \in [n] \setminus \{v\}]$ . Then

$$\frac{S'}{J_{G\setminus v}} \cong \frac{S_1}{J_{G_1}} \bigotimes_K \cdots \bigotimes_K \frac{S_c}{J_{G_c}}$$

where  $S_i = K[\{x_j, y_j\} : j \in V(G_i)]$  for i = 1, ..., c. This implies that

$$\operatorname{reg}(S/J_{G\setminus v} + (x_v, y_v)) = \operatorname{reg}(S'/J_{G\setminus v}) = \sum_{i=1}^{c} \operatorname{reg}(S_i/J_{G_i})$$

$$\leq \sum_{i=1}^{c} (|V(G_i)| - 1) = (n-1) - c \leq n-3.$$

If v has at least 4 neighbors in G, then the graph  $G_v \setminus v$  has a maximal clique with at least 4 vertices, thus, by Theorem 2.1, we get

$$reg(S/J_{G_v \setminus v} + (x_v, y_v))) = reg(S'/J_{G_v \setminus v}) \le (n-1) - 3 = n - 4.$$

Therefore, from inequality (9), we get  $reg(S/J_G) \le n-3$ .

**Lemma 3.4.** Let G be a connected indecomposable graph on  $n \ge 4$  vertices with the following properties:

- (a)  $J_G$  is unmixed;
- (b) G has a vertex v with exact two neighbors  $u_1, u_2$  and  $\{u_1, u_2\} \in E(G)$ . Then  $reg(S/J_G) \le n - 3$ .

*Proof.* If n = 4, then there are only two graphs which satisfy the condition (b), namely two triangles which share an edge and a triangle with an edge attached to one of its vertices; see Figure 1.




FIGURE 1. 4 vertices

The first graph does not satisfy the condition (a), while the second graph is decomposable. Thus, we may consider  $n \geq 5$ .

Let us consider an indecomposable graph G with  $n \geq 5$  vertices satisfying the conditions (a) and (b). We claim that  $\deg u_1 \geq 4$  or  $\deg u_2 \geq 4$ . Let us assume that this is not the case, thus  $\deg u_1 \leq 3$  and  $\deg u_2 \leq 3$ . Since G is indecomposable, it follows that  $\deg u_1 = 3$ ,  $\deg u_2 = 3$ , and there exists a path connecting  $u_1$  and  $u_2$  different from the edge  $\{u_1, u_2\}$  and the path  $u_1, v, u_2$ . But, in this case, the set

 $S = \{u_1, u_2\}$  is a cut set of G with c(S) = |S|, which is impossible since  $J_G$  is an unmixed ideal.

Without loss of generality, we may assume that  $\deg u_2 \geq 4$ .

We set  $e = \{u_1, v\}$ . By Proposition 2.3 (a), we have

(10) 
$$\operatorname{reg} \frac{S}{J_G} \le \max \left\{ \operatorname{reg} \frac{S}{J_{G \setminus e}}, \operatorname{reg} \frac{S}{J_{G \setminus e} : f_e} + 1 \right\}.$$

In the graph  $G \setminus e$ ,  $u_2$  is a cut vertex with at least 4 neighbors. Thus, by Lemma 3.3, it follows that

$$\operatorname{reg} \frac{S}{J_{G \setminus e}} \le n - 3.$$

Now we look at  $J_{G \setminus e}: f_e$ . By applying Theorem 2.4, we obtain

$$J_{G \setminus e} : f_e = J_{(G \setminus e)_e} + (x_{u_2}, y_{u_2})$$

since all the paths connecting  $u_1$  and v pass trough  $u_2$ . Therefore, since v becomes an isolated vertex in the graph  $(G \setminus e)_e \setminus u_2$ , we get

$$\operatorname{reg} \frac{S}{J_{G \setminus e} : f_e} = \operatorname{reg} \frac{S}{J_{(G \setminus e)_e} + (x_{u_2}, y_{u_2})} = \operatorname{reg} \frac{S'}{J_{(G \setminus e)_e \setminus \{u_2, v\}}}$$

where  $S' = K[\{x_j, y_j\} : j \in [n] \setminus \{u_2, v\}]$ . If the graph  $(G \setminus e)_e \setminus \{u_2, v\}$  is a path, as deg  $u_2 \geq 4$ , the graph G looks like in Figure 2, that is, there are some edges connecting  $u_2$  to some vertices of the the path  $(G \setminus e)_e \setminus \{u_2, v\}$  different from  $u_1$ . But then  $J_G$  is not unmixed since  $S = \{u_1, u_2\}$  is a cut set of G with c(S) = |S|, a contradiction. Therefore, the graph  $(G \setminus e)_e \setminus \{u_2, v\}$  is not a path. Thus, by Theorem 1.5, we obtain

$$\operatorname{reg} \frac{S}{J_{G \setminus e} : f_e} = \operatorname{reg} \frac{S'}{J_{(G \setminus e) \setminus \{u_2, v\}}} \le (n-2) - 2 = n-4,$$

which implies that

$$\operatorname{reg} \frac{S}{J_{G\backslash e}: f_e} + 1 \le n - 3.$$

and the proof of the lemma is completed.




FIGURE 2. The graph G when  $(G \setminus e)_e \setminus \{u_2, v\}$  is a path

We can now state the main result of this section.

**Theorem 3.5.** Let G be a connected graph on the vertex set [n]. Then the following statements are equivalent:

- (i)  $(J_G)_{\mathfrak{m}} \subset R$  is licci.
- (ii)  $J_G$  is Cohen-Macaulay and  $n-2 \leq \operatorname{reg}(S/J_G) \leq n-1$ .
- (iii) G is a path graph or it is isomorphic to one of the graphs depicted in Figure 3 where r, s, t are non-negative integers. In other words, G is a triangle with possibly some paths connected to some of its vertices.




Figure 3. Licci graphs

Proof. (i)  $\Rightarrow$  (ii). Let  $(J_G)_{\mathfrak{m}} \subset R$  be licci. By Theorem 1.1, it follows that  $\operatorname{reg}(S/J_G) \geq \operatorname{height}(J_G) - 1$ . Since  $J_G$  is Cohen-Macaulay, thus unmixed, we have  $\operatorname{height}(J_G) = \operatorname{height} P_{\emptyset}(G) = n - 1$ , by (2). Therefore, if G is connected and  $(J_G)_{\mathfrak{m}}$  is licci, then  $J_G$  is Cohen-Macaulay and  $\operatorname{reg}(S/J_G) \geq n - 2$ . But we know from [15] that  $\operatorname{reg}(S/J_G) \leq n - 1$ .

Let us prove that (ii)  $\Rightarrow$  (iii). Since, by Theorem 1.5, we have  $\operatorname{reg}(S/J_G) = n - 1$  if and only if G is a path graph, it remains to consider  $\operatorname{reg}(S/J_G) = n - 2$ . By using Lemma 3.2, we may reduce the problem to considering only the case when G is indecomposable. Therefore, in order to get (iii), by taking into account Lemma 3.2, it is enough to show that there is no indecomposable graph G with  $|V(G)| \geq 4$  such that  $J_G$  is Cohen-Macaulay and  $\operatorname{reg}(S/J_G) = n - 2$ . There is no such graph among those with 4 vertices. Thus, we may consider  $n = |V(G)| \geq 5$ .

Let us assume that such a graph does exist. By [1, Remark 5.3], since  $J_G$  is Cohen-Macaulay, the graph G must have a cut vertex, say v. Since G is indecomposable, v has at least 3 neighbors in G. If v has at least 4 neighbors, by Lemma 3.3, it follows that  $reg(S/J_G) \leq n-3$ , a contradiction. Thus, v has exactly 3 neighbors, say  $w, u_1, u_2$ . Since G is indecomposable and v is a cut vertex in G, it follows that none of the edges  $\{u_1, u_2\}, \{u_1, w\}, \{u_2, w\}$  belongs to E(G). On the other hand, as  $J_G$  is unmixed, the graph  $G \setminus v$  has exactly two connected components, say  $G_1$  and  $G_2$ . We may assume that  $u_1, u_2$  are vertices in  $G_1$  and w is a vertex in  $G_2$ . Let

 $e = \{v, w\}$ . By Proposition 2.3 (a), we have

(11) 
$$n-2 = \operatorname{reg} \frac{S}{J_G} \le \max \left\{ \operatorname{reg} \frac{S}{J_{G \setminus e}}, \operatorname{reg} \frac{S}{J_{G \setminus e} : f_e} + 1 \right\}.$$

We observe that  $G \setminus e$  has two connected components, namely G' with  $V(G') = V(G_1) \cup \{v\}$  and  $E(G') = E(G_1) \cup \{\{u_1, v\}, \{u_2, v\}\}\}$  and  $G'' = G_2$ . Obviously, G' is not a path graph since  $G_1$  is connected, thus there exists at least one path connecting  $u_1$  and  $u_2$  in  $G_1$  which does not contain v and is not the edge  $\{u_1, u_2\}$ . On the other hand, if  $G_2$  does not consist only of the isolated vertex w, then  $G_2$  cannot be a path since the graph G is indecomposable. Let  $S' = K[\{x_j, y_j\} : j \in V(G')]$  and  $S'' = K[\{x_i, y_i\} : j \in V(G'')]$ . Then, by Theorem 1.5, we have

$$\operatorname{reg} \frac{S'}{J_{G'}} + \operatorname{reg} \frac{S''}{J_{G''}} \le (|V(G')| - 2) + (|V(G'')| - 2) = n - 4.$$

Therefore,

$$\operatorname{reg} \frac{S}{J_{G \setminus e}} = \operatorname{reg} \frac{S'}{J_{G'}} + \operatorname{reg} \frac{S''}{J_{G''}} < n - 3.$$

If  $G_2$  consist only of the isolated vertex w, then we get

$$\operatorname{reg} \frac{S}{J_{G \setminus e}} = \operatorname{reg} \frac{S'}{J_{G'}} \le |V(G')| - 2 = n - 3.$$

Thus, in any case we have

(12) 
$$\operatorname{reg} \frac{S}{J_{G \setminus e}} \le n - 3.$$

Now we look at the term  $\operatorname{reg}(S/J_{G\setminus e}:f_e)$  of inequality (11). By Theorem 2.4, it follows that  $J_{G\setminus e}:f_e=J_{(G\setminus e)_e}$  since there is no path in G connecting v and w except the edge  $e=\{v,w\}$ . This is due to the fact that when we remove the cut vertex v from G, we get two connected components by the unmixedness of  $J_G$ . The graph  $(G\setminus e)_e$  consists as well of two connected components, say  $H_1$  which contains v and  $H_2$  which contains w. If  $H_2$  contains some other vertices together with w, then  $H_2$  cannot be a path since G is indecomposable. The component  $H_1$  is not a path since it contains at least the triangle with vertices  $u_1, u_2, v$ . Therefore, if  $S_i = K[\{x_j, y_j\}: j \in V(H_i)]$  for i = 1, 2, by Theorem 1.5, we obtain

$$\operatorname{reg} \frac{S}{J_{G \setminus e} : f_e} = \operatorname{reg} \frac{S}{J_{(G \setminus e)_e}} = \operatorname{reg} \frac{S_1}{J_{H_1}} + \operatorname{reg} \frac{S_2}{J_{H_2}} \le (|V(H_1)| - 2) + (|V(H_2)| - 2) = n - 4.$$

This inequality and (12) contradicts inequality (11).

It remains to analyze the case when  $H_2$  consists of the isolated vertex w. In this case we have

(13) 
$$\operatorname{reg} \frac{S}{J_{(G \setminus e)_e}} = \operatorname{reg} \frac{S_1}{J_{H_1}}.$$

We claim that  $H_1$  satisfies the conditions of Lemma 3.4. Clearly,  $H_1$  satisfies the condition (b). It remains to prove that  $J_{H_1}$  is an unmixed ideal because if  $H_1$  is decomposable in  $u_1$  or  $u_2$ , then G is decomposable, and this is impossible by our

hypotheses on G. We first observe that any non-empty cut set of  $H_1$  does not contain the vertex v which is a simplicial vertex in  $H_1$ . Let us assume that there exists a non-empty cut set  $S \subset V(H_1)$  such that  $c_{H_1}(S) \neq |S| + 1$ . The set S is obviously a cut set for the graph G as well. Moreover, if  $H_1, \ldots, H_{c_{H_1}(S)}$  are the connected components of the restriction of  $H_1$  to the vertex set  $V(H_1) \setminus S$ , with  $v \in V(H_1)$ , then the connected components of the restriction of G to  $V(G) \setminus S$  are  $H_1 \cup \{v, w\}, H_2, \ldots, H_{c_{H_1}(S)}$ . Hence  $c_G(S) = c_{H_1}(S) \neq |S| + 1$ , a contradiction to the unmixedness of  $J_G$ . Since  $H_1$  is a graph on n-1 vertices which satisfies the conditions of Lemma 3.4, we get  $\operatorname{reg}(S_1/J_{H_1}) \leq (n-1) - 3 = n-4$ . Thus, we have proved that

$$\operatorname{reg} \frac{S}{J_{G \setminus e} : f_e} = \operatorname{reg} \frac{S}{J_{(G \setminus e)_e}} = \operatorname{reg} \frac{S_1}{J_{H_1}} \le n - 4.$$

This inequality together with (12) contradicts inequality (11) and the proof of (ii)  $\Rightarrow$  (iii) is completed.

Finally, we prove the implication (iii)  $\Rightarrow$  (i).

As it was observed in the proof of [8, Proposition 3], if  $G = G_1 \cup G_2$  is a decomposable graph, then we have  $\operatorname{Tor}_i(S/J_{G_1}, S/J_{G_2}) = 0$  for all i > 0. In particular, it follows that  $J_{G_1}$  and  $J_{G_2}$  are transversal ideals in the sense of [11, Section 2]. Now, let  $G_1$  be a triangle with the vertices  $v_1, v_2, v_3$ . Then  $J_{G_1}$  is a Cohen-Macaulay ideal of height 2, thus it is licci by [19]. If we attach a path  $G_2$  to  $G_1$  in one of its vertices, say  $v_1$ , the resulting graph G is decomposable in  $v_1$  and  $J_{G_2}$  is a complete intersection ideal. According to [11, Theorem 2.6], it follows that  $(J_G)_{\mathfrak{m}}$  is a licci ideal. We repeat this argument by attaching a path in the vertex  $v_2$  to G and, next another path in the vertex  $v_3$ . In each step, we get a licci ideal.

An immediate consequence of the above theorem is the following.

Corollary 3.6. Let G be a connected bipartite graph. Then the ideal  $(J_G)_{\mathfrak{m}} \subset R = S_{\mathfrak{m}}$  is licci if and only if G is a path graph, or equivalently,  $J_G$  is a complete intersection.

We now turn to the disconnected graphs.

**Proposition 3.7.** Let G be a graph with the connected components  $G_1, \ldots, G_c$  where  $c \geq 2$ . Then  $(J_G)_{\mathfrak{m}} \subset R = S_{\mathfrak{m}}$  is licci if and only if either all the connected components of G are paths or one component of G is isomorphic to a graph of Figure 3 and all the other components are paths.

*Proof.* We first remark that, by [11, Theorem 2.6], if the components of G satisfy the conditions of the proposition, then  $(J_G)_{\mathfrak{m}}$  is licci since the ideals  $J_{G_i}$  are pairwise transversal by [4, Lemma 3.1].

For the converse, let  $(J_G)_{\mathfrak{m}}$  be a licci ideal. Then  $J_G$  is Cohen-Macaulay which implies that all the ideals  $J_{G_i}$  are Cohen-Macaulay and

$$\operatorname{reg}(S/J_G) \ge \operatorname{height}(J_G) - 1 = \operatorname{height}(J_{G_1}) + \dots + \operatorname{height}(J_{G_c}) - 1 = n - c - 1.$$

On the other hand, we have

$$reg(S/J_G) = \sum_{i=1}^{c} reg(S_i/J_{G_i}) \le \sum_{i=1}^{c} (|V(G_i)| - 1) = n - c.$$

Here  $S_i = K[\{x_j, y_j\} : j \in V(G_i)]$  for  $1 \leq i \leq c$ . The above inequalities imply that  $\operatorname{reg}(S/J_G) = n - c$  or  $\operatorname{reg}(S/J_G) = n - c - 1$ . In the first case, it follows that  $\operatorname{reg}(S_i/J_{G_i}) = |V(G_i)| - 1$  for all i, which implies that all the connected components of G are path graphs.

Let  $\operatorname{reg}(S/J_G) = n - c - 1$ . This means that for one of the connected components, say  $G_1$ , we have  $\operatorname{reg}(S_1/J_{G_1}) = |V(G_1)| - 2$  and all the other components of G are path graphs. Then, by Theorem 3.5, it follows that  $G_1$  is isomorphic to one of the graphs displayed in Figure 3.

# 4. LICCI BINOMIAL EDGE IDEALS OF CHORDAL GRAPHS

In this section we show that if we restrict to chordal graphs, we may relax the condition (ii) in Theorem 3.5, namely, we may ask that  $J_G$  is only unmixed instead of being Cohen-Macaulay. Before proving the main theorem of this section, we need a preparatory result. We first recall that for a graph G, c(G) denotes the number of maximal cliques of G, that is, the number of facets of the clique complex  $\Delta(G)$ .

**Lemma 4.1.** Let G be a connected chordal graph with n vertices. Then c(G) = n-2 if and only if the following conditions hold:

- (i) the maximal cliques of G have at most 3 vertices;
- (ii) G has at least one maximal clique with 3 vertices;
- (iii) G has exactly one maximal clique with 3 vertices or, for any two triangles  $F_1, F_2$  of  $\Delta(G)$ , there is a sequence of triangles  $F_1 = F_{i_1}, \ldots, F_{i_r} = F_2$  such that for any  $1 \leq j \leq r-1$ ,  $F_{i_j}$  and  $F_{i_{j+1}}$  share an edge.

*Proof.* Let c(G) = n - 2. Then (i) follows by [23, Proposition 3.1]. If G has no maximal clique with 3 vertices, then G is a tree, thus c(G) = n - 1, contradiction. Therefore, condition (ii) holds.

We prove (iii) by induction on n. Since G is chordal, by Dirac's theorem, we may order the facets of  $\Delta(G)$  as  $F_1, \ldots, F_c$  where c = c(G) such that  $F_i$  is a leaf of  $\langle F_1, \ldots, F_i \rangle$  for all i. If  $F_c$  is an edge, say  $F_c = \{v, w\}$  with deg w = 1, then the graph  $G \setminus w$  has n-1 vertices and n-3 cliques, thus, by induction, it satisfies (iii), and, consequently, G satisfies (iii) as well.

Let  $F_c$  be a triangle with the vertices u, v, w and assume that  $F_j$  with j < c is a branch of  $F_c$ . If  $F_j \cap F_c$  consists of just one vertex, say  $F_j \cap F_c = \{v\}$ , then the subgraph  $G' = G \setminus \{u, w\}$  has n-2 vertices and n-3 maximal cliques, therefore G' is a tree. This implies that  $\Delta(G)$  has exactly one facet with 3 elements, and condition (iii) is automatically fulfilled. Let us now assume that the branch  $F_j$  intersects  $F_c$  in the edge  $\{v, w\}$ . We consider the graph  $G \setminus u$ . This is a graph on n-1 vertices with n-3 maximal cliques, thus, by the inductive hypothesis, it satisfies (iii). Let us choose two triangles F, F' in  $\Delta(G)$ . If they are facets in  $\Delta(G \setminus u)$ , then they satisfy (iii). Otherwise, we may assume that  $F' = F_c$ . But then, by the inductive

hypothesis on  $G \setminus u$  there is a sequence of triangles  $F = F_{i_1}, \ldots, F_{i_r} = F_j$  such that for any  $1 \leq s \leq r-1$ ,  $F_{i_s}$  and  $F_{i_{s+1}}$  share an edge. Then the sequence  $F = F_{i_1}, \ldots, F_{i_r} = F_j, F_{i_{r+1}} = F_c$  satisfies the required condition for G.

For the converse, let us assume that G is a connected chordal graph with n vertices, which satisfies the three conditions of the statement. By condition (ii) and [23, Proposition 3.1], it follows that  $c(G) \leq n-2$ .

Let us assume that there exists a connected chordal graph G satisfying conditions (i)–(iii) and such that c(G) < n-2 and choose one with the minimal number of vertices. We consider again the leaf order  $F_1, \ldots, F_c$  on the facets of  $\Delta(G)$  and take  $F_j$  with j < c a branch of  $F_c$ . If  $F_c$  is an edge,  $F_c = \{v, w\}$  with deg w = 1, then the graph  $G \setminus w$  has n-1 vertices and satisfies conditions (i)–(iii), thus, by our assumption on G we have  $c(G \setminus w) = n-3$ , which implies that c(G) = n-2, contradiction.

If  $F_c$  is a triangle,  $F_c = \{u, v, w\}$ , and  $F_j$  intersects  $F_c$  in just one vertex, say v, then we have the following cases.

Case 1. The facet  $F_c$  is the only triangle in  $\Delta(G)$ . Then, the subgraph  $G \setminus \{u, w\}$  is a tree on n-2 vertices, thus  $\Delta(G \setminus \{u, w\})$  has n-3 maximal cliques, which implies that c(G) = n-2, contradiction.

Case 2. There exists a triangle  $F \in \Delta(G \setminus \{u, w\})$ . Then, as G satisfies condition (iii), there exists a triangle  $F' \neq F_c$  which intersects  $F_c$  along an edge. But this is impossible since the branch  $F_j$  intersects  $F_c$  in one vertex.

Finally, we have to consider that  $F_j$  shares an edge with  $F_c$ , say  $F_j \cap F_j = \{v, w\}$ . Since  $F_j$  is a branch of  $F_c$ , there is no other facet F of  $\Delta(G)$  with  $F \cap F_c = \{u, w\}$  or  $F \cap F_c = \{u, v\}$ . Then the graph  $G \setminus u$  obviously satisfies conditions (i)–(iii) and has n-1 vertices. By the choice of G, we have  $c(G \setminus u) = n-3$ , thus c(G) = n-2, contradiction.

**Theorem 4.2.** Let G be a connected chordal graph on the vertex set [n]. Then the following statements are equivalent:

- (i)  $(J_G)_{\mathfrak{m}} \subset R$  is licci.
- (ii)  $J_G$  is Cohen-Macaulay and  $n-2 \leq \operatorname{reg}(S/J_G) \leq n-1$ .
- (iii)  $J_G$  is unmixed and  $n-2 \le \operatorname{reg}(S/J_G) \le n-1$ .
- (iv) G is a path graph or it is isomorphic to a graph depicted in Figure 3.

*Proof.* We have to prove only the implication (iii)  $\Rightarrow$  (iv). Let  $J_G$  be unmixed and let  $\operatorname{reg}(S/J_G) = n-1$ . Then, by Theorem 1.5, G is a path graph. Let us now discuss the case when  $\operatorname{reg}(S/J_G) = n-2$ . By [23, Theorem 3.5], we have  $\operatorname{reg}(S/J_G) \leq c(G)$ . Thus, we get  $c(G) \geq n-2$ . If c(G) = n-1, then G is a tree, but since  $J_G$  is unmixed, by [3, Corollary 1.2], it follows that G is a path graph.

As in the proof of Theorem 3.5, it is enough to show that there is no indecomposable chordal graph with  $n \geq 4$  vertices which satisfies the conditions  $J_G$  unmixed and  $\operatorname{reg}(S/J_G) = c(G) = n - 2$ . Let us assume that such a graph G does exists.

By Theorem 2.1, it follows that the maximal cliques of G have at most three vertices. As G is a chordal graph, by Dirac's theorem, it follows that the facets of the clique complex  $\Delta(G)$  of G have a leaf order, say  $F_1, \ldots, F_{n-2}$ . In particular, this means that  $F_{n-2}$  has a branch. Let  $F_j$  with  $j \leq n-3$  be a branch of  $F_{n-2}$ .

Case 1. Assume that the intersection  $F_j \cap F_{n-2}$  consists of only one vertex of G, say  $F_j \cap F_{n-2} = \{v\}$ . If  $F_{n-2}$  has only the branch  $F_j$ , then G is decomposable which contradicts our assumption on G. Thus  $F_{n-2}$  has  $q \geq 2$  branches, say  $F_{j_1}, \ldots, F_{j_q}$ . Then, as  $J_G$  is unmixed, it follows that the induced subgraph of  $G \setminus v$  on the vertex set  $\bigcup_{i=1}^q F_{j_i} \setminus v$  is connected. This implies that all the facets  $F_{j_1}, \ldots, F_{j_q}$  are triangles. If  $F_{n-2}$  is also a triangle, we get a contradiction to Lemma 4.1. Thus,  $F_{n-2}$  must be an edge and then v is a cut vertex of G with  $\deg_G(v) \geq 4$ . By Lemma 3.3, it follows that  $\operatorname{reg}(S/J_G) \leq n-3$ , a contradiction.

Case 2. Assume that the intersection  $F_j \cap F_{n-2}$  consists of two vertices of G, say  $F_j \cap F_{n-2} = \{v, w\}$ . In this case,  $F_{n-2}$  is a triangle with the vertices u, v, w. Since  $J_G$  is unmixed, there must be other facets of  $\Delta(G)$  whose intersection with  $F_{n-2}$  is contained in  $\{v, w\}$  or equal to  $\{v, w\}$ . Let  $F_{j_1}, \ldots, F_{j_q}$  with  $q \geq 2$  and  $j_q = j$  be the facets of  $\Delta(G)$  with  $F_{j_s} \cap F_{n-2} \subseteq \{v, w\}$  for  $1 \leq s \leq q$ . As v is not a simplicial vertex in G, we may apply again [17, Lemma 4.8] and get

$$J_G = J_{G_v} \cap (J_{G \setminus v} + (x_v, y_v)).$$

We use the following exact sequence of S-modules:

$$0 \to \frac{S}{J_G} \to \frac{S}{J_{G_v}} \bigoplus \frac{S}{J_{G \setminus v} + (x_v, y_v)} \to \frac{S}{J_{G_v \setminus v} + (x_v, y_v)} \to 0.$$

to derive that

(14) 
$$\operatorname{reg} \frac{S}{J_G} \le \max\{\operatorname{reg} \frac{S}{J_{G_v}}, \operatorname{reg} \frac{S}{J_{G \setminus v} + (x_v, y_v)}, \operatorname{reg} \frac{S}{J_{G_v \setminus v} + (x_v, y_v)} + 1\}.$$

By [23, Lemma 3.4], it follows that  $c(G_v) \leq c(G) - q$ , hence, by our assumption on q, we get  $c(G_v) \leq n-4$ . On the other hand, by [23, Lemma 3.3], we have  $c(G_v \setminus v) \leq c(G_v)$ , thus  $c(G_v \setminus v) \leq n-4$ . In particular, it follows that

(15) 
$$\operatorname{reg}(S/J_{G_v}) \le n - 4 \text{ and } \operatorname{reg}(S/J_{G_v \setminus v} + (x_v, y_v)) \le n - 4.$$

Therefore, by (14), we must have

$$\operatorname{reg} \frac{S}{J_{G \setminus v} + (x_v, y_v)} = \operatorname{reg} \frac{S'}{J_{G \setminus v}} \ge n - 2.$$

where  $S' = K[\{x_j, y_j\} : j \in [n] \setminus \{v\}]$ . As  $G \setminus v$  has n-1 vertices, it follows by Theorem 1.5 that  $G \setminus v$  is a path graph. But in this case,  $\mathcal{S} = \{v, w\}$  is a cut set of G because G is indecomposable. In addition, the restriction of G to the vertex set  $[n] \setminus \{v, w\}$  has two connected components, which is a contradiction to the unmixedness of  $J_G$ .

## References

- [1] D. Bolognini, A. Macchia, F. Strazzanti, *Binomial edge ideals of bipartite graphs*, European J. Combin. **70** (2018), 1–25.
- [2] G. A. Dirac, On rigid circuit graphs, Abh. Math. Semin. Univ. Hambg. 38 (1961), 71âÅ?6.
- [3] V. Ene, J. Herzog, T. Hibi, Cohen-Macaulay binomial edge ideals, Nagoya Math. J. **204** (2011), 57–68
- [4] H. T. Hà, H. D. Nguyen, N. V. Trung, T. N. Trung, Symbolic powers of sums of ideals, Math. Z. https://doi.org/10.1007/s00209-019-02323-8

- [5] J. Herzog, T. Hibi, Monomial Ideals, Grad. Texts in Math. 260, Springer, London, 2010.
- [6] J. Herzog, T. Hibi, F. Hreinsdotir, T. Kahle, J. Rauh, Binomial edge ideals and conditional independence statements, Adv. Appl. Math. 45 (2010), 317–333.
- [7] J. Herzog, T. Hibi, H. Ohsugi, Binomial ideals, Grad. Texts in Math. 279, Springer, London, 2018.
- [8] J. Herzog, G. Rinaldo, On the extremal Betti numbers of binomial edge ideals of block graphs, Electron. J. Combin. **25**(1) (2018), #P1.63
- [9] C. Huneke, Linkage and Koszul homology of ideals, Amer. J. Math. 104 (1982), 1043–1062.
- [10] C. Huneke, B. Ulrich, The structure of linkage, Ann. of Math. 126 (1987), 277–334.
- [11] M. R. Johnson, Linkage and sums of ideals, Trans. Amer. Math. Soc. 350 (5) (1998), 1913– 1930.
- [12] D. Kiani, S. Saeedi Madani, Some Cohen-Macaulay and unmixed binomial edge ideals, Comm. Algebra 43 (2015), 5434–5453.
- [13] D. Kiani, S. Saeedi Madani, *The Castelnuovo–Mumford regularity of binomial edge ideals*, J. Combin. Theory Serie A **139** (2016), 80–86.
- [14] K. Kimura, N. Terai, K. Yoshida, Licci squarefree monomial ideals generated in degree two or with deviation two, J. Algebra **390** (2013), 264–289.
- [15] K. Matsuda, S. Murai, Regularity bounds for binomial edge ideals, J. Commut. Algebra 5(1), 141–149 (2013).
- [16] F. Mohammadi, L. Sharifan, Hilbert function of binomial edge ideals, Comm. Algebra 42 (2014), 688–703.
- [17] M. Ohtani, Graphs and ideals generated by some 2-minors, Comm. Algebra, 39 (2011), 905–917.
- [18] I. Peeva, Graded syzygies, Algebr. Appl. 14 Springer, 2010.
- [19] C. Peskine, L. Szpiro, Liaison des variétés algébriques, I, Invent. Math. 26 (1974), 271–302.
- [20] A. Rauf, G. Rinaldo, Construction of Cohen-Macaulay binomial edge ideals, Comm. Algebra 42 (2014), 238–252.
- [21] G. Rinaldo, Cohen-Macaulay binomial edge ideals of small deviation, Bull. Math. Soc. Sci. Math. Roumanie **56**(104)(4) (2013), 497–503.
- [22] G. Rinaldo, Cohen-Macaulay binomial edge ideals of cactus graphs, J. Algebra Appl. 18 No. 4 (2019) 1950072, pp. 1-18.
- [23] M. Rouzbahani Malayeri, S. Saeedi Madani, D. Kiani, Regularity of binomial edge ideals of chordal graphs, arXiv:1810.03119
- [24] S. Saeedi Madani, *Binomial Edge Ideals: A Survey*, in Multigraded Algebras and Applications (V. Ene, E. Miller Eds.) Springer Proceedings in Mathematics & Statistics (2018), 83–94.

VIVIANA ENE, FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, OVIDIUS UNIVERSITY, BD. MAMAIA 124, 900527 CONSTANTA, ROMANIA

E-mail address: vivian@univ-ovidius.ro

GIANCARLO RINALDO, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TRENTO, VIA SOM-MARIVE, 14 38123 POVO (TRENTO), ITALY

E-mail address: giancarlo.rinaldo@unitn.it

NAOKI TERAI, FACULTY OF EDUCATION, SAGA UNIVERSITY, SAGA 840-8502, JAPAN *E-mail address*: terai@cc.saga-u.ac.jp