Maximising the number of induced cycles in a graph^{*}

Natasha Morrison[†] Alex Scott[†]

April 13, 2017

Abstract

We determine the maximum number of induced cycles that can be contained in a graph on $n \ge n_0$ vertices, and show that there is a unique graph that achieves this maximum. This answers a question of Chvátal and Tuza from the 1980s. We also determine the maximum number of odd or even induced cycles that can be contained in a graph on $n \ge n_0$ vertices and characterise the extremal graphs. This resolves a conjecture of Chvátal and Tuza from 1988.

1 Introduction

What is the maximum number of induced cycles in a graph on n vertices? For cycles of fixed length, this problem has been extensively studied. Indeed, for any fixed graph H, let the *induced density* of H in a graph G be the number of induced copies of H in G divided by $\binom{|G|}{|H|}$; let I(H; n) be the maximum induced density of H over all graphs G on n vertices; and let the *inducibility* of H be the limit $\lim_{n\to\infty} I(H; n)$. In 1975, Pippinger and Golumbic [12] made the following conjecture.

Conjecture 1.1. [12] For $k \ge 5$, the inducibility of the cycle C_k is $k!/(k^k - k)$.

Balogh, Hu, Lidický and Pfender [2] recently proved this conjecture in the case k = 5 via a flag algebra method, and showed that the maximum density was achieved by a unique graph. Apart from this case, the problem remains open (though see [3, 4, 5, 6, 7, 8, 9] for results on inducibility of other graphs).

In this paper, we consider the total number of induced cycles, without restriction on length. This problem was raised in the 1980s by Chvátal and Tuza (see [15] and [16]), who asked for the maximum possible number of induced cycles in a graph with n vertices. The problem was investigated independently in unpublished work of Robson, who showed in the

^{*}Work partially done during the 2015 Barbados Workshop on Structural Graph Theory. The attendance of the first author was possible due to the generous support of Merton College Oxford.

[†]Mathematical Institute, University of Oxford, Woodstock Road, Oxford, OX2 6GG, United Kingdom. E-mail: {morrison,scott}@maths.ox.ac.uk.

1980s that a graph on n vertices has at most $3^{(1+o(1))n/3}$ induced cycles ([10, 13]). Tuza also raised a second, closely related problem on induced cycles. In 1988 he conjectured with Chvátal (see [14], [15] and [16]) that the maximum possible number of odd induced cycles in a graph on n vertices is $3^{n/3}$.

In this paper we resolve both problems, proving exact bounds for all sufficiently large n, and determining the extremal graphs. Our methods work for a number of problems of this type: thus we will determine, for sufficiently large n, the graphs with n vertices that maximize the number of induced cycles (Theorem 1.2); we will also determine the graphs with the maximum number of even induced cycles, the maximum number of odd induced cycles (Theorem 1.6), and in Theorem 1.7, the maximum number of odd holes (i.e. induced odd cycles of length at least 5).

In order to state our results, it is helpful to have a couple of definitions. As usual, for G a graph define the *neighbourhood* of x to be $N_G(x) := \{y \in V(G) : xy \in E(G)\}$. A graph B is called a *cyclic braid* if there exists $k \geq 3$ and a partition B_1, \ldots, B_k of V(B) such that for every $1 \leq i \leq k$ and every $x \in B_i$, we have $B_{i-1} \cup B_{i+1} \subseteq N_B(x) \subseteq B_{i-1} \cup B_i \cup B_{i+1}$ where indices are taken modulo k. For such a partition, the notation $B = (B_1, \ldots, B_k)$ is used. The sets B_i are called *clusters* of B; the *length* of the cyclic braid is the number of clusters. If a cyclic braid contains no edges within its clusters, it is called an *empty cyclic braid*. Observe that an empty cyclic braid is k-partite and, when k > 3, is triangle free. If a cyclic braid contains every possible edge within each cluster, then it is called *full*. A pair of clusters B_1 and B_2 are *adjacent* in G if $v_1v_2 \in E(G)$ for all $v_1 \in B_1$ and $v_2 \in B_2$. A triple of clusters B_1, B_2, B_3 are *consecutive* if B_1 is adjacent to B_2 and B_2 is adjacent to B_3 .

As it turns out, the structure of the extremal graph depends on the value of n modulo 3. For $n \ge 8$ define an n-vertex graph H_n separately for each value of n modulo 3. Let $k \ge 3$. Define H_{3k} to be the empty cyclic braid of length k where every cluster has size 3. Define H_{3k+1} to be the empty cyclic braid containing k-1 clusters of size 3 and one of size 4. Finally, define H_{3k-1} to be the empty cyclic braid containing k-1 clusters of size 3 and one of size 2.

Let m(n) be the maximum number of induced cycles that can be contained in a graph on *n* vertices. The main result of our paper is the following.

Theorem 1.2. There exists n_0 such that, for all $n \ge n_0$, H_n is the unique graph on n vertices containing m(n) induced cycles.

More precisely,

Corollary 1.3. There exists n_0 such that, for all $n \ge n_0$:

$$m(n) = \begin{cases} 3^{n/3} + 12n & \text{for } n \equiv 0 \mod 0 \ 3; \\ 4 \cdot 3^{(n-4)/3} + 12n + 51 & \text{for } n \equiv 1 \mod 0 \ 3; \\ 2 \cdot 3^{(n-2)/3} + 12n - 36 & \text{for } n \equiv 2 \mod 0 \ 3. \end{cases}$$

Corollary 1.3 implies that $m(n) = \Theta(3^{n/3})$.

A hole is an induced cycle of length at least 4. For $n \ge 10$, the graph H_n is triangle free and each induced cycle is a hole, so Theorem 1.2 implies the following. **Corollary 1.4.** There exists n_0 such that, for all $n \ge n_0$, H_n is the unique graph on n vertices with the maximum number of holes.

Using similar arguments to those in the proof of Theorem 1.2 we also prove a stability-type result.

Theorem 1.5. Fix $0 < \alpha < 1$. There exists a constant $C = C(\alpha)$ and $n_0 = n_0(\alpha)$ such that for any $n \ge n_0$, if a graph F on n vertices contains at least $\alpha \cdot m(n)$ induced cycles, then by adding or deleting edges incident to at most $C(\alpha)$ vertices of F, the graph F can be transformed into a cyclic braid with the same cluster sizes as H_n .

The arguments used to prove Theorem 1.2 can be adapted to give results about other sets of induced cycles, for instance induced cycles of given parity. We say that a path or cycle is *odd* if it contains an odd number of vertices (*even* if it contains an even number).

Let $m_o(n)$ be the maximum number of induced odd cycles that can be contained in a graph on *n* vertices. The value of $m_o(n)$ and the structure of the extremal graphs depend on the value of *n* modulo 6.

Define G_n to be the full cyclic braid on n vertices whose clusters all have size 3 except for:

- three consecutive clusters of size 2, when $n \equiv 0$ modulo 6;
- two adjacent clusters of size 2, when $n \equiv 1 \mod 6$;
- one cluster of size 2, when $n \equiv 2 \mod 6$;
- one cluster of size 4, when $n \equiv 4 \mod 6$;
- two adjacent clusters of size 4, when $n \equiv 5 \mod 6$.

We will prove the following.

Theorem 1.6. There exists n_0 such that, for all $n \ge n_0$, G_n is the unique n-vertex graph containing $m_o(n)$ induced odd cycles.

This resolves the conjecture of Chvátal and Tuza (see [14], [15] and [16]), showing that $m_o(n)$ is within an O(n) additive term of the conjectured $3^{n/3}$ when $n \equiv 3$ modulo 6 and within a constant factor when $n \not\equiv 3$ modulo 6.

If we consider odd holes (induced odd cycles of length at least 5), we get the same bound but a larger family of extremal graphs. Let $m_o^h(n)$ be the maximum number of odd holes that can be contained in a graph on n vertices. Define \mathcal{G}_n to be the family of cyclic braids on nvertices whose cluster sizes are the same as the cluster sizes in G_n , but with no restrictions on which clusters are adjacent or consecutive, or on which edges are present inside the clusters; in addition, when $n \equiv 5$ modulo 6, we also include the cyclic braids whose clusters all have size 3 except for four clusters of size 2.

A modification of the proof of Theorem 1.6 gives the following.

Theorem 1.7. There exists n_0 such that, for all $n \ge n_0$, the family of n-vertex graphs that contain $m_o^h(n)$ odd holes is \mathcal{G}_n .

Let $m_e(n)$ be the maximum number of induced even cycles that can be contained in a graph on n vertices, and define E_n to be the empty cyclic braid on n vertices whose clusters all have size 3 except for:

- one cluster of size 4, when $n \equiv 1 \mod 6$;
- two adjacent clusters of size 4, when $n \equiv 2 \mod 6$;
- three consecutive clusters of size 2, when $n \equiv 3 \mod 6$;
- two adjacent clusters of size 2, when $n \equiv 4 \mod 6$; and
- one cluster of size 2, when $n \equiv 5 \mod 6$.

The proof of Theorem 1.6, adapted to consider even rather than odd induced cycles, gives the following.

Theorem 1.8. There exists n_0 such that, for all $n \ge n_0$, E_n is the unique n-vertex graph containing $m_e(n)$ induced even cycles.

As in the case of $m_o(n)$, we have $m_e(n) = \Theta(3^{n/3})$.

The paper is structured as follows. In Section 2 we prove a preliminary result (Theorem 2.1) determining the structure of the *n*-vertex graphs that maximise the number of induced paths between a particular pair of vertices. During the proof of this result we will introduce several of the key ideas needed later. The proof of the main theorem (Theorem 1.2, maximising the number of induced cycles) is given in Section 3. The proof uses Theorem 2.1 from Section 2, but otherwise is entirely contained in Section 3. A number of lemmas proved in Section 3 are proved in more generality than is strictly needed. This is because the more general versions will be used in Section 4, where we prove Theorem 1.5. The proofs of Theorem 1.6, Theorem 1.7 and Theorem 1.8 are given in Section 5. However, all the key ideas for the proofs are the same as those in the proof of Theorem 1.2. Finally, in Section 6 we conclude by discussing some open questions.

2 Induced paths between a pair of vertices

Let G be a finite graph and let x and y be distinct vertices in V(G). Define $p_2(G; x, y)$ to be the number of induced paths in G beginning at x and ending at y. Also define:

$$p_2(G) := \max\{p_2(G; x, y) : x, y \in V(G)\},\$$

and

$$p_2(n) := \max\{p_2(G) : |V(G)| = n\}.$$

We use the notation $p_2(\cdot)$, as it indicates that we are counting the maximum number of induced paths between *two* fixed vertices.

Our first goal in this section is to determine the structure of the *n*-vertex graphs that contain $p_2(n)$ induced paths between some pair of vertices. We show that these extremal graphs have a particular structure that depends on the value of *n* modulo 3. We then prove analogous results for odd and even length paths. The strategy and notation used in this section for paths are later developed for induced cycles in Section 3.

Let F be a graph and let B_1, \ldots, B_k be disjoint subsets of V(F). A subgraph $B \subseteq F$ is a braid in F if there exists $k \geq 2$ and a partition B_1, \ldots, B_k of V(B) such that for each $2 \leq i \leq k-1$ and for every $x \in B_i$ we have $B_{i-1} \cup B_{i+1} \subseteq N_F(x) \subseteq B_{i-1} \cup B_i \cup B_{i+1}$. If $V(F) = \bigcup_{i=1}^k B_i$, we say that F is a braid. For such a partition, the notation $B = (B_1, \ldots, B_k)$ is used. The sets B_i are called *clusters*. If $i \in \{1, k\}$ we say B_i is an *end cluster*; otherwise we say B_i is a *central cluster*. The *length* of a braid is the number of clusters it contains.

Let $n \geq 4$. Define \mathcal{F}_n to be the set of all braids B with the following properties:

- |V(B)| = n.
- *B* has end clusters of size one.
- If $n \equiv 0$ modulo 3, then either one central cluster has size 4 and the rest have size 3, or two have size two and the rest have size 3.
- If $n \equiv 1$ modulo 3, then one central cluster has size 2 and the rest have size 3.
- If $n \equiv 2 \mod 3$, all clusters have size 3.

Observe that there are no conditions on whether clusters in the braid contain edges. See Figure 1 for an example of a braid in \mathcal{F}_{10} .




Figure 1: An example of a braid in \mathcal{F}_{10} .

Let the end clusters be $\{x\}$ and $\{y\}$. Every graph in \mathcal{F}_n contains the same number of induced paths between x and y and so we define

$$f_2(n) = \begin{cases} 3^{(n-2)/3} & \text{for } n \equiv 2 \text{ modulo } 3\\ 4 \cdot 3^{(n-6)/3} & \text{for } n \equiv 0 \text{ modulo } 3\\ 2 \cdot 3^{(n-4)/3} & \text{for } n \equiv 1 \text{ modulo } 3 \end{cases}$$

and observe that $p_2(F) = p_2(F; x, y) = f_2(n)$ for all $F \in \mathcal{F}_n$.

In this section we prove the following.

Theorem 2.1. Let G be a finite graph on $n \ge 4$ vertices and let x and y be distinct vertices of G. Suppose that G contains $p_2(n)$ induced paths between x and y. Then G is isomorphic to a graph in \mathcal{F}_n with end clusters $\{x\}$ and $\{y\}$.

In particular, this gives the following.

Corollary 2.2. For all $n \ge 4$, we have $p_2(n) = f_2(n)$.

Before proving Theorem 2.1, we introduce some preliminary notation and definitions.

Definition 2.3. Define $N[v] := N(v) \cup \{v\}$. Also, for a set $X \subseteq V(G)$, let $N(X) := \bigcup_{x \in X} N(x)$, and $N[X] := \bigcup_{x \in X} N[x]$. Note that $X \subseteq N[X]$, and $X \cap N(X)$ may or may not be empty. For a subgraph $H \subseteq G$, define N(H) := N(V(H)) and N[H] := N[V(H)].

In order to prove Theorem 2.1 (counting induced paths), the following definition is used. A similar definition is given in Definition 3.6 to prove Theorem 1.2 (counting induced cycles).

Definition 2.4. Let G be a finite graph and fix $x, y \in V(G)$, with $y \notin N[x]$. The x-y-path tree of G is a tree T = T(x, y) together with a function $t : V(T) \to V(G)$ defined as follows.

- T is a tree with vertex set V(T) disjoint from V(G).
- The vertices of T correspond to induced paths $P := x, x_1 \dots, x_j$ in G such that $y \notin N_G(\{x_1, \dots, x_{j-2}\})$ and $y \in N(x_{j-1})$ only if $x_j = y$. For every such P define a vertex w_P in T, and set $t(w_P) = x_j$. We say that P is the G-path of w_P . These vertices are the only vertices in T. Define the root of the tree to be $v_0 := w_x$.
- Given a vertex $w \in V(T)$ with *G*-path x, x_1, \ldots, x_j we define C(w), the *children* of w, to be the set of vertices in T whose *G*-path is x, x_1, \ldots, x_j, z , for some $z \in G$. Define $N_T(v_0) = C(v_0)$. For $w \in V(T) \setminus \{v_0\}$ define $N_T(w) := C(w) \cup \{u\}$ where u is the unique vertex in T with *G*-path x, x_1, \ldots, x_{j-1} . (So two vertices are adjacent in T precisely when one of their *G*-paths extends the other by one vertex.)

We write $t(S) := \{t(s) : x \in S\}$ for any subset $S \subseteq V(T)$ and $t(H) := G[\{t(x) : x \in V(H)\}]$ for any subgraph $H \subseteq T$. Given a set $S \subseteq V(T)$, we say that S sees a vertex $w \in V(G)$ (or w is seen by S) if $w \in N_G[t(S)]$. An empty set does not see any vertex. If $w \notin N_G[t(S)]$, we say that w is unseen by S (or that S does not see w).

See Figure 2 for an example of an x-y-path tree. We get the following proposition as an easy consequence of Definition 2.3.

Proposition 2.5. Let G be a graph containing vertices x and y such that $y \notin N[x]$. Let T be the x-y-path tree of G rooted at v_0 . Let P be a path in T starting at v_0 . If V(P) sees a vertex $w \in V(G)$, then there exists a unique $u \in N_T[V(P)]$ such that t(u) = w.




Figure 2: A graph and its x-y-path tree. Each vertex w in the tree is labelled with t(w).

Proof. This follows immediately from the construction of T: u is a child in T of the first vertex v in P such that t(v) is adjacent to w in G.

The following terminology will also be used later during the proof of Theorem 1.2 (maximising induced cycles), as well as in this section for the proof of Theorem 2.1 (maximising induced paths between a pair of vertices).

Definition 2.6. Let G be a graph containing vertices x and y such that $y \notin N[x]$. Let T be the x-y-path tree of G. For any $z \in V(T)$, z is the child of a unique vertex w. Define B(z), the branch rooted at z, to be the component of $T \setminus \{wz\}$ containing z. Also, define L(z) to be the number of leaves of T that are contained in B(z) and define $L_y(z)$ to be the number of leaves l in T contained in B(z) such that t(l) = y. If it is unclear which tree we are considering, we will write $B_T(z)$, $L_T(z)$, etc.

Observe that for $w \in T$, t(w) = y only if w is a leaf. It directly follows from Definition 2.3 that

$$p_2(G; x, y) = L_y(v_0).$$
 (2.1)

In order to prove Theorem 2.1, we use the following lemma about x-y-path trees. For $w \in V(T)$, we write D(w) := |C(w)|.

Lemma 2.7. Let G be a graph on $n \ge 4$ vertices. Let x and y be distinct vertices in V(G), with $y \notin N[x]$ and $p_2(G; x, y) > 0$. Let T be the x-y-path tree rooted at v_0 and $P := v_0, \ldots, v_k$ be any path in T where v_k is a leaf. Then:

- (i) $L_y(v_0) \le f_2(n)$.
- (*ii*) If $L_y(v_0) = f_2(n)$, then:
 - (a) for any v_j and for all $u, w \in C(v_j)$, we have $L_y(u) = L_y(w)$;
 - (b) $V(P) \setminus \{v_k\}$ sees every vertex of G and $t(v_k) = y$.

Proof. Sequentially choose a path $v_0, v_1, \ldots, v_k \subseteq V(T)$, where v_k is a leaf. At vertex v_j we choose v_{j+1} to be some $z \in C(v_j)$ such that $L_y(z) = \max\{L_y(x) : x \in C(v_j)\}$. Let \mathcal{P} be the set of paths that can be obtained in this manner and fix $P := v_0, \ldots, v_k \in \mathcal{P}$.

We first show that $t(v_k) = y$. Suppose otherwise that $t(v_k) \neq y$. As $p_2(G; x, y) > 0$, by construction of P we have $L_y(v_{k-1}) > 0$. If v_{k-1} had a child u with t(u) = y, by construction of T this would be the only child of v_{k-1} . Thus v_{k-1} has no such child. This fact, along with the fact $L_y(v_{k-1}) > 0$ implies that v_{k-1} has a child z with $L_y(z) > 0$. As $L_y(v_k) = 0$, we have $L_y(z) > L_y(v_k)$, a contradiction.

For $0 \le i \le k - 1$, we have

$$L_y(v_i) = \sum_{z \in C(v_i)} L_y(z) \le D(v_i) \max\{L_y(z) : z \in C(v_i)\} = D(v_i)L_y(v_{i+1}).$$
(2.2)

By repeatedly applying (2.2) we get

$$L_y(v_0) \le D(v_0) \max\{L_y(z) : z \in C(v_0)\} \le \ldots \le L_y(v_{k-1}) \prod_{i=0}^{k-2} D(v_i).$$
(2.3)

As $t(v_k) = y$, v_k is the only child of v_{k-1} (by construction of T) and $L_y(v_{k-1}) = 1$. Thus

$$L_y(v_0) \le \prod_{i=0}^{k-2} D(v_i),$$
 (2.4)

where $\sum_{i=0}^{k-2} D(v_i) \leq n-2$, as v_0, \ldots, v_{k-2} have disjoint sets of children in $G \setminus \{x, y\}$ by Propostion 2.5.

A quick check shows that the maximal value of $\prod_{i=0}^{k-2} D(v_i)$ subject to $\sum_{i=0}^{k-2} D(v_i) \le n-2$ occurs only in the following cases:

- If $n \equiv 2$ modulo 3, we have $D(v_i) = 3$ for all *i*.
- If $n \equiv 0$ modulo 3, we have either $D(v_i) = 4$ for exactly one *i* and $D(v_j) = 3$ for all $j \neq i$; or there are i_1, i_2 such that $D(v_i) = 2$ for $i = i_1, i_2$, and $D(v_j) = 3$ for all $i \notin \{i_1, i_2\}$.
- If $n \equiv 1$ modulo 3, we have $D(v_i) = 2$ for exactly one *i*, and $D(v_j) = 3$ for all $j \neq i$.

Thus we see that the maximal possible value of $\prod_{i=0}^{k-2} D(v_i)$ is $f_2(n)$, and so $L_y(v_0) \leq f_2(n)$ as required for (i).

When $L_y(v_0) = f_2(n)$ we have

$$\prod_{i=0}^{k-2} D(v_i) = f_2(n).$$
(2.5)

This is only possible if $\sum_{i=0}^{k-2} D(v_i) = n-2$ and the $D(v_i)$ take the values defined in the above cases. In addition, we have equality in (2.4) and hence in (2.2) for each value of $0 \le i \le k-1$. Therefore, for each *i* and for all $z, w \in C(v_i)$, we have $L_y(z) = L_y(w)$.

Suppose there exists a path $X := x_0, \ldots, x_k$, where $x_0 = v_0$ and x_k is a leaf, such that $X \notin \mathcal{P}$. We will derive a contradiction and hence conclude that every such path is in \mathcal{P} .

Choose $P' := y_0, \ldots, y_k \in \mathcal{P}$ so that it coincides with X on the longest possible initial segment, i.e., so that *i* is maximal such that $y_0, \ldots, y_i = x_0, \ldots, x_i$. As $X \notin \mathcal{P}$, for some *j* we have $L_y(x_j) \neq L_y(y_j)$, but $x_i = y_i$ for i < j. But by the argument of the previous paragraph, as $P' \in \mathcal{P}$, we have that for each *i*, $L_y(z) = L_y(w)$ for all $z, w \in C(y_i)$. Thus as $x_{j-1} = y_{j-1}$, we have $x_j \in C(y_{j-1})$ and $L_y(x_j) = L_y(y_j)$, a contradiction. So $X \in \mathcal{P}$, as required. Again by the argument of the previous paragraph, for any x_j and any $u, w \in C(x_j)$ we have $L_y(u) = L_y(w)$. This concludes part (ii a)

As $X \in \mathcal{P}$, (2.5) holds for X (our choice of $P \in \mathcal{P}$ was arbitrary). But then we have $\sum_{j=0}^{k-2} D(x_j) = n-2$ and so $X \setminus \{x_k\}$ sees every vertex of G as required for (ii b). As $X \in \mathcal{P}$, $t(x_k) = y$. This concludes the proof of (ii b).

We now complete the proof of Theorem 2.1.

Proof of Theorem 2.1. Let T := T(x, y) be the x-y-path tree of G rooted at v_0 . By (2.1), the number of induced paths between x and y is precisely the number of leaves $l \in T$ such that t(l) = y. As by Lemma 2.7(i) we know $L_y(v_0) \leq f_2(n)$ and moreover we know there exist graphs H such that $p_2(H) = f_2(n)$ (just pick $H \in \mathcal{F}_n$), then $L_y(v_0) = f_2(n)$. We will show that G is in \mathcal{F}_n . First we will show that G is a braid.

Let $P := v_0, \ldots, v_k$ be the shortest path in T such that $t(v_0) = x$ and $t(v_k) = y$. For $i \in \{0, \ldots, k-1\}$, define $C_{i+1} := C(v_i)$ to be the set of children of v_i in T (note that $v_{i+1} \in C_{i+1}$). Define $V_0 := \{x\}$ and define $V_i := t(C_i)$ for $1 \le i \le k$. Therefore, $V_k = \{y\}$. The sets V_i are disjoint by Proposition 2.5. We also have that $\bigcup_{i=0}^k V_i = V(G)$, as V(P) sees every vertex in G by Lemma 2.7(ii b). Theorem 2.1 will follow immediately from the next claim.

Claim 2.8. G is the braid $(\{x\}, V_1, \ldots, V_{k-1}, \{y\})$.

Proof. We prove by reverse induction on j that the graph induced by $\bigcup_{i=j}^{k} V_i$ is a braid (V_j, \ldots, V_k) in G. First, note that by Lemma 2.7(ii b), every leaf $l \in T$ satisfies t(l) = y. Thus no vertex in C_{k-1} can be a leaf of T (else we would have a shorter path to y in G) and every vertex in C_{k-1} has a child. Since, by Lemma 2.7(ii b), $V(P) \setminus \{v_k\}$ sees every vertex of G, all vertices except y have been seen by v_0, \ldots, v_{k-2} . Thus every child z of a vertex in C_{k-1} satisfies t(z) = y, otherwise it would contradict Proposition 2.5. Therefore every vertex in C_{k-1} has exactly one child z and t(z) = y. Therefore $(V_{k-1}, \{y\})$ is a braid, completing our base case.

We now show that $N_G(y) = V_{k-1}$. Suppose there exist some $0 \le j < k-1$ and some $u \in V_j$ such that $u \in N(y)$. Then there exists a vertex w in T such that w is a child of $v_{j-1}, t(w) = u$ and w has a child z with t(z) = y. The path $v_0, \ldots, v_{j-1}, w, z \subseteq T$ is shorter than P, a contradiction. So y is adjacent to no vertex in $\bigcup_{i=0}^{k-2} V_i$. As by Lemma 2.7(ii b) $V(P) \setminus \{v_k\}$ sees every vertex of $G \setminus \{y\}, V(G) \subseteq \bigcup_{i=0}^{k-1} V_i$ and $N_G(y) = V_{k-1}$.

Now suppose that for $1 \leq s+1 \leq k-1$, the inductive hypothesis holds for j = s+1. So (V_{s+1}, \ldots, V_k) is a braid in G. We will show that (V_s, \ldots, V_k) is a braid in G. In order to do

this, we need to show that for each $s + 1 \le r \le k - 1$, and every $u \in V_r$:

$$V_{r-1} \cup V_{r+1} \subseteq N(u) \subseteq V_{r-1} \cup V_r \cup V_{r+1}.$$
(2.6)

As by our inductive hypothesis (V_{s+1}, \ldots, V_k) is a braid in G, we know that (2.6) is satisfied for each $s + 2 \leq r \leq k - 1$ and every $u \in V_r$. We also know $V_{s+1} \subseteq N(u)$ for any $u \in V_{s+2}$. So it suffices to show for each $u \in V_{s+1}$,

$$V_s \subseteq N(u) \subseteq V_s \cup V_{s+1} \cup V_{s+2}$$

We first show there are no edges between $\bigcup_{i=0}^{s-1} V_i$ and V_{s+1} . Suppose, for some $i \leq s-1$, there exists a vertex $v \in C_i$ with a child z such that $t(z) \in V_{s+1}$. Then there exists a shorter path in T from v_0 to v_k : the path $v_0, \ldots, v_{i-1}, v, z, v_{s+2} \ldots, v_k$. This contradicts our choice of P as the shortest such path. Thus no such vertex v exists. So there are no edges between $\bigcup_{i=0}^{s-1} V_i$ and V_{s+1} . As (V_{s+1}, \ldots, V_k) is a braid in G, there are no edges between $\bigcup_{i=s+3}^k V_i$ and V_{s+1} . Thus $N(V_{s+1}) \subseteq V_s \cup V_{s+1} \cup V_{s+2}$.

It remains to show that $\{uw : u \in V_s, w \in V_{s+1}\} \subseteq E(G)$. Suppose there exists some $v \in C_s \setminus \{v_s\}$ and some $z \in C_{s+1}$ such that t(v) is not adjacent to $t(z) \in V_{s+1}$.

We know $t(v) \neq y$, it would contradict the choice of P otherwise as s < k - 1. As by Lemma 2.7 every leaf l satisfies t(l) = y, v is not a leaf and has a child u. We know that t(u) is a neighbour of t(v) in G and that:

- $t(u) \notin \bigcup_{i=0}^{s} V_i$, as t(u) is unseen by $\{v_0, \ldots, v_{s-1}\}$ by construction of T;
- $t(u) \notin \bigcup_{i=s+2}^{k} V_i$, as V_{s+1}, \ldots, V_k forms a braid in G.

Thus $t(u) \in V_{s+1}$.

If s + 2 = k (and so $V_{s+2} = V_k = \{y\}$) then consider the path $v_0, v_1, \ldots, v_{s-1}, v \in T$. We have $L_y(v) = D(v) < D(v_s) = L_y(v_s)$, contradicting Lemma 2.7(ii a).

Therefore s + 2 < k. Since every leaf $l \in T$ satisfies t(l) = y, by construction of T any induced path x, x_1, \ldots, x_j in G such that $y \notin N(x_j)$ can be extended to an induced path terminating at y.

We consider two cases (see Figure 3 for an illustration).

First suppose that t(u) is adjacent to t(z). Consider $P := t(v_0), \ldots, t(v_{s-1}), t(v), t(u), t(z)$, an induced path in G. As $t(z) \in V_{s+1}$ and s + 1 < k - 1, y is not adjacent to t(z) and so it is possible to extend P to an induced path terminating at y. As (V_{s+1}, \ldots, V_k) is a braid in G, any extension of P to an induced path that terminates at y contains a vertex from V_{s+2} . However, $V_{s+2} \subseteq N_G(t(u))$ (and so V_{s+2} has been seen by $V(P_u)$). It is therefore impossible to extend P to an induced path terminating at y, a contradiction.

Now suppose that t(u) is not adjacent to t(z). Let w be a neighbour of t(u) in V_{s+2} . Consider the induced path $P := t(v_0), \ldots, t(v_{s-1}), t(v), t(u), w, t(z)$. Observe that $y \notin N(t(z))$. As $t(z) \in V_{s+1}$ and (V_{s+1}, \ldots, V_k) is a braid in G, any extension of P from t(z) to an induced path that terminates at y passes through V_{s+2} . However, $V_{s+2} \subseteq N_G(t(u))$ and so has been seen by $V(P_{t(u)})$. It is therefore impossible to extend this P from t(z) to an induced path terminating at y, a contradiction.




Figure 3: Examples of the cases we get if t(v) is not adjacent to every vertex in V_{s+1} . The upper picture is the case t(u) is adjacent to t(z). The lower picture is the case t(u) is not adjacent to t(z). The dashed lines represent non-edges. The blue lines are the induced path we take from t(v) in each case. The red edges depict which vertices have been seen by $P \setminus \{t(z)\}$.

Thus $\{uw : u \in V_s, w \in V_{s+1}\} \subseteq E(G)$. We conclude that the graph induced by $\bigcup_{i=s}^k V_i$ is indeed a braid (V_s, \ldots, V_k) in G. Claim 2.8 now follows by induction.

We have $|V_i| = D(v_{i-1})$. As $\prod_{i=0}^{k-2} D(v_i) = p_2(F(n))$, a straightforward check shows that the braid is in \mathcal{F}_n . Hence Theorem 2.1 follows.

2.1 Odd and even induced paths between a pair of vertices

Let G be a graph and let x and y be distinct vertices in V(G). We will define similar notions for odd and even paths as we did for paths in general at the start of Section 2. Define $p_2^o(G; x, y)$ to be the number of induced odd paths in G beginning at x and ending at y. Also define:

$$p_2^o(G) := \max\{p_2^o(G; x, y) : x, y \in V(G)\},\$$

and

$$p_2^o(n) := \max\{p_2^o(G) : |V(G)| = n\}.$$

In addition, define $p_2^e(G; x, y)$ to be the number of induced even paths in G beginning at x and ending at y. In the even case, define $p_2^e(G)$ and $p_2^e(n)$ analogously to $p_2^o(G)$ and $p_2^o(n)$.

In this subsection we will determine the structure of the *n*-vertex graphs that contain $p_2^o(n)$ induced odd paths (or $p_2^e(n)$ induced even paths) between some pair of vertices (proving Theorems 2.9 and 2.10). Theorem 2.9 will be used to prove Theorem 1.6. The extremal graphs for this path problem will have a certain structure that depends on the value of n modulo 6. For $n \ge 10$, define \mathcal{F}_n^o to be the set of all braids B with the following properties.

- |V(B)| = n.
- *B* has end clusters of size 1.
- All central clusters of *B* have size three except:
 - a single cluster of size 4, when $n \equiv 0$ modulo 6;
 - either two clusters of size 4 or four clusters of size 2, when $n \equiv 1 \mod 6$;
 - three clusters of size 2, when $n \equiv 2 \mod 6$;
 - two clusters of size 2, when $n \equiv 3 \mod 6$; and
 - one cluster of size 2, when $n \equiv 4 \mod 6$.

Let $F \in \mathcal{F}_n^o$ and suppose that the end clusters are $\{x\}$ and $\{y\}$. Observe that every induced path between x and y is odd. It is not difficult to check that for all $F \in \mathcal{F}_n^o$ we have $p_2^o(F) = p_2^o(F; x, y)$. Every graph in \mathcal{F}_n^o contains the same number of induced paths between x and y and so we define:

$$f_2^o(n) = \begin{cases} 4 \cdot 3^{(n-6)/3} & \text{for } n \equiv 0 \mod 0 \ 6\\ 2^4 \cdot 3^{(n-10)/3} & \text{for } n \equiv 1 \mod 0 \ 6\\ 2^3 \cdot 3^{(n-8)/3} & \text{for } n \equiv 2 \mod 0 \ 6\\ 2^2 \cdot 3^{(n-6)/3} & \text{for } n \equiv 3 \mod 0 \ 6\\ 2 \cdot 3^{(n-4)/3} & \text{for } n \equiv 4 \mod 0 \ 6\\ 3^{(n-2)/3} & \text{for } n \equiv 5 \mod 0 \ 6. \end{cases}$$

The following is a theorem for odd paths analogous to Theorem 2.1.

Theorem 2.9. Let G be a finite graph on $n \ge 10$ vertices and let x and y be distinct vertices of G. Suppose that $p_2^o(G; x, y) = p_2^o(n)$. Then G is isomorphic to a graph in \mathcal{F}_n^o with end clusters $\{x\}$ and $\{y\}$.

The proof of this theorem is very similar to the proof of Theorem 2.1 and a sketch will be given later in this subsection.

We will also state a version of Theorem 2.9 for even length paths (Theorem 2.10). As one would expect, the extremal graphs differ from those in the odd case. Thus for $n \ge 10$, define \mathcal{F}_n^e to be the set of all braids B with the following properties.




Figure 4: An example of a braid in \mathcal{F}_{13}^e and a braid in \mathcal{F}_{10}^o . There may or may not be edges within the clusters.

- |V(B)| = n.
- *B* has end clusters of size 1.
- All central clusters of *B* have size three except:
 - two clusters of size 2, when $n \equiv 0$ modulo 6;
 - one cluster of size 2, when $n \equiv 1 \mod 6$;
 - a single cluster of size 4, when $n \equiv 3 \mod 6$;
 - either two clusters of size 4 or four clusters of size 2, when $n \equiv 4 \mod 6$; and
 - three clusters of size 2, when $n \equiv 5 \mod 6$.

Observe that the extremal graphs in the odd and even cases are essentially the same (shifting by 3 modulo 6), as when $n \ge 13$ we can delete a cluster of size 3 to get from an extremal graph for the odd case to an extremal graph for the even case (or vice versa). See Figure 4 for an example.

Theorem 2.10. Let G be a finite graph on $n \ge 10$ vertices and let x and y be distinct vertices of G. Suppose that $p_2^e(G; x, y) = p_2^e(n)$. Then G is isomorphic to a graph in \mathcal{F}_n^e with end clusters $\{x\}$ and $\{y\}$.

To obtain the proof of Theorem 1.6, we only need Theorem 2.9. We remark that the proof of Theorem 2.9 can easily be adapted to prove Theorem 2.10, so we omit the proof of Theorem 2.10.

Sketch proof of Theorem 2.9. Fix x and y to be distinct vertices of G with $y \notin N[x]$. Let T be the x-y-path tree rooted at v_0 . For $z \in V(T)$ define $L_o(z)$ to be the number of leaves l contained in B(z) such that t(l) = y and the path from v_0 to l is odd.

We first prove an *odd-path version* of Lemma 2.7.

Claim 2.11 (Odd-path version of Lemma 2.7). Let G be a graph on $n \ge 10$ vertices. Let x and y be distinct vertices in V(G) with $y \notin N[x]$ and $p_2^o(G; x, y) > 0$. Let T be the x-y-path tree rooted at v_0 and $P := x_1, \ldots, x_k$ be any path in T where $x_0 = v_0$ and x_k is a leaf. Then:

- (i) $L_o(v_0) \le f_2^o(n)$.
- (ii) If $L_o(v_0) = f_2^o(n)$, then:
 - (a) for any x_i and for all $u, w \in C(x_i)$, we have $L_o(u) = L_o(w)$;
 - (b) k is even;
 - (c) $V(P) \setminus \{x_k\}$ sees every vertex of G and $t(x_k) = y$.

Proof of Claim. We essentially mimic the proof of Lemma 2.7, replacing \mathcal{F}_n with \mathcal{F}_n^o and $L_y(z)$ with $L_o(z)$ for any $z \in T$.

Sequentially choose a path $v_0, \ldots, v_k \subseteq V(T)$, where v_k is a leaf. At vertex v_j we choose v_{j+1} to be some $z \in C(v_j)$ such that $L_o(z) = \max\{L_o(x) : x \in C(v_j)\}$. Let \mathcal{P} be the set of paths that can be obtained in this manner and fix $P := v_0, \ldots, v_k \in \mathcal{P}$.

We now show that k is even and $t(v_k) = y$. Suppose first that k is odd. Thus any path from v_0 to some leaf neighbour of v_{k-1} is even. As $p_2^o(G; x, y) > 0$, by construction of P we have $L_o(v_{k-1}) > 0$. So v_{k-1} has some non-leaf child z with $L_o(z) > 0$. But then $L_o(z) > L_o(v_k) = 0$, a contradiction. So k is even. The fact that $t(v_k) = y$ follows from exactly the same argument as in Lemma 2.7.

Arguing as in (2.3), we see that

$$L_o(v_0) \le \prod_{i=0}^{k-2} D(v_i),$$
 (2.7)

where k is even and $\sum_{i=0}^{k-2} D(v_i) \leq n-2$, as v_0, \ldots, v_{k-2} have disjoint sets of children in $G \setminus \{x, y\}$ by proposition 2.5.

It is not difficult to check that the maximal value of $\prod_{i=0}^{k-2} D(v_i)$ subject to $\sum_{i=0}^{k-2} D(v_i) \le n-2$, where k is even is $f_2^o(n)$. This concludes the proof of (i).

When $L_o(v_0) = f_2^o(n)$ we have for even k:

$$\prod_{i=0}^{k-2} D(v_i) = f_2^o(n)$$

Statement (ii a) follows from an analogous argument to the proof of (ii a) in Lemma 2.7. Also (using an identical argument to the one used in Lemma 2.7) we have that any path $X := x_0, \ldots, x_j$, where $x_0 = v_0$ and x_k is a leaf, is in \mathcal{P} . Thus j is even, as required for (ii b), and $t(v_k) = y$. The other claim in statement (ii c) follows an analogous argument to the proof of (ii b) in Lemma 2.7. This completes the proof of the claim.

In particular, we know that any path $v_0 \ldots v_k$, where v_k is a leaf, is odd and satisfies $t(v_k) = y$. We now use analogous arguments to those used in the proof of Theorem 2.1 replacing \mathcal{F}_n with \mathcal{F}_n^o , replacing $L_y(z)$ with $L_o(z)$ for any $z \in T$ and applying Claim 2.11 in the place of Lemma 2.7, to show that G is a braid in \mathcal{F}_n^o .

3 Proof of Theorem 1.2

We fix a large constant n_0 and let G_{max} be a graph on $n \ge n_0$ vertices, that contains m(n) induced cycles. In what follows we will take n_0 (and thus n) to be sufficiently large when required and we will make no attempts to optimise the constants in our arguments. We will show that the graph G_{max} is isomorphic to H_n . As it turns out, Theorem 1.5 (the stability result) will follow almost immediately from the arguments required for the proof of Theorem 1.2. Therefore, in this section several lemmas are proved in more generality than is needed for the proof of Theorem 1.2: they will be used in their more general form in the next section.

Given a graph H, let f(H) denote the number of induced cycles in H and for a vertex $v \in H$, let $f_v(H)$ denote the number of induced cycles in H containing v. Observe that we have:

$$f(G_{max}) = m(n) \ge f(H_n) \ge \begin{cases} 3^{n/3} & \text{if } n \equiv 0 \mod 0 \ 3 \\ 4 \cdot 3^{(n-4)/3} & \text{if } n \equiv 1 \mod 0 \ 3 \\ 2 \cdot 3^{(n-2)/3} & \text{if } n \equiv 2 \mod 0 \ 3. \end{cases}$$
(3.1)

Any G_{max} is connected (if it were disconnected we could add edges between two components to increase the number of induced cycles). We begin by proving several lemmas which determine information about the structure of G_{max} .

Lemma 3.1. Let F be an n-vertex graph. For $v \in V(F)$, we have $f_v(F) \leq {\binom{d(v)}{2}} 3^{(n-d(v)-1)/3}$.

Proof. Each induced cycle containing v contains exactly two neighbours of v. Fix a pair of vertices $u, w \in N(v)$. By Corollary 2.2 there are at most $3^{(n-d(v)-1)/3}$ induced paths between u and w in $(F \setminus N[v]) \cup \{u, w\}$. Thus there can be at most $3^{(n-d(v)-1)/3}$ induced cycles in F containing $\{v, u, w\}$. As there are $\binom{d(v)}{2}$ distinct pairs of neighbours of v, we have,

$$f_v(F) \le {d(v) \choose 2} 3^{(n-d(v)-1)/3}$$

as required.

The next lemma tells us that any vertex in G_{max} is contained in a constant proportion of $f(G_{max})$ induced cycles.

Lemma 3.2. Let $0 < c \leq 1$ and let $\alpha = 0.11$. Let F be an n-vertex graph with $f(F) \geq c \cdot m(n)$. Then:

(i) (1 - o(1))f(F) induced cycles in F have length at least αn .

- (ii) F contains a vertex v such that $f_v(F) \geq \frac{c}{10}m(n)$.
- (iii) Every vertex $w \in V(G_{max})$ satisfies $f_w(G_{max}) \geq \frac{c}{20}m(n)$.

Proof. F contains at most $\sum_{i=1}^{\lfloor \alpha n \rfloor} {n \choose i}$ induced cycles of length at most αn (for any $W \subseteq V(F)$, there exists at most one induced cycle C such that V(C) = W). Using Stirling's approximation, we get

$$\sum_{i=1}^{\lfloor \alpha n \rfloor} \binom{n}{i} \le \alpha n \cdot \binom{n}{\alpha n} \le (1+o(1)) \frac{\sqrt{\alpha n}}{\sqrt{2\pi(1-\alpha)}} \left[\frac{1}{\alpha^{\alpha}(1-\alpha)^{1-\alpha}}\right]^n.$$
$$\frac{1}{\alpha^{\alpha}(1-\alpha)^{1-\alpha}} < 3^{1/3},$$

we get

As

$$\sum_{i=1}^{\lfloor \alpha n \rfloor} \binom{n}{i} = o\left(3^{n/3}\right).$$

By (3.1), $f(F) = \Omega(3^{n/3})$, so (1 - o(1))f(F) induced cycles in F have length at least αn , as required for (i).

Provided n_0 is sufficiently large, we have for all $n > n_0$,

$$\sum_{i=1}^{\lfloor \alpha n \rfloor} \binom{n}{i} < \frac{c}{1000} \cdot 3^{n/3} < \frac{c}{100} m(n),$$

where the second inequality follows from (3.1). Thus there exists a vertex v such that

$$f_v(F) \ge \frac{99\alpha c}{100} m(n) \ge \frac{c}{10} m(n),$$

proving (ii).

Now suppose that there exists some vertex $w \in V(G_{max})$ with $f_w(G_{max}) < \frac{c}{20}m(n)$. Consider the graph G' obtained from G_{max} by duplicating the vertex v and removing the vertex w. We have that

$$f(G') \ge f(G_{max}) + f_v(G_{max}) - 2f_w(G_{max}) > f(G_{max}),$$

a contradiction. This proves (iii).

Now consider the graph obtained from G_{max} by duplicating a vertex. By applying Lemma 3.2, we have

$$m(n+1) \ge \left(1 + \frac{1}{20}\right) m(n).$$
 (3.2)

We now use this to show that the graph G_{max} has maximum degree bounded by a constant.

Lemma 3.3. $\Delta(G_{max}) < 30.$

Proof. Let v be a vertex of maximal degree. Given v, split the induced cycles in G_{max} into those contained in $G_{max} \setminus \{v\}$, and those containing v. Using Lemma 3.1 we have

$$m(n) \le m(n-1) + {d(v) \choose 2} 3^{(n-d(v)-1)/3}$$

Using (3.2) to bound m(n-1) gives,

$$m(n) \le m(n) \left(1 + \frac{1}{20}\right)^{-1} + {d(v) \choose 2} 3^{(n-d(v)-1)/3}.$$

This expression rearranges to give

$$m(n) \le 21 \binom{d(v)}{2} 3^{(n-d(v)-1)/3}$$

For $d(v) \ge 30$, this implies $m(n) < 3^{(n-6)/3}$, a contradiction (as $m(n) \ge f(H_n) > 3^{(n-6)/3}$).

Combining Lemma 3.1 with Lemma 3.3 shows, for any $v \in V(G_{max})$, we have $f_v(G_{max}) \leq {\binom{30}{2}}3^{(n-3)/3}$. By Lemma 3.2 (i), we know (1-o(1))m(n) induced cycles in G_{max} have length at least 0.11*n*. Thus

$$(1 - o(1))m(n) \le \frac{1}{0.11n} \sum_{v \in V(G)} f_v(G_{max}) = O(3^{n/3}).$$

This along with (3.1) implies

$$m(n) = \Theta(3^{n/3}). \tag{3.3}$$

The next stage of our proof involves showing that all but a constant number of vertices v in our graph have the property that their closed second neighbourhood $N^2[v]$ has the same local structure as the closed second neighbourhood of a vertex in H_n . We introduce some preliminary definitions.

Given a graph F and a set $S \subseteq V(F)$, we say that a vertex $v \in V(F)$ is seen by S if $v \in N[S]$ and v is unseen by S otherwise. Given a subgraph $H \subseteq F$, we say v is seen by H if $v \in N[H]$. When it is clear which set/subgraph we are referring to, we will just say v is (un)seen. For a vertex $v \in V(G)$, let $N^i(v)$ be the set of points at distance exactly i from v. Define $N^k[v]$ to be the set of points within distance k of v (for example $N^3[v] = \{v\} \cup \bigcup_{i=1}^3 N^i(v)$ and $N^1[v] := N[v]$).

In order to determine what sort of local structure a 'typical' vertex in G_{max} should have, we define a game on F.

Definition 3.4. Let F be a finite graph, let $v \in V(F)$ and let $w \in V(F) \setminus N^4[v]$. We define the *w*-typical-game on (F, v) as follows. There are two players, Adversary and Builder. The game starts at vertex $u_1 := v$ and the players choose a sequence of vertices $\{u_2, \ldots, u_k\}$ under the following set of rules. At vertex u_i :

- If $u_i \in N^4[w]$, then Adversary is the *active player*, otherwise Builder is.
- The active player chooses a neighbour u_{i+1} of u_i that is unseen by $\{u_1, \ldots, u_{i-1}\}$.
- The game terminates when a vertex u_j is chosen such that u_j has no neighbours unseen by $\{u_1, \ldots, u_{j-1}\}$.
- If, for some j, the vertex u_j does not have exactly 3 neighbours unseen by $\{u_1, \ldots, u_{j-1}\}$, we call u_j bad; we call u_j good otherwise.
- Adversary wins if either:
 - for some j, the vertex u_j is in $N^4[w]$ and is bad; or
 - upon termination of the game at vertex u_k , there exists a vertex in $N^4[w]$ that is unseen by $\{u_1, \ldots, u_k\}$.

Builder wins otherwise.

A vertex $w \in V(F) \setminus N^4[v]$ is *v*-typical in F if there exists a winning strategy for Builder in the *w*-typical-game on (F, v). A vertex is *v*-atypical otherwise. When it is clear which vertex has been chosen to play the role of v, we simply say that w is (a)typical. Note that the set of vertices $\{u_1, \ldots, u_k\}$ chosen during the game induces a path in F. Also, if we play this game on H_n starting at any vertex, most of the chosen vertices are good.

The next lemma shows a v-typical vertex has the required local structure (see Figure 5).

Lemma 3.5. Let F be a graph and let v be a vertex in F. Suppose that $z = z_1 \in V(F) \setminus N^4[v]$ is a v-typical vertex. Then there exist disjoint sets of vertices $Z := \{z_1, z_2, z_3\}, V := \{v_1, v_2, v_3\}$, and $W := \{w_1, w_2, w_3\}$ such that:

- (i) for $i, j \in \{1, 2, 3\}$, we have $N(v_i) \cap N(w_j) = Z$;
- (ii) for all i, we have $V \cup W \subseteq N(z_i) \subseteq V \cup W \cup Z$;
- (iii) there are no edges between V and W.

Proof. We play as Adversary in the z-typical-game on (F, v). As z is v-typical, Builder has a winning strategy σ . We assume that Builder uses strategy σ , and deduce information about the structure of F from the results of our choices of vertices as Adversary (we know we cannot win so whatever choices we make have certain consequences). For each vertex u_i that is chosen, let P_{u_i} denote the subgraph induced by $\{u_1, \ldots, u_i\}$, where $u_1 = v$. So P_{u_i} is an induced path between v and u_i .

Suppose that u_k is the first vertex chosen such that $u_k \in N^4(z)$ (as z is typical, at some point such a vertex will be chosen). We arbitrarily choose the next two vertices $u_{k+1} \in N^3(z) \cap N(u_k)$ and $u_{k+2} \in N^2(z) \cap N(u_{k+1})$. Let $x := u_{k+2}$. This vertex is unseen by P_{u_k} as u_{k+1} was the first vertex we chose in $N^3(z)$. We also have, by choice of u_k , that $x \notin N(z)$. As z is typical, x has 3 neighbours $V := \{v_1, v_2, v_3\}$ unseen by $P_{u_{k+1}}$.




Figure 5: The local structure around a v-typical vertex z_1 . Note that we have not yet determined the edges within the sets V, Z and W.

As $x \in N^2(z)$, for some *i* we have $v_i \in N(x) \cap N(z)$. Without loss of generality suppose $v_1 \in N(x) \cap N(z)$. Since we could choose u_{k+3} to be v_1 , the vertex v_1 has 3 neighbours unseen by P_x , one of which is *z*, so let the set of neighbours of v_1 unseen by P_x be $Z := \{z_1, z_2, z_3\}$.

If we choose $u_{k+3} := v_1$ and then $u_{k+4} := z$, we have that z has 3 neighbours unseen by P_{v_1} . Thus z has at most 5 neighbours unseen by P_x (as z could be adjacent to z_2 or z_3).

We now prove that $N(x) \cap N(z) = V$. Suppose otherwise, so without loss of generality we have $v_3 \notin N(z)$. Now we describe the set of choices we make for the remainder of the game (recall that Builder always plays by strategy σ). Choose $u_{k+3} := v_3$. Now consider a later step in the game, but before z has been chosen, and suppose the most recently chosen vertex is $u_i \in N^4[z]$, where $i \geq k+3$. Then:

- (1) If there is no vertex in $N(u_i) \cap N(z)$ that is unseen by $P_{u_{i-1}}$, choose u_{i+1} arbitrarily.
- (2) If, for some r, the vertex z has r neighbours unseen by $P_{u_{i-1}}$ (by the argument above, $r \leq 5$) and u_i is adjacent to $j \geq 1$ of these neighbours:
 - (i) If r j < 3, choose $u_{i+1} \in N(u_i) \cap N(z)$ and $u_{i+2} := z$. As z now has at most 2 neighbours unseen by $P_{u_{i+1}}$ we reach a contradiction.
 - (ii) If $r j \ge 3$, then $j \le 2$ and u_i has an unseen neighbour s that is not adjacent to z (otherwise u_i is bad, which contradicts the fact that Builder has a winning strategy). Choose $u_{i+1} := s$. Observe that the vertex z is not seen by P_s .

Once we have chosen z, we play arbitrarily.

We now analyse the results of making these choices. As z is typical, we will at some point enter case (2). If we are in case (2ii), we pick u_{i+1} and the number of neighbours unseen by z decreases, so eventually we enter case (2i) where we reach a contradiction. Thus $N(x) \cap N(z) = V$, as required. We now know that $N(x) \cap N(z) = V$. As in the first part, if we choose $u_{k+3} := v_1$ and $u_{k+4} := z$, the vertex z has 3 neighbours unseen by P_{v_1} . Call these neighbours W := $\{w_1, w_2, w_3\}$. Thus $N(z) \supseteq V \cup W$ as required. Observe that by choice of W, there are no edges between V and W, thus proving the third statement of the lemma. From the argument above we have $N(z) \subseteq V \cup W \cup Z$.

Let $w_j \in W$, for $j \in \{1, 2, 3\}$. We will first show that $N(v_1) \cap N(w_j) = Z$. Again we play as Adversary in a new z-typical-game on (F, v) and assume that Builder uses the same winning strategy σ as above.

By playing the same strategy as in the game above, u_{k+1} is the first vertex chosen in $N^3(z)$ and that the first vertex chosen in $N^2(z)$ is x. We know that $N(x) \cap N(z) = V$. Now we choose $u_{k+3} := v_1$. Let y be the first vertex chosen in $N^2(w_j)$. Observe that as $w_j \in N(z)$, we have $N^2(w_j) \subseteq N^3[z]$. So $y \in \{u_{k+1}, x, v_1\}$. We will show that $y = v_1$.

By repeating the same argument as above (where we showed $N(x) \cap N(z) = V$) with yin place of x and w_j in place of z, we see that y and w_j have exactly 3 common neighbours. The assumptions we needed to apply the argument above will hold here: we require that y is the first vertex chosen in $N^2(w_j)$ and that we are able to make moves at vertices in $N^2[w_j]$. The latter holds as $N^2[w_j] \subseteq N^4[z]$ and, in the z-typical game on (F, v), Adversary makes moves at vertices in $N^4[z]$.

Suppose that $y = u_{k+1}$. Then $|N(u_{k+1}) \cap N(w_j)| = 3$ and in particular w_j is adjacent to x. This is a contradiction as $w_j \notin V$. Now suppose that y = x. This implies that w_j is adjacent to v_1 . As $w_j \notin Z$, this is a contradiction. Therefore $y = v_1$ and $N(v_1) \cap N(w_j) = Z$ for each $j \in \{1, 2, 3\}$ (by the analogous argument to where we show $N(x) \cap N(z) = V$, with v_1 in place of x, w_j in place of z and Z in place of V).

We now show that, for $i, j \in \{2, 3\}$, each v_i is adjacent to each z_j . Suppose that v_i is not adjacent to z_j for some $i, j \in \{2, 3\}$. Again we play as Adversary in a new z-typical-game on (F, v). We assume that Builder uses the same winning strategy σ as above. By playing the same strategy as in both games above, the first vertex chosen in $N^2[z]$ is x and $x = u_{k+2}$ in the sequence of vertices chosen. It is deduced from analogous arguments to those above that the only neighbours of z_j that are unseen by P_x are contained in $V \cup Z \cup W$. We know that $N(x) \cap N(z) = V$. Now choose $u_{k+3} := v_i$ and $u_{k+4} := z$. If z is adjacent to z_j , pick $u_{k+5} := z_j$. Otherwise, pick $u_{k+5} := w_1$ and $u_{k+6} := z_j$. In both cases, all vertices of $W \cup V$ have been seen when z_j is chosen. Therefore z_j has at most one neighbour (the vertex in $Z \setminus \{z, z_j\}$) unseen by $P_{z_j} \setminus \{z_j\}$. This contradicts z being typical. Thus we have $N(v_i) \cap N(w_i) = Z$, completing the proof of (i).

It remains to show that $N(z_i) \subseteq V \cup W \cup Z$ for $i \in \{2, 3\}$. Suppose otherwise, that z_i has a neighbour $u \notin V \cup Z \cup W$. By the above arguments, u is seen by P_x . u is not a neighbour of x, and as $u \in N^3[z]$, u is either a neighbour of u_k or u_{k+1} . Now consider a new z-typical game on (F, v). Builder still plays by the winning strategy σ . This time, we play as before until we are the first vertex u' which is a neighbour of u (so u' is either u_k or u_{k+1}). At u', we choose u and then z_i . Observe that $N(z_i) \supseteq W \cup V$. By the arguments above, no vertices in $W \cup V$ are seen by $P_{u'}$. As z_i is typical it has exactly 3 neighbours unseen by P_u , this implies that at least 3 vertices of $W \cup V$ are neighbours of u. However, this means that u has 4 neighbours unseen by $P_{u'}$ (these 3 and z_i), contradicting z being typical. Thus for $i \in \{2,3\}, N(z_i) \subseteq V \cup W \cup Z$. This completes the proof of the lemma.

We will now show that, for any $v \in V(G_{max})$, all but a bounded number of vertices in G_{max} are v-typical. We do this in the following manner. For each vertex v in G_{max} we define a tree T(v) that will 'explore' the graph G_{max} outwards from v. As we will see, leaves on this tree will correspond to induced paths or cycles in G_{max} containing v. Every vertex on T represents some vertex in G_{max} (and many vertices in T may represent the same vertex of G_{max}). Our proof proceeds by showing that T has a particular structure, which in turn implies conditions on the structure of G_{max} .

The next definition contains similar concepts to those introduced in the definition of an x-y-path-tree in Section 2. The main difference is that previously the vertex y played a special role in the creation of leaves of our tree. Now there is no such significant vertex.

Definition 3.6. For F a finite graph and $v \in V(F)$, the *exploration tree from* v is a tree T = T(v) together with a function $t : V(T) \to V(F)$ defined as follows.

- T is a tree with vertex set V(T) disjoint from V(F).
- The vertices in T correspond to sets $S := \{v, v_1, \ldots, v_j\} \subseteq V(F)$ such that F[S] is an induced path v, v_1, \ldots, v_j or induced cycle v, v_1, \ldots, v_j, v . For each S where F[S] is an induced path, define a vertex $w_S \in T$ and set $t(w_S) = v_j$. For each S where F[S] is an induced cycle (where an edge is not considered to be a cycle), define two vertices $w_S^1, w_S^2 \in T$ and set $t(w_S^1) = v_1$ and $t(w_S^2) = v_j$. We call S the F-set of w (or F-set of w_S^1 and w_S^2 , if F[S] is a cycle). These vertices are the only vertices in T. Define the root of the tree to be $v_0 := w_x$.
- Given a vertex $w \in V(T)$ with F-set $\{v, v_1, \ldots, v_j\}$ we define C(w), the children of w, to be the set of vertices in T whose F-set is $\{v, v_1, \ldots, v_j, z\}$ for some $z \in V(F)$. Define $N_T(v_0) := C(v_0)$. For $w \in V(T) \setminus \{v_0\}$ define $N_T(w) := C(w) \cup \{u\}$, where u is the unique vertex in T with F-set $\{v, v_1, \ldots, v_{j-1}\}$. (So two vertices are adjacent in T precisely when one of their F-sets extends the other by one vertex. A vertex whose F-set induces a cycle in F will be a leaf of T.)

We write $t(S) := \{t(x) : x \in S\}$ for any subset $S \subseteq V(T)$ and $t(H) := \{t(x) : x \in V(H)\}$ for any subgraph $H \subseteq T$. Given a set $P \subseteq V(T)$ we say that it sees a vertex $w \in V(F)$ if $w \in N_F[t(P)]$. If $w \notin N_F[t(P)]$ we say w is unseen by P. Note that if some set P sees wthen there exists $u \in N_T[P]$ such that t(u) = w.

As in Definition 2.6, define a *branch* and L(u) for $u \in V(T)$ with respect to this tree. We now describe a correspondence between certain leaves on T and induced cycles in F. For $z \in T$, let L(z) be the number of leaves of T contained in B(z).

See Figure 6 for an example of an exploration tree.




Figure 6: A graph and its exploration tree (from v_1). Each vertex w in the tree is labelled by t(w).

Lemma 3.7. For F a finite graph and $v \in V(F)$, let T be the exploration tree from v rooted at v_0 . We have $f_v(F) \leq \frac{1}{2}L(v_0)$.

Proof. In the construction of T, for every induced cycle containing v in F we define two vertices of T. Again by construction, these two vertices are leaves of T. The result follows.

Call a vertex $v \in T$ is good if it has exactly three children: call it bad otherwise. We now define a game on T, as we did previously in this section for a graph. We use the game to define vertices that are '(a)typical' for T. The following definition is the analogue in T of Definition 3.4 for a graph.

Definition 3.8. Let F be a finite graph and let T be the exploration tree from v in F. Let w be a vertex in $V(F) \setminus N^4[v]$. We define the *w*-typical-game on T as follows. There are two players, Adversary and Builder. The game starts at vertex $u_0 := v_0$ ($v_0 = t(v)$) and the players choose a sequence of vertices $\{u_1, u_2, \ldots, u_k\} \subseteq V(T)$ under the following set of rules. At vertex u_i :

- If $t(u_i) \in N^4[w]$, then Adversary is the *active player*, otherwise Builder is.
- The active player chooses a child u_{i+1} of u_i .

- The game terminates when a vertex u_j is chosen such that u_j is a leaf.
- Adversary wins if either for some j, we have that $t(u_j) \in N^4[w]$ and u_j is bad, or if upon termination of the game at vertex u_k there exists a vertex in $N^4[w]$ that is unseen by $\{u_0, \ldots, u_k\}$. Builder wins otherwise.

A vertex $w \in V(F) \setminus N^4[v]$ is typical for T if there exists a winning strategy for Builder in the w-typical-game on T. A vertex is atypical for T otherwise. Observe that a vertex in $V(F) \setminus N^4[v]$ is atypical for T(v) if and only if it is v-atypical in F. Also, note that a vertex wbeing atypical for T means that Adversary has a strategy to ensure that, whatever strategy Builder chooses, either a bad vertex in $N^4[w]$ is chosen, or that there exists some vertex in $N^4[w]$ that remains unseen by $\{u_1, \ldots, u_k\}$ upon termination at vertex u_k .

Now let c > 0 and F be any *n*-vertex graph with $f(F) \ge c \cdot 3^{n/3}$ and $\Delta(F) \le \Delta$, for some constant Δ (G_{max} satisfies these conditions as $\Delta(G_{max})$ is bounded by Lemma 3.3). Our next aim is to prove that, for any vertex $v \in V(F)$, only a bounded number of vertices are atypical for T(v). Using this fact with Lemma 3.5 implies that the majority of the structure of G_{max} is close to the structure of H_n . The remainder of the proof consists of 'cleaning' G_{max} to show that it is in fact isomorphic to H_n .

We first outline how the proof will proceed before giving the details. We assume (in order to get a contradiction) that there is a large set $A \subseteq V(F)$ of vertices atypical for T(v), such that for each $a, a' \in A$ we have $N^4[a] \cap N^4[a'] = \emptyset$ (for any set of atypical vertices there exists a subset of constant proportion with this property as $\Delta(F)$ is bounded). We will sequentially choose a path in T of vertices u_0, \ldots, u_k where $u_0 := v_0$ and u_k is a leaf.

For each $a \in A$, there exists a winning strategy τ_a for Adversary in the *a*-typical game on T(v). This means that whatever vertices u_i with $t(u_i) \notin \bigcup_{a \in A} N^4[a]$, are chosen in the path, for every $a \in A$ we are able to ensure that either:

- (i) we choose a bad vertex u_i with $t(u_i) \in N^4[a]$, or
- (ii) there is some vertex in $N^4[a]$ that remains unseen by $\{u_0, \ldots, u_k\}$.

We assume at the start that $L(v_0)$ is bounded below by $c \cdot 3^{n/3}$, for some constant c. As we move down the tree we keep track of the number of leaves that the branch we are in contains. If we are at a vertex u_i , such that $t(u_i)$ is not in $N^4[a]$ for any $a \in A$, we choose the branch that has the most leaves. When $t(u_i)$ is in $N^4[a]$ for some $a \in A$, we play the winning strategy τ_a to move towards the outcomes (i) or (ii), unless there is a sub-branch that contains a large proportion of the leaves in our current branch. These outcomes mean that the tree is 'unbalanced' in some way, and the strategy that achieves these outcomes picks branches that contain more leaves than average. As it turns out, when we reach a leaf and the process ends, if |A| was too large we find that the branch we are in ought to contain more than one leaf, a contradiction.

Lemma 3.9. Fix c > 0. Let F be an n-vertex graph with $\Delta := \Delta(F) > 1$ and let $\epsilon := 2^{-\Delta^{100}}$. Let $v \in V(F)$ and let T = T(v) be the exploration tree from v in F with root v_0 . Let $A \subseteq V(F) \setminus N^4[v]$ be a set of atypical vertices for T such that for all $a, a' \in A$, we have $N^4[a] \cap N^4[a'] = \emptyset$. If $L(v_0) \ge c \cdot 3^{n/3}$, then |A| < M, where M is the smallest integer such that $c \cdot 3^{1/3}(1+\epsilon)^M > \Delta$.

Proof. Suppose, in order to obtain a contradiction, that $|A| \ge M$. For each $a \in A$, Adversary has a winning strategy τ_a played on vertices of $N^4[a]$ in the *a*-typical game on *T*. As for all $a, a' \in A$, we have $N^4[a] \cap N^4[a'] = \emptyset$, these strategies are played on disjoint sets of vertices.

We sequentially choose a path v_0, u_1, \ldots, u_k of vertices through the tree where u_k is a leaf. At each stage i, we choose a vertex u_i and define A_i (the subset of A that we still care about tracking). We also define $C_i(a)$ for each $a \in A$. For each $a \in A$ define $C_1(a) := 1$. Also define

$$C_i := \frac{c \cdot 3^{1/3}}{\Delta} \prod_{a \in A} C_i(a) \text{ and } q_i := C_i 3^{\frac{n - m_i - 1}{3}}, \tag{3.4}$$

where m_i is the number of vertices of $V(F) \setminus \{v\}$ seen by $V(P_{u_{i-1}}) \setminus \{u_0\}$ (thus $n - m_i - 1$ vertices of $V(F) \setminus \{v\}$ are unseen by $V(P_{u_{i-1}}) \setminus \{u_0\}$). So $q_1 = \frac{c \cdot 3^{1/3}}{\Delta} 3^{(n-1)/3}$. Throughout the process we maintain the property that $L(u_i) \geq q_i$ for each i.

We now describe an algorithm that determines our choice of vertices. For $r \ge 1$, let $\epsilon_r = 2^{2(r-1)}\epsilon$.

Vertex Choice Algorithm

We pick $u_1 \in N(v_0)$ such that $L(u_1)$ is maximised, and define $A_1 := A$. Suppose the most recently chosen vertex is u_i and that m_i vertices of $V(F) \setminus \{v\}$ have been seen by $\{u_1, \ldots, u_{i-1}\}$. If u_i is not a leaf; we have two cases:

Case 1: $t(u_i) \in N^4[a]$ for some $a \in A_i$.

Suppose it is the r-th time we have chosen a vertex y such that $t(y) \in N^4[a]$. We have two subcases:

Subcase 1: u_i is good.

In this case u_i has exactly three children y_1, y_2, y_3 .

- (i) If there exists j such that $L(y_j) \ge \frac{1}{3}(1+\epsilon_r)C_i 3^{(n-m_i-1)/3}$ then choose $u_{i+1} := y_j$. - Set $C_{i+1}(a) := (1+\epsilon_r)C_i(a)$ and $C_{i+1}(y) := C_i(y)$, for all $y \in A \setminus \{a\}$. - Set $A_{i+1} := A_i \setminus \{a\}$.
- (ii) Else, every y_j satisfies $L(y_j) > \frac{1}{3}(1 2\epsilon_r)C_i 3^{(n-m_i-1)/3}$. In this case, choose u_{i+1} according to strategy τ_a .

- Set
$$C_{i+1}(a) := (1 - 2\epsilon_r)C_i(a)$$
 and $C_{i+1}(y) := C_i(y)$, for all $y \in A \setminus \{a\}$.

- Set $A_{i+1} := A_i$.

Subcase 2: u_i is bad.

In this case u_i does not have exactly 3 children. Suppose u_i has children y_1, \ldots, y_k for some $k \neq 3$. Pick j such that $L(y_j)$ is maximised and set $u_{i+1} := y_j$.

- Set $C_{i+1}(a) = (1 + \epsilon_r)C_i(a)$ and $C_{i+1}(y) := C_i(y)$, for all $y \in A \setminus \{a\}$.
- Set $A_{i+1} := A_i \setminus \{a\}$.

Case 2: $t(u_i) \notin N^4[a]$ for any $a \in A_i$:

Then v has children $y_1, \ldots y_k$ for some $k \ge 1$. Pick j such that $L(y_j)$ is maximised and set $u_{i+1} = y_j$.

- Set $C_{i+1}(y) := C_i(y)$, for all $y \in A$.
- Set $A_{i+1} := A_i$.

The process terminates when u_i is a leaf.

We now analyse the consequences of choosing vertices in this manner.

Claim 3.10. For each vertex u_i chosen during the Vertex Choice Algorithm, we have $L(u_i) \ge q_i$.

Proof of Claim 3.10. We argue by induction on i; the case i = 1 holds as we chose $u_1 \in N(v_0)$ to maximise $L(u_1)$. Suppose $L(u_i) \ge q_i = C_i 3^{(n-m_i-1)/3}$. Now for the inductive step: we consider each case of the algorithm separately, and prove that the statement holds there.

In Subcase 1(i) we have:

$$L(u_{i+1}) \ge \frac{1}{3}(1+\epsilon_r)C_i 3^{\frac{n-m_i-1}{3}} = C_{i+1} 3^{\frac{n-m_{i+1}-1}{3}} = q_{i+1}.$$

In Subcase 1(ii) we have:

$$L(u_{i+1}) > \frac{1}{3}(1 - 2\epsilon_r)C_i 3^{\frac{n - m_i - 1}{3}} = C_{i+1} 3^{\frac{n - m_{i+1} - 1}{3}} = q_{i+1}.$$

In Subcase 2, recall that u_i has neighbours y_1, \ldots, y_k (for $k \neq 3$) and we pick u_{i+1} to be the y_j which maximises $L(y_j)$. Thus we have:

$$L(u_{i+1}) \ge \frac{C_i}{k} 3^{\frac{n-m_i-1}{3}} = C_i \frac{3^{k/3}}{k} 3^{\frac{n-m_i-k-1}{3}}.$$

The value of $\frac{3^{k/3}}{k}$ is minimised for $k \neq 3$ at k = 2. Thus,

$$L(u_{i+1}) \ge C_i \frac{3^{2/3}}{2} 3^{\frac{n-m_i-k-1}{3}} \ge C_i (1+\epsilon_r) 3^{\frac{n-m_i-k-1}{3}} = C_{i+1} 3^{\frac{n-m_{i+1}-1}{3}} = q_{i+1}.$$

In Case 2, recall that u_i has neighbours y_1, \ldots, y_k and we pick u_{i+1} to be the y_j which maximises $L(y_j)$. Thus we have:

$$L(u_{i+1}) \ge \frac{C_i}{k} 3^{\frac{n-m_i-1}{3}} = C_i \frac{3^{k/3}}{k} 3^{\frac{n-m_i-k-1}{3}} \ge C_i 3^{\frac{n-m_i-k-1}{3}} = C_{i+1} 3^{\frac{n-m_i+1-1}{3}} = q_{i+1},$$

where the last inequality is strict unless k = 3.

We are now equipped to analyse what remains once the algorithm terminates at a leaf u_k . For each $a \in A$ at least one of the following outcomes occurs upon termination of the algorithm.

- (O1) During the algorithm, at a vertex $a \in N^4[a]$, we either chose a branch with a large proportion of leaves via Case 1(i) or we chose a bad vertex via Subcase 2.
- (O2) There is some vertex $w \in N^4[a]$ that is unseen by $V(P_{u_k})$ upon termination of the algorithm.

First observe that for all $s \leq \Delta^5 + 1$,

$$(1 - 2\epsilon_1)(1 - 2\epsilon_2)\dots(1 - 2\epsilon_s)(1 + \epsilon_{s+1}) > (1 + \epsilon).$$
(3.5)

Our choice of ϵ ensures that each factor on the left hand side is greater than zero.

Suppose $a \notin A_k$. Then there exists some j such that $a \in A_j$ but $a \notin A_{j+1}$. Thus at the *j*th stage of the algorithm we had $t(u_j) \in N^4[a]$ and we either chose a branch with a large proportion of leaves via Subcase 1(i) or we chose a bad vertex via Subcase 2. Let $t := |t(\{u_1, \ldots, u_j\}) \cap N^4[a]|$ be the number of vertices, $x \in T$ with $t(x) \in N^4[a]$, chosen up to the *j*th stage. From the algorithm we see

$$C_k(a) = C_{j+1}(a) = (1 - 2\epsilon_1)(1 - 2\epsilon_2)\dots(1 - 2\epsilon_t)(1 + \epsilon_{t+1}).$$
(3.6)

As $|N^4[a]| \leq \Delta^5$, we have $t \leq \Delta^5$, and so by (3.5) we have for $a \notin A_k$

$$C_k(a) > (1+\epsilon). \tag{3.7}$$

Now suppose $a \in A_k$. Let $t := |t(\{u_1, \ldots, u_{k-1}\}) \cap N^4[a]|$. By following the algorithm we see that whenever we are at a vertex $u \in N^4[a]$, we do not pass through Subcase 1(i) or Subcase 2, as this would imply $a \notin A_k$. Thus $C_k(a) = \prod_{i=1}^t (1 - 2\epsilon_i)$. By choice of ϵ , for all $s \leq \Delta^5$ we have

$$3^{1/3} > 1 + 2^{2s}\epsilon$$

so by (3.5) and the observation that $t \leq \Delta^5$,

$$3^{1/3} \cdot C_k(a) > (1+\epsilon).$$
 (3.8)

For each $a \in A_k$, the set $V(P_{u_k})$ does not see all of $N^4[a]$, as we either achieve outcome (O1) or (O2) for a, and if we achieved (O1), then a would not be in A_k . So at termination we have $n - m_k - 1 \ge |A_k|$ and so by the definition of q_k (3.4) we have:

$$q_k \ge C_k 3^{\frac{|A_k|}{3}}.$$
 (3.9)

By (3.7), we have:

$$\prod_{a \in A} C_k(a) \ge (1+\epsilon)^{|A \setminus A_k|} \prod_{a \in A_k} C_k(a),$$

and so by substituting for C_k in (3.9) and applying (3.8), we have:

$$q_k \ge \frac{c \cdot 3^{1/3}}{\Delta} 3^{\frac{|A_k|}{3}} \prod_{a \in A} C_k(a) \ge \frac{c \cdot 3^{1/3}}{\Delta} (1+\epsilon)^{M-|A_k|} 3^{\frac{|A_k|}{3}} \prod_{a \in A_k} C_k(a) \ge \frac{c \cdot 3^{1/3}}{\Delta} (1+\epsilon)^M > 1.$$

This contradicts Claim 3.10, as u_k is a leaf and thus $L(u_k) = 1$. Thus |A| < M, concluding the proof of Lemma 3.9.

Corollary 3.11. Fix c > 0. Let F be an n-vertex graph with $f(F) \ge 2c \cdot 3^{n/3}$ and $\Delta := \Delta(F) > 1$. Then for any $v \in V(F)$, there exists a constant $C = C(\Delta, c)$ such that at most C vertices are atypical for T(v), the exploration tree from v in F.

Proof. Let A be the set of vertices that are atypical for T. Let U be the largest subset of A such that for all $a, a' \in U$, we have $N^4[a] \cap N^4[a'] = \emptyset$. As $|N^4[x]| \leq \Delta^5$ for every $x \in V(F)$, we have

$$|U| \ge \frac{|A|}{\Delta^5}.\tag{3.10}$$

We wish to apply Lemma 3.9 to F, T and U.

As $f(F) \ge 2c \cdot m(n)$, by Lemma 3.2 $f_v(F) \ge \frac{c}{10}m(n)$ for all $v \in F$. Combining this with Lemma 3.7 gives

$$L(v_0) \ge 2f_v(F) \ge \frac{c}{5}m(n) \ge \frac{c}{20}3^{n/3},$$

for some constant c' and where the last inequality follows from (3.3).

Applying Lemma 3.9 shows that |U| < M, where M is the smallest integer such that $\frac{c}{20} \cdot 3^{1/3} (1 + 2^{-\Delta^{100}})^M > \Delta$. Combining this with (3.10) gives the required result.

Let $B = (B_1, \ldots, B_k)$ be a braid in G_{max} . If $|B_i| = 3$ for all *i*, we call *B* a 3-braid. For a braid *B* of length at least 4, we say that an induced cycle passes through *B* if it contains a vertex from every cluster of *B*. Call a braid maximal if it is not contained in any longer braid. The following simple deduction will be used.

Lemma 3.12. There exists a constant C such that G_{max} contains at most C maximal 3braids and a 3-braid B such that $|V(B)| = \Omega(n)$. Moreover, for any 3-braid B' on rn vertices, at least $f(H_n) (1 - 3^{-rn/6})$ induced cycles in G_{max} pass through B'.

Proof. Let $v \in V(G_{max})$. The only vertices which can be contained in more than one maximal 3-braid lie in end clusters. By Lemma 3.5, every v-typical vertex is contained in a central cluster of exactly one maximal 3-braid. So any vertex in the end cluster of a maximal 3-braid is v-atypical. By Corollary 3.11, there exists a constant c such that at most c vertices are

v-atypical for G_{max} . Each of these vertices is contained in at most $\Delta(G_{max}) \leq 30$ maximal 3braids. So G_{max} contains (crudely) at most 30*c* maximal 3-braids, proving the first statement of the lemma.

The union of the maximal 3-braids in G_{max} contains all the typical vertices and so it contains at least n/2 vertices for large n. Therefore, when n is sufficiently large, as there are at most 30c maximal 3-braids some 3-braid $B = (B_1, \ldots, B_k)$ contains $\Omega(n)$ vertices.

For the final claim, observe that if an induced cycle does not pass through a 3-braid $\mathcal{B}' = (B'_1, \ldots, B'_k)$ on rn vertices, then it is either a C_4 contained in B (there are at most $O(n^4)$ of these), or it is contained in $V(G_{max}) \setminus \bigcup_{i=3}^{k-2} B'_i$ (by Lemma 3.1, there are at most $[(1-r)n+12]\binom{30}{2}3^{[(1-r)n+9]/3}$ of these). Therefore at most

$$[(1-r)n+12]\binom{30}{2}3^{[(1-r)n+9]/3} + O(n^4)$$

induced cycles of G_{max} do not pass through B. So for n_0 sufficiently large, at least

$$f(H_n) \left(1 - 3^{-rn/6}\right)$$

induced cycles pass through B.

The next lemma shows that G_{max} is a cyclic braid. It will remain to determine the cluster sizes and whether there are edges within the clusters of G_{max} .

Lemma 3.13. G_{max} is a cyclic braid.

Proof. Let $B := (B_1, \ldots, B_{Cn/3})$ be the longest 3-braid in G_{max} . Let Q be the number of induced cycles in G_{max} that pass through B. By Lemma 3.12,

$$Q \ge f(H_n) \left(1 - 3^{-Cn/6} \right). \tag{3.11}$$

Now let $G' := G[V(G_{max}) \setminus \bigcup_{i=2}^{Cn/3-1} B_i]$. Let x and y be two new vertices and define H to be the graph on vertex set $V(H) := V(G') \cup \{x, y\}$, and edge set

 $E(H) := E(G') \cup \{xb : b \in B_1\} \cup \{yb : b \in B_{Cn/3}\}.$

We have

$$Q = 3^{(Cn-6)/3} p_2(H; x, y).$$
(3.12)

Combining (3.11) and (3.12) gives

$$p_2(H; x, y) \ge 3^{-(Cn-6)/3} \cdot f(H_n) \left(1 - 3^{-Cn/6}\right).$$
 (3.13)

We now focus on the structure of H. Let us call a central cluster C of a maximal 3-braid *B* supercentral if for any $x \in C$ and y in an end cluster of B, $d(x, y) \geq 5$. Define a new graph H' via the following process.

- Set $F_1 := H$.
- Let *i* be maximal such that we have defined F_i . Suppose there exists a vertex $v_i \in F_i$, contained in a supercentral cluster C_i of a maximal 3-braid M_i , where C_i is adjacent to clusters C_1^i and C_2^i . Then define F_{i+1} to be the graph obtained from F_i by deleting C_i and adding every edge $\{uw : u \in C_1^i, w \in C_2^i\}$.
- If there exists no such vertex v_i , define $H' := F_i$.

The process will terminate as H has a finite number of vertices. Observe that F_{i+1} is a braid if and only if F_i is a braid. In addition, when H' is a braid this process can be reversed to find H. We now show that H' is a braid.

Any v-typical vertex in F_i that does not get deleted during the process is v-typical in F_{i+1} . Hence any v-typical vertex in G_{max} that does not get deleted is v-typical in H'. By Lemma 3.12, there exists a constant a such that G_{max} contains a maximal 3-braids. O(a) vertices from each of these braids will remain in H' when the process terminates. Any vertex not contained in a 3-braid in G_{max} is v-atypical in G_{max} . By Corollary 3.11 there exists a constant b such that there are at most b such vertices. As H' contains all the atypical vertices of G_{max} and at most O(a) vertices from each 3-braid in G_{max} , there exists a constant β such that $|V(H')| \leq \beta$.

At stage *i* of the process, F_{i+1} contains all induced cycles of F_i that do not pass through M_i and a third of the number of cycles in F_i that do pass through M_i . Thus we have

$$p_2(F_{i+1}; x, y) \ge 3^{-1} \cdot p_2(F_i; x, y),$$
(3.14)

and so

$$p_2(H'; x, y) \ge 3^{-(|V(H)| - |V(H')|)/3} \cdot p_2(H; x, y),$$
(3.15)

Combining (3.13) and (3.15) and observing that |V(H)| = (1 - C)n + 8 gives

$$p_2(H'; x, y) \ge 3^{(-n-2+|V(H')|)/3} \cdot f(H_n) \left(1 - 3^{-Cn/6}\right).$$
(3.16)

As

$$3^{(-n-2+|V(H')|)/3}f(H_n) = f_2(|V(H')|) + o(1),$$

when n_0 is sufficiently large we have

$$3^{(-n-2+|V(H')|)/3} \cdot f(H_n) \left(1 - 3^{-Cn/6}\right) > f_2(|V(H')|) - 1.$$
(3.17)

As $p_2(H'; x, y)$ is an integer, by taking n_0 to be sufficiently large, (3.16) and (3.17) give

$$p_2(H'; x, y) \ge f_2(|V(H')|)$$

Therefore, by Theorem 2.1, H' is isomorphic to a graph in $\mathcal{F}_{|V(H')|}$. Thus H' is a braid. By reversing the process applied above (adding back in the supercentral clusters) to recreate H from H', we see that H is a graph in $\mathcal{F}_{|V(H)|}$, and hence G_{max} is a cyclic braid. **Corollary 3.14.** We have the following:

- when $n \equiv 0$ modulo 3, G_{max} has exactly n/3 clusters of size 3;
- when n ≡ 1 modulo 3, G_{max} has either one cluster of size 4 and (n-4)/3 of size 3, or two of size two and (n-4)/3 of size 3;
- when $n \equiv 2 \mod 3$, G_{\max} has exactly one cluster of size 2 and (n-2)/3 of size 3.

We are now in a position to complete the proof of Theorem 1.2. The next lemma shows that the clusters in G_{max} do not contain any edges, and thus we will prove the required result for $n \equiv 0$ or 2 modulo 3. In the remaining case we will need a short argument to decide whether the graph contains two clusters of size two, or one of size four. In both cases, the arguments are essentially routine checks.

Lemma 3.15. When $n \equiv 0, 2$ modulo 3, no cluster of G_{max} contains edges.

Proof. First observe, that if e is an edge within a cluster, the only induced cycles containing e can be triangles, either contained within the cluster, or containing exactly one vertex from a neighbouring cluster; or induced copies of C_4 within the cluster (in the case that the cluster contains 4 vertices).

Let B be a cluster adjacent to clusters A and C. Suppose there exists an edge e = uvwhere $u, v \in V(B)$. The edge e is contained in at most |A| + |C| + (|B| - 2) induced cycles within G_{max} . The graph $G' = G_{max} \setminus \{e\}$ will contain at least |A||C| induced copies of C_4 (for any $x \in A, y \in C$, the set $\{x, y, u, v\}$ induces a C_4) that are not induced cycles in G_{max} .

As G_{max} does not contain both a cluster of size 2 and a cluster of size 4, we have

$$|A||C| > |A| + |C| + (|B| - 2),$$

unless |B| = 3 and at least one of |A| or |C| is equal to 2. Except for this case, the number of induced cycles in $G' = G_{max} \setminus \{e\}$ is greater than the number of induced cycles in G_{max} , a contradiction.

Now suppose |B| = 3 and suppose without loss of generality that $A = \{a_1, a_2\}$. First consider the case where |C| = 3. Suppose B contains an edge e = uv. This edge is contained in at most 6 triangles in G_{max} . By the above argument, A does not contain an edge. The graph $G' = G_{max} \setminus \{e\}$ will contain at least 7 induced copies of C_4 that are not induced cycles in G_{max} (for any $x \in A, y \in C$, the sets $\{x, y, u, v\}$ and $\{a_1, a_2, u, v\}$ induce copies of C_4). Thus $f(G') > f(G_{max})$, a contradiction.

The remaining case to consider is when $C = \{c_1, c_2\}$. If *B* contains an edge *e*, this edge is contained in at most 5 triangles in G_{max} . The graph $G' = G_{max} \setminus \{e\}$ contains at least 6 induced copies of C_4 that are not induced cycles in G_{max} . Thus $f(G') > f(G_{max})$, a contradiction. So no cluster in G_{max} contains an edge.

We have proved Theorem 1.2 in the cases where $n \equiv 0$ or 2 modulo 3. It remains to prove the result in the case $n \equiv 1$ modulo 3.

Lemma 3.16. When $n \equiv 1$ modulo 3, G_{max} is isomorphic to H_n .

Proof. By Corollary 3.14 and Lemma 3.15, we know that G_{max} is one of two empty cyclic braids. One possibility is that it is isomorphic to H_n . The other possibility is that G_{max} is an empty cyclic braid G_2 with exactly two clusters of size 2, and the rest of size 3. An induced cycle in H_n or G_2 either contains exactly one vertex from each cluster, or is an induced copy of C_4 . In both H_n and G_2 , the number of induced cycles containing exactly one vertex from each cluster is $4 \cdot 3^{(n-4)/3}$. Thus the only difference in $f(H_n)$ and $f(G_2)$ comes from the number of induced copies of C_4 .

There are two types of C_4 . Type 1 contains vertices from exactly two clusters. Type 2 contains vertices from three clusters. The graph H_n contains 3(n+5) induced type 1 cycles; G_2 contains at most 3n-14 of this form (fewer if the two clusters of size 2 are not adjacent). The graph H_n contains 9(n+4) induced type 2 cycles; G_2 contains at most 9n-42 of this form. Thus H_n contains more induced cycles than G_2 and therefore G_{max} is isomorphic to H_n .

4 Proof of Theorem 1.5

The proof of Theorem 1.5 follows the same lines as the proof of Theorem 1.2. Before proceeding with the details of the proof, we first give an outline of what is to come. Let $0 < \alpha < 1$ be any constant and let F be an *n*-vertex graph containing at least $\alpha \cdot m(n)$ induced cycles. We will show that it is possible to delete a constant number of vertices from F to give a graph F' with maximum degree bounded by a constant. Applying Lemma 3.9 to F' then shows that the number of atypical vertices in F' is bounded by a constant. The result will immediately follow. We cannot simply apply Lemma 3.9 to F, as F may contain vertices of arbitrarily large degree.

Lemma 4.1. Fix $0 < \alpha < 1$ and define $\Delta^* = \Delta^*(\alpha)$ to be the smallest integer such that $6\binom{\Delta^*}{2}3^{(1-\Delta^*)/3} < \alpha \cdot 4 \cdot 3^{-4/3}$. For n sufficiently large, let F be an n-vertex graph with $f(F) \ge \alpha \cdot m(n)$. Then there exists a constant $C = C(\alpha)$ such that deleting edges incident to C vertices of F gives a graph H with $\Delta(H) \le \Delta^*$. Moreover, $f(H) \ge \frac{\alpha}{2} \cdot m(n)$.

Proof. Suppose that $\Delta(F) > \Delta^*$ (else we are trivially done). We create a new graph H, with $\Delta(H) \leq \Delta^*$, in the following manner. Define $F_1 := F$. Let *i* be maximal such that F_i has been defined. If there exists a vertex $v_i \in V(F_i)$ with $d(v_i) > \Delta^*$ then define $F_{i+1} := F_i \setminus \{v_i\}$. This process will terminate as F has a finite number of vertices. Suppose the process terminates at a graph F_j . We have $\Delta(F_j) \leq \Delta^*$. Define $H := F_j$.

To prove the first statement of the lemma it suffices to show that there exists some constant $C := C(\alpha)$ such that j = C. To prove this, we will bound the size of f(F) - f(H) and use this to show that when j > C, we have $f(F) < \alpha \cdot m(n)$, a contradiction.

By Lemma 3.1,

$$f_{v_i}(F_i) \le {\binom{d_{F_i}(v_i)}{2}} 3^{(n-i-d_{F_i}(v_i)-1)/3} \le {\binom{\Delta^*}{2}} 3^{(n-i-\Delta^*-1)/3},$$

where the second inequality follows as the function $\binom{x}{2}3^{-x/3}$ is decreasing for $x \ge 6$. So

$$f(H) = f(F) - \sum_{i=1}^{j-1} f_{v_i}(F_i)$$

$$\geq f(F) - {\Delta^* \choose 2} 3^{(n-\Delta^*+1)/3} \sum_{i=1}^{j-1} 3^{-i/3}$$

$$\geq f(F) - 3{\Delta^* \choose 2} 3^{(n-\Delta^*+1)/3}.$$
(4.1)

As |V(H)| = n - j + 1, for some constant c we have $f(H) \le c \cdot 3^{(n-j+1)/3}$ by (3.3). Combining this with (4.1) gives:

$$f(F) \le c \cdot 3^{(n-j+1)/3} + 3\binom{\Delta^*}{2} 3^{(n-\Delta^*+1)/3}.$$
 (4.2)

There exists a constant C such that, whenever j > C and n is sufficiently large:

$$c \cdot 3^{(n-j+1)/3} < \frac{1}{2} (\alpha \cdot 4 \cdot 3^{-4/3}) 3^{n/3}.$$
 (4.3)

Suppose that $j \ge C$ and let n be sufficiently large. Using the definition of Δ^* and substituting (4.3) into (4.2), gives

$$f(F) < \alpha \cdot 4 \cdot 3^{-4/3} 3^{n/3} < \alpha \cdot m(n),$$

where the final inequality is implied by (3.1). This contradicts the hypothesis that $f(F) \ge \alpha \cdot m(n)$. Therefore j < C, completing the proof of the first statement of the lemma.

We now prove the second statement. By (4.1) we have

$$f(H) \ge f(F) - 3\binom{\Delta^*}{2} 3^{(n-\Delta^*+1)/3}$$

Given the definition of Δ^* , we have $f(H) \geq \frac{\alpha}{2}m(n)$.

Proof of Theorem 1.5. Let F be an n-vertex graph containing at least $\alpha \cdot m(n)$ induced cycles. By Lemma 4.1, there exist constants $c = c(\alpha)$ and $\Delta^* = \Delta^*(\alpha)$ such that deleting cvertices from F gives a graph F' with $\Delta(F') \leq \Delta^*$ and $f(F') \geq \frac{\alpha}{2}m(n)$. For any $v \in V(F')$, by Lemma 3.2 we have $f_v(F) \geq \frac{\alpha}{20}$. Let A be the set of v-atypical vertices in F'. By applying Lemma 3.9 we deduce that |A| < M (where $M = M(\alpha)$ is defined as in Lemma 3.9). By Lemma 3.5, every v-typical vertex in F' is contained in a central cluster of exactly one maximal 3-braid. We obtain H from F' by adding and deleting edges incident to vertices in A. The result follows with $C(\alpha) = c + M$.

5 Induced odd or even cycles

In this section we prove Theorem 1.6. The proofs of Theorem 1.7 and Theorem 1.8 closely follow that of Theorem 1.6.

Given a graph G, define $f_o(G)$ to be the number of induced odd cycles contained within G. Similarly, for $v \in G$, define $f_o^v(G)$ to be the number of induced odd cycles in G that contain v. We have:

$$m_o(n) \ge f_o(G_n) = \Omega(3^{n/3}),$$
(5.1)

where G_n is defined as in Section 1. The proof of Theorem 1.6 follows from Theorem 1.5 and some arguments analogous to those used in Theorem 1.2. For the latter, we refer back to Sections 2 and 3 where necessary. The main difference is that, instead of applying Theorem 2.1, we use Theorem 2.9.

We fix a large constant n_0 and let G be a graph on $n \ge n_0$ vertices that contains $m_o(n)$ induced odd cycles. In what follows we let n_0 be sufficiently large when required and we make no attempts to optimise the constants given in our argument.

Sketch proof of Theorem 1.6. We first show that $\Delta(G) \leq 35$ using analogous arguments to those in Theorem 1.2.

Lemma 3.2 holds (as (5.1) gives us the analogous bound to (3.1) that we need). Thus every vertex is contained in at least $\frac{1}{20}m_o(n)$ induced odd cycles. Thus we have

$$m_o(n+1) \ge \left(1 + \frac{1}{20}\right) m_o(n),$$
 (5.2)

as in (3.2).

We use the same argument as in Lemma 3.3, replacing m(n) with $m_o(n)$, to show that $\Delta(G) \leq 35$ (we get a different value for Δ as we use the lower bound $m_o(G_n) \geq 3^{(n-8)/3}$ and this differs from the lower bound used for m(n)).

By (5.1) we have $f_o(G_n) = \Omega(3^{n/3})$. Thus applying Theorem 1.5 shows that there exists a constant c such that adding and deleting edges incident to c vertices of G gives a cyclic braid H with the same cluster sizes as H_n . Using this and the knowledge that $\Delta(G)$ is bounded by a fixed constant, it is seen that G contains a 3-braid B of even length such that |V(B)| = rn for some constant r.

We now show that G is a cyclic braid. We use essentially the same argument as in Lemma 3.13 with $f_o(G_n)$ in place of $f(H_n)$. However, when applying the process of deleting central clusters, we delete a pair of adjacent clusters at a time (to maintain the count of odd cycles). We again reach a graph H' such that there exists a constant β with $|V(H')| = \beta$. We make the analogous deductions from there to reach the bound

$$f_2^o(n) \le p_2^o(H'; x, y).$$

We then apply Theorem 2.9 to determine that $H' \in \mathcal{F}^{o}_{|V(H')|}$. Reversing the process of deleting central clusters to obtain H from H', we get that $H \in \mathcal{F}^{o}_{|V(H)|}$. Therefore G is a cyclic braid, with clusters all of size 3 except:

- three clusters of size 2, when $n \equiv 0 \mod 6$;
- two clusters of size 2, when $n \equiv 1 \mod 6$;
- one cluster of size 2, when $n \equiv 2 \mod 6$;
- a single cluster of size 4, when $n \equiv 4 \mod 6$; and
- either two clusters of size 4 or four clusters of size 2, when $n \equiv 5 \mod 6$.

It remains to determine whether there are edges within the clusters, the relative positions of the clusters in the cyclic braid (in the cases where more than one cluster does not have size 3) and, in the case $n \equiv 5$ modulo 6, to determine the precise cluster sizes. Using arguments of a similar nature to those in Lemma 3.15 and Lemma 3.16, it can be checked that $G \cong G_n$ for every value of $n \ge n_0$.

Theorem 1.6 determines which n-vertex graphs contain the maximum number of odd cycles. Following essentially the same argument we prove Theorem 1.7 and Theorem 1.8, which determine the family of n-vertex graphs that contain the maximum number of odd holes or even holes respectively.

Sketch proof of Theorem 1.7 and Theorem 1.8. We use the same argument as in the proof of Theorem 1.6. In the case of Theorem 1.8, the argument can be modified to consider even induced cycles rather than odd. The main difference is at the final stage, where we know G is a cyclic braid and the possible cluster sizes in G. Changing the positions of clusters and edges within clusters can only affect the holes that do not contain a vertex from every cluster. Thus any hole that can be affected has size 3 or 4.

For the odd hole case, the positions of the clusters and the existence of edges within clusters will not alter the number of odd holes (as any induced cycle with size 3 or 4 is not an odd hole). In the even case, a simple check shows the number of holes of size 4 (given the cluster sizes) is maximised when G is isomorphic to E_n .

6 Conclusion

For sufficiently large n, we have determined precisely which graphs on n vertices contain the maximum number of induced cycles, the maximum number of odd or even induced cycles, and the maximum number of holes. However, there are a number of interesting related questions.

In our proofs above we make no attempts to optimise the value of n_0 . We know that in some small cases, H_n does not contain the maximum number of induced cycles [10, 13]. We believe Theorem 1.2 ought to be true for $n_0 = 30$, but our proof gives a much larger number. There are several places where we could improve the bound, most notably by choosing a more careful strategy in Lemma 3.9. However we omit the details as the bound would still be extremely large. It is natural to consider induced cycles of some length that depends on n. Let c(n, l) be the maximum number of length l induced cycles that can be contained in a graph on n vertices. Let C(n, l) be the set of graphs containing c(n, l) induced cycles of length l.

Question 6.1. For l = l(n), what is C(n, l)?

When l is linear we believe the following should hold.

Conjecture 6.2. Fix $c \in (0,1)$. If $l(n) = \lceil cn \rceil$, then for sufficiently large n the only graphs in C(n,l) are cyclic braids of length l.

Perhaps a similar result holds down to cycles of length $\Omega(\sqrt{n})$.

Question 6.3. Suppose $l(n) > \sqrt{n}$. For sufficiently large *n*, are all graphs in C(n, l) cyclic braids?

Another related question is to ask about induced subgraphs which are subdivisions of some fixed graph H.

Question 6.4. Given a fixed finite graph H, what is the maximum number of induced subdivisions of H that can be contained in a graph on n vertices (and which graphs realise this maximum)?

Theorem 1.2 answers this question for $H = C_3$, but what happens for other graphs? For instance, which graphs maximise the number of induced subdivisions of $K_{1,3}$? For large n, are the extremal graphs always blowups of some subdivision of H? The rooted version of the question is also interesting, where we consider induced subdivisions of H where the branch vertices are fixed (for instance Theorem 2.1 is a result of this form for $H = K_2$).

Finally, we remark that the related problem of finding the graph on n vertices that contains the most cycles (not necessarily induced) is trivial as K_n is the extremal graph. However, the problem becomes interesting when we forbid certain subgraphs (see Arman, Gunderson and Tsaturian [1] and Morrison, Roberts and Scott [11]).

7 Acknowledgments

We would like to thank Brendan McKay and Mike Robson for helpful discussions. We would also like to thank Andrew Treglown for bringing [15] to our attention and the referees for their helpful comments.

References

[1] A. Arman, D. S. Gunderson, and S. Tsaturian, *Triangle-free graphs with the maximum number of cycles*, Discrete Math. **339** (2016), no. 2, 699–711.

- [2] J. Balogh, P. Hu, B. Lidický, and F. Pfender, Maximum density of induced 5-cycle is achieved by an iterated blow-up of 5-cycle, European J. Combin. 52 (2016), no. part A, 47–58.
- [3] B. Bollobás, Y. Egawa, A. Harris, and G. P. Jin, *The maximal number of induced r-partite subgraphs*, Graphs Combin. **11** (1995), no. 1, 1–19.
- [4] B. Bollobás, C. Nara, and S. Tachibana, The maximal number of induced complete bipartite graphs, Discrete Math. 62 (1986), no. 3, 271–275.
- [5] J. I. Brown and A. Sidorenko, The inducibility of complete bipartite graphs, J. Graph Theory 18 (1994), no. 6, 629–645.
- [6] C. Even-Zohar and N. Linial, A note on the inducibility of 4-vertex graphs, Graphs Combin. 31 (2015), no. 5, 1367–1380.
- [7] G. Exoo, Dense packings of induced subgraphs, Ars Combin. 22 (1986), 5–10.
- [8] H. Hatami, J. Hirst, and S. Norine, *The inducibility of blow-up graphs*, J. Combin. Theory Ser. B **109** (2014), 196–212.
- [9] J. Hirst, The inducibility of graphs on four vertices, J. Graph Theory 75 (2014), no. 3, 231–243.
- [10] Brendan McKay, personal communication.
- [11] N. Morrison, A. Roberts, and A. Scott, *Maximising the number of cycles in H-free graphs*, In preparation.
- [12] N. Pippenger and M. C. Golumbic, *The inducibility of graphs*, J. Combinatorial Theory Ser. B **19** (1975), no. 3, 189–203.
- [13] Mike Robson, personal communication.
- [14] Z. Tuza, Problems and results on graph and hypergraph colorings, Matematiche (Catania) 45 (1990), no. 1, 219–238 (1991), Graphs, designs and combinatorial geometries (Catania, 1989).
- [15] _____, Unsolved combinatorial problems., BRICS Lecture Series, LS-01-1 (2001), (available from http://www.brics.dk/publications/).
- [16] _____, *Problems on cycles and colorings*, Discrete Math. **313** (2013), no. 19, 2007–2013.