

# Minimal Stable Sets in Tournaments

Felix Brandt

Technische Universität München

85748 Garching bei München, Germany

brandtf@in.tum.de

We propose a systematic methodology for defining tournament solutions as extensions of maximality. The central concepts of this methodology are maximal qualified subsets and minimal stable sets. We thus obtain an infinite hierarchy of tournament solutions, encompassing the top cycle, the uncovered set, the Banks set, the minimal covering set, the tournament equilibrium set, the Copeland set, and the bipartisan set. Moreover, the hierarchy includes a new tournament solution, the *minimal extending set*, which is conjectured to refine both the minimal covering set and the Banks set.

## 1 Introduction

Given a finite set of alternatives and choices between all pairs of alternatives, how to choose from the entire set in a way that is faithful to the pairwise comparisons? This simple, yet captivating, problem is studied in the literature on tournament solutions. A tournament solution thus seeks to identify the “best” elements according to some binary dominance relation, which is usually assumed to be asymmetric and complete. Since the ordinary notion of maximality may return no elements due to cyclical dominations, numerous alternative solution concepts have been devised and axiomatized (see, e.g., Moulin, 1986; Laslier, 1997). Tournament solutions have numerous applications throughout economics, most prominently in social choice theory where the dominance relation is typically defined via majority rule (e.g., Fishburn, 1977; Bordes, 1983). Other application areas include multi-criteria decision analysis (e.g., Arrow and Raynaud, 1986; Bouyssou et al., 2006), non-cooperative game theory (e.g., Fisher and Ryan, 1995; Laffond et al., 1993b; Duggan and Le Breton, 1996), and cooperative game theory (Gillies, 1959; Brandt and Harrenstein, 2010).

In this paper, we approach the tournament choice problem using a methodology consisting of two layers: *qualified subsets* and *stable sets*. Our framework captures most known tournament solutions (notable omissions are the Slater set and the Markov set)

and allows us to provide unified proofs of properties and inclusion relationships between tournament solutions.

In Section 2, we introduce the terminology and notation required to handle tournaments and define six standard properties of tournament solutions: monotonicity (MON), independence of unchosen alternatives (IUA), the weak superset property (WSP), the strong superset property (SSP), composition-consistency (COM), and irregularity (IRR). The remainder of the paper is then divided into four sections.

**Qualified Subsets** (Section 3) The point of departure for our methodology is to collect the maximal elements of so-called qualified subsets, i.e., distinguished subsets that admit a maximal element. In general, families of qualified subsets are characterized by three properties (closure, independence, and fusion). Examples of families of qualified subsets are all subsets with at most two elements, all subsets that admit a maximal element, or all transitive subsets. Each family yields a corresponding tournament solution and we thus obtain an infinite hierarchy of tournament solutions. The tournament solutions corresponding to the three examples given above are the set of *all alternatives except the minimum*, the *uncovered set* (Fishburn, 1977; Miller, 1980), and the *Banks set* (Banks, 1985). Our methodology allows us to easily establish a number of inclusion relationships between tournament solutions defined via qualified subsets (Proposition 2) and to prove that all such tournament solutions satisfy WSP and MON (Proposition 1). Based on an axiomatic characterization using minimality and a new property called strong retentiveness, we show that the Banks set is the finest tournament solution definable via qualified subsets (Theorem 1).

**Stable Sets** (Section 4) Generalizing an idea by Dutta (1988), we then propose a method for refining any suitable solution concept  $S$  by defining minimal sets that satisfy a natural stability criterion with respect to  $S$ . A crucial property in this context is whether  $S$  always admits a *unique* minimal stable set. For tournament solutions defined via qualified subsets, we show that this is the case if and only if no tournament contains two disjoint stable sets (Lemma 2). As a consequence of this characterization and a theorem by Dutta (1988), we show that an infinite number of tournament solutions (defined via qualified subsets) always admit a unique minimal stable set (Theorem 3). Moreover, we show that all tournament solutions defined as unique minimal stable sets satisfy WSP and IUA (Proposition 4), SSP and various other desirable properties if the original tournament solution is defined via qualified subsets (Theorem 4), and MON and COM if the original tournament solution satisfies these properties (Proposition 5 and Proposition 6). The minimal stable sets with respect to the three tournament solutions mentioned in the paragraph above are the *minimal dominant set*, better known as the *top cycle* (Good, 1971; Smith, 1973), the *minimal covering set* (Dutta, 1988), and a new tournament solution that we call the *minimal extending set (ME)*. Whether *ME* satisfies uniqueness turns out to be a highly non-trivial combinatorial problem and remains open. If true, *ME* would be contained in both the minimal covering set and the Banks set while satisfying a number of desirable properties. We conclude the section

by axiomatically characterizing all tournament solutions definable via unique minimal stable sets (Proposition 7).

**Retentiveness and Stability** (Section 5) *ME* bears some resemblance to Schwartz’s *tournament equilibrium set TEQ* (Schwartz, 1990), which is defined as the minimal *retentive* set of a tournament. There are some similarities between retentiveness and stability and, as in the case of *ME*, the uniqueness of a minimal retentive set and thus the characteristics of *TEQ* remain an open problem (Laffond et al., 1993a; Houy, 2009). We show that Schwartz’s conjecture is stronger than ours (Theorem 6) and has a number of interesting consequences such as that *TEQ* can also be represented as a minimal stable set (Theorem 7) and is strictly contained in *ME* (Corollary 2).

**Quantitative Concepts** (Section 6) We also briefly discuss a quantitative version of our framework, which considers qualified subsets that are maximal in terms of cardinality rather than set inclusion. We thus obtain the *Copeland set* (Copeland, 1951) and—using a slightly modified definition of stability—the *bipartisan set* (Laffond et al., 1993b). This completes our picture of tournament solutions and their corresponding minimal stable sets as given in Table 1 on page 25.

## 2 Preliminaries

The core of the problem studied in the literature on tournament solutions is how to extend choices in sets consisting of only two elements to larger sets. Thus, our primary objects of study will be functions that select one alternative from any pair of alternatives. Any such function can be conveniently represented by a tournament, i.e., a binary relation on the entire set of alternatives. Tournament solutions then advocate different views on how to choose from arbitrary subsets of alternatives based on these pairwise comparisons (see, e.g., Laslier, 1997, for an excellent overview of tournament solutions).

### 2.1 Tournaments

Let  $X$  be a universe of alternatives. The set of all *finite* subsets of set  $X$  will be denoted by  $\mathcal{F}_0(X)$  whereas the set of all *non-empty* finite subsets of  $X$  will be denoted by  $\mathcal{F}(X)$ . A (finite) *tournament*  $T$  is a pair  $(A, \succ)$ , where  $A \in \mathcal{F}(X)$  and  $\succ$  is an asymmetric and complete (and thus irreflexive) binary relation on  $X$ , usually referred to as the *dominance relation*.<sup>1</sup> Intuitively,  $a \succ b$  signifies that alternative  $a$  is preferable to  $b$ . The dominance relation can be extended to sets of alternatives by writing  $A \succ B$  when  $a \succ b$  for all  $a \in A$  and  $b \in B$ . When  $A$  or  $B$  are singletons, we omit curly braces to improve readability. We further write  $\mathcal{T}(X)$  for the set of all tournaments on  $X$ .

---

<sup>1</sup>This definition slightly diverges from the common graph-theoretic definition where  $\succ$  is defined on  $A$  rather than  $X$ . However, it facilitates the sound definition of tournament functions (such as tournament solutions or concepts of qualified subsets).

For a set  $B$ , a relation  $R$ , and an element  $a$ , we denote by  $D_{B,R}(a)$  the *dominion* of  $a$  in  $B$ , i.e.,

$$D_{B,R}(a) = \{b \in B \mid a R b\},$$

and by  $\overline{D}_{B,R}(a)$  the *dominators* of  $a$  in  $B$ , i.e.,

$$\overline{D}_{B,R}(a) = \{b \in B \mid b R a\}.$$

Whenever the tournament  $(A, \succ)$  is known from the context and  $R$  is the dominance relation  $\succ$  or  $B$  is the set of all alternatives  $A$ , the respective subscript will be omitted to improve readability.

The *order* of a tournament  $T = (A, \succ)$  refers to the cardinality of  $A$ . A tournament  $T = (A, \succ)$  is called *regular* if  $|D(a)| = |D(b)|$  for all  $a, b \in A$ .

Let  $T = (A, \succ)$  and  $T' = (A', \succ')$  be two tournaments. A *tournament isomorphism* of  $T$  and  $T'$  is a bijective mapping  $\pi : A \rightarrow A'$  such that  $a \succ b$  if and only if  $\pi(a) \succ' \pi(b)$ . A tournament  $T = (A, \succ)$  is called *homogeneous* (or vertex-transitive) if for every pair of alternatives  $a, b \in A$  there is a tournament isomorphism  $\pi : A \rightarrow A$  of  $T$  such that  $\pi(a) = b$ .

## 2.2 Components and Decompositions

An important structural concept in the context of tournaments is that of a *component*. A component is a subset of alternatives that bear the same relationship to all alternatives not in the set.

Let  $T = (A, \succ)$  be a tournament. A non-empty subset  $B$  of  $A$  is a *component* of  $T$  if for all  $a \in A \setminus B$  either  $B \succ a$  or  $a \succ B$ . A *decomposition* of  $T$  is a set of pairwise disjoint components  $\{B_1, \dots, B_k\}$  of  $T$  such that  $A = \bigcup_{i=1}^k B_i$ . Given a particular decomposition  $\tilde{B} = \{B_1, \dots, B_k\}$  of  $T$ , the *summary* of  $T$  is defined as the tournament on the individual components rather than the alternatives. Formally, the summary  $\tilde{T} = (\tilde{B}, \tilde{\succ})$  of  $T$  is the tournament such that for all  $i, j \in \{1, \dots, k\}$  with  $i \neq j$ ,

$$B_i \tilde{\succ} B_j \quad \text{if and only if} \quad B_i \succ B_j.$$

Conversely, a new tournament can be constructed by replacing each alternative with a component. For notational convenience, we tacitly assume that  $\mathbb{N} \subseteq X$ . For pairwise disjoint sets  $B_1, \dots, B_k \subseteq X$  and tournaments  $\tilde{T} = (\{1, \dots, k\}, \tilde{\succ})$ ,  $T_1 = (B_1, \succ_1)$ ,  $\dots$ ,  $T_k = (B_k, \succ_k)$ , the *product* of  $T_1, \dots, T_k$  with respect to  $\tilde{T}$ , denoted by  $\Pi(\tilde{T}, T_1, \dots, T_k)$ , is a tournament  $(A, \succ)$  such that  $A = \bigcup_{i=1}^k B_i$  and for all  $b_1 \in B_i, b_2 \in B_j$ ,

$$b_1 \succ b_2 \quad \text{if and only if} \quad i = j \text{ and } b_1 \succ_i b_2, \text{ or } i \neq j \text{ and } i \tilde{\succ} j.$$

Components can also be used to simplify the graphical representation of tournaments. We will denote components by gray circles. Furthermore, omitted edges in figures that depict tournaments are assumed to point downwards or from left to right by convention (see, e.g., Figure 1).

### 2.3 Tournament Functions

A central aspect of this paper will be functions that, for a given tournament, yield one or more subsets of alternatives. We will therefore define the notion of a *tournament function*. A function on tournaments is a tournament function if it is independent of outside alternatives and stable with respect to tournament isomorphisms. A tournament function may yield a (non-empty) subset of alternatives—as in the case of tournament solutions—or a set of subsets of alternatives—as in the case of qualified or stable sets.

**Definition 1.** Let  $Z \in \{\mathcal{F}_0(X), \mathcal{F}(X), \mathcal{F}(\mathcal{F}(X))\}$ . A function  $f : \mathcal{T}(X) \rightarrow Z$  is a *tournament function* if

- (i)  $f(T) = f(T')$  for all tournaments  $T = (A, \succ)$  and  $T' = (A, \succ')$  such that  $\succ|_A = \succ'|_A$ , and
- (ii)  $f((\pi(A), \succ')) = \pi(f((A, \succ)))$  for all tournaments  $(A, \succ)$ ,  $(A', \succ')$ , and tournament isomorphisms<sup>2</sup>  $\pi : A \rightarrow A'$  of  $(A, \succ)$  and  $(A', \succ')$ .

For a given set  $B \in \mathcal{F}(X)$  and tournament function  $f$ , we overload  $f$  by also writing  $f(B)$ , provided the dominance relation is known from the context. For two tournament functions  $f$  and  $f'$ , we write  $f' \subseteq f$  if  $f'(T) \subseteq f(T)$  for all tournaments  $T$ .

### 2.4 Tournament Solutions

The first tournament function we consider is  $\max_{\prec} : \mathcal{T}(X) \rightarrow \mathcal{F}_0(X)$ , which returns the undominated alternatives of a tournament. Formally,

$$\max_{\prec}((A, \succ)) = \{a \in A \mid \overline{D}_{\succ}(a) = \emptyset\}.$$

Due to the asymmetry of the dominance relation, this function returns at most one alternative in any tournament. Moreover, maximal—i.e., undominated—and maximum—i.e., dominant—elements coincide. In social choice theory, the maximum of a majority tournament is commonly referred to as the *Condorcet winner*. Obviously, the dominance relation may contain cycles and thus fail to have a maximal element. For this reason, a variety of alternative concepts to single out the “best” alternatives of a tournament have been suggested. Formally, a *tournament solution*  $S$  is defined as a function that associates with each tournament  $T = (A, \succ)$  a non-empty subset  $S(T)$  of  $A$ .

**Definition 2.** A *tournament solution*  $S$  is a tournament function  $S : \mathcal{T}(X) \rightarrow \mathcal{F}(X)$  such that  $\max_{\prec}(T) \subseteq S(T) \subseteq A$  for all tournaments  $T = (A, \succ)$ .<sup>3</sup>

The set  $S(T)$  returned by a tournament solution for a given tournament  $T$  is called the *choice set* of  $T$  whereas  $A \setminus S(T)$  consists of the *unchosen alternatives*. Since tournament solutions always yield non-empty choice sets, they have to select all alternatives in homogeneous tournaments. If  $S' \subseteq S$  for two tournament solutions  $S$  and  $S'$ , we say that  $S'$  is a *refinement* of  $S$  or that  $S'$  is *finer* than  $S$ .

<sup>2</sup> $\pi(A)$  is a shorthand for the set  $\{\pi(a) \mid a \in A\}$ .

<sup>3</sup>Laslier (1997) is slightly more stringent here as he requires the maximum be the only element in  $S(T)$  whenever it exists.

## 2.5 Properties of Tournament Solutions

The literature on tournament solutions has identified a number of desirable properties for tournament solutions. In this section, we will define six of the most common properties.<sup>4</sup> In a more general context, Moulin (1988) distinguishes between *monotonicity* and *independence* conditions, where a monotonicity condition describes the positive association of the solution with some parameter and an independence condition characterizes the invariance of the solution under the modification of some parameter.

In the context of tournament solutions, we will further distinguish between properties that are defined in terms of the dominance relation and properties defined in terms of the set inclusion relation. With respect to the former, we consider *monotonicity* and *independence of unchosen alternatives*. A tournament solution is monotonic if a chosen alternative remains in the choice set when extending its dominion and leaving everything else unchanged.

**Definition 3.** A tournament solution  $S$  satisfies *monotonicity* (MON) if  $a \in S(T)$  implies  $a \in S(T')$  for all tournaments  $T = (A, \succ)$ ,  $T' = (A, \succ')$ , and  $a \in A$  such that  $\succ|_{A \setminus \{a\}} = \succ'|_{A \setminus \{a\}}$  and  $D_\succ(a) \subseteq D_{\succ'}(a)$ .

A solution is independent of unchosen alternatives if the choice set is invariant under any modification of the dominance relation between unchosen alternatives.

**Definition 4.** A tournament solution  $S$  is *independent of unchosen alternatives* (IUA) if  $S(T) = S(T')$  for all tournaments  $T = (A, \succ)$  and  $T' = (A, \succ')$  such that  $D_\succ(a) = D_{\succ'}(a)$  for all  $a \in S(T)$ .

With respect to set inclusion, we consider a monotonicity property to be called the *weak superset property* and an independence property known as the *strong superset property*. A tournament solution satisfies the weak superset property if an unchosen alternative remains unchosen when other unchosen alternatives are removed.

**Definition 5.** A tournament solution  $S$  satisfies the *weak superset property* (WSP) if  $S(B) \subseteq S(A)$  for all tournaments  $T = (A, \succ)$  and  $B \subseteq A$  such that  $S(A) \subseteq B$ .

The strong superset property states that a choice set is invariant under the removal of unchosen alternatives.

**Definition 6.** A tournament solution  $S$  satisfies the *strong superset property* (SSP) if  $S(B) = S(A)$  for all tournaments  $T = (A, \succ)$  and  $B \subseteq A$  such that  $S(A) \subseteq B$ .

The difference between WSP and SSP is precisely another independence condition called *idempotency*. A solution is *idempotent* if the choice set is invariant under repeated application of the solution concept, i.e.,  $S(S(T)) = S(T)$  for all tournaments  $T$ . When  $S$  is not idempotent, we define  $S^k(T) = S(S^{k-1}(T))$  inductively by letting  $S^1(T) = S(T)$  and  $S^\infty(T) = \bigcap_{k \in \mathbb{N}} S^k(T)$ .

---

<sup>4</sup>Our terminology slightly differs from the one by Laslier (1997) and others. *Independence of unchosen alternatives* is also called *independence of the losers* or *independence of non-winners*. The *weak superset property* has been referred to as  $\epsilon^+$  or the *Aizerman property*.

The four properties defined above (MON, IUA, WSP, and SSP) will be called *basic* properties of tournament solutions. The conjunction of MON and SSP implies IUA. It is therefore sufficient to show MON and SSP in order to prove that a tournament solution satisfies all four basic properties.

Two further properties considered in this paper are *composition-consistency* and *irregularity*. A tournament solution is composition-consistent if it chooses the “best” alternatives from the “best” components.

**Definition 7.** A tournament solution  $S$  is *composition-consistent* (COM) if for all tournaments  $T, T_1, \dots, T_k$ , and  $\tilde{T}$  such that  $T = \Pi(\tilde{T}, T_1, \dots, T_k)$ ,  $S(T) = \bigcup_{i \in S(\tilde{T})} S(T_i)$ .

Finally, a tournament solution is irregular if it is capable of excluding alternatives in regular tournaments.

**Definition 8.** A tournament solution  $S$  satisfies *irregularity* (IRR) if there exists a regular tournament  $T = (A, \succ)$  such that  $S(T) \neq A$ .

### 3 Qualified Subsets

In this section, we will define a class of tournament solutions that is based on identifying significant subtournaments of the original tournament, such as subtournaments that admit a maximal alternative.

#### 3.1 Concepts of Qualified Subsets

A *concept of qualified subsets* is a tournament function that, for a given tournament  $T = (A, \succ)$ , returns subsets of  $A$  that satisfy certain properties. Each such set of sets will be referred to as a *family of qualified subsets*. Two natural examples of concepts of qualified subsets are  $\mathcal{M}$ , which yields all subsets that admit a maximal element, and  $\mathcal{M}^*$ , which yields all non-empty transitive subsets. Formally,

$$\begin{aligned}\mathcal{M}((A, \succ)) &= \{B \subseteq A \mid \max_{\prec}(B) \neq \emptyset\} \text{ and} \\ \mathcal{M}^*((A, \succ)) &= \{B \subseteq A \mid \max_{\prec}(C) \neq \emptyset \text{ for all non-empty } C \subseteq B\}.\end{aligned}$$

$\mathcal{M}$  and  $\mathcal{M}^*$  are examples of concepts of qualified subsets, which are formally defined as follows.

**Definition 9.** Let  $\mathcal{Q} : \mathcal{T}(X) \rightarrow \mathcal{F}(\mathcal{F}(X))$  be a tournament function such that  $\mathcal{M}_1(T) \subseteq \mathcal{Q}(T) \subseteq \mathcal{M}(T)$ .  $\mathcal{Q}$  is a *concept of qualified subsets* if it meets the following three conditions for every tournament  $T = (A, \succ)$ .

(Closure)  $\mathcal{Q}(T)$  is downward closed with respect to  $\mathcal{M}$ : Let  $Q \in \mathcal{Q}(T)$ . Then,  $Q' \in \mathcal{Q}(T)$  if  $Q' \subseteq Q$  and  $Q' \in \mathcal{M}(T)$ .

(Independence) Qualified sets are independent of outside alternatives: Let  $A' \in \mathcal{F}(X)$  and  $Q \subseteq A \cap A'$ . Then,  $Q \in \mathcal{Q}(A)$  if and only if  $Q \in \mathcal{Q}(A')$ .

(Fusion) Qualified sets may be merged under certain conditions: Let  $Q_1, Q_2 \in \mathcal{Q}(T)$  and  $Q_1 \setminus Q_2 \succ Q_2$ . Then  $Q_1 \cup Q_2 \in \mathcal{Q}(T)$  if there is a tournament  $T' \in \mathcal{T}(X)$  and  $Q \in \mathcal{Q}(T')$  such that  $|Q_1 \cup Q_2| \leq |Q|$ .

Whether a set is qualified only depends on its internal structure (due to independence and the isomorphism condition of Definition 1). While closure, independence, and the fact that all singletons are qualified are fairly natural, the fusion condition is slightly more technical. Essentially, it states that if a qualified subset dominates another qualified subset, then the union of these subsets is also qualified. The additional cardinality restriction is only required to enable *bounded* qualified subsets. For every concept of qualified subsets  $\mathcal{Q}$  and every given  $k \in \mathbb{N}$ ,  $\mathcal{Q}_k : \mathcal{T}(X) \rightarrow \mathcal{F}(\mathcal{F}(X))$  is a tournament function such that  $\mathcal{Q}_k(T) = \{B \in \mathcal{Q}(T) \mid |B| \leq k\}$ . It is easily verified that  $\mathcal{Q}_k$  is a concept of qualified subsets. Furthermore,  $\mathcal{M}$  and  $\mathcal{M}^*$  (and thus also  $\mathcal{M}_k$  and  $\mathcal{M}_k^*$ ) are concepts of qualified subsets. Since only tournaments of order 4 or more may be intransitive and admit a maximal element at the same time,  $\mathcal{M}_k = \mathcal{M}_k^*$  for  $k \in \{1, 2, 3\}$ .

### 3.2 Maximal Elements of Maximal Qualified Subsets

For every concept of qualified subsets, we can now define a tournament solution that yields the maximal elements of all inclusion-maximal qualified subsets, i.e., all qualified subsets that are not contained in another qualified subset.

**Definition 10.** Let  $\mathcal{Q}$  be a concept of qualified subsets. Then, the tournament solution  $S_{\mathcal{Q}}$  is defined as

$$S_{\mathcal{Q}}(T) = \{\max_{\subseteq}(B) \mid B \in \max_{\subseteq}(\mathcal{Q}(T))\}.$$

Since any family of qualified subsets contains all singletons,  $S_{\mathcal{Q}}(T)$  is guaranteed to be non-empty and contains the Condorcet winner whenever one exists. As a consequence,  $S_{\mathcal{Q}}$  is well-defined as a tournament solution.

The following tournament solutions can be restated via appropriate concepts of qualified subsets.

**Condorcet non-losers.**  $S_{\mathcal{M}_2}$  is arguably the largest non-trivial tournament solution. In tournaments of order two or more, it chooses every alternative that dominates at least one other alternative. We will refer to this concept as *Condorcet non-losers (CNL)* as it selects everything except the minimum (or *Condorcet loser*) in such tournaments.

**Uncovered set.**  $S_{\mathcal{M}}(T)$  returns the *uncovered set*  $UC(T)$  of a tournament  $T$ , i.e., the set consisting of the maximal elements of inclusion-maximal subsets that admit a maximal element. The uncovered set is usually defined in terms of a subrelation of the dominance relation called the *covering relation* (Fishburn, 1977; Miller, 1980).

**Banks set.**  $S_{\mathcal{M}^*}(T)$  yields the *Banks set*  $BA(T)$  of a tournament  $T$  (Banks, 1985).  $\mathcal{M}^*(T)$  contains subsets that not only admit a maximum, but can be completely ordered from maximum to minimum such that all of their non-empty subsets admit a maximum.  $S_{\mathcal{M}^*}(T)$  thus returns the maximal elements of inclusion-maximal *transitive* subsets.

In the remainder of this section, we will prove various statements about tournament solutions defined via qualified subsets. For a set  $B$  and an alternative  $a \notin B$ , the short notation  $[B, a]$  will be used to denote the set  $B \cup \{a\}$  and the fact that  $\max_{\prec}(B \cup \{a\}) = \{a\}$ . In several proofs, we will make use of the fact that whenever  $a \notin S_Q(T)$ , there is some  $b \in S_Q(T)$  for every qualified subset  $[Q, a]$  such that  $[Q \cup \{a\}, b] \in \mathcal{Q}(T)$ . We start by showing that every tournament solution defined via qualified subsets satisfies the weak superset property and monotonicity.

**Proposition 1.** *Let  $\mathcal{Q}$  be a concept of qualified subsets. Then,  $S_Q$  satisfies WSP and MON.*

*Proof.* Let  $T = (A, \succ)$  be a tournament,  $a \notin S_Q(A)$ , and  $A' \subseteq A$  such that  $S_Q(T) \cup \{a\} \subseteq A'$ . For WSP, we need to show that  $a \notin S_Q(A')$ . Let  $[Q, a] \in \mathcal{Q}(A')$ . Due to independence,  $[Q, a] \in \mathcal{Q}(A)$ . Since  $a \notin S_Q(A)$ , there has to be some  $b \in S_Q(A)$  such that  $[Q \cup \{a\}, b] \in \mathcal{Q}(A)$ . Again, independence implies that  $[Q \cup \{a\}, b] \in \mathcal{Q}(A')$ . Hence,  $a \notin S_Q(A')$ .

For MON, observe that  $a \in S_Q$  implies that there exists  $[Q, a] \in \max_{\subseteq}(\mathcal{Q}(T))$ . Define  $T' = (A, \succ')$  by letting  $T'|_{A \setminus \{a\}} = T|_{A \setminus \{a\}}$  and  $a \succ' b$  for some  $b \in A$  with  $b \succ a$ . Clearly,  $[Q, a]$  is contained in  $\mathcal{Q}(T')$  due to independence and the fact that  $b \notin Q$ . Now, assume for contradiction that there is some  $c \in A$  such that  $[Q \cup \{a\}, c] \in \mathcal{Q}(T')$ . Since  $a \succ' b$ ,  $c \neq b$ . Independence then implies that  $[Q \cup \{a\}, c] \in \mathcal{Q}(T)$ , a contradiction.  $\square$

Proposition 1 implies several known statements such as that *CNL*, *UC*, and *BA* satisfy MON and WSP. All three concepts are known to fail idempotency (and thus *SSP*). *CNL* trivially satisfies *IUA* whereas this is not the case for *UC* and *BA* (see Laslier, 1997). We also obtain some straightforward inclusion relationships, which define an infinite hierarchy of tournament solutions ranging from *CNL* to *BA*.

**Proposition 2.**  $S_{\mathcal{M}^*} \subseteq S_{\mathcal{M}}$ ,  $S_{\mathcal{M}_k^*} \subseteq S_{\mathcal{M}_k}$ ,  $S_{\mathcal{Q}_{k+1}} \subseteq S_{\mathcal{Q}_k}$ , and  $S_Q \subseteq S_{\mathcal{Q}_k}$  for every concept of qualified subsets  $\mathcal{Q}$  and  $k \in \mathbb{N}$ .

*Proof.* All inclusion relationships follow from the following observation. Let  $T$  be a tournament and  $\mathcal{Q}$  and  $\mathcal{Q}'$  concepts of qualified subsets such that for every  $[Q, a] \in \max_{\subseteq}(\mathcal{Q}(T))$ , there is  $[Q', a] \in \max_{\subseteq}(\mathcal{Q}'(T))$ . Then,  $S_Q \subseteq S_{Q'}$ .  $\square$

It turns out that the Banks set is the finest tournament solution definable via qualified subsets. In order to show this, we introduce a new property called *strong retentiveness* that prescribes that the choice set of every dominator set is contained in the original choice set. Alternatively, it can be seen as a variant of WSP because it states that a choice set may not grow when an alternative and its entire dominion are removed from the tournament.

**Definition 11.** A tournament solution  $S$  satisfies *strong retentiveness* if  $S(\overline{D}(a)) \subseteq S(A)$  for all tournaments  $T = (A, \succ)$  and  $a \in A$ .

**Lemma 1.** Let  $\mathcal{Q}$  be a concept of qualified subsets. Then,  $S_{\mathcal{Q}}$  satisfies strong retentiveness.

*Proof.* Let  $(A, \succ)$  be a tournament,  $a \in A$  an alternative, and  $B = \overline{D}(a)$ . We show that  $b \in S_{\mathcal{Q}}(B)$  implies that  $b \in S_{\mathcal{Q}}(A)$ . Let  $[Q, b]$  be a maximal qualified subset in  $B$ , i.e.,  $[Q, b] \in \max_{\subseteq}(\mathcal{Q}(B))$ . If  $[Q, b] \in \max_{\subseteq}(\mathcal{Q}(A))$ , we are done. Otherwise, there has to be some  $c \in A$  such that  $[Q \cup \{b\}, c] \in \mathcal{Q}(A)$ . Furthermore,  $[Q, b] \succ a$  and  $a \succ c$  because otherwise  $[Q \cup \{b\}, c]$  would be qualified in  $B$  as well. We can now merge the qualified subsets  $[Q, b]$  and  $[\{a\}, b]$  according to the fusion condition. We claim that  $[Q \cup \{a\}, b] \in \max_{\subseteq}(\mathcal{Q}(A))$ . Assume for contradiction that there is some  $d \in A$  such that  $[Q \cup \{a, b\}, d] \in \mathcal{Q}(A)$ . Since  $d \in \overline{D}(a)$ , independence implies that  $[Q \cup \{a, b\}, d] \in \mathcal{Q}(B)$ . This is a contradiction because  $[Q, b]$  was assumed to be a *maximal* qualified subset of  $B$ .  $\square$

**Theorem 1.** The Banks set is the finest tournament solution satisfying strong retentiveness and thus the finest tournament solution definable via qualified subsets.

*Proof.* Let  $S$  be a tournament solution that satisfies strong retentiveness and  $T = (A, \succ)$  a tournament. We first show that  $BA(A) \subseteq S(A)$ . For every  $a \in BA(A)$ , there has to be maximal transitive set  $[Q, a] \subseteq A$ . Let  $Q = \{q_1, q_2, \dots, q_n\}$ . We show that  $B = \bigcap_{i=1}^n \overline{D}(q_i) = \{a\}$ . Since  $a \succ Q$ ,  $a \in B$ . Assume for contradiction that  $b \in B$  with  $b \neq a$ . Then  $b \succ Q$  and either  $[Q \cup \{b\}, a]$  or  $[Q \cup \{a\}, b]$  is a transitive set, which contradicts the maximality of  $[Q, a]$ . Since  $S(B) = S(\{a\}) = \{a\}$ , the repeated application of strong retentiveness implies that  $a \in S(A)$ . The statement now follows from Lemma 1.  $\square$

## 4 Stable Sets

In this section, we propose a general method for refining any suitable solution concept  $S$  by formalizing the stability of sets of alternatives with respect to  $S$ . This method is based on the notion of stable sets (von Neumann and Morgenstern, 1944) and generalizes covering sets as introduced by Dutta (1988).

### 4.1 Stability and Directedness

The reason why we are interested in maximal—i.e., undominated—alternatives is that dominated alternatives can be upset by other alternatives; they are unstable. The rationale behind stable sets is that this instability is only meaningful if an alternative is upset by something which itself is stable. Hence, a set of alternatives  $B$  is said to be stable if it consists precisely of those alternatives not upset by  $B$ . In von Neumann and Morgenstern's original definition,  $a$  is upset by  $B$  if some element of  $B$  dominates  $a$ .

In our generalization,  $a$  is upset by  $B$  if  $a \notin S(B \cup \{a\})$  for some underlying solution concept  $S$ .<sup>5</sup>

As an alternative to this fixed-point definition, which will be formalized in Corollary 1, stable sets can be seen as sets that comply with internal and external stability in some well-defined way. First, there should be no reason to restrict the selection by excluding some alternative from it and, secondly, there should be an argument against each proposal to include an outside alternative into the selection.<sup>6</sup> In our context, external stability with respect to some tournament solution  $S$  is defined as follows.

**Definition 12.** Let  $S$  be a tournament solution and  $T = (A, \succ)$  a tournament. Then,  $B \subseteq A$  is *externally stable* in  $T$  with respect to tournament solution  $S$  (or  *$S$ -stable*) if  $a \notin S(B \cup \{a\})$  for all  $a \in A \setminus B$ . The set of  $S$ -stable sets for a given tournament  $T = (A, \succ)$  will be denoted by  $\mathcal{S}_S(T) = \{B \subseteq A \mid B \text{ is } S\text{-stable in } T\}$ .

Externally stable sets are guaranteed to exist since the set of all alternatives  $A$  is trivially  $S$ -stable in  $(A, \succ)$  for every  $S$ . We say that a set  $B \subseteq A$  is *internally stable* with respect to  $S$  if  $S(B) = B$ . We will focus on external stability for now because we will see later that certain conditions imply the existence of a unique minimal externally stable set, which also satisfies internal stability. We define  $\hat{S}(T)$  to be the tournament solution that returns the union of all *inclusion-minimal*  $S$ -stable sets in  $T$ , i.e., the union of all  $S$ -stable sets that do not contain an  $S$ -stable set as a proper subset.

**Definition 13.** Let  $S$  be a tournament solution. Then, the tournament solution  $\hat{S}$  is defined as

$$\hat{S}(T) = \bigcup_{\subseteq} \min(\mathcal{S}_S(T)).$$

It is easily verified that  $\hat{S}$  is well-defined as a tournament solution as there are no  $S$ -stable sets that do not contain the Condorcet winner whenever one exists. We will only be concerned with tournament solutions  $S$  that (presumably) admit a *unique* minimal  $S$ -stable set in any tournament. It turns out it is precisely this property that is most difficult to prove for all but the simplest tournament solutions. A tournament  $T$  contains a *unique* minimal  $S$ -stable set if and only if  $\mathcal{S}_S(T)$  is a *directed* set with respect to set inclusion, i.e., for all sets  $B, C \in \mathcal{S}_S(T)$  there is a set  $D \in \mathcal{S}_S(T)$  contained in both  $B$  and  $C$ . We say that  $\mathcal{S}_S$  is directed when  $\mathcal{S}_S(T)$  is a directed set for all tournaments  $T$ . Throughout this paper, directedness of a set of sets  $\mathcal{S}$  is shown by proving the stronger property of closure under intersection, i.e.,  $B \cap C \in \mathcal{S}$  for all  $B, C \in \mathcal{S}$ . A set of sets  $\mathcal{S}$  *pairwise intersects* if  $B \cap C \neq \emptyset$  for all  $B, C \in \mathcal{S}$ . We will prove that, for every concept of qualified subsets  $\mathcal{Q}$ ,  $\mathcal{S}_{S_{\mathcal{Q}}}$  is closed under intersection if and only if  $\mathcal{S}_{S_{\mathcal{Q}}}$  pairwise intersects. In order to improve readability, we will use the short notation  $\mathcal{S}_{\mathcal{Q}}$  for  $\mathcal{S}_{S_{\mathcal{Q}}}$ .

<sup>5</sup>Von Neumann and Morgenstern's definition can be seen as the special case where  $a$  is upset by  $B$  if  $a \notin \max_{\prec}(B \cup \{a\})$ .

<sup>6</sup>A large number of solution concepts in the social sciences spring from similar notions of internal and/or external stability (see, e.g., von Neumann and Morgenstern, 1944; Nash, 1951; Shapley, 1964; Schwartz, 1986; Dutta, 1988; Basu and Weibull, 1991; Duggan and Le Breton, 1996). Wilson (1970) refers to stability as the *solution property*.

**Lemma 2.** *Let  $\mathcal{Q}$  be a concept of qualified subsets. Then,  $\mathcal{S}_{\mathcal{Q}}$  is closed under intersection if and only if  $\mathcal{S}_{\mathcal{Q}}$  pairwise intersects.*

*Proof.* The direction from left to right is straightforward since the empty set is not stable. The opposite direction is shown by contraposition, i.e., we prove that  $\mathcal{S}_{\mathcal{Q}}$  does not pairwise intersect if  $\mathcal{S}_{\mathcal{Q}}$  is not closed under intersection. Let  $T = (A, \succ)$  be a tournament and  $B_1, B_2 \in \mathcal{S}_{\mathcal{Q}}(T)$  be two sets such that  $C = B_1 \cap B_2 \notin \mathcal{S}_{\mathcal{Q}}(T)$ . Since  $C$  is not  $\mathcal{S}_{\mathcal{Q}}$ -stable, there has to be  $a \in A \setminus C$  such that  $a \in \mathcal{S}_{\mathcal{Q}}(C \cup \{a\})$ . In other words, there has to be a set  $Q \subseteq C$  such that  $[Q, a] \in \max_{\subseteq}(\mathcal{Q}(C \cup \{a\}))$ . Define

$$B'_1 = \{b \in B_1 \mid b \succ Q\} \text{ and } B'_2 = \{b \in B_2 \mid b \succ Q\}.$$

Clearly,  $(B'_1 \setminus B'_2) \cap C = \emptyset$  and  $(B'_2 \setminus B'_1) \cap C = \emptyset$ . Assume without loss of generality that  $a \notin B_1$ . It follows from the stability of  $B_1$ , that  $B_1$  has to contain an alternative  $b_1$  such that  $b_1 \succ [Q, a]$ . Hence,  $B'_1$  is not empty. Next, we show that  $B'_1 \cap B'_2 = \emptyset$ . Assume for contradiction that there is some  $b \in B'_1 \cap B'_2$ . If  $b \succ a$ , independence implies that  $[Q \cup \{a\}, b] \in \mathcal{Q}(C \cup \{a\})$ , which contradicts the fact that  $[Q, a]$  is a maximal qualified subset in  $C \cup \{a\}$ . If, on the other hand,  $a \succ b$ , the set  $[Q \cup \{b\}, a]$  is isomorphic to  $[Q \cup \{a\}, b_1]$ , which is a qualified subset of  $B_1 \cup \{a\}$ . Thus,  $[Q \cup \{b\}, a] \in \mathcal{Q}(C \cup \{a\})$ , again contradicting the maximality of  $[Q, a]$ . Independence, the isomorphism of  $[Q, a]$  and  $[Q, b_1]$ , and the stability of  $B_2$  further require that there has to be an alternative  $b_2 \in B_2$  such that  $b_2 \succ [Q, b_1]$ . Hence,  $B'_1$  and  $B'_2$  are disjoint and non-empty.

Let  $a' \in B'_2$  and  $R$  be a maximal subset of  $B'_1 \cup Q$  such that  $[R, a'] \in \mathcal{Q}(B'_1 \cup Q \cup \{a'\})$ . We claim that  $Q$  has to be contained in  $R$ . Assume for contradiction that there exists some  $b \in Q \setminus R$ . Clearly,  $[Q, a]$  and  $[Q, a']$  are isomorphic. It therefore follows from independence that  $[Q, a'] \in \mathcal{Q}(B'_1 \cup Q \cup \{a'\})$  and from closure that  $[(Q \cap R) \cup \{b\}, a'] \in \mathcal{Q}(B'_1 \cup Q \cup \{a'\})$ . Due to the stability of  $B_1$ ,  $[R, a']$  is not a maximal qualified subset in  $B_1 \cup \{a'\}$ , i.e., there exists a qualified subset that contains more elements. We may thus merge the qualified subsets  $[R, a']$  and  $[(Q \cap R) \cup \{b\}, a']$  according to the fusion condition because  $R \setminus Q \succ Q$  and consequently  $R \setminus Q \succ (Q \cap R) \cup \{b\}$ . We then have that  $[R \cup \{b\}, a'] \in \mathcal{Q}(B'_1 \cup Q \cup \{a'\})$ , which yields a contradiction because  $R$  was assumed to be a maximal set such that  $[R, a'] \in \mathcal{Q}(B'_1 \cup Q \cup \{a'\})$ . Hence,  $Q \subseteq R$ . Due to the stability of  $B_1$  in  $T$ , there has to be a  $c \in B_1$  such that  $c \succ [R, a']$ . Since  $B'_1$  contains all alternatives in  $B_1$  that dominate  $Q \subseteq R$ , it also contains  $c$ . Independence then implies that  $[R, a'] \notin \max_{\subseteq}(\mathcal{Q}_k(B'_1 \cup Q \cup \{a'\}))$ .

Thus,  $B'_1 \cup Q$  is stable in  $B'_1 \cup B'_2 \cup Q$ . Since  $Q$  is contained in every maximal set  $R \subseteq B'_1 \cup Q$  such that  $[R, a'] \in \mathcal{Q}(B'_1 \cup Q \cup \{a'\})$  for some  $a' \in B'_2$ ,  $B'_1$  (and by an analogous argument  $B'_2$ ) remains stable when removing  $Q$ . This completes the proof because  $B'_1$  and  $B'_2$  are two disjoint  $\mathcal{S}_{\mathcal{Q}}$ -stable sets in  $B'_1 \cup B'_2$ .  $\square$

Dutta has shown by induction on the tournament order that tournaments admit no disjoint  $\mathcal{S}_{\mathcal{M}}$ -stable sets (so-called *covering sets*).

**Theorem 2** (Dutta, 1988).  $\mathcal{S}_{\mathcal{M}}$  pairwise intersects.

Dutta (1988) also showed that covering sets are closed under intersection, which now also follows from Lemma 2.<sup>7</sup>

Naturally, finer solution concepts also yield smaller minimal stable sets (if their uniqueness is guaranteed).

**Proposition 3.** *Let  $S$  and  $S'$  be two tournament solutions such that  $\mathcal{S}_{S'}$  is directed and  $S' \subseteq S$ . Then,  $\widehat{S}' \subseteq \widehat{S}$  and  $\mathcal{S}_S$  pairwise intersects.*

*Proof.* The statements follow from the simple fact that every  $S$ -stable set is also  $S'$ -stable. Let  $B \subseteq A$  be a minimal  $S$ -stable set in tournament  $(A, \succ)$ . Then,  $a \notin S(B \cup \{a\})$  for every  $a \in A \setminus B$  and, due to the inclusion relationship,  $a \notin S'(B \cup \{a\}) \subseteq S(B \cup \{a\})$ . As a consequence,  $B$  is  $S'$ -stable and has to contain the unique minimal  $S'$ -stable set since  $\mathcal{S}_{S'}$  is directed.  $\mathcal{S}_S$  pairwise intersects because two disjoint  $S$ -stable sets would also be  $S'$ -stable, which contradicts the directedness of  $\mathcal{S}_{S'}$ .  $\square$

As a corollary of the previous statements, the set of  $S_{\mathcal{M}_k}$ -stable sets for every  $k$  is closed under intersection.

**Theorem 3.**  *$\mathcal{S}_{\mathcal{M}_k}$  is closed under intersection for all  $k \in \mathbb{N}$ .*

*Proof.* Let  $k \in \mathbb{N}$ . We know from Proposition 2 that  $S_{\mathcal{M}} \subseteq S_{\mathcal{M}_k}$  and from Theorem 2 and Lemma 2 that  $\mathcal{S}_{\mathcal{M}}$  is directed. Proposition 3 implies that  $\mathcal{S}_{\mathcal{M}_k}$  pairwise intersects. The statement then straightforwardly follows from Lemma 2.  $\square$

Interestingly,  $\mathcal{S}_{\mathcal{M}_2}$ , the set of all dominant sets, is not only closed under intersection, but in fact totally ordered with respect to set inclusion.

We conjecture that the set of all  $S_{\mathcal{M}^*}$ -stable sets also pairwise intersects and thus admits a unique minimal element. However, the combinatorial structure of transitive subtournaments within tournaments is extraordinarily rich (see, e.g., Woeginger, 2003; Gaspers and Mnich, 2010) and it seems that a proof of the conjecture would be significantly more difficult than Dutta's. As will be shown in Section 5, the conjecture that  $S_{\mathcal{M}^*}$ -stable sets pairwise intersect is a weakened version of a conjecture by Schwartz (1990).

**Conjecture 1.**  *$\mathcal{S}_{\mathcal{M}^*}$  is closed under intersection.*

Using Lemma 2, the conjecture entails that  $\mathcal{S}_{\mathcal{M}_k^*}$  for all  $k \in \mathbb{N}$  is also closed under intersection. Since tournaments with less than four alternatives may not contain a maximal element and a cycle at the same time, this trivially holds for  $k \leq 3$ . The weakest version of Conjecture 1 that is not implied by Theorem 2 is that  $\mathcal{S}_{\mathcal{M}_4^*}$  is closed under intersection. We were able to show this by reducing it to a large, but finite, number of cases that were checked using a computer. Unfortunately, this exercise did not yield any insight on how to prove Conjecture 1.

Two well-known examples of minimal stable sets are the top cycle of a tournament, which is the minimal stable set with respect to  $S_{\mathcal{M}_2}$ , and the minimal covering set, which is the minimal stable set with respect to  $S_{\mathcal{M}}$ .

---

<sup>7</sup>As Dutta's definition requires a stable set to be internally and externally stable, he actually proves that the intersection of any pair of coverings sets *contains* a covering set. A simpler proof, which shows that *externally*  $S_{\mathcal{M}}$ -stable sets are closed under intersection, is given by Laslier (1997).

**Minimal dominant set.** The *minimal dominant set* (or top cycle) of a tournament  $T = (A, \succ)$  is given by  $TC(T) = \widehat{S}_{\mathcal{M}_2}(T) = \widehat{CNL}$ , i.e., it is the smallest set  $B$  such that  $B \succ A \setminus B$  (Good, 1971; Smith, 1973).

**Minimal covering set.** The *minimal covering set* of a tournament  $T$  is given by  $MC(T) = \widehat{S}_{\mathcal{M}}(T) = \widehat{UC}$ , i.e., it is the smallest set  $B$  such that for all  $a \in A \setminus B$ , there exists  $b \in B$  so that every alternative in  $B$  that is dominated by  $a$  is also dominated by  $b$  (Dutta, 1988).

The proposed methodology also suggests the definition of a new tournament solution that has not been considered before in the literature.

**Minimal extending set.** The *minimal extending set* of a tournament  $T$  is given by  $ME(T) = \widehat{S}_{\mathcal{M}^*}(T) = \widehat{BA}$ , i.e., it is the smallest set  $B$  such that no  $a \in A \setminus B$  is the maximal element of a maximal transitive subset in  $B \cup \{a\}$ .

The minimal extending set will be further analyzed in Section 4.3.

## 4.2 Properties of Minimal Stable Sets

If  $\mathcal{S}_S$  is directed—and we will only be concerned with tournament solutions  $S$  for which this is (presumably) the case— $\widehat{S}$  satisfies a number of desirable properties.

**Proposition 4.** *Let  $S$  be a tournament solution such that  $\mathcal{S}_S$  is directed. Then,  $\widehat{S}$  satisfies WSP and IUA.*

*Proof.* Clearly, any minimal  $S$ -stable set  $B$  remains  $S$ -stable when losing alternatives are removed or when edges between losing alternatives are modified. In the latter case,  $B$  also remains minimal. In the former case, the minimal  $S$ -stable set is contained in  $B$ .  $\square$

It can be shown that sets that are stable within a stable set are also stable in the original tournament when the underlying tournament solution is defined via a concept of qualified subsets  $\mathcal{Q}$ . This lemma will prove very useful when analyzing  $\widehat{S}_{\mathcal{Q}}$ .

**Lemma 3.** *Let  $T = (A, \succ)$  be a tournament and  $\mathcal{Q}$  a concept of qualified subsets. Then,  $\mathcal{S}_{\mathcal{Q}}(B) \subseteq \mathcal{S}_{\mathcal{Q}}(A)$  for all  $B \in \mathcal{S}_{\mathcal{Q}}(A)$ .*

*Proof.* We prove the statement by showing that the following implication holds for all  $B \subseteq A$ ,  $C \in \mathcal{S}_{\mathcal{Q}}(B)$ , and  $a \in A$ :

$$\text{if } a \notin \mathcal{S}_{\mathcal{Q}}(B \cup \{a\}) \text{ then } a \notin \mathcal{S}_{\mathcal{Q}}(C \cup \{a\}).$$

To see this, let  $a \notin \mathcal{S}_{\mathcal{Q}}(B \cup \{a\})$  and assume for contradiction that there exist  $[Q, a] \in \max_{\subseteq} \mathcal{Q}(C \cup \{a\})$ . Then there has to be  $b \in B$  such that  $[Q \cup \{a\}, b] \in \mathcal{Q}(B \cup \{a\})$  because  $[Q, a] \notin \max_{\subseteq} \mathcal{Q}(B \cup \{a\})$ . Now, if  $b \in C$ , closure and independence imply that  $[Q \cup \{a\}, b] \in \mathcal{Q}(C \cup \{a\})$ , contradicting the maximality of  $[Q, a]$ . If, on the other hand,

$b \in B \setminus C$ , then there has to be  $c \in C$  such that  $[Q \cup \{b\}, c] \in \mathcal{Q}(C \cup \{b\})$ . No matter whether  $c \succ a$  or  $a \succ c$ ,  $Q \cup \{a, c\}$  is isomorphic to  $[Q \cup \{b\}, c]$  and thus also a qualified subset, which again contradicts the assumption that  $[Q, a]$  was maximal.  $\square$

We are now ready to show a number of appealing properties of *unique* minimal stable sets when the underlying solution concept is defined via qualified subsets.

**Theorem 4.** *Let  $\mathcal{Q}$  be a concept of qualified subsets such that  $\mathcal{S}_{\mathcal{Q}}$  is directed. Then,*

- (i)  $\widehat{\mathcal{S}}_{\mathcal{Q}} \subseteq S_{\mathcal{Q}}^{\infty}$ ,
- (ii)  $S_{\mathcal{Q}}(\widehat{\mathcal{S}}_{\mathcal{Q}}(T) \cup \{a\}) = \widehat{\mathcal{S}}_{\mathcal{Q}}(T)$  for all tournaments  $T = (A, \succ)$  and  $a \in A$  (in particular,  $\widehat{\mathcal{S}}_{\mathcal{Q}}(T)$  is internally stable),
- (iii)  $\widehat{\mathcal{S}}_{\mathcal{Q}}$  satisfies SSP, and
- (iv)  $\widehat{\widehat{\mathcal{S}}}_{\mathcal{Q}} = \widehat{\mathcal{S}}_{\mathcal{Q}}$ .

*Proof.* Let  $T = (A, \succ)$  be a tournament. The first statement of the theorem is shown by proving by induction on  $k$  that  $S_{\mathcal{Q}}^k(T)$  is an  $S_{\mathcal{Q}}$ -stable set. For the basis, let  $B = S_{\mathcal{Q}}(T)$ . Then,  $S_{\mathcal{Q}}(B \cup \{a\}) \subseteq B$  for every  $a \in A \setminus B$  due to WSP of  $S_{\mathcal{Q}}$  (Proposition 1) and thus  $B$  is  $S_{\mathcal{Q}}$ -stable. Now, assume that  $B = S_{\mathcal{Q}}^k(T)$  is  $S_{\mathcal{Q}}$ -stable and let  $C = S_{\mathcal{Q}}(B)$ . Again, WSP implies that  $a \notin S_{\mathcal{Q}}(C \cup \{a\})$  for every  $a \in B \setminus C$ , i.e.,  $C \in \mathcal{S}_{\mathcal{Q}}(B)$ . We can thus directly apply Lemma 3 to obtain that  $C = S_{\mathcal{Q}}^{k+1}(T) \in \mathcal{S}_{\mathcal{Q}}(T)$ . As the minimal  $S_{\mathcal{Q}}$ -stable set is contained in every  $S_{\mathcal{Q}}$ -stable set, the statement follows.

Regarding internal stability, assume for contradiction that  $S_{\mathcal{Q}}(\widehat{\mathcal{S}}_{\mathcal{Q}}(T)) \subset \widehat{\mathcal{S}}_{\mathcal{Q}}(T)$ . However, Lemma 3 implies that  $S_{\mathcal{Q}}(\widehat{\mathcal{S}}_{\mathcal{Q}}(T))$  is  $S_{\mathcal{Q}}$ -stable, contradicting the minimality of  $\widehat{\mathcal{S}}_{\mathcal{Q}}(T)$ . The remainder of the second statement follows straightforwardly from internal stability. If  $S_{\mathcal{Q}}(\widehat{\mathcal{S}}_{\mathcal{Q}}(T) \cup \{a\}) = C \subset \widehat{\mathcal{S}}_{\mathcal{Q}}(T)$  for some  $a \in A \setminus \widehat{\mathcal{S}}_{\mathcal{Q}}(T)$ , WSP implies that  $S_{\mathcal{Q}}(\widehat{\mathcal{S}}_{\mathcal{Q}}(T)) \subseteq C$ , contradicting internal stability.

Regarding SSP, let  $B = \widehat{\mathcal{S}}_{\mathcal{Q}}(T)$  and assume for contradiction that  $C = \widehat{\mathcal{S}}_{\mathcal{Q}}(A') \subset B$  for some  $A'$  with  $B \subseteq A' \subset A$ . Clearly,  $C$  is  $S_{\mathcal{Q}}$ -stable not only in  $A'$  but also in  $B$ , which implies that  $C \in \mathcal{S}_{\mathcal{Q}}(B)$ . According to Lemma 3,  $C$  is also contained in  $\mathcal{S}_{\mathcal{Q}}(A)$ , contradicting the minimality of  $\widehat{\mathcal{S}}_{\mathcal{Q}}(T)$ .

Finally, for  $\widehat{\widehat{\mathcal{S}}}_{\mathcal{Q}}(T) = \widehat{\mathcal{S}}_{\mathcal{Q}}(T)$ , we show that every  $S_{\mathcal{Q}}$ -stable set is  $\widehat{\mathcal{S}}_{\mathcal{Q}}$ -stable and that every *minimal*  $\widehat{\mathcal{S}}_{\mathcal{Q}}$ -stable set is  $S_{\mathcal{Q}}$ -stable. The former follows from  $\widehat{\mathcal{S}}_{\mathcal{Q}}(T) \subseteq S_{\mathcal{Q}}(T)$ , which is a consequence of the first statement of this theorem. For the latter statement, let  $B \in \min_{\subseteq}(\mathcal{S}_{\widehat{\mathcal{S}}_{\mathcal{Q}}}(T))$ . We first show that  $\widehat{\mathcal{S}}_{\mathcal{Q}}(B \cup \{a\}) = B$  for all  $a \in A \setminus B$ . Assume for contradiction that  $\widehat{\mathcal{S}}_{\mathcal{Q}}(B \cup \{a\}) = C \subset B$  for some  $a \in A \setminus B$ . Since  $\widehat{\mathcal{S}}_{\mathcal{Q}}$  satisfies SSP,  $\widehat{\mathcal{S}}_{\mathcal{Q}}(B \cup \{a\}) = C$  for all  $a \in A \setminus B$ . As a consequence,  $C$  is  $\widehat{\mathcal{S}}_{\mathcal{Q}}$ -stable in  $B \cup \{a\}$  for all  $a \in A \setminus B$  and, due to the definition of stability, also in  $A$ . This contradicts the assumption that  $B$  was the *minimal*  $\widehat{\mathcal{S}}$ -stable set. Hence,  $\widehat{\mathcal{S}}_{\mathcal{Q}}(B \cup \{a\}) = B$  for all  $a \in A \setminus B$ . By definition of  $\widehat{\mathcal{S}}_{\mathcal{Q}}$ , this implies that  $a \notin S_{\mathcal{Q}}(B \cup \{a\})$  and thus that  $B$  is  $S_{\mathcal{Q}}$ -stable.  $\square$

The second statement of Theorem 4 allows us to characterize stable sets using the fixed-point formulation mentioned at the beginning of this section, which unifies internal and external stability.

**Corollary 1.** *Let  $\mathcal{Q}$  be a concept of qualified subsets such that  $S_{\mathcal{Q}}$  is directed and  $T = (A, \succ)$  a tournament. Then,*

$$\widehat{S}_{\mathcal{Q}}(T) = \min_{\subseteq} \{B \subseteq A \mid B = \bigcup_{a \in A} S_{\mathcal{Q}}(B \cup \{a\})\}.$$

There may very well be more than one internally and externally  $S_{\mathcal{Q}}$ -stable set in a tournament. For example, the proof of Theorem 4 implies that  $S_{\mathcal{Q}}^{\infty}(T)$  is internally and externally  $S_{\mathcal{Q}}$ -stable.

We have already seen that  $\widehat{S}_{\mathcal{Q}}$  satisfies some of the basic properties defined in Section 2.5. It further turns out that  $\widehat{S}$  inherits monotonicity and composition-consistency from  $S$ .

**Proposition 5.** *Let  $S$  be a tournament solution such that  $S_S$  is directed and  $S$  satisfies MON. Then,  $\widehat{S}$  satisfies MON as well.*

*Proof.* Let  $T = (A, \succ)$  be a tournament with  $a, b \in A$ ,  $a \in \widehat{S}(T)$ , and  $b \succ a$ , and let the relation  $\succ'$  be identical to  $\succ$  except that  $a \succ' b$ . Denote  $T' = (A, \succ')$  and assume for contradiction that  $a \notin \widehat{S}(T')$ . Then, there has to be a minimal  $S$ -stable set  $B \subseteq A \setminus \{a\}$  in  $T'$ . We show that  $B$  is also  $S$ -stable in  $T$ , a contradiction. If  $b \notin B$ , this would clearly be the case because  $\widehat{S}$  satisfies IUA. If, on the other hand,  $b \in B$ , the only reason for  $B$  not to be  $S$ -stable in  $T$  is that  $a \in S((B \cup \{a\}), \succ)$ . However, monotonicity of  $S$  then implies that  $a \in S((B \cup \{a\}), \succ')$ , which is a contradiction because  $B$  is  $S$ -stable in  $T'$ .  $\square$

**Proposition 6.** *Let  $S$  be a tournament solution that satisfies COM. Then,  $\widehat{S}$  satisfies COM as well.*

*Proof.* Let  $S$  be a composition-consistent tournament solution and  $T = (A, \succ) = \Pi(\tilde{T}, T_1, \dots, T_k)$  a product tournament with  $T = (\{1, \dots, k\}, \tilde{\succ})$ ,  $T_1 = (B_1, \succ_1)$ ,  $\dots$ ,  $T_k = (B_k, \succ_k)$ . For a subset  $C$  of  $A$ , let  $C_i = C \cap B_i$  for all  $i \in \{1, \dots, k\}$  and  $\tilde{C} = \bigcup_{i: C_i \neq \emptyset} \{i\}$ . We will prove that  $C \subseteq A$  is  $S$ -stable if and only if

- (i)  $\tilde{C}$  is  $S$ -stable in  $\tilde{T}$ , and
- (ii)  $C_i$  is  $S$ -stable in  $T_i$  for all  $i \in \{1, \dots, k\}$ .

Consider an arbitrary alternative  $a \in A \setminus C$ . For  $C$  to be  $S$ -stable,  $a$  should not be contained in  $S(C \cup \{a\})$ . Since  $S$  is composition-consistent,  $a$  may be excluded for two reasons. First,  $a$  may be contained in an unchosen component, i.e.,  $a \in B_i$  such that  $i \notin S(\tilde{C} \cup \{i\})$ . Secondly,  $a$  may not be selected despite being in a chosen component, i.e.,  $a \in B_i$  such that  $i \in S(\tilde{C} \cup \{i\})$  and  $a \notin S(C_i \cup \{a\})$ . This directly establishes the claim above and consequently that  $\widehat{S}$  is composition-consistent.  $\square$

The previous propositions and theorems allow us to deduce several known statements about  $TC$  and  $MC$ , in particular that both concepts satisfy all basic properties and that  $MC$  is a refinement of  $UC^\infty$  and satisfies **COM**.

We conclude this section by generalizing the axiomatization of the minimal covering set (Dutta, 1988) to abstract minimal stable sets. One of the cornerstones of the axiomatization is  $S$ -exclusivity, which prescribes under which circumstances a single element may be dismissed from the choice set.<sup>8</sup>

**Definition 14.** A tournament solution  $S'$  satisfies  $S$ -exclusivity if, for every tournament  $T = (A, \succ)$ ,  $S'(T) = A \setminus \{a\}$  implies that  $a \notin S(A)$ .

If  $S$  always admits a *unique* minimal  $S$ -stable set and  $\hat{S}$  satisfies **SSP**, which is always the case if  $S$  is defined via qualified subsets, then  $\hat{S}$  can be characterized by **SSP**,  $S$ -exclusivity, and inclusion-minimality.

**Proposition 7.** Let  $S$  be a tournament solution such that  $\mathcal{S}_S$  is directed and  $\hat{S}$  satisfies **SSP**. Then,  $\hat{S}$  is the finest tournament solution satisfying **SSP** and  $S$ -exclusivity.

*Proof.* Let  $S$  be a tournament solution as desired and  $S'$  a tournament solution that satisfies **SSP** and  $S$ -exclusivity. We first prove that  $\hat{S} \subseteq S'$  by showing that  $S'(T)$  is  $S$ -stable for every tournament  $T = (A, \succ)$ . Let  $B = S'(T)$  and  $a \in A \setminus B$ . It follows from **SSP** that  $S'(B \cup \{a\}) = B$  and from  $S$ -exclusivity that  $a \notin S(B \cup \{a\})$ , which implies that  $B$  is  $S$ -stable. Since  $\hat{S}(T)$  is the unique inclusion-minimal  $S$ -stable set, it has to be contained in all  $S$ -stable sets. The statement now follows from the fact that  $\hat{S}$  satisfies **SSP** and  $S$ -exclusivity.  $\square$

Hence,  $TC$  is the finest tournament solution satisfying **SSP** and  $CNL$ -exclusivity,  $MC$  is the finest tournament solution satisfying **SSP** and  $UC$ -exclusivity, and  $ME$  is the finest tournament solution satisfying **SSP** and  $BA$ -exclusivity if Conjecture 1 holds.

### 4.3 The Minimal Extending Set

As mentioned in Section 4.1, the minimal extending set is a new tournament solution that has not been considered before. In analogy to  $UC$ -stable sets, which are known as covering sets, we will call  $BA$ -stable sets *extending sets*.  $B$  is an extending set of tournament  $T = (A, \succ)$  if, for all  $a \notin B$ , every transitive path (or so-called Banks trajectory) in  $B \cup \{a\}$  with maximal element  $a$  can be extended, i.e., there is  $b \in B$  such that  $b$  dominates every element on the path. In other words,  $B \subseteq A$  is an *extending set* if for all  $a \in A \setminus B$ ,  $a \notin BA(B \cup \{a\})$ .

If Conjecture 1 is correct,  $ME$  satisfies all properties defined in Section 2.5 and is a refinement of  $BA$  due to Propositions 4 and 5 and Theorem 4. Assuming that Conjecture 1 holds, Proposition 3 furthermore implies that  $ME$  is a refinement of  $MC$  since every covering set is also an extending set. We refer to Figure 1 for an example tournament  $T$  where  $ME(T)$  happens to be *strictly* contained in  $MC(T)$ .<sup>9</sup>

<sup>8</sup>  $UC$ -exclusivity is the property  $\gamma^{**}$  used in the axiomatization of  $MC$  (Laslier, 1997).

<sup>9</sup> This is also the case for a tournament of order eight given by Dutta (1990).




Figure 1: Example tournament  $T = (A, \succ)$  where  $MC$  and  $ME$  differ ( $MC(T) = A$  and  $ME(T) = A \setminus \{a_{10}\}$ ).  $a_{10}$  only dominates  $a_3$ ,  $a_6$ , and  $a_9$ .

A remarkable property of  $ME$  is that, just like  $BA$ , it is capable of ruling out alternatives in *regular* tournaments, i.e., it satisfies IRR. No irregular tournament solution is known to satisfy all four basic properties. However, if Conjecture 1 were true,  $ME$  would be such a concept.

## 5 Retentiveness and Stability

Motivated by cooperative majority voting, Schwartz (1990) introduced a tournament solution based on a notion he calls *retentiveness*. It turns out that retentiveness bears some similarities to stability. For example, the top cycle can be represented as a minimal stable set as well as a minimal retentive set, albeit using different underlying tournament solutions.

### 5.1 The Tournament Equilibrium Set

The intuition underlying retentive sets is that alternative  $a$  is only “properly” dominated by alternative  $b$  if  $b$  is chosen among  $a$ ’s dominators by some underlying tournament solution  $S$ . A set of alternatives is then called  $S$ -retentive if none of its elements is properly dominated by some outside alternative with respect to  $S$ .<sup>10</sup>

**Definition 15.** Let  $S$  be a tournament solution and  $T = (A, \succ)$  a tournament. Then,  $B \subseteq A$  is *retentive* in  $T$  with respect to tournament solution  $S$  (or  *$S$ -retentive*) if  $B \neq \emptyset$  and  $S(\overline{D}(b)) \subseteq B$  for all  $b \in B$  such that  $\overline{D}(b) \neq \emptyset$ . The set of  $S$ -retentive sets for a given tournament  $T = (A, \succ)$  will be denoted by  $\mathcal{R}_S(T) = \{B \subseteq A \mid B \text{ is } S\text{-retentive in } T\}$ .

<sup>10</sup>In analogy to the discussion at the beginning of Section 4.1, a set of alternatives  $B$  may be called retentive if it consists precisely of those alternatives not upset by  $B$ . Here,  $a$  is upset by  $B$  if  $a \notin \bigcup_{b \in B} S(\overline{D}(b))$  for some underlying solution concept  $S$ .

$S$ -retentive sets are guaranteed to exist since the set of all alternatives  $A$  is trivially  $S$ -retentive in  $(A, \succ)$  for all tournament solutions  $S$ . In analogy to Definition 13, the union of minimal  $S$ -retentive sets defines a tournament solution.

**Definition 16.** Let  $S$  be a tournament solution. Then, the tournament solution  $\mathring{S}$  is defined as

$$\mathring{S}(T) = \bigcup_{\subseteq} \min(\mathcal{R}_S(T)).$$

It is easily verified that  $\mathring{S}$  is well-defined as a tournament solution as there are no  $S$ -retentive sets that do not contain the Condorcet winner whenever one exists.

As an example, consider the tournament solution  $S_{\mathcal{M}_1}$  that always returns all alternatives, i.e.,  $S_{\mathcal{M}_1}((A, \succ)) = A$ . The unique minimal retentive set with respect to  $S_{\mathcal{M}_1}$  is the top cycle, that is  $TC = \widehat{S}_{\mathcal{M}_2} = \mathring{S}_{\mathcal{M}_1}$ .

Schwartz introduced retentiveness in order to recursively define the *tournament equilibrium set* ( $TEQ$ ) as the union of minimal  $TEQ$ -retentive sets. This recursion is well-defined because the order of the dominator set of any alternative is strictly smaller than the order of the original tournament.

**Definition 17** (Schwartz, 1990). The *tournament equilibrium set* ( $TEQ$ ) of a tournament  $T$  is defined recursively as  $TEQ(T) = T\mathring{E}Q(T)$ .

In other words,  $TEQ$  is the unique fixed point of the  $\circ$ -operator.

Schwartz conjectured that every tournament admits a *unique* minimal  $TEQ$ -retentive set. Despite several attempts to prove or disprove this statement (e.g., Laffond et al., 1993a; Houy, 2009), the statement has remained a conjecture. A recent computer analysis failed to find a counter-example in all tournaments of order 12 or less and a fairly large number of random tournaments (Brandt et al., 2010).

**Conjecture 2** (Schwartz, 1990).  $\mathcal{R}_{TEQ}$  is directed.

It is easily appreciated that the non-empty intersection of two  $S$ -retentive sets is also  $S$ -retentive. As a consequence, Conjecture 2 is equivalent to the statement that there are no two disjoint  $TEQ$ -retentive sets in any tournament. Unfortunately, and somewhat surprisingly, it is not known whether  $TEQ$  satisfies *any* of the basic properties defined in Section 2.5. However, Laffond et al. (1993a) and Houy (2009) have shown that  $TEQ$  satisfies any of the basic properties if and only if  $\mathcal{R}_{TEQ}$  is directed, and is strictly contained in  $MC$  if  $\mathcal{R}_{TEQ}$  is directed. We will strengthen the last statement by showing that  $TEQ$  is also strictly contained in  $ME$  if  $\mathcal{R}_{TEQ}$  is directed.

## 5.2 Inclusion of $TEQ$ in $ME$

In Section 3.2, the Banks set was characterized as the finest tournament solution satisfying strong retentiveness. It turns out that, if  $\mathcal{R}_{TEQ}$  is directed,  $TEQ$  is the finest tournament solution satisfying a very natural weakening of strong retentiveness, where the inclusion of the choice sets of dominator sets is only required to hold for alternatives contained in the original choice set.

**Definition 18.** A tournament solution  $S$  satisfies (weak) *retentiveness* if  $S(\overline{D}(a)) \subseteq S(T)$  for all tournaments  $T$  and  $a \in S(T)$ .

In other words, a tournament solution  $S$  satisfies retentiveness if and only if  $S(T)$  is  $S$ -retentive for all tournaments  $T$ . It follows from Schwartz's axiomatization of  $TEQ$  that  $TEQ$  is the finest tournament solution satisfying retentiveness, given that Conjecture 2 is true.

**Theorem 5** (Schwartz, 1990). *TEQ is the finest tournament solution satisfying retentiveness if  $\mathcal{R}_{TEQ}$  is directed.*

The fact that  $S_Q$  satisfies strong retentiveness for every concept of qualified subsets  $\mathcal{Q}$  can be used to show that every  $S_Q$ -stable set is  $\hat{S}_Q$ -retentive, which has a number of useful consequences.

**Lemma 4.** *Let  $\mathcal{Q}$  be a concept of qualified subsets such that  $\mathcal{S}_{S_Q}$  is directed for all tournaments of order  $n$  or less. Then,  $\mathcal{S}_{S_Q}(T) \subseteq \mathcal{R}_{\hat{S}_Q}(T)$  for all tournaments  $T$  of order  $n + 1$ .*

*Proof.* Let  $T = (A, \succ)$  be a tournament of order  $n + 1$ ,  $B \in \mathcal{S}_{S_Q}(T)$ ,  $b \in B$ , and  $C = \overline{D}_B(b)$ . We show that  $C$  is  $S_Q$ -stable in  $\overline{D}(b)$ . Let  $a \in \overline{D}(b) \setminus B$  and consider the tournament restricted to  $C \cup \{a\}$ . We know from  $B$ 's stability that  $S_Q(B \cup \{a\}) \subseteq B$ . Furthermore, since  $S_Q$  satisfies strong retentiveness (Lemma 1),  $S_Q(C \cup \{a\}) \subseteq C$ , which shows that  $C$  is  $S_Q$ -stable in  $C \cup \{a\}$ .  $C \cup \{a\}$  is of order  $n$  or less. Hence, there is a unique minimal  $S_Q$ -stable set in  $C \cup \{a\}$ , which is contained in  $C$ .  $\square$

**Theorem 6.** *Let  $\mathcal{Q}$  be a concept of qualified subsets. Then,*

- (i)  $\hat{S}_Q$  satisfies retentiveness if  $\mathcal{S}_Q$  is directed,
- (ii)  $\mathcal{S}_Q$  is directed if  $\mathcal{R}_{TEQ}$  is directed, and
- (iii)  $TEQ \subseteq \hat{S}_Q$  if  $\mathcal{R}_{TEQ}$  is directed.

*Proof.* We prove all statements simultaneously by induction on the tournament order  $n$ . The basis is straightforward. Now, assume that all three implications hold for tournaments of order  $n$  or less. If  $\mathcal{S}_Q$  is directed for tournaments of order  $n + 1$  or less, we can apply Lemma 4 to show that every  $S_Q$ -stable set is  $\hat{S}_Q$ -retentive in such tournaments. Hence,  $\hat{S}_Q$  satisfies retentiveness and the first statement holds for tournaments of order  $n + 1$ .

For the second statement, assume for contradiction that there is a tournament  $T = (A, \succ)$  of order  $n + 1$  that contains two disjoint  $S_Q$ -stable sets  $B_1$  and  $B_2$  (while  $\mathcal{S}_Q$  is directed for all tournaments of order  $n$  or less and  $\mathcal{R}_{TEQ}$  is directed for all tournaments of order  $n + 1$  or less, including  $T$ ). It follows from Lemma 4 that  $B_1$  and  $B_2$  are also  $\hat{S}_Q$ -retentive. Moreover, the induction hypothesis of the third statement implies that  $TEQ(\overline{D}(a)) \subseteq \hat{S}_Q(\overline{D}(a))$  for all  $a \in A$ , which implies that  $B_1$  and  $B_2$  are  $TEQ$ -retentive, a contradiction.

In order to show the third statement, let  $\mathcal{R}_{TEQ}$  be directed for tournaments of order  $n + 1$  or less. It follows from the second implication that  $\mathcal{S}_Q$  of such tournaments is directed as well and from the first that  $\hat{S}_Q$  satisfies retentiveness for such tournaments. We know from Theorem 5 that  $TEQ$  is contained in all tournament solutions that satisfy retentiveness. Hence,  $TEQ(T) \subseteq \hat{S}_Q(T)$  for all tournaments of order  $n + 1$ .  $\square$

In other words,  $TC$  and  $MC$  satisfy retentiveness and  $ME$  satisfies retentiveness if Conjecture 1 holds. Furthermore, Conjecture 2 is at least as strong as Conjecture 1. Similarly, the directedness of  $\mathcal{S}_M$ , which was proved by Dutta (1988) (see Theorem 2), also follows from Conjecture 2. Finally, given that Conjecture 2 holds,  $TEQ$  is a refinement of all tournament solutions  $\hat{S}_Q$  where  $Q$  is a concept of qualified subsets. In particular, we have the following.

**Corollary 2.**  *$TEQ \subseteq ME$  if  $\mathcal{R}_{TEQ}$  is directed*

The remaining question is whether  $TEQ$  and  $ME$  are actually different solution concepts. The tournament given in Figure 2 demonstrates that this is indeed the case.




Figure 2: Example tournament  $T = (A, \succ)$  where  $ME$  and  $TEQ$  differ ( $ME(T) = A$  and  $TEQ(T) = A \setminus \{a_5\}$ ). In particular,  $A \setminus \{a_5\}$  is no extending set since  $a_5 \in BA(A)$  via the non-extendable transitive set  $\{a_5, a_6, a_7, a_8\}$ .

### 5.3 TEQ as a Minimal Stable Set

A natural question is whether  $TEQ$  itself can be represented as a minimal stable set. The following two lemmas establish that this is indeed the case if  $\mathcal{R}_{TEQ}$  is directed. We first show that every  $S$ -retentive set is  $\hat{S}$ -stable if  $S$  satisfies WSP for strictly smaller tournaments.

**Lemma 5.** *Let  $S$  be a tournament solution that satisfies WSP for all tournaments of order  $n$  or less. Then,  $\mathcal{R}_S(T) \subseteq \mathcal{S}_{\hat{S}}(T)$  for all tournaments  $T$  of order  $n + 1$ .*

*Proof.* Let  $T = (A, \succ)$  be a tournament of order  $n + 1$  and  $B$  an  $S$ -retentive set in  $A$ . If  $B = A$ , the statement is trivially satisfied. Otherwise, let  $a \in A \setminus B$ . We first show that  $B$  is  $S$ -retentive in  $B \cup \{a\}$ . Let  $b$  be an arbitrary alternative in  $B$ .  $S$ -retentiveness implies  $S(\overline{D}(b)) \subseteq B$  and WSP implies  $S(\overline{D}_{B \cup \{a\}}(b)) \subseteq S(\overline{D}(b))$ . As a consequence,  $S(\overline{D}_{B \cup \{a\}}(b)) \subseteq B$ , and thus  $B$  is  $S$ -retentive in  $B \cup \{a\}$ . It remains to be shown

that  $a$  is not contained in another minimal  $S$ -retentive subset of  $B \cup \{a\}$ . Assume for contradiction that  $a$  is contained in some minimal  $S$ -retentive set. If  $\{a\}$  itself were an  $S$ -retentive set,  $a$  would be the Condorcet winner in  $B \cup \{a\}$ , contradicting the fact that  $B$  is  $S$ -retentive in  $B \cup \{a\}$ . Now let  $C \subset B \cup \{a\}$  with  $a \in C$  and  $|C| > 1$  be a minimal  $S$ -retentive set. This implies that  $B \cap C$  is also  $S$ -retentive, contradicting the minimality of  $C$ . It follows that  $B$  is  $\dot{S}$ -stable.  $\square$

The next lemma shows that every  $TEQ$ -stable set is also  $TEQ$ -retentive, assuming the directedness of  $\mathcal{R}_{TEQ}$ .

**Lemma 6.**  $\mathcal{S}_{TEQ} \subseteq \mathcal{R}_{TEQ}$  if  $\mathcal{R}_{TEQ}$  is directed.

*Proof.* We prove the statement by induction on the tournament order  $n$ . We may assume that  $\mathcal{R}_{TEQ}$  is directed for all tournaments of order  $n + 1$  or less and that  $\mathcal{S}_{TEQ} \subseteq \mathcal{R}_{TEQ}$  for all tournaments of order  $n$  or less. Let  $T = (A, \succ)$  be a tournament of order  $n + 1$  and  $B \subseteq A$  a  $TEQ$ -stable set in  $T$ . In other words, if we let  $A \setminus B = \{a_1, \dots, a_k\}$ , then  $TEQ(B \cup \{a_i\}) \subseteq B$  for all  $1 \leq i \leq k$ . Now, in order to show that  $B$  is  $TEQ$ -retentive, consider an arbitrary  $b \in B$  and let  $C = \overline{D}_B(b)$ . By definition of  $TEQ$ ,  $TEQ(C \cup \{a_i\}) \subseteq C$  for all  $a_i \in \overline{D}_{A \setminus B}(b)$ , i.e.,  $C$  is  $TEQ$ -stable in  $\overline{D}(b)$ . Since  $\mathcal{S}_{TEQ}(\overline{D}(b)) \subseteq \mathcal{R}_{TEQ}(\overline{D}(b))$ ,  $C$  is also  $TEQ$ -retentive in  $\overline{D}(b)$  and, due to the directedness of  $\mathcal{R}_{TEQ}(\overline{D}(b))$ ,  $TEQ(\overline{D}(b)) \subseteq C$ . Hence,  $B$  is  $TEQ$ -retentive in  $T$ .  $\square$

We have now cleared the ground for the main result of this section.

**Theorem 7.**  $\mathcal{S}_{TEQ}$  is directed if and only if  $\mathcal{R}_{TEQ}$  is directed.  $TEQ = \widehat{TEQ}$  if  $\mathcal{R}_{TEQ}$  is directed.

*Proof.* We first prove the following two implications by induction on the tournament order  $n$ : (i) if  $\mathcal{R}_{TEQ}$  is directed, then  $TEQ$  satisfies WSP and  $\mathcal{R}_{TEQ} \subseteq \mathcal{S}_{TEQ}$ ; (ii) if  $\mathcal{S}_{TEQ}$  is directed, then  $\mathcal{R}_{TEQ}$  is directed.

The basis is straightforward. Assume that both implications hold for tournaments of order  $n$  and let  $T = (A, \succ)$  be a tournament of order  $n + 1$ . In order to prove the first statement, assume that  $\mathcal{R}_{TEQ}$  is directed for all tournaments of order  $n + 1$  or less and let  $B \subseteq A$  such that  $TEQ(T) \subseteq B$ . The induction hypothesis implies that  $TEQ$  satisfies WSP in all dominator sets  $\overline{D}(a)$  for  $a \in A$ . Hence,  $TEQ(T)$  is  $TEQ$ -retentive in  $B$  and, due to the directedness of  $\mathcal{R}_{TEQ}(T)$ ,  $TEQ(B) \subseteq TEQ(T)$ . Moreover, Lemma 5 shows that  $\mathcal{R}_{TEQ}(T) \subseteq \mathcal{S}_{TEQ}(T)$  since  $TEQ = \dot{TEQ}$  by definition.

For the second statement, assume that  $\mathcal{S}_{TEQ}$  is directed for tournaments of order  $n + 1$  or less. It follows from the induction hypothesis that  $\mathcal{R}_{TEQ}$  is directed for tournaments of order  $n$  or less and from the induction hypothesis of the first statement that  $TEQ$  satisfies WSP in these tournaments. Now, assume for contradiction that there two disjoint  $TEQ$ -retentive sets in  $T$ . According to Lemma 5, these sets are also  $TEQ$ -stable, which contradicts the directedness of  $\mathcal{S}_{TEQ}(T)$ .

Finally, assume that  $\mathcal{R}_{TEQ}$  is directed. The first statement and Lemma 6 establish that  $\mathcal{R}_{TEQ} = \mathcal{S}_{TEQ}$  and hence that  $TEQ = \widehat{TEQ}$ . As a consequence,  $\mathcal{S}_{TEQ}$  has to be directed as well, which concludes the proof.  $\square$

Combining the previous theorem and the definition of  $TEQ$ , Conjecture 2 entails that

$$TEQ = T\overset{\circ}{E}Q = \widehat{TEQ}.$$

## 6 Quantitative Concepts

In Section 3, several solution concepts were defined by collecting the maximal elements of inclusion-maximal qualified subsets. In this section, we replace maximality with respect to set inclusion by maximality with respect to *cardinality*, i.e., we look at qualified subsets containing the largest number of elements.

### 6.1 Maximal Qualified Subsets

For every set of finite sets  $\mathcal{S}$ , define  $\max_{\leq}(\mathcal{S}) = \{S \in \mathcal{S} \mid |S| \geq |S'| \text{ for all } S' \in \mathcal{S}\}$ . In analogy to Definition 10, we can now define a solution concept that yields the maximal elements of the largest qualified subsets.

**Definition 19.** Let  $\mathcal{Q}$  be a concept of qualified subsets. Then, the tournament solution  $S_{\mathcal{Q}}^{\#}$  is defined as

$$S_{\mathcal{Q}}^{\#}(T) = \{\max_{\prec}(B) \mid B \in \max_{\leq}(\mathcal{Q}(T))\}.$$

Obviously,  $S_{\mathcal{Q}}^{\#} \subseteq S_{\mathcal{Q}}$  for all concepts of qualified subsets  $\mathcal{Q}$ . For the concept of qualified subsets  $\mathcal{M}$ , i.e., the set of subsets that admit a maximal element, we obtain the Copeland set.

**Copeland set.**  $S_{\mathcal{M}}^{\#}(T)$  returns the *Copeland set*  $CO(T)$  of a tournament  $T$ , i.e., the set of all alternatives whose dominion is of maximal size (Copeland, 1951).<sup>11</sup>

### 6.2 Minimal Stable Sets

When the Copeland set is taken as the basis for stable sets, some tournaments contain more than one inclusion-minimal externally stable set and, even worse, do not admit a set that satisfies both internal *and* external stability (see Figure 3 for an example).

However, as it turns out, every tournament is the *summary* of some tournament consisting only of homogeneous components that admits a unique internally and externally *CO*-stable set. For example, when replacing  $a_4$  and  $a_5$  in the tournament given in Figure 3 with 3-cycle components, the set of all alternatives is internally and externally stable. The following definition captures this strengthened notion of stability.

**Definition 20.** Let  $S$  be a tournament solution and  $\tilde{T} = (\{1, \dots, k\}, \tilde{\succ})$  a tournament. Then,  $\tilde{B} \subseteq \{1, \dots, k\}$  is *strongly stable* with respect to tournament solution

<sup>11</sup>This set is usually attributed to Copeland despite the fact that Zermelo (1929) and Llull (as early as 1283, see Hägele and Pukelsheim 2001) have suggested equivalent concepts much earlier.




Figure 3: Example tournament  $T = (A, \succ)$  that does not contain an internally and externally  $CO$ -stable set. There are eight externally  $CO$ -stable sets (e.g.,  $A$ ,  $\{a_2, a_3, a_4, a_5\}$ , or  $\{a_1, a_2, a_5\}$ ), none of which is internally stable.

$S$  (or *strongly  $S$ -stable*) if there exist homogeneous tournaments  $T_1, \dots, T_k$  on  $k$  disjoint sets  $B_1, \dots, B_k \subseteq X$  such that  $B = \bigcup_{i \in \tilde{B}} B_i$  is internally and externally  $S$ -stable in  $T = \Pi(\tilde{T}, T_1, \dots, T_k)$ . The set of strongly  $S$ -stable sets for a given tournament  $T = (A, \succ)$  will be denoted by  $\tilde{\mathcal{S}}_S(T) = \{B \subseteq A \mid B \text{ is strongly } S\text{-stable in } T\}$ .

Now, in analogy to  $\hat{S}$ , we can define a tournament solution that yields the minimal strongly  $S$ -stable set with respect to some underlying tournament solution  $S$ .

**Definition 21.** Let  $S$  be a tournament solution. Then, the tournament solution  $\hat{S}$  is defined as

$$\hat{S}(T) = \bigcup_{\subseteq} \min(\tilde{\mathcal{S}}_S(T)).$$

The following result follows from observations made independently by Laffond et al. (1993b) and Fisher and Ryan (1995) (see also Laslier, 1997, 2000).

**Theorem 8** (Laffond et al., 1993b).  $|\tilde{\mathcal{S}}_{S_{\mathcal{M}}}^{\#}(T)| = 1$  for all tournaments  $T$ .

The unique strongly stable set with respect to the Copeland set is known as the *bipartisan set*.

**Bipartisan set.** The *bipartisan set* of a tournament  $T$  is given by  $BP(T) = \hat{S}_{\mathcal{M}}^{\#}(T)$ . It was originally defined as the set of alternatives corresponding to the support of the unique Nash equilibrium of the tournament game (Laffond et al., 1993b). The tournament game of a tournament is the two-player zero-sum game given by its adjacency matrix.

$BP$  satisfies all basic properties, composition-consistency, and is contained in  $MC$ . Its relationship with  $BA$  is unknown (Laffond et al., 1993b).

Interestingly, minimality is not required for  $\hat{S}_{\mathcal{M}}^{\#}$ , because there is always exactly one strongly  $S_{\mathcal{M}}^{\#}$ -stable set. Further observe that internally and externally stable sets and strongly stable sets coincide when the underlying solution concept satisfies **COM**. This is the case for  $MC$ ,  $ME$ , and  $TEQ$ . Even though,  $TC$  does not satisfy **COM**, it is easily verified that replacing alternatives with homogeneous components does not affect the top cycle of a tournament. It is thus possible to define all mentioned concepts using minimal strongly stable sets instead of stable sets (see Table 1).

$S$	$\widehat{S}$
Condorcet non-losers ( $CNL$ )	Top cycle ( $TC$ )
Copeland set ( $CO$ )	Bipartisan set ( $BP$ )
Uncovered set ( $UC$ )	Minimal covering set ( $MC$ )
Banks set ( $BA$ )	Minimal extending set ( $ME$ )
Tournament equilibrium set ( $TEQ$ )	Tournament equilibrium set ( $TEQ$ )

Table 1: Tournament solutions and their minimal strongly stable counterparts. The representation of  $TEQ$  as a stable set relies on Conjecture 2.

## 7 Conclusion

We proposed a unifying treatment of tournament solutions based on maximal qualified subsets and minimal stable sets. Given the results of Section 4 and Section 5, a central role in the theory of tournament solutions may be ascribed to Conjecture 2, a statement of considerable mathematical depth. Conjecture 2 has a number of appealing consequences on minimal stable sets, some of which have been proved already.

- (i) Every tournament  $T$  admits a unique *minimal dominant set*  $TC(T)$  (as shown by Good, 1971).  $TC$  satisfies all basic properties and is the finest solution concept satisfying  $SSP$  and  $CNL$ -exclusivity.
- (ii) Every tournament  $T$  admits a unique *minimal covering set*  $MC(T)$  (as shown by Dutta, 1988).  $MC$  satisfies all basic properties and is the finest solution concept satisfying  $SSP$  and  $UC$ -exclusivity.
- (iii) Every tournament  $T$  admits a unique *minimal extending set*  $ME(T)$  (open problem).  $ME$  satisfies all basic properties and is the finest solution concept satisfying  $SSP$  and  $BA$ -exclusivity.
- (iv) Every tournament  $T$  admits a unique *minimal  $TEQ$ -retentive set*  $TEQ(T)$  (open problem).  $TEQ$  satisfies all basic properties and is the finest solution concept satisfying retentiveness and the finest solution concept  $S$  such that  $S$  satisfies  $SSP$  and, for all tournaments  $T = (A, \succ)$ ,  $S(A) = A \setminus \{a\}$  only if  $a \notin S(\overline{D}(b))$  for every  $b \in A$ .
- (v) The following inclusion relationships hold:  $TEQ \subseteq ME \subseteq MC \subseteq TC$  and  $ME \subseteq BA$ .<sup>12</sup>

Conjecture 1 is a weaker version of Conjecture 2, which implies all of the above statements except those that involve  $TEQ$ .

<sup>12</sup>A consequence of these inclusions is that deciding whether an alternative is contained in the minimal extending set of a tournament is NP-hard. This follows from a proof by Brandt et al. (2010), which establishes hardness of all solution concepts that are sandwiched between  $BA$  and  $TEQ$ .

Table 2 and Figure 4 summarize the properties and set-theoretic relationships of the considered tournament solutions, respectively.

Solution Concept	Origin	MON	IUA	WSP	SSP	COM	IRR
$S_{M_2}$ (CNL)		✓	✓	✓	–	–	–
$S_M$ (UC)	Fishburn (1977); Miller (1980)	✓	–	✓	–	✓	–
$S_{M^*}$ (BA)	Banks (1985)	✓	–	✓	–	✓	✓
$\hat{S}_{M_2}$ (TC)	Good (1971); Smith (1973)	✓	✓	✓	✓	–	–
$\hat{S}_M$ (MC)	Dutta (1988)	✓	✓	✓	✓	✓	–
$\hat{S}_{M^*}$ (ME)		✓ <sup>a</sup>	✓ <sup>a</sup>	✓ <sup>a</sup>	✓ <sup>a</sup>	✓	✓
$\overline{TEQ}$ (TEQ)	Schwartz (1990)	✓ <sup>b</sup>	✓ <sup>b</sup>	✓ <sup>b</sup>	✓ <sup>b</sup>	✓	✓
$S_M^\#$ (CO)	Copeland (1951)	✓	–	–	–	–	–
$\hat{S}_M^\#$ (BP)	Laffond et al. (1993b)	✓	✓	✓	✓	✓	–

<sup>a</sup>This statement relies on Conjecture 1.

<sup>b</sup>This statement relies on Conjecture 2.

Table 2: Properties of solution concepts (MON: monotonicity, IUA: independence of unchosen alternatives, WSP: weak superset property, SSP: strong superset property, COM: composition-consistency, IRR: irregularity). See Laslier (1997) for all results not shown in this paper.

## Acknowledgments

I am indebted to Paul Harrenstein for countless valuable discussions. Furthermore, I thank Haris Aziz, Markus Brill, and Felix Fischer for providing helpful feedback on drafts of this paper. This material is based on work supported by the Deutsche Forschungsgemeinschaft under grants BR 2312/3-2, BR 2312/3-3, and BR 2312/7-1.

Preliminary results of this paper were presented at the 9th International Meeting of the Society of Social Choice and Welfare (Montreal, June 2008).

## References

- K. J. Arrow and H. Raynaud. *Social Choice and Multicriterion Decision-Making*. MIT Press, 1986.
- J. S. Banks. Sophisticated voting outcomes and agenda control. *Social Choice and Welfare*, 3:295–306, 1985.
- K. Basu and J. Weibull. Strategy subsets closed under rational behavior. *Economics Letters*, 36:141–146, 1991.
- G. Bordes. On the possibility of reasonable consistent majoritarian choice: Some positive results. *Journal of Economic Theory*, 31:122–132, 1983.




Figure 4: Set-theoretic relationships between qualitative tournament solutions.  $BA$  and  $MC$  are not included in each other, but they always intersect. The inclusion of  $TEQ$  in  $ME$  relies on Conjecture 2 and that of  $ME$  in  $MC$  on Conjecture 1 (which is implied by Conjecture 2).  $CO$  is contained in  $UC$  but may be disjoint from  $MC$  and  $BA$ . The exact location of  $BP$  in this diagram is unknown ( $BP$  is contained in  $MC$  and is a superset or subset of  $TEQ$  in all known instances (Laslier, 1997)).

- D. Bouyssou, T. Marchant, M. Pirlot, A. Tsoukiàs, and P. Vincke. *Evaluation and Decision Models: Stepping Stones for the Analyst*. Springer-Verlag, 2006.
- F. Brandt and P. Harrenstein. Characterization of dominance relations in finite coalitional games. *Theory and Decision*, 69(2):233–256, 2010.
- F. Brandt, F. Fischer, P. Harrenstein, and M. Mair. A computational analysis of the tournament equilibrium set. *Social Choice and Welfare*, 34(4):597–609, 2010.
- A. H. Copeland. A ‘reasonable’ social welfare function. Mimeographed, University of Michigan Seminar on Applications of Mathematics to the Social Sciences, 1951.
- J. Duggan and M. Le Breton. Dutta’s minimal covering set and Shapley’s saddles. *Journal of Economic Theory*, 70:257–265, 1996.
- B. Dutta. Covering sets and a new Condorcet choice correspondence. *Journal of Economic Theory*, 44:63–80, 1988.
- B. Dutta. On the tournament equilibrium set. *Social Choice and Welfare*, 7(4):381–383, 1990.
- P. C. Fishburn. Condorcet social choice functions. *SIAM Journal on Applied Mathematics*, 33(3):469–489, 1977.

- D. C. Fisher and J. Ryan. Tournament games and positive tournaments. *Journal of Graph Theory*, 19(2):217–236, 1995.
- S. Gaspers and M. Mnich. Feedback vertex sets in tournaments. In *Proceedings of the 18th European Symposium on Algorithms (ESA)*, Lecture Notes in Computer Science (LNCS). Springer-Verlag, 2010. Forthcoming.
- D. B. Gillies. Solutions to general non-zero-sum games. In A. W. Tucker and R. D. Luce, editors, *Contributions to the Theory of Games IV*, volume 40 of *Annals of Mathematics Studies*, pages 47–85. Princeton University Press, 1959.
- I. J. Good. A note on Condorcet sets. *Public Choice*, 10:97–101, 1971.
- G. Hägele and F. Pukelsheim. The electoral writings of Ramon Llull. *Studia Lulliana*, 41(97):3–38, 2001.
- N. Houy. Still more on the tournament equilibrium set. *Social Choice and Welfare*, 32:93–99, 2009.
- G. Laffond, J.-F. Laslier, and M. Le Breton. More on the tournament equilibrium set. *Mathématiques et sciences humaines*, 31(123):37–44, 1993a.
- G. Laffond, J.-F. Laslier, and M. Le Breton. The bipartisan set of a tournament game. *Games and Economic Behavior*, 5:182–201, 1993b.
- J.-F. Laslier. *Tournament Solutions and Majority Voting*. Springer-Verlag, 1997.
- J.-F. Laslier. Aggregation of preferences with a variable set of alternatives. *Social Choice and Welfare*, 17:269–282, 2000.
- N. R. Miller. A new solution set for tournaments and majority voting: Further graph-theoretical approaches to the theory of voting. *American Journal of Political Science*, 24(1):68–96, 1980.
- H. Moulin. Choosing from a tournament. *Social Choice and Welfare*, 3:271–291, 1986.
- H. Moulin. *Axioms of Cooperative Decision Making*. Cambridge University Press, 1988.
- J. F. Nash. Non-cooperative games. *Annals of Mathematics*, 54(2):286–295, 1951.
- T. Schwartz. *The Logic of Collective Choice*. Columbia University Press, 1986.
- T. Schwartz. Cyclic tournaments and cooperative majority voting: A solution. *Social Choice and Welfare*, 7:19–29, 1990.
- L. Shapley. Some topics in two-person games. In M. Dresher, L. S. Shapley, and A. W. Tucker, editors, *Advances in Game Theory*, volume 52 of *Annals of Mathematics Studies*, pages 1–29. Princeton University Press, 1964.

- J. H. Smith. Aggregation of preferences with variable electorate. *Econometrica*, 41(6): 1027–1041, 1973.
- J. von Neumann and O. Morgenstern. *Theory of Games and Economic Behavior*. Princeton University Press, 1944.
- R. B. Wilson. The finer structure of revealed preference. *Journal of Economic Theory*, 2(4):348–353, 1970.
- G. J. Woeginger. Banks winners in tournaments are difficult to recognize. *Social Choice and Welfare*, 20:523–528, 2003.
- E. Zermelo. Die Berechnung der Turnier-Ergebnisse als ein Maximumproblem der Wahrscheinlichkeitsrechnung. *Mathematische Zeitschrift*, 29(1):436–460, 1929.