

# Some Remarks on Sign-Balanced and Maj-Balanced Posets

Richard P. Stanley<sup>1</sup>

Department of Mathematics  
Massachusetts Institute of Technology  
Cambridge, MA 02139  
*e-mail:* rstan@math.mit.edu

version of 26 October 2002

## 1 Introduction.

Let  $P$  be an  $n$ -element poset (partially ordered set), and let  $\omega : P \rightarrow [n] = \{1, 2, \dots, n\}$  be a bijection, called a *labeling* of  $P$ . We call the pair  $(P, \omega)$  a *labelled poset*. A *linear extension* of  $P$  is an order-preserving bijection  $f : P \rightarrow [n]$ . We can regard  $f$  as defining a permutation  $\pi = \pi(f)$  of the set  $[n]$  given by  $\pi(i) = j$  if  $f(\omega^{-1}(j)) = i$ . We write  $\pi$  in the customary way as a word  $a_1 a_2 \cdots a_n$ , where  $\pi(i) = a_i = \omega(f^{-1}(i))$ . We will say for instance that  $f$  is an *even linear extension* of  $(P, \omega)$  if  $\pi$  is an even permutation (i.e., an element of the alternating group  $\mathfrak{A}_n$ ). Let  $\mathcal{E}_P$  denote the set of linear extensions of  $P$ , and set  $\mathcal{L}_{P, \omega} = \{\pi(f) : f \in \mathcal{E}_P\}$ .

We say that  $(P, \omega)$  is *sign-balanced* if  $\mathcal{L}_{P, \omega}$  contains the same number of even permutations as odd permutations. Note that the parity of a linear extension  $f$  depends on the labeling  $\omega$ . However, the notion of sign-balanced depends only on  $P$ , since changing the labeling of  $P$  simply multiplies the elements of  $\mathcal{L}_{P, \omega}$  by a fixed permutation in  $\mathfrak{S}_n$ , the symmetric group of all permutations of  $[n]$ . Thus we can simply say that  $P$  is sign-balanced without specifying  $\omega$ .

We say that a function  $\vartheta : \mathcal{E}_P \rightarrow \mathcal{E}_P$  is *parity-reversing* (respectively, *parity-preserving*) if for all  $f \in \mathcal{E}_P$ , the permutations  $\pi(f)$  and  $\pi(\vartheta(f))$  have

---

<sup>1</sup>Partially supported by NSF grant #DMS-9988459.

opposite parity (respectively, the same parity). Note that the properties of parity-reversing and parity-preserving do not depend on  $\omega$ ; indeed,  $\vartheta$  is parity-reversing (respectively, parity-preserving) if and only if for all  $f \in \mathcal{E}_P$ , the permutation  $\vartheta f \circ f^{-1} \in \mathfrak{S}_n$  is odd (respectively, even),

Sign-balanced posets were first considered by Ruskey [16]. He established the following result, which shows that many combinatorially occurring classes of posets, such as geometric lattices and Eulerian posets, are sign-balanced.

**1.1 Theorem.** *Suppose  $\#P \geq 2$ . If every nonminimal element of the poset  $P$  is greater than at least two minimal elements, then  $P$  is sign-balanced.*

**Proof.** Let  $\pi = a_1 a_2 a_3 \cdots a_n \in \mathcal{L}_{P,\omega}$ . Let  $\pi' = \pi(1, 2) = a_2 a_1 a_3 \cdots a_n \in \mathfrak{S}_n$ . (We always multiply permutations from right to left.) By the hypothesis on  $P$ , we also have  $\pi' \in \mathcal{L}_{P,\omega}$ . The map  $\pi \mapsto \pi'$  is a parity-reversing involution (i.e., exactly one of  $\pi$  and  $\pi'$  is an even permutation) on  $\mathcal{L}_{P,\omega}$ , and the proof follows.  $\square$

The above proof illustrates what will be our basic technique for showing that a poset  $P$  is sign-balanced, viz., giving a bijection  $\sigma : \mathcal{L}_{P,\omega} \rightarrow \mathcal{L}_{P,\omega}$  such that  $\pi$  and  $\sigma(\pi)$  have opposite parity for all  $\pi \in \mathcal{L}_{P,\omega}$ . Equivalently, we are giving a parity-reversing bijection  $\vartheta : \mathcal{E}_P \rightarrow \mathcal{E}_P$ .

In 1992 Ruskey [17, §5, item 6] conjectured as to when the product  $\mathbf{m} \times \mathbf{n}$  of two chains of cardinalities  $m$  and  $n$  is sign-balanced, viz.,  $m, n > 1$  and  $m \equiv n \pmod{2}$ . Ruskey proved this when  $m$  and  $n$  are both even by giving a simple parity-reversing involution, which we generalize in Proposition 4.1 and Corollary 4.2. Ruskey's conjecture for  $m$  and  $n$  odd was proved by D. White [27], who also computed the “imbalance” between even and odd linear extensions in the case when exactly one of  $m$  and  $n$  is even (stated here as Theorem 3.5). None of our theorems below apply to the case when  $m$  and  $n$  are both odd. Ruskey [17, §5, item 5] also asked what order ideals  $I$  (defined below) of  $\mathbf{m} \times \mathbf{n}$  are sign-balanced. Such order ideals correspond to integer partitions  $\lambda$  and will be denoted  $P_\lambda$ ; the linear extensions of  $P_\lambda$  are equivalent to standard Young tableaux (SYT) of shape  $\lambda$ . White [27] also determined some additional  $\lambda$  for which  $P_\lambda$  is sign-balanced, and our results below will give some further examples. In Sections 5 and 6 we consider some analogous

questions for the parity of the major index of a linear extension of a poset  $P$ .

Given  $\pi = a_1 a_2 \cdots a_n \in \mathcal{L}_{P,\omega}$ , let  $\text{inv}(f)$  denote the number of *inversions* of  $\pi$ , i.e.,

$$\text{inv}(\pi) = \#\{(i, j) : i < j, a_i > a_j\}.$$

Let

$$I_{P,\omega}(q) = \sum_{\pi \in \mathcal{L}_{P,\omega}} q^{\text{inv}(f)}, \quad (1)$$

the generating function for linear extensions of  $(P, \omega)$  by number of inversions. Since  $f$  is an even linear extension if and only if  $\text{inv}(f)$  is an even integer, we see that  $P$  is sign-balanced if and only if  $I_{P,\omega}(-1) = 0$ . In general  $I_{P,\omega}(q)$  seems difficult to understand, even when  $P$  is known to be sign-balanced.

## 2 Promotion and evacuation.

Promotion and evacuation are certain bijections on the set  $\mathcal{E}_P$  of linear extensions of a finite poset  $P$ . They were originally defined by M.-P. Schützenberger [18] and have subsequently arisen in many different situations (e.g., [5, §5][8, §8][9, §4][12, §3]). To be precise, the original definitions of promotion and evacuation require an insignificant reindexing to become bijections. We will incorporate this reindexing into our definition. Let  $f : P \rightarrow [n]$  be a linear extension of the poset  $P$ . Define a maximal chain  $u_0 < u_1 < \cdots < u_\ell$  of  $P$ , called the *promotion chain* of  $f$ , as follows. Let  $u_0 = f^{-1}(1)$ . Once  $u_i$  is defined let  $u_{i+1}$  be that element  $u$  covering  $u_i$  (i.e.,  $u_i < u_{i+1}$  and no  $s \in P$  satisfies  $u_i < s < u_{i+1}$ ) for which  $f(u)$  is minimal. Continue until reaching a maximal element  $u_\ell$  of  $P$ . Now define the *promotion*  $g = \partial f$  of  $f$  as follows. If  $t \neq u_i$  for any  $i$ , then set  $g(t) = f(t) - 1$ . If  $1 \leq i \leq \ell - 1$ , then set  $g(u_i) = f(u_{i+1}) - 1$ . Finally set  $g(u_\ell) = n$ . Figure 1 gives an example, with the elements in the promotion chain of  $f$  circled. (The vertex labels in Figure 1 are the values of a linear extension and are unrelated to the (irrelevant) labeling  $\omega$ .) It is easy to see that  $\partial f \in \mathcal{E}_P$  and that the map  $\partial : \mathcal{E}_P \rightarrow \mathcal{E}_P$  is a bijection.




Figure 1: The promotion operator  $\partial$

**2.1 Lemma.** *Let  $P$  be an  $n$ -element poset. Then the promotion operator  $\partial : \mathcal{E}_P \rightarrow \mathcal{E}_P$  is parity-reversing if and only if the length  $\ell$  (or cardinality  $\ell+1$ ) of every maximal chain of  $P$  satisfies  $n \equiv \ell \pmod{2}$ . Similarly,  $\partial$  is parity-preserving if and only if the length  $\ell$  of every maximal chain of  $P$  satisfies  $n \equiv \ell + 1 \pmod{2}$ .*

**Proof.** Let  $f \in \mathcal{E}_P$ , and let  $u_0 < u_1 < \dots < u_\ell$  be the promotion chain of  $f$ . Then  $(\partial f)f^{-1}$  is a product of two cycles, viz.,

$$(\partial f)f^{-1} = (n, n-1, \dots, 1)(b_0, b_1, \dots, b_\ell),$$

where  $b_i = f(u_i)$ . This permutation is odd if and only if  $n \equiv \ell \pmod{2}$ , and the proof follows since every maximal chain of  $P$  is the promotion chain of some linear extension.  $\square$

**2.2 Corollary.** *Let  $P$  be an  $n$ -element poset, and suppose that the length  $\ell$  of every maximal chain of  $P$  satisfies  $n \equiv \ell \pmod{2}$ . Then  $P$  is sign-balanced.*

**Proof.** By the previous lemma,  $\partial$  is parity-reversing. Since it is also a bijection,  $\mathcal{E}_P$  must contain the same number of even linear extensions as odd linear extensions.  $\square$




Figure 2: The evacuation operator  $\text{evac}$ .

We now consider a variant of promotion known as evacuation. For any linear extension  $g$  of an  $m$ -element poset  $Q$ , let  $u_0 < u_1 < \dots < u_\ell$  be the promotion chain of  $g$ , so  $\partial g(u_\ell) = m$ . Define  $\rho_g(Q) = Q - \{u_\ell\}$ . The restriction of  $\partial g$  to  $\rho_g(Q)$ , which we also denote by  $\partial g$ , is a linear extension of  $\rho_g(Q)$ . Let

$$\mu_{g,k}(Q) = \rho_{\partial^k g} \rho_{\partial^{k-1} g} \cdots \rho_{\partial g} \rho_g(Q).$$

Now let  $\#P = n$  and define the *evacuation*  $\text{evac}(f)$  of  $f$  to be the linear extension of  $P$  whose value at the unique element of  $\mu_{g,k-1}(P) - \mu_{g,k}(P)$  is  $n - k + 1$ , for  $1 \leq k \leq n$ . Figure 2 gives an example of  $\text{evac}(f)$ , where we circle the values of  $\text{evac}(f)$  as soon as they are determined. A remarkable theorem of Schützenberger [18] asserts that  $\text{evac}$  is an involution (and hence a bijection  $\mathcal{E}_P \rightarrow \mathcal{E}_P$ ).

We say that the poset  $P$  is *consistent* if for all  $t \in P$ , the lengths of all maximal chains of the principal order ideal  $\Lambda_t := \{s \in P : s \leq t\}$  have the same parity. Let  $\nu(t)$  denote the length of the longest chain of  $\Lambda_t$ , and set

$$\Gamma(P) = \sum_{t \in P} \nu(t).$$

We also say that a permutation  $\sigma$  of a finite set has *parity*  $k \in \mathbb{Z}$  if either  $\sigma$  and  $k$  are both even or  $\sigma$  and  $k$  are both odd. Equivalently,  $\text{inv}(\sigma) \equiv k \pmod{2}$ .

**2.3 Proposition.** *Suppose that  $P$  is consistent. Then  $\text{evac}: \mathcal{E}_P \rightarrow \mathcal{E}_P$  is parity-preserving if  $\binom{n}{2} - \Gamma(P)$  is even, and parity-reversing if  $\binom{n}{2} - \Gamma(P)$  is odd.*

**Proof.** The evacuation of a linear extension  $f$  of an  $n$ -element poset  $P$  consists of  $n$  promotions  $\delta_1, \dots, \delta_n$ , where  $\delta_i$  is applied to a certain subposet

$P_{i-1}$  of  $P$  with  $n - i + 1$  elements. Let  $f_i$  be the linear extension of  $P$  whose restriction to  $P_i$  agrees with  $\delta_i \delta_{i-1} \cdots \delta_1$ , and whose value at the unique element of  $P_{j-1} - P_j$  for  $j \leq i$  is  $n - i + 1$ . Thus  $f_0 = f$  and  $f_n = \text{evac}(f)$ . (Figure 2 gives an example of the sequence  $f_0, \dots, f_5$ .) Let  $u_i$  be the end (top) of the promotion chain for the promotion  $\delta_i$ . Thus  $\{u_1, u_2, \dots, u_n\} = P$ . Lemma 2.1 shows that if  $P$  is consistent, then  $f_i f_{i-1}^{-1}$  has parity  $n - i + 1 - (\nu(u_i) + 1)$ . Hence the parity of  $\text{evac}(f) f^{-1}$  is given by

$$\sum_{i=1}^n (n - i - \nu(u_i)) = \binom{n}{2} - \sum_{t \in P} \nu(P) = \binom{n}{2} - \Gamma(P),$$

from which the proof follows.  $\square$

**2.4 Corollary.** *Suppose that  $P$  is consistent and  $\binom{n}{2} - \Gamma(P)$  is odd. Then  $P$  is sign-balanced.*

NOTE. In [20, pp. 50–51][21, Cor. 19.5] it was shown using the theory of  $P$ -partitions that the number  $e(P)$  of linear extensions of  $P$  is even if  $P$  is graded of rank  $\ell$  (i.e., every maximal chain of  $P$  has length  $\ell$ ) and  $n - \ell$  is even, and it was stated that it would be interesting to give a direct proof. Our Corollary 2.2 gives a direct proof of a stronger result. Similarly in [20, Cor. 4.6][21, Cor. 19.6] it was stated (in dual form) that if for all  $t \in P$  all maximal chains of  $\Lambda_t$  have the same length, and if  $\binom{n}{2} - \Gamma(P)$  is odd, then  $e(P)$  is even. Corollary 2.4 gives a direct proof of a stronger result.

### 3 Partitions.

In this section we apply our previous results and obtain some new results for certain posets corresponding to (integer) partitions. We first review some notation and terminology concerning partitions. Further details may be found in [24, Ch. 7]. Let  $\lambda = (\lambda_1, \lambda_2, \dots)$  be a partition of  $n$ , denoted  $\lambda \vdash n$  or  $|\lambda| = n$ . Thus  $\lambda_1 \geq \lambda_2 \geq \cdots \geq 0$  and  $\sum \lambda_i = n$ . We can identify  $\lambda$  with its *diagram*  $\{(i, j) \in \mathbb{P} \times \mathbb{P} : 1 \leq j \leq \lambda_i\}$ . Let  $\mu$  be another partition such that  $\mu \subseteq \lambda$ , i.e.,  $\mu_i \leq \lambda_i$  for all  $i$ . Define the *skew partition* or *skew diagram*  $\lambda/\mu$  by

$$\lambda/\mu = \{(i, j) \in \mathbb{P} \times \mathbb{P} : \mu_i + 1 \leq j \leq \lambda_i\}.$$

Write  $|\lambda/\mu| = n$  to denote that  $|\lambda| - |\mu| = n$ , i.e.,  $n$  is the number of squares in the shape  $\lambda/\mu$ , drawn as a Young diagram [22, p. 29]. We can regard  $\lambda/\mu$  as a subposet of  $\mathbb{P} \times \mathbb{P}$  (with the usual coordinatewise ordering). We write  $P_{\lambda/\mu}$  for this poset. As a set it is the same as  $\lambda/\mu$ , but the notation  $P_{\lambda/\mu}$  emphasizes that we are considering it to be a poset. In this section we will only be concerned with “ordinary” shapes  $\lambda$ , but in Section 5 skew shapes  $\lambda/\mu$  will arise as a special case of Proposition 5.3.

The posets  $P_\lambda$  are consistent for any  $\lambda$ , so we can ask for which  $P_\lambda$  is evacuation parity-reversing, i.e.,  $\binom{n}{2} - \Gamma(P_\lambda)$  is odd. To this end, the *content*  $c(i, j)$  of the cell  $(i, j)$  is defined by  $c(i, j) = j - i$  [24, p. 373]. Also let  $\mathcal{O}(\mu)$  denote the number of odd parts of the partition  $\mu$ . An *order ideal* of a poset  $P$  is a subset  $K \subseteq P$  such that if  $t \in K$  and  $s < t$ , then  $s \in K$ . Similarly a *dual order ideal* or *filter* of  $P$  is a subset  $F \subseteq P$  such that if  $s \in F$  and  $t > s$ , then  $t \in F$ . If we successively remove two-element chains from  $P_\lambda$  which are dual order ideals of the poset from which they are removed, then eventually we reach a poset  $\text{core}_2(P_\lambda)$ , called the *2-core* of  $P_\lambda$ , that contains no dual order ideals which are two-element chains. The 2-core is *unique*, i.e., independent of the order in which the dual order ideals are removed, and is given by  $P_{\delta_k}$  for some  $k \geq 1$ , where  $\delta_k$  denotes the “staircase shape”  $(k-1, k-2, \dots, 1)$ . For further information see [24, Exer. 7.59].

**3.1 Proposition.** Let  $\lambda \vdash n$ . The following numbers all have the same parity.

- (a)  $\Gamma(P_\lambda)$
- (b)  $\sum_{t \in P_\lambda} c(t)$
- (c)  $\frac{1}{2}(\mathcal{O}(\lambda) - \mathcal{O}(\lambda'))$
- (d)  $\frac{1}{2}(n - \binom{k}{2})$ , where  $\binom{k}{2} = \#\text{core}_2(P_\lambda)$

Hence if  $a_\lambda$  denotes any of the above four numbers, then evacuation is parity-reversing on  $P_\lambda$  if and only if  $\binom{n}{2} - a_\lambda$  is odd.

**Proof.** It is easy to see that if  $t \in P_\lambda$ , then  $\nu(t) \equiv c(t) \pmod{2}$ . Hence (a) and (b) have the same parity. It is well-known and easy to see [13, Exam.

3, p. 11] that

$$\sum_{t \in P_\lambda} c(t) = \sum \binom{\lambda_i}{2} - \sum \binom{\lambda'_i}{2}.$$

Since  $\sum \lambda_i = \sum \lambda'_i$ , we have

$$\sum_{t \in P_\lambda} c(t) = \frac{1}{2} \left( \sum \lambda_i^2 - \sum (\lambda'_i)^2 \right).$$

Since  $a^2 \equiv 0, 1 \pmod{4}$  depending on whether  $a$  is even or odd, we see that (b) and (c) have the same parity. If we remove from  $P_\lambda$  a 2-element dual order ideal which is also a chain, then we remove exactly one element with an odd content. A 2-core is self-conjugate and hence has an even content sum. Hence the number of odd contents of  $P_\lambda$  is equal to the number of dominos that must be removed from  $P_\lambda$  in order to reach  $\text{core}_2(P_\lambda)$ . It follows that (b) and (c) have the same parity, completing the proof.  $\square$

It can be shown [25] that if  $t(n)$  denotes the number of partitions  $\lambda \vdash n$  for which  $a_\lambda$  is even, then  $t(n) = \frac{1}{2}(p(n) + f(n))$ , where  $p(n)$  denotes the total number of partitions of  $n$  and

$$\sum_{n \geq 0} f(n)x^n = \prod_{i \geq 1} \frac{1 + x^{2i-1}}{(1 - x^{4i})(1 + x^{4i-2})^2}.$$

Hence the number  $g(n)$  of partitions  $\lambda \vdash n$  for which  $\text{evac}$  is parity-reversing on  $P_\lambda$  is given by

$$g(n) = \begin{cases} \frac{1}{2}(p(n) + f(n)), & \text{if } \binom{n}{2} \text{ is odd} \\ \frac{1}{2}(p(n) - f(n)), & \text{if } \binom{n}{2} \text{ is even} \end{cases}$$

We conclude this section with some applications of the theory of domino tableaux. A *standard domino tableau* (SDT) of shape  $\lambda \vdash 2n$  is a sequence

$$\emptyset = \lambda^0 \subset \lambda^1 \subset \cdots \subset \lambda^n = \lambda$$

of partitions such that each skew shape  $\lambda^i / \lambda^{i-1}$  is a *domino*, i.e., two squares with an edge in common. Each of these dominos is either horizontal (two squares in the same row) or vertical (two squares in the same column). Let



$\text{Dom}_\lambda$  denote the set of all SDT of shape  $\lambda$ . Given  $D \in \text{Dom}_\lambda$ , define  $\text{ev}(D)$  to be the number of vertical dominos in even columns of  $D$ , where an *even column* means the  $2i$ th column for some  $i \in \mathbb{P}$ . For the remainder of this section, fix the labeling  $\omega$  of  $P_\lambda$  to be the usual “reading order,” i.e., the first row of  $\lambda$  is labelled  $1, 2, \dots, \lambda_1$ ; the second row is labelled  $\lambda_1 + 1, \lambda_1 + 2, \dots, \lambda_1 + \lambda_2$ , etc. We write  $I_\lambda(q)$  for  $I_{P_\lambda, \omega}(q)$  and set  $I_\lambda = I_\lambda(-1)$ , the *imbalance* of the partition  $\lambda$ . It is shown in [27, Thm. 12] (by analyzing the formula that results from setting  $q = -1$  in (9)) that

$$I_\lambda = \sum_{D \in \text{Dom}_\lambda} (-1)^{\text{ev}(D)}.$$

Let  $\lambda \vdash n$ . Lascoux, Leclerc and Thibon [11, (27)] define a certain class of symmetric functions  $\tilde{G}_\lambda^{(k)}$  (defined earlier by Carré and Leclerc [3] for the special case  $k = 2$  and  $\lambda = 2\mu$ ). We will only be concerned with the case  $k = 2$ , for which we write  $G_\lambda = \tilde{G}_\lambda^{(2)}$ . The symmetric function  $G_\lambda$  is homogeneous of degree  $n$ . We will not define it here but only recall the properties relevant to us. The connection with the imbalance  $I_\lambda$  is provided by the formula (immediate from the definition of  $G_\lambda$  in [11] together with [27, Thm. 12])

$$[x_1 \cdots x_n] G_\lambda = (-1)^{d(\lambda)} I_\lambda, \quad (2)$$

where  $[x_1 \cdots x_n] F$  denotes the coefficient of  $x_1 \cdots x_n$  in the symmetric function  $F$ , and

$$d(\lambda) = \sum_{d \geq 1} \left\lfloor \frac{1}{2} \lambda'_{2d} \right\rfloor.$$

If there exists an SDT of shape  $\lambda$ , then  $d(\lambda)$  is the maximum value of  $\text{ev}(D)$  among all such SDT.

**3.2 Theorem.** (a) *We have*

$$\sum_{\mu \vdash n} I_{2\mu} = 1$$

for all  $n \geq 1$ .

(b) *Let  $v(\lambda)$  denote the maximum number of disjoint vertical dominos that fit in the shape  $\lambda$ . Equivalently,*

$$v(\lambda) = \sum_{i \geq 1} \left\lfloor \frac{1}{2} \lambda'_i \right\rfloor.$$

Then

$$\sum_{\lambda \vdash 2n} (-1)^{v(\lambda)} I_\lambda^2 = 0.$$

**Proof.** (a) Carré and Leclerc [3, Def. 9.1] define a symmetric function  $H_\mu(x; q)$  which satisfies  $H_\mu(x, -1) = G_{2\mu}$ . In [10, Thm. 1] they prove the identity

$$\sum_{\mu} H_\mu(x; q) = \prod_i \frac{1}{1 - x_i} \prod_{i < j} \frac{1}{1 - x_i x_j} \prod_{i \geq j} \frac{1}{1 - q x_i x_j}.$$

Setting  $q = -1$  gives

$$\sum_{\mu} G_{2\mu} = \prod_i \frac{1}{(1 - x_i)(1 + x_i^2)} \prod_{i < j} \frac{1}{1 - x_i^2 x_j^2}.$$

Taking the coefficient of  $x_1 \cdots x_n$  on both sides and using (2) completes the proof.

(b) Shimozono and White [19, Thm. 30] define a bijection between elements  $\pi$  of the hyperoctahedral group  $B_n$ , regarded as signed permutations of  $1, 2, \dots, n$ , and pairs  $(P, Q)$  of SDT of the same shape  $\lambda \vdash 2n$ . In this bijection,

$$\text{tc}(\pi) = \frac{1}{2}(v(P) + v(Q)),$$

where  $\text{tc}(\pi)$  denotes the number of minus signs in  $\pi$  and  $v(R)$  denotes the number of vertical dominos in the SDT  $R$ . It is easy to see that for any SDT  $D$  we have

$$v(D) = v(\lambda) - 2d(\lambda) + 2\text{ev}(D).$$

Thus

$$\begin{aligned} 0 &= \sum_{\pi \in B_n} (-1)^{\text{tc}(\pi)} \\ &= \sum_{P, Q} (-1)^{\frac{1}{2}(v(P) + v(Q))} \\ &= \sum_{\lambda \vdash 2n} \left( \sum_{D \in \text{Dom}_\lambda} (-1)^{\frac{1}{2}v(D)} \right)^2 \end{aligned}$$

$$\begin{aligned}
&= \sum_{\lambda \vdash 2n} (-1)^{v(\lambda)} \left( \sum_{D \in \text{Dom}_\lambda} (-1)^{d(D)} \right)^2 \\
&= \sum_{\lambda \vdash 2n} (-1)^{v(\lambda)} I_\lambda^2. \quad \square
\end{aligned}$$

In the same spirit as Theorem 3.2 we have the following conjecture, which has been verified for  $n \leq 20$ .

**3.3 Conjecture.** For all  $n \geq 0$  we have

$$\begin{aligned}
\sum_{\lambda \vdash n} q^{v(\lambda)} t^{d(\lambda)} I_\lambda &= (1+q)^{\lfloor n/2 \rfloor} \\
\sum_{\lambda \vdash n} (-1)^{v(\lambda)} t^{d(\lambda)} I_\lambda^2 &= 0 \\
\sum_{\mu \vdash n} q^{v(\mu)} t^{d(\mu)} I_{2\mu} &= 1 \\
\sum_{\mu \vdash n} q^{v(\mu)} t^{d(\mu)} I_{\mu \cup \mu} &= 1.
\end{aligned} \tag{3}$$

The case  $t = 0$  of equation (3) follows from the following proposition, which in a sense “explains” where the right-hand side  $(1+q)^{\lfloor n/2 \rfloor}$  comes from.

**3.4 Proposition.** For all  $n \geq 0$  we have

$$\sum_{\lambda = \langle n-k, 1^k \rangle} q^{v(\lambda)} I_\lambda = (1+q)^{\lfloor n/2 \rfloor}, \tag{4}$$

where  $\lambda$  ranges over all “hook shapes”  $\langle n-k, 1^k \rangle$ ,  $0 \leq k \leq n-1$ . (Note that  $d(\lambda) = 0$  if and only if  $\lambda$  is a hook.)

**First proof.** Let  $\lambda = \langle n-k, 1^k \rangle$ . Let  $\omega$  denote the “reading order” labeling of  $P_\lambda$  as above. The set  $\mathcal{L}_{P,\omega}$  consists of all permutations  $1, a_2, \dots, a_n$ , where  $a_2, \dots, a_n$  is a *shuffle* of the permutations  $2, 3, \dots, n-k$  and  $n-k+1, n-k+2, \dots, n$ . It follows e.g. from [22, Prop. 1.3.17] that

$$I_\lambda(q) = \begin{bmatrix} n-1 \\ k \end{bmatrix},$$

a  $q$ -binomial coefficient.

Suppose first that  $n = 2m + 1$ . By [22, Exer. 3.45(b)],

$$(1+x)(1+qx)\cdots(1+q^{n-2}x) = \sum_{k=0}^{n-1} q^{\binom{k}{2}} \begin{bmatrix} n-1 \\ k \end{bmatrix} x^k.$$

Setting  $q = -1$  gives

$$\begin{bmatrix} n-1 \\ k \end{bmatrix}_{q=-1} = \begin{cases} \binom{m}{j}, & k = 2j \\ 0, & k = 2j+1. \end{cases}$$

Note that if  $\lambda = \langle n-2j, 1^{2j} \rangle$ , then  $v(\lambda) = j$ . Hence

$$\begin{aligned} \sum_{\lambda=\langle n-k, 1^k \rangle} q^{v(\lambda)} I_\lambda &= \sum_{j=0}^m q^j \binom{m}{j} \\ &= (1+q)^m, \end{aligned}$$

as desired. The proof for  $n$  even is similar and will be omitted.  $\square$

**Second proof.** Assume first that  $n = 2m$ . We use an involution argument analogous to the proof of Theorem 1.1 or to arguments in [27, §5] and Section 4 of this paper. Let  $T$  be an SYT of shape  $\lambda = \langle n-k, 1^k \rangle$ , which can be regarded as an element of  $\mathcal{L}_{P_\lambda, \omega}$ . Let  $i$  be the least positive integer (if it exists) such that  $2i-1$  and  $2i$  appear in different rows and in different columns of  $T$ . Let  $T'$  denote the SYT obtained from  $T$  by transposing  $2i-1$  and  $2i$ . Since multiplying by a transposition changes the sign of a permutation, we have  $(-1)^{\text{inv}(T)} + (-1)^{\text{inv}(T')} = 0$ . The surviving SYT are obtained by first placing  $1, 2$  in the same row or column, then  $3, 4$  in the same row or column, etc. If  $k = 2j$  or  $2j+1$ , then the number of survivors is easily seen to be  $\binom{m-1}{j}$ . Because the entries of  $T$  come in pairs  $2i-1, 2i$ , the number of inversions of each surviving SYT is even. Moreover, if  $k = 2j$  then  $v(\lambda) = j$ , while if  $k = 2j+1$  then  $v(\lambda) = j+1$ . Hence

$$\begin{aligned} \sum_{\lambda=\langle n-k, 1^k \rangle} q^{v(\lambda)} I_\lambda &= \sum_{j=0}^{m-1} (1+q) \binom{m-1}{j} q^j \\ &= (1+q)^m, \end{aligned}$$

as desired.

The proof is similar for  $n = 2m + 1$ . Let  $i$  be the least positive integer (if it exists) such that  $2i$  and  $2i + 1$  (rather than  $2i - 1$  and  $2i$ ) appear in different rows and in different columns of  $T$ . There are now no survivors when  $k = 2j + 1$  and  $\binom{m}{j}$  survivors when  $k = 2j$ . Other details of the proof remain the same, so we get

$$\begin{aligned} \sum_{\lambda=\langle n-k, 1^k \rangle} q^{v(\lambda)} I_\lambda &= \sum_{j=0}^m \binom{m-1}{j} q^j \\ &= (1+q)^m, \end{aligned}$$

completing the proof.  $\square$

There are some additional properties of the symmetric functions  $G_\lambda$  that yield information about  $I_\lambda$ . For instance, there is a product formula in [10, Thm. 2] for  $\sum_\mu G_{2\mu \cup 2\mu}$ , where  $\mu$  ranges over all partitions and

$$2\mu \cup 2\mu = (2\mu_1, 2\mu_1, 2\mu_2, 2\mu_2, \dots),$$

which implies that  $\sum_{\mu \vdash n} I_{2\mu \cup 2\mu} = 0$ . In fact, in [3, Cor. 9.2] it is shown that  $G_{2\mu \cup 2\mu}(x) = \pm s_\mu(x_1^2, x_2^2, \dots)$ , from which it follows easily that in fact  $I_{2\mu \cup 2\mu} = 0$ . However, this result is just a special case of Corollary 2.2 and of Proposition 2.3, so we obtain nothing new.

Also relevant to us is an expansion of  $G_\lambda$  into Schur functions due to Shimozono (see [27, Thm. 18]) for certain shapes  $\lambda$ , namely, those whose 2-quotient (in the sense e.g. of [13, Exam. I.1.8]) is a pair of rectangles. This expansion was used by White [27, Cor. 20] to evaluate  $I_\lambda$  for such shapes. White [27, §8] also gives a combinatorial proof, based on a sign-reversing involution, in the special case that  $\lambda$  itself is a rectangle. We simply state here White's result for rectangles.

**3.5 Theorem.** *Let  $\lambda$  be an  $m \times n$  rectangle. Then*

$$I_\lambda = \begin{cases} 1, & \text{if } m = 1 \text{ or } n = 1 \\ 0, & \text{if } m \equiv n \pmod{2} \text{ and } m, n > 1 \\ g^\mu, & m \not\equiv n \pmod{2}, \end{cases}$$

where  $g^\mu$  denotes the number of shifted standard tableaux (as defined e.g. in [13, Exam. III.8.12]) of shape

$$\mu = \left( \frac{m+n-1}{2}, \frac{m+n-3}{2}, \dots, \frac{|n-m|+3}{2}, \frac{|n-m|+2}{2} \right).$$

(An explicit “hook length formula” for any  $g^\mu$  appears e.g. in the reference just cited.)

It seems unlikely that a “nice” formula exists for  $I_\lambda(q)$ , or even  $I_\lambda(-1)$ , for an arbitrary partition  $\lambda$ . We can nevertheless give some general properties of  $I_\lambda(q)$ . Let  $C(\lambda)$  denote the set of *corner squares* of  $\lambda$ , i.e., those squares of the Young diagram of  $\lambda$  whose removal still gives a Young diagram. Equivalently, Pieri’s formula [24, Thm. 7.15.7] implies that

$$s_{\lambda/1} = \sum_{t \in C(\lambda)} s_{\lambda-t}. \quad (5)$$

Let  $f^\lambda$  denote the number of SYT of shape  $\lambda$  [24, Prop. 7.10.3], so

$$f^\lambda = \sum_{t \in C(\lambda)} f^{\lambda-t}. \quad (6)$$

Note that  $I_\lambda(1) = f^\lambda$ , so  $I_\lambda(q)$  is a (nonstandard)  $q$ -analogue of  $f^\lambda$ . The  $q$ -analogue of equation (6) is the following result.

**3.6 Proposition.** *We have*

$$I_\lambda(q) = \sum_{t \in C(\lambda)} q^{b_\lambda(t)} I_{\lambda-t}(q),$$

where  $b_\lambda(t)$  denotes the number of squares in the diagram of  $\lambda$  in a lower row than that of  $t$ .

**Proof.** We have by definition

$$I_\lambda(q) = \sum_T q^{\text{inv}(\pi(T))},$$

where  $T$  ranges over all SYT of shape  $\lambda$  and  $\pi(T)$  is the permutation obtained by reading the entries of  $T$  in the usual reading order, i.e., left-to-right and

top-to-bottom when  $T$  is written in “English notation” as in [13][22][24]. Suppose  $\lambda \vdash n$ . If  $T$  is an SYT of shape  $\lambda$ , then the square  $t$  occupied by  $n$  is a corner square. The number of inversions  $(i, j)$  of  $\pi(T) = a_1 \cdots a_n$  such that  $a_i = n$  is equal to  $b_\lambda(t)$ , and the proof follows.  $\square$

Now let  $D_1$  denote the linear operator on symmetric functions defined by  $D_1(s_\lambda) = s_{\lambda/1}$ . We then have the commutation relation [24, Exercise 7.24(a)]

$$D_1 s_1 - s_1 D_1 = I, \quad (7)$$

the identity operator. This leads to many enumerative consequences, discussed in [23]. There is an analogue of (7) related to  $I_\lambda$ , though we don’t know of any applications. Define a linear operator  $D(q)$  on symmetric functions by

$$D(q)s_\lambda = \sum_{t \in C(\lambda)} q^{b_\lambda(t)} s_{\lambda-t}.$$

Let  $U(q)$  denote the adjoint of  $D(q)$  with respect to the basis  $\{s_\lambda\}$  of Schur functions, so

$$U(q)s_\mu = \sum_t q^{b_{\mu+t}(t)} s_{\mu+t},$$

where  $t$  ranges over all boxes that we can add to the diagram of  $\mu$  to get the diagram of a partition  $\mu + t$  (for which necessarily  $t \in C(\mu + t)$ ). Note that  $U(1) = s_1$  (i.e., multiplication by  $s_1$ ) and  $D(1) = D_1$  as defined above. It follows from Proposition 3.6 that

$$U(q)^n \cdot 1 = \sum_{\lambda \vdash n} I_\lambda(q) s_\lambda,$$

where  $U(q)^n \cdot 1$  denotes  $U(q)^n$  acting on the symmetric function  $1 = s_\emptyset$ . Write  $U = U(-1)$  and  $D = D(-1)$ . Let  $A$  be the linear operator on symmetric functions given by  $As_\lambda = (2k(\lambda) + 1)s_\lambda$ , where  $k(\lambda) = \#C(\lambda)$ , the number of corner boxes of  $\lambda$ .

**3.7 Proposition.** *We have  $DU + UD = A$ .*

**Proof.** The proof is basically a brute force computation. Write  $\bar{\lambda}_i = \lambda_i + \lambda_{i+1} + \cdots$ . Suppose  $\mu$  is obtained from  $\lambda$  by adding a box in row  $r - 1$

and deleting a box in row  $s - 1$ , where  $r < s$ . Then the coefficient of  $s_\mu$  in  $(D(q)U(q) + U(q)D(q))s_\lambda$  is given by

$$\langle s_\mu, (D(q)U(q) + U(q)D(q))s_\lambda \rangle = q^{\bar{\lambda}_r} q^{\bar{\lambda}_s} + q^{\bar{\lambda}_s} q^{\bar{\lambda}_r - 1},$$

which vanishes when  $q = -1$ . Similarly if  $r > s$  we get

$$\langle s_\mu, (D(q)U(q) + U(q)D(q))s_\lambda \rangle = q^{\bar{\lambda}_s} q^{\bar{\lambda}_r + 1} + q^{\bar{\lambda}_r} q^{\bar{\lambda}_s},$$

which again vanishes when  $q = -1$ . On the other hand, if  $\lambda = \mu$  we have

$$\begin{aligned} \langle s_\lambda, (D(q)U(q) + U(q)D(q))s_\lambda \rangle &= (c(\lambda) + 1)q^{2\bar{\lambda}_r} + c(\lambda)q^{2\bar{\lambda}_r} \\ &= (2c(\lambda) + 1)q^{2\bar{\lambda}_r}. \end{aligned}$$

When  $q = -1$  the right-hand side become  $2c(\lambda) + 1$ , completing the proof.  $\square$

## 4 Chains of order ideals.

Suppose that  $P$  is an  $n$ -element poset, and let  $\alpha = (\alpha_1, \dots, \alpha_k)$  be a composition of  $n$ , i.e.,  $\alpha_i \in \mathbb{P} = \{1, 2, \dots\}$  and  $\sum \alpha_i = n$ . Define an  $\alpha$ -chain of order ideals of  $P$  to be a chain

$$\emptyset = K_0 \subset K_1 \subset \dots \subset K_k = P \tag{8}$$

of order ideals satisfying  $\#(K_i - K_{i-1}) = \alpha_i$  for  $1 \leq i \leq k$ . The following result is quite simple but has a number of consequences.

**4.1 Proposition.** *Let  $P$  be an  $n$ -element poset and  $\alpha$  a fixed composition of  $n$ . Suppose that for every  $\alpha$ -chain (8) of order ideals of  $P$ , at least one subposet  $K_i - K_{i-1}$  is sign-balanced. Then  $P$  is sign-balanced.*

**Proof.** Let  $\mathcal{C}$  be the  $\alpha$ -chain (8). We say that a linear extension  $f$  is  $\mathcal{C}$ -compatible if

$$K_1 = f^{-1}(\{1, \dots, \alpha_1\}), \quad K_2 - K_1 = f^{-1}(\{\alpha_1 + 1, \dots, \alpha_1 + \alpha_2\}),$$



etc. Let  $\text{inv}(\mathcal{C})$  be the minimum number of inversions of a  $\mathcal{C}$ -compatible linear extension. Clearly

$$\sum_f q^{\text{inv}(f)} = q^{\text{inv}(\mathcal{C})} \prod_{i=1}^k I_{K_i - K_{i-1}}(q),$$

where the sum is over all  $\mathcal{C}$ -compatible  $f$ . Since every linear extension is compatible with a unique  $\alpha$ -chain, there follows

$$I_{P,\omega}(q) = \sum_{\mathcal{C}} q^{\text{inv}(\mathcal{C})} \prod_{i=1}^k I_{K_i - K_{i-1}}(q), \quad (9)$$

where  $\mathcal{C}$  ranges over all  $\alpha$ -chains of order ideals of  $P$ . The proof now follows by setting  $q = -1$ .  $\square$

Define a finite poset  $P$  with  $2m$  elements to be *tilable by dominos* if there is a chain  $\emptyset = K_0 \subset K_1 \subset \cdots \subset K_m = P$  of order ideals such that each subposet  $K_i - K_{i-1}$  is a two-element chain. Similarly, if  $\#P = 2m + 1$  and  $1 \leq j \leq m + 1$  then we say that  $P$  is  *$j$ -tilable by dominos* if there is a chain  $\emptyset = K_0 \subset K_1 \subset \cdots \subset K_{m+1} = P$  of order ideals such that  $\#(K_i - K_{i-1}) = 2$  if  $1 \leq i \leq m + 1$  and  $i \neq j$  (so  $\#(K_j - K_{j-1}) = 1$ ). Note that being tilable by dominos is stronger than the existence of a partition of  $P$  into cover relations (or two element saturated chains). For instance, the poset  $P$  with cover relations  $a < c, b < c, a < d, b < d$  can be partitioned into the two cover relations  $a < c$  and  $b < d$ , but  $P$  is not tilable by dominos. When  $n = 2m$ , we define a  *$P$ -domino tableau* to be a chain  $\emptyset = K_0 \subset K_1 \subset \cdots \subset K_m = P$  of order ideals such that  $K_i - K_{i-1}$  is a two-element chain for  $1 \leq i \leq m$ . Similarly, when  $n = 2m + 1$ , we define a (standard)  *$P$ -domino tableau* to be a chain  $\emptyset = K_0 \subset K_1 \subset \cdots \subset K_{m+1} = P$  of order ideals such that  $K_i - K_{i-1}$  is a two-element chain for  $1 \leq i \leq m$  (so that  $K_{m+1} - K_m$  consists of a single point). Thus for  $\lambda \vdash 2n$ , a  $P_\lambda$ -domino tableau coincides with our earlier definition of an SDT of shape  $\lambda$ .

**4.2 Corollary.** Let  $\#P = 2m$ , and assume that  $P$  is not tilable by dominos. Then  $P$  is sign-balanced. Similarly if  $\#P = 2m + 1 \geq 3$  and  $P$  is not  $j$ -tilable by dominos for some  $j$ , then  $P$  is sign-balanced.

**Proof.** Let  $\alpha = (2, 2, \dots, 2)$  ( $m$  2's). If  $\#P = 2m$  and  $P$  is not tilable by dominos, then for any  $\alpha$ -chain (8) there is an  $i$  for which  $K_i - K_{i-1}$

consists of two disjoint points. Since a poset consisting of two disjoint points is sign-balanced, it follows from Proposition 4.1 that  $P$  is sign-balanced. The argument is similar for  $\#P = 2m + 1$ .  $\square$

Corollary 4.2 was proved in a special case (the product of two chains with an even number of elements, with the  $\hat{0}$  and  $\hat{1}$  removed), using essentially the same proof as we have given, by Ruskey [17, §5, item 6].

Corollary 4.2 is particularly useful for the posets  $P_\lambda$ . From this corollary and the definition of  $\text{core}_2(\lambda)$  we conclude the following.

**4.3 Corollary.** *If  $\text{core}_2(P_\lambda)$  consists of more than one element, then  $P_\lambda$  is sign-balanced.*

It follows from [24, Exer. 7.59(e)] that if  $f(n)$  denotes the number of partitions  $\lambda \vdash n$  such that  $\#\text{core}_2(\lambda) \leq 1$ , then

$$\sum_{n \geq 0} f(n)x^n = \frac{1+x}{\prod_{i \geq 1} (1-x^{2i})^2}.$$

Standard partition asymptotics (e.g., [1, Thm. 6.2]) shows that

$$f(n) \sim \frac{C}{n^{5/4}} \exp\left(\pi\sqrt{2n/3}\right)$$

for some  $C > 0$ . Since the total number  $p(n)$  of partitions of  $n$  satisfies

$$p(n) \sim \frac{C'}{n} \exp\left(\pi\sqrt{2n/3}\right),$$

it follows that  $\lim_{n \geq 0} f(n)/p(n) = 0$ . Hence as  $n \rightarrow \infty$ ,  $P_\lambda$  is sign-balanced for almost all  $\lambda \vdash n$ .

## 5 Maj-balanced posets.

If  $\pi = a_1 a_2 \cdots a_n$  is a permutation of  $[n]$ , then the *descent set*  $D(\pi)$  of  $\pi$  is defined as

$$D(\pi) = \{i : a_i > a_{i+1}\}.$$

An element of  $D(\pi)$  is called a *descent* of  $\pi$ , and *major index*  $\text{maj}(\pi)$  is defined as

$$\text{maj}(\pi) = \sum_{i \in D(\pi)} i.$$

The major index has many properties analogous to the number of inversions, e.g., a classic theorem of MacMahon states that  $\text{inv}$  and  $\text{maj}$  are equidistributed on the symmetric group  $\mathfrak{S}_n$  [6][7]. Thus it is natural to try to find “maj analogues” of the results of the preceding sections. In general, the major index of a linear extension of a poset can be more tractable or less tractable than the number of inversions. Thus, for example, in Theorem 5.1 we are able to completely characterize naturally labelled maj-balanced posets. An analogous result for sign-balanced partitions seems very difficult. On the other hand, since multiplying a permutation by a fixed permutation has no definite effect on the parity of the major index, many of the results for sign-balanced posets are false (Theorem 1.1, Lemma 2.1, Proposition 2.3).

Let  $f$  be a linear extension of the labelled poset  $(P, \omega)$ , and let  $\pi = \pi(f)$  be the associated permutation of  $[n]$ . In analogy to our definition of  $\text{inv}(f)$ , define  $\text{maj}(f) = \text{maj}(\pi)$  and

$$W_{P, \omega}(q) = \sum_{f \in \mathcal{E}_P} q^{\text{maj}(f)} = \sum_{\pi \in \mathcal{L}_{P, \omega}} q^{\text{maj}(\pi)}.$$

We say that  $(P, \omega)$  is *maj-balanced* if  $W_{P, \omega}(-1) = 0$ , i.e., if the number of linear extensions of  $P$  with even major index equals the number with odd major index. Unlike the situation for sign-balanced posets, the property of being maj-balanced can depend on the labeling  $\omega$ . Thus an interesting special case is that of *natural labelings*, for which  $\omega(s) < \omega(t)$  whenever  $s < t$  in  $P$ . We write  $W_P(q)$  for  $W_{P, \omega}(q)$  when  $\omega$  is natural. It is a basic consequence of the theory of  $P$ -partitions [22, Thm. 4.5.8] that  $W_P(q)$  does not depend on the choice of natural labeling of  $P$ .

Figures 3(a) and (b) show two different labelings of a poset  $P$ . The first labeling (which is natural) is not maj-balanced, while the second one is. Moreover, the dual poset  $P^*$  to the poset  $P$  in Figure 3(b), whether naturally labelled or labelled the same as  $P$ , is maj-balanced. Contrast that with the trivial fact that the dual of a sign-balanced poset is sign-balanced. As a further example of the contrast between sign and maj-balanced posets,




Figure 3: Some counterexamples

Figure 3(c) shows a naturally labelled maj-balanced poset  $Q$ . However, if we adjoin an element  $\hat{0}$  below every element of  $Q$  and label it 0 (thus keeping the labeling natural) then we get a poset which is no longer maj-balanced. On the other hand, it is clear that such an operation has no effect on whether a poset is sign-balanced. (In fact, it leaves  $I_{Q,\omega}(q)$  unchanged.)

Corollary 4.2 carries over to the major index in the following way.

**5.1 Theorem.** (a) *Let  $P$  be naturally labelled. Then  $W_P(-1)$  is equal to the number of  $P$ -domino tableaux. In particular,  $P$  is maj-balanced if and only if there does not exist a  $P$ -domino tableau.*

(b) *A labelled poset  $(P, \omega)$  is maj-balanced if there does not exist a  $P$ -domino tableau.*

**Proof.** (a) Let  $\pi = a_1 \cdots a_n \in \mathcal{L}_{P,\omega}$ . Let  $i$  be the least number (if it exists) for which  $\pi' = a_1 \cdots a_{2i} a_{2i+2} a_{2i+1} a_{2i+3} \cdots a_n \in \mathcal{L}_{P,\omega}$ . Note that  $(\pi')' = \pi$ . Now exactly one of  $\pi$  and  $\pi'$  has a descent at  $2i+1$ . The only other differences in the descent sets of  $\pi$  and  $\pi'$  occur (possibly) for the even numbers  $2i$  and  $2i+2$ . Hence  $(-1)^{\text{maj}(\pi)} + (-1)^{\text{maj}(\pi')} = 0$ . The surviving permutations  $\sigma = b_1 \cdots b_n$  in  $\mathcal{L}_{P,\omega}$  are exactly those for which  $\emptyset \subset \{b_1, b_2\} \subset \{b_1, \dots, b_4\} \subset \cdots$  is a  $P$ -domino tableau with  $\omega^{-1}(b_{2i-1}) < \omega^{-1}(b_{2i})$  in  $P$ . (If  $n$  is even, then the  $P$ -domino tableau ends as  $\{b_1, \dots, b_{n-2}\} \subset P$ , while if  $n$  is odd it ends as  $\{b_1, \dots, b_{n-1}\} \subset P$ .) Since  $\omega$  is natural we have  $b_{2i-1} < b_{2i}$  for all  $i$ , so  $\text{maj}(\sigma)$  is even. Hence  $W_P(-1)$  is equal exactly to the number of  $P$ -domino tableaux.




Figure 4: A maj-balanced labelled poset tilable by dominos

(b) Regardless of the labeling  $\omega$ , if there does not exist a  $P$ -domino tableau then there will be no survivors in the argument of (a), so  $W_P(-1) = 0$ .  $\square$

The converse to Theorem 5.1(b) is false. The labelled poset  $(P, \omega)$  of Figure 4 is tilable by dominos and is maj-balanced.

Given an  $n$ -element poset  $P$  with dual  $P^*$ , set  $\Delta(P) = \Gamma(P^*)$ . In [20, Thm 4.4][21, Prop. 18.4][22, Thm. 4.5.2] it is shown that the following two conditions are equivalent:

- (i) For all  $t \in P$ , all maximal chains of the principal dual order ideal  $V_t = \{s \in P : s \geq t\}$  have the same length.
- (ii)  $q^{\binom{n}{2} - \Delta(P)} W_P(1/q) = W_P(q)$ .

It follows by setting  $q = -1$  that if (i) holds and  $\binom{n}{2} - \Delta(P)$  is odd, then  $P$  is maj-balanced. Corollary 2.4 suggests in fact the following stronger result.

**5.2 Corollary.** *Suppose that  $P$  is naturally labelled and dual consistent (i.e.,  $P^*$  is consistent). If  $\binom{n}{2} - \Delta(P)$  is odd, then  $P$  is maj-balanced.*

**Proof.** By Theorem 5.1 we need to show that there does not exist a  $P$ -domino tableau. Given  $t \in P$ , let  $\delta(t)$  denote the length of the longest

chain of  $V_t$ , so  $\Delta(P) = \sum_{t \in P} \delta(t)$ . First suppose that  $n = 2m$ , and assume to the contrary that  $\emptyset = I_0 \subset I_1 \subset \cdots \subset I_m = P$  is a  $P$ -domino tableau. If  $s, t \in I_i - I_{i-1}$  then by dual consistency  $\delta(s) + \delta(t) \equiv 1 \pmod{2}$ . Hence  $\Delta(P) \equiv m \pmod{2}$ , so

$$\binom{n}{2} - \Delta(P) \equiv m(2m - 1) - m \equiv 0 \pmod{2},$$

a contradiction.

Similarly if  $n = 2m + 1$ , then the existence of a  $P$ -domino tableau implies  $\Delta(P) \equiv m \pmod{2}$ , so

$$\binom{n}{2} - \Delta(P) \equiv m(2m + 1) - m \equiv 0 \pmod{2},$$

again a contradiction.  $\square$

Now let  $\mathcal{S}$  be a finite subset of solid unit squares with integer vertices in  $\mathbb{R} \times \mathbb{R}$  such that the set  $|\mathcal{S}| = \bigcup_{S \in \mathcal{S}} S$  is simply-connected. For  $S, T \in \mathcal{S}$ , define  $S < T$  if the center vertices  $(s_1, s_2)$  of  $S$  and  $(t_1, t_2)$  of  $T$  satisfy either (a)  $t_1 = s_1$  and  $t_2 = s_2 + 1$  or (b)  $t_1 = s_1 + 1$  and  $t_2 = s_2$ . Regard  $\mathcal{S}$  as a poset, denoted  $P_{\mathcal{S}}$ , under the transitive (and reflexive) closure of the relation  $<$ . Figure 5 gives an example, where (a) shows  $\mathcal{S}$  as a set of squares and (b) as a poset. Note that the posets  $P_{\lambda/\mu}$  are a special case.

A *Schur labelling*  $\omega$  of  $P_{\mathcal{S}}$  is a labeling that increases along rows and decreases along columns, as illustrated in Figure 5. For the special case  $P_{\lambda/\mu}$ , Schur labelings play an important role in the expansion of skew Schur functions  $s_{\lambda/\mu}$  in terms of quasisymmetric functions [24, pp. 360–361]. Suppose that  $\#P_{\mathcal{S}}$  is even and that  $P_{\mathcal{S}}$  is tilable by dominos. Then  $\mathcal{S}$  itself is tilable by dominos in the usual sense. It is known (implicit, for instance, in [26], and more explicit in [4]) that any two domino tilings of  $\mathcal{S}$  can be obtained from each other by “ $2 \times 2$  flips,” i.e., replacing two horizontal dominos in a  $2 \times 2$  square by two vertical dominos or *vice versa*. It follows that if  $D$  is a domino tiling of  $\mathcal{S}$  with  $v(D)$  vertical dominos, then  $(-1)^{v(D)}$  depends only on  $\mathcal{S}$ . Set  $\text{sgn}(\mathcal{S}) = (-1)^{v(D)}$  for any domino tiling of  $\mathcal{S}$ .

**5.3 Proposition.** *Let  $\mathcal{S}$  be as above, and let  $\omega$  be a Schur labeling of*




Figure 5: A set  $\mathcal{S}$  of squares and the Schur labelled poset  $P_{\mathcal{S}}$

$P_{\mathcal{S}}$ , where  $\#P_{\mathcal{S}}$  is even, say  $\#P_{\mathcal{S}} = n$ . Then  $\text{sgn}(\mathcal{S})W_{P_{\mathcal{S}}}(-1)$  is the number of  $P_{\mathcal{S}}$ -domino tableaux.

**Proof.** The proof parallels that of Theorem 5.1. Define the involution  $\pi \mapsto \pi'$  as in the proof of Theorem 5.1. Each survivor  $\sigma = b_1 \cdots b_n$  corresponds to a  $P_{\mathcal{S}}$ -domino tableau  $D$ . We have  $b_{2i-1} > b_{2i}$  if and only if the domino labelled with  $b_{2i-1}$  and  $b_{2i}$  is vertical. As noted above,  $(-1)^{v(D)} = \text{sgn}(\mathcal{S})$ , independent of  $D$ . Hence  $(-1)^{\text{maj}(\sigma)} = \text{sgn}(\sigma)$ , and the proof follows as in Theorem 5.1(a).  $\square$

A result analogous to Proposition 5.3 holds for  $\#P_{\mathcal{S}}$  odd (with essentially the same proof) provided  $P_{\mathcal{S}}$  has a  $\hat{0}$  or  $\hat{1}$ . The special case  $P_{\lambda/\mu}$  of Proposition 5.3 (and its analogue for  $\#P_{\mathcal{S}}$  odd) can also be proved using the theory of symmetric functions, notably, [24, Prop. 7.19.11] and the Murnaghan-Nakayama rule ([24, Cor. 7.17.5]).

## 6 Hook lengths.

In this section we briefly discuss a class of posets  $P$  for which  $W_P(q)$ , and sometimes even  $I_{P,\omega}(q)$ , can be explicitly computed. For this class of posets

we get a simple criterion for being maj balanced and, if applicable, sign balanced.

Following [21, p. 84], an  $n$ -element poset  $P$  is called a *hook length poset* if there exist positive integers  $h_1, \dots, h_n$ , the *hook lengths* of  $P$ , such that

$$W_P(q) = \frac{[n]!}{(1 - q^{h_1}) \cdots (1 - q^{h_n})}, \quad (10)$$

where  $[n]! = (1 - q)(1 - q^2) \cdots (1 - q^n)$ . It is easy to see that if  $P$  is a hook length poset, then the multiset of hook lengths is unique. In general, if  $P$  is an “interesting” hook length poset, then each element of  $P$  should have a hook length associated to it in a “natural” combinatorial way.

NOTE. We could just as easily have extended our definition to *labelled* posets  $(P, \omega)$ , where now

$$W_{P, \omega}(q) = \frac{q^c [n]!}{(1 - q^{h_1}) \cdots (1 - q^{h_n})}$$

for some  $c \in \mathbb{N}$ . However, little is known about the labelled situation except when we can reduce it to the case of natural labelings by subtracting certain constants from the values of  $\sigma$ .

The following result is an immediate consequence of equation (10).

**6.1 Proposition.** *Suppose that  $P$  is a hook length poset with hook lengths  $h_1, \dots, h_n$ . Then  $P$  is maj-balanced if and only if the number of even hook lengths is less than  $\lfloor n/2 \rfloor$ . If  $P$  isn't maj-balanced, then the maj imbalance is given by*

$$W_P(-1) = \frac{\lfloor n/2 \rfloor!}{\prod_{h_i \text{ even}} (h_i/2)}.$$

It is natural to ask at this point what are the known hook length posets. The strongest work in this area is due to Proctor [14][15]. We won't state his remarkable results here, but let us note that his *d-complete* posets encompass all known “interesting” examples of hook length posets. These include forests



(i.e., posets for which every element is covered by at most one element) and the duals  $P_\lambda^*$  of the posets  $P_\lambda$  of Section 3.

Björner and Wachs [2, Thm. 1.1] settle the question of what naturally labelled posets  $(P, \omega)$  satisfy

$$I_{P,\omega}(q) = W_{P,\omega}(q). \quad (11)$$

Namely,  $P$  is a forest and  $\omega$  is a post ordered labeling. Hence for post ordered labelled forests, Proposition 6.1 holds also for  $I_{P,\omega}(-1)$ . Björner and Wachs also obtain less definitive results for arbitrary labelings, whose relevance to sign and maj imbalance we omit.

## References

- [1] G. E. Andrews, *The Theory of Partitions*, Cambridge University Press, Cambridge, 1998.
- [2] A. Björner and M. Wachs, Permutation statistics and linear extensions of posets, *J. Combinatorial Theory (A)* **58** (1991), 85–114.
- [3] C. Carré and B. Leclerc, Splitting the square of a Schur function into its symmetric and antisymmetric parts, *J. Algebraic Combin.* **4** (1995), 201–231.
- [4] T. Chaboud, Domino tiling in planar graphs with regular and bipartite dual, *Theoret. Comput. Sci.* **159** (1996), 137–142.
- [5] P. Edelman and C. Greene, Balanced tableaux, *Advances in Math.* **63** (1987), 42–99.
- [6] D. Foata, On the Netto inversion number of a sequence, *Proc. Amer. Math. Soc.* **19** (1968), 236–240.
- [7] D. Foata and M.-P. Schützenberger, Major index and inversion number of permutations, *Math. Nachr.* **83** (1978), 143–159.
- [8] M. Haiman, On mixed insertion, symmetry, and shifted Young tableaux, *J. Combinatorial Theory (A)* **50** (1989), 196–225.

- [9] M. Haiman, Dual equivalence with applications, including a conjecture of Proctor, *Discrete Math.* **99** (1992), 79–113.
- [10] A. N. Kirillov, A. Lascoux, B. Leclerc, and J.-Y. Thibon, Séries génératrices pour les tableaux de dominos, *C. R. Acad. Sci. Paris, Série I* **318** (1994), 395–400.
- [11] A. Lascoux, B. Leclerc, and J.-Y. Thibon, Ribbon tableaux, Hall-Littlewood symmetric functions, quantum affine algebras, and unipotent varieties, *J. Math. Phys.* **38**(2) (1997), 1041–1068.
- [12] M. van Leeuwen, Flag varieties and interpretations of Young tableaux algorithms, *J. Algebra* **224** (2000), 397–426; math.CO/9908041.
- [13] I. G. Macdonald, *Symmetric Functions and Hall Polynomials*, second ed., Oxford University Press, New York, 1995.
- [14] R. A. Proctor, Minuscule elements of Weyl groups, the numbers game, and  $d$ -complete posets, *J. Algebra* **213** (1999), 272–303.
- [15] R. A. Proctor, Dynkin diagram classification of lambda-minuscule Bruhat lattices and of  $d$ -complete posets, *J. Algebraic Combinatorics* **9** (1999), 61–94.
- [16] R. Ruskey, Research problem 90, *Discrete Math.* **70** (1989), 111–112.
- [17] F. Ruskey, Generating linear extensions of posets by transpositions, *J. Combinatorial Theory Ser. B* **54** (1992), 77–101.
- [18] M.-P. Schützenberger, Promotion des morphisms d’ensemble ordonnés, *Discrete Math.* **2** (1972), 73–94.
- [19] M. Shimozono and D. E. White, A color-to-spin domino Schensted algorithm, *Electron. J. Combinatorics*, to appear.
- [20] R. Stanley, Ordered Structures and Partitions, Ph.D. thesis, Harvard University, 1971.
- [21] R. Stanley, Ordered structures and partitions, *Memoirs of the Amer. Math. Soc.*, no. 119, 1972.

- [22] R. Stanley, *Enumerative Combinatorics*, vol. 1, Wadsworth and Brooks/Cole, Pacific Grove, CA, 1986; second printing, Cambridge University Press, Cambridge, 1996.
- [23] R. Stanley, Differential posets, *J. Amer. Math. Soc.* **1** (1988), 919–961.
- [24] R. Stanley, *Enumerative Combinatorics*, vol. 2, Cambridge University Press, New York/Cambridge, 1999.
- [25] R. Stanley, Problem 10969, *Amer. Math. Monthly* **109** (2002), 760.
- [26] W. Thurston, Conway’s tiling group, *Amer. Math. Monthly* **97** (1990), 757–773.
- [27] D. White, Sign-balanced posets, *J. Combinatorial Theory Ser. A*, to appear.