Generalizations of Cauchy's Determinant and Schur's Pfaffian

Masao Ishikawa*

Soichi Okada[†]

Hiroyuki Tagawa[‡]

Jiang Zeng[§]

Abstract

We present several identities of Cauchy-type determinants and Schur-type Pfaffians involving generalized Vandermonde determinants, which generalize Cauchy's determinant det $(1/(x_i + y_j))$ and Schur's Pfaffian Pf $((x_j - x_i)/(x_j + x_i))$. Some special cases of these identities are given by S. Okada and T. Sundquist. As an application, we give a relation for the Littlewood– Richardson coefficients involving a rectangular partition.

Keywords : Pfaffian, Cauchy's Determinant, Plücker relations, Schur functions.

1 Introduction

Computations of determinants and Pfaffians are of great importance not only in many branches of mathematics but also in physics. Some people need relations among minors or subpfaffians of a general matrix, others have to evaluate special determinants or Pfaffians. In enumerative combinatorics and representation theory, a central role is played by Cauchy's determinant identity [2]

$$\det\left(\frac{1}{x_i + y_j}\right)_{1 \le i,j \le n} = \frac{\prod_{1 \le i < j \le n} (x_j - x_i)(y_j - y_i)}{\prod_{i,j=1}^n (x_i + y_j)},$$
(1.1)

and Schur's Pfaffian identity [17]

$$\operatorname{Pf}\left(\frac{x_j - x_i}{x_j + x_i}\right)_{1 \le i, j \le 2n} = \prod_{1 \le i < j \le 2n} \frac{x_j - x_i}{x_j + x_i}.$$
(1.2)

The reader is referred to [4, 6, 9, 13, 14, 18, 19] for some recent variations and generalizations with their applications of (1.2) and (1.1). Besides, Krattenthaler [8] has given a comprehensive survey of determinant evaluations.

In the same vein, we shall give several identities of Cauchy-type determinants and Schurtype Pfaffians whose entries involve two kinds of generalized Vandermonde determinants. Let $\boldsymbol{x} = (x_1, \dots, x_n)$ and $\boldsymbol{a} = (a_1, \dots, a_n)$ be two vectors of variables of length n. Let p and q be two nonnegative integers such that p + q = n. Denote by $V^{p,q}(\boldsymbol{x}; \boldsymbol{a})$ the $n \times n$ matrix with *i*th row

$$(1, x_i, \cdots, x_i^{p-1}, a_i, a_i x_i, \cdots, a_i x_i^{q-1}),$$

and $W^n(\boldsymbol{x}; \boldsymbol{a})$ the $n \times n$ matrix with *i*th row

$$(1 + a_i x_i^{n-1}, x_i + a_i x_i^{n-2}, \cdots, x_i^{n-1} + a_i).$$

For example, if q = 0, then $V^{n,0}(\boldsymbol{x}; \boldsymbol{a}) = \left(x_i^{j-1}\right)_{1 \le i,j \le n}$ is the usual Vandermonde matrix and $\det V^{n,0}(\boldsymbol{x}; \boldsymbol{a}) = \prod_{1 \le i < j \le n} (x_j - x_i)$. If p = q = 1, then $\det V^{1,1}(\boldsymbol{x}; \boldsymbol{a}) = a_2 - a_1$, while the

^{*}Faculty of Education, Tottori University, e-mail:ishikawa@fed.tottori-u.ac.jp

[†]Graduate School of Mathematics, Nagoya University, e-mail:okada@math.nagoya-u.ac.jp

[‡]Faculty of Education, Wakayama University, e-mail:tagawa@math.edu.wakayama-u.ac.jp

[§]Institut Girard Desargues, Université Claude Bernard Lyon-I, e-mail:**zeng@igd.univ-lyon1.fr**

matrices $V^{3,2}(\boldsymbol{x};\boldsymbol{a})$ and $W^5(\boldsymbol{x};\boldsymbol{a})$ can be visualized as follows:

$$W^{5}(\boldsymbol{x};\boldsymbol{a}) = \begin{pmatrix} 1 & x_{1} & x_{1}^{2} & a_{1} & a_{1}x_{1} \\ 1 & x_{2} & x_{2}^{2} & a_{2} & a_{2}x_{2} \\ 1 & x_{3} & x_{3}^{2} & a_{3} & a_{3}x_{2} \\ 1 & x_{4} & x_{4}^{2} & a_{4} & a_{4}x_{4} \\ 1 & x_{5} & x_{5}^{2} & a_{5} & a_{5}x_{5} \end{pmatrix},$$

$$W^{5}(\boldsymbol{x};\boldsymbol{a}) = \begin{pmatrix} 1 + a_{1}x_{1}^{4} & x_{1} + a_{1}x_{1}^{3} & x_{1}^{2} + a_{1}x_{1}^{2} & x_{1}^{3} + a_{1}x_{1} & x_{1}^{4} + a_{1} \\ 1 + a_{2}x_{2}^{4} & x_{2} + a_{2}x_{2}^{3} & x_{2}^{2} + a_{2}x_{2}^{2} & x_{2}^{3} + a_{2}x_{2} & x_{2}^{4} + a_{2} \\ 1 + a_{3}x_{3}^{4} & x_{3} + a_{3}x_{3}^{3} & x_{3}^{2} + a_{3}x_{3}^{2} & x_{3}^{3} + a_{3}x_{3} & x_{3}^{4} + a_{4} \\ 1 + a_{5}x_{5}^{4} & x_{5} + a_{5}x_{5}^{3} & x_{5}^{2} + a_{5}x_{5}^{2} & x_{5}^{3} + a_{5}x_{5} & x_{5}^{4} + a_{5} \end{pmatrix}.$$

The main purpose of this paper is to prove the following identities for the determinants and Pfaffians whose entries involve these generalized Vandermonde determinants.

Theorem 1.1. (a) Let n be a positive integer and let p and q be nonnegative integers. For six vectors of variables

$$\boldsymbol{x} = (x_1, \cdots, x_n), \ \boldsymbol{y} = (y_1, \cdots, y_n), \ \boldsymbol{a} = (a_1, \cdots, a_n), \ \boldsymbol{b} = (b_1, \cdots, b_n),$$

 $\boldsymbol{z} = (z_1, \cdots, z_{p+q}), \ \boldsymbol{c} = (c_1, \cdots, c_{p+q}),$

we have

$$\det\left(\frac{\det V^{p+1,q+1}(x_i, y_j, \boldsymbol{z}; a_i, b_j, \boldsymbol{c})}{y_j - x_i}\right)_{1 \le i, j \le n} = \frac{(-1)^{n(n-1)/2}}{\prod_{i,j=1}^n (y_j - x_i)} \det V^{p,q}(\boldsymbol{z}; \boldsymbol{c})^{n-1} \det V^{n+p,n+q}(\boldsymbol{x}, \boldsymbol{y}, \boldsymbol{z}; \boldsymbol{a}, \boldsymbol{b}, \boldsymbol{c}).$$
(1.3)

(b) Let n be a positive integer and let p, q, r, s be nonnegative integers. For seven vectors of variables

we have

$$\Pr\left(\frac{\det V^{p+1,q+1}(x_i, x_j, \boldsymbol{z}; a_i, a_j, \boldsymbol{c}) \det V^{r+1,s+1}(x_i, x_j, \boldsymbol{w}; b_i, b_j, \boldsymbol{d})}{x_j - x_i}\right)_{1 \le i,j \le 2n} = \frac{1}{\prod_{1 \le i < j \le 2n} \det V^{p,q}(\boldsymbol{z}; \boldsymbol{c})^{n-1} \det V^{r,s}(\boldsymbol{w}; \boldsymbol{d})^{n-1}} \times \det V^{n+p,n+q}(\boldsymbol{x}, \boldsymbol{z}; \boldsymbol{a}, \boldsymbol{c}) \det V^{n+r,n+s}(\boldsymbol{x}, \boldsymbol{w}; \boldsymbol{b}, \boldsymbol{d}). \quad (1.4)$$

(c) Let n be a positive integer and let p be a nonnegative integer. For six vectors of variables

$$x = (x_1, \dots, x_n), \ y = (y_1, \dots, y_n), \ a = (a_1, \dots, a_n), \ b = (b_1, \dots, b_n),$$

 $z = (z_1, \dots, z_p), \ c = (c_1, \dots, c_p),$

we have

$$\det\left(\frac{\det W^{p+2}(x_i, y_j, \boldsymbol{z}; a_i, b_j, \boldsymbol{c})}{(y_j - x_i)(1 - x_i y_j)}\right)_{1 \le i, j \le n} = \frac{1}{\prod_{i,j=1}^n (y_j - x_i)(1 - x_i y_j)} \det W^p(\boldsymbol{z}; \boldsymbol{c})^{n-1} \det W^{2n+p}(\boldsymbol{x}, \boldsymbol{y}, \boldsymbol{z}; \boldsymbol{a}, \boldsymbol{b}, \boldsymbol{c}).$$
(1.5)

(d) Let n be a positive integer and let p and q be nonnegative integers. For seven vectors of variables

we have

$$Pf\left(\frac{\det W^{p+2}(x_i, x_j, \boldsymbol{z}; a_i, a_j, \boldsymbol{c}) \det W^{q+2}(x_i, x_j, \boldsymbol{w}; b_i, b_j, \boldsymbol{d})}{(x_j - x_i)(1 - x_i x_j)}\right)_{1 \le i, j \le 2n} = \frac{1}{\prod_{1 \le i < j \le 2n} (x_j - x_i)(1 - x_i x_j)} \det W^p(\boldsymbol{z}; \boldsymbol{c})^{n-1} \det W^q(\boldsymbol{w}; \boldsymbol{d})^{n-1} \times \det W^{2n+p}(\boldsymbol{x}, \boldsymbol{z}; \boldsymbol{a}, \boldsymbol{c}) \det W^{2n+q}(\boldsymbol{x}, \boldsymbol{w}; \boldsymbol{b}, \boldsymbol{d}). \quad (1.6)$$

These identities were conjectured by one of the authors [15]. If we put p = q = 0 in (1.3) or p = q = r = s = 0 in (1.4), then the identities read

$$\det\left(\frac{b_j - a_i}{y_j - x_i}\right)_{1 \le i, j \le n} = \frac{(-1)^{n(n-1)/2}}{\prod_{i,j=1}^n (y_j - x_i)} \det V^{n,n}(\boldsymbol{x}, \boldsymbol{y}; \boldsymbol{a}, \boldsymbol{b}),$$
(1.7)

$$\Pr\left(\frac{(a_j - a_i)(b_j - b_i)}{x_j - x_i}\right)_{1 \le i, j \le 2n} = \frac{1}{\prod_{1 \le i < j \le 2n} (x_j - x_i)} \det V^{n,n}(\boldsymbol{x}; \boldsymbol{a}) \det V^{n,n}(\boldsymbol{x}; \boldsymbol{b}).$$
(1.8)

These particular cases, as well as the identities (1.5) with p = 0 and (1.6) with p = q = 0, are first given by S. Okada [13, Theorems 4.2, 4.7, 4.3, 4.4] in his study of rectangular-shaped representations of classical groups. Another special case of the identity (1.5) with p = 1 is given in [14] and applied to the enumeration of vertically and horizontally symmetric alternating sign matrices. These special cases are the starting point of our study.

Under the specialization

 $x_i \leftarrow x_i^2, \quad y_i \leftarrow y_i^2, \quad z_i \leftarrow z_i^2, \quad w_i \leftarrow w_i^2, \quad a_i \leftarrow x_i, \quad b_i \leftarrow y_i, \quad c_i \leftarrow z_i, \quad d_i \leftarrow w_i,$

one can deduce from (1.3) and (1.4) the following identities:

$$\det\left(\frac{s_{\delta(k)}(x_{i}, y_{j}, \boldsymbol{z})}{x_{i} + y_{j}}\right)_{1 \leq i, j \leq n} = \frac{\prod_{1 \leq i < j \leq n} (x_{j} - x_{i})(y_{j} - y_{i})}{\prod_{i, j = 1}^{n} (x_{i} + y_{j})} s_{\delta(k)}(\boldsymbol{z})^{n-1} s_{\delta(k)}(\boldsymbol{x}, \boldsymbol{y}, \boldsymbol{z}), \quad (1.9)$$

$$\operatorname{Pf}\left(\frac{x_{j} - x_{i}}{x_{j} + x_{i}} s_{\delta(k)}(x_{i}, x_{j} \boldsymbol{z}) s_{\delta(l)}(x_{i}, x_{j}, \boldsymbol{w})\right)_{1 \leq i, j \leq 2n}$$

$$= \prod_{1 \leq i < j \leq 2n} \frac{x_{j} - x_{i}}{x_{j} + x_{i}} s_{\delta(k)}(\boldsymbol{z})^{n-1} s_{\delta(l)}(\boldsymbol{w})^{n-1} s_{\delta(k)}(\boldsymbol{x}, \boldsymbol{z}) s_{\delta(l)}(\boldsymbol{x}, \boldsymbol{w}), \quad (1.10)$$

where s_{λ} denotes the Schur function corresponding to a partition λ and $\delta(k) = (k, k - 1, \dots, 1)$ denotes the staircase partition. If we take k = 0 in (1.9) and k = l = 0 in (1.10), we obtain Cauchy's determinant identity (1.1) and Schur's Pfaffian identity (1.2). Another special case of (1.9) with k = 1 is the rational case of Frobenius' identity [3]. Also, if we take k = l = 1 in (1.10), we obtain the rational case of an elliptic generalization of (1.2) given in [16].

This paper is organized as follows. Sections 2 and 3 are devoted to the proof of Theorem 1.1. In Section 2, we prove the identity (1.4) by using the Pfaffian version of Desnanot–Jacobi formula and induction. In Section 3, we give a homogeneous version of the identity (1.4) and derive the other three identities (1.3), (1.5) and (1.6). A variation of the main identities is given in Section 4, and another instance of a Cauchy-type determinant identity is presented in Section 5. Also we present a formula expressing the determinant of $V^{n,n}$ in terms of the hyperpfaffian. In the last section, we give an application of the identity (1.4) to the Littlewood–Richardson coefficients.

Here we recall the definition of Pfaffians. Given a $2n \times 2n$ skew-symmetric matrix $A = (a_{ij})_{1 \le i,j \le 2n}$, the Pfaffian of A is defined by

$$Pf(A) = \sum_{\sigma} sgn(\sigma) a_{\sigma(1),\sigma(2)} a_{\sigma(3),\sigma(4)} \cdots a_{\sigma(2n-1),\sigma(2n)},$$

where σ runs over all permutations of 2n letters $1, 2, \dots, 2n$ satisfying

$$\sigma(1) < \sigma(2), \quad \sigma(3) < \sigma(4), \quad \cdots, \quad \sigma(2n-1) < \sigma(2n),$$

$$\sigma(1) < \sigma(3) < \cdots < \sigma(2n-1).$$

2 Proof of the identity (1.4) in Theorem 1.1

In this section, we give a proof of the identity (1.4). First we show (1.4) in the special case where n = 2 by using induction. Then we apply the Desnanot–Jacobi formula for Pfaffians to reduce the proof of the general case to this special case.

First we prove the case of n = 2 by induction on p + q + r + s.

Proposition 2.1. Let p, q, r and s be nonnegative integers. For vectors of variables x, a, b of length 4, z, c of length p + q, and w, d of length r + s, we have

$$\operatorname{Pf}\left(\frac{\det V^{p+1,q+1}(x_i, x_j, \boldsymbol{z}; a_i, a_j, \boldsymbol{c}) \det V^{r+1,s+1}(x_i, x_j, \boldsymbol{w}; b_i, b_j, \boldsymbol{d})}{x_j - x_i}\right)_{1 \le i,j \le 4}$$
$$= \frac{1}{\prod_{1 \le i < j \le 4} (x_j - x_i)} \det V^{p,q}(\boldsymbol{z}; \boldsymbol{c}) \det V^{r,s}(\boldsymbol{w}; \boldsymbol{d})$$
$$\times \det V^{p+2,q+2}(\boldsymbol{x}, \boldsymbol{z}; \boldsymbol{a}, \boldsymbol{c}) \det V^{r+2,s+2}(\boldsymbol{x}, \boldsymbol{w}; \boldsymbol{b}, \boldsymbol{d}). \quad (2.1)$$

In the induction step of the proof, we need relations between det $V^{p,q}$ and det $V^{p-1,q}$ (or det $V^{q,p}$).

Lemma 2.2. (1) If $p \ge q$ and $p \ge 1$, then we have

$$\det V^{p,q}(\boldsymbol{x};\boldsymbol{a}) = \prod_{i=1}^{p+q-1} (x_{p+q} - x_i) \cdot \det V^{p-1,q}(x_1,\cdots,x_{p+q-1};a'_1,\cdots,a'_{p+q-1}), \quad (2.2)$$

where we put

$$a'_i = \frac{a_i - a_{p+q}}{x_i - x_{p+q}}$$
 $(1 \le i \le p+q-1).$

(2) For nonnegative integers p and q, we have

det
$$V^{p,q}(\boldsymbol{x}; \boldsymbol{a}) = (-1)^{pq} \prod_{i=1}^{p+q} a_i \cdot \det V^{q,p}(\boldsymbol{x}; \boldsymbol{a}^{-1}),$$
 (2.3)

where $\boldsymbol{a}^{-1} = (a_1^{-1}, \cdots, a_{p+q}^{-1}).$

Proof. (1) We put m = p + q. By subtracting the *i*th column multiplied by a_m from the (p+i)th column for $i = 1, \dots, q$, and by subtracting the *i*th column multiplied by x_m from the (i + 1)th column for $i = p - 1, \dots, 1$, we obtain

$$\det V^{p,q}(\boldsymbol{x};\boldsymbol{a}) = \det \begin{pmatrix} 1 & x_1 - x_m & \cdots & (x_1 - x_m)x_1^{p-2} & a_1 - a_m & \cdots & (a_1 - a_m)x_1^{q-1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & x_{m-1} - x_m & \cdots & (x_{m-1} - x_m)x_{m-1}^{p-2} & a_{m-1} - a_m & \cdots & (a_{m-1} - a_m)x_{m-1}^{q-1} \\ 1 & 0 & \cdots & 0 & 0 & \cdots & 0 \end{pmatrix}$$
$$= (-1)^{m+1} \prod_{k=1}^{m-1} (x_k - x_m) \cdot \det \begin{pmatrix} 1 & x_1 & \cdots & x_1^{p-2} & a_1' & a_1'x_1 & \cdots & a_1'x_1^{q-1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & x_{m-1} & \cdots & x_{m-1}^{p-2} & a_{m-1}' & a_{m-1}'x_{m-1} & \cdots & a_{m-1}'x_{m-1}^{q-1} \end{pmatrix}$$
$$= \prod_{k=1}^{p+q-1} (x_{p+q} - x_k) \cdot \det V^{p-1,q}(x_1, \cdots, x_{p+q-1}; a_1', \cdots, a_{p+q-1}').$$

(2) As before we put m = p + q. By performing an appropriate permutation of the columns and by dividing the *i*th row by a_i , we obtain

$$\det V^{p,q}(\boldsymbol{x};\boldsymbol{a}) = (-1)^{pq} \det \begin{pmatrix} a_1 & a_1x_1 & \cdots & a_1x_1^{q-1} & 1 & x_1 & \cdots & x_1^{p-1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ a_m & a_mx_m & \cdots & a_mx_m^{q-1} & 1 & x_m & \cdots & x_m^{p-1} \end{pmatrix}$$
$$= (-1)^{pq} \prod_{k=1}^{p+q} a_k \cdot \det \begin{pmatrix} 1 & x_1 & \cdots & x_1^{q-1} & a_1^{-1} & a_1^{-1}x_1 & \cdots & a_1^{-1}x_1^{p-1} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & x_m & \cdots & x_m^{q-1} & a_m^{-1} & a_m^{-1}x_m & \cdots & a_m^{-1}x_m^{q-1} \end{pmatrix}$$
$$= (-1)^{pq} \prod_{k=1}^{p+q} a_k \cdot \det V^{q,p}(\boldsymbol{x};\boldsymbol{a}^{-1}).$$

This completes the proof of the lemma.

Proof of Proposition 2.1. We prove (2.1) by induction on p + q + r + s. If p + q + r + s = 0, i.e., p = q = r = s = 0, then one can easily check the equality in (2.1) by a direct computation.

Suppose p + q + r + s > 0. By symmetry, we may assume p + q > 0 without loss of generality. First we consider the case where $p \ge q$. Using the relation (2.2) in Lemma 2.2, we have

$$\det V^{p+1,q+1}(x_i, x_j, \boldsymbol{z}; a_i, a_j, \boldsymbol{c}) = (z_{p+q} - x_i)(z_{p+q} - x_j) \prod_{k=1}^{p+q-1} (z_{p+q} - z_k) \cdot \det V^{p,q+1}(x_i, x_j, \boldsymbol{\tilde{z}}; a'_i, a'_j, \boldsymbol{c'}),$$

where $\widetilde{\boldsymbol{z}} = (z_1, \cdots, z_{p+q-1})$ and a'_i, a'_j and $\boldsymbol{c'} = (c'_1, \cdots, c'_{p+q-1})$ are given by

$$a'_{k} = \frac{a_{k} - c_{p+q}}{x_{k} - z_{p+q}}$$
 $(k = i, j), \quad c'_{l} = \frac{c_{l} - c_{p+q}}{z_{l} - z_{p+q}}$ $(1 \le l \le p + q - 1).$

Hence we have

$$\Pr\left(\frac{\det V^{p+1,q+1}(x_i, x_j, \boldsymbol{z}; a_i, a_j, \boldsymbol{c}) \det V^{r+1,s+1}(x_i, x_j, \boldsymbol{w}; b_i, b_j, \boldsymbol{d})}{x_j - x_i}\right)_{1 \le i,j \le 4}$$

$$= \prod_{i=1}^4 (z_{p+q} - x_i) \prod_{i=1}^{p+q-1} (z_{p+q} - z_i)^2$$

$$\times \Pr\left(\frac{\det V^{p,q+1}(x_i, x_j, \boldsymbol{\tilde{z}}; a_i', a_j', \boldsymbol{c'}) \det V^{r+1,s+1}(x_i, x_j, \boldsymbol{w}; b_i, b_j, \boldsymbol{d})}{x_j - x_i}\right)_{1 \le i,j \le 4}$$

by the induction hypothesis,

$$= \prod_{i=1}^{4} (z_{p+q} - x_i) \prod_{i=1}^{p+q-1} (z_{p+q} - z_i)^2 \\\times \frac{1}{\prod_{1 \le i < j \le 4} (x_j - x_i)} \det V^{p-1,q}(\widetilde{\boldsymbol{z}}; \boldsymbol{c'}) \det V^{r,s}(\boldsymbol{w}; \boldsymbol{d}) \\\times \det V^{p+1,q+2}(x_1, x_2, x_3, x_4, \widetilde{\boldsymbol{z}}; a'_1, a'_2, a'_3, a'_4, \boldsymbol{c'}) \det V^{r+2,s+2}(\boldsymbol{x}, \boldsymbol{w}; \boldsymbol{b}, \boldsymbol{d})$$

by using the relation (2.2) again,

$$= \frac{1}{\prod_{1 \le i < j \le 4} (x_j - x_i)} \det V^{p,q}(\boldsymbol{z}; \boldsymbol{c}) \det V^{r,s}(\boldsymbol{w}; \boldsymbol{d}) \det V^{p+2,q+2}(\boldsymbol{x}, \boldsymbol{z}; \boldsymbol{a}, \boldsymbol{c}) \det V^{r+2,s+2}(\boldsymbol{x}, \boldsymbol{w}; \boldsymbol{b}, \boldsymbol{d}).$$

If p < q, then we use the relation (2.3) in Lemma 2.2 and the case we have just proven. Then we see that

$$\begin{aligned} &\Pr\left(\frac{\det V^{p+1,q+1}(x_i, x_j, \boldsymbol{z}; a_i, a_j, \boldsymbol{c}) \det V^{r+1,s+1}(x_i, x_j, \boldsymbol{w}; b_i, b_j, \boldsymbol{d})}{x_j - x_i}\right)_{1 \le i,j \le 4} \\ &= \Pr\left(\frac{a_i a_j \prod_{k=1}^{p+q} c_k \cdot \det V^{q+1,p+1}(x_i, x_j, \boldsymbol{z}; a_i^{-1}, a_j^{-1}, \boldsymbol{c}^{-1}) \det V^{r+1,s+1}(x_i, x_j, \boldsymbol{w}; b_i, b_j, \boldsymbol{d})}{x_j - x_i}\right)_{1 \le i,j \le 4} \\ &= \prod_{i=1}^4 a_i \prod_{k=1}^{p+q} c_k^2 \\ &\times \frac{1}{\prod_{1 \le i < j \le 2n} (x_j - x_i)} \det V^{q,p}(\boldsymbol{z}; \boldsymbol{c}^{-1}) \det V^{r,s}(\boldsymbol{w}; \boldsymbol{d}) \det V^{q+2,p+2}(\boldsymbol{x}, \boldsymbol{z}; \boldsymbol{a}^{-1}, \boldsymbol{c}^{-1}) \det V^{r+2,s+2}(\boldsymbol{x}, \boldsymbol{w}; \boldsymbol{b}, \boldsymbol{d}) \\ &= \frac{1}{\prod_{1 \le i < j \le 4} (x_j - x_i)} \det V^{p,q}(\boldsymbol{z}; \boldsymbol{c}) \det V^{r,s}(\boldsymbol{w}; \boldsymbol{d}) \det V^{p+2,q+2}(\boldsymbol{x}, \boldsymbol{z}; \boldsymbol{a}, \boldsymbol{c}) \det V^{r+2,s+2}(\boldsymbol{x}, \boldsymbol{w}; \boldsymbol{b}, \boldsymbol{d}). \end{aligned}$$

This completes the proof of Proposition 2.1.

Here we recall the Desnanot-Jacobi formula for determinants and Pfaffians. Given a square matrix A and indices $i_1, \dots, i_r, j_1, \dots, j_r$, we denote by $A_{j_1, \dots, j_r}^{i_1, \dots, i_r}$ the matrix obtained by removing the rows i_1, \dots, i_r and the columns j_1, \dots, j_r of A.

Lemma 2.3. (1) If A is a square matrix, then we have

$$\det A_1^1 \cdot \det A_2^2 - \det A_2^1 \cdot \det A_1^2 = \det A \cdot \det A_{1,2}^{1,2}.$$
(2.4)

(2) If A is a skew-symmetric matrix A, then we have

$$\Pr A_{1,2}^{1,2} \cdot \Pr A_{3,4}^{3,4} - \Pr A_{1,3}^{1,3} \cdot \Pr A_{2,4}^{2,4} + \Pr A_{1,4}^{1,4} \cdot \Pr A_{2,3}^{2,3} = \Pr A \cdot \Pr A_{1,2,3,4}^{1,2,3,4}.$$
 (2.5)

This Pfaffian analogue of the Desnanot-Jacobi formula is given in [7] and [6].

If $\boldsymbol{x} = (x_1, \ldots, x_n)$ is a vector of variables and $1 \leq i_1 < \cdots < i_r \leq n$ are indices, we denote by $\boldsymbol{x}^{(i_1, \cdots, i_r)}$ the vector obtained from \boldsymbol{x} by removing the variables x_{i_1}, \cdots, x_{i_r} .

Proof of the identity in (1.4) in Theorem 1.1. We proceed by induction on n. If n = 1, then there is nothing to prove, and, if n = 2, we already proved (1.4) in Proposition 2.1.

Suppose $n \ge 3$. Apply the Desnanot–Jacobi formula for Pfaffians (2.5) in Lemma 2.3 to the skew-symmetric matrix

$$A = \left(\frac{\det V^{p+1,q+1}(x_i, x_j, \boldsymbol{z}; a_i, a_j, \boldsymbol{c}) \det V^{r+1,s+1}(x_i, x_j, \boldsymbol{w}; b_i, b_j, \boldsymbol{d})}{x_j - x_i}\right)_{1 \le i, j \le 2n}.$$

Then the induction hypothesis tells us that, for $1 \le k < l \le 4$, we have

$$\Pr A_{k,l}^{k,l} = \frac{1}{(x_{l'} - x_{k'}) \prod_{i=5}^{2n} (x_i - x_{k'}) (x_i - x_{l'}) \prod_{5 \le i < j \le 2n} (x_j - x_i)} \times \det V^{p,q}(\boldsymbol{z}; \boldsymbol{c})^{n-2} \det V^{r,s}(\boldsymbol{w}; \boldsymbol{d})^{n-2} \times \det V^{n+p-1,n+q-1}(\boldsymbol{x}^{(k,l)}, \boldsymbol{z}; \boldsymbol{a}^{(k,l)}, \boldsymbol{c}) \det V^{n+r-1,n+s-1}(\boldsymbol{x}^{(k,l)}, \boldsymbol{w}; \boldsymbol{b}^{(k,l)}, \boldsymbol{d})$$

where k' and l' are the indices satisfying $\{k, l, k', l'\} = \{1, 2, 3, 4\}$ and k < l, k' < l', and

$$\begin{aligned} & \operatorname{Pf} A_{1,2,3,4}^{1,2,3,4} \\ &= \frac{1}{\prod_{5 \le i < j \le 2n} (x_j - x_i)} \det V^{p,q}(\boldsymbol{z}; \boldsymbol{c})^{n-3} \det V^{r,s}(\boldsymbol{w}; \boldsymbol{d})^{n-3} \\ & \times \det V^{n+p-2,n+q-2}(\boldsymbol{x}^{(1,2,3,4)}, \boldsymbol{z}; \boldsymbol{a}^{(1,2,3,4)}, \boldsymbol{c}) \det V^{n+r-2,n+s-2}(\boldsymbol{x}^{(1,2,3,4)}, \boldsymbol{w}; \boldsymbol{b}^{(1,2,3,4)}, \boldsymbol{d}). \end{aligned}$$

Hence, by applying (2.5) and cancelling the common factors, we see that, in order to prove (1.4), it suffices to show

$$\begin{split} & \frac{\det V^{n+p-1,n+q-1}(\boldsymbol{x}^{(1,2)},\boldsymbol{z};\boldsymbol{a}^{(1,2)},\boldsymbol{c})\det V^{n+r-1,n+s-1}(\boldsymbol{x}^{(1,2)},\boldsymbol{w};\boldsymbol{b}^{(1,2)},\boldsymbol{d})}{x_2 - x_1} \\ & \times \frac{\det V^{n+p-1,n+q-1}(\boldsymbol{x}^{(3,4)},\boldsymbol{z};\boldsymbol{a}^{(3,4)},\boldsymbol{c})\det V^{n+r-1,n+s-1}(\boldsymbol{x}^{(3,4)},\boldsymbol{w};\boldsymbol{b}^{(3,4)},\boldsymbol{d})}{x_4 - x_3} \\ & - \frac{\det V^{n+p-1,n+q-1}(\boldsymbol{x}^{(1,3)},\boldsymbol{z};\boldsymbol{a}^{(1,3)},\boldsymbol{c})\det V^{n+r-1,n+s-1}(\boldsymbol{x}^{(1,3)},\boldsymbol{w};\boldsymbol{b}^{(1,3)},\boldsymbol{d})}{x_3 - x_1} \\ & \times \frac{\det V^{n+p-1,n+q-1}(\boldsymbol{x}^{(2,4)},\boldsymbol{z};\boldsymbol{a}^{(2,4)},\boldsymbol{c})\det V^{n+r-1,n+s-1}(\boldsymbol{x}^{(2,4)},\boldsymbol{w};\boldsymbol{b}^{(2,4)},\boldsymbol{d})}{x_4 - x_2} \\ & + \frac{\det V^{n+p-1,n+q-1}(\boldsymbol{x}^{(1,4)},\boldsymbol{z};\boldsymbol{a}^{(1,4)},\boldsymbol{c})\det V^{n+r-1,n+s-1}(\boldsymbol{x}^{(1,4)},\boldsymbol{w};\boldsymbol{b}^{(1,4)},\boldsymbol{d})}{x_4 - x_1} \\ & \times \frac{\det V^{n+p-1,n+q-1}(\boldsymbol{x}^{(2,3)},\boldsymbol{z};\boldsymbol{a}^{(2,3)},\boldsymbol{c})\det V^{n+r-1,n+s-1}(\boldsymbol{x}^{(2,3)},\boldsymbol{w};\boldsymbol{b}^{(2,3)},\boldsymbol{d})}{x_3 - x_2} \\ & = \det V^{n+p-2,n+q-2}(\boldsymbol{x}^{(1,2,3,4)},\boldsymbol{z};\boldsymbol{a}^{(1,2,3,4)},\boldsymbol{c})\det V^{n+r-2,n+s-2}(\boldsymbol{x}^{(1,2,3,4)},\boldsymbol{w};\boldsymbol{b}^{(1,2,3,4)},\boldsymbol{d}) \\ & \times \frac{\det V^{n+p,n+q}(\boldsymbol{x},\boldsymbol{z};\boldsymbol{a},\boldsymbol{c})\det V^{n+r,n+s}(\boldsymbol{x},\boldsymbol{w};\boldsymbol{b},\boldsymbol{d})}{\prod_{1 \le i \le j \le 4}(x_j - x_i)}. \end{split}$$

This is equivalent to the identity (2.1) with $\boldsymbol{z}, \boldsymbol{c}, \boldsymbol{w}, \boldsymbol{d}$ replaced by

$$m{z} \leftarrow (m{x}^{(1,2,3,4)}, m{z}), \quad m{c} \leftarrow (m{a}^{(1,2,3,4)}, m{c}), \quad m{w} \leftarrow (m{x}^{(1,2,3,4)}, m{w}), \quad m{d} \leftarrow (m{b}^{(1,2,3,4)}, m{d}),$$

respectively, and it is proven in Proposition 2.1. This completes the proof of (1.4).

Remark 2.4. We can also reduce the proof of the other identities (1.3), (1.5) and (1.6) in Theorem 1.1 to the case of n = 2 with the help of the Desnanot-Jacobi formulae. It is easy to show the case of n = 2 of (1.3) by using the relations in Lemma 2.2 and induction on p + q. Also we can prove (1.5) (resp. (1.6)) in the case of n = 2, by regarding both sides as polynomials in z_p and showing that the values coincide at (2p+3) distinct points $z_1, \dots, z_{p-1}, z_1^{-1}, \dots, z_{p-1}^{-1}x_1, x_2, y_1, y_2$ and -1 (resp. $z_1, \dots, z_{p-1}, z_1^{-1}, \dots, z_{p-1}^{-1}x_1, x_2, x_3, x_4$ and -1) with the help of induction. But, in this paper, we adopt another method, namely, we "homogenize" the identity (1.4) and

derive the other identities from this homogeneous version (3.3).

Proof of the identities (1.3), (1.4) and (1.5)3

In this section, we give a homogeneous version of the identity (1.4), which is shown in the previous section, and derive the identities (1.3), (1.5) and (1.6) from this homogeneous version.

Throughout the remaining of this paper, we use the following notation for vectors $\boldsymbol{x} = (x_1, \cdots, x_n)$ and $y = (y_1, \dots, y_n)$:

$$\boldsymbol{x} + \boldsymbol{y} = (x_1 + y_1, \dots, x_n + y_n), \quad \boldsymbol{x} \boldsymbol{y} = (x_1 y_1, \dots, x_n y_n),$$

and, for integers k and l,

$$\boldsymbol{x}^k = (x_1^k, \dots, x_n^k), \quad \boldsymbol{x}^k \boldsymbol{y}^l = (x_1^k y_1^l, \dots, x_n^k y_n^l).$$

We introduce a homogeneous version of the matrix $V^{p,q}(x; a)$ as follows. For vectors x, y, a, **b** of length *n* and nonnegative integers *p*, *q* with p + q = n, we define a matrix $U^{p,q}\begin{pmatrix} x & a \\ y & b \end{pmatrix}$ to be the $n \times n$ matrix with *i*th row

$$(a_i x_i^{p-1}, a_i x_i^{p-2} y_i, \cdots, a_i y_i^{p-1}, b_i x_i^{q-1}, b_i x_i^{q-2} y_i, \cdots, b_i y_i^{q-1}).$$

Then the following relations between det $V^{p,q}$ and det $U^{p,q}$ are easily shown by elementary transformations, so we omit their proofs.

Lemma 3.1.

$$\det U^{p,q}\begin{pmatrix} \boldsymbol{x} & \boldsymbol{a} \\ \boldsymbol{y} & \boldsymbol{b} \end{pmatrix} = \prod_{k=1}^{p+q} a_k x_k^{p-1} \cdot \det V^{p,q} \left(\boldsymbol{x}^{-1} \boldsymbol{y}; \boldsymbol{a}^{-1} \boldsymbol{b} \boldsymbol{x}^{q-p} \right).$$
(3.1)

In particular,

$$\det V^{p,q}(\boldsymbol{x};\boldsymbol{a}) = \det U^{p,q}\begin{pmatrix} \boldsymbol{1} & | & \boldsymbol{1} \\ \boldsymbol{x} & | & \boldsymbol{a} \end{pmatrix},$$
(3.2)

where 1 = (1, ..., 1).

Now we give a homogeneous version of the identity (1.4).

Theorem 3.2. Let *n* be a positive integer and let *p*, *q*, *r* and *s* be nonnegative integers. Suppose that the vectors $\boldsymbol{x}, \boldsymbol{y}, \boldsymbol{a}, \boldsymbol{b}, \boldsymbol{c}, \boldsymbol{d}$ have length 2n, the vectors $\boldsymbol{\xi}, \boldsymbol{\eta}, \boldsymbol{\alpha}, \boldsymbol{\beta}$ have length p + q, and the vectors $\boldsymbol{\zeta}, \boldsymbol{\omega}, \boldsymbol{\gamma}, \boldsymbol{\delta}$ have length r + s. Then we have

$$\operatorname{Pf}\left(\frac{\det U^{p+1,q+1}\begin{pmatrix}x_{i},x_{j},\boldsymbol{\xi} & a_{i},a_{j},\boldsymbol{\alpha} \\ y_{i},y_{j},\boldsymbol{\eta} & b_{i},b_{j},\boldsymbol{\beta} \end{pmatrix} \det U^{r+1,s+1}\begin{pmatrix}x_{i},x_{j},\boldsymbol{\zeta} & c_{i},c_{j},\boldsymbol{\gamma} \\ y_{i},y_{j},\boldsymbol{\omega} & d_{i},d_{j},\boldsymbol{\delta} \end{pmatrix}}{\det \begin{pmatrix}x_{i} & x_{j} \\ y_{i} & y_{j} \end{pmatrix}}\right)_{1 \leq i < j \leq 2n}$$

$$=\frac{1}{\prod_{1 \leq i < j \leq 2n} \det \begin{pmatrix}x_{i} & x_{j} \\ y_{i} & y_{j} \end{pmatrix}} \det U^{p,q}\begin{pmatrix}\boldsymbol{\xi} & \boldsymbol{\alpha} \\ \boldsymbol{\beta} \end{pmatrix}^{n-1} \det U^{r,s}\begin{pmatrix}\boldsymbol{\zeta} & \boldsymbol{\gamma} \\ \boldsymbol{\omega} & \boldsymbol{\delta} \end{pmatrix}^{n-1}}_{\times \det U^{n+p,n+q}\begin{pmatrix}x,\boldsymbol{\xi} & \boldsymbol{\alpha},\boldsymbol{\alpha} \\ y,\boldsymbol{\eta} & \boldsymbol{b},\boldsymbol{\beta} \end{pmatrix}} \det U^{n+r,n+s}\begin{pmatrix}x,\boldsymbol{\zeta} & c,\boldsymbol{\gamma} \\ y,\boldsymbol{\omega} & d,\boldsymbol{\delta} \end{pmatrix}.$$
 (3.3)

The special case of p = q = r = s = 0 of this identity (3.3) is given by M. Ishikawa [4, Theorem 3.1], and is one of the key ingredients of his proof of Stanley's conjecture.

The identity (1.4) is equivalent to (3.3). It follows from the relation (3.2) that (3.3) specialize to (1.4) by setting

$$oldsymbol{x} = oldsymbol{a} = oldsymbol{c} = oldsymbol{1}_{2n}, \quad oldsymbol{\xi} = oldsymbol{\alpha} = oldsymbol{1}_{p+q}, \quad oldsymbol{\zeta} = oldsymbol{\gamma} = oldsymbol{1}_{r+s}$$

and renaming the variables. As we see in the following proof, we derive (3.3) from (1.4).

Proof of Theorem 3.2. In the identity (1.4), we substitute as follows:

$$\begin{aligned} \boldsymbol{x} \leftarrow \boldsymbol{x}^{-1} \boldsymbol{y}, \quad \boldsymbol{z} \leftarrow \boldsymbol{\xi}^{-1} \boldsymbol{\eta}, \quad \boldsymbol{w} \leftarrow \boldsymbol{\zeta}^{-1} \boldsymbol{\omega}, \\ \boldsymbol{a} \leftarrow \boldsymbol{a}^{-1} \boldsymbol{b} \boldsymbol{x}^{q-p}, \quad \boldsymbol{c} \leftarrow \boldsymbol{\alpha}^{-1} \boldsymbol{\beta} \boldsymbol{\xi}^{q-p}, \quad \boldsymbol{b} \leftarrow \boldsymbol{c}^{-1} \boldsymbol{d} \boldsymbol{x}^{s-r}, \quad \boldsymbol{d} \leftarrow \boldsymbol{\gamma}^{-1} \boldsymbol{\delta} \boldsymbol{\zeta}^{s-r}. \end{aligned}$$
(3.4)

By using the relations (3.1) and

$$\frac{y_j}{x_j} - \frac{y_i}{x_i} = x_i^{-1} x_j^{-1} \det \begin{pmatrix} x_i & x_j \\ y_i & y_j \end{pmatrix},$$

we see that the (i, j) entry of the left-hand side of (1.4) under the substitution (3.4) becomes

$$\frac{\left(a_{i}a_{j}x_{i}^{p}x_{j}^{p}\prod_{k=1}^{p+q}\alpha_{k}\xi_{k}^{p}\right)^{-1}\left(c_{i}c_{j}x_{i}^{r}x_{j}^{r}\prod_{k=1}^{r+s}\gamma_{k}\zeta_{k}^{r}\right)^{-1}}{x_{i}^{-1}x_{j}^{-1}} \times \frac{\det U^{p+1,q+1}\left(x_{i},x_{j},\boldsymbol{\xi} \mid a_{i},a_{j},\boldsymbol{\alpha}\right)}{\det \left(y_{i},y_{j},\boldsymbol{\eta} \mid b_{i},b_{j},\boldsymbol{\beta}\right)}\det U^{r+1,s+1}\left(x_{i},x_{j},\boldsymbol{\zeta} \mid c_{i},c_{j},\boldsymbol{\gamma}\right)}{y_{i},y_{j},\boldsymbol{\omega} \mid d_{i},d_{j},\boldsymbol{\delta}}.$$

Hence, by noting the linearity of Pfaffian

$$\operatorname{Pf}\left(\lambda \alpha_{i} \alpha_{j} a_{ij}\right)_{1 \leq i < j \leq 2n} = \lambda^{n} \prod_{i=1}^{2n} \alpha_{i} \cdot \operatorname{Pf}\left(a_{ij}\right)_{1 \leq i < j \leq 2n},$$

we can see the Pfaffian on the left-hand side of (1.4) becomes

$$\left(\prod_{i=1}^{2n} a_i c_i x_i^{p+r-1} \right)^{-1} \left(\prod_{i=1}^{p+q} \alpha_i \xi_i^p \prod_{i=1}^{r+s} \gamma_i \zeta_i^r \right)^{-n} \\ \times \operatorname{Pf} \left(\frac{\det U^{p+1,q+1} \begin{pmatrix} x_i, x_j, \boldsymbol{\xi} \\ y_i, y_j, \boldsymbol{\eta} \\ \end{bmatrix} \begin{pmatrix} a_i, a_j, \boldsymbol{\alpha} \\ b_i, b_j, \boldsymbol{\beta} \\ b_i, b_j, \boldsymbol{\beta} \\ \end{bmatrix} \det \begin{pmatrix} x_i \cdot x_j \\ y_i \cdot y_j \end{pmatrix} \det U^{r+1,s+1} \begin{pmatrix} x_i, x_j, \boldsymbol{\zeta} \\ y_i, y_j, \boldsymbol{\omega} \\ d_i, d_j, \boldsymbol{\delta} \\ d_i, d_j, \boldsymbol{\delta} \\ \end{bmatrix} \right)_{1 \le i < j \le 2n} .$$

On the other hand, using the relations (3.1) and

$$\prod_{1 \le i < j \le 2n} \left(\frac{y_j}{x_j} - \frac{y_i}{x_i} \right) = \prod_{k=1}^{2n} x_k^{-2n+1} \prod_{1 \le i < j \le 2n} \det \begin{pmatrix} x_i & x_j \\ y_i & y_j \end{pmatrix},$$

we see that the right-hand side of (1.4) becomes

$$\frac{1}{\left(\prod_{k=1}^{2n} x_{k}\right)^{-2n+1}} \cdot \frac{1}{\prod_{1 \leq i < j \leq 2n} \det \begin{pmatrix} x_{i} & x_{j} \\ y_{i} & y_{j} \end{pmatrix}} \times \left(\prod_{i=1}^{p+q} \alpha_{i} \xi_{i}^{p-1}\right)^{-n+1} \left(\prod_{i=1}^{r+s} \gamma_{i} \zeta_{i}^{r-1}\right)^{-n+1} \det U^{p,q} \begin{pmatrix} \boldsymbol{\xi} & \boldsymbol{\alpha} \\ \boldsymbol{\eta} & \boldsymbol{\beta} \end{pmatrix}^{n-1} \det U^{r,s} \begin{pmatrix} \boldsymbol{\zeta} & \boldsymbol{\gamma} \\ \boldsymbol{\omega} & \boldsymbol{\delta} \end{pmatrix}^{n-1} \times \left(\prod_{i=1}^{2n} a_{i} x_{i}^{n+p-1} \prod_{i=1}^{p+q} \alpha_{i} \xi_{i}^{n+p-1}\right)^{-1} \left(\prod_{i=1}^{2n} c_{i} x_{i}^{n+r-1} \prod_{i=1}^{r+s} \gamma_{i} \zeta_{i}^{n+r-1}\right)^{-1} \times \det U^{n+p,n+q} \begin{pmatrix} \boldsymbol{x}, \boldsymbol{\xi} & \boldsymbol{\alpha}, \boldsymbol{\alpha} \\ \boldsymbol{y}, \boldsymbol{\eta} & \boldsymbol{b}, \boldsymbol{\beta} \end{pmatrix} \det U^{n+r,n+s} \begin{pmatrix} \boldsymbol{x}, \boldsymbol{\zeta} & \boldsymbol{c}, \boldsymbol{\gamma} \\ \boldsymbol{y}, \boldsymbol{\omega} & \boldsymbol{d}, \boldsymbol{\delta} \end{pmatrix}.$$

Comparing the both sides and cancelling the common factors, we obtain the desired identity (3.3). \Box

In this setting, a homogeneous version of (1.3) is a direct consequence of (3.3). A key is the following relation between determinant and Pfaffian. If A is any $m \times (2n - m)$ matrix, then we have

$$\operatorname{Pf}\begin{pmatrix} O & A \\ -{}^{t}\!A & O \end{pmatrix} = \begin{cases} (-1)^{n(n-1)/2} \det A & \text{if } m = n, \\ 0 & \text{if } m \neq n. \end{cases}$$
(3.5)

Corollary 3.3. Let *n* be a positive integer and let *p* and *q* be fixed nonnegative integers. For vectors $\boldsymbol{x}, \boldsymbol{y}, \boldsymbol{z}, \boldsymbol{w}, \boldsymbol{a}, \boldsymbol{b}, \boldsymbol{c}, \boldsymbol{d}$ of length *n*, and vectors $\boldsymbol{\xi}, \boldsymbol{\eta}, \boldsymbol{\alpha}, \boldsymbol{\beta}$ of length p + q, we have

$$\det\left(\frac{\det U^{p+1,q+1}\begin{pmatrix}x_i, z_j, \boldsymbol{\xi} & a_i, c_j, \boldsymbol{\alpha}\\y_i, w_j, \boldsymbol{\eta} & b_i, d_j, \boldsymbol{\beta}\end{pmatrix}}{\det\begin{pmatrix}x_i & z_j\\y_i & w_j\end{pmatrix}}\right)_{1 \le i, j \le n}$$
$$= \frac{(-1)^{n(n-1)/2}}{\prod_{1 \le i, j \le n} \det\begin{pmatrix}x_i & z_j\\y_i & w_j\end{pmatrix}} \det U^{p,q}\begin{pmatrix}\boldsymbol{\xi} & \boldsymbol{\alpha}\\\boldsymbol{\beta}\end{pmatrix}^{n-1} \det U^{n+p,n+q}\begin{pmatrix}\boldsymbol{x}, \boldsymbol{z}, \boldsymbol{\xi} & \boldsymbol{a}, \boldsymbol{c}, \boldsymbol{\alpha}\\\boldsymbol{y}, \boldsymbol{w}, \boldsymbol{\eta} & \boldsymbol{b}, \boldsymbol{d}, \boldsymbol{\beta}\end{pmatrix}. \quad (3.6)$$

Proof. In (3.3), we take r = s = 0 and put

$$c_1 = \dots = c_n = 1, \qquad c_{n+1} = \dots = c_{2n} = 0, d_1 = \dots = d_n = 0, \qquad d_{n+1} = \dots = d_{2n} = 1.$$
(3.7)

Under this substitution (3.7), we have

$$\det U^{1,1}\begin{pmatrix} x_i, x_j \\ y_i, y_j \end{pmatrix} \begin{pmatrix} c_i, c_j \\ d_i, d_j \end{pmatrix} = \det \begin{pmatrix} c_i & d_i \\ c_j & d_j \end{pmatrix} = \begin{cases} 0 & \text{if } 1 \le i, j \le n \text{ or } n+1 \le i, j \le 2n, \\ 1 & \text{if } 1 \le i \le n \text{ and } n+1 \le j \le 2n, \\ -1 & \text{if } n+1 \le i \le 2n \text{ and } 1 \le j \le n. \end{cases}$$

Hence, by (3.5), we see that the left-hand side of (3.3) becomes

$$(-1)^{n(n-1)/2} \det \left(\frac{\det U^{p+1,q+1} \begin{pmatrix} x_i, x_{n+j}, \boldsymbol{\xi} & a_i, a_{n+j}, \boldsymbol{\alpha} \\ y_i, y_{n+j}, \boldsymbol{\eta} & b_i, b_{n+j}, \boldsymbol{\beta} \end{pmatrix}}{\det \begin{pmatrix} x_i & x_{n+j} \\ y_i & y_{n+j} \end{pmatrix}} \right)_{1 \le i,j \le n}.$$

On the other hand, under the specialization (3.7), we have

$$\det U^{n,n}\begin{pmatrix} \boldsymbol{x} & \boldsymbol{c} \\ \boldsymbol{y} & \boldsymbol{d} \end{pmatrix} = \det \begin{pmatrix} \begin{pmatrix} x_i^{n-j}y_i^{j-1} \end{pmatrix}_{1 \le i,j \le n} & O \\ O & \begin{pmatrix} x_{n-j}^{n-j}y_{n+i}^{j-1} \end{pmatrix}_{1 \le i,j \le n} \end{pmatrix}$$
$$= \prod_{1 \le i < j \le n} \det \begin{pmatrix} x_i & x_j \\ y_i & y_j \end{pmatrix} \prod_{n+1 \le i < j \le 2n} \det \begin{pmatrix} x_i & x_j \\ y_i & y_j \end{pmatrix}.$$

Thus the right-hand side of (3.3) becomes

$$\frac{1}{\prod_{1\leq i,j\leq n}\det\begin{pmatrix}x_i & x_{n+j}\\y_i & y_{n+j}\end{pmatrix}}\det U^{p,q}\begin{pmatrix}\boldsymbol{\xi} & \boldsymbol{\alpha}\\\boldsymbol{\eta} & \boldsymbol{\beta}\end{pmatrix}^{n-1}\det U^{n+p,n+q}\begin{pmatrix}x,\boldsymbol{\xi} & \boldsymbol{\alpha},\boldsymbol{\alpha}\\y,\boldsymbol{\eta} & \boldsymbol{b},\boldsymbol{\beta}\end{pmatrix}.$$

Lastly, if we replace the variables as $x_{n+i} = z_i$, $y_{n+i} = w_i$, $a_{n+i} = c_i$ and $b_{n+i} = d_i$ for $1 \le i \le n$, then we obtain the desired identity (3.6). This completes our proof.

Now it is easy to derive (1.3) from (3.5) by using the relation (3.1). To prove the remaining identities (1.5) and (1.6) in Theorem 1.1, we need the following lemma.

Lemma 3.4. Let n be a nonnegative integer.

(1) For vectors \boldsymbol{x} , \boldsymbol{a} and $\boldsymbol{1} = (1, \dots, 1)$ of length 2n, we have

$$\det U^{n,n}\begin{pmatrix} \boldsymbol{x} & | \boldsymbol{1} + \boldsymbol{a}\boldsymbol{x} \\ \boldsymbol{1} + \boldsymbol{x}^2 & | \boldsymbol{x} + \boldsymbol{a} \end{pmatrix} = (-1)^{n(n-1)/2} \det W^{2n}(\boldsymbol{x}; \boldsymbol{a}).$$
(3.8)

(2) For vectors \boldsymbol{x} , \boldsymbol{a} and $\boldsymbol{1} = (1, \dots, 1)$ of length 2n + 1, we have

$$\det U^{n,n+1}\begin{pmatrix} x \\ 1+x^2 \\ 1+a \end{pmatrix} = (-1)^{n(n-1)/2} \det W^{2n+1}(x;a).$$
(3.9)

Proof. By definition, we have

$$\det U^{p,q} \begin{pmatrix} \mathbf{x} & | & \mathbf{c} \\ \mathbf{1} + \mathbf{x}^2 & | & \mathbf{d} \end{pmatrix}$$

=
$$\det \left(\begin{cases} c_i x_i^{p-j} (1 + x_i^2)^{j-1} & \text{if } 1 \le i \le p+q \text{ and } 1 \le j \le p, \\ d_i x_i^{p+q-j} (1 + x_i^2)^{j-p-1} & \text{if } 1 \le i \le p+q \text{ and } p+1 \le j \le p+q. \end{cases} \right).$$

By performing appropriate elementary column transformations, one can see that this determinant is equal to

$$\det \begin{pmatrix} c_i x_i^{p-1} & \text{if } 1 \le i \le p+q \text{ and } j=1, \\ c_i x_i^{p-j} \left(1+x_i^{2(j-1)}\right) & \text{if } 1 \le i \le p+q \text{ and } 2 \le j \le p, \\ d_i x_i^{q-1} & \text{if } 1 \le i \le p+q \text{ and } j=p+1, \\ d_i x_i^{p+q-j} \left(1+x_i^{2(j-p-1)}\right) & \text{if } 1 \le i \le p+q \text{ and } p+2 \le j \le p+q. \end{pmatrix}.$$
(3.10)

(1) First put q = p = n, $c_i = 1 + a_i x_i$ and $d_i = x_i + a_i$ for $1 \le i \le 2n$. Then the above determinant (3.10) is equal to

$$\det \begin{pmatrix} x_i^{n-1} + a_i x_i^n & \text{if } 1 \le i \le 2n \text{ and } j = 1, \\ x_i^{n-j} + a_i x_i^{n+j-1} + x_i^{n+j-2} + a_i x_i^{n-j+1} & \text{if } 1 \le i \le 2n \text{ and } 2 \le j \le n, \\ x_i^n + a_i x_i^{n-1} & \text{if } 1 \le i \le 2n \text{ and } j = n+1, \\ x_i^{2n-j+1} + a_i x_i^{j-2} + x_i^{j-1} + a_i x_i^{2n-j} & \text{if } 1 \le i \le 2n \text{ and } n+2 \le j \le 2n. \end{pmatrix}.$$

Subtract the first column from the (n + 2)th column and subtract the (n + 1)th column from the second column, then subtract the second column from the (n + 3)th column and subtract the (n + 2)th column from the third column, and so on. We continue these elementary column transformations until we obtain

$$\det\left(\begin{cases} x_i^{n-j} + a_i x_i^{n+j-1} & \text{if } 1 \le i \le 2n \text{ and } 1 \le j \le n, \\ x_i^{j-1} + a_i x_i^{2n-j} & \text{if } 1 \le i \le 2n \text{ and } n+1 \le j \le 2n. \end{cases}\right),$$

which is $(-1)^{n(n-1)/2} \det W^{2n}(x; a)$.

(2) Next we take p = n, q = n + 1, $c_i = 1 + a_i x_i^2$ and $d_i = 1 + a_i$ $(1 \le i \le 2n + 1)$ in (3.10), then we obtain

$$\det \begin{pmatrix} x_i^{n-1} + a_i x_i^{n+1} & \text{if } 1 \le i \le 2n+1 \text{ and } j = 1, \\ x_i^{n-j} + a_i x_i^{n+j} + x_i^{n+j-2} + a_i x_i^{n-j+2} & \text{if } 1 \le i \le 2n+1 \text{ and } 2 \le j \le n, \\ x_i^n + a_i x_i^n & \text{if } 1 \le i \le 2n+1 \text{ and } j = n+1, \\ x_i^{2n+1-j} + a_i x_i^{j-1} + x_i^{j-1} + a_i x_i^{2n+1-j} & \text{if } 1 \le i \le 2n+1 \text{ and } n+2 \le j \le 2n+1. \end{pmatrix}$$

We subtract the first column from the (n+2)th column and subtract the (n+1)th column from the second column, and then subtract the second column from the (n+3)th column and subtract the (n+2)th column from the third column, and so on. We continue these elementary transformations until we obtain

$$\det\left(\begin{cases} x_i^{n-i} + a_i x_i^{n+j} & \text{if } 1 \le i \le 2n+1 \text{ and } 1 \le j \le n, \\ x_i^{j-1} + a_i x_i^{2n+1-j} & \text{if } 1 \le i \le 2n+1 \text{ and } n+1 \le j \le 2n+1. \end{cases}\right),$$

which is equal to $(-1)^{n(n-1)/2} \det W^{2n+1}(\boldsymbol{x}; \boldsymbol{a})$. This completes our proof.

Now we can finish our proof of Theorem 1.1.

Proof of the identities (1.3), (1.5) and (1.6) in Theorem 1.1. As we mentioned before, the identity (1.3) follows from (3.6) by virtue of (3.1).

We derive (1.6) from (3.3). First we consider the case where both p = 2l and q = 2m are even. In (3.3), we take p = q = l and r = s = m, and perform the following substitutions:

By using the relation

$$\det \begin{pmatrix} x_i & x_j \\ 1 + x_i^2 & 1 + x_j^2 \end{pmatrix} = (x_i - x_j)(1 - x_i x_j),$$

and (3.8) in Lemma 3.4, we see that the identity (3.3) becomes

$$\begin{split} & \operatorname{Pf}\left(\frac{(-1)^{l(l+1)/2+m(m+1)/2} \det W^{2l+2}(x_i, x_j, \boldsymbol{z}; a_i, a_j, \boldsymbol{c}) \det W^{2m+2}(x_i, x_j, \boldsymbol{w}; b_i, b_j, \boldsymbol{d})}{(x_i - x_j)(1 - x_i x_j)}\right)_{1 \leq i < j \leq 2n} \\ & = (-1)^{(n-1)\binom{l}{2} + (n-1)\binom{m}{2} + \binom{n+l}{2} + \binom{n+m}{2}} \\ & \times \frac{1}{\prod_{1 \leq i < j \leq 2n} (x_i - x_j)(1 - x_i x_j)} \\ & \times \det W^{2l}(\boldsymbol{z}; \boldsymbol{c})^{n-1} \det W^{2m}(\boldsymbol{w}; \boldsymbol{d})^{n-1} \det W^{2n+2l}(\boldsymbol{x}, \boldsymbol{z}; \boldsymbol{a}, \boldsymbol{c}) \det W^{2n+2m}(\boldsymbol{x}, \boldsymbol{w}; \boldsymbol{b}, \boldsymbol{d}). \end{split}$$

Since we have

$$\begin{array}{l} (n-1)l(l-1)/2 + (n-1)m(m-1)/2 + (n+l)(n+l-1)/2 + (n+m)(n+m-1)/2 \\ \\ = n\left\{l(l+1)/2 + m(m+1)/2\right\} + n(n-1), \end{array}$$

we obtain the identity (1.6) when p and q are both even. We can prove the other cases similarly by using (3.8) or (3.9) according as p and q are even or odd.

Also the remaining identity (1.5) can be derived from (3.6) by using (3.8) or (3.9). The details are left to the reader.

4 A variation of the determinant and Pfaffian identities

In this section, we give a variation of the identities in Theorem 1.1, which can be regarded as a generalization of an identity of T. Sundquist [19]. This variation is proposed by one of the authors.

Let *n* be a positive integer and let *p* and *q* be nonnegative integers with p + q = n. Let $\boldsymbol{x} = (x_1, \dots, x_n)$ and $\boldsymbol{a} = (a_1, \dots, a_n)$ be vectors of variables. For partitions λ and μ with $l(\lambda) \leq p$ and $l(\mu) \leq q$, we define a matrix $V_{\lambda,\mu}^{p,q}(\boldsymbol{x};\boldsymbol{a})$ to be the $n \times n$ matrix with *i*th row

$$(x_i^{\lambda_p}, x_i^{\lambda_{p-1}+1}, x_i^{\lambda_{p-2}+2}, \cdot, x_i^{\lambda_1+p-1}, a_i x_i^{\mu_q}, a_i x_i^{\mu_{q-1}+1}, a_i x_i^{\mu_{q-2}+2}, \cdots, a_i x_i^{\mu_1+q-1}).$$

For example, if $\lambda = \mu = \emptyset$, then we have $V_{\emptyset,\emptyset}^{p,q}(\boldsymbol{x};\boldsymbol{a}) = V^{p,q}(\boldsymbol{x};\boldsymbol{a})$. Let P_n denote the set of integer partitions of the form $(\alpha_1, \ldots, \alpha_r | \alpha_1 + 1, \ldots, \alpha_r + 1)$ in the Frobenius notation with $\alpha_1 + 2 \leq n$. (So $n \geq 2$). We define

$$F^{p,q}(\boldsymbol{x};\boldsymbol{a}) = \sum_{\lambda \in \mathcal{P}_p, \ \mu \in \mathcal{P}_q} (-1)^{(|\lambda|+|\mu|)/2} \det V^{p,q}_{\lambda,\mu}(\boldsymbol{x};\boldsymbol{a}).$$

For example, if p = q = 1, then $F^{1,1}(x; a) = a_2 - a_1$, and, if p = q = 2, then $\mathcal{P}_2 = \{\emptyset, (1, 1)\}$ and

$$F^{2,2}(\boldsymbol{x};\boldsymbol{a}) = \det \begin{pmatrix} 1 & x_1 & a_1 & a_1 x_1 \\ 1 & x_2 & a_2 & a_2 x_2 \\ 1 & x_3 & a_3 & a_3 x_3 \\ 1 & x_4 & a_4 & a_4 x_4 \end{pmatrix} - \det \begin{pmatrix} x_1 & x_1^2 & a_1 & a_1 x_1 \\ x_2 & x_2^2 & a_2 & a_2 x_2 \\ x_3 & x_3^2 & a_3 & a_3 x_3 \\ x_4 & x_4^2 & a_4 & a_4 x_4 \end{pmatrix} - \det \begin{pmatrix} 1 & x_1 & a_1 x_1 & a_1 x_1^2 \\ 1 & x_2 & a_2 x_2 & a_2 x_2^2 \\ 1 & x_3 & a_3 x_3 & a_3 x_3^2 \\ 1 & x_4 & a_4 x_4 & a_4 x_4^2 \end{pmatrix} + \det \begin{pmatrix} x_1 & x_1^2 & a_1 x_1 & a_1 x_1^2 \\ x_2 & x_2^2 & a_2 x_2 & a_2 x_2^2 \\ x_3 & x_3^2 & a_3 x_3 & a_3 x_3^2 \\ x_4 & x_4^2 & a_4 x_4 & a_4 x_4^2 \end{pmatrix}$$

The aim of this section is to prove the following theorem:

Theorem 4.1. (a) Let n be a positive integer and let p and q be nonnegative integers. For six vectors of variables

$$x = (x_1, \dots, x_n), \quad y = (y_1, \dots, y_n), \quad z = (z_1, \dots, z_{p+q}),$$

 $a = (a_1, \dots, a_n), \quad b = (b_1, \dots, b_n), \quad c = (c_1, \dots, c_{p+q})$

we have

$$\det\left(\frac{F^{p+1,q+1}(x_i, y_j, \boldsymbol{z}; a_i, b_j, \boldsymbol{c})}{(y_j - x_i)(1 - x_i y_j)}\right)_{1 \le i, j \le n} = \frac{(-1)^{n(n-1)/2}}{\prod_{i,j=1}^n (y_j - x_i)(1 - x_i y_j)} F^{p,q}(\boldsymbol{z}; \boldsymbol{c})^{n-1} F^{n+p,n+q}(\boldsymbol{x}, \boldsymbol{y}, \boldsymbol{z}; \boldsymbol{a}, \boldsymbol{b}, \boldsymbol{c}).$$
(4.1)

(b) Let n be a positive integer and let p, q, r, s be nonnegative integers. For seven vectors of variables

we have

$$Pf\left(\frac{F^{p+1,q+1}(x_i, x_j, \boldsymbol{z}; a_i, a_j, \boldsymbol{c})F^{r+1,s+1}(x_i, x_j, \boldsymbol{w}; b_i, b_j, \boldsymbol{d})}{(x_j - x_i)(1 - x_i x_j)}\right)_{1 \le i,j \le 2n} = \frac{1}{\prod_{1 \le i < j \le 2n} (x_j - x_i)(1 - x_i x_j)}}F^{p,q}(\boldsymbol{z}; \boldsymbol{c})^{n-1}F^{r,s}(\boldsymbol{w}; \boldsymbol{d})^{n-1} \times F^{n+p,n+q}(\boldsymbol{x}, \boldsymbol{z}; \boldsymbol{a}, \boldsymbol{c})F^{n+r,n+s}(\boldsymbol{x}, \boldsymbol{w}; \boldsymbol{b}, \boldsymbol{d}).$$
(4.2)

In particular, by putting p = q = r = s = 0 and $b_i = x_i$ for $1 \le i \le 2n$ in (4.2), we obtain Sundquist's identity [19, Theorem 2.1].

Corollary 4.2. (Sundquist)

$$\Pr\left(\frac{a_j - a_i}{1 - x_i x_j}\right)_{1 \le i < j \le 2n} = \frac{(-1)^{n(n-1)/2}}{\prod_{1 \le i < j \le 2n} (1 - x_i x_j)} \sum_{\lambda, \mu \in \mathcal{P}_n} (-1)^{(|\lambda| + |\mu|)/2} \det V_{\lambda, \mu}^{n, n}(\boldsymbol{x}; \boldsymbol{a}).$$
(4.3)

In order to prove Theorem 4.1 and Corollary 4.2, we need a relation between $F^{p,q}(\boldsymbol{x}; \boldsymbol{a})$ and det $V^{p,q}(\boldsymbol{y}; \boldsymbol{b})$.

Proposition 4.3. We have

$$F^{p,q}(\boldsymbol{x};\boldsymbol{a}) = (-1)^{\binom{p}{2} + \binom{q}{2}} \prod_{i=1}^{p+q} x_i^{p-1} \cdot \det V^{p,q}(\boldsymbol{x} + \boldsymbol{x}^{-1}; \boldsymbol{a}\boldsymbol{x}^{q-p}),$$

$$= (-1)^{\binom{p}{2} + \binom{q}{2}} \det U^{p,q} \begin{pmatrix} \boldsymbol{x} & | \mathbf{1} \\ \mathbf{1} + \boldsymbol{x}^2 & | \mathbf{a} \end{pmatrix},$$
(4.4)

where

$$\boldsymbol{x} + \boldsymbol{x}^{-1} = (x_1 + x_1^{-1}, \cdots, x_{p+q} + x_{p+q}^{-1}),$$
$$\boldsymbol{a} \boldsymbol{x}^{q-p} = (a_1 x_1^{q-p}, \cdots, a_{p+q} x_{p+q}^{q-p}),$$
$$\boldsymbol{1} + \boldsymbol{x}^2 = (1 + x_1^2, \cdots, 1 + x_{p+q}^2).$$

Here we give a proof by using the Cauchy-Binet formula. Let $A = (a_{ij})_{1 \le i \le m, 1 \le j \le n}$ be an m by n matrix. For any subsets $I = \{i_1 < \cdots < i_r\} \subset [m]$, and $J = \{j_1 < \cdots < j_r\} \subset [n]$, let $\Delta_J^I(A)$ denote the submatrix obtained by selecting the rows indexed by I and the columns indexed by J. If all rows or columns are selected, i.e., if I = [m] or J = [n], then we simply write $\Delta_J(A)$ or $\Delta^I(A)$ for $\Delta_J^{[m]}(A)$ or $\Delta_{[n]}^I(A)$.

Lemma 4.4. Let X and Y be any $n \times N$ matrix and A be any $N \times N$ matrix. Then we have

$$\det(XA^{t}Y) = \sum_{I,J} \det \Delta_{J}^{I}(A) \det \Delta_{I}(X) \det \Delta_{J}(Y), \qquad (4.5)$$

where the sum is taken over all pairs (I, J) of *n*-element subsets of [N].

For a partition λ with length $\leq r$, we put

$$I(\lambda) = \{\lambda_r, \lambda_{r-1} + 1, \lambda_{r-2} + 2, \cdots, \lambda_1 + r - 1\}.$$

Then a key of the proof of Proposition 4.3 is is the following lemma.

Lemma 4.5. Let D_r be the following $r \times (2r-1)$ matrix with columns indexed by $0, 1, \dots, 2r-2$:

$$D_r = \begin{pmatrix} 0 & r-2 & r-1 & r & 2r-2 \\ & 1 & & \\ & 1 & 1 & \\ & \ddots & & \ddots & \\ 1 & & & & 1 \end{pmatrix}.$$

Then the minor of D_r corresponding to a partition λ is given by

$$\det \Delta_{I(\lambda)} (D_r) = \begin{cases} (-1)^{r(r-1)/2 + |\lambda|/2} & \text{if } \lambda \in \mathcal{P}_r, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. First we show that det $\Delta_{I(\lambda)}(D_r) = 0$ unless $\lambda \in \mathcal{P}_r$. Suppose that det $\Delta_{I(\lambda)}(D_r) \neq 0$. Since the first row of the matrix D_r has only 1 in the (r-1)th column, we must have $r-1 \in I(\lambda)$. If we denote by $p = p(\lambda)$ the length of the main diagonal of the Young diagram of λ , then we have

$$p = \#\{i : \lambda_i \ge i\} = \#\{i : \lambda_i + r - i \ge r\} = \#\{k \in I(\lambda) : k \ge r\}.$$

Hence the elements $\lambda_1 + r - 1 > \cdots > \lambda_p + r - p$ are the largest p elements belonging in $I(\lambda)$. Since the Frobenius' Lemma ([12, (1.7)]) says that

$$\{\lambda_i + r - i : 1 \le i \le r\} \cup \{r - 1 + j - {}^t\lambda_j : 1 \le j \le r - 1\} = \{0, 1, \cdots, 2r - 2\},\$$

we see that the elements $r - t\lambda_1 < \cdots < r - 1 + p - t\lambda_p$ are the smallest p elements not belonging in $I(\lambda)$. On the other hand, by noting that the kth column of D_r is identical with the (2r - 2 - k)th column, we have, if $k \neq r - 1$, then $k \in I(\lambda)$ if and only if $2r - 2 - k \notin I(\lambda)$. Therefore we have

$$\lambda_i + r - i + (r - 1 + i - {}^t\lambda_i) = 2r - 2 \quad (1 \le i \le p).$$

So we have ${}^{t}\lambda_{i} = \lambda_{i} + 1$ for $1 \leq i \leq p$, which implies $\lambda \in \mathcal{P}_{r}$.

Next we show that, if $\lambda = (\alpha | \alpha + 1) \in \mathcal{P}_r$, then det $\Delta_{I(\lambda)}(D_r) = (-1)^{r(r-1)/2 + |\lambda|/2} = (-1)^{r(r-1)/2 + |\alpha| + p}$. Note that $\Delta_{I(\lambda)}(D_r)$ is a permutation matrix. Let σ be the permutation corresponding to $\Delta_{I(\lambda)}(D_r)$. Then we have

$$\sigma(1) > \sigma(2) > \dots > \sigma(r-p-1),$$

$$\sigma(r-p) = 1, \quad \sigma(r-p+1) = \alpha_p + 2, \quad \dots, \quad \sigma(r) = \alpha_1 + 2.$$

The number of pairs (i, j) such that i < j and $\sigma(i) < \sigma(j)$ is equal to

$$(\alpha_1 + 1) + \dots + (\alpha_p + 1) = |\alpha| + p,$$

so we have

$$\det \Delta_{I(\lambda)} \left(D_r \right) = \operatorname{sgn}(\sigma) = (-1)^{r(r-1)/2 - |\alpha| - p}.$$

Proof of Proposition 4.3. In this proof we put m = p + q for brevity. Apply the Cauchy-Binet formula (4.4) to the following $(p + q) \times (2p + 2q - 2)$ matrices X and Y (and the identity matrix A):

$$X = \begin{pmatrix} D_p & O \\ O & D_q \end{pmatrix}, \quad Y = \begin{pmatrix} 1 & x_1 & \cdots & x_1^{2p-2} & a_1 & a_1x_1 & \cdots & a_1x_1^{2q-2} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & x_m & \cdots & x_m^{2p-2} & a_m & a_mx_m & \cdots & a_mx_m^{2q-2} \end{pmatrix}$$

Let $C \cup C' = \{0, 1, \dots, 2p - 2\} \cup \{0', 1', \dots, (2q - 2)'\}$ be the set indexing the columns of X and Y. Let I be a subset of $C \cup C'$ which has cardinality p + q and consider the minors of X and Y obtained by choosing the columns with indices in I. Then we have

det
$$\Delta_I(X) = 0$$
 unless $\#(I \cap C) = p$ and $\#(I \cap C') = q$.

Suppose that $\#(I \cap C) = p$ and $\#(I \cap C') = q$ and that the subsets $I \cap C$ and $I \cap C'$ determine partitions λ and μ , i.e., $I(\lambda) = I \cap C$ and $I(\mu) = I \cap C'$. Then, by Lemma 4.5 and the definition of $V^{p,q}_{\lambda,\mu}(\boldsymbol{x};\boldsymbol{a})$, we have

$$\det \Delta_I (X) = \begin{cases} (-1)^{\binom{p}{2} + \binom{q}{2} + |\lambda|/2 + |\mu|/2} & \text{if } \lambda \in \mathcal{P}_p \text{ and } \mu \in \mathcal{P}_q, \\ 0 & \text{otherwise,} \end{cases}$$
$$\det \Delta_I (Y) = \det V^{p,q}_{\lambda,\mu}(\boldsymbol{x}; \boldsymbol{a}).$$

Hence the Cauchy-Binet formula gives

$$\det(X^{t}Y) = \sum_{I \subset C \cup C'} \det \Delta_{I}(X) \det \Delta_{I}(Y)$$
$$= \sum_{\lambda \in \mathcal{P}_{p}, \ \mu \in \mathcal{P}_{q}} (-1)^{\binom{p}{2} + \binom{q}{2} + |\lambda|/2 + |\mu|/2} \det V_{\lambda,\mu}^{p,q}(\boldsymbol{x}; \boldsymbol{a}).$$

On the other hand, we have

$$\det(X^tY)$$

$$= \det \begin{pmatrix} x_1^{p-1} & x_1^p + x_1^{p-2} & \cdots & x_1^{2p-2} + 1 & a_1 x_1^{q-1} & a_1 (x_1^q + x_1^{q-2}) & \cdots & a_1 (x_1^{2q-2} + 1) \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ x_m^{p-1} & x_m^p + x_m^{p-2} & \cdots & x_m^{2p-2} + 1 & a_m x_m^{q-1} & a_m (x_m^q + x_m^{q-2}) & \cdots & a_m (x_m^{2q-2} + 1) \end{pmatrix}.$$

Now, by applying elementary transformations and by using the relations

$$(x+x^{-1})^{2k} = \sum_{i=0}^{k-1} \binom{2k}{i} (x^{2k-2i} + x^{-2k+2i}) + \binom{2k}{k},$$
$$(x+x^{-1})^{2k+1} = \sum_{i=0}^{k} \binom{2k+1}{i} (x^{2k-2i+1} + x^{-2k+2i-1}),$$

we have

$$\det(X^{t}Y) = \prod_{i=1}^{m} x_{i}^{p-1} \cdot \det\left(\begin{cases} (x_{i} + x_{i}^{-1})^{j-1} & \text{if } 1 \leq i \leq m \text{ and } 1 \leq j \leq p, \\ a_{i}x_{i}^{q-p}(x_{i} + x_{i}^{-1})^{j-p-1} & \text{if } 1 \leq i \leq m \text{ and } p+1 \leq j \leq p+q. \end{cases}\right)$$
$$= \prod_{i=1}^{m} x_{i}^{p-1} \cdot \det V^{p,q}(\boldsymbol{x} + \boldsymbol{x}^{-1}; \boldsymbol{a}\boldsymbol{x}^{q-p})$$
$$= U^{p,q}\begin{pmatrix} \boldsymbol{x} & | & \mathbf{1} \\ \mathbf{1} + \boldsymbol{x}^{2} & | & \mathbf{a} \end{pmatrix}.$$

Remark 4.6. By the above argument in the case of q = 0, we actually show one of the Littlewood's formula

$$\sum_{\lambda \in \mathcal{P}_n} s_{\lambda}(x_1, \cdots, x_n) = \prod_{1 \le i < j \le n} (1 - x_i x_j).$$

Now we are in position to derive Theorem 4.1 from Theorem 3.2.

Proof of Theorem 4.1. Note that

$$\det \begin{pmatrix} x & 1+x^2 \\ y & 1+y^2 \end{pmatrix} = (x-y)(1-xy).$$
(4.6)

First we prove (4.2). From the above relation (4.6) and (4.4), we have

$$\frac{F^{p+1,q+1}(x_i, x_j, \boldsymbol{z}; a_i, a_j, \boldsymbol{c})F^{r+1,s+1}(x_i, x_j, \boldsymbol{w}; b_i, b_j, \boldsymbol{d})}{(x_j - x_i)(1 - x_i x_j)} = \frac{(-1)^{1+\binom{p+1}{2} + \binom{q+1}{2} + \binom{r+1}{2} + \binom{r+1}{2}}{\det \begin{pmatrix} x_i & 1 + x_i^2 \\ x_j & 1 + x_j^2 \end{pmatrix}} \times \det U^{p+1,q+1} \begin{pmatrix} x_i, x_j, \boldsymbol{z} \\ 1 + x_i^2, 1 + x_j^2, \boldsymbol{1} + \boldsymbol{z}^2 & a_i, a_j, \boldsymbol{c} \end{pmatrix} \times \det U^{r+1,s+1} \begin{pmatrix} x_i, x_j, \boldsymbol{w} \\ 1 + x_i^2, 1 + x_j^2, \boldsymbol{1} + \boldsymbol{w}^2 & b_i, b_j, \boldsymbol{d} \end{pmatrix}.$$

Hence we apply (3.6) to obtain

$$\begin{split} & \operatorname{Pf}\left(\frac{F^{p+1,q+1}(x_i, x_j, \boldsymbol{z}; a_i, a_j, \boldsymbol{c})F^{r+1,s+1}(x_i, x_j, \boldsymbol{w}; b_i, b_j, \boldsymbol{d})}{(x_j - x_i)(1 - x_i x_j)}\right)_{1 \leq i,j \leq 2n} \\ &= \frac{(-1)^{n+n\binom{p+1}{2} + n\binom{q+1}{2} + n\binom{r+1}{2} + n\binom{s+1}{2}}{\prod_{1 \leq i < j \leq 2n} \det \begin{pmatrix} x_i & 1 + x_i^2 \\ x_j & 1 + x_j^2 \end{pmatrix}}}{(x_j - 1 + z^2) \operatorname{extrm{def}} \left(\begin{array}{c} \boldsymbol{x} \\ \boldsymbol{x}_j \\$$

If we use the relation

$$n\binom{p+1}{2} - (n-1)\binom{p}{2} - \binom{p+n}{2} = -\binom{n}{2},$$

then we obtain the desired identity. This proves (4.2).

The determinant identity (4.1) can be proven by the same method by using (4.6), (4.4) and (3.3), so we omit the detailed proof.

Proof of Corollary 4.2. If we put p = q = r = s = 0 and $b_i = x_i$ $(1 \le i \le 2n)$ in (4.2), then we have

$$\Pr\left(\frac{a_j - a_i}{1 - x_i x_j}\right)_{1 \le i, j \le 2n} = \frac{1}{\prod_{1 \le i < j \le 2n} (x_j - x_i)(1 - x_i x_j)} F^{n,n}(\boldsymbol{x}; \boldsymbol{a}) F^{n,n}(\boldsymbol{x}; \boldsymbol{x}).$$

From the relation (4.4), we see that

$$F^{n,n}(\boldsymbol{x};\boldsymbol{x}) = \prod_{i=1}^{2n} x_i^{n-1} \det V^{n,n}(\boldsymbol{x}+\boldsymbol{x}^{-1};\boldsymbol{x}).$$

And, by applying appropriate elementary column transformations, we obtain

$$F^{n,n}(\boldsymbol{x};\boldsymbol{x}) = (-1)^{n(n-1)/2} \det\left(x_i^{j-1}\right)_{1 \le i,j \le 2n} = (-1)^{n(n-1)/2} \prod_{1 \le i < j \le 2n} (x_j - x_i).$$

This completes the proof of the corollary.

In Theorem 4.1, we can replace $F^{p,q}(\boldsymbol{x};\boldsymbol{a})$ by the following linear combination of det $V^{p,q}_{\lambda,\mu}(\boldsymbol{x};\boldsymbol{a})$:

$$\begin{split} G^{p,q}(\boldsymbol{x};\boldsymbol{a}) &= \sum_{\lambda \in \mathcal{Q}_p, \, \mu \in \mathcal{Q}_q} (-1)^{(|\lambda|+|\mu|)/2} \det V^{p,q}_{\lambda,\mu}(\boldsymbol{x};\boldsymbol{a}), \\ H^{p,q}(\boldsymbol{x};\boldsymbol{a}) &= \sum_{\lambda \in \mathcal{R}_p, \, \mu \in \mathcal{R}_q} (-1)^{(|\lambda|+p(\lambda)+|\mu|+p(\mu))/2} \det V^{p,q}_{\lambda,\mu}(\boldsymbol{x};\boldsymbol{a}), \end{split}$$

where Q_n (resp. \mathcal{R}_n) is the set of partitions λ with length $\leq n$ which is of the form $\lambda = (\alpha + 1|\alpha)$ (resp. $\lambda = (\alpha|\alpha)$) in the Frobenius notation.

The following Lemma and Proposition can be proven by the same idea as Lemma 4.5 and Proposition 4.3, so we leave the proof to the reader.

Lemma 4.7. Let C_r (resp. B_r) be the $r \times (2r+1)$ (resp. $r \times 2r$) matrix given by

$$B_r = \begin{pmatrix} 0 & r-2 & r-1 & r & r+1 & 2r-1 \\ & -1 & 1 & & \\ & \ddots & & & \ddots & \\ -1 & & & & 1 \end{pmatrix},$$
$$B_r = \begin{pmatrix} -1 & 1 & & \\ & \ddots & & & \ddots & \\ 0 & r-2 & r-1 & r & r+1 & r+2 & 2r \\ & -1 & 0 & 1 & & \\ & & -1 & & 1 & \\ & \ddots & & & & \ddots & \\ -1 & & & & & 1 \end{pmatrix}.$$

Then we have

(1) For a partition λ of length $\leq r$, we have

$$\det \Delta_{I(\lambda)} (B_r) = \begin{cases} (-1)^{\binom{r+1}{2} + (|\lambda| + p(\lambda))/2} & \text{if } \lambda \in \mathcal{R}_r, \\ 0 & \text{otherwise.} \end{cases}$$

(2) For a partition λ of length $\leq r$, we have

$$\det \Delta_{I(\lambda)} (C_r) = \begin{cases} (-1)^{\binom{r+1}{2} + |\lambda|/2} & \text{if } \lambda \in \mathcal{Q}_r, \\ 0 & \text{otherwise.} \end{cases}$$

Proposition 4.8.

$$\begin{aligned} G^{p,q}(\boldsymbol{x};\boldsymbol{a}) &= (-1)^{\binom{p}{2} + \binom{q}{2}} \prod_{i=1}^{p+q} x_i^{p-1} (1-x_i^2) \cdot \det V^{p,q}(\boldsymbol{x}+\boldsymbol{x}^{-1};\boldsymbol{a}\boldsymbol{x}^{q-p}), \\ &= (-1)^{\binom{p}{2} + \binom{q}{2}} \prod_{i=1}^{p+q} (1-x_i^2) \cdot \det U^{p,q} \begin{pmatrix} \boldsymbol{x} & | \ \boldsymbol{1} \\ \boldsymbol{1}+\boldsymbol{x}^2 & | \ \boldsymbol{a} \end{pmatrix}, \\ H^{p,q}(\boldsymbol{x};\boldsymbol{a}) &= (-1)^{\binom{p}{2} + \binom{q}{2}} \prod_{i=1}^{p+q} x_i^{p-1} (1-x_i) \cdot \det V^{p,q}(\boldsymbol{x}+\boldsymbol{x}^{-1};\boldsymbol{a}\boldsymbol{x}^{q-p}), \\ &= (-1)^{\binom{p}{2} + \binom{q}{2}} \prod_{i=1}^{p+q} (1-x_i) \cdot \det U^{p,q} \begin{pmatrix} \boldsymbol{x} & | \ \boldsymbol{1} \\ \boldsymbol{1}+\boldsymbol{x}^2 & | \ \boldsymbol{a} \end{pmatrix}. \end{aligned}$$

In particular, we have

$$G^{p,q}(\boldsymbol{x};\boldsymbol{a}) = \prod_{i=1}^{p+q} (1 - x_i^2) \cdot F^{p,q}(\boldsymbol{x};\boldsymbol{a}),$$
$$H^{p,q}(\boldsymbol{x};\boldsymbol{a}) = \prod_{i=1}^{p+q} (1 - x_i) \cdot F^{p,q}(\boldsymbol{x};\boldsymbol{a}).$$

From these relations, we have determinant and Pfaffian identities involving $G^{p,q}(\boldsymbol{x};\boldsymbol{a})$ and $H^{p,q}(\boldsymbol{x};\boldsymbol{a})$ similar to (4.1) and (4.2). More generally, we can consider, for example,

$$\sum_{\lambda \in \mathcal{P}_p, \, \mu \in \mathcal{Q}_q} (-1)^{(|\lambda|+|\mu|)/2} \det V^{p,q}_{\lambda,\mu}(\boldsymbol{x};\boldsymbol{a}),$$

which can be expressed in terms of $\det V^{p,q}$ or $\det U^{p,q}$.

Another generalization of Cauchy's determinant identity 5

In this section, we give another type of generalized Cauchy's determinant identities involving det $V^{p,q}$ and det W^p .

Theorem 5.1.

$$\det\left(\frac{1}{\det V^{p+1,q+1}(x_i, y_j, \boldsymbol{z}; a_i, b_j, \boldsymbol{c})}\right)_{1 \le i,j \le n}$$

$$= \frac{(-1)^{n(n-1)/2} \prod_{1 \le i < j \le n} \det V^{p+1,q+1}(x_i, x_j, \boldsymbol{z}; a_i, a_j, \boldsymbol{c}) \det V^{p+1,q+1}(y_i, y_j, \boldsymbol{z}; b_i, b_j, \boldsymbol{c})}{\prod_{i,j=1}^n \det V^{p+1,q+1}(x_i, y_j, \boldsymbol{z}; a_i, b_j, \boldsymbol{c})}, \quad (5.1)$$

$$\det\left(\frac{1}{\det W^{p+2}(x_i, y_j, \boldsymbol{z}; a_i, b_j, \boldsymbol{c})}\right)_{1 \le i,j \le n}$$

$$= \frac{(-1)^{n(n-1)/2} \prod_{1 \le i < j \le n} \det W^{p+2}(x_i, x_j, \boldsymbol{z}; a_i, a_j, \boldsymbol{c}) \det W^{p+2}(y_i, y_j, \boldsymbol{z}; b_i, b_j, \boldsymbol{c})}{\prod_{i,j=1}^n \det W^{p+2}(x_i, y_j, \boldsymbol{z}; a_i, b_j, \boldsymbol{c})}. \quad (5.2)$$

If p = q = 0, then the identity (5.1) becomes

$$\det\left(\frac{1}{b_j - a_i}\right)_{1 \le i, j \le n} = \frac{(-1)^{n(n-1)/2} \prod_{1 \le i < j \le n} (a_j - a_i)(b_j - b_i)}{\prod_{i, j=1}^n (b_j - a_i)},$$

which is equivalent to Cauchy's determinant identity (1.1).

In order to prove Theorem 5.1, we put

$$f(x, y; a, b) = \det V^{p+1, q+1}(x, y, \boldsymbol{z}; a, b, \boldsymbol{c}), \quad \text{or} \quad \det W^{p+2}(x, y, \boldsymbol{z}; a, b, \boldsymbol{c}).$$

The proof is based on the quadratic relations among f(x, y; a, b)'s, which follows from the Plücker relation for determinants.

Lemma 5.2. Let $(a_1, a_2, a_3, a_4, x_1, \dots, x_m)$ be a $(m+2) \times (m+4)$ matrix. If we put

,

$$D(i,j) = \det(a_i, a_j, x_1, \cdots, x_m),$$

then we have

$$D(1,2)D(3,4) - D(1,3)D(2,4) + D(1,4)D(2,3) = 0.$$

Proposition 5.3. The polynomials f(x, y; a, b)'s satisfy

$$f(x_1, x_2; a_1, a_2)f(y_1, y_2; b_1, b_2) - f(x_1, y_1; a_1, b_1)f(x_2, y_2; a_2, b_2) + f(x_1, y_2; a_1, b_2)f(x_2, y_1; a_2, b_1) = 0.$$
 (5.3)

Proof. Apply the Plücker relation to the transposes of the matrices

$$\begin{pmatrix} 1 & x_1 & \cdots & a_1 x_1^q \\ 1 & x_2 & \cdots & a_2 x_2^q \\ 1 & y_1 & \cdots & b_1 y_1^q \\ 1 & y_2 & \cdots & b_2 y_2^q \\ 1 & z_1 & \cdots & c_1 z_1^q \\ \vdots & \vdots & & \vdots \\ 1 & z_m & \cdots & c_m z_m^q \end{pmatrix} \quad \text{or} \quad \begin{pmatrix} 1 + a_1 x_1^{p+1} & x_1 + a_1 x_1^p & \cdots & x_1^{p+1} + a_1 \\ 1 + a_2 x_2^{p+1} & x_2 + a_2 x_2^p & \cdots & x_2^{p+1} + a_2 \\ 1 + b_1 y_1^{p+1} & y_1 + b_1 y_1^p & \cdots & y_1^{p+1} + b_1 \\ 1 + b_2 y_2^{p+1} & y_2 + b_2 y_2^p & \cdots & y_2^{p+1} + b_2 \\ 1 + c_1 z_1^{p+1} & z_1 + c_1 z_1^p & \cdots & z_1^{p+1} + c_1 \\ \vdots & \vdots & \vdots & & \vdots \\ 1 + c_p z_p^{p+1} & z_p + c_p z_p^p & \cdots & z_p^{p+1} + c_p \end{pmatrix},$$

where m = p + q.

Proof of Theorem 5.1. We proceed by induction on n. In this proof, we write $f(x_i, x_i)$ (resp. $f(x_i, y_j), f(y_i, y_j)$ instead of $f(x_i, x_j; a_i, a_j)$ (resp. $f(x_i, y_j; a_i, b_j), f(y_i, y_j; b_i, b_j)$).

If n = 1, then there is nothing to prove, and, if n = 2, then the desired identities are equivalent to the quadratic relations (5.3).

Suppose $n \geq 3$. Applying the Desnanot–Jacobi formula (2.4) to the matrix

$$A = \left(\frac{1}{f(x_i, y_j)}\right)_{1 \le i, j \le n}.$$

Then, from the induction hypothesis, we have

$$\det A_l^k = \frac{(-1)^{(n-1)(n-2)/2} \prod_{i=3}^n f(x_{k'}, x_i) f(y_{l'}, y_i) \prod_{3 \le i < j \le n} f(x_i, x_j) f(y_i, y_j)}{f(x_{k'}, y_{l'}) \prod_{i=3}^n f(x_{k'}, y_i) f(x_i, y_{l'}) \prod_{i,j=3}^n f(x_i, y_j)}$$

where $1 \le k, l \le 2$ and the indices k' and l' are determined by the condition $\{k, k'\} = \{l, l'\} = \{1, 2\}$, and

$$\det A_{1,2}^{1,2} = \frac{(-1)^{(n-2)(n-3)/2} \prod_{3 \le i < j \le n} f(x_i, x_j) \prod_{3 \le i < j \le n} f(y_i, y_j)}{\prod_{i,j=3}^n f(x_i, y_j)}.$$

By cancelling the common factors, we see that, in order to prove the identity, it is enough to show

$$-f(x_1, x_2)f(y_1, y_2) = f(x_1, y_2)f(x_2, y_1) - f(x_1, y_1)f(x_2, y_2)$$

This is equivalent to (5.3).

6 A hyperpfaffian expression

H. Tagawa finds that det $V^{n,n}(x; a)$ is expressed by a hyperpfaffian. The aim of this section is to prove this expression.

First we recall the definition of hyperpfaffians (see [11]). Let n and r be positive integers. Define a subset $\mathcal{E}_{rn,n}$ of the symmetric groups \mathcal{S}_{rn} by

$$\mathcal{E}_{rn,n} = \{ \sigma \in \mathcal{S}_{rn} : \sigma(n(i-1)+1) < \sigma(n(i-1)+2) < \dots < \sigma(ni) \text{ for } 1 \le i \le r \}.$$

For example, if n = r = 2, then $\mathcal{E}_{4,2}$ is composed of the following 6 elements:

$$\mathcal{E}_{4,2} = \{(1,2,3,4), (1,3,2,4), (1,4,2,3), (3,4,1,2), (2,4,1,3), (2,3,1,4)\}$$

Let $a = (a_{i_1...i_n})_{1 \leq i_1 < \cdots < i_n \leq nr}$ be an alternating tensor, i.e. $a_{i_{\sigma(1)}...i_{\sigma(n)}} = \operatorname{sgn}(\sigma)a_{i_1...i_n}$ for any permutations $\sigma \in S_{nr}$. The hyperpfaffian of a is, by definition,

$$\operatorname{Pf}^{[n]}(a) = \frac{1}{r!} \sum_{\sigma \in \mathcal{E}_{nr,n}} \operatorname{sgn}(\sigma) \prod_{i=1}^{r} a_{\sigma(n(i-1)+1),\dots,\sigma(ni)}.$$

An alternating 2 tensor a is a skew-symmetric matrix and the hyperpfaffian $Pf^{[2]}(a)$ is the usual Pfaffian of the skew-symmetric matrix. J.-G. Luque and J.-Y. Thibon [11] computed the following composition of hyperpfaffians by using the Grassmann algebra.

Proposition 6.1. ([11]) Let n and r be positive integers and assume n = 2m is even. Given a skew-symmetric matrix $A = (a_{ij})_{1 \le i,j \le nr}$, we define an alternating *n*-tensor $A^{[n]}$ by putting

$$(A^{[n]})_{i_1,\dots,i_n} = \Pr(a_{i_k,i_l})_{1 \le k,l \le n} \quad \text{for } 1 \le i_1 < \dots < i_n \le nr.$$

Then we have

$$Pf^{[n]}(A^{[n]}) = \frac{(mr)!}{(m!)^r r!} Pf(A).$$
(6.1)

The main theorem in this section is the following.

Theorem 6.2. If n is even, then

$$\det V^{n,n}(\boldsymbol{x};\boldsymbol{a}) = \Pr^{[n]} \left[\left(1 + \prod_{s=1}^{n} a_{i_s} \right) \prod_{1 \le s < t \le n} (x_{i_t} - x_{i_s}) \right]_{1 \le i_1 < \dots < i_n \le 2n}, \quad (6.2)$$

$$\det U^{n,n}\begin{pmatrix} \boldsymbol{x} & \boldsymbol{a} \\ \boldsymbol{y} & \boldsymbol{b} \end{pmatrix} = \Pr^{[n]} \left[\left(\prod_{s=1}^{n} a_{i_s} + \prod_{s=1}^{n} b_{i_s} \right) \prod_{1 \le s < t \le n} \det \begin{pmatrix} y_{i_s} & x_{i_s} \\ y_{i_t} & x_{i_t} \end{pmatrix} \right]_{1 \le i_1 < \dots < i_n \le 2n}.$$
 (6.3)

To prove this theorem, we need to compute the following special Pfaffian and hyperpfaffian. Lemma 6.3. Let n and r be positive integers and assume n = 2m is even. Then we have

$$\Pr\left(\frac{(x_j^m - x_i^m)^2}{x_j - x_i}\right)_{1 \le i, j \le nr} = \begin{cases} \prod_{1 \le i < j \le n} (x_j - x_i) & \text{if } r = 1, \\ 0 & \text{if } r \ge 2, \end{cases}$$
(6.4)

$$\Pr[n] \left[\prod_{1 \le s < t \le n} (x_{i_t} - x_{i_s}) \right]_{1 \le i_1 < \dots < i_n \le nr} = \begin{cases} \prod_{1 \le i < j \le n} (x_j - x_i) & \text{if } r = 1, \\ 0 & \text{if } r \ge 2. \end{cases}$$
(6.5)

Proof. There are several ways to prove the identity (6.4). Here we appeal to Theorem 1.1 (1.4). If we take p = q = r = s = 0 and put $a_i = b_i = x_i^m$ $(1 \le i \le nr)$ in (1.4), we have

$$\Pr\left(\frac{(x_j^m - x_i^m)^2}{x_j - x_i}\right)_{1 \le i, j \le nr} = \frac{1}{\prod_{1 \le i < j \le nr} (x_j - x_i)} \det V^{mr, mr}(\boldsymbol{x}; \boldsymbol{x}^m)^2.$$

If r = 1, then det $V^{m,m}(\boldsymbol{x}; \boldsymbol{x}^m)$ is the usual Vandermonde determinant and det $V^{m,m}(\boldsymbol{x}; \boldsymbol{x}^m) = \prod_{1 \le i < j \le n} (x_j - x_i)$. If $r \ge 2$, then the (m + 1)st column of $V^{mr,mr}(\boldsymbol{x}; \boldsymbol{x}^m)$ is the same as the (rm + 1)st column, so we have det $V^{mr,mr}(\boldsymbol{x}; \boldsymbol{x}^m) = 0$. Hence we obtain (6.4).

Next we prove (6.5). Apply Proposition 6.1 to the matrix $A = \left((x_j^m - x_i^m)^2 / (x_j - x_i) \right)_{1 \le i,j \le nr}$. Then it follows from (6.4) that

$$\Pr f^{[n]} \left[\prod_{1 \le s < t \le n} (x_{i_t} - x_{i_s}) \right]_{1 \le i_1 < \dots < i_n \le nr} = \Pr f^{[n]} \left[\Pr \left(\frac{(x_{i_l}^m - x_{i_k}^m)^2}{x_{i_l} - x_{i_k}} \right)_{1 \le k, l \le n} \right]_{1 \le i_1 < \dots < i_n \le nr}$$
$$= \frac{(mr)!}{(m!)^r r!} \Pr \left(\frac{(x_j^m - x_i^m)^2}{x_j - x_i} \right)_{1 \le i, j \le nr}.$$

Again using (6.4), we obtain the desired identity.

Now we are in position to prove Theorem 6.2.

Proof of Theorem 6.2. The identity (6.3) immediately follows from (6.2) by noting the relation (3.2), so we prove (6.2).

Let $\binom{[2n]}{n}$ denote the set of all *n*-element subsets of $[2n] = \{1, 2, \dots, 2n\}$. For a subset $I \in \binom{[2n]}{n}$, we put

$$a_I = \prod_{i \in I} a_i, \quad \Delta(\boldsymbol{x}_I) = \prod_{\substack{i,j \in I \\ i < j}} (x_j - x_i).$$

By the Laplace expansion formula and Vandermonde determinant formula, we have

$$V^{n,n}(\boldsymbol{x}; \boldsymbol{a}) = \sum_{I \in \binom{[2n]}{n}} (-1)^{|I| + \binom{n+1}{2}} a_I \Delta(\boldsymbol{x}_I) \Delta(\boldsymbol{x}_{I^c}),$$

where we write $|I| = \sum_{i \in I} i$ and denote by I^c the complementary subset I in [2n].

On the other hand, by the definition of hyperpfaffians, we see that

$$\begin{aligned} \operatorname{Pf}^{[n]}\left[(1+a_{I})\Delta(\boldsymbol{x}_{I})\right]_{I} &= \frac{1}{2!} \sum_{I \in \binom{[2n]}{n}} (-1)^{|I| + \binom{n+1}{2}} (1+a_{I})\Delta(\boldsymbol{x}_{I})(1+a_{I^{c}})\Delta(\boldsymbol{x}_{I^{c}}) \\ &= \left(1 + \prod_{i=1}^{2n} a_{i}\right) \operatorname{Pf}^{[n]}\left[\prod_{1 \leq s < t \leq n} (x_{i_{t}} - x_{i_{s}})\right]_{1 \leq i_{1} < \cdots < i_{n} \leq 2n} \\ &+ \sum_{I \in \binom{[2n]}{n}} (-1)^{|I| + \binom{n+1}{2}} a_{I}\Delta(\boldsymbol{x}_{I})\Delta(\boldsymbol{x}_{I^{c}}). \end{aligned}$$

By (6.5), the first term vanishes, and we obtain the desired formula.

At the end of this section, we should remark that H. Tagawa has a similar hyperpfaffian expression for the case that p = q is odd. It has slightly different from the case p = q is even, but we don't have any general formula when $p \neq q$.

7 Application to Littlewood–Richardson coefficients

In this section, we use the Pfaffian identity (1.5) in Theorem 1.1 and the minor-summation formula [5] to derive a relation between Littlewood–Richardson coefficients.

For three partitions λ , μ and ν , we denote by $LR^{\lambda}_{\mu,\nu}$ the Littlewood-Richardson coefficient. These numbers $LR^{\lambda}_{\mu,\nu}$ appear in the following expansions (see [12]) :

$$\begin{split} s_{\mu}(X)s_{\nu}(X) &= \sum_{\lambda} \mathrm{LR}^{\lambda}_{\mu,\nu} \, s_{\lambda}(X), \\ s_{\lambda/\mu}(X) &= \sum_{\nu} \mathrm{LR}^{\lambda}_{\mu,\nu} \, s_{\nu}(X), \\ s_{\lambda}(X,Y) &= \sum_{\mu,\nu} \mathrm{LR}^{\lambda}_{\mu,\nu} \, s_{\mu}(X)s_{\nu}(Y). \end{split}$$

We are concerned with the Littlewood–Richardson coefficients involving rectangular partitions. Let $\Box(a, b)$ denote the partition whose Young diagram is the rectangle $a \times b$, i.e.

$$\Box(a,b) = (b^a) = (\underbrace{b,\dots,b}_a).$$

For a partition $\lambda \subset \Box(a, b)$, we define a partition $\lambda^{\dagger} = \lambda^{\dagger}(a, b)$ by

$$\lambda_i^{\dagger} = b - \lambda_{a+1-i} \quad (1 \le i \le a).$$

This partition λ^{\dagger} is the complement of λ in the rectangle $\Box(a, b)$.

Okada [13] used the special case of the identities (1.3) and (1.4) (i.e., the case of p = q = 0 and p = q = r = s = 0) to prove the following proposition.

Proposition 7.1. Let n be a positive integer and let e and f be nonnegative integers.

(1) For partitions μ , ν , we have

$$LR^{\Box(n,e)}_{\mu,\nu} = \begin{cases} 1 & \text{if } \nu = \mu^{\dagger}(n,e), \\ 0 & \text{otherwise.} \end{cases}$$
(7.1)

(2) For a partition λ of length $\leq 2n$, we have

$$\mathrm{LR}^{\lambda}_{\Box(n,e),\Box(n,f)} = \begin{cases} 1 & \text{if } \lambda_{n+1} \leq \min(e,f) \text{ and } \lambda_i + \lambda_{2n+1-i} = e+f \ (1 \leq i \leq n), \\ 0 & \text{otherwise.} \end{cases}$$
(7.2)

The main result of this section is the following theorem, which generalizes (7.2).

Theorem 7.2. Let *n* be a positive integer and let *e* and *f* be nonnegative integers. Let λ and μ be partitions such that the length $l(\lambda) \leq 2n$ and $\mu \subset \Box(n, e)$. Then we have

(1) $LR^{\lambda}_{\mu,\Box(n,f)} = 0$ unless

$$\lambda_n \ge f \quad \text{and} \quad \lambda_{n+1} \le \min(e, f).$$
 (7.3)

(2) If λ satisfies the above condition (7.3) and we define two partitions α and β by

$$\alpha_i = \lambda_i - f, \quad \beta_i = e - \lambda_{2n+1-i}, \quad (1 \le i \le n), \tag{7.4}$$

then we have

$$\mathrm{LR}^{\lambda}_{\mu,\Box(n,f)} = \mathrm{LR}^{\beta}_{\alpha,\mu^{\dagger}(n,e)}$$

In particular, $LR^{\lambda}_{\mu,\Box(n,f)} = 0$ unless $\alpha \subset \beta$.

In particular, if $\mu = \Box(n, e)$ is a rectangle, then this theorem reduces to (7.2), because $LR^{\alpha}_{\beta,\emptyset} = \delta_{\alpha,\beta}$. If μ is a near-rectangle, then we have the following corollary by using Pieri's rule [12, (5.16), (5.17)].

Corollary 7.3. Suppose that a partition $\lambda \subset \Box(2n, e+f)$ satisfies the condition (7.3) in Theorem 7.2. Define two partitions α and β by (7.4). Then we have

$$\mathrm{LR}^{\lambda}_{(e^{n-1},e-k),(f^n)} = \begin{cases} 1 & \text{if } \beta/\alpha \text{ is a horizontal strip of length } k \\ 0 & \text{otherwise,} \end{cases}$$

$$\mathrm{LR}^{\lambda}_{(e^{n-k},(e-1)^k),(f^n)} = \begin{cases} 1 & \text{if } \beta/\alpha \text{ is a vertical strip of length } k, \\ 0 & \text{otherwise.} \end{cases}$$

In order to prove Theorem 7.2, we substitute

$$a_i = x_i^{e+p+n}, \quad c_i = z_i^{e+p+n}, \quad d_i = w_i^{f+r+n}$$
 (7.5)

in the Pfaffian identity (1.4). By the bi-determinant definition of Schur functions, we have

$$\det V^{p,q}(\boldsymbol{x};\boldsymbol{x}^k) = \begin{cases} s_{\Box(q,k-p)}(\boldsymbol{x})\Delta(\boldsymbol{x}) & \text{if } k \ge p, \\ 0 & \text{if } k < p, \end{cases}$$

where $\Delta(\mathbf{x}) = \prod_{1 \leq i < j \leq n} (x_j - x_i)$. Hence, under the substitution (7.5), the identity (1.4) give us the following Pfaffian identity.

Proposition 7.4. We have

$$\frac{1}{\Delta(\boldsymbol{x})} \operatorname{Pf} \left((x_j - x_i) s_{\Box(q+1,e+n-1)}(x_i, x_j, \boldsymbol{z}) s_{\Box(s+1,f+n-1)}(x_i, x_j, \boldsymbol{w}) \right)_{1 \le i,j \le 2n} \\ = s_{\Box(q,e+n)}(\boldsymbol{z})^{n-1} s_{\Box(s,f+n)}(\boldsymbol{w})^{n-1} s_{\Box(n+q,e)}(\boldsymbol{x}, \boldsymbol{z}) s_{\Box(n+s,f)}(\boldsymbol{x}, \boldsymbol{w}).$$
(7.6)

Remark 7.5. If we substitute

$$a_i = x_i^{e+p+n}, \quad b_i = y_i^{e+p+n} \quad (1 \le i \le n)$$

in the determinant identity (1.3), then we have

$$\frac{1}{\Delta(\boldsymbol{x})\Delta(\boldsymbol{y})} \det \left(s_{\Box(q+1,e+n-1)}(x_i, y_j, \boldsymbol{z}) \right)_{1 \le i,j \le n} = (-1)^{n(n-1)/2} s_{\Box(q,e+n)}(\boldsymbol{z})^{n-1} s_{\Box(q+n,e)}(\boldsymbol{x}, \boldsymbol{y}, \boldsymbol{z}).$$
(7.7)

The special case (q = e + n - 1) of this identity is given in [10, Proposition 8.4.3], and the proof there works in the general case.

If we take q = s = 0 in (7.6), we have

$$\frac{1}{\Delta(\boldsymbol{x})} \operatorname{Pf}\left((x_j - x_i)h_{e+n-1}(x_i, x_j, \boldsymbol{z})h_{f+n-1}(x_i, x_j, \boldsymbol{w})\right)_{1 \le i,j \le 2n} = s_{\Box(n,e)}(\boldsymbol{x}, \boldsymbol{z})s_{\Box(n,f)}(\boldsymbol{x}, \boldsymbol{w}). \quad (7.8)$$

We use the minor-summation formula [5] to expand the left hand side in the Schur function bases $\{s_{\lambda}(\boldsymbol{x})\}$.

Lemma 7.6. Let $b_{k,l}$ be the coefficient of $x^k y^l$ in

$$(y-x)h_{e+n-1}(x,y,z)h_{f+n-1}(x,y,w).$$

Then we have $b_{kl} = -b_{lk}$, and b_{kl} , k < l, is given by

$$b_{kl} = \sum_{i,j} h_i(\boldsymbol{z}) h_j(\boldsymbol{w}),$$

where the sum is taken over all pairs of integers (i, j) satisfying

$$i+j = (e+n-1) + (f+n-1) + 1 - k - l, \quad 0 \le i \le (e+n-1) - k, \quad 0 \le j \le (f+n-1) - k.$$

Note that $b_{kl} = 0$ unless $0 \le k, l \le e + f + 2n - 1$.

Proof. By using the relation

$$h_r(x, y, \boldsymbol{z}) = \sum_{a, b \ge 0} x^a y^b h_{r-a-b}(\boldsymbol{z}),$$

we see that

$$b_{kl} = \left(\sum_{\substack{0 \le a, b \le e+n-1, \ 0 \le c, d \le f+n-1 \\ a+c=k, \ b+d=l-1}} -\sum_{\substack{0 \le a, b \le e+n-1, \ 0 \le c, d \le f+n-1 \\ a+c=k-1, \ b+d=l}}\right) h_{(e+n-1)-a-b}(\boldsymbol{z})h_{(f+n-1)-c-d}(\boldsymbol{w})$$

Let $b_{kl}(i,j)$ be the coefficient of $h_i(\boldsymbol{z})h_j(\boldsymbol{w})$ in b_{kl} . Then, by considering the homogeneous degree, we see that $b_{kl}(i,j) = 0$ unless $i + j = (e + n - 1) + (f + n - 1) + 1 - k - l, 0 \le i \le e + n - 1$ and $0 \le j \le f + n - 1.$

Now we assume

$$i+j = (e+n-1) + (f+n-1) + 1 - k - l, \quad 0 \le i \le e+n-1, \quad 0 \le j \le f+n-1.$$

If we put

$$S_{1} = \left\{ \begin{pmatrix} a \\ b \\ c \\ d \end{pmatrix} \in \mathbb{N}^{4} : \begin{array}{c} b + d = l - 1, \\ a + b = (e + n - 1) - i, \\ c + d = (f + n - 1) - j \end{array} \right\},$$
$$S_{2} = \left\{ \begin{pmatrix} a' \\ b' \\ c' \\ d' \end{pmatrix} \in \mathbb{N}^{4} : \begin{array}{c} a' + c' = k - 1, \\ b' + d' = l, \\ c' + d' = (e + n - 1) - i, \\ c' + d' = (f + n - 1) - j \end{array} \right\},$$

where $\mathbb N$ denotes the set of nonnegative integers, then we have

$$b_{kl}(i,j) = \#S_1 - \#S_2.$$

The solutions to the equations in S_1 and S_2 are given by

$$\begin{pmatrix} a \\ b \\ c \\ d \end{pmatrix} = \begin{pmatrix} t + (e+n-1) - i - l + 1 \\ -t + l - 1 \\ -t + k + l - 1 - (e+n-1) + i \\ t \end{pmatrix}, \qquad \begin{pmatrix} a' \\ b' \\ c' \\ d' \end{pmatrix} = \begin{pmatrix} t + (e+n-1) - i - l \\ -t + l \\ -t + l \\ -t + k + l - 1 - (e+n-1) + i \\ t \end{pmatrix}.$$

Hence we see that

$$#S_1 = \#\{t \in \mathbb{Z} : t \ge a_0, \ t \le b_0, \ t \le c_0, \ t \ge d_0\}, #S_2 = \#\{t \in \mathbb{Z} : t \ge a_0', \ t \le b_0', \ t \le c_0', \ t \ge d_0'\},$$

where

$$\begin{pmatrix} a_0 \\ b_0 \\ c_0 \\ d_0 \end{pmatrix} = \begin{pmatrix} l-1+i-(e+n-1) \\ l-1 \\ k+l-1-(e+n-1)+i \\ 0 \end{pmatrix}, \qquad \begin{pmatrix} a'_0 \\ b'_0 \\ c'_0 \\ d'_0 \end{pmatrix} = \begin{pmatrix} l+i-(e+n-1) \\ l \\ k+l-1-(e+n-1)+i \\ 0 \end{pmatrix}.$$

We compute $\#S_1$ and $\#S_2$ in the following four cases :

- (a) $i \le (e+n-1) k$ and $j \le (f+n-1) k$.
- (b) $i \le (e+n-1) k$ and j > (f+n-1) k.
- (c) i > (e + n 1) k and $j \le (f + n 1) k$. (d) i > (e + n 1) k and j > (f + n 1) k.

Here we note that

$$j \le (f+n-1) - k$$
 if and only if $l+i - (e+n-1) - 1 \ge 0$,

and that

$$a_0 - d_0 = l + i - (e + n - 1) - 1, \quad b_0 - c_0 = (e + n - 1) - i - k,$$

$$a'_0 - d'_0 = l + i - (e + n - 1) = a_0 - d_0 + 1, \quad b'_0 - c'_0 = (e + n - 1) - i - k + 1 = b_0 - c_0 + 1.$$

Hence we see that, if $i \leq (e+n-1)-k$, then $b_0 \geq c_0$ and $b'_0 > c'_0$, and that, if $i \leq l-k$, then $a_0 \ge d_0$ and $a'_0 > d'_0$.

In Case (a), we have $a_0 \ge d_0, b_0 \ge c_0, a'_0 > d'_0$ and $b'_0 > c'_0$, so

$$#S_1 = #\{t \in \mathbb{Z} : t \ge a_0, \ t \le c_0\} = c_0 - a_0 + 1 = k + 1, #S_2 = #\{t \in \mathbb{Z} : t \ge a'_0, \ t \le c'_0\} = c'_0 - a'_0 + 1 = k$$

Hence we have $b_{kl}(p,q) = 1$. (This argument holds if k = 0.) In Case (b), we have $b_0 \ge c_0, b'_0 > c'_0$, $a_0 < d_0$ and $a'_0 \le d'_0$, so

$$#S_1 = #\{t \in \mathbb{Z} : t \ge d_0, \ t \le c_0\} = c_0 - d_0 + 1 = k + l + i - (e + n - 1), #S_2 = #\{t \in \mathbb{Z} : t \ge d'_0, \ t \le c'_0\} = c'_0 - d'_0 + 1 = k + l + i - (e + n - 1).$$

Hence we have $b_{kl}(i,j) = 0$. Similarly, in Case (c), we have $b_{kl}(i,j) = 0$. In Case (d), we have i+j > (e+n-1) + (f+n-1) - k - l + 1, which contradicts to the assumption i+j = 1(e+n-1) + (f+n-1) - k - l + 1.

This completes the proof.

Here we recall the minor summation formula [5].

Lemma 7.7. Let X be a $2n \times N$ matrix and A be an $N \times N$ skew-symmetric matrix. Then we have

$$\sum_{I} \operatorname{Pf} \Delta_{I}^{I}(A) \det \Delta_{I}(X) = \operatorname{Pf}(XA^{t}X),$$

where I runs over all 2n-element subsets of [N].

By applying this minor-summation formula, we obtain

Proposition 7.8. Let $B = (b_{ij})_{i,j>0}$ be the skew-symmetric matrix, whose entries b_{ij} are given in Lemma 7.6. Then, for a partition λ of length $\leq 2n$, we have

$$\sum_{\substack{\mu \subset \square(n,e)\\\nu \subset \square(n,f)}} \operatorname{LR}_{\mu,\nu}^{\lambda} s_{\mu^{\dagger}(n,e)}(\boldsymbol{z}) s_{\nu^{\dagger}(n,f)}(\boldsymbol{w}) = \operatorname{Pf} \Delta_{I(\lambda)}^{I(\lambda)}(B).$$
(7.9)

Proof. Apply Lemma 7.7 to the matrix $X = (x_i^k)_{1 \le i \le 2n, k \ge 0}$ and the skew-symmetric matrix B. Since det $\Delta_{I(\lambda)}(X)/\Delta(x) = s_{\lambda}(x)$, the left hand side of (7.6) becomes

$$\frac{1}{\Delta(\boldsymbol{x})}\operatorname{Pf}\left((x_j - x_i)h_{e+n-1}(x_i, x_j, \boldsymbol{z})h_{f+n-1}(x_i, x_j, \boldsymbol{w})\right)_{1 \le i, j \le 2n} = \sum_{\lambda}\operatorname{Pf}\Delta_{I(\lambda)}^{I(\lambda)}(B)s_{\lambda}(\boldsymbol{x}).$$

where λ runs over all partitions of length $\leq 2n$. Here we note that $\operatorname{Pf} \Delta_{I(\lambda)}^{I(\lambda)}(B) = 0$ unless $\lambda \subset \Box(2n, e+f).$

On the other hand, the right hand side of (7.6) is expanded in the Schur function basis $\{s_{\lambda}(x)\}$ as follows. It follows from (7.1) that

$$egin{aligned} &s_{\Box(n,e)}(oldsymbol{x},oldsymbol{z}) &= \sum_{\mu \subset \Box(n,e)} s_{\mu}(oldsymbol{x}) s_{\mu^{\dagger}(n,e)}(oldsymbol{z}), \ &s_{\Box(n,f)}(oldsymbol{x},oldsymbol{w}) &= \sum_{
u \subset \Box(n,f)} s_{
u}(oldsymbol{x}) s_{
u^{\dagger}(n,f)}(oldsymbol{w}). \end{aligned}$$

Hence we see that the right hand side of (7.8) becomes

$$egin{aligned} &s_{\Box(n,e)}(oldsymbol{x},oldsymbol{z})s_{\Box(n,f)}(oldsymbol{x},oldsymbol{w}) &= \sum_{\substack{\mu \subset \Box(n,e) \
u \subset \Box(n,f)}} s_{\mu}(oldsymbol{x})s_{
u^{\dagger}(n,e)}(oldsymbol{z})s_{
u^{\dagger}(n,f)}(oldsymbol{w}) & \ &= \sum_{\lambda} \left(\sum_{\substack{\mu \subset \Box(n,e) \
u \subset \Box(n,f)}} \operatorname{LR}^{\lambda}_{\mu,
u} s_{\mu^{\dagger}(n,e)}(oldsymbol{z})s_{
u^{\dagger}(n,f)}(oldsymbol{w})
ight) s_{\lambda}(oldsymbol{x}) \end{aligned}$$

Comparing the coefficient of $s_{\lambda}(x)$ on both sides of (7.8) completes the proof of (7.9).

Now we can finish the proof of Theorem 7.2.

Proof of Theorem 7.2. In the above argument, we take $p \ge n$ and r = 0. In this case, the variables \boldsymbol{w} disappear and we see that

$$b_{kl} = \begin{cases} h_{(e+n-1)+(f+n-1)+1-k-l}(z) & \text{if } 0 \le k \le \min(e+n-1, f+n-1) \text{ and } l \ge f+n-1, \\ 0 & \text{otherwise} \end{cases}$$

and the equation (7.9) becomes

$$\sum_{\mu \subset \Box(n,e)} \operatorname{LR}^{\lambda}_{\mu,\Box(n,f)} s_{\mu^{\dagger}(n,e)}(\boldsymbol{z}) = \operatorname{Pf} \Delta^{I(\lambda)}_{I(\lambda)}(B).$$

The skew-symmetric matrix B has the form

$$B = \begin{pmatrix} O & C & O \\ -{}^{t}C & O & O \\ O & O & O \end{pmatrix}, \quad C = (h_{e+n-1-i-j}(\boldsymbol{z}))_{0 \le i \le f+n-1, 0 \le j \le e+n-1}.$$

¿From the relation (3.5), we see that the subpfaffian Pf $\Delta_{I(\lambda)}^{I(\lambda)}(B)$ vanishes unless

$$\lambda_{n+1} + n - 1 \le \min(e + n - 1, f + n - 1), \quad \lambda_n + n \ge f + n,$$

i.e.,

$$\lambda_{n+1} \le \min(e, f), \quad \lambda_n \ge f.$$

If these conditions are satisfied, then we have

$$Pf \Delta_{I(\lambda)}^{I(\lambda)}(B) = (-1)^{n(n-1)/2} \det \left(h_{\beta_i - \alpha_{n+1-j} - i + (n+1-j)}(\boldsymbol{z}) \right)_{1 \le i,j \le n}$$

= $(-1)^{n(n-1)/2} (-1)^{n(n-1)/2} \det \left(h_{\beta_i - \alpha_j - i + j}(\boldsymbol{z}) \right)_{1 \le i,j \le n}$
= $s_{\beta/\alpha}(\boldsymbol{z}).$

Hence we have

$$\sum_{\mu \subset \Box(n,e)} \operatorname{LR}^{\lambda}_{\mu,\Box(n,f)} s_{\mu^{\dagger}(n,e)}(\boldsymbol{z}) = s_{\beta/\alpha}(\boldsymbol{z}).$$

Comparing the coefficients of $s_{\mu^{\dagger}(n,e)}(z)$ completes the proof.

References

- [1] D. M. Bressoud, "Proofs and Confirmations : The Story of the Alternating Sign Matrix Conjecture", Cambridge University Press, 1999.
- [2] A. L. Cauchy, Mémoire sur les fonctions alternées et sur les sommes alternées, Exercices Anal. et Phys. Math. 2 (1841), 151–159.

- [3] G. Frobenius, Uber die elliptischen Funktionen zweiter Art, J. Reine und Angew. Math. 93 (1882), 53-68.
- [4] M. Ishikawa, Minor summation formula and a proof of Stanley's open problem, arXiv:math.CO/0408204.
- [5] M. Ishikawa and M. Wakayama, Minor summation formulas of Pfaffians, Linear and Multilinear Algebra 39 (1995), 285–305.
- [6] M. Ishikawa and M. Wakayama, Applications of the minor summation formula III: Plücker relations, lattice paths and Pfaffians, to appear in J. Combin. Theory Ser. A, arXiv:math.CO/0312358.
- [7] D. Knuth, Overlapping Pfaffians, Electron. J. Combin. 3 (1996), no. 2, R5, 13 pp.
- [8] C. Krattenthaler, Advanced determinant calculus, Sem. Lothar. Combin. 42 (1999), B42q.
- [9] D. Laksov, A. Lascoux and A. Thorup, On Giambelli's theorem on complete correlations, Acta Math. 162 (1989), 143–199.
- [10] A. Lascoux, Symmetric Functions and Combinatorial Operators on Polynomials, CBMS Regional Conference Series in Mathematics, 99, AMS, 2003.
- [11] J.-G. Luque and J.-Y. Thibon, Pfaffian and Hafnian identities in shuffle algebras, Adv. in Appl. Math. 29 (2002), 620–646.
- [12] I. G. Macdonald, "Symmetric Functions and Hall Polynomials (2nd ed.)", Oxford University Press, 1995.
- [13] S. Okada, Applications of minor-summation formulas to rectangular-shaped representations of classical groups, J. Algebra 205 (1998), 337–367.
- [14] S. Okada, Enumeration of symmetry classes of alternating sign matrices and characters of classical groups, to appear in J. Algebraic Combin., arXiv:math.CO/0408234.
- [15] S. Okada, Determinant and Pfaffian formulae of Cauchy type and their applications, in "Aspects of Combinatorial Representation Theory," Sūrikaisekikenkyūsho Kōkyūroku 1382 (2004), 198–215 (in Japanese).
- [16] S. Okada, An elliptic generalization of Schur's Pfaffian identity, to appear in Adv. Math., arXiv:math.CA/0412038.
- [17] I. Schur, Über die Darstellung der symmetrischen und der alternirenden Gruppe durch gebrochene lineare Substitutuionen, J. Reine Angew. Math. 139 (1911), 155–250.
- [18] J. R. Stembridge, Non-intersecting paths, Pfaffians and plane partitions, Adv. Math. 83 (1990), 96–131.
- [19] T. Sundquist, Two variable Pfaffian identities and symmetric functions, J. Algebraic Combin. 5 (1996), 135–148.