Infinitely many homoclinic solutions for a class of subquadratic second-order Hamiltonian systems *

Xiang Lv[†]

Department of Mathematics, Shanghai Normal University, Shanghai 200234, PR China

Abstract :

In this paper, we mainly consider the existence of infinitely many homoclinic solutions for a class of subquadratic second-order Hamiltonian systems $\ddot{u} - L(t)u + W_u(t, u) = 0$, where L(t) is not necessarily positive definite and the growth rate of potential function W can be in (1, 3/2). Using the variant fountain theorem, we obtain the existence of infinitely many homoclinic solutions for the second-order Hamiltonian systems.

MSC: primary 34C37; 70H05; 58E05

Keywords: Homoclinic solutions; Hamiltonian systems; Variational methods

1 Introduction and main results

The aim of this paper is to study the following second-order Hamiltonian systems

$$\ddot{u} - L(t)u + W_u(t, u) = 0, \quad \forall \ t \in \mathbb{R}$$
(HS)

where $u = (u_1, u_2, \ldots, u_N) \in \mathbb{R}^N$, $W \in C^1(\mathbb{R} \times \mathbb{R}^N, \mathbb{R})$ and $L \in C(\mathbb{R}, \mathbb{R}^{N \times N})$ is a symmetric matrix-valued function. We usually say that a solution u of (HS) is homoclinic (to 0) if $u \in C^2(\mathbb{R}, \mathbb{R}^N)$, $u(t) \to 0$ and $\dot{u}(t) \to 0$ as $t \to \pm \infty$. Furthermore, if $u \not\equiv 0$, then u is called nontrivial.

In the applied sciences, Hamiltonian systems can be used in many practical problems regarding gas dynamics, fluid mechanics and celestial mechanics. It is clear that the existence of homoclinic solutions is one

^{*}Supported by the National Natural Science Foundation of China (NSFC) under Grants No.11371252 and No.11501369; Research and Innovation Project of Shanghai Education Committee under Grant No.14zz120; Yangfan Program of Shanghai (14YF1409100); Chen Guang Project(14CG43) of Shanghai Municipal Education Commission, Shanghai Education Development Foundation and the Research Program of Shanghai Normal University (SK201403) and Shanghai Gaofeng Project for University Academic Program Development.

[†]E-mail:lvxiang@shnu.edu.cn.

of the most important problems in the theory of Hamiltonian systems. Recently, more and more mathematicians have paid their attention to the existence and multiplicity of homoclinic orbits for Hamiltonian systems, see [1-21].

For the case of that L(t) and W(t, x) are either independent of t or periodic in t, there have been several excellent results, see [1–3,7,8,12–16]. More precisely, in the paper [16], Rabinowitz has proved the existence of homoclinic orbits as a limit of 2kT-periodic solutions of (HS). Later, using the same method, several results for general Hamiltonian systems were obtained by Izydorek and Janczewska [8], Lv et al. [12].

If L(t) and W(t, x) are not periodic with respect to t, it will become more difficult to consider the existence of homoclinic orbits for (HS). This problem is quite different from the case mentioned above, due to the lack of compactness of the Sobolev embedding. In [17], Rabinowitz and Tanaka investigated system (HS) without periodicity, both for L and W. Specifically, they assumed that the smallest eigenvalue of L(t) tends to $+\infty$ as $|t| \to \infty$, and showed that system (HS) admits a homoclinic orbit by using a variant of the Mountain Pass theorem without the Palais-Smale condition. Inspired by the work of Rabinowitz and Tanaka [17], many results [4, 6, 10, 11, 14, 15, 18, 20, 21] were obtained for the case of aperiodicity. Most of them were presented under the following condition that L(t) is positive definite for all $t \in \mathbb{R}$,

$$(L(t)u, u) > 0, \quad \forall t \in \mathbb{R} \text{ and } u \in \mathbb{R}^N \setminus \{0\}.$$

Motivated by [6,20], in this article we will study the existence of infinitely many homoclinic solutions for (HS), where L(t) is not necessarily positive definite for all $t \in \mathbb{R}$ and the growth rate of potential function W can be in (1,3/2). The main tool is the variant fountain theorem established in [22]. Our main results are the following theorems.

Theorem 1.1. Assume that L and W satisfy the following conditions: (L1) There exists an $\alpha < 1$ such that

$$l(t)|t|^{\alpha-2} \to \infty \quad as \ |t| \to \infty$$

where $l(t) := \inf_{\substack{|u|=1, u \in \mathbb{R}^N}} (L(t)u, u)$ is the smallest eigenvalue of L(t); (L2) There exist constants $\bar{a} > 0$ and $\bar{r} > 0$ such that

(i) $L \in C^1(\mathbb{R}, \mathbb{R}^{N \times N})$ and $|L'(t)u| \leq \bar{a}|L(t)u|, \quad \forall |t| > \bar{r} \text{ and } u \in \mathbb{R}^N$, or

(ii) $L \in C^2(\mathbb{R}, \mathbb{R}^{N \times N})$ and $\left((L^{''}(t) - \bar{a}L(t))u, u \right) \leq 0, \quad \forall |t| > \bar{r} \text{ and } u \in \mathbb{R}^N$,

where $L^{'}(t) = (d/dt)L(t)$ and $L^{''}(t) = (d^2/dt^2)L(t);$

(W) $W(t, u) = a(t)|u|^{\nu}$ where $a : \mathbb{R} \to \mathbb{R}^+$ is a continuous function such that $a \in L^{\mu}(\mathbb{R}, \mathbb{R}), 1 < \nu < 2$ is a constant, $2 \le \mu \le \overline{\nu}$ and

$$\bar{\nu} = \begin{cases} \frac{2}{3 - 2\nu}, & 1 < \nu < \frac{3}{2} \\ \infty, & \frac{3}{2} \le \nu < 2 \end{cases}$$

Then (HS) possesses infinitely many homoclinic solutions.

Remark 1.2. When we choose $\nu \in (1, \frac{3}{2})$, it is easy to see that W satisfies the condition (W) of Theorem 1.1 but does not satisfy the corresponding conditions in [6, 20]. Furthermore, the constant μ can be change in $[2, \bar{\nu}]$.

2 Preliminaries

In this section, for the purpose of readability and making this paper self-contained, we will show the variational setting for (HS), which can be found in [6,20]. In what follows, we will always assume that L(t) satisfies (L1). Let \mathcal{A} be the selfadjoint extension of the operator $-(d^2/dt^2) + L(t)$ with domain $\mathscr{D}(\mathcal{A}) \subset L^2 \equiv L^2(\mathbb{R}, \mathbb{R}^N)$. Let us write $\{E(\lambda) : -\infty < \lambda < +\infty\}$ and $|\mathcal{A}|$ for the spectral resolution and the absolute value of \mathcal{A} respectively, and denote by $|\mathcal{A}|^{1/2}$ the square root of $|\mathcal{A}|$. Define U = I - E(0) - E(-0). Then U commutes with \mathcal{A} , $|\mathcal{A}|$ and $|\mathcal{A}|^{1/2}$, and $\mathcal{A} = U|\mathcal{A}|$ is the polar decomposition of \mathcal{A} (see [9]). We write $E = \mathscr{D}(|\mathcal{A}|^{1/2})$ and introduce the following inner product on E

$$(u, v)_0 = (|\mathcal{A}|^{1/2}u, |\mathcal{A}|^{1/2}v)_2 + (u, v)_2$$

and norm

$$||u||_0 = (u, u)_0^{1/2}$$

Here, $(\cdot, \cdot)_2$ denotes the usual L^2 -inner product. Therefore, E is a Hilbert space. Since $C_0^{\infty}(\mathbb{R}, \mathbb{R}^N)$ is dense in E, it is obvious that E is continuous embedded in $H^1(\mathbb{R}, \mathbb{R}^N)$ (see [6]). Furthermore, we have the following lemmas by [6].

Lemma 2.1. If L satisfies (L1), then E is compactly embedded in $L^p \equiv L^p(\mathbb{R}, \mathbb{R}^N)$ for all $1 \le p \in (2/(3 - \alpha), \infty]$.

Lemma 2.2. Let L satisfies (L1) and (L2), then $\mathscr{D}(\mathcal{A})$ is continuously embedded in $W^{2,2}(\mathbb{R},\mathbb{R}^N)$, and consequently, we have

$$|u(t)| \to 0 \quad and \quad |\dot{u}(t)| \to 0 \quad as \ |t| \to \infty, \quad \forall \ u \in \mathscr{D}(\mathcal{A}).$$

From [6], combining (L1) and Lemma 2.1, we can prove that \mathcal{A} possesses a compact resolvent. Consequently, the spectrum $\sigma(\mathcal{A})$ consists of eigenvalues, which can be arranged as $\lambda_1 \leq \lambda_2 \leq \cdots \rightarrow \infty$ (counted with multiplicity), and the corresponding system of eigenfunctions $\{e_n : n \in \mathbb{N}\}$, $\mathcal{A}e_n = \lambda_n e_n$, which forms an orthogonal basis in L^2 . Next, we define

$$n^- = \#\{i|\lambda_i < 0\}, \ n^0 = \#\{i|\lambda_i = 0\}, \ \bar{n} = n^- + n^0$$

and

$$E^{-} = \operatorname{span}\{e_1, \cdots, e_{n^-}\}, \ E^0 = \operatorname{span}\{e_{n^-+1}, \cdots, e_{\bar{n}}\} = \operatorname{Ker}\mathcal{A}, \ E^+ = \overline{\operatorname{span}\{e_{\bar{n}+1}, \cdots\}}.$$

Here, the closure is taken in E with respect to the norm $\|\cdot\|_0$. Then

$$E = E^- \oplus E^0 \oplus E^+$$

Furthermore, we define on E the following inner product

$$(u,v) = (|\mathcal{A}|^{1/2}u, |\mathcal{A}|^{1/2}v)_2 + (u^0, v^0)_2$$

and norm

$$\|u\|^2 = (u, u) = \||\mathcal{A}|^{1/2}u\|_2^2 + \|u^0\|_2^2,$$

where $u = u^- + u^0 + u^+$ and $v = v^- + v^0 + v^+ \in E^- \oplus E^0 \oplus E^+$. It is clear that the norms $\|\cdot\|_0$ and $\|\cdot\|$ are equivalent by [6]. From now on, we will take $(E, \|\cdot\|)$ instead of $(E, \|\cdot\|_0)$ as the working space without loss of generality.

Remark 2.3. We note that the decomposition $E = E^- \oplus E^0 \oplus E^+$ is also orthogonal with respect to inner products (\cdot, \cdot) and $(\cdot, \cdot)_2$. Moreover, we will denote by $E = E^- \oplus E^0 \oplus E^+$ the orthogonal decomposition with respect to the inner products (\cdot, \cdot) unless otherwise stated.

Remark 2.4. Since the norms $\|\cdot\|_0$ and $\|\cdot\|$ are equivalent, by Lemma 2.1, for any $1 \le p \in (2/(3-\alpha), \infty]$, there exists a constant $\beta_p > 0$ such that

$$\|u\|_p \le \beta_p \|u\|, \quad \forall \ u \in E, \tag{2.1}$$

where $||u||_p$ denotes the usual norm of L^p and β_p is independent of u.

Let

$$\mathcal{O}(u,v) = (|\mathcal{A}|^{1/2} Uu, |\mathcal{A}|^{1/2} v), \quad \forall \ u, v \in E$$

be the quadratic form associated with \mathcal{A} , where U is the polar decomposition of \mathcal{A} . Given any $u \in \mathscr{D}(\mathcal{A})$ and $v \in E$, we can get

$$\mathcal{O}(u,v) = \int_{\mathbb{R}} \left((\dot{u}, \dot{v}) + (L(t)u, v) \right) dt.$$
(2.2)

Note that $\mathscr{D}(\mathcal{A})$ is dense in E, we have (2.2) holds for all $u, v \in E$. Furthermore, by definition, it follows that

$$\mathcal{O}(u,v) = \left((P^+ - P^-)u, v \right) = \|u^+\|^2 - \|u^-\|^2$$
(2.3)

for all $u = u^- + u^0 + u^+ \in E$, where $P^{\pm} : E \to E^{\pm}$ are the respective orthogonal projections.

Combining (2.2) and (2.3), we define the functional Φ on E by

$$\Phi(u) = \frac{1}{2} \int_{\mathbb{R}} \left(\|\dot{u}\|^2 + (L(t)u, u) \right) dt - \int_{\mathbb{R}} W(t, u) dt$$

$$= \frac{1}{2} \|u^+\|^2 - \frac{1}{2} \|u^-\|^2 - \int_{\mathbb{R}} W(t, u) dt$$

$$= \frac{1}{2} \|u^+\|^2 - \frac{1}{2} \|u^-\|^2 - \Psi(u),$$

(2.4)

where $\Psi(u) = \int_{\mathbb{R}} W(t,u) dt = \int_{\mathbb{R}} a(t) |u|^{\nu} dt$ for all $u = u^- + u^0 + u^+ \in E = E^- \oplus E^0 \oplus E^+$.

Remark 2.5. From (W) with Lemma 2.1, we can easily see that Φ and Ψ are well defined. We will consider two cases as follows.

Case (i) If $2 \le \mu < \infty$, then

$$\begin{split} \Psi(u)| &= \left| \int_{\mathbb{R}} W(t, u) dt \right| = \left| \int_{\mathbb{R}} a(t) |u|^{\nu} dt \right| \\ &\leq \left(\int_{\mathbb{R}} |a(t)|^{\mu} dt \right)^{\frac{1}{\mu}} \left(\int_{\mathbb{R}} |u|^{\nu \mu^{*}} dt \right)^{\frac{1}{\mu^{*}}} \\ &= \|a\|_{\mu} \|u\|_{\nu \mu^{*}}^{\nu} < \infty \end{split}$$

where $\frac{1}{\mu} + \frac{1}{\mu^*} = 1$, $\nu \mu^* \ge 1$. Case (ii) If $\mu = \infty$, then $|\Psi(u)| \le ||a||_{\infty} ||u||_{\nu}^{\nu} < \infty$.

Lemma 2.6. Let (L1), (L2) and (W) hold. Then $\Psi \in C^1(E, \mathbb{R})$ and $\Psi' : E \to E^*$ is compact, and consequently $\Phi \in C^1(E, \mathbb{R})$. Moreover,

$$\Psi'(u)v = \int_{\mathbb{R}} \left(W_u(t,u), u \right) dt = \int_{\mathbb{R}} \left(\nu a(t) |u|^{\nu-2} u, v \right) dt$$
(2.5)

$$\Phi'(u)v = (u^+, v^+) - (u^-, v^-) - \Psi'(u)v$$

= $(u^+, v^+) - (u^-, v^-) - \int_{\mathbb{R}} (W_u(t, u), v) dt$ (2.6)

for all $u = u^- + u^0 + u^+$ and $v = v^- + v^0 + v^+ \in E^- \oplus E^0 \oplus E^+$. Moreover, all critical points of Φ on E are homoclinic solutions of (HS) satisfying $u(t) \to 0$ and $\dot{u}(t) \to 0$ as $|t| \to \infty$.

Proof. We first show that (2.5) holds by definition. If $2 \le \mu < \infty$, then $1 < \mu^* \le 2$, where $\frac{1}{\mu} + \frac{1}{\mu^*} = 1$. For any given $u, v \in E$, by the Mean Value Theorem and the Hölder inequality, we have

$$\begin{aligned} \left| \int_{\mathbb{R}} \left[W(t, u+v) - W(t, u) - \left(W_{u}(t, u), v \right) \right] dt \right| \\ &= \left| \int_{\mathbb{R}} \left[\int_{0}^{1} \left(W_{u}(t, u+\theta v) - W_{u}(t, u), v \right) d\theta \right] dt \right| \\ &\leq 2\nu \int_{\mathbb{R}} |a(t)| (|u|+|v|)^{\nu-1} |v| dt \\ &\leq 2\nu \int_{\mathbb{R}} |a(t)| (|u|^{\nu-1} + |v|^{\nu-1}) |v| dt \\ &\leq 2\nu \left(\int_{\mathbb{R}} |a(t)|^{\mu} dt \right)^{\frac{1}{\mu}} \left(\int_{\mathbb{R}} |u|^{\mu^{*}(\nu-1)} |v|^{\mu^{*}} dt \right)^{\frac{1}{\mu^{*}}} \\ &+ 2\nu \left(\int_{\mathbb{R}} |a(t)|^{\mu} dt \right)^{\frac{1}{\mu}} \left(\int_{\mathbb{R}} |v|^{\mu^{*}\nu} dt \right)^{\frac{1}{\mu^{*}}} \\ &\leq 2\nu ||a||_{\mu} \left(\int_{\mathbb{R}} |u|^{2} dt \right)^{\frac{\nu-1}{2}} \left(\int_{\mathbb{R}} |v|^{\frac{2\mu^{*}}{2+\mu^{*}-\mu^{*}\nu}} dt \right)^{\frac{2+\mu^{*}-\mu^{*}\nu}{2\mu^{*}}} + 2\nu ||a||_{\mu} ||v||_{\mu^{*}\nu}^{\nu} \\ &= 2\nu ||a||_{\mu} ||u||_{2}^{\nu-1} ||v||_{\frac{2\mu^{*}}{2+\mu^{*}-\mu^{*}\nu}} + 2\nu ||a||_{\mu} ||v||_{\mu^{*}\nu}^{\nu} \\ &\leq 2\nu \beta_{\frac{2\mu^{*}}{2+\mu^{*}-\mu^{*}\nu}} ||a||_{\mu} ||u||_{2}^{\nu-1} ||v|| + 2\nu \beta_{\mu^{*}\nu}^{\mu} ||a||_{\mu} ||v||^{\nu} \to 0, \quad \text{as } v \to 0 \text{ in } E \end{aligned}$$

where $\frac{2\mu^*}{2+\mu^*-\mu^*\nu} \ge 1$ and the second inequality holds by the fact that if $0 , then <math>(|a|+|b|)^p \le |a|^p+|b|^p$, $\forall a, b \in \mathbb{R}$. If $\mu = \infty$, then similar to the proof of (2.7), we can obtain

$$\begin{aligned} \left| \int_{\mathbb{R}} \left[W(t, u + v) - W(t, u) - \left(W_u(t, u), v \right) \right] dt \right| \\ &\leq 2\nu \|a\|_{\infty} (\|u\|_{\infty}^{\nu-1} + \|v\|_{\infty}^{\nu-1}) \int_{\mathbb{R}} |v| dt \\ &\leq 2\nu \|a\|_{\infty} \beta_{\infty}^{\nu-1} \beta_1 (\|u\|^{\nu-1} + \|v\|^{\nu-1}) \|v\| \to 0, \quad \text{as } v \to 0 \text{ in } E \end{aligned}$$

$$(2.8)$$

where the last inequality holds by (2.1) and β_{∞} , β_1 are constants there. Combining (2.7) and (2.8), (2.5) holds immediately by the definition of Fréchet derivatives. Consequently, (2.6) also holds due to the definition of Φ .

Next, we verify that $\Psi' : E \to E^*$ is compact. Let $u_n \rightharpoonup u_0$ (weakly) in E, by Lemma 2.1, we have $u_n \to u_0$ in L^p for all $1 \le p \in (2/(3-\alpha), \infty]$. If $2 \le \mu < \infty$, using the Hölder inequality, we can obtain

$$\begin{aligned} \|\Psi'(u_{n}) - \Psi'(u_{0})\|_{E^{*}} &= \sup_{\|v\|=1} \|(\Psi'(u_{n}) - \Psi'(u_{0}))v\| \\ &= \sup_{\|v\|=1} \left| \int_{\mathbb{R}} (W_{u}(t, u_{n}) - W_{u}(t, u_{0}), v) dt \right| \\ &\leq \sup_{\|v\|=1} \left[\left(\int_{\mathbb{R}} |W_{u}(t, u_{n}) - W_{u}(t, u_{0})|^{\mu} dt \right)^{\frac{1}{\mu}} \|v\|_{\mu^{*}} \right] \\ &\leq \beta_{\mu^{*}} \left(\int_{\mathbb{R}} |W_{u}(t, u_{n}) - W_{u}(t, u_{0})|^{\mu} dt \right)^{\frac{1}{\mu}}, \quad \forall n \in \mathbb{N} \end{aligned}$$

$$(2.9)$$

where the last inequality holds by (2.1) and β_{μ}^* is the constant there, $\frac{1}{\mu} + \frac{1}{\mu^*} = 1$. Next, we will prove that $W_u(t, u_n) \to W_u(t, u_0)$ in $L^{\mu}(\mathbb{R}, \mathbb{R}^N)$. Observing that u_n is bounded in L^{∞} , then by the Jensen inequality, we have

$$\int_{\mathbb{R}} |W_{u}(t, u_{n}) - W_{u}(t, u_{0})|^{\mu} dt
\leq 2^{\mu-1} \nu^{\mu} \int_{\mathbb{R}} |a(t)|^{\mu} (|u_{n}|^{\mu} + |u_{0}|^{\mu}) dt
\leq 2^{\mu-1} \nu^{\mu} \int_{\mathbb{R}} |a(t)|^{\mu} (||u_{n}||_{\infty}^{\mu} + ||u_{0}||_{\infty}^{\mu}) dt
\leq 2^{\mu-1} \nu^{\mu} M \int_{\mathbb{R}} |a(t)|^{\mu} dt$$

where $M = 2 \max\{\|u_0\|_{\infty}^{\mu}, \|u_n\|_{\infty}^{\mu}, \forall n \in \mathbb{N}\}$. Combining the fact that $u_n \to u_0$ in L^{∞} and the Lebesgue's Dominated Convergence Theorem,

$$\left(\int_{\mathbb{R}} |W_u(t, u_n) - W_u(t, u_0)|^{\mu} dt\right)^{\frac{1}{\mu}} \to 0, \quad \text{as} \quad n \to \infty$$

Next, we will deal with the case of $\mu = \infty$ (i.e. $\nu > \frac{3}{2}$), this part is mainly motivated by the proof of Lemma 2 in [14]. By the Hölder inequality, we have

$$\begin{aligned} \|\Psi'(u_n) - \Psi'(u_0)\|_{E^*} &\leq \sup_{\|v\|=1} \left[\left(\int_{\mathbb{R}} |W_u(t, u_n) - W_u(t, u_0)|^2 dt \right)^{\frac{1}{2}} \|v\|_2 \right] \\ &\leq \beta_2 \left(\int_{\mathbb{R}} |W_u(t, u_n) - W_u(t, u_0)|^2 dt \right)^{\frac{1}{2}}, \quad \forall n \in \mathbb{N} \end{aligned}$$
(2.10)

We note that by Lemma 2.1, $u_n \to u_0$ in $L^{2(\nu-1)}$ for $\nu > \frac{3}{2}$, passing to a subsequence if necessary, it can be assumed that

$$\sum_{n=1}^{\infty} \|u_n - u_0\|_{2(\nu-1)} < +\infty,$$

which implies that

$$\sum_{n=1}^{\infty} |u_n(t) - u_0(t)| = g(t) \in L^{2(\nu-1)}(\mathbb{R}, \mathbb{R})$$

Since $\nu > \frac{3}{2}$, then

$$\begin{split} &\int_{\mathbb{R}} |W_u(t,u_n) - W_u(t,u_0)|^2 dt \\ &\leq \int_{\mathbb{R}} 2\nu^2 |a(t)|^2 (|u_n|^{2(\nu-1)} + |u_0|^{2(\nu-1)}) dt \\ &\leq \int_{\mathbb{R}} 2\nu^2 |a(t)|^2 (2^{2\nu-3}|u_n - u_0|^{2(\nu-1)} + (2^{2\nu-3} + 1)|u_0|^{2(\nu-1)}) dt \\ &\leq 2^{2\nu-1}\nu^2 \|a\|_{\infty}^2 \int_{\mathbb{R}} (|g(t)|^{2(\nu-1)} + |u_0|^{2(\nu-1)}) dt \\ &\leq 2^{2\nu-1}\nu^2 \|a\|_{\infty}^2 (\|g\|_{2(\nu-1)}^{2(\nu-1)} + \beta_{2(\nu-1)}^{2(\nu-1)}\|u_0\|^{2(\nu-1)}) \end{split}$$

Applying the Lebesgue's Dominated Convergence Theorem, we have

$$\left(\int_{\mathbb{R}} |W_u(t, u_n) - W_u(t, u_0)|^2 dt\right)^{\frac{1}{2}} \to 0, \quad \text{as} \quad n \to \infty.$$

Consequently, Ψ' is weakly continuous, and so Ψ' is continuous. Therefore $\Psi \in C^1(E, \mathbb{R})$ and hence $\Phi \in C^1(E, \mathbb{R})$. Moreover, Ψ' is compact due to the weak continuity of Ψ' and the fact that E is a Hilbert Space.

Finally, we will prove that all critical points of Φ on E are homoclinic solutions of (HS). By the standard procedure, we can see that any critical points of Φ on E satisfy (HS) and $u \in C^2(\mathbb{R}, \mathbb{R}^N)$. We note that if $1 < \nu < \frac{3}{2}$, then $2 \le \mu \le \frac{2}{3-2\nu}$. For $\mu = 2$, by (HS), we have

$$\begin{aligned} \|\mathcal{A}u\|_{2}^{2} &= \int_{\mathbb{R}} |W_{u}(t,u)|^{2} dt \\ &\leq \nu^{2} \|u\|_{\infty}^{2(\nu-1)} \int_{\mathbb{R}} |a(t)|^{2} dt \\ &\leq \nu^{2} \beta_{\infty}^{2(\nu-1)} \|u\|^{2(\nu-1)} \int_{\mathbb{R}} |a(t)|^{\mu} dt < \infty. \end{aligned}$$
(2.11)

In the case of $2 < \mu \leq \frac{2}{3-2\nu}$, then

$$\begin{aligned} \|\mathcal{A}u\|_{2}^{2} &= \int_{\mathbb{R}} |W_{u}(t,u)|^{2} dt \\ &\leq \nu^{2} \left(\int_{\mathbb{R}} |a(t)|^{\mu} dt \right)^{\frac{2}{\mu}} \left(\int_{\mathbb{R}} |u|^{2\bar{\mu}(\nu-1)} dt \right)^{\frac{1}{\bar{\mu}}} \\ &\leq \nu^{2} \|u\|_{2\bar{\mu}(\nu-1)}^{2(\nu-1)} \left(\int_{\mathbb{R}} |a(t)|^{\mu} dt \right)^{\frac{2}{\mu}} \\ &\leq \nu^{2} \beta_{2\bar{\mu}(\nu-1)}^{2(\nu-1)} \|u\|^{2(\nu-1)} \left(\int_{\mathbb{R}} |a(t)|^{\mu} dt \right)^{\frac{2}{\mu}} < \infty, \end{aligned}$$

$$(2.12)$$

where $\frac{2}{\mu} + \frac{1}{\bar{\mu}} = 1$ and $2\bar{\mu}(\nu - 1) \ge 1$ because of $\mu \le \frac{2}{3-2\nu}$. If $\frac{3}{2} \le \nu < 2$, combining the fact that $2(\nu - 1) \ge 1$ and Hölder inequality, similar to the proof of (2.11) and (2.12), we can get the same result. Consequently, $u \in \mathscr{D}(\mathcal{A})$ and hence u is a homoclinic solution of (HS) by Lemma 2.2. The proof is complete.

In the next argument, the following variant fountain theorem will be used to prove our main results. Let E be a Banach space with the norm $\|\cdot\|$ and $E = \overline{\bigoplus_{j \in \mathbb{N}} X_j}$ with $\dim X_j < \infty$ for any $j \in \mathbb{N}$. We write $Y_k = \bigoplus_{j=1}^k X_j$ and $Z_k = \overline{\bigoplus_{j=k} X_j}$. The C^1 -functional $\Phi_{\lambda} : E \to \mathbb{R}$ is given by

$$\Phi_{\lambda}(u) := A(u) - \lambda B(u), \qquad \lambda \in [1, 2]$$

Theorem 2.7. ([22, Theorem2.2.]) Assume that the functional Φ_{λ} defined above satisfies (F1) Φ_{λ} maps bounded sets to bounded sets uniformly for $\lambda \in [1, 2]$. Furthermore, $\Phi_{\lambda}(-u) = \Phi_{\lambda}(u)$ for all $(\lambda, u) \in [1, 2] \times E$;

(F2) $B(u) \ge 0$; $B(u) \to \infty$ as $||u|| \to \infty$ on any finite dimensional subspace of E;

(F3) There exist $\rho_k > r_k > 0$ such that

$$a_k(\lambda) := \inf_{u \in Z_k, \|u\| = \rho_k} \Phi_\lambda(u) \ge 0 > b_k(\lambda) := \max_{u \in Y_k, \|u\| = r_k} \Phi_\lambda(u), \quad \forall \ \lambda \in [1, 2]$$

and

$$d_k(\lambda) := \inf_{u \in Z_k, \|u\| \le \rho_k} \Phi_{\lambda}(u) \to 0 \quad as \ k \to \infty \ uniformly \ for \ \lambda \in [1, 2].$$

Then there exist $\lambda_n \to 1$, $u_{\lambda_n} \in Y_n$ such that

$$\Phi_{\lambda_n}'|_{Y_n}(u_{\lambda_n}) = 0, \ \Phi_{\lambda_n}(u_{\lambda_n}) \to c_k \in [d_k(2), b_k(1)] \quad as \quad n \to \infty.$$

In particular, if $\{u_{\lambda_n}\}$ has a convergent subsequence for every k, then Φ_1 has infinitely many nontrivial critical points $\{u_k\} \in E \setminus \{0\}$ satisfying $\Phi_1(u_k) \to 0^-$ as $k \to \infty$.

In order to make use of Theorem 2.7, we consider the functionals A, B and Φ_{λ} on the working space defined $E = \mathscr{D}(|\mathcal{A}|^{1/2})$ by

$$A(u) = \frac{1}{2} \|u^+\|^2, \qquad B(u) = \frac{1}{2} \|u^-\|^2 + \int_{\mathbb{R}} W(t, u) dt,$$
(2.13)

and

$$\Phi_{\lambda}(u) = A(u) - \lambda B(u) = \frac{1}{2} ||u^{+}||^{2} - \lambda \left(\frac{1}{2} ||u^{-}||^{2} + \int_{\mathbb{R}} W(t, u) dt\right)$$
(2.14)

for all $u = u^- + u^0 + u^+ \in E$ and $\lambda \in [1, 2]$. By Lemma 2.6, it is clear that $\Phi_{\lambda} \in C^1(E, \mathbb{R})$ for all $\lambda \in [1, 2]$. Let $X_j := \mathbb{R}e_j = \operatorname{span}\{e_j\}, j \in \mathbb{N}$, where $\{e_j, j \in \mathbb{N}\}$ is the system of eigenfunctions and the orthogonal basis in L^2 below Lemma 2.2. Furthermore, it is evident that $\Phi_1 = \Phi$, where Φ is the functional defined in (2.4).

3 Proof of theorems

Lemma 3.1. Let (L1), (L2) and (W) hold, then $B(u) \ge 0$. Moreover, $B(u) \to \infty$ as $||u|| \to \infty$ on any finite dimensional subspace of E.

Proof. By definitions of the functional B and (W), $B(u) \ge 0$ holds obviously. Next we will prove that $B(u) \to \infty$ as $||u|| \to \infty$ on any finite dimensional subspace of E. First we claim that for any finite dimensional subspace $F \subset E$, there exists $\varepsilon > 0$ such that

$$meas\{t \in \mathbb{R} : a(t)|u(t)|^{\nu} \ge \varepsilon ||u||^{\nu}\} \ge \varepsilon, \qquad \forall \ u \in F \setminus \{0\}.$$

$$(3.1)$$

The proof of (3.1) is very similar as that of [18]. We omit it here. Now, let

$$\Omega_u = \{ t \in \mathbb{R} : a(t) | u(t) |^{\nu} \ge \varepsilon || u ||^{\nu} \}, \qquad \forall \ u \in F \setminus \{0\},$$
(3.2)

where ε is given in (3.1). From (3.1), we can obtain that

$$meas(\Omega_u) \ge \varepsilon, \qquad \forall \ u \in F \setminus \{0\},\tag{3.3}$$

Combining (W) and (3.3), for all $u \in F \setminus \{0\}$, we can see that

$$B(u) = \frac{1}{2} ||u^-||^2 + \int_{\mathbb{R}} W(t, u) dt$$

$$\geq \int_{\Omega_u} a(t) |u(t)|^{\nu} dt$$

$$\geq \varepsilon ||u||^{\nu} meas(\Omega_u) \geq \varepsilon^2 ||u||^{\nu}.$$
(3.4)

This implies $B(u) \to \infty$ as $||u|| \to \infty$ on any finite dimensional subspace of E. If $\mu = \infty$, similar to the case of $2 \le \mu < \infty$, by the standard procedure, we can prove that there exists $\varepsilon_1 > 0$ such that

$$meas\{t \in \mathbb{R} : a(t)|u(t)|^{\nu} \ge \varepsilon_1 ||u||^{\nu}\} \ge \varepsilon_1, \qquad \forall \ u \in F \setminus \{0\}.$$

$$(3.5)$$

Therefore, by (3.4), we can conclude that $B(u) \to \infty$ as $||u|| \to \infty$ on any finite dimensional subspace of E. The proof is complete.

Lemma 3.2. Under the conditions in Theorem 1.1, then there exists a sequence $\rho_k \to 0^+$ as $k \to \infty$ such that

$$a_k(\lambda) := \inf_{u \in Z_k, \|u\| = \rho_k} \Phi_\lambda(u) \ge 0, \quad \forall \ \lambda \in [1, 2], \ k \ge \bar{n} + 1,$$

and

$$d_k(\lambda) := \inf_{u \in Z_k, \|u\| \le \rho_k} \Phi_{\lambda}(u) \to 0 \quad as \ k \to \infty \ uniformly \ for \ \lambda \in [1, 2].$$

where $Z_k = \overline{\bigoplus_{j=k} X_j}$ for all $k \in \mathbb{N}$.

Proof. By the definition of \bar{n} below the Lemma 2.2, we can know that $Z_k \subset E^+$ for all $k \ge \bar{n}+1$. Therefore, for all $k \ge \bar{n}+1$, from (W) and (2.14), it follows that

$$\Phi_{\lambda}(u) = \frac{1}{2} \|u\|^{2} - \lambda \int_{\mathbb{R}} W(t, u) dt
\geq \frac{1}{2} \|u\|^{2} - 2 \int_{\mathbb{R}} W(t, u) dt
= \frac{1}{2} \|u\|^{2} - 2 \int_{\mathbb{R}} a(t) |u|^{\nu} dt, \qquad \forall (\lambda, u) \in [1, 2] \times Z_{k}.$$
(3.6)

If $2 \le \mu < \infty$, let $\eta_k := \sup_{u \in Z_k, ||u||=1} ||u||_{\nu\mu^*}$, where $\frac{1}{\mu} + \frac{1}{\mu^*} = 1$. By Lemma 2.1, we can conclude that $\eta_k \to 0$ as $k \to \infty$. Therefore, combining (3.6) with (W), we have

$$\Phi_{\lambda}(u) \ge \frac{1}{2} \|u\|^2 - 2\|a\|_{\mu} \|u\|_{\nu\mu^*}^{\nu} \ge \frac{1}{2} \|u\|^2 - 2\eta_k^{\nu} \|a\|_{\mu} \|u\|^{\nu}, \qquad \forall \ (\lambda, u) \in [1, 2] \times Z_k.$$
(3.7)

Let $\rho_k := (8\eta_k^{\nu} ||a||_{\mu})^{1/(2-\nu)}$, the rest of proof is very similar as that of [18]. We omit it here. For the case of $\mu = \infty$, similar to the above procedure, the same result can be obtained. We omit it here. The proof is complete.

Lemma 3.3. Assume that (L1), (L2) and (W) hold, then for the sequence $\{\rho_k\}_{k\in\mathbb{N}}$ obtained in Lemma 3.2, there exists a sequence $\{r_k\}_{k\in\mathbb{N}}$ such that $\rho_k > r_k > 0$ for $\forall k \in \mathbb{N}$ and

$$b_k(\lambda) := \max_{u \in Y_k, \|u\| = r_k} \Phi_\lambda(u) < 0, \quad \forall \ \lambda \in [1, 2].$$

$$(3.8)$$

where $Y_k = \bigoplus_{j=1}^k X_j = span\{e_1, \dots, e_k\}$ for $\forall k \in \mathbb{N}$.

Proof. For $\forall k \in \mathbb{N}$, it is clear that Y_k is a finite dimensional subspace of E. Therefore, for $\forall \lambda \in [1, 2]$, from (W), (3.2), (3.3) and (3.5), let $\varepsilon_0 = \min\{\varepsilon, \varepsilon_1\}$, we have

$$\Phi_{\lambda}(u) = \frac{1}{2} ||u^{+}||^{2} - \lambda \left(\frac{1}{2} ||u^{-}||^{2} + \int_{\mathbb{R}} W(t, u) dt \right) \\
\leq \frac{1}{2} ||u||^{2} - \int_{\mathbb{R}} W(t, u) dt \\
\leq \frac{1}{2} ||u||^{2} - \int_{\Omega_{u}} a(t) |u|^{\nu} dt \\
\leq \frac{1}{2} ||u||^{2} - \varepsilon_{0} ||u||^{\nu} \operatorname{meas}(\Omega_{u}) \\
\leq \frac{1}{2} ||u||^{2} - \varepsilon_{0}^{2} ||u||^{\nu}, \qquad \forall u \in Y_{k}, k \in \mathbb{N}.$$
(3.9)

For $\forall k \in \mathbb{N}$, we choose $0 < r_k < \min\{\rho_k, \varepsilon_0^{\frac{2}{2-\nu}}\}$. From (3.9), an easy computation shows that

$$b_k(\lambda) := \max_{u \in Y_k, \|u\| = r_k} \Phi_{\lambda}(u) \le -\frac{r_k^2}{2} < 0, \qquad \forall \ k \in \mathbb{N}.$$

The proof is complete.

Next we will present the proof of our main result.

Proof of Theorem 1.1. Combining Remark 2.5 and (2.14), it is clear that the condition (F1) in Theorem 2.7 holds obviously. By Lemma 3.1, 3.2 and 3.3, we can easily see that conditions (F2) and (F3) in Theorem 2.7 hold for all $k \ge \bar{n} + 1$. Consequently, from Theorem 2.7, for all $k \ge \bar{n} + 1$, there exist $\lambda_n \to 1$, $u_{\lambda_n} \in Y_n$ such that

$$\Phi_{\lambda_n}'|_{Y_n}(u_{\lambda_n}) = 0, \ \Phi_{\lambda_n}(u_{\lambda_n}) \to c_k \in [d_k(2), b_k(1)] \quad \text{as } n \to \infty.$$
(3.10)

In what follows, the first step is to show that $\{u_{\lambda_n}\}$ is bounded in E. For the case of $2 \leq \mu < \infty$, since $\Phi'_{\lambda_n}|_{Y_n}(u_{\lambda_n}) = 0$, by (2.6) and (2.14), we have

$$\Phi_{\lambda_n}'|_{Y_n}(u_{\lambda_n})u_{\lambda_n}^+ = \|u_{\lambda_n}^+\|^2 - \lambda_n \int_{\mathbb{R}} \left(W_u(t, u_{\lambda_n}), u_{\lambda_n}^+ \right) dt = 0.$$
(3.11)

Therefore, using (W) and the Hölder inequality, it follows that

$$\begin{aligned} \|u_{\lambda_{n}}^{+}\|^{2} &= \lambda_{n} \int_{\mathbb{R}} \left(W_{u}(t, u_{\lambda_{n}}), u_{\lambda_{n}}^{+} \right) dt \\ &\leq 2 \int_{\mathbb{R}} |a(t)| |u_{\lambda_{n}}|^{\nu-1} |u_{\lambda_{n}}^{+}| dt \\ &\leq 2 \left(\int_{\mathbb{R}} |a(t)|^{\mu} dt \right)^{\frac{1}{\mu}} \left(\int_{\mathbb{R}} |u_{\lambda_{n}}|^{\mu^{*}(\nu-1)} |u_{\lambda_{n}}^{+}|^{\mu^{*}} dt \right)^{\frac{1}{\mu^{*}}} \\ &\leq 2 \nu \|a\|_{\mu} \left(\int_{\mathbb{R}} |u_{\lambda_{n}}|^{2} dt \right)^{\frac{\nu-1}{2}} \left(\int_{\mathbb{R}} |u_{\lambda_{n}}^{+}|^{\frac{2\mu^{*}}{2+\mu^{*}-\mu^{*}\nu}} dt \right)^{\frac{2+\mu^{*}-\mu^{*}\nu}{2\mu^{*}}} \\ &= 2\nu \|a\|_{\mu} \|u_{\lambda_{n}}\|_{2}^{\nu-1} \|u_{\lambda_{n}}^{+}\|_{\frac{2\mu^{*}}{2+\mu^{*}-\mu^{*}\nu}} \\ &\leq M_{1} \|a\|_{\mu} \|u_{\lambda_{n}}\|^{\nu} \end{aligned}$$
(3.12)

for some $M_1 > 0$, where $\frac{1}{\mu} + \frac{1}{\mu^*} = 1$, $\frac{2\mu^*}{2+\mu^*-\mu^*\nu} \ge 1$ and the last inequality holds because of (2.1). Furthermore, combing (2.6) with (3.10) and the Hölder inequality, we have

$$-\Phi_{\lambda_{n}}(u_{\lambda_{n}}) = \frac{1}{2} \Phi_{\lambda_{n}}'|_{Y_{n}}(u_{\lambda_{n}})u_{\lambda_{n}} - \Phi_{\lambda_{n}}(u_{\lambda_{n}}) = \lambda_{n}(1-\frac{\nu}{2}) \int_{\mathbb{R}} |a(t)||u_{\lambda_{n}}|^{\nu} dt \geq \frac{1}{2^{\nu-1}} \lambda_{n}(1-\frac{\nu}{2}) \int_{\mathbb{R}} |a(t)||u_{\lambda_{n}}^{-} + u_{\lambda_{n}}^{0}|^{\nu} dt - \lambda_{n}(1-\frac{\nu}{2}) \int_{\mathbb{R}} |a(t)||u_{\lambda_{n}}^{+}|^{\nu} dt \geq \frac{\varepsilon^{2}}{2^{\nu-1}} \lambda_{n}(1-\frac{\nu}{2}) ||u_{\lambda_{n}}^{-} + u_{\lambda_{n}}^{0}||^{\nu} - \lambda_{n}(1-\frac{\nu}{2}) ||a||_{\mu} ||u_{\lambda_{n}}^{+}||_{\mu^{*}\nu}^{\nu}$$
(3.13)

where the last inequality holds by the fact that $\dim(E^- \oplus E^0) < \infty$ and (3.1). Note that $1 < \nu < 2$, then (3.12) and (3.13) implies that $\{\|u_{\lambda_n}^+\|\}$ is bounded. Next, we just have to show that $\{\|u_{\lambda_n}^- + u_{\lambda_n}^0\|\}$ is also bounded. Consequently, from (3.13) and (2.1), we get

$$\|u_{\lambda_n}^- + u_{\lambda_n}^0\|^{\nu} \le -M_2 \Phi_{\lambda_n}(u_{\lambda_n}) + M_3 \|u_{\lambda_n}^+\|_{\mu^*\nu}^{\nu} \le -M_2 \Phi_{\lambda_n}(u_{\lambda_n}) + M_4 \|u_{\lambda_n}^+\|^{\nu}$$
(3.14)

for some positive constants M_2 , M_3 and M_4 . Notice that $\{\|u_{\lambda_n}^+\|\}$ is bounded, by (3.10), we can conclude that $\{\|u_{\lambda_n}^-+u_{\lambda_n}^0\|\}$ is also bounded. Therefore, there exists $M_5 > 0$ such that $\|u_{\lambda_n}\|^2 = \|u_{\lambda_n}^+\|^2 + \|u_{\lambda_n}^-+u_{\lambda_n}^0\|^2 \le M_5$, i.e. $\{u_{\lambda_n}\}$ is bounded in E.

Finally, we prove that $\{u_{\lambda_n}\}$ has a strong convergent subsequence in E. The proof of this assertion can be accomplished as that of [18]. We omit it here.

Now by the last conclusion of Theorem 2.7, we obtain that $\Phi = \Phi_1$ has infinitely many nontrivial critical points. Consequently, (HS) possesses infinitely many homoclinic solutions by Lemma 2.6. The proof of Theorem 1.1 is complete. \Box

Remark 3.4. In this paper, we have considered the existence of infinitely many homoclinic solutions for a class of subquadratic second-order Hamiltonian systems, where $1 < \nu < \frac{3}{2}$ is allowed. We view this result as merely one first step in the theory for the case of $1 < \nu < \frac{3}{2}$, there are still many problems to pursue. For example, when $1 < \nu < \frac{3}{2}$, the upper bound of μ whether can be ∞ , what we will discuss in the future study.

Acknowledgements

The authors would like to thank the reviewer for their valuable comments and suggestions.

References

- A. Ambrosetti, V. Coti Zelati, Multiple homoclinic orbits for a class of conservative systems, Rend. Sem. Mat. Univ. Padova 89 (1993) 177-194.
- [2] A. Ambrosetti, P.H. Rabinowitz, Dual variational methods in critical point theory and applications, J. Funct. Anal. 14 (1973) 349-381.
- [3] P.C. Carrião, O.H. Miyagaki, Existence of homoclinic solutions for a class of time-dependent Hamiltonian systems, J. Math. Anal. Appl. 230 (1999) 157-172.
- [4] V. Coti Zelati, I. Ekeland, E. Séré, A variational approach to homoclinic orbits in Hamiltonian systems, Math. Ann. 288 (1990) 133-160.
- [5] V. Coti Zelati, P.H. Rabinowitz, Homoclinic orbits for second order Hamiltonian systems possessing superquadratic potentials, J. Amer. Math. Soc. 4 (1991) 693-727.
- Y. Ding, Existence and multiplicity results for homoclinic solutions to a class of Hamiltonian systems, Nonlinear Anal. 25 (1995) 1095-1113.
- [7] Y. Ding, M. Girardi, Periodic and homoclinic solutions to a class of Hamiltonian systems with the potentials changing sign, Dynam. Systems Appl. 2 (1993) 131-145.
- [8] M. Izydorek, J. Janczewska, Homoclinic solutions for a class of second order Hamiltonian systems, J. Differential Equations 219 (2005) 375-389.

- [9] T. Kato, Perturbation Theory for Linear Operators Springer-Verlag, New York, 1980.
- [10] P. Korman, A.C. Lazer, Homoclinic orbits for a class of symmetric Hamiltonian systems, Electron. J. Differential Equations 1 (1994) 1-10.
- [11] X. Lv, S. Lu, P. Yan, Existence of homoclinic solutions for a class of second-order Hamiltonian systems, Nonlinear Anal. 72 (2010) 390-398.
- [12] X. Lv, S. Lu, J. Jiang, Homoclinic solutions for a class of second-order Hamiltonian systems, Nonlinear Anal. RWA. 13 (2012) 176-185.
- [13] J. Mawhin, M. Willem, Critical Point Theory and Hamiltonian Systems, Appl. Math. Sci, vol. 74, Springer-Verlag, New York, 1989.
- [14] W. Omana, M. Willem, Homoclinic orbits for a class of Hamiltonian systems, Differential Integral Equations 5 (1992) 1115-1120.
- [15] Z. Qu, C.L. Tang, Existence of homoclinic orbits for the second order Hamiltonian systems, J. Math. Anal. Appl. 291 (2004) 203-213.
- [16] P.H. Rabinowitz, Homoclinic orbits for a class of Hamiltonian systems, Proc. Roy. Soc. Edinburgh Sect. A 114 (1990) 33-38.
- [17] P.H. Rabinowitz, K. Tanaka, Some results on connecting orbits for a class of Hamiltonian systems, Math. Z. 206 (1991) 473-499.
- [18] J. Sun, H. Chen, J.J. Nieto, Homoclinic solutions for a class of subquadratic second-order Hamiltonian systems, J. Math. Anal. Appl. 373 (2011) 20-29.
- [19] M. Willem, Minimax Theorems. Boston: Birkhäuser, 1996.
- [20] Q. Zhang, C. Liu, Infinitely many homoclinic solutions for second order Hamiltonian systems, Nonlinear Anal. 72 (2010) 894-903.
- [21] Z. Zhang, R. Yuan, Homoclinic solutions for a class of non-autonomous subquadratic second-order Hamiltonian systems, Nonlinear Anal. 71 (2009) 4125-4130.
- [22] W. Zou, Variant fountain theorems and their applications, Manuscripta Math. 104 (2001) 343-358.