O-minimal residue fields of o-minimal fields

Jana Maříková Dept. of Math., Western Illinois University* J-Marikova@wiu.edu

April 29, 2022

Abstract

Let R be an o-minimal field with a proper convex subring V. We axiomatize the class of all structures (R, V) such that \mathbf{k}_{ind} , the corresponding residue field with structure induced from R via the residue map, is o-minimal. More precisely, in [8] it was shown that certain first order conditions on (R, V) are sufficient for the o-minimality of \mathbf{k}_{ind} . Here we prove that these conditions are also necessary.

1 Introduction

Throughout this paper we let R be an o-minimal field, that is, an o-minimal expansion of a real closed field, and V a proper convex subring (hence a valuation ring). Let st: $V \to \mathbf{k}$ be the corresponding residue (standard part) map with kernel \mathfrak{m} and residue field $\mathbf{k} = V/\mathfrak{m}$. For $X \subseteq R^n$ we set st $X := \operatorname{st}(X \cap V^n)$. By $\mathbf{k}_{\operatorname{ind}}$ we denote the residue field expanded by all sets st $X \subseteq \mathbf{k}^n$ with definable $X \subseteq R^n$, for some n. "Definable" means "definable (with parameters) in R", unless indicated otherwise.

Here are some notational conventions we use: By $\mathbb{N} = \{0, 1, ...\}$ we denote the set of natural numbers, and the letters i, j, k, l, m, n denote natural numbers. For $a \in \mathbb{R}^n$ and i = 1, ..., n, we let a_i be the *i*-th coordinate of a,

^{*}Part of this work was done while the author was a Fields Ontario postdoctoral fellow, during the thematic program on o-minimality and real analytic geometry at the Fields Institute, Toronto, spring 2009.

unless stated otherwise. We let $I := [-1, 1] \subseteq R$, and for a definable set $X \subseteq R^{1+n}$ and $r \in R$, we set

$$X(r) := \{ x \in R^n \colon (r, x) \in X \}.$$

By d(x, y) we denote the euclidean distance between x and y. For definable $Y \subseteq \mathbb{R}^n$ and $x \in \mathbb{R}^n$,

$$d(x, Y) := \inf\{d(x, y) : y \in Y\}.$$

If $X, Y \subseteq \mathbb{R}^n$ are definable, then $X \cong Y$ means that there is a definable homeomorphism $X \to Y$. For a definable C^1 -map $\phi \colon X \to \mathbb{R}^n$, with X an open subset of \mathbb{R}^m , and $a \in X$, we denote by $D\phi(a)$ the Jacobian matrix of ϕ at a. If $X \subseteq \mathbb{R}^n$, then we denote by cl(X) the closure of X in \mathbb{R}^n , by X^o the interior of X in \mathbb{R}^n , and we let $\partial X := cl(X) \setminus X$, and $bd(X) := cl(X) \setminus X^o$. For 0 < k < n, p_k^n denotes the projection map

$$R^n \to R^k \colon (x_1, \dots, x_n) \mapsto (x_1, \dots, x_k).$$

For a definable map $\phi: X \to \mathbb{R}^n$, $X \subseteq \mathbb{R}^m$, and $i = 1, \ldots, m$, we denote by ϕ_i the *i*-th coordinate function of ϕ . If $a, b \in \mathbb{R}^n$, then [a, b] denotes the set $\{(t-1)a+tb: t \in [0, 1]\}$, and (a, b) denotes the set $\{(1-t)a+tb: t \in (0, 1)\}$. We recall now some definitions and the main result from [8]:

 $(R, V) \models \Sigma(n)$ means that for every definable $X \subseteq I^{1+n}$ there is $\epsilon_0 \in \mathfrak{m}^{>0}$ such that st $X(\epsilon_0) = \operatorname{st} X(\epsilon)$ for all $\epsilon \in \mathfrak{m}^{>\epsilon_0}$.

 $(R, V) \models \Sigma$ means that $(R, V) \models \Sigma(n)$ for every n.

Theorem 1.1 ([8]) If $(R, V) \models \Sigma$, then \mathbf{k}_{ind} is o-minimal, and the subsets of \mathbf{k}^n definable in \mathbf{k}_{ind} are exactly the finite unions of sets st $X \setminus \text{st } Y$, where $X, Y \subseteq R^n$ are definable.

The question, whether, on the other hand, o-minimality of \mathbf{k}_{ind} implies $(R, V) \models \Sigma$, was left unanswered. Here, we give a positive answer to it, yielding the remarkable fact, that the o-minimality of \mathbf{k}_{ind} is equivalent to (R, V) satisfying a first-order axiom scheme. More concretely, we obtain:

Theorem 1.2 The following conditions on (R, V) are equivalent:

1. \mathbf{k}_{ind} is o-minimal

2. Σ(1)
 3. Σ.

The implication $3 \Rightarrow 1$ is Theorem 1.1, and $1 \Rightarrow 2$ can be found in [5]. So it is left to show $\Sigma(1) \Rightarrow \Sigma$. Note that by cell decomposition, and using the definable homeomorphism

$$\tau \colon R^2 \to (-1,1)^2 \colon (x_1,x_2) \mapsto (\frac{x_1}{\sqrt{1+x_1^2}},\frac{x_2}{\sqrt{1+x_2^2}})$$

 $\Sigma(1)$ is equivalent to: for every definable $f: R \to R$ there is $\epsilon_0 \in \mathfrak{m}^{>0}$ such that st $f(\epsilon_0) = \operatorname{st} f(\epsilon)$ for all $\epsilon \in \mathfrak{m}^{>\epsilon_0}$.

Towards proving $\Sigma(1) \Rightarrow \Sigma$ we use a reduction to definable families of one-dimensional subsets of I^2 from [5] (where it is stated for the case when \mathbf{k}_{ind} is o-minimal, but the proof does not use this assumption):

Lemma 1.3 ([5]) Let $n \ge 1$, and suppose that for all definable $X \subseteq I^{1+n}$ with dim $X(r) \le 1$ for all $r \in I$ there is $\epsilon_0 \in \mathfrak{m}^{>0}$ such that st $X(\epsilon_0) = \operatorname{st} X(\epsilon)$ for all $\epsilon \in \mathfrak{m}^{>\epsilon_0}$. Then $(R, V) \models \Sigma(n)$.

The above lemma will enable us to work mainly in the plane, but at the cost of having to handle level curves of definable functions. The main result on level curves is Lemma 2.4, which is proved in the first section. Its proof uses the (two-variable version of the) Invariance of Domain Theorem for o-minimal fields by Woerheide (see [10]):

Theorem 1.4 ([10]) Every injective definable, continuous map $f: X \to \mathbb{R}^n$, where $X \subseteq \mathbb{R}^n$, is open.

We remark that it is well-known that \mathbf{k}_{ind} is always weakly o-minimal (this follows from a result by Baisalov and Poizat in [1]). An example to the effect that \mathbf{k}_{ind} is not always o-minimal is given in [8]. Results of van den Dries and Lewenberg (see [4]) show that \mathbf{k}_{ind} is o-minimal if V is T-convex, where T is the theory of R. Hrushovski, Peterzil and Pillay observe in [6] that \mathbf{k}_{ind} is o-minimal when R is sufficiently saturated and V is the convex hull of \mathbb{Q} in R.

Here is a related open question: Assume that $(R, V) \models \Sigma(1)$. Are the sets st X, with $X \subseteq R^n$ definable in (R, V), definable in \mathbf{k}_{ind} ? (A positive answer was given in [8] for the special case when R is ω -saturated and V is the convex hull of \mathbb{Q} in R.)

2 Level curves of definable functions

Definition 2.1 Let $f: X \to R$ be definable with $X \subseteq \mathbb{R}^n$. We say that ϵ_0 is good for f if $\epsilon_0 \in \mathfrak{m}^{>0}$ and st $f^{-1}(\epsilon_0) = \operatorname{st} f^{-1}(\epsilon)$ for all $\epsilon \in \mathfrak{m}^{>\epsilon_0}$.

Lemma 2.2 Suppose $(R, V) \models \Sigma(1)$, and let $f: X \to R$ be definable, with $X \subseteq R^n$ of dimension one. Then there is ϵ_0 good for f.

PROOF: The case when there is no $\delta \in \mathfrak{m}^{>0}$ and $q > \mathfrak{m}$ such that $[\delta, q] \subseteq \operatorname{im}(f)$ is trivial. So suppose $\delta \in \mathfrak{m}^{>0}$ and $q > \mathfrak{m}$ are such that $[\delta, q] \subseteq \operatorname{im}(f)$. Then $f^{-1}([\delta, q])$ has finitely many definably connected components X_1, \ldots, X_k . After possibly shrinking $[\delta, q]$ subject to the conditions $\delta \in \mathfrak{m}^{>0}$ and $q > \mathfrak{m}$, we may assume that for every i, either $f|_{X_i}$ is injective or constant.

If $f|_{X_i}$ is constant and $f(x) \in \mathfrak{m}^{>0}$, where $x \in X_i$, then set $\epsilon_i := f(x)$. If $f|_{X_i}$ is constant and $f(x) \notin \mathfrak{m}^{>0}$, for $x \in X_i$, then set $\epsilon_i = 0$. If $f|_{X_i}$ is injective, then $f|_{X_i}$ has an inverse, f^{-1} . Apply $\Sigma(1)$ to each coordinate function $(f^{-1})_j$ of f^{-1} to obtain a positive infinitesimal ϵ_{ij} such that st $f^{-1}(\epsilon) = \operatorname{st} f^{-1}(\epsilon_{ij})$ for every $\epsilon \in \mathfrak{m}^{>\epsilon_{ij}}$. Then put $\epsilon_i := \max_{j \in \{1, \dots, n\}} \epsilon_{ij}$. Now set

$$\epsilon_0 := \max_{i=1,\dots,k} \{\epsilon_i\}.$$

It is easy to see that ϵ_0 is good for f.

The following lemma is obvious.

Lemma 2.3 Let $f: X \to R$, where $X \subseteq R^n$, be definable, and let X_1, \ldots, X_k be a partition of X into definable sets. If $\epsilon_1, \ldots, \epsilon_k$ are good for $f|_{X_1}, \ldots, f|_{X_k}$, then $\max{\epsilon_1, \ldots, \epsilon_k}$ is good for f.

The proof of the next lemma is a little lengthy, but the idea is fairly straight-forward, so here's a rough outline: In order to find ϵ_0 good for $G: X \to R^{\geq 0}$ as in the statement of Lemma 2.4, we first show that, essentially, the level curves of G are all uniformly definably homeomorphic to I. Then we find a definable curve γ passing through the points of X where $|\nabla G|$ is minimal (o-minimality enables us to reduce to the case when G is sufficiently nice). We use $\Sigma(1)$ to find ϵ_0 good for $G|_{im(\gamma)}$, and for $G|_{bd(X)}$. Then we proceed to show that this ϵ_0 is also good for G. We assume towards a contradiction that it is not, hence some $a \in G^{-1}(\epsilon)$, where $\epsilon \in \mathfrak{m}^{>\epsilon_0}$, is at noninfinitesimal distance from $G^{-1}(\epsilon_0)$. We would be done if this assumption would enable us to find a definable curve in X, of length $> \mathfrak{m}$, such

-	_	_
L		
L		
L		

that at every point of the curve, the tangent vector to this curve points in the direction of ∇G , and such that the image of G restricted to the curve is $[\epsilon_0, \epsilon]$. This would put us into the paradoxical situation that the steepest path down a hill is much longer than a path which is less steep. To find a curve as described above is not necessarily possible, but we define a curve α (long, steep path down the hill) with the above properties, except that, at every point x of α , the tangent vector to α is only reasonably close to $\nabla G(x)$.

Lemma 2.4 Suppose $(R, V) \models \Sigma(1)$, and let $X \subseteq V^2$ be a closed definable set of dimension two, and let $G: X \to R^{\geq 0}$ be definable and continuous. Then there is ϵ_0 good for G.

PROOF: The points (0,1), $(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$, (1,0), $(\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}})$, (0,-1), $(-\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}})$, (-1,0), and $(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$ determine a partition of $S^1 \subseteq R^2$ into definable S_1, \ldots, S_{16} , such that $S_i \cong (0,1)$ for $i = 1, \ldots, 8$, and S_i is a singleton for $i = 9, \ldots, 16$.

By cell decomposition and Lemmas 2.2 and 2.3, we may assume that X is the closure of a cell of dimension two such that G is C^1 on X^o , and either $|\nabla G| = 0$ on X^o , or $|\nabla G| > 0$ on X^o and there is $i \in \{1, \ldots, 8\}$ satisfying $-\frac{\nabla G(x)}{|\nabla G(x)|} \in S_i$ for all $x \in X^o$. If $|\nabla G| = 0$ on X^o , then G is constant on X, so we only need to consider the second option.

The case when G(X) does not contain an interval $[\delta, q]$, with $\delta \in \mathfrak{m}^{>0}$ and $q > \mathfrak{m}$ is trivial. So let $\delta \in \mathfrak{m}^{>0}$ and $q > \mathfrak{m}$ be such that $[\delta, q] \subseteq G(X)$. Using the Trivialization Theorem (1.7, p.147 in [2]), and after possibly shrinking $[\delta, q]$, subject to $\delta \in \mathfrak{m}^{>0}$ and $q > \mathfrak{m}$, and replacing X by $G^{-1}[\delta, q]$, we obtain a definable homeomorphism

$$h: X \to [\delta, q] \times G^{-1}(\delta)$$

such that the diagram



commutes, where $X \to [\delta, q]$ is given by G and $[\delta, q] \times G^{-1}(\delta) \to [\delta, q]$ is the projection map on the first factor. So all $G^{-1}(t)$, for $t \in [\delta, q]$, are uniformly definably homeomorphic, and we may assume that dim $G^{-1}(\delta) = 1$.

By Lemma 2.3, we can reduce to the case when X is definably connected. Claim 1. $G^{-1}(\delta)$ is definably connected.

The set X is closed and definably homeomorphic to $G^{-1}(\delta) \times I$. Assume towards a contradiction that $G^{-1}(\delta) = Y_1 \dot{\cup} Y_2$, where Y_1, Y_2 are definable, nonempty and closed in $G^{-1}(\delta)$. Then Y_1, Y_2 are also closed in X, and

$$X \cong (Y_1 \dot{\cup} Y_2) \times I \cong (Y_1 \times I) \dot{\cup} (Y_2 \times I),$$

a contradiction with X being definably connected. For $x = (x_1, x_2) \in V^2$ and r > 0 we let

$$B_r(x) := [x_1 - r, x_1 + r] \times [x_2 - r, x_2 + r].$$

Claim 2. Let $Y \subseteq V^2$ be definable, definably connected, and closed. If for every $y \in Y$, there is r > 0 such that $B_r(y) \cap Y \cong I$, then either $Y \cong S^1$ or $Y \cong I$.

Let S be a stratification of Y. Then Y is a finite union of 0-dimensional cells $p_1, \ldots, p_k \in S$, and 1-dimensional cells $q_1, \ldots, q_l \in S$, where $k \ge 2$ and $l \ge 1$. Every q_i is definably homeomorphic to an open interval, and with every q_i , Y contains its endpoints among p_1, \ldots, p_k . On the other hand, every p_j is the endpoint of one or at most two of the q_i 's. Now use the fact that Y is definably connected.

Claim 3. $G^{-1}(\delta) \cong S^1$ or $G^{-1}(\delta) \cong I$.

It suffices to prove $G^{-1}(t) \cong S^1$ or $G^{-1}(t) \cong I$ for $t \in (\delta, q)$. So let $t \in (\delta, q)$, and assume towards a contradiction that $G^{-1}(t) \ncong S^1$ and $G^{-1}(t) \ncong I$. By Claims 1 and 2, we can find $a \in G^{-1}(t)$ and $r_0 > 0$ with $G^{-1}(t) \cap B_r(a) \ncong I$ for all $r \in (0, r_0)$. Hence, for all $r \in (0, r_0)$, $G^{-1}(t) \cap B_r(a)$ is definably homeomorphic to the union of $k \ge 3$ closed line segments l_1, \ldots, l_k such that each l_i has a as an endpoint, and $\bigcap_{i=1}^k l_i = \{a\}$ (we identify a here with its homeomorphic image). So let $r \in (0, r_0)$, and let

$$\phi \colon (\bigcup_{i=1}^k l_i) \times I \to X \cap B_r(a)$$

be a definable homeomorphism. We can map $(-1,1)^2$ into $\phi((l_1 \cup l_2) \times I)$ via a definable, continuous, injective map $\theta: (-1,1)^2 \to X$ such that $\theta(0,0) = a$. Then $\operatorname{im}(\theta)$ is not open in X. Now $X^o \cong (-1,1)^2$ via a definable homeomorphism ψ , so $\psi \circ \theta: (-1,1)^2 \to R^2$ yields a definable, injective,

continuous map which does not map $(-1, 1)^2$ onto an open subset of \mathbb{R}^2 , a contradiction with Theorem 1.4.

We assume from now on that $G^{-1}(t) \cong I$ for all $t \in [\delta, q]$. The case when $G^{-1}(t) \cong S^1$ is similar, but simpler, and left to the reader.

Construction of ϵ_0 good for G: Let $\{\phi_t : t \in [\delta, q]\}$ be a definable family of homeomorphisms $\phi_t : [0, 1] \to G^{-1}(t)$. For $t \in (\delta, q)$ we set

$$|\nabla G(\phi_t(0))| := \lim_{s \to 0} |\nabla G(\phi_t(s))|$$
, and $|\nabla G(\phi_t(1))| := \lim_{s \to 1} |\nabla G(\phi_t(s))|$,

where

$$\nabla G(\phi_t(s)) = \left(\frac{\partial G}{\partial x_1}(\phi_t(s)), \frac{\partial G}{\partial x_2}(\phi_t(s))\right),$$

for $s \in (0, 1)$. The above limits exist in $R \cup \{-\infty, \infty\}$ by o-minimality. Now set

$$\begin{aligned} x_t &:= \min\{|\nabla G(x)| : x \in G^{-1}(t)\}, \\ H_t &:= \{x \in G^{-1}(t) : |\nabla G(x)| = x_t\}, \end{aligned}$$

where $t \in (\delta, q)$, and let $H := \bigcup_{t \in (\delta,q)} H_t$.

By definable choice, and after replacing δ by a bigger $\delta \in \mathfrak{m}^{>0}$, and replacing q by a possibly smaller $q > \mathfrak{m}$, we can find a definable map $\gamma : [\delta, q] \to H$ with $G(\gamma(t)) = t$ for every $t \in [\delta, q]$. Next, find $\epsilon' \in \mathfrak{m}^{>0}$ good for $G|_{\mathrm{im}(\gamma)}$, and $\epsilon'' \in \mathfrak{m}^{>0}$ good for $G|_{\mathrm{bd}X}$, and set

$$\epsilon_0 := \max\{\epsilon', \epsilon''\}.$$

The rest of the proof consists of showing that ϵ_0 is indeed good for G.

First, assume towards a contradiction that $\epsilon \in \mathfrak{m}^{>\epsilon_0}$ and $a \in G^{-1}(\epsilon)$ are such that $d(a, G^{-1}(\epsilon_0)) > \mathfrak{m}$. By continuity of G, we may as well assume that $a \notin \operatorname{bd} X$. We now proceed to define what we called in the remark just before Lemma 2.4 "a long, steep path down the hill".

Construction of α : It is clear that ∇G is defined and continuous on $X^o = (G^{-1}([\delta, q]))^o$. Let *i* be such that $-\frac{\nabla G(x)}{|\nabla G(x)|} \in S_i$ on X^o , and set $w := -\frac{\nabla G(a)}{|\nabla G(a)|}$. Since $a \notin \operatorname{bd} X$, for all sufficiently small t > 0, $a + tw \in (G^{-1}([\epsilon_0, \epsilon]))^o$. Let t_0 be the first t > 0 such that $a + t_0 w \in \operatorname{bd} G^{-1}([\epsilon_0, \epsilon])$. We set $b := a + t_0 w$, and define $\alpha : [0, 1] \to X$ by

$$\alpha(t) = (1-t)a + tb.$$

From now on let α be as in "Construction of α ".

Claim 4. $G(\alpha)$ is strictly decreasing.

Assume towards a contradiction that we can find $t_1 \in (0,1)$ so that $(G \circ \alpha)'(t_1) \ge 0$. By the definition of α , $G(\alpha(0)) > G(\alpha(t))$ for all sufficiently small t > 0. Then, by the Mean Value Theorem, $(G \circ \alpha)'(t_2) = 0$ for some $t_2 \in (0, t_1]$. But

$$(G \circ \alpha)'(t_2) = \nabla G(\alpha(t_2)) \cdot D\alpha(t_2),$$

hence $\nabla G(\alpha(t_2))$ and $D\alpha(t_2)$ are orthogonal, a contradiction with the definition of α .

Claim 5. $d(a, b) > \mathfrak{m}$.

To see this, first observe that either $b \in G^{-1}(\epsilon_0)$, or $b \in \mathrm{bd}X$. If $b \in G^{-1}(\epsilon_0)$, then the claim is just the assumption that $d(a, G^{-1}(\epsilon_0)) > \mathfrak{m}$. If $b \in \mathrm{bd}X$, then use $d(a, b) + d(b, G^{-1}(\epsilon_0)) > \mathfrak{m}$ and $d(b, G^{-1}(\epsilon_0)) \in \mathfrak{m}^{>0}$ (since ϵ_0 is good for $G|_{\mathrm{bd}X}$).

Construction of β : Note that $|\nabla G(x)| > 0$ for all $x \in \operatorname{im}(\alpha|_{[0,1)})$, since $\operatorname{im}(\alpha|_{[0,1)}) \subseteq X^o$. If $|\nabla G(b)| = 0$, then replace b by an element in $\operatorname{im}(\alpha)$ which is infinitely close to the original b. Now define

$$m_0 := \min\{|\nabla G(x)| : x \in \operatorname{im}(\alpha)\} > 0.$$

For $t \in [\epsilon_0, \epsilon]$ let

$$K_t := \{ x \in G^{-1}(t) \cap X^o : |\nabla G(x)| - x_t < m_0 \text{ and } d(x, \gamma(t)) < \epsilon_0 \},\$$

note that every K_t is nonempty, and put $K := \bigcup_{t \in [\epsilon_0, \epsilon]} K_t$. By curve selection, we can find a definable map $\beta : [\epsilon_0, \epsilon] \to K$ with $G(\beta(t)) = t$.

Let $\epsilon_1, \epsilon_2 \in [\epsilon_0, \epsilon]$ be such that

$$G(b) \le \epsilon_1 < \epsilon_2 \le \epsilon,$$

 $d(c,d) > \mathfrak{m}$, for $c,d \in \operatorname{im}(\alpha)$ with $G(c) = \epsilon_1$, $G(d) = \epsilon_2$, and $\beta|_{(\epsilon_1,\epsilon_2)}$ is C^1 . By Lemma 10.8, p.204 in [3]¹ we may further assume that s is an orthogonal linear transformation of R^2 such that $s(\operatorname{im}(\beta|_{(\epsilon_1,\epsilon_2)})) = \Gamma \phi$, where $\phi: p_1^2 s(\operatorname{im}(\beta|_{(\epsilon_1,\epsilon_2)})) \to R$ is C^1 , and $|\frac{\partial \phi}{\partial x_1}| < 1$ on $p_1^2 s(\Gamma \phi)$.

¹Lemma 10.8 in [3] is an adaptation of Kurdyka and Raby's subanalytic Proposition 1.4 in [7]. In [3] it is stated for an elementary extension of an o-minimal expansion of the real field, but the proof goes through for any o-minimal field.

Let $x \in (c, d)$ and t = G(x). Then, by the definition of α ,

$$\frac{|\nabla G(x)|}{\sqrt{2}} \le |\nabla G(x) \cdot w| \le |\nabla G(x)|. \tag{1}$$

By the definition of β ,

$$|\nabla G(\beta(t))| - m_0 \le x_t \le |\nabla G(x)|,$$

 \mathbf{SO}

$$|\nabla G(\beta(t))| \le |\nabla G(x)| + m_0,$$

and thus

$$|\nabla G(\beta(t))| \le 2|\nabla G(x)|,$$

and, using inequality (1), we obtain

$$|\nabla G(\beta(t)) \cdot \tau| \le 2\sqrt{2} |\nabla G(x) \cdot w|, \tag{2}$$

where τ is the unit tangent vector to $im(\beta)$ at $\beta(t)$.

We now use α and β to define two one-variable functions, h_{α} and h_{β} respectively, which behave in a non-permissible way.

Let $h_{\beta} \colon p_1^2 \Gamma \phi \to (\epsilon_1, \epsilon_2)$ be the function

$$h_{\beta}(x) = G(s^{-1}(x, \phi(x))).$$

By the definition of β , ϵ_0 is good for $G|_{\operatorname{im}(\beta)}$, hence any two points in $\operatorname{im}(\beta)$ are infinitely close to each other. It follows that $|p_1^2\Gamma\phi| \in \mathfrak{m}^{>0}$. For $x \in p_1^2\Gamma\phi$ we have

$$|h'_{\beta}(x)| = |\nabla G(s^{-1}(x,\phi(x))) \cdot Ds^{-1}(x,\phi(x)) \cdot D(id_R,\phi)(x)|.$$

Since $Ds^{-1}(x, \phi(x)) \cdot D(id_R, \phi)(x)$ is just $c\tau$, where $c \in [1, \sqrt{2})$, and τ is the unit tangent vector to $im(\beta)$ at $s^{-1}(x, \phi(x))$, it follows that for every $x \in p_1^2 \Gamma \phi$ there is $c \in [1, \sqrt{2})$ such that

$$|h'_{\beta}(x)| = c |\nabla G(s^{-1}(x, \phi(x))) \cdot \tau|.$$
(3)

Next, note that either the projection $p_1: \mathbb{R}^2 \to \mathbb{R}: (x_1, x_2) \mapsto x_1$ or the projection $p_2: \mathbb{R}^2 \to \mathbb{R}: (x_1, x_2) \mapsto x_2$ maps (c, d) onto an open interval of

finite, noninfinitesimal length. Say $|p_1(c,d)| > \mathfrak{m}$ (the other case is similar), and assume for simplicity $c_1 < d_1$. Define $h_{\alpha}: p_1(c,d) \to (\epsilon_1, \epsilon_2)$ by

$$h_{\alpha}(x) = G((p_1|_{(c,d)})^{-1}(x)).$$

Then for all $x \in p_1(c, d)$,

$$|h'_{\alpha}(x)| = |\nabla G((p_1|_{(c,d)})^{-1}(x)) \cdot \begin{pmatrix} 1\\ \frac{d_2-c_2}{d_1-c_1} \end{pmatrix}|,$$

hence

$$|h'_{\alpha}(x)| = \lambda |\nabla G((p_1|_{(c,d)})^{-1}(x)) \cdot w|, \qquad (4)$$

for some $\lambda \in V^{>\mathfrak{m}}$.

Let $x \in (c, d)$ and $y \in im(\beta)$ be such that G(x) = G(y). Then inequality (2) and equations (3) and (4) yield that for some $\lambda \in V^{>0}$, $|h'_{\beta}(y)| \leq \lambda |h'_{\alpha}(x)|$.

Then, by Claim 4, the inverses of h_{α} and h_{β} yield two definable, continuous, strictly monotone functions $(\epsilon_1, \epsilon_2) \to R$, with the image of h_{α}^{-1} an interval of length $> \mathfrak{m}$, the image of h_{β}^{-1} an interval of infinitesimal length, and such that for every $t \in (\epsilon_1, \epsilon_2)$ there is $\lambda \in V^{>0}$ with $\lambda |(h_{\beta}^{-1}(t))'| \ge |(h_{\alpha}^{-1}(t))'|$, a contradiction.

We have shown st $G^{-1}(\epsilon) \subseteq$ st $G^{-1}(\epsilon_0)$. It is left to see that st $G^{-1}(\epsilon_0) \subseteq$ st $G^{-1}(\epsilon)$. Assume towards a contradiction that we can find $x \in G^{-1}(\epsilon_0)$ such that $d(x, G^{-1}(\epsilon)) > \mathfrak{m}$. Then we construct a definable map $[0, 1] \to X$ whose image at 0 is x, the image at 1 lies in $\mathrm{bd}G^{-1}([\epsilon_0, \epsilon])$, and the restriction of G to the image of this curve is strictly increasing, just as we constructed α , except that we set $w := \frac{\nabla G(x)}{|\nabla G(x)|}$, and we obtain a contradiction similarly as in the proof of the other inclusion.

3 $\Sigma(1) \Rightarrow \Sigma$

Definition 3.1 A small path is a definable, continuous map $\gamma: [0, r] \to X$, with definable $X \subseteq \mathbb{R}^n$ and $r \in \mathbb{R}^{>0}$, such that $\operatorname{st} \gamma(0) = \operatorname{st} \gamma(t)$ for all $t \in [0, r]$.

Lemma 3.2 Let $C = (f,g) \subseteq V^2$ be a cell of dimension two, and let $a \in \partial C$. Then there is $a' \in C$ and a small path $\gamma \colon [0,r] \to \operatorname{cl}(C)$ such that $\gamma(0) = a$, $\gamma(r) = a'$, and $\gamma(0,1] \subseteq C$. PROOF: By curve selection, we can find a definable and continuous map $\gamma: [0, r] \to X$, for some r > 0, such that $\gamma(0) = a$, and $d(a, \gamma(t)) = t$ for every $t \in [0, r]$. We may assume that $r \in \mathfrak{m}^{>0}$, and we set $a' = \gamma(r)$. Then a' and γ have the required properties.

Lemma 3.3 Let $C = (f,g) \subseteq V^2$ be a cell of dimension two, and assume that f, g are C^1 and f', g' have constant sign on p_1^2C .

- a) If $a, b \in C$ are such that st a = st b, then there is a small path $\gamma : [0, 1] \to C$ with $\gamma(0) = a$ and $\gamma(1) = b$.
- b) If $a, b \in cl(C)$ are such that st a = st b, then there is a small path $\gamma \colon [0, 1] \to cl(C)$ with $\gamma(0) = a, \gamma(1) = b$, and such that $\gamma(0, 1) \subseteq C$.

PROOF: First, let $a, b \in C$ with st $a = \operatorname{st} b$. If $a_1 = b_1$, then we define $\gamma(t) = (1-t)a + tb$, for $t \in [0,1]$. So assume that $a_1 \neq b_1$, say $a_1 < b_1$ and define $\gamma: [a_1, b_1] \to I^2$ by $\gamma(t) = (t, \frac{b_2 - a_2}{b_1 - a_1}(t - a_1) + a_2)$. Let

$$a_1 < t_1 < \dots < t_k < b_1$$

be such that

$$\gamma(t_1, t_2), \gamma(t_3, t_4), \ldots, \gamma(t_{k-1}, t_k)$$

have empty intersection with C, and

$$\gamma(a_1, t_1), \gamma(t_2, t_3), \ldots, \gamma(t_k, b_1)$$

are subsets of C. Note that then k is even, and let $i \in \{1, \ldots, \frac{k}{2}\}$. Then either $\gamma(t_{2i-1}), \gamma(t_{2i}) \in \operatorname{cl}(\Gamma f)$, or $\gamma(t_{2i-1}), \gamma(t_{2i}) \in \operatorname{cl}(\Gamma g)$. Assume $\gamma(t_{2i-1}), \gamma(t_{2i}) \in \operatorname{cl}(\Gamma f)$ (the other case is similar). Note that st $a_2 = \operatorname{st} f(t_{2i-1}) = \operatorname{st} f(t)$ for every $t \in [t_{2i-1}, t_{2i}]$, since f' has constant sign on $p_1^2 C$. Find $\delta \in \mathfrak{m}^{>0}$ such that

$$\delta < \min\{t_{2i-1} - t_{2i-2}, t_{2i+1} - t_{2i}\},\$$

and

$$\gamma_2(t_{2i-1}-\delta) - f(t_{2i-1}-\delta) \in \mathfrak{m}^{>0}, \ \gamma_2(t_{2i}+\delta) - f(t_{2i}+\delta) \in \mathfrak{m}^{>0}$$

Then g(x) - f(x) > 0 on $[t_{2i-1} - \delta, t_{2i} + \delta]$. Take $\epsilon \in \mathfrak{m}^{>0}$ such that

$$\epsilon < \min\{g(x) - f(x) : x \in [t_{2i-1} - \delta, t_{2i} + \delta]\}.$$

Define $\alpha_i \colon [t_{2i-1} - \delta, t_{2i} + \delta] \to C$ by

$$\begin{aligned} \alpha_{i}(t) &= (t, f(t) + \epsilon) \text{ for } t \in (t_{2i-1} - \frac{\delta}{2}, t_{2i} + \frac{\delta}{2}), \\ \alpha_{i}(t) &= (t, (1 - \frac{2}{\delta}(t - t_{2i-1} + \delta))\gamma_{2}(t_{2i-1} - \delta) + \frac{2}{\delta}(t - t_{2i-1} + \delta)(f(t_{2i-1} - \frac{\delta}{2}) + \epsilon)) \\ &\qquad \text{when } t \in [t_{2i-1} - \delta, t_{2i-1} - \frac{\delta}{2}], \\ \alpha_{i}(t) &= (t, (1 - \frac{2}{\delta}(t - t_{2i} - \frac{\delta}{2}))(f(t_{2i} + \frac{\delta}{2}) + \epsilon) + \frac{2}{\delta}(t - t_{2i} - \frac{\delta}{2})\gamma_{2}(t_{2i} + \delta)). \end{aligned}$$

$$\alpha_{i}(t) = (t, (1 - \frac{2}{\delta}(t - t_{2i} - \frac{\delta}{2}))(f(t_{2i} + \frac{\delta}{2}) + \epsilon) + \frac{2}{\delta}(t - t_{2i} - \frac{\delta}{2})\gamma_{2}(t_{2i} + \delta))$$

whenever $t \in [t_{2i} + \frac{\delta}{2}, t_{2i} + \delta].$

(So $\alpha_i|_{[t_{2i-1}-\delta,t_{2i-1}-\frac{\delta}{2}]}$ is a parametrization of the line segment

$$[\gamma_2(t_{2i-1}-\delta), f(t_{2i-1}-\frac{\delta}{2})+\epsilon],$$

and $\alpha_i|_{[t_{2i}+\frac{\delta}{2},t_{2i}+\delta]}$ is a parametrization of $[f(t_{2i}+\frac{\delta}{2})+\epsilon,\gamma_2(t_{2i}+\delta)]$.) For every $i \in \{1,\ldots,\frac{k}{2}\}$ replace $\gamma|_{[t_{2i}-\delta,t_{2i}+\delta]}$ by α_i . The resulting map is, after a linear change of variables, as required.

To prove the second part of the lemma, use Lemma 3.2 to find $a', b' \in C$ and small paths $\gamma_1 \colon [0, r] \to \operatorname{cl}(C), \gamma_2 \colon [0, r] \to \operatorname{cl}(C)$ such that $\gamma_1(0) = a$, $\gamma_1(r) = a', \gamma_2(0) = b, \gamma_2(r) = b'$, and $\gamma_1(0, r] \subseteq C, \gamma_2(0, r] \subseteq C$. Then use the first part of the lemma to find a small path $\gamma_3 \colon [0, 1] \to C$ with $\gamma_3(0) = a', \gamma_3(1) = b'$, and $\operatorname{im}(\gamma) \subseteq C$. After a linear changes of variables, $\gamma_1 \colon [0, \frac{1}{3}] \to \operatorname{cl}(C), \gamma_3 \colon [\frac{1}{3}, \frac{2}{3}] \to C$, and $\gamma_2 \colon [\frac{2}{3}, 1] \to \operatorname{cl}(C)$. Then define $\gamma \colon [0, 1] \to \operatorname{cl}(C)$ by $\gamma(t) = \gamma_1(t)$ if $t \in [0, \frac{1}{3}], \gamma(t) = \gamma_3(t)$ if $t \in (\frac{1}{3}, \frac{2}{3})$, and $\gamma(t) = \gamma_2(t)$ if $t \in [\frac{2}{3}, 1]$. Clearly, γ satisfies the requirements.

Lemma 3.4 Let $(f,g) \subseteq V^2$ be a cell of dimension two such that f, g are C^1 and f', g' have constant sign on $p_1^2(f,g)$. Let $h: (f,g) \to R$ be definable and C^1 and such that $\left|\frac{\partial h}{\partial x_1}\right|, \left|\frac{\partial h}{\partial x_2}\right| < 1$. Then st $h(a) = \operatorname{st} h(b)$ whenever $a, b \in (f,g)$ are such that st $a = \operatorname{st} b$. If moreover h extends continuously to clh: $\operatorname{cl}(f,g) \to R$, then st $\operatorname{clh}(a) = \operatorname{st} \operatorname{clh}(b)$ whenever $a, b \in \operatorname{cl}(f,g)$ with st $a = \operatorname{st} b$.

PROOF: Let $a, b \in (f, g)$ be such that st(a) = st(b). Then, by Lemma 3.3, we can find a small path $\gamma: [0, 1] \to (f, g)$ with $\gamma(0) = a$ and $\gamma(1) = b$.

By Lemma 10.8 in [3], we can partition $\operatorname{im}(\gamma)$ into definable X_1, \ldots, X_{k+1} , where dim $X_{k+1} = 0$, and we can find orthogonal linear transformations s_1, \ldots, s_k of R^2 , such that for $i = 1, \ldots, k$, $s_i X_i$ is the graph of a definable C^1 -function $\phi_i \colon p_1^2 s_i X_i \to V$ on an open $p_1^2 s_i X \subseteq V$ with $|\phi'_i| < 1$ on $p_1^2 s_i X_i$. We may assume that each $\Gamma \phi_i$ is a cell, and we let $\operatorname{cl}\phi_i$ be the continuous extension of ϕ_i to $\operatorname{cl}(p_1^2 s_i X_i) \to R$.

Define $H_i: \operatorname{cl}(p_1^2 s_i X_i) \to R$ by

$$H_i(x) = h(s_i^{-1}(x, \operatorname{cl}\phi_i(x))).$$

Assume towards a contradiction that $\operatorname{st} h(a) \neq \operatorname{st} h(b)$. For $i = 1, \ldots, k$, let $x_i, y_i \in R$ be such that $(x_i, y_i) = p_1^2 s_i X_i$. Then for some i, $\operatorname{st} H_i(x_i) \neq \operatorname{st} H_i(y_i)$. However, $|x_i - y_i| \in \mathfrak{m}^{>0}$ (as s_i is an isometry), and

$$H'_{i}(t)| = |\nabla h(s_{i}^{-1}(t,\phi_{i}(t))) \cdot Ds_{i}^{-1}(t,\phi_{i}(t)) \cdot D(id_{R},\phi_{i})(t)| \in V^{>0},$$

for $t \in (x_i, y_i)$, a contradiction with the mean value theorem.

Assume now that h extends continuously to $clh: cl(f,g) \to R$ and let $a, b \in cl(f,g)$ be such that st a = st b. By continuity of clh and the proof of Lemma 3.2 we can find $a', b' \in (f,g)$ infinitely close to a and to b respectively, and small paths $\alpha: [0,r] \to cl(f,g), \beta: [0,r] \to cl(f,g)$, where r > 0, so that $\alpha(0) = a, \alpha(r) = a', \beta(0) = b, \beta(r) = b', \alpha(0,r], \beta(0,r] \subseteq (f,g)$, and

$$|\operatorname{cl} h(a) - h(\alpha(t))|, |\operatorname{cl} h(b) - h(\beta(t))| \in \mathfrak{m}^{>0}$$

for all $t \in (0, r]$. Now use the first part of the lemma to find a small path $\gamma: [0, 1] \to (f, g)$ with $\gamma(0) = a'$ and $\gamma(1) = b'$. It is now obvious how to construct the desired map from α, β , and γ .

Lemma 3.5 Let $C = (f,g) \subseteq V^2$ be a cell of dimension two such that f, g are C^1 and f', g' have constant sign on p_1^2C . Let further $h: C \to R$ be definable and C^1 , with $|\frac{\partial h}{\partial x_1}|$, $|\frac{\partial h}{\partial x_2}| < 1$, and $\Gamma h \subseteq V^3$. Then h extends continuously to a definable function $clh: cl(C) \to R$.

PROOF: We first show that $cl(\Gamma h)$ is the graph of a function. By Lemma 3.4, st Γh is the graph of a function, and st $\Gamma h = st cl(\Gamma h)$. So if there are $x, y \in cl(\Gamma h)$ with $p_2^3 x = p_2^3 y$ and $x_3 \neq y_3$, then $|x_3 - y_3| \in \mathfrak{m}^{>0}$. Set $\delta := |x_3 - y_3|$, let \mathfrak{m}' be the maximal ideal

$$\{x \in R : |x| < \delta q \text{ for all } q \in \mathbb{Q}^{>0}\},\$$

and let V' be the corresponding valuation ring with residue map st': $V' \to \mathbf{k}'$. Then $V \subseteq V'$. Since Lemma 3.4 holds for any choice of V, st' $\Gamma h = \text{st'} \operatorname{cl}(\Gamma h)$ is again the graph of a function, a contradiction with $\delta \notin \mathfrak{m}'$.

By Lemma 1.7, p.95 in [2], $\operatorname{cl}(C) = p_2^3 \operatorname{cl}(\Gamma h)$. We let $\operatorname{cl} h$ be the function $\operatorname{cl}(C) \to R$ whose graph is the set $\operatorname{cl}(\Gamma h)$. To see that $\operatorname{cl} h$ is continuous, assume towards a contradiction that $a \in \operatorname{cl}(C)$ and $\epsilon > 0$ are such that for every $\delta > 0$, we can find $x_{\delta} \in \operatorname{cl} C$ with $\operatorname{d}(x_{\delta}, a) < \delta$ and $|\operatorname{cl} h(a) - \operatorname{cl} h(x_{\delta})| > \epsilon$. Then curve selection yields a definable and continuous $\gamma \colon (0, r] \to \operatorname{cl}(C)$, r > 0, with $\lim_{t\to 0} \gamma(t) = a$, $\operatorname{d}(\gamma(t), a) = t$, and $|\operatorname{cl} h(\gamma(t)) - \operatorname{cl} h(a)| > \epsilon$. Then, since $\operatorname{cl}(\Gamma h)$ is bounded, $\lim_{t\to 0} \operatorname{cl} h(\gamma(t))$ exists in R, and since $\operatorname{cl}(\Gamma h)$ is closed, $\lim_{t\to 0} \operatorname{cl} h(\gamma(t)) \in \Gamma \operatorname{cl} h$, a contradicition with $\operatorname{cl} h$ being a function. \Box

Theorem 3.6 $\Sigma(1) \Rightarrow \Sigma$.

PROOF: Assume that $(R, V) \models \Sigma(1)$, and let $X \subseteq I^{1+n}$ be definable. By Lemmas 1.3 and 2.2, towards proving $\Sigma(n)$, we may assume that X is of dimension two, and $p_1^{n+1}X = (0, 1)$.

By Lemma 10.8 from [3], we can find a partition of X into definable sets X_1, \ldots, X_{k+1} , where dim $X_{k+1} < 2$, and orthogonal linear transformations s_1, \ldots, s_k of R^{n+1} , such that each $s_i X_i$ is the graph of a definable C^1 -map $\phi_i: p_2^{n+1} s_i X_i \to R^{n-1}$ on an open $p_2^{n+1} s_i X_i \subseteq R^2$ and $|\frac{\partial \phi_{ij}}{\partial x_l}| < 1$ on $p_2^{n+1} s_i X_i$ for l = 1, 2 and $j = 1, \ldots, n-1$.

Let $i \in \{1, \ldots, k\}$, and set, for the sake of simplicity, $X := X_i$, $s := s_i$, and $\phi := \phi_i$. By cell decomposition, we may assume that sX is a cell such that $p_2^{n+1}sX = (g, h)$, where $g, h : p_1^2(g, h) \to R$ are definable and C^1 , and g', h' have constant sign on $p_1^2(g, h)$.

By Lemma 3.5 each coordinate function ϕ_j of ϕ extends continuously to

$$\operatorname{cl}\phi_j \colon \operatorname{cl}(p_2^{n+1}sX) \to R.$$

We set

$$\mathrm{cl}\phi := (\mathrm{cl}\phi_1, \ldots, \mathrm{cl}\phi_{n-1}),$$

and one easily checks, using $cl(\Gamma\phi_j) = \Gamma(cl\phi_j)$ and the continuity of $cl\phi_j$ for all j, that

$$\operatorname{cl}(X) = s^{-1}(\operatorname{cl}(sX)) = s^{-1}(\Gamma \operatorname{cl}\phi).$$

Let $F: cl(X) \to [0,1]$ be the coordinate projection $(x_1, \ldots, x_{n+1}) \mapsto (x_1)$, and define $G: cl(p_2^{n+1}sX) \to R^{\geq 0}$ by

$$G(x) := F(s^{-1}(x, \operatorname{cl}\phi(x))).$$

Since G is continuous, Lemma 2.4 yields ϵ_0 good for G. We claim that ϵ_0 is also good for F.

To see that st $F^{-1}(\epsilon) \subseteq$ st $F^{-1}(\epsilon_0)$, let $x \in F^{-1}(\epsilon)$, for some $\epsilon \in \mathfrak{m}^{>\epsilon_0}$. Our aim is to show that $d(x, F^{-1}(\epsilon_0)) \in \mathfrak{m}^{>0}$. We have $p_2^{n+1}s(x) \in G^{-1}(\epsilon)$, hence $d(p_2^{n+1}s(x), G^{-1}(\epsilon_0)) = \delta$ for some $\delta \in \mathfrak{m}^{>0}$. Take $y \in G^{-1}(\epsilon_0)$ such that $d(p_2^{n+1}s(x), y) = \delta$. Then, by the assumption on g' and h', and since $|\frac{\partial \phi_j}{\partial x_1}|, |\frac{\partial \phi_j}{\partial x_2}| < 1$ for $j = 1, \ldots, n-1$, Lemma 3.4 applied to each coordinate function of ϕ yields $d(sx, (y, \operatorname{cl}\phi(y))) \in \mathfrak{m}^{>0}$. Now $F(s^{-1}(y, \operatorname{cl}\phi(y)) = \epsilon_0$, and, since s is an isometry,

$$d(x, s^{-1}(y, cl\phi(y))) \in \mathfrak{m}^{>0}.$$

Similarly, one shows that st $F^{-1}(\epsilon_0) \subseteq \operatorname{st} F^{-1}(\epsilon)$.

By Lemma 2.2, we can find ϵ_{k+1} good for $X_{k+1} \to (0, 1) : x \mapsto x_1$, and by the above, we can find ϵ_i good for $X_i \to (0, 1) : x \mapsto x_1$ for each $i = 1, \ldots, k$. Then $\epsilon_0 := \max_{i=1,\ldots,k+1} \{\epsilon_i\}$ is such that st $X(\epsilon_0) = \operatorname{st} X(\epsilon)$ for all $\epsilon \in \mathfrak{m}^{>\epsilon_0}$.

Theorem 1.2 from the introduction now follows.

ACKNOWLEDGMENTS An earlier version of the proof of Lemma 2.4 uses a consequence of Shiota's Hauptvermutung for o-minimal fields (see [9]), instead of the Invariance of Domain Theorem. The present version is shorter, but nevertheless inspired by Shiota's striking result.

I thank Lou van den Dries for his interest in and comments on this paper.

References

- Y. Baisalov, B. Poizat, Paires de Structures O-Minimales. J. Symb. Log. 63(2): 570-578 (1998).
- [2] L. van den Dries, *Tame topology and o-minimal structures*. London Mathematical Society Lecture Note Series, vol. 248, Cambridge University Press (1998).
- [3] L. van den Dries, *Limit sets in o-minimal structures*. Proceedings of the RAAG summer school, O-minimal structures, Lisbon 2003.

- [4] L. van den Dries, A. H. Lewenberg, *T*-convexity and tame extensions. J. Symb. Log. 60: 74-102 (1995).
- [5] L. van den Dries, J. Maříková, Triangulation in o-minimal fields with standard part map, preprint (arXiv:0901.2339).
- [6] E. Hrushovski, Y. Peterzil, A. Pillay, Groups, measures and the NIP, J. Amer. Math. Soc. 21 (2008), 563-596.
- [7] K. Kurdyka, G. Raby, Densité des ensembles sous-analytiques, Ann. Inst. Fourier 139 (1989), 753-771.
- [8] J. Maříková, *O-minimal fields with standard part map*, preprint (arXiv:0901.2337).
- [9] M. Shiota, *PL* and differential topology in o-minimal structures, preprint.
- [10] A. Woerheide, O-minimal homology, PhD. Thesis (1996), University of Illinois at Urbana-Champaign.