

BANACH-STONE-LIKE RESULTS FOR COMBINATORIAL BANACH SPACES

C. BRECH AND C. PIÑA

ABSTRACT. We show that under a certain topological assumption on two compact hereditary families \mathcal{F} and \mathcal{G} on some infinite cardinal κ , the corresponding combinatorial spaces $X_{\mathcal{F}}$ and $X_{\mathcal{G}}$ are isometric if and only if there is a permutation of κ inducing a homeomorphism between \mathcal{F} and \mathcal{G} . We also prove that two different regular families \mathcal{F} and \mathcal{G} on ω cannot be permuted one to the other. Both these results strengthen the main result of [4].

The classical Banach-Stone theorem states that, given two compact spaces K and L , $C(K)$ and $C(L)$ are isometric Banach spaces if and only if K and L are homeomorphic (see e.g. [12, Theorem 7.8.4]). Our main purpose in this paper is to prove versions of this result in the context of combinatorial Banach spaces.

Given a family \mathcal{F} of finite subsets of an infinite cardinal κ , it can be seen as a topological subspace of 2^κ . If \mathcal{F} contains all singletons of κ and is closed under subsets, the combinatorial Banach space $X_{\mathcal{F}}$ (also called the generalized Schreier space) is the completion of $c_{00}(\kappa)$, the vector space of finitely supported scalar sequences, with respect to the norm:

$$\|x\| = \sup \left\{ \sum_{\alpha \in s} |x(\alpha)| : s \in \mathcal{F} \right\}.$$

Notice that the sequence of unit vectors (e_α) forms a (long) Schauder basis of $X_{\mathcal{F}}$, which is considered to be the canonical one and is 1-unconditional. Moreover, if \mathcal{F} is compact, the canonical basis is shrinking.

Both spaces $X_{\mathcal{F}}$ and $C(\mathcal{F})$ can be seen as Banach spaces counterparts of the family \mathcal{F} and they share some properties: for instance, both contain homeomorphic copies of \mathcal{F} in the dual unit sphere equipped with the weak* topology. Banach-Stone theorem guarantees that the isometric structure of $C(\mathcal{F})$ carries exactly the topological structure of \mathcal{F} . In Section 2 we prove our first main result, Theorem 7, and give in Corollary 14 sufficient topological assumptions on two families \mathcal{F} and \mathcal{G} , so that $X_{\mathcal{F}}$ and $X_{\mathcal{G}}$ are isometric if and only if \mathcal{F} and \mathcal{G} are π -homeomorphic (that is, there is a homeomorphism between \mathcal{F} and \mathcal{G} which is induced by a permutation of κ , see Section 1). Hence, the isometric structure of $X_{\mathcal{F}}$ carries more than the topological structure of \mathcal{F} .

Isometries between combinatorial spaces have been recently analysed in [1, 4]. In [1, Theorem 5.1], the authors prove that any isometry of the real combinatorial space of a Schreier family of finite order

1991 *Mathematics Subject Classification.* Primary 46B45; Secondary 03E75.

Key words and phrases. Banach-Stone Theorem, combinatorial Banach spaces, compact and spreading families.

The first author was supported by CNPq grant (308183/2018-5) and by FAPESP grant (2016/25574-8). The second author was supported by the postdoctoral program of Vicerrectoría de Investigación y Extensión de la UIS.

is determined by a change of signs of the elements of the canonical basis. This has been generalized in [4] to real and complex combinatorial spaces of regular families on the set of the integers ω (see Definition 10). Our proof follows similar lines as theirs, but we extracted a sufficient topological condition which made it possible to enlarge the class of families for which any isometry between two combinatorial spaces is given by a signed permutation of the canonical bases: it includes now families on ω which are not spreading, as well as families on uncountable cardinals κ . Moreover, Arens and Kelley [2] gave a simple proof of the Banach-Stone theorem using the fact that isometries take extreme points to extreme points. It was improved to a noncommutative version for C^* -algebras by Kadison [7]. Our proof also uses this fact, as does that of [4].

Theorem 7 states that any isometry between $X_{\mathcal{F}}$ and $X_{\mathcal{G}}$ is induced by a permutation of κ and a sequence of signs for a class of families. If we take \mathcal{F} to be the family of singletons of ω , we get that $X_{\mathcal{F}} = c$, the Banach space of convergent sequences of scalars, with the supremum norm. Taking \mathcal{F} to be the family of all finite subsets of κ makes $X_{\mathcal{F}}$ to be $\ell_1(\kappa)$. Hence, similarly to what is mentioned in [4], our results can be compared to the fact that isometries of the spaces c_0 or ℓ_p , $1 \leq p < \infty$, $p \neq 2$, are all signed permutations of the canonical unit basis (see e.g. [8, Theorem 2.f.14]).

Our second main result, Theorem 13, states that two different regular families on ω are not π -homeomorphic. A permutation of ω which takes \mathcal{F} onto \mathcal{G} and a sequence of signs induce an isometry between $X_{\mathcal{F}}$ and $X_{\mathcal{G}}$. [4, Theorem 10] shows that the converse is also true for regular families \mathcal{F} and \mathcal{G} : any isometry between $X_{\mathcal{F}}$ and $X_{\mathcal{G}}$ is induced by a permutation of ω and a sequence of signs. Our Theorem 13 shows that under any of these two equivalent assumptions - \mathcal{G} being the image of \mathcal{F} under a permutation of ω or $X_{\mathcal{F}}$ and $X_{\mathcal{G}}$ being isometric - \mathcal{F} and \mathcal{G} are just the same family. Meaning that these morphisms preserve not only the structure of the family, but the family itself.

The crucial difference between the hypotheses of theorems 7 and 13 is the assumption that the families are spreading (see Definition 10). A compact family can only happen to be spreading for families on ω . The properties of being compact or hereditary are preserved under permutations of κ , as a permutation induces a homeomorphism between a family and its image. However, this is not the case for spreading. This lead us to suspect that being spreading is a strong assumption for the classification of the isometries between combinatorial spaces obtained in [4] and this is indeed the case, as guarantees Theorem 13.

Another implication of a family being spreading is that the Cantor-Bendixson index of the singletons $\{n\}$ is a monotone nondecreasing sequence of ordinals. This fact is explored in more detail in the last section, where we also give further examples and remarks related to both the countable and the general setting.

The paper is organized as follows. In Section 1 we introduce definitions, notations and basic facts to be used in the following ones. In Section 2 we present Theorem 7, which characterizes the isometries

on combinatorial spaces of families in a broader class than the class of regular families, including families on uncountable cardinals. In Section 3, we show that two different regular families cannot be permuted one to the other: this is Theorem 13 and its proof is purely combinatorial. We finish the paper exploring more about the topology and giving examples and remarks presented in Section 4.

1. PRELIMINARIES

Given an infinite cardinal κ , we will denote by $[\kappa]^{<\omega}$ the set of all finite subsets of κ . Given $n \in \omega$ and $M \subseteq \kappa$, we denote by $[M]^n$ the collection of all subsets of M of size n , by $[M]^{<n}$ the collection of all subsets of M of size less than n , and by $[M]^{\leq n}$ the union $[M]^n \cup [M]^{<n}$. If $s \in [\kappa]^n$, we write $s = \{\alpha_1, \dots, \alpha_n\}_<$ to indicate that the enumeration is strictly increasing.

We introduce now some notions about families:

Definition 1. A family \mathcal{F} on κ is a set $\mathcal{F} \subseteq [\kappa]^{<\omega}$ and we will always assume that $[\kappa]^{\leq 1} \subseteq \mathcal{F}$.

- We say that a family \mathcal{F} is hereditary if it is closed under subsets: $s \in \mathcal{F}$ whenever $t \in \mathcal{F}$ and $s \subseteq t$.
- A family \mathcal{F} on κ is a topological space when endowed with the subspace topology inherited from 2^κ , where each $s \in \mathcal{F}$ is identified with its characteristic function and 2^κ has the product topology.
- Any topological assumption about \mathcal{F} refers to this topology. For instance, \mathcal{F} is compact if it is a closed subspace of 2^κ .

Given a family \mathcal{F} , let \mathcal{F}^{MAX} denote the set of maximal elements of \mathcal{F} with respect to the inclusion and $\overline{\mathcal{F}}^\subseteq$ denote its downwards closure, that is:

$$\mathcal{F}^{MAX} = \{s \in \mathcal{F} : \nexists t \in \mathcal{F} \text{ such that } s \subsetneq t\} \quad \text{and} \quad \overline{\mathcal{F}}^\subseteq = \{s : \exists t \in \mathcal{F} \text{ such that } s \subseteq t\}.$$

A permutation of κ is simply a bijection $\pi : \kappa \rightarrow \kappa$ and if $s \subseteq \kappa$, then $\pi[s] = \{\pi(\alpha) : \alpha \in s\}$. Any permutation π of κ induces a homeomorphism $\hat{\pi} : 2^\kappa \rightarrow 2^\kappa$, defined by $\hat{\pi}(x)(\alpha) = x(\pi^{-1}(\alpha))$. We chose to use here π^{-1} instead of π so that we have the following: if $\pi[\mathcal{F}]$ denotes the image of \mathcal{F} by $\hat{\pi}$, then it coincides with $\{\pi[s] : s \in \mathcal{F}\}$. We say that two families \mathcal{F} and \mathcal{G} on κ are π -homeomorphic if there is a permutation π of κ such that $\mathcal{G} = \pi[\mathcal{F}]$.

By an isometry between Banach spaces we mean a linear bijective operator which preserves the norm. If X is a Banach space, X^* denotes its topological dual, B_X is the unit ball and B_{X^*} is the dual unit ball.

Throughout the paper we will be concerned about conditions on two families \mathcal{F} and \mathcal{G} which guarantee that all or some of the following statements are equivalent and examples to show when they are not:

- (i) $\mathcal{F} = \mathcal{G}$.

- (ii) \mathcal{F} and \mathcal{G} are π -homeomorphic.
- (iii) $X_{\mathcal{F}}$ and $X_{\mathcal{G}}$ are isometric.
- (iv) $X_{\mathcal{F}}^*$ and $X_{\mathcal{G}}^*$ are isometric.

Notice that these statements are decreasing in strength. (i) clearly implies (ii) and if π is a permutation of κ witnessing (ii), then, given any sequence (θ_α) of scalars such that $|\theta_\alpha| = 1$, there is an isometry $T : X_{\mathcal{F}} \rightarrow X_{\mathcal{G}}$ satisfying $T(e_\alpha) = \theta_\alpha e_{\pi(\alpha)}$ for all $\alpha \in \kappa$, so that (ii) implies (iii). (iii) implies (iv) easily, as the adjoint operator of any isometry between two Banach spaces is an isometry between their dual spaces. Moreover, in case T is such that $T(e_\alpha) = \theta_\alpha e_{\pi(\alpha)}$ for every $\alpha \in \kappa$, some sequence of scalars (θ_α) and a permutation π , then clearly $T^*(e_\alpha^*) = \theta_\alpha e_{\pi^{-1}(\alpha)}^*$ for all $\alpha \in \kappa$.

Also, we will be mostly working with hereditary families, so that conditions (i) and (ii) are clearly equivalent to conditions (i') and (ii') below, and we will use them interchangeably:

- (i') $\mathcal{F}^{MAX} = \mathcal{G}^{MAX}$.
- (ii') \mathcal{F}^{MAX} and \mathcal{G}^{MAX} are π -homeomorphic.

2. ISOMETRIES BETWEEN COMBINATORIAL SPACES

The purpose of this section is to (essentially) improve the results of [4] by getting that two families are π -homeomorphic from weaker assumptions on them. Our first purpose was to replace the spreading condition by some condition which does not take the order into account. This lead us to a much more general version, for compact and hereditary families, on any infinite cardinal κ , satisfying some topological condition. The existence of nontrivial compact and hereditary families on uncountable cardinals was already known in [9], and much more complex families were explored in [5]. We recall some of these examples and discuss these briefly in Section 4.

Our proof follows the path of the proof of Theorem 10 of [4]. In particular, we will also make use of extreme points in our proof. Given a Banach space X , recall that x is an extreme point of a convex set $B \subseteq X$ if there are no $x_1 \neq x_2$ in B such that $x = \alpha x_1 + (1 - \alpha)x_2$ for some $0 < \alpha < 1$. Let $Ext(B_{X^*})$ be the set of extreme points of B_{X^*} .

We will use the characterization of the extreme points of $B_{X_{\mathcal{F}}^*}$ stated in [6], and proved and used in [1] and [4] in more particular cases. Actually, despite being stated for regular families in [4, Proposition 5], a careful analysis of its proof shows that the characterization holds for any compact and hereditary family \mathcal{F} . For completeness purposes, we reproduce their proof here with the required adaptations.

Let \mathbb{K} be the field of scalars \mathbb{R} or \mathbb{C} . Given a Banach space X , we say that a subset $N \subseteq B_{X^*}$ is *norming* if $\|x\| = \sup\{|x^*(x)| : x^* \in N\}$ for every $x \in X$, and N is *sign invariant* if for any sign $\theta \in \mathbb{K}$, we have $\theta N = N$. The following classical lemma was stated as Lemma 4 of [4]:

Lemma 2. *Let X be a Banach space over \mathbb{K} , and let $N \subseteq B_{X^*}$ be a sign invariant norming set for X . Then $B_{X^*} = \overline{\text{conv}(N)}^{w*}$.*

To prove the characterization of the extreme points in our more general setting, we replace weak* convergence of sequences by the following auxiliary lemma:

Lemma 3. *Let \mathcal{F} be a compact and hereditary family on κ . Then*

$$\left\{ \sum_{\xi \in s} \theta_\xi e_\xi^* : s \in \mathcal{F} \text{ and } (\theta_\xi)_{\xi \in s} \text{ is a sequence of scalars with } |\theta_\xi| = 1 \right\}$$

is weak-closed.*

Proof. Let

$$M = \left\{ \sum_{\xi \in s} \theta_\xi e_\xi^* : s \in \mathcal{F} \text{ and } (\theta_\xi)_{\xi \in s} \text{ is a sequence of scalars with } |\theta_\xi| = 1 \right\}$$

and let $x^* \in \overline{M}^{w^*}$. For each $\xi \in \kappa$, notice that

$$|x^*(e_\xi)| \in \overline{\{|y^*(e_\xi)| : y^* \in M\}} = \{0, 1\}.$$

Hence, there is $s \subseteq \kappa$ and a sequence $(\theta_\xi)_{\xi \in s}$ in \mathbb{K} with $|\theta_\xi| = 1$ such that $x^* = \sum_{\xi \in s} \theta_\xi e_\xi^*$. It remains to prove that $s \in \mathcal{F}$.

Let us show that $[s]^{<\omega} \subseteq \mathcal{F}$. For each finite $t \subseteq s$, there is $y_t^* \in M$ such that $|x^*(e_\xi) - y_t^*(e_\xi)| < \frac{1}{2}$ for every $\xi \in t$. In particular, this implies that $|y_t^*(e_\xi)| = 1$ for every $\xi \in t$, so that $t \subseteq \text{supp } y_t^* \in \mathcal{F}$. Since \mathcal{F} is hereditary, we get that $t \in \mathcal{F}$.

Finally, s has to be finite since \mathcal{F} is compact. In particular, $s \in \mathcal{F}$. \square

We are now ready to reproduce the proof given in [4, Proposition 5] of the following characterization:

Proposition 4. *If \mathcal{F} is a compact and hereditary family on an infinite cardinal κ , then*

$$\text{Ext}(B_{X_{\mathcal{F}}^*}) = \left\{ \sum_{\alpha \in s} \theta_\alpha e_\alpha^* : s \in \mathcal{F}^{MAX} \text{ and } (\theta_\alpha)_{\alpha \in s} \text{ is a sequence of scalars with } |\theta_\alpha| = 1 \right\}.$$

Proof. Take $N := \{\sum_{\alpha \in s} \theta_\alpha e_\alpha^* : s \in \mathcal{F}^{MAX}, |\theta_\alpha| = 1\}$ and $M := \{\sum_{\alpha \in s} \theta_\alpha e_\alpha^* : s \in \mathcal{F}, |\theta_\alpha| = 1\}$. M is clearly norming for $X_{\mathcal{F}}$ and sign invariant. It follows from Lemma 2 that $B_{X_{\mathcal{F}}^*} = \overline{\text{conv}(M)}^{w^*}$. By Lemma 3, M is w^* -closed, so that both M and $B_{X_{\mathcal{F}}^*} = \overline{\text{conv}(M)}^{w^*}$ are compact in the locally convex space $(X_{\mathcal{F}}^*, w^*)$. It follows from Milman's theorem (see [11], Theorem 3.25) that every extreme point of $B_{X_{\mathcal{F}}^*}$ lies in M . Also, $(M \setminus N) \cap \text{Ext}(B_{X_{\mathcal{F}}^*}) = \emptyset$, as any $x \in M \setminus N$ can be written as $x = \frac{(x+e_\alpha^*)+(x-e_\alpha^*)}{2}$, where $\text{supp } x \cup \{\alpha\} \in \mathcal{F}$. Since $N \subseteq \text{Ext}(B_{X_{\mathcal{F}}^*})$ we conclude that $\text{Ext}(B_{X_{\mathcal{F}}^*}) = N$. \square

We prove another auxiliary lemma before getting our improved version of [4, Proposition 10]

Lemma 5. *Let \mathcal{F} be a compact and hereditary family on κ . Given $\alpha \in \kappa$, if $\{\alpha\} \in \overline{\mathcal{F}^{MAX}}$, then $e_\alpha^* \in \overline{\text{Ext}(B_{X_{\mathcal{F}}^*})}^{w^*}$.*

Proof. Fix $\alpha \in \kappa$ such that $\{\alpha\} \in \overline{\mathcal{F}^{MAX}}$ and let $\varepsilon > 0$ and $x_1, \dots, x_n \in X_{\mathcal{F}}$. We have to show that there is $x^* \in \text{Ext}(B_{X_{\mathcal{F}}}^*)$ such that $|e_{\alpha}^*(x_i) - x^*(x_i)| < \varepsilon$ for all $1 \leq i \leq n$.

Take $y_1, \dots, y_n \in c_{00}(\kappa)$ such that $\|y_i - x_i\| < \frac{\varepsilon}{2}$ for every $1 \leq i \leq n$ and take $F = \bigcup \{\text{supp } y_i : 1 \leq i \leq n\}$. Since F is finite and $\{\alpha\} \in \overline{\mathcal{F}^{MAX}}$, there is $s \in \mathcal{F}^{MAX}$ such that $\alpha \in s$ and $s \cap F \subseteq \{\alpha\}$.

Let $x^* = \sum_{\xi \in s} e_{\xi}^*$ and notice that $x^* \in \text{Ext}(B_{X_{\mathcal{F}}}^*)$ by Proposition 4. Also, for every $1 \leq i \leq n$, since $\text{supp } y_i \cap s \subseteq \{\alpha\}$, we have that

$$|e_{\alpha}^*(y_i) - x^*(y_i)| = \left| \sum_{\xi \in s \setminus \{\alpha\}} e_{\xi}^*(y_i) \right| = 0.$$

Finally, since $\|e_{\alpha}^*\| = 1$ and $\|x^*\| = 1$, we conclude that

$$\begin{aligned} |e_{\alpha}^*(x_i) - x^*(x_i)| &\leq |e_{\alpha}^*(x_i) - e_{\alpha}^*(y_i)| + |e_{\alpha}^*(y_i) - x^*(y_i)| + |x^*(y_i) - x^*(x_i)| \\ &\leq \|e_{\alpha}^*\| \cdot \|x_i - y_i\| + \|x^*\| \cdot \|x_i - y_i\| < \varepsilon, \end{aligned}$$

and this concludes the proof that $e_{\alpha}^* \in \overline{\text{Ext}(B_{X_{\mathcal{F}}}^*)}^{w^*}$. \square

Proposition 6. *Let \mathcal{F} and \mathcal{G} be compact and hereditary families on an infinite cardinal κ with the additional property that $[\kappa]^1 \subseteq \overline{\mathcal{F}^{MAX}}$. If $T : X_{\mathcal{F}} \rightarrow X_{\mathcal{G}}$ is weak*-weak* continuous and preserves extreme points, then for every $\alpha \in \kappa$ there is $s_{\alpha} \in \mathcal{G}$ and a sequence $(\theta_{\xi}^{\alpha})_{\xi \in s_{\alpha}}$ with $|\theta_{\xi}^{\alpha}| = 1$ such that $T(e_{\alpha}^*) = \sum_{\xi \in s_{\alpha}} \theta_{\xi}^{\alpha} e_{\xi}^*$.*

Proof. Fix $\alpha \in \kappa$ and since $\{\alpha\} \in \overline{\mathcal{F}^{MAX}}$, then by Proposition 4 and Lemma 5,

$$e_{\alpha}^* \in \overline{\text{Ext}(B_{X_{\mathcal{F}}}^*)}^{w^*} \subseteq \overline{B_{X_{\mathcal{F}}}^*}^{w^*} = B_{X_{\mathcal{F}}}^*.$$

Since T is weak*-weak*-continuous and takes extreme points to extreme points, we get that:

$$T(e_{\alpha}^*) \in T[\overline{\text{Ext}(B_{X_{\mathcal{F}}}^*)}^{w^*}] \subseteq T[\overline{\text{Ext}(B_{X_{\mathcal{F}}}^*)}]^{w^*} \subseteq \overline{\text{Ext}(B_{X_{\mathcal{G}}}^*)}^{w^*}.$$

By Lemma 3 for \mathcal{G} we get that there is $s_{\alpha} \in \mathcal{G}$ and a sequence $(\theta_{\xi}^{\alpha})_{\xi \in s_{\alpha}}$ with $|\theta_{\xi}^{\alpha}| = 1$ such that $T(e_{\alpha}^*) = \sum_{\xi \in s_{\alpha}} \theta_{\xi}^{\alpha} e_{\xi}^*$. \square

We can now establish the main result of this section:

Theorem 7. *Let \mathcal{F} and \mathcal{G} be compact and hereditary families on κ with the additional property that*

$$[\kappa]^1 \subseteq \overline{\mathcal{F}^{MAX}} \cap \overline{\mathcal{G}^{MAX}}.$$

If $T : X_{\mathcal{F}} \rightarrow X_{\mathcal{G}}$ is an isometry, then there exists a permutation π of κ and a sequence of signs $(\theta_{\alpha})_{\alpha}$ such that $Te_{\alpha} = \theta_{\alpha} e_{\pi(\alpha)}$ for all $\alpha \in \kappa$.

Proof. Given an isometry $T : X_{\mathcal{F}} \rightarrow X_{\mathcal{G}}$, the adjoint operator $T^* : X_{\mathcal{G}}^* \rightarrow X_{\mathcal{F}}^*$ is weak*-weak* continuous and is also an isometry, so that T^* takes the extreme points of $B_{X_{\mathcal{G}}^*}$ to the extreme points of $B_{X_{\mathcal{F}}^*}$. By Proposition 6, for each $\alpha \in \kappa$, there is $s_{\alpha} \in \mathcal{F}$ and a sequence $(\theta_{\xi}^{\alpha})_{\xi \in s_{\alpha}}$ such that $|\theta_{\xi}^{\alpha}| = 1$ and $T^*(e_{\alpha}^*) = \sum_{\xi \in s_{\alpha}} \theta_{\xi}^{\alpha} e_{\xi}^*$. The conclusion now follows similarly as Theorem 10 of [4]. \square

Let us now compare the results of this section to those of [4]. Notice that both in the previous Theorem 7 and Proposition 6, we have a weaker assumption on the families \mathcal{F} and \mathcal{G} : they suppose them to be regular families and we replace the spreading condition by the strictly weaker topological condition that the $\overline{\mathcal{F}^{MAX}}$ and $\overline{\mathcal{G}^{MAX}}$ contain the singletons. On the other hand, we start with a slightly stronger assumption on the operator: in case of Proposition 6 we ask $T : X_{\mathcal{F}}^* \rightarrow X_{\mathcal{G}}^*$ to be weak*-weak* continuous and to preserve extreme points, while in [4] they require only the latter. As a consequence, we start in Theorem 7 from an isometry $T : X_{\mathcal{F}} \rightarrow X_{\mathcal{G}}$, while they start from an isometry $T : X_{\mathcal{F}}^* \rightarrow X_{\mathcal{G}}^*$.

Given two compact and hereditary families \mathcal{F} and \mathcal{G} such that $[\kappa]^1 \subseteq \overline{\mathcal{F}^{MAX}} \cap \overline{\mathcal{G}^{MAX}}$, we do not know the answer to the following question:

Question 8. *Is every isometry $T : X_{\mathcal{F}}^* \rightarrow X_{\mathcal{G}}^*$ weak*-weak* continuous? Or, equivalently, is every isometry $T : X_{\mathcal{F}}^* \rightarrow X_{\mathcal{G}}^*$ the adjoint operator of an isometry $S : X_{\mathcal{G}} \rightarrow X_{\mathcal{F}}$?*

If the answer to these questions is positive, we could get a full version of Corollary 12 of [4] with the weaker assumption on the families. Nevertheless, given a π -homeomorphism between two hereditary families \mathcal{F} and \mathcal{G} induced by a permutation π of κ , the unique linear continuous operator taking e_{α} to $e_{\pi(\alpha)}$ is clearly an isometry between $X_{\mathcal{F}}$ and $X_{\mathcal{G}}$. Hence, we have the following corollary from our previous results:

Corollary 9. *Let \mathcal{F} and \mathcal{G} be compact and hereditary families on κ such that $[\kappa]^1 \subseteq \overline{\mathcal{F}^{MAX}} \cap \overline{\mathcal{G}^{MAX}}$. Then TFAE:*

- (ii) \mathcal{F} and \mathcal{G} are π -homeomorphic.
- (iii) $X_{\mathcal{F}}$ and $X_{\mathcal{G}}$ are isometric.

3. UNIQUENESS OF REGULAR FAMILIES UNDER PERMUTATIONS

In the previous section we improved the results of [4] by getting weaker assumptions on families which are sufficient to guarantee that the combinatorial spaces are isometric iff the families are π -homeomorphic. In this section we deal with families on the countable infinite cardinal ω and improve their results in another direction. We show that two π -homeomorphic regular families are actually the same. Let us consider the following definitions:

Definition 10. *Given $s, t \in [\omega]^{<\omega}$, we say that t is a spread of s if there is an injective mapping $\sigma : s \rightarrow \omega$ such that $\sigma(i) \geq i$ for every $i \in s$, and $\sigma[s] = t$; we will abbreviate this fact by saying that $\sigma[s]$ is a spread of s .*

We call a family \mathcal{F} spreading if $\sigma[t] \in \mathcal{F}$ for every $t \in \mathcal{F}$ and every spread $\sigma[t]$ of t ; and we say that a family \mathcal{F} is regular if it is compact, hereditary and spreading.

The main purpose of this section is to show Theorem 13, which states that hereditary and spreading families on ω are invariable under permutations. In particular, different regular families on ω cannot be permuted one to the other. As a consequence of Theorem 10 of [4], we get that the combinatorial spaces of two different regular families are not isometric.

We start with the following two simple lemmas, which will be used in the proof.

Lemma 11. *Given $s \in [\omega]^{<\omega}$ and a spread $\sigma[s]$ of s , there is a spread $\sigma'[s]$ of s such that $\sigma'[s] = \sigma[s]$ and $\sigma'(i) = i$ for every $i \in s \cap \sigma[s]$.*

Proof. Let $s \in [\omega]^{<\omega}$ and let $\sigma[s]$ be a spread of s . Define $\Delta = s \cap \sigma[s]$ and notice that, since $|s| = |\sigma[s]|$, we have that $|s \setminus \Delta| = |\sigma[s] \setminus \Delta|$. Let $s \setminus \Delta = \{m_1, \dots, m_k\}_{<}$ and $\sigma[s] \setminus \Delta = \{n_1, \dots, n_k\}_{<}$.

Now notice that if there was $1 \leq j < k$ such that $m_j < n_j < n_{j+1} < m_{j+1}$, then $\sigma[s]$ would not be a spread of s . Hence, $m_j < n_j$ for every $1 \leq j \leq k$.

Define $\sigma' : s \rightarrow \omega$ by $\sigma'(i) = i$ if $i \in \Delta$ and $\sigma'(m_j) = n_j$ if $1 \leq j \leq k$. Notice that $\sigma'[s] = \Delta \cup \{n_1, \dots, n_k\} = \sigma[s]$, as wanted. \square

Lemma 12. *If $\mathcal{F} \subseteq [\omega]^{<\omega}$ is a spreading family, then so is $\overline{\mathcal{F}}^\subseteq$.*

Proof. Let $\mathcal{F} \subseteq [\omega]^{<\omega}$ be a spreading family, $s \in \overline{\mathcal{F}}^\subseteq$ and $\sigma[s]$ a spread of s . We must show that $\sigma[s] \in \overline{\mathcal{F}}^\subseteq$.

Let $t \in \mathcal{F}$ be such that $s \subseteq t$ and let $t \setminus s = \{m_1, m_2, \dots, m_k\}_{<}$. Then, consider integers $n_k > \dots > n_2 > n_1 > \max(t \cup \sigma[s])$ and define $\sigma' : t \rightarrow \omega$ by $\sigma'(j) = \sigma(j)$ if $j \in s$; and $\sigma'(m_i) = n_i$ for $1 \leq i \leq k$. It follows that $\sigma'[t]$ is a spread of t and that $\sigma[s] = \sigma'[s] \subseteq \sigma'[t]$. By being \mathcal{F} spreading, we get that $\sigma'[t] \in \mathcal{F}$ and, therefore, $\sigma[s] \in \overline{\mathcal{F}}^\subseteq$. \square

We are now ready to prove our main result:

Theorem 13. *If \mathcal{F} and \mathcal{G} are hereditary, spreading and π -homeomorphic families¹ on ω , then $\mathcal{F} = \mathcal{G}$.*

Proof. We prove, by induction on n , that for any two hereditary and spreading families \mathcal{F} and \mathcal{G} , if there is a permutation $\pi : \omega \rightarrow \omega$ such that $\mathcal{G} = \pi[\mathcal{F}]$, then $\mathcal{F} \cap [\omega]^{\leq n} = \mathcal{G} \cap [\omega]^{\leq n}$.

For $n = 1$, let

$$I = \{i \in \omega : \{i\} \notin \mathcal{F}\} \text{ and } J = \{i \in \omega : \{i\} \notin \mathcal{G}\}.$$

Notice that, from spreading, I and J are initial segments of ω . Moreover, $i \in I$ iff $\pi(i) \in J$, so that $|I| = |J|$. It follows that $I = J$ and, therefore, $\mathcal{F} \cap [\omega]^1 = \{\{i\} : i \notin I\} = \{\{i\} : i \notin J\} = \mathcal{G} \cap [\omega]^1$.

Assume that the statement holds for n and let us prove it for $n + 1$. Given $s \in \mathcal{F} \cap [\omega]^n$, let

$$I_s = \{i \in \omega : \{i\} \cup s \notin \mathcal{F}\}.$$

¹In this result we do not actually need to assume that \mathcal{F} and \mathcal{G} contain all singletons. If they contain, the proof for $n = 1$ in the induction would be simpler, but we opted to present a proof which makes sense both under this assumption or without it.

Let us prove some properties of I_s . It is easy to see, from spreading, that I_s is an initial segment of $\omega \setminus s$. Moreover, we have the following:

Claim. *If $s \in \mathcal{F} \cap [\omega]^n$ and $t \in [\omega]^n$ is a spread of s (in particular, $t \in \mathcal{F}$), then $|I_t| \leq |I_s|$.*

Proof of the claim. First notice that

$$I_t \setminus s = I_t \setminus (s \cup t) \subseteq I_s \setminus (s \cup t) = I_s \setminus t,$$

where the two equalities follow from the fact that $I_s \cap s = I_t \cap t = \emptyset$ and the inclusion follows from spreading of \mathcal{F} .

Second, let us show that $|I_t \cap s| \leq |I_s \cap t|$. Indeed, by Lemma 11, consider $\sigma : s \rightarrow t$ a bijection such that $\sigma(i) \geq i$ for every $i \in s$ and, additionally, $\sigma(i) = i$ for every $i \in s \cap t$. We show that if $j \in I_t \cap s$, then $\sigma(j) \in I_s \cap t$. Indeed, fix $j \in I_t \cap s$ and let $\sigma' : \{\sigma(j)\} \cup s \rightarrow \{j\} \cup t$ be defined by $\sigma'(i) = \sigma(i)$ for $i \in s \setminus \{j, \sigma(j)\}$, $\sigma'(j) = j$ and $\sigma'(\sigma(j)) = \sigma(j)$. σ' is well-defined (since $\sigma(i) = i$ for $i \in s \cap t$) and it witnesses that $\{j\} \cup t$ is a spread of $\{\sigma(j)\} \cup s$.

Since $j \in I_t$, then $\sigma'[\{\sigma(j)\} \cup s] = \{j\} \cup t \notin \mathcal{F}$. Also, \mathcal{F} is spreading, so that $\{\sigma(j)\} \cup s \notin \mathcal{F}$, that is, $\sigma(j) \in I_s$. Hence, $\sigma(j) \in I_s \cap t$ and we conclude that $|I_t \cap s| \leq |I_s \cap t|$. Summing up we get that $|I_t| \leq |I_s|$ and this concludes the proof of the claim. \square

Now, for each $k \in \omega$, let

$$\mathcal{F}_{n,k} = \{s \in \mathcal{F} \cap [\omega]^n : |I_s| \leq k\}.$$

It follows from the claim that $\mathcal{F}_{n,k}$ is a spreading family, for each $k \in \omega$.

For each $t \in \mathcal{G} \cap [\omega]^n$ and each $k \in \omega$, define

$$J_t = \{i \in \omega : \{i\} \cup t \notin \mathcal{G}\} \text{ and } \mathcal{G}_{n,k} = \{t \in \mathcal{G} \cap [\omega]^n : |J_t| \leq k\},$$

and get similar properties as for I_s and $\mathcal{F}_{n,k}$. In particular, $\mathcal{G}_{n,k}$ is a spreading family for every $k \in \omega$.

Since both $\mathcal{F}_{n,k}$ and $\mathcal{G}_{n,k}$ are spreading and $\mathcal{G}_{n,k} = \pi[\mathcal{F}_{n,k}]$, then by Lemma 12, the downward closures $\overline{\mathcal{F}}_{n,k}^\subseteq$ and $\overline{\mathcal{G}}_{n,k}^\subseteq$ are hereditary and spreading families, and $\overline{\mathcal{G}}_{n,k}^\subseteq = \pi[\overline{\mathcal{F}}_{n,k}^\subseteq]$. It follows from the inductive hypothesis that $\mathcal{G}_{n,k} = \overline{\mathcal{G}}_{n,k}^\subseteq \cap [\omega]^n = \overline{\mathcal{F}}_{n,k}^\subseteq \cap [\omega]^n = \mathcal{F}_{n,k}$, for every $k \in \omega$.

Finally,

$$\begin{aligned} \mathcal{F} \cap [\omega]^{n+1} &= \{\{i\} \cup s : i \notin s \text{ and } \exists k \in \omega, s \in \mathcal{F}_{n,k} \setminus \mathcal{F}_{n,k-1} \text{ and } i \notin \tau_k(\omega \setminus s)\} \\ &= \{\{i\} \cup s : i \notin s \text{ and } \exists k \in \omega, s \in \mathcal{G}_{n,k} \setminus \mathcal{G}_{n,k-1} \text{ and } i \notin \tau_k(\omega \setminus s)\} = \mathcal{G} \cap [\omega]^{n+1}, \end{aligned}$$

where $\tau_k : \wp(\omega) \rightarrow [\omega]^k$ associates to any set X its initial segment $\tau_k(X)$ of size k . \square

The following corollary follows immediately from [4, Corollary 12] and Theorem 13 above, our crucial contribution being the equivalence between (i) and (ii).

Corollary 14. *If \mathcal{F} and \mathcal{G} are regular families on ω , then TFAE:*

(i) $\mathcal{F} = \mathcal{G}$.

- (ii) \mathcal{F} and \mathcal{G} are π -homeomorphic
- (iii) $X_{\mathcal{F}}$ and $X_{\mathcal{G}}$ are isometric.
- (iv) $X_{\mathcal{F}}^*$ and $X_{\mathcal{G}}^*$ are isometric.

4. EXAMPLES AND FURTHER REMARKS

The purpose of this section is to give some examples and prove several facts which put our main results into a bigger picture. In particular, a topological approach is useful.

For most of the results in this section we will use the Cantor-Bendixson rank. We recall that if X is a topological space, then $X' = \{x \in X : x \text{ is not isolated in } X\}$ and given an ordinal α , the α -th Cantor-Bendixson derivative of X is defined recursively by:

$$\begin{aligned} X^{(0)} &= X \\ X^{(\beta+1)} &= (X^{(\beta)})' \\ X^{(\lambda)} &= \bigcap_{\beta < \lambda} X^{(\beta)}, \text{ if } \lambda > 0 \text{ is a limit ordinal.} \end{aligned}$$

The Cantor-Bendixson rank of X is the minimal ordinal such that $X^{(\alpha)} = X^{(\alpha+1)}$ and it is denoted by $\text{rk}(X)$.

If $\mathcal{F} \subseteq [\kappa]^{<\omega}$ is a compact family, then its Cantor-Bendixson index is the smallest ordinal α such that $\mathcal{F}^{(\alpha)} = \emptyset$ and it is therefore a successor ordinal. Moreover, $\text{rk}(\mathcal{F}) < \kappa^+$.

4.1. Injective Cantor-Bendixson index. Given a compact family $\mathcal{F} \subseteq [\kappa]^{<\omega}$, we can define a function $\text{rk}_{\mathcal{F}} : \mathcal{F} \rightarrow \kappa^+$ which associates to $s \in \mathcal{F}$ the biggest ordinal β for which $s \in \mathcal{F}^{(\beta)}$. Notice that the rank of a point coincides with the rank of its image under a homeomorphism between two topological spaces. So, in particular, the rank of a singleton coincides with the rank of its image under a π -homeomorphism between two families.

When a compact family \mathcal{F} on ω is spreading, the corresponding function $\text{rk}_{\mathcal{F}} : [\omega]^1 \rightarrow \omega_1$ is nondecreasing. In particular, for the Schreier family

$$\mathcal{S} := \{s \in [\omega]^{<\omega} : |s| \leq \min(s) + 1\} \cup \{\emptyset\}$$

(known to be regular) $\text{rk}_{\mathcal{S}}(\{n\}) = n$ for every $n \in \omega$, so that $\text{rk}_{\mathcal{S}} \upharpoonright [\omega]^1$ is injective. This imposes restrictions on which permutations induce a homeomorphism of a given family:

Remark 15. *If \mathcal{F} is a compact family on κ such that $\text{rk}_{\mathcal{F}} \upharpoonright [\kappa]^1$ is injective, then the only permutation π of κ such that $\pi[\mathcal{F}] = \mathcal{F}$ is the identity map.*

It follows from this remark and Theorem 7 that if, moreover, $[\kappa]^1 \subseteq \overline{\mathcal{F}^{MAX}}$, then every isometry of $X_{\mathcal{F}}$ is given by a change of signs of the elements of its basis. This generalizes [4, Theorem 15].

Also, for families which have strictly increasing rank function, we can easily get the conclusion of Theorem 13:

Proposition 16. *If \mathcal{F} and \mathcal{G} are π -homeomorphic compact families on κ such that $\text{rk}_{\mathcal{F}}$ is strictly increasing and $\text{rk}_{\mathcal{G}}$ is nondecreasing, then the only permutation of κ inducing an homeomorphism between them is the identity map. In particular, $\mathcal{F} = \mathcal{G}$.*

Proof. Let \mathcal{F} and \mathcal{G} be as in the hypothesis, and let π be a permutation of κ such that $\mathcal{G} = \pi[\mathcal{F}]$. We will prove that π is increasing and since it is a permutation, it has to be the identity map. Assume towards a contradiction that there are $\alpha < \beta < \kappa$ such that $\pi(\alpha) \geq \pi(\beta)$. Then

$$\text{rk}_{\mathcal{F}}(\{\alpha\}) = \text{rk}_{\mathcal{G}}(\{\pi(\alpha)\}) \geq \text{rk}_{\mathcal{G}}(\{\pi(\beta)\}) = \text{rk}_{\mathcal{F}}(\{\beta\}),$$

contradicting the fact that $\text{rk}_{\mathcal{F}}$ is strictly increasing. \square

As an application, we get an example of distinct compact and hereditary families which are π -homeomorphic. It guarantees that Theorem 13 cannot be improved by dropping the spreading condition:

Example 17. *Let $\pi \neq \text{Id}$ be any permutation of ω and $\mathcal{F} = \pi[\mathcal{S}]$. From Remark 15 we get that $\mathcal{F} \neq \mathcal{S}$. Since \mathcal{F} is π -homeomorphic to \mathcal{S} , \mathcal{F} is also compact and hereditary and it follows from Corollary 9 that $X_{\mathcal{S}}$ and $X_{\mathcal{F}}$ are isometric.*

4.2. More on spreading families. In the context of compact spreading families on ω , the hypotheses of Proposition 16 are almost complete, the only missing one being that the rank of one of the families has to be not only nondecreasing, but strictly increasing. The next example shows that Proposition 16 does not necessarily hold when $\text{rk}_{\mathcal{F}}$ is not strictly increasing. It also shows that two families may be π -homeomorphic with only one of them being spreading.

Example 18. *Consider $\mathcal{F} = [\omega]^{\leq 2} \setminus \{\{2, 3\}\}$ and $\mathcal{G} = [\omega]^{\leq 2} \setminus \{\{1, 2\}\}$. It is clear that \mathcal{F} is compact, that $\text{rk}_{\mathcal{F}}(\{n\}) = 1$ for every $n \in \omega$, and that \mathcal{G} is spreading. The permutation $\pi : \omega \rightarrow \omega$ given by $\pi(1) = 3$, $\pi(2) = 1$, $\pi(3) = 2$, and $\pi(n) = n$ for every $n > 3$ witnesses that $\mathcal{G} = \pi[\mathcal{F}]$. Nevertheless, $\mathcal{F} \neq \mathcal{G}$.*

It is well known that for every limit ordinal $\alpha < \omega_1$ there is a compact family \mathcal{F} on ω with rank $\alpha + 1$ such that $\text{rk}_{\mathcal{F}}$ is strictly increasing; for example, the Schreier families (see [3]) or the closure of an α -uniform front (introduced in [10]). We should also mention the existence of a non hereditary and non spreading compact family \mathcal{F} on ω with strictly increasing rank: let for example

$$\mathcal{F} = \{s \in \mathcal{S} : \{n, n+1\} \text{ is not an initial segment of } s, \text{ for every } n \in \omega\};$$

then \mathcal{F} is not hereditary nor spreading since $\{3, 5, 6\} \in \mathcal{F}$ but $\{5, 6\}, \{4, 5, 6\} \notin \mathcal{F}$. Thus, although we cannot apply Theorem 13 here to conclude that any spreading family π -homeomorphic to \mathcal{F} is \mathcal{F} itself, we can instead get that conclusion from Proposition 16.

As a consequence of Theorem 13, if \mathcal{F} and \mathcal{G} are two different hereditary and spreading families, they are not π -homeomorphic. A natural question to ask is whether \mathcal{F} and \mathcal{G} turn out to be π -homeomorphic if in addition we ask them to be homeomorphic. The following example answers negatively this question.

Example 19. Let $\mathcal{F} = [\omega]^{\leq 1} \cup [\omega \setminus \{1\}]^2$ and $\mathcal{G} = [\omega]^{\leq 2}$. Consider the mapping $\varphi : \mathcal{F} \rightarrow \mathcal{G}$ given by $\varphi(\emptyset) = \emptyset$, $\varphi(\{1\}) = \{1, 2\}$, $\varphi(\{n\}) = \{n-1\}$ if $n > 1$, $\varphi(\{2, m\}) = \{1, m\}$ for $m > 2$, and $\varphi(\{n, m\}_{<}) = \{n-1, m-1\}$ for every $n > 2$. It is clear that φ is a homeomorphism. Nevertheless, there is no permutation $\pi : \omega \rightarrow \omega$ such that $\mathcal{G} = \{\pi[s] : s \in \mathcal{F}\}$. Indeed, if such a permutation π exists, then $\pi(\{1\})$ should be an isolated point in \mathcal{G} with size 1, which is impossible. Hence, \mathcal{F} and \mathcal{G} are homeomorphic regular families which are not π -homeomorphic. In particular, $X_{\mathcal{F}}$ and $X_{\mathcal{G}}$ are not isometric.

Since each regular family determines a class of compact and hereditary families on ω which have isometric combinatorial spaces, it would be nice if every compact and hereditary family on ω would be π -homeomorphic to a regular one, so that regular families would classify all isometric-types of separable combinatorial spaces. However, this is not the case, as shows the following example.

Example 20. Consider $\mathcal{F} = [\mathbb{N}]^{\leq 2} \setminus \{\{n, n+1\} : n \in \mathbb{N}\}$ which is clearly compact, hereditary and not spreading. We claim that \mathcal{F} is not π -homeomorphic to any regular family. Indeed, assume towards a contradiction that there is a regular family \mathcal{G} and a permutation of ω such that $\mathcal{G} = \pi[\mathcal{F}]$. Then, $\{\pi(n), \pi(n+1)\} \notin \mathcal{G}$ for every $n < \omega$.

We will now see that $\pi(n)$ and $\pi(n+1)$ is one the successor of the other, for every $n < \omega$. Indeed, if there are $n, k < \omega$ such that $\pi(n) < k < \pi(n+1)$, let $p < \omega$ be such that $\pi(p) = k$. Since $p \neq n, n+1$, then $\{n, p\} \in \mathcal{F}$ and $\{\pi(n), k\} \in \mathcal{G}$. By being \mathcal{G} spreading, we have that $\{\pi(n), \pi(n+1)\} \in \mathcal{G}$, which is a contradiction. Analogously, we see that there are not $n, k < \omega$ such that $\pi(n+1) < k < \pi(n)$.

Let $m < \omega$ be such that $\pi(m) = 1$. Then, $\pi(m+1) = 2$ as it is the successor of $\pi(m)$. It follows that $\pi[\{m, m+1, m+2, \dots\}] = \omega$. Thus, π is the identity map, contradicting that \mathcal{F} is not spreading.

4.3. Examples in the uncountable setting. Our purpose now is to explore some examples of families on a given uncountable cardinal κ which satisfy the hypotheses of Theorem 7. It is clear that $\mathcal{F} = [\kappa]^{\leq n}$ is such a family, but we consider this to be somewhat trivial, as the norm looks very much like the supremum norm and any permutation of κ fixes \mathcal{F} .

On the other hand, if we assume \mathcal{F} on some uncountable cardinal κ not to be contained in any $[\kappa]^{\leq n}$, then it cannot be spreading: let $(s_n)_n \subseteq \mathcal{F}$ with $|s_n| \geq n$ and fix $\alpha \in \kappa$ such $s_n \subseteq \alpha$ for every $n \in \omega$. Spreading and hereditary would imply that the intervals $[\alpha, \alpha + n] \in \mathcal{F}$. In turn, this would contradict compactness, since this sequence converges to $[\alpha, \alpha + \omega)$, which is infinite and does not belong to \mathcal{F} .

Example 21. Given a cardinal κ , let $\mathcal{S}(\kappa)$ be the family formed by copies of the Schreier family \mathcal{S} in each interval of the form $[\lambda, \lambda + \omega)$, where $\lambda < \kappa$ is a limit ordinal (or zero). Formally,

$$\mathcal{S}(\kappa) = \{s \in [\kappa]^{<\omega} : s \subseteq [\lambda, \lambda + \omega) \text{ for some limit ordinal } \lambda < \kappa \text{ and } |s| \leq \min\{n \in \omega : \lambda + n \in s\} + 1\}.$$

Notice that $\mathcal{S}(\kappa)$ is a compact hereditary family such that $\{\alpha\} \in \overline{\mathcal{S}(\kappa)}^{MAX}$ for each $\alpha \in \kappa$, so that it satisfies the hypothesis of Theorem 7. Moreover, $\mathcal{S}(\kappa)$ contains arbitrarily large (finite) elements. The corresponding combinatorial space $X_{\mathcal{S}(\kappa)}$ is isometric to an ℓ_∞ -sum of κ many copies of $X_{\mathcal{S}}$.

To get an example which is a bit more involving we use trees. A tree is a partially ordered set (T, \leq) such that for every $f \in T$, the set $\{g \in T : g \leq f\}$ is well-ordered. Hence, we can define the height $ht_T(f)$ of f in T as the only ordinal which is order-isomorphic to $\{g \in T : g \leq f\}$. The height $ht(T)$ of T is the supremum of all $ht_T(f)$, $f \in T$. A chain in T is a chain with respect to \leq , that is, a subset $C \subseteq T$ such that for every $f, g \in C$, either $f \leq g$ or $g \leq f$. A branch in T is a maximal chain in T .

Finally, we have the following proposition, which is implicit in [5, Lemma 2.32]:

Proposition 22. Let κ be an infinite cardinal and let T be a tree of height κ . If there is a compact and hereditary family \mathcal{F} on κ such that $[\kappa]^1 \subseteq \overline{\mathcal{F}}^{MAX}$, then there is such a family on $\lambda = |T|$.

Proof. First notice that the existence of a compact and hereditary family on some cardinal κ is equivalent to the existence of a compact and hereditary family on some index set I of cardinality κ , since being compact and hereditary doesn't depend on the order.

Hence, without loss of generality, we may assume that \mathcal{F} is a compact and hereditary family on $\kappa + 1$ (instead of κ) such that $[\kappa + 1]^1 \subseteq \overline{\mathcal{F}}^{MAX}$. We define

$$\mathcal{G} = \{C \subseteq T : C \text{ is a chain and } \{ht_T(f) : f \in C\} \in \mathcal{F}\}.$$

It is easy to see that \mathcal{G} is a compact and hereditary family on T . Given $f \in T$, we fix a branch C which contains f and, given $(s_n)_n \subseteq \mathcal{F}^{MAX}$ converging to $\{ht_T(f)\}$, it is easy to build $(t_n)_n$ converging to $\{f\}$, where each $t_n \subseteq C$ and $\{ht_T(g) : g \in t_n\} = s_n$. Clearly each $t_n \in \mathcal{G}^{MAX}$, which concludes the proof that $[T]^1 \subseteq \overline{\mathcal{G}}^{MAX}$. From the first paragraph of the proof, it follows that the desired family on $\lambda = |T|$ exists. \square

Example 21 can be seen as a particular case of a family given by Proposition 22, taking T to be κ -many incomparable copies of ω . Also, if we have a family on κ , we get a family on 2^κ , by taking T the complete binary tree of height κ . This way we get more interesting families for every cardinal below the first inaccessible cardinal.

Motivated by Remark 15 and Proposition 16, we asked ourselves if there is a compact hereditary family \mathcal{F} on ω_1 such that $rk(\{\alpha\}) = \alpha$ for every $\alpha < \omega_1$. The following example answers this question positively:

Example 23. Given $\alpha < \omega_1$, consider \mathcal{F}_α a compact and hereditary family on ω with $\text{rk}(\mathcal{F}_\alpha) = \alpha + 1$ (the existence of these families is guaranteed, for example, in [3]).

Given a limit ordinal $\lambda < \omega_1$, and $s \in [\omega_1]^{<\omega}$ such that $s \subseteq [\lambda, \lambda + \omega)$, we will denote by $s - \lambda$ the set $\{\gamma - \lambda : \gamma \in s\}$ which belongs to $[\omega]^{<\omega}$. In a similar way as we have defined $\mathcal{S}(\kappa)$ in Example 21, we now define the families $\mathcal{F}_\alpha(\kappa)$, for κ a tail of ω_1 and $\alpha < \omega_1$, by

$$\mathcal{F}_\alpha(\kappa) = \{s \in [\kappa]^{<\omega} : \text{there is } \lambda < \kappa \text{ a limit ordinal or zero such that } s \subseteq [\lambda, \lambda + \omega) \text{ and } s - \lambda \in \mathcal{F}_\alpha\}.$$

Notice that $\mathcal{F}_\alpha(\kappa)$ is a compact and hereditary family on κ with $\text{rk}(\mathcal{F}_\alpha(\kappa)) = \alpha + 1$. Consider

$$\mathcal{F} := \mathcal{S} \cup \bigcup_{\omega \leq \alpha < \omega_1} \{\{\alpha\} \cup s : s \in \mathcal{F}_\alpha(\omega_1 \setminus \alpha), \text{ and } \alpha < \gamma \text{ for every } \gamma \in s\}.$$

It is clear that \mathcal{F} is a compact and hereditary family on ω_1 . Moreover, notice that $\text{rk}_{\mathcal{F}}(\{\alpha\}) = \text{rk}_{\mathcal{F}_\alpha(\omega_1 \setminus \alpha)}(\emptyset) = \text{rk}(\mathcal{F}_\alpha(\omega_1 \setminus \alpha)) - 1 = \alpha$, for every $\omega \leq \alpha < \omega_1$.

REFERENCES

- [1] L. Antunes, K. Beanland, and H. Viet Chu, *On the geometry of higher order Schreier spaces*, preprint.
- [2] R. F. Arens, J. L. Kelley, *Characterization of the space of continuous functions over a compact Hausdorff space*, Trans. Amer. Math. Soc. 62 (1947), 499–508.
- [3] S. A. Argyros, S. Todorćević, *Ramsey methods in analysis*, Advanced Courses in Mathematics. CRM Barcelona. Birkhäuser Verlag, Basel, 2005.
- [4] C. Brech, V. Ferenczi and A. Tcaciuc, *Isometries of combinatorial Banach spaces*, accepted to Proc. Amer. Math. Soc.
- [5] C. Brech, J. Lopez-Abad, S. Todorćević, *Homogeneous families on trees and subsymmetric basic sequences*, Adv. Math. 334 (2018), 322–388.
- [6] W. Gowers, *Must an “explicitly defined” Banach space contain c_0 or l_p ?*, Feb. 17, 2009, Gowers’s Weblog: Mathematics related discussions.
- [7] E. V. Kadison, *Isometries of Operator Algebras*, Annals of Mathematics Second Series, Vol. 54, No. 2 (Sep., 1951), pp. 325–338.
- [8] J. Lindenstrauss, L. Tzafriri, *Classical Banach spaces. I. Sequence spaces*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Vol. 92. Springer-Verlag, Berlin-New York, 1977.
- [9] J. Lopez-Abad, S. Todorćević, *Positional graphs and conditional structure of weakly null sequences*, Adv. Math. 242 (2013), 163–186.
- [10] P. Pudlák, V. Rödl, *Partition theorems for systems of finite subsets of integers*, Discrete Math. 39 (1) (1982) 67–73.
- [11] W. Rudin, *Functional analysis*, International Series in Pure and Applied Mathematics, McGraw-Hill, Inc., New York, 1991.
- [12] Z. Semadeni, *Banach spaces of continuous functions*, Monografie Matematyczne, Tom 55, Państwowe Wydawnictwo Naukowe, Warsaw, 1971.

DEPARTAMENTO DE MATEMÁTICA, INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE DE SÃO PAULO, RUA DO MATÃO, 1010 - CEP 05508-090 - SÃO PAULO - SP - BRAZIL.

E-mail address: `brech@ime.usp.br`

DEPARTAMENTO DE MATEMÁTICAS, FACULTAD DE CIENCIAS, UNIVERSIDAD DE LOS ANDES, C.P. 5101, MÉRIDA-VENEZUELA.

ESCUELA DE MATEMÁTICAS, UNIVERSIDAD INDUSTRIAL DE SANTANDER, C.P. 680001, BUCARAMANGA - COLOMBIA.

E-mail address: `cpinarangel@gmail.com`