

Adding quadric fillets to quador lattice structures

Fehmi Cirak*, Malcolm Sabin

Department of Engineering, University of Cambridge, Trumpington Street, Cambridge CB2 1PZ, UK

Abstract

Gupta et al. [1, 2] describe a very beautiful application of algebraic geometry to lattice structures composed of quadric of revolution (quador) implicit surfaces. However, the shapes created have concave edges where the stubs meet, and such edges can be stress-raisers which can cause significant problems with, for instance, fatigue under cyclic loading. This note describes a way in which quadric fillets can be added to these models, thus relieving this problem while retaining their computational simplicity and efficiency.

Keywords: lattice structures, quadrics of revolution, quadors, algebraic geometry

Throughout, we use an upper case letter to denote both a surface (e.g., a sphere S with the centre $(x_c, y_c, z_c)^T$ and radius r) and the implicit function (i.e., $S(x, y, z) = (x - x_c)^2 + (y - y_c)^2 + (z - z_c)^2 - r^2$) which is zero on that surface. No confusion should arise. Let S denote the quadratic function with the zero set on the central sphere of a quador hub, increasing outward from the centre.




Figure 1: A hub (lattice joint) with two attached stubs (beams leading to other joints). This figure is diagrammatic only. Its purpose is to identify what surfaces the various letters denote. The arrow on a surface indicates the gradient vector of the expression whose zero set is the surface.

Let H_1 and H_2 denote the quadratic functions of two quadors tangent to the central sphere S . Then, these two functions have the form

$$H_1 = S - G_1^2 \quad \text{and} \quad H_2 = S - G_2^2$$

for linear functions G_1 and G_2 which are zero on the respective planes of tangency, so that $\nabla H_1 = \nabla S$ and $\nabla H_2 = \nabla S$.

We can construct planes, for constants $\alpha > 0$ and $\beta > 0$,

$$E_1 = \alpha F_+ + \beta F_- \quad \text{and} \quad E_2 = \alpha F_+ - \beta F_-$$

with the functions $F_- = G_2 - G_1$ and $F_+ = G_2 + G_1$.

Consider the quadrics whose equations are $H_1 - E_1^2 = 0$ (which is tangential to H_1 along the curve of intersection with E_1) and $H_2 - E_2^2 = 0$ (which is tangential to H_2 along the curve of intersection with E_2). These two will be the same quadric, providing a fillet between H_1 and H_2 if

$$\begin{aligned} H_1 - E_1^2 &= H_2 - E_2^2 \\ \text{or} \quad H_1 - H_2 &= E_1^2 - E_2^2 \\ \text{but} \quad H_1 - H_2 &= S - G_1^2 - S + G_2^2 = G_2^2 - G_1^2 \\ &= (G_2 + G_1)(G_2 - G_1) \\ &= F_+ F_- \\ \text{and} \quad E_1^2 - E_2^2 &= (E_1 + E_2)(E_1 - E_2) \\ &= (2\alpha F_+)(2\beta F_-) \\ &= 4\alpha\beta F_+ F_- \end{aligned}$$

so we get a single fillet quadric if $\alpha\beta = 1/4$. We can choose either α or β and then the other is fixed. The ratio between the two controls the angles of the planes E_1 and E_2 either side of F_- . Increasing β slowly from zero increases the size of the fillet and its smallest radius of curvature, but this increases the length of the stub.

The above shows that there exists a fan of possible quadrics providing a tangent continuous join between adjacent stubs. Each piece of surface has an exact implicit form, and an exact parametric form. The curves of tangency are all conics with exact parametric curves, exact implicit curves within their planes, and exact p-curves (i.e., trimming curves in parameter space) within both surfaces. Surfaces can be separated by the planes which contain the curves of tangency, and so all the important properties in [1] and [2] still apply.

References

- [1] A. Gupta, G. Allen, J. Rossignac, *QUADOR: QUADric-Of-Revolution beams for lattices* CAD 102, 160–170 (2018)
- [2] A. Gupta, G. Allen, J. Rossignac, *Exact Representations and Geometric Queries for Lattice Structures with Quador Beams* CAD 115, 64–77 (2019)

*Corresponding author