

KEK-TH-1203, FERMILAB-PUB-07-651-T
 CERN-PH-TH/2007-258, MAN/HEP/2007/43
 ANL-HEP-PR-07-105, EFI-07-38
 arXiv:0712.2360
 December 2007

CP superH 2.0 :

an Improved Computational Tool for Higgs Phenomenology
 in the MSSM with Explicit CP Violation

J. S. Lee^{a,b}, M. Carena^c, J. Ellis^d, A. Pilaftsis^e and C. E. M. Wagner^{f,g}

^aTheory Group, KEK, Oho 1-1 Tsukuba, 305-0801, Japan

^bDepartment of Physics, National Central University, Chung-Li, Taiwan 32054

^cFermilab, P.O. Box 500, Batavia IL 60510, U.S.A.

^dTheory Division, CERN, CH-1211 Geneva 23, Switzerland

^eSchool of Physics and Astronomy, University of Manchester,
 Manchester M13 9PL, United Kingdom

^fHEP Division, Argonne National Laboratory, 9700 Cass Ave., Argonne, IL 60439, USA

^gEnrico Fermi Institute, Univ. of Chicago, 5640 Ellis Ave., Chicago, IL 60637, USA

ABSTRACT

We describe the Fortran code CPsuperH2.0, which contains several improvements and extensions of its predecessor CPsuperH. It implements improved calculations of the Higgs-boson pole masses, notably a full treatment of the 4×4 neutral Higgs propagator matrix including the Goldstone boson and a more complete treatment of threshold effects in self-energies and Yukawa couplings, improved treatments of two-body Higgs decays, some important three-body decays, and two-loop Higgs-mediated contributions to electric dipole moments. CPsuperH2.0 also implements an integrated treatment of several B-meson observables, including the branching ratios of $B_s \rightarrow \mu^+ \mu^-$, $B_d \rightarrow \mu^+ \mu^-$, $B_u \rightarrow \mu^+ \mu^-$, $B \rightarrow X_s$ and the latter's CP-violating asymmetry A_{CP} , and the supersymmetric contributions to the $B_{s,d}^0 \rightarrow B_{s,d}^0$ mass differences. These additions make CPsuperH2.0 an attractive integrated tool for analyzing supersymmetric CP and flavour physics as well as searches for new physics at high-energy colliders such as the Tevatron, LHC and linear colliders.

The program may be obtained from <http://www.hep.man.ac.uk/u/jslee/CPsuperH.html>.

1 Introduction

With the imminent advent of the LHC, particle physics experiments are poised to explore the TeV energy range directly for the first time. There are several reasons to expect new physics in this energy range, such as the origin of particle masses and electroweak symmetry breaking, the hierarchy problem and the nature of dark matter. In parallel with the direct exploration of the TeV scale, precision experiments at low energies continue to place important constraints on the possible flavour and CP-violating structure of any TeV-scale physics. Prominent examples include experiments on B and K mesons, and probes of electric dipole moments [1]. It is clearly desirable to develop computational tools that can be used to calculate consistently observables for both low- and high-energy experiments in a coherent numerical framework. This is particularly desirable in view of the possibility that the dominance of matter over antimatter in the Universe may be due to CP-violating interactions at the TeV scale [2].

Supersymmetry is one of the most prominent possibilities for new TeV-scale physics, and the minimal supersymmetric extension of the Standard Model (MSSM) provides a natural cold dark matter candidate as well as stabilizing the electroweak scale and facilitating the unification of the fundamental interactions. There are many computational tools available for calculations within the MSSM. The first to include CP-violating phases was CPsuperH [3] based on the renormalization-group-(RG-)improved effective potential approach. The Higgs-boson pole mass shifts are calculated by employing the RG-improved diagrammatic approach. The recent versions of FeynHiggs [4] are based the Feynman diagrammatic approach. There are merits in both approaches and the difference between two programs may be attributed to some unknown higher-order corrections.

Some of us have recently published an analysis of several B-physics observables taking into account the most general set of CP-violating parameters allowed under the assumption of minimal flavour violation in the supersymmetric sector [5]. For this purpose we used an updated and extended computational tool, CPsuperH2.0, which we introduce and describe in this paper.

The main new features of CPsuperH2.0 are its inclusion of a number of B observables, including the branching ratios of $B_s \rightarrow \mu^+ \mu^-$, $B_d \rightarrow \mu^+ \mu^-$, $B_u \rightarrow \mu^+ \mu^-$, $B \rightarrow X_s \mu^+ \mu^-$ and the latter's CP-violating asymmetry A_{CP} , and the supersymmetric contributions to the $B_{s,d}^0 \rightarrow B_{s,d}^0$ mass differences. In addition, CPsuperH2.0 includes a more complete treatment of Higgs-boson pole masses, based on a full treatment of the 4×4 neutral Higgs propagator matrix including the Goldstone boson and a more complete treatment of threshold effects in self-energies and Yukawa couplings. It also includes improved treatments of two-body Higgs decays, some important three-body decays, and two-loop Higgs-mediated contributions to electric dipole moments. Therefore, CPsuperH2.0 provides an essentially complete, self-contained and consistent computational tool for evaluating flavour and CP-violating physics

at energies up to the TeV scale.

The structure of this paper is as follows. Several updated features of CPsuperH2.0 are described in Section 2. In particular, in Subsection 2.1 we introduce the improved treatment of Higgs-boson pole masses, and Section 2.2 contains a description of the improvements in the treatment of Higgs decay modes. Then, in Section 3 we describe the CPsuperH2.0 treatment of two-loop Higgs effects on electric dipole moments. The most important new features are described in Section 4, where we discuss its treatment of B observables. In each Section, we illustrate in figures some typical results obtained using CPsuperH2.0.

2 Updated Features of CPsuperH2.0

It is to be understood that, throughout this paper, we follow the notations and conventions defined and adopted in CPsuperH for the fixing matrices of neutral Higgs bosons, charginos, neutralinos and third-generation sfermions, as well as their masses and couplings, etc. The updates to the original version of CPsuperH [3] that are presented here reflect, in part, feedback from users, as well as extending it to B observables.

New common blocks /HC_RAUX/ and /HC_CAUX/ have been introduced for the general purpose of storing new numerical outputs which are available in CPsuperH2.0:

```
COMMON /HC_RAUX/ RAUX_H
```

```
COMMON /HC_CAUX/ CAUX_H
```

The two arrays RAUX_H and CAUX_H are NAUX=999 dimensional and only parts of them are being used presently as shown in Tables 1 and 2. The contents of these two new arrays are explained in the corresponding following subsections. These common blocks can also be used by users for their specific purposes.

2.1 Improved Treatment of Higgs-Boson Masses and Propagators

In CPsuperH2.0 we make three main improvements in the calculation of the Higgs-boson pole masses.

The finite threshold corrections induced by the exchanges of gluinos and charginos have been included in the top- and bottom-quark self-energies of the neutral and charged Higgs bosons. For the explicit expressions of the self-energies, we refer to Eqs. (B.14), (B.15), and (B.16) of Ref. [6]^y.

^yWe find that overall minus signs are missing in the expressions of $P_{11;22}^{i(c)}(s)$.




Figure 1: The masses of the neutral Higgs bosons as functions of $\tan\beta$ for the CPX scenario [8] taking $\mu_3 = A_{\tau\beta\tau} = 90$ in the convention $\mu = 0$, $M_{\text{SUSY}} = 0.5 \text{ TeV}$, and the charged Higgs-boson pole mass $M_{H^\pm} = 160 \text{ GeV}$. In each frame, the dashed line is for the case $\text{IFLAG_H}(12)=1$ and the solid line for other case indicated.

Also included are the threshold corrections to the Yukawa couplings $y_{t,j}$ in the one-loop running quartic couplings, $\lambda_i^{(1)}(Q = m_t^{\text{pole}})$ with $i = 1 \dots 4$. For the explicit expressions of $\lambda_i^{(1)}$, we refer to Eqs. (3.3)–(3.6) of Ref. [7].

An improved iterative method has been employed for the calculation of the pole masses.

As a help in assessing the improvements in the calculation of Higgs sector, new flags $\text{IFLAG_H}(12)$ and $\text{IFLAG_H}(60)$ have been introduced as follows:

$\text{IFLAG_H}(12)$:




Figure 2: The masses of the neutral Higgs bosons as functions of the common phase Φ_A for the trimixing scenario [9] taking $\beta_3 = 90^\circ$. Specially, in this scenario, $\tan\beta = 50$ and $M_H = 155$ GeV. The lines are the same as in Fig. 1.

- { IFLAG_H(12)= 1: Gives the same result as that obtained by the older version of CPsuperH.
- { IFLAG_H(12)= 2: Includes only the threshold corrections to the neutral and charged Higgs-boson quark self-energies.
- { IFLAG_H(12)= 3: Includes only the threshold corrections to $\tilde{m}_i^{(1)}$.
- { IFLAG_H(12)= 4: Includes only the iterative method for the pole masses.
- { IFLAG_H(12)= 5 or 0: All the improvements are fully included.

IFLAG_H(60)= 1: This is an error message that appears when the iterative method for the pole masses fails.

The improvement in the threshold corrections to the top- and bottom-quark Yukawa couplings is important when $\tan\beta$ is large and the charged Higgs boson is light. In Figs. 1 and 2, we show the pole masses of the neutral Higgs bosons for the CPX [8] and trimixing [9] scenarios, respectively, when $\text{IFLAG_H}(12) = 2-5$ as indicated. In each frame, the old calculation with $\text{IFLAG_H}(12) = 1$ (dashed line) is also shown for comparison.

Finally, $\text{RAUX_H}(1-6)$, $\text{RAUX_H}(10-36)$, and $\text{CAUX_H}(1-2)$ are allocated for numerical information on the Higgs-sector calculation based on a renormalization-group-improved diagrammatic approach including dominant higher-order logarithmic and threshold corrections [6,7], see Tables 1 and 2.




Figure 3: The absolute value of each component of the neutral Higgs-boson propagator matrix $D^{H^0}(\hat{s})$ with (red solid lines) and without (black dashed lines) including off-diagonal absorptive parts in the trimixing scenario with $\beta_A = \beta_3 = 90$ and $\text{IFLAG_H}(12) = 5$. We note that $|D_{44}^{H^0}(\hat{s})| = 1$. The three Higgs-boson pole masses are indicated by thin vertical lines.

In situations where two or more MSSM Higgs bosons contribute simultaneously to a process, the transitions between the Higgs-boson mass eigenstates need to be considered before their decays. For this reason, we include the complete 4 × 4-dimensional propagator matrix $D^{H^0}(\hat{s})$ spanned by the basis $(H_1; H_2; H_3; G^0)$ [10], including off-diagonal absorptive parts [9]. The dimensionless neutral Higgs-boson propagator matrix is given by

$$D^{H^0}(\hat{s}) = \begin{array}{cccc} \hat{s} & M_{H_1}^2 + i\text{Im} b_{11}(\hat{s}) & i\text{Im} b_{12}(\hat{s}) & i\text{Im} b_{13}(\hat{s}) & i\text{Im} b_{14}(\hat{s}) \\ i\text{Im} b_{21}(\hat{s}) & \hat{s} & M_{H_2}^2 + i\text{Im} b_{22}(\hat{s}) & i\text{Im} b_{23}(\hat{s}) & i\text{Im} b_{24}(\hat{s}) \\ i\text{Im} b_{31}(\hat{s}) & i\text{Im} b_{32}(\hat{s}) & \hat{s} & M_{H_3}^2 + i\text{Im} b_{33}(\hat{s}) & i\text{Im} b_{34}(\hat{s}) \\ i\text{Im} b_{41}(\hat{s}) & i\text{Im} b_{42}(\hat{s}) & i\text{Im} b_{43}(\hat{s}) & \hat{s} & i\text{Im} b_{44}(\hat{s}) \end{array} \quad \begin{array}{l} 1 \\ 1 \\ C \\ C \\ C \\ A \end{array} \quad ; \quad (1)$$

where $M_{H_{1,2,3}}$ are the one-loop Higgs-boson pole masses, and higher-order absorptive effects on $M_{H_{1,2,3}}$ have been ignored [6]. The label ‘4’ refers to the would-be Goldstone boson of the Z boson. The absorptive part of the Higgs-boson propagator matrix receives contributions from loops of fermions, vector bosons, associated pairs of Higgs and vector bosons, Higgs-boson pairs, and sfermions:

$$i\text{Im} b_{ij}(\hat{s}) = i\text{Im} b_{ij}^{\text{ff}}(\hat{s}) + i\text{Im} b_{ij}^{\text{VV}}(\hat{s}) + i\text{Im} b_{ij}^{\text{HV}}(\hat{s}) + i\text{Im} b_{ij}^{\text{HH}}(\hat{s}) + i\text{Im} b_{ij}^{\text{ff}'}(\hat{s}); \quad (2)$$

respectively. We refer to Ref. [9] for their explicit expressions. For the Goldstone-Higgs mixings, $i\text{Im} b_{i4}$ and $i\text{Im} b_{44}$, we take the leading contributions ignoring all gauge-coupling mediated parts. We also include the 2 × 2-dimensional propagator matrix for the charged Higgs bosons $D^H(\hat{s})$ spanned by the basis $(H^\pm; G^\pm)$, including off-diagonal absorptive parts:

$$D^H(\hat{s}) = \begin{array}{cc} \hat{s} & M_H^2 + i\text{Im} b_{HH}(\hat{s}) & i\text{Im} b_{HG}(\hat{s}) \\ i\text{Im} b_{GH}(\hat{s}) & \hat{s} & i\text{Im} b_{GG}(\hat{s}) \end{array} \quad ; \quad (3)$$

The relevant Goldstone-boson couplings are given in Appendix B. For the 16 elements of the neutral Higgs-boson propagator matrix $D^{H^0}(\hat{s})$ and for the 4 elements of the charged Higgs-boson propagator matrix $D^H(\hat{s})$, the slots CAUX_H(100–119) are used as shown in Table 2. In Fig. 3, as an example, we show the absolute value of all components of the Higgs-boson propagator matrix $D^{H^0}(\hat{s})$ as functions of $\sqrt{\hat{s}}$ for the mixing scenario with $\alpha = \beta = 90^\circ$.

It is important to remark that the 4 × 4 propagator matrix (1) is sufficient to encode all H_i –Z- and G^0 –Z mixing effects within the Pinch Technique (PT) framework [10,11], which has been adopted here to remove consistently gauge-dependent and high-energy unitarity-violating terms from $i\text{Im} b_{ij}(\hat{s})$ [9]. For example, the self-energy transition $H_i \rightarrow Z$, $b_{ZH_i} = p b_{ZH_i}$, is related to $b_{G^0H_i}$ through

$$\hat{s} b_{ZH_i}(\hat{s}) = iM_Z^2 b_{G^0H_i}(\hat{s}); \quad (4)$$

with $\hat{s} = p^2$. We recall that the self-energy transitions $H_i \rightarrow G^0$ and $G^0 \rightarrow H_i$ are completely absent within the PT framework. More details may be found in [10].

Note that the elements of the propagator matrix depend on the center-of-mass energy, denoted by $\sqrt{\hat{s}}$, which is stored in RAUX_H(101), see Table 1. Along with $D^{H^0 \neq i}(\hat{s})$, the \hat{s} -dependent couplings of the neutral Higgs bosons to two gluons, $S_i^g(\sqrt{\hat{s}})$ and $P_i^g(\sqrt{\hat{s}})$, and two photons, $S_i(\sqrt{\hat{s}})$ and $P_i(\sqrt{\hat{s}})$, are needed when we consider the production of the neutral Higgs bosons and study its CP properties at the LHC [9,12,13] and a collider [14,16]. They are calculated and stored in CAUX_H(130–135) and CAUX_H(140–145) as shown in Table 2. We have included the dominant contributions coming from the tan-enhanced loops of sbottoms and gluinos and the subdominant ones coming from the stop-Higgsino mediated diagrams. Also included are the resummed corrections to Yukawa couplings. For the electroweak corrections, see next subsection. For the next-to-leading-order QCD corrections, appropriately calculated K factors should be taken into account separately in the calculation of production cross sections [17,18].

Two additional flags are used to control the inclusion of the off-diagonal absorptive parts and print out the \hat{s} -dependent propagator matrix and the \hat{s} -dependent Higgs couplings to two photons and gluons:

IFLAG_H(13)= 1: Does not include the off-diagonal absorptive parts in the propagator matrices $D^{H^0 \neq i}(\hat{s})$.

IFLAG_H(14)= 1: Prints out each component of the Higgs-boson propagator matrices $D^{H^0 \neq i}(\hat{s})$ and the \hat{s} -dependent couplings $S_i^g(\sqrt{\hat{s}})$ and $P_i^g(\sqrt{\hat{s}})$.

2.2 Improved Treatment of Higgs-Boson Couplings and Decays

The main updates include:

The electroweak corrections to the neutral Higgs couplings to pairs of tau leptons and b-quarks [19]. The explicit formulae used in the code for the corrections, including non-vanishing CP phase effects, could be found in Ref. [9] and Eqs.(A.1)–(A.2) of Ref. [3].

The three-body decay $H^+ \rightarrow tb + W^+ bb$. Some three-body decays play an important role in Higgs searches [20]. In addition to the three-body decays involving more than one massive gauge boson considered previously, we include the three-body decay $H^+ \rightarrow tb + W^+ bb$ in the new version. The decay width is given by

$$\Gamma(H^+ \rightarrow W^+ bb) = N_c \frac{g^2 g_{tb}^2 M_H}{512 \pi^3} \int_0^{1-w} dx_1 \int_{1-w-x_1}^{1-\frac{w}{x_1}} dx_2 \frac{F(x_1; x_2)}{(1-x_2-t+b)^2 + t^2}; \quad (5)$$

where $x_b = m_b^2/M_H^2$, $x_t = m_t^2/M_H^2$ and $x_i = 2E_i/M_H$ with E_1 and E_2 being the energies of the b and \bar{b} quarks, respectively. In the charged Higgs-boson rest frame, the function $F(x_1; x_2)$ is given by

$$\begin{aligned}
 F(x_1; x_2) = & \frac{1}{2} \left[\frac{(1-x_1)(1-x_2)}{W} + 2x_1 + 2x_2 + \frac{3+2W}{2} \frac{1}{x_b x_t} \right. \\
 & + \frac{x_2^3 + x_1 x_2^2 - 3x_2^2 - 2x_1 x_2 + 3x_2 + x_1 - 1}{W} \\
 & \left. + (x_2^2 + 2x_1 x_2 - 4x_2 - 2x_1 + 3 - 2W) \right. \\
 & \left. + \frac{2x_1 + 3 + 2W}{2} + \frac{2x_2^2 - x_1 x_2 + 5x_2 + x_1 - 3}{W} \right] \frac{1}{2} \frac{1}{x_b^2} \\
 & + 2 \frac{P}{x_b x_t} < e(g_L g_R) \frac{(x_2 - 1)^2}{W} + (x_2 + 1 - 2W) + 2 \frac{1}{x_b} ; \quad (6)
 \end{aligned}$$

where $g_L = g_{H^+tb}^S + ig_{H^+tb}^P$ and $g_R = g_{H^+tb}^S + ig_{H^+tb}^P$.

The contributions from tau-lepton and charm-quark loops to the couplings $S_i(M_{H_i})$ and $P_i(M_{H_i})$.

A new msg IFLAG H(57)= 1: This is an error message that appears when one of the magnitudes of the complex input parameters is negative.

The CPsuperH homepage has been continuously brought up to date after its first appearance to include the updates discussed in this subsection and others not mentioned here. We refer to the file OLIST_V1 for a full list of updates to the original version which can be found in the CPsuperH homepage.

3 Higgs-Mediated Two-Loop Electric Dipole Moments

The CP phases in the MSSM are significantly constrained by measurements of Electric Dipole Moments (EDMs). In particular, the EDM of the Thallium atom may provide currently the most stringent constraint on MSSM scenarios with explicit CP violation. The atomic EDM of ^{205}Tl gets its main contributions from two terms [21,22]:

$$\begin{aligned}
 d_{T1}[\text{cm}] = & 585 d[\text{cm}] + 8.5 \cdot 10^{19} [\text{cm}] (\mathcal{G} \text{TeV}^2) + \dots ; \\
 & (d_{T1})^e[\text{cm}] + (d_{T1})^{Cs}[\text{cm}] + \dots ; \quad (7)
 \end{aligned}$$

where d_e denotes the electron EDM and C_S is the coefficient of the CP-odd electron-nucleon interaction $L_{C_S} = C_S e i_5 e N N$. The dots denote sub-dominant contributions from 6-dimensional tensor and higher-dimensional operators.




Figure 4: The Thallium EDM $\hat{d}_{Tl}^H = d_{Tl}^H / 10^{24}$ [e cm] in the CPX scenario with $\mu = 90$ and $M_{SUSY} = 0.5$ TeV taking $IFLAG_H(12) = 5$ [25]. The different shaded regions correspond to different ranges of $|\hat{d}_{Tl}^H|$ as shown. Specially, in the narrow region denoted by black squares, one has $|\hat{d}_{Tl}^H| < 1$, consistent with the current Thallium EDM constraint.

The contributions of the first- and second-generation phases, $\mathcal{A}_{e;}$ and $\mathcal{A}_{d;}$, to EDMs can be drastically reduced either by assuming that these phases are sufficiently small, or if the first- and second-generation squarks and sleptons are sufficiently heavy. However, even when the contributions of the first and second generation phases to EDMs are suppressed, there are still sizeable contributions to EDMs from Higgs-mediated two-loop diagrams [23].

The Higgs-mediated two-loop Thallium (d_{Tl}^H), electron (d_e^H), and muon (d^μ) EDMs are calculated and stored in `RAUX_H(111-120)` as shown in Table 1. The Thallium and electron EDMs consist of:

$$\begin{aligned}
 d_{Tl}^H &= (d_{Tl}^H)^e + (d_{Tl}^H)^{Cs} ; \\
 d_e^H &= (d_e^H)^{\tau} + (d_e^H)^{\beta} + (d_e^H)^{\tau} + (d_e^H)^{\beta} + (d_e^H)^{\sim} :
 \end{aligned} \tag{8}$$

The explicit expressions for the EDMs in the CPsuperH conventions and notations may be

found in Ref. [24]. A flag IFLAG_H(15)= 1 is used to print out the results of the EDM calculations:

IFLAG_H(15)= 1: Print out EDM s.

In Fig. 4, we show the rescaled Thallium EDM $\hat{d}_{T1}^H = d_{T1}^H / 10^{24}$ in units of ecm in the $\tan \beta - M_{H_1}$ plane, in the CPX scenario with IFLAG_H(12)= 5. We observe, when $\tan \beta < 5$ and $M_{H_1} < 10$ GeV, one may have $|\hat{d}_{T1}^H| < 1$ only in the narrow region denoted by black squares which is consistent with the current 2-loop upper bound on the Thallium EDM [26]: $|\hat{d}_{T1}^H| < 1.3 \times 10^{-24}$ [ecm]. We note that the region $8 \text{ GeV} < M_{H_1} < 10 \text{ GeV}$ with $\tan \beta < 10$ has not been excluded by the combined constraints from the LEP searches [27] and the $(1S) \rightarrow H_1$ decay [28].

The Thallium EDM constraint can be evaded by assuming cancellations between the two-loop contributions considered here and possible one-loop contributions which depend on different CP-odd phases related to the first and second generations of squarks and sleptons. For example, assuming cancellation of less than 1 part in 10, the region with $|\hat{d}_{T1}^H| < 10$ in Fig. 4 is allowed. In the future, this treatment of the most important two-loop contributions to the Thallium EDM will be supplemented by a more complete implementation of calculations of the well-known 1-loop contributions to this and other EDMs.

4 B-Meson Observables

An important innovation in CPsuperH2.0 is the inclusion of the following important Higgs-mediated B-meson observables:

The branching ratio of B_s meson into a pair of muons: $B(B_s \rightarrow \mu\mu)$,

The branching ratio of B_d meson into a pair of tau leptons: $B(B_d \rightarrow \tau\tau)$,

The SUSY contribution to the $B_d^0 - \bar{B}_d^0$ mass difference: $M_{B_d}^{\text{SUSY}}$,

The SUSY contribution to the $B_s^0 - \bar{B}_s^0$ mass difference: $M_{B_s}^{\text{SUSY}}$,

The ratio of the branching ratio $B(B_u \rightarrow \mu\mu)$ to the SM value:

$$R_B = \frac{B(B_u \rightarrow \mu\mu)}{B^{\text{SM}}(B_u \rightarrow \mu\mu)}$$

,

The branching ratio $B(B \rightarrow X_s \gamma)$ and the direct CP asymmetry $A_{\text{CP}}(B \rightarrow X_s \gamma)$.

We adopt the most recent gauge-invariant and flavour-covariant formalism to calculate the flavour-changing effective Lagrangian for the interactions of the neutral and charged Higgs fields to the up- and down-type quarks including a new class of dominant subleading contributions [5]. In the current version, the single-Higgs insertion approximation is used.

For the calculations of B-meson observables, the array SMPARA_H for the SM parameters has been extended to include information on the CKM matrix, parameterized via θ , A , δ , and ϕ , as seen in Table 3. The CKM matrix is constructed as [29]

$$V = \begin{pmatrix} 1 & 0 & 0 \\ s_{12} c_{23} & c_{12} c_{23} & s_{12} c_{13} \\ s_{12} s_{23} & c_{12} s_{23} & s_{12} s_{23} s_{13} e^{i\delta} \\ s_{12} s_{23} & c_{12} c_{23} s_{13} e^{i\delta} & c_{12} s_{23} s_{13} e^{i\delta} \\ c_{12} c_{23} & s_{12} c_{23} & s_{12} c_{23} s_{13} e^{i\delta} \\ c_{12} s_{23} & s_{12} s_{23} & s_{12} s_{23} s_{13} e^{i\delta} \\ c_{23} c_{13} & c_{23} s_{13} e^{i\delta} & c_{23} s_{13} e^{i\delta} \end{pmatrix} ; \quad (9)$$

where $s_{ij} = \sin \theta_{ij}$, $c_{ij} = \cos \theta_{ij}$, and δ is the KM phase with $s_{ij}, c_{ij} \geq 0$. In terms of θ , A , δ , and ϕ , they are given by

$$s_{12} = \frac{A}{\sqrt{1+A^2}} ; \quad s_{23} = A^2 ; \quad s_{13} e^{i\delta} = \frac{A^3 (1+i)^{\frac{1}{2}} \sqrt{1-A^2}}{1+A^2} ; \quad (10)$$

and $c_{ij} = \frac{1}{\sqrt{1+s_{ij}^2}}$. The SUSY parameter array SSPARA_H is also extended to include the hierarchy factors $\rho_{\tilde{Q}, \tilde{U}, \tilde{D}, \tilde{L}, \tilde{E}}$ between the first two and third generations [30], see Table 4. In the super-CKM basis, the 3×3 squark mass matrices squared are taken to be diagonal:

$$\begin{aligned} \tilde{M}_Q^2 &= m_{\tilde{Q}_3}^2 \text{diag} \left(\frac{\rho_{\tilde{Q}}^2}{\rho_{\tilde{Q}}}; \frac{\rho_{\tilde{Q}}^2}{\rho_{\tilde{Q}}}; 1 \right); \\ \tilde{M}_U^2 &= m_{\tilde{U}_3}^2 \text{diag} \left(\frac{\rho_{\tilde{U}}^2}{\rho_{\tilde{U}}}; \frac{\rho_{\tilde{U}}^2}{\rho_{\tilde{U}}}; 1 \right); \\ \tilde{M}_D^2 &= m_{\tilde{D}_3}^2 \text{diag} \left(\frac{\rho_{\tilde{D}}^2}{\rho_{\tilde{D}}}; \frac{\rho_{\tilde{D}}^2}{\rho_{\tilde{D}}}; 1 \right); \\ \tilde{M}_L^2 &= m_{\tilde{L}_3}^2 \text{diag} \left(\frac{\rho_{\tilde{L}}^2}{\rho_{\tilde{L}}}; \frac{\rho_{\tilde{L}}^2}{\rho_{\tilde{L}}}; 1 \right); \\ \tilde{M}_E^2 &= m_{\tilde{E}_3}^2 \text{diag} \left(\frac{\rho_{\tilde{E}}^2}{\rho_{\tilde{E}}}; \frac{\rho_{\tilde{E}}^2}{\rho_{\tilde{E}}}; 1 \right); \end{aligned} \quad (11)$$

Finally, the results for the B-meson observables are stored in RAUX_H(130–136) as shown in Table 1. The SUSY contributions to the $B \rightarrow 2$ transition amplitudes are stored in CAUX_H(150) and CAUX_H(151), see Table 1. Note the relations $\text{RAUX_H}(132) = 2 \sqrt{\text{CAUX_H}(150)}$ and $\text{RAUX_H}(133) = 2 \sqrt{\text{CAUX_H}(151)}$. Two flags IFLAG_H(16) and IFLAG_H(17) are used to print out the results of the calculation of B-meson observables:

IFLAG_H(16)= 1: Print out B-meson observables.

IFLAG_H(17)= 1: Print out details of the $B \rightarrow X_s$ calculation.

For numerical examples of B-meson observables, we take the CPX scenario [8] with $M_{\text{SUSY}} = 0.5 \text{ TeV}$ and the common A-term phase $A_{\tilde{Q}} = A_{\tilde{U}} = A_{\tilde{D}} = A_{\tilde{L}} = A_{\tilde{E}} = A$ in the convention

$= 0$. We take account of the dependence on the hierarchy factors $q_{\mu, \nu, \rho}$ between the first two and the third generations, taking a common value for the three of them.

Figure 5 shows the dependence of the branching ratio $B(B_s \rightarrow \tau^+ \nu_\tau)$ on the phase of the gluino mass parameter ϕ_3 for four values of $\tan \beta$. The charged Higgs-boson pole mass is fixed at $M_{H^\pm} = 200$ GeV. In each frame, two sets of three lines are shown. The upper lines are for higher $\tan \beta = 10$ and the lower ones for $\tan \beta = 1$. For fixed ϕ_3 , three lines show the cases of $\phi_A = 0$ (solid), 90° (dashed), and 180° (dash-dotted). The dependence is shown in Fig. 6. We clearly see the GIM operative point mechanism discussed in Ref. [30] around $\phi_3 \approx 1.2$ when $(\phi_3; \phi_A) = (0; 180^\circ)$ (solid lines). Figure 7 shows the rescaled branching ratio $\tilde{B} = B(B_s \rightarrow \tau^+ \nu_\tau) \cdot 10^7$ in the M_{H_1} - $\tan \beta$ plane when the phases are fixed at $\phi_A = \phi_3 = 90^\circ$. The unshaded region is not theoretically allowed. Only the region with $\tilde{B} < 0.58$ is consistent with the current experimental upper limit at 95% C.L., corresponding to $\tan \beta < 20$ (8) for $\tan \beta = 1$ (10).

The rescaled branching ratio $\tilde{B}_s = B(B \rightarrow X_s) \cdot 10^4$ is shown in Fig. 8. In contrast to the $B_s \rightarrow \tau^+ \nu_\tau$ case, we observe that higher $\tan \beta$ region is experimentally allowed: $\tan \beta > 35$ (20) for $\tan \beta = 1$ (10). This is because the charged-Higgs contribution is suppressed due to the threshold corrections when $\tan \beta$ is large. The charged-Higgs contribution to $B \rightarrow X_s$ is proportional to $1/(1 + |j|^2 \tan^2 \beta)$ [33], where j represents the threshold corrections with $|j| \approx 0.05$ for the parameters chosen [34].

Figure 9 shows the ratio of the branching ratio $B(B_u \rightarrow \tau^+ \nu_\tau)$ to its SM value, R_B . In the left frame with $\tan \beta = 1$, we see two connected bands of the experimentally allowed 1- region, $0.62 < R_B < 1.38$. If we consider the 2- limit, only the upper-left region with $M_{H_1} < 95$ GeV and $\tan \beta > 35$ is not allowed. For larger $\tan \beta = 10$, the allowed region becomes narrower.

In Fig. 10, we show the region satisfying the experimental constraints from $B(B_s \rightarrow \tau^+ \nu_\tau)$ (95%), $B(B \rightarrow X_s)$ (2%), and R_B (1%). First we observe that there is no region that satisfies the $B_s \rightarrow \tau^+ \nu_\tau$ and $B \rightarrow X_s$ constraints simultaneously for both $\tan \beta = 1$ and 10. If one neglects the constraint from $B(B_s \rightarrow \tau^+ \nu_\tau)$, only the high- $\tan \beta$ region would remain. Taking account of $B_u \rightarrow \tau^+ \nu_\tau$ constraint, the region with $\tan \beta > 36$ and $M_{H_1} > 80$ GeV is allowed when $\tan \beta = 1$. On the other hand, neglecting the constraint from $B(B \rightarrow X_s)$, the allowed region is constrained in the parameter space with $\tan \beta < 20$ and $M_{H_1} > 10$ GeV for $\tan \beta = 1$. For $\tan \beta = 10$, the $B \rightarrow X_s$ constraint is relaxed but those from $B(B_s \rightarrow \tau^+ \nu_\tau)$ and R_B become more stringent.

Finally, in Fig. 11, we show the region allowed experimentally by the measurement $B(B_d \rightarrow \tau^+ \nu_\tau) < 4.1 \cdot 10^{-3}$ (90%) [36] (upper frames) and the regions where the SUSY contribution is smaller than the measured values of $B_s^0 - B_s^0$ mass difference [37] (middle frames) and $B_d^0 - B_d^0$ mass difference [29] (lower frames). We see that the $B(B_d \rightarrow \tau^+ \nu_\tau)$ constraint has the least impact on these parameter planes, whereas the impacts of the $B_s^0 - B_s^0$ and $B_d^0 - B_d^0$ mass differences are similar.

These examples illustrate the possible interplays between the different B-meson observables, and how they may vary significantly with the values of the CP-violating phases. CPsuperH2.0 provides a unique tool for combining these constraints and pursuing their implications for other observables. In the future, the CPsuperH2.0 treatment of these important B-meson observables will be supplemented by the implementation of calculations of other flavour observables, including the K sector.

5 Summary and Outlook

We have presented in this paper a description of the new features of the Fortran code CPsuperH2.0. In addition to improved calculations of the Higgs-boson pole masses with more complete treatment of threshold effects in self-energies and Yukawa couplings, the complete 4×4 (2 \times 2) neutral (charged) Higgs-boson propagator matrices with the Goldstone-Higgs mixing effects have been consistently implemented. Specially, the neutral Higgs-boson propagator matrix constitutes a necessary ingredient for the studies of a system of strongly-mixed Higgs bosons at colliders together with the center-of-mass dependent Higgs-boson couplings to gluons and photons. It also provides the improved Higgs-boson couplings to tau leptons, b quarks, and two photons. The important three-body decay $H^+ \rightarrow t b \rightarrow W^+ b b$ is included.

In order to provide a more complete, consistent tool for calculating CP-violating observables in the MSSM, and specially to incorporate the important constraints coming from precision experiments at low energies, CPsuperH2.0 has been extended to include a number of B-meson observables, as well as the Higgs-mediated two-loop contributions to EDMs of the Thallium atom, electron and muon. The currently available B-meson observables are the branching ratios of $B_s \rightarrow \pi^+ \pi^-$, $B_d \rightarrow \pi^+ \pi^-$, $B_u \rightarrow \pi^+ \pi^-$, $B \rightarrow X_s$ and the latter's CP-violating asymmetry A_{CP} , and the supersymmetric contributions to the $B_{s,d}^0 - \bar{B}_{s,d}^0$ mass differences. Further low-energy observables are to be included in future updates.

The improved Fortran code CPsuperH2.0 provides a coherent and complete numerical framework in which one can calculate consistently observables in both low- and high-energy experiments probing physics beyond the SM.

Acknowledgements

The work of J.S.L. was supported in part by the Korea Research Foundation and the Korean Federation of Science and Technology Societies Grant funded by the Korea Government (M.O.E.H.R.D., Basic Research Promotion Fund) and in part by the National Science Council of Taiwan, R.O.C. under Grant No. NSC 96-2811-M-008-068. The work of A.P. was supported in part by the STFC research grant: PP/D000157/1. Work at ANL is supported in part

by the U S D O E , D iv . of H E P , Contract D E - A C 02-06C H 11357 . Ferm ilab is operated by U niversities R esearch A ssociation I nc. under contract no. D E - A C 02-76C H 02000 w ith the D O E . W e thank S . Y . C hoi and M . D rees for past collaboration on C PsuperH , and for discussions on this updated version .

A List of changes

Here we summarize the improved features introduced in CPsuperH2.0 compared to the prior version of CPsuperH.

New common blocks:

- { COMMON /HC_RAUX/ RAUX_H(NAUX= 999), see Table 1
- { COMMON /HC_CAUX/ CAUX_H(NAUX= 999), see Table 2

Extended arrays for input parameters:

- { SMPARA_H(NSMIN= 19), see Table 3
- { SMPARA_H(NSSIN= 26), see Table 4

New names for improved FORTRAN files:

- { cpsuperh.f ! cpsuperh2.f
- { fillpara.f ! fillpara2.f
- { fillhiggs.f ! fillhiggs2.f
- { fillcoupl.f ! fillcoupl2.f
- { fillgambr.f ! fillgambr2.f

New FORTRAN files:

- { filldhpq.f is to calculate the full propagator matrices $D^{H^0 \#i}(\hat{s})$ and the \hat{s} -dependent couplings $S_i^{g'j}(\hat{s})$ and $P_i^{g'j}(\hat{s})$.
- { higgsedm.f is to calculate Higgs-mediated two-loop EDMs of Thallium, electron, and muon.
- { fillbobs.f is to calculate the B-meson observables: $B(B_s \rightarrow \dots)$, $B(B_d \rightarrow \dots)$, $M_{B_d}^{SU(2)_C}$, $M_{B_s}^{SU(2)_C}$, R_B , $B(B \rightarrow X_s)$, and $A_{CP}(B \rightarrow X_s)$.

New flags:

- { IFLAG_H(12) = 0 5: For the level of improvement in the calculation of the Higgs-boson pole masses.
- { IFLAG_H(13) = 1: Not to include the off-diagonal absorptive parts in the propagator matrices $D^{H^0 \#i}(\hat{s})$.
- { IFLAG_H(14) = 1: Print out the elements of the full propagator matrices $D^{H^0 \#i}(\hat{s})$ and the \hat{s} -dependent couplings $S_i^{g'j}(\hat{s})$ and $P_i^{g'j}(\hat{s})$.

- { IFLAG_H(15) = 1: Print out EDM s.
- { IFLAG_H(16) = 1: Print out B -m eson observables.
- { IFLAG_H(17) = 1: Print out B ! X_s details.
- { IFLAG_H(57) = 1: This is an error m essage that appears when one of the m agnitudes of the com plex SU SY input param eters is negative.
- { IFLAG_H(60) = 1: This is an error m essage that appears when the iterative m ethod for the neutral H iggs-boson pole m asses fails.

B Goldstone-boson couplings to third-generation ferm ions and sferm ions

Here we present the Goldstone-(s)ferm ion-(s)ferm ion couplings in the CPsuperH convention.

G⁰-f-f

$$L_{G^0 f f} = \sum_{f=t,b} \frac{g m_f}{2M_W} G^0 f i g_{G^0 f f}^P \gamma_5 f ; \quad (B.1)$$

where

$$g_{G^0 t t}^P = 1 ; \quad g_{G^0 b b}^P = g_{G^0}^P = +1 ; \quad (B.2)$$

G⁻-f-f⁰

$$\begin{aligned} L_{G^- f f^0} &= \frac{g}{2M_W} \sum_{(f^*, f_\#) = (t^*, b); (\tilde{f}, \tilde{f}^*)} G^+ f^* m_{f^*} P_L m_{f_\#} P_R f_\# + h.c. ; \quad (B.3) \\ &= g_{tb} G^+ t (g_{G^+ tb}^S + i g_{G^+ tb}^P \gamma_5) b + g G^+ (g_{G^+}^S + i g_{G^+}^P \gamma_5) + h.c. ; \end{aligned}$$

where

$$\begin{aligned} g_{tb} &= \frac{g m_t}{2M_W} ; \quad g_{G^+ tb}^S = \frac{1 - m_b m_t}{2} ; \quad g_{G^+ tb}^P = i \frac{1 + m_b m_t}{2} ; \\ g &= \frac{g m}{2M_W} ; \quad g_{G^+}^S = \frac{1}{2} ; \quad g_{G^+}^P = i \frac{1}{2} ; \quad (B.4) \end{aligned}$$

G⁰-f^c-f^c

$$L_{G^0 f^c f^c} = \sum_{f=t,b} v \sum_{f_1, f_2} g_{G^0 f_1 f_2} (G^0 f_1^c f_2^c) ; \quad (B.5)$$

where

$$v g_{G^0 f_1 f_2} = G^0 f^c f^c U_{f_1}^{f^c} U_{f_2}^{f^c} ; \quad (B.6)$$

The couplings in the weak-interaction basis are given by

$$\begin{aligned}
G^0_{\tau\tau} &= \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & ih_t(s A_t c) \\ ih_t(s A_t c) & 0 \end{pmatrix}; \\
G^0_{\mathbb{B}\mathbb{B}} &= \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & ih_b(c A_b s) \\ ih_b(c A_b s) & 0 \end{pmatrix}; \\
G^0_{\tilde{\nu}\tilde{\nu}} &= \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & ih(c A s) \\ ih(c A s) & 0 \end{pmatrix}; \quad (B.7)
\end{aligned}$$

G_{ff^0}

$$L_{G_{ff^0}} = v g_{G^+ \tau_i \mathbb{B}_j} (G^+ \tau_i \mathbb{B}_j) + v g_{G^+ \tilde{\nu}_i} (G^+ \tilde{\nu}_i) + h.c.; \quad (B.8)$$

where

$$v g_{G^+ \tau_i \mathbb{B}_j} = G^+ \tau \mathbb{B} U_i^\tau U_j^\mathbb{B} \quad \text{and} \quad v g_{G^+ \tilde{\nu}_i} = G^+ \tilde{\nu} U_i^{\tilde{\nu}}; \quad (B.9)$$

The couplings in the weak-interaction basis are given by

$$\begin{aligned}
G^+_{\tau\mathbb{B}} &= \frac{1}{2} (h_u f s^2 - h_d f c^2) v + \frac{1}{2\sqrt{2}} g^2 c_2 v \begin{pmatrix} h_d(c A_d s) \\ h_u(s A_u c) \end{pmatrix}; \\
G^+_{\tilde{\nu}\tilde{L}} &= \frac{1}{2} h f c^2 v + \frac{1}{2\sqrt{2}} g^2 c_2 v; \\
G^+_{\tilde{\nu}\tilde{R}} &= h(c A s); \quad (B.10)
\end{aligned}$$

C Sample new outputs

Here we show the new outputs of CPsuperH2.0 for the CPX scenario with $\tan\beta = 5$, $M_H = 300 \text{ GeV}$, $M_{\text{SUSY}} = 500 \text{ GeV}$, and $A = \mu = 3 = 90$.

IFLAG H(1) = 1: In the new version, we are using $m_b(m_t^{\text{pole}}) = 3.155 \text{ GeV}$ and $m_c(m_t^{\text{pole}}) = 0.735 \text{ GeV}$ as defaults. Note also that the list of the SM and SUSY input parameters is extended to include the CKM matrix and the diagonal sfermion mass matrices.

Standard Model Parameters in /HC_SMPARA/

```

AEM_H   = 0.7812E-02 : alpha_em(MZ)
ASMZ_H  = 0.1185E+00 : alpha_s(MZ)
MZ_H    = 0.9119E+02 : Z boson mass in GeV
SW_H    = 0.4808E+00 : sinTheta_W
ME_H    = 0.5000E-03 : electron mass in GeV

```

MMU_H = 0.1065E+00 : muon mass in GeV
MTAU_H = 0.1777E+01 : tau mass in GeV
MDMT_H = 0.4000E-02 : d-quark mass at M_t^{pole} in GeV
MSMT_H = 0.9000E-01 : s-quark mass at M_t^{pole} in GeV
MBMT_H = 0.3155E+01 : b-quark mass at M_t^{pole} in GeV
MUMT_H = 0.2000E-02 : u-quark mass at M_t^{pole} in GeV
MCMT_H = 0.7350E+00 : c-quark mass at M_t^{pole} in GeV
MTPOLE_H = 0.1743E+03 : t-quark pole mass in GeV
GAMW_H = 0.2118E+01 : Γ_W in GeV
GAMZ_H = 0.2495E+01 : Γ_Z in GeV
EEM_H = 0.3133E+00 : $e = (4\pi\alpha_{\text{em}})^{1/2}$
ASMT_H = 0.1084E+00 : $\alpha_s(M_t^{\text{pole}})$
CW_H = 0.8768E+00 : $\cos\theta_W$
TW_H = 0.5483E+00 : $\tan\theta_W$
MW_H = 0.7996E+02 : W boson mass $M_W = M_Z \cdot C_W$
GW_H = 0.6517E+00 : SU(2) gauge coupling $g_W = e/s_W$
GP_H = 0.3573E+00 : U(1)_Y gauge coupling $g_P = e/c_W$
V_H = 0.2454E+03 : $V = 2 M_W / g_W$
GF_H = 0.1174E-04 : $G_F = \sqrt{2} \cdot g_W^2 / 8 M_W^2$ in GeV^{-2}
MTMT_H = 0.1666E+03 : t-quark mass at M_t^{pole} in GeV

CKM Matrix :

$|V_{ud}| = |(0.9738E+00 \ 0.0000E+00)| = 0.9738E+00$
 $|V_{us}| = |(0.2272E+00 \ 0.0000E+00)| = 0.2272E+00$
 $|V_{ub}| = |(0.2174E-02 \ -.3349E-02)| = 0.3993E-02$
 $|V_{cd}| = |(-.2271E+00 \ -.1377E-03)| = 0.2271E+00$
 $|V_{cs}| = |(0.9730E+00 \ -.3213E-04)| = 0.9730E+00$
 $|V_{cb}| = |(0.4222E-01 \ 0.0000E+00)| = 0.4222E-01$
 $|V_{td}| = |(0.7478E-02 \ -.3259E-02)| = 0.8157E-02$
 $|V_{ts}| = |(-.4161E-01 \ -.7602E-03)| = 0.4162E-01$
 $|V_{tb}| = |(0.9991E+00 \ 0.0000E+00)| = 0.9991E+00$

Real SUSY Parameters in /HC_RSUSYPARA/

TB_H = 0.5000E+01 : $\tan(\beta)$
CB_H = 0.1961E+00 : $\cos(\beta)$
SB_H = 0.9806E+00 : $\sin(\beta)$
MQ3_H = 0.5000E+03 : $M_{\tilde{Q}_3}$ in GeV
MU3_H = 0.5000E+03 : $M_{\tilde{U}_3}$ in GeV
MD3_H = 0.5000E+03 : $M_{\tilde{D}_3}$ in GeV

ML3_H = 0.5000E+03 : M_tildeL3 in GeV
ME3_H = 0.5000E+03 : M_tildeE3 in GeV

Complex SUSY Parameters in /HC_CSUSYPARA/

|MU_H| = 0.2000E+04:Mag. of MU parameter in GeV
|M1_H| = 0.5000E+02:Mag. of M1 parameter in GeV
|M2_H| = 0.1000E+03:Mag. of M2 parameter in GeV
|M3_H| = 0.1000E+04:Mag. of M3 parameter in GeV
|AT_H| = 0.1000E+04:Mag. of AT parameter in GeV
|AB_H| = 0.1000E+04:Mag. of AB parameter in GeV
|ATAU_H| = 0.1000E+04:Mag. of ATAU parameter in GeV
ARG(MU_H) = 0.0000E+00:Arg. of MU parameter in Degree
ARG(M1_H) = 0.0000E+00:Arg. of M1 parameter in Degree
ARG(M2_H) = 0.0000E+00:Arg. of M2 parameter in Degree
ARG(M3_H) = 0.9000E+02:Arg. of M3 parameter in Degree
ARG(AT_H) = 0.9000E+02:Arg. of AT parameter in Degree
ARG(AB_H) = 0.9000E+02:Arg. of AB parameter in Degree
ARG(ATAU_H) = 0.9000E+02:Arg. of ATAU parameter in Degree

Diagonal Sfermion Mass Matrices [GeV] (Not squared) :

M_Q = 0.5000E+03 x Diag(0.1000E+01 0.1000E+01 0.1000E+01)
M_U = 0.5000E+03 x Diag(0.1000E+01 0.1000E+01 0.1000E+01)
M_D = 0.5000E+03 x Diag(0.1000E+01 0.1000E+01 0.1000E+01)
M_L = 0.5000E+03 x Diag(0.1000E+01 0.1000E+01 0.1000E+01)
M_E = 0.5000E+03 x Diag(0.1000E+01 0.1000E+01 0.1000E+01)

Charged Higgs boson pole mass : 0.3000E+03 GeV

IFLAG H(2) = 1: The masses and mixing matrix of the neutral Higgs boson change due to the improvement in their calculations and the new input for the b-quark mass.

Masses and Mixing Matrix of Higgs bosons :

HMASS_H(I) and OMIX_H(A,I)

H1 Pole Mass = 0.1193E+03 GeV
H2 Pole Mass = 0.2718E+03 GeV
H3 Pole Mass = 0.2983E+03 GeV

Charged Higgs Pole Mass = 0.3000E+03 GeV [SSPARA_H(2)]

	[H1]	[H2]	[H3]	
[phi_1] /	0.2457E+00	0.3360E+00	0.9093E+00	n
O(IA,IH)= [phi_2]	0.9693E+00	-.7551E-01	-.2340E+00	
[a] n	-.9973E-02	0.9388E+00	-.3442E+00	/

IFLAG H(14) = 1: The elements of the propagator matrices $D^{H^0 \#i}(\hat{s})$ and the \hat{s} -dependent couplings of the neutral Higgs bosons to two photons, $S_i(\hat{s})$ and $P_i(\hat{s})$, and two gluons, $S_i^g(\hat{s})$ and $P_i^g(\hat{s})$, taking $\hat{s} = M_{H_i}^2$. The couplings are compared to their values at the Higgs-boson pole masses: $S_i(\hat{s} = M_{IH}) = \text{NHC_H}(88;IH)$, $P_i(\hat{s} = M_{IH}) = \text{NHC_H}(89;IH)$, $S_i^g(\hat{s} = M_{IH}) = \text{NHC_H}(84;IH)$, $P_i^g(\hat{s} = M_{IH}) = \text{NHC_H}(85;IH)$.

DNH4 at sqrt(s) = 0.2718E+03 GeV

DNH4[H1,H1]:	(0.1238E+01 0.2290E-01)	= 0.1238E+01
DNH4[H2,H2]:	(0.5542E-01 -.1611E+04)	= 0.1611E+04
DNH4[H3,H3]:	(-.4876E+01 -.2128E-01)	= 0.4876E+01
DNH4[H1,H2]:	(-.1607E+00 -.2973E-02)	= 0.1608E+00
DNH4[H1,H3]:	(-.5956E-05 0.2606E-03)	= 0.2606E-03
DNH4[H2,H3]:	(-.4893E-01 -.2377E-03)	= 0.4893E-01
DNH4[G0,H1]:	(0.3403E-06 -.1825E-04)	= 0.1825E-04
DNH4[G0,H2]:	(0.1872E+00 0.2446E-05)	= 0.1872E+00
DNH4[G0,H3]:	(-.2222E-06 0.5181E-04)	= 0.5182E-04
DNH4[G0,G0]:	(0.1000E+01 -.2365E-05)	= 0.1000E+01

DCH2 at sqrt(s) = 0.2718E+03 GeV

DCH2[H+,H+]:	(-.4576E+01 -.2790E-01)	= 0.4576E+01
DCH2[H+,G+]:	(-.1256E-03 0.2294E-01)	= 0.2294E-01
DCH2[G+,H+]:	(-.1256E-03 0.2294E-01)	= 0.2294E-01
DCH2[G+,G+]:	(0.1000E+01 -.6202E-03)	= 0.1000E+01

Comparisons of the H-photon-photon couplings at MH^pole and those at sqrt(s) = 0.2718E+03 GeV

	S couplings	P couplings
H1PP(M):	(-.6615E+01 0.6386E-01)	(0.1303E-01 0.7314E-03)


```

B(B_d -> tau tau)   x 10^7 = 0.2294E+00
ACP(B -> X_s gamma) x 10^2 = -.7954E-01 [%]
Delta M [B_d] (SUSY)      = 0.6659E-04 [1/ps]
Delta M [B_s] (SUSY)      = 0.1982E-01 [1/ps]

```

IFLAG H(17) = 1: The details of the $B \rightarrow X_s \gamma$ calculation. As a default, we use $m_c(m_c = m_c^{\text{pole}})$ to capture a part of NNLO corrections [38]. The case when only the charged-Higgs contribution is added to the SM prediction is also shown.

```

                B -> X_s gamma
delta and E_gamma^cut [GeV]: 0.3333E+00  0.1601E+01

```

```

b-q masses [GeV] (pole,      @mb^pole,  @mt^pole):
                    0.4802E+01  0.4415E+01  0.3155E+01
c-q masses [GeV] (pole,      @mc^pole,  @mb^pole):
                    0.1415E+01  0.1250E+01  0.1029E+01
mu_b and mu_c [GeV]      : 0.4802E+01  0.1415E+01

```

```

BR   x 10^4: 0.4396E+01 (SM+Charged Higgs+Chargino)
          [0.4471E+01 (SM+Charged Higgs)]
          [0.3351E+01 (SM)]
ACP  x 10^2: -.7954E-01 %

```

References

- [1] For a recent review, see T. Ibrahim and P. Nath, arXiv:0705.2008 [hep-ph].
- [2] M. Trodden, In the Proceedings of 32nd SLAC Summer Institute on Particle Physics (SSI 2004): Natures Greatest Puzzles, Menlo Park, California, 2-13 Aug 2004, pp L018 [arXiv:hep-ph/0411301]; M. Quiros, J. Phys. A 40 (2007) 6573; W. Buchmuller, arXiv:0710.5857 [hep-ph].
- [3] J. S. Lee, A. Pilaftsis, M. Carena, S. Y. Choi, M. Drees, J. R. Ellis and C. E. M. Wagner, Comput. Phys. Commun. 156 (2004) 283 [arXiv:hep-ph/0307377].
- [4] S. Heinemeyer, W. Hollik and G. Weiglein, Comp. Phys. Comm. 124 2000 76, hep-ph/9812320; T. Hahn, S. Heinemeyer, W. Hollik, H. Rzehak and G. Weiglein, arXiv:0710.4891 [hep-ph].
- [5] J. Ellis, J. S. Lee and A. Pilaftsis, arXiv:0708.2079 [hep-ph].
- [6] M. Carena, J. R. Ellis, A. Pilaftsis and C. E. M. Wagner, Nucl. Phys. B 625 (2002) 345 [arXiv:hep-ph/0111245].
- [7] M. Carena, J. R. Ellis, A. Pilaftsis and C. E. M. Wagner, Nucl. Phys. B 586 (2000) 92 [arXiv:hep-ph/0003180].
- [8] M. Carena, J. R. Ellis, A. Pilaftsis and C. E. M. Wagner, Phys. Lett. B 495 (2000) 155 [arXiv:hep-ph/0009212].
- [9] J. R. Ellis, J. S. Lee and A. Pilaftsis, Phys. Rev. D 70 (2004) 075010 [arXiv:hep-ph/0404167].
- [10] A. Pilaftsis, Nucl. Phys. B 504 (1997) 61.
- [11] J. M. Cornwall and J. Papavassiliou, Phys. Rev. D 40 (1989) 3474; J. Papavassiliou, Phys. Rev. D 41 (1990) 3179; D. Binosi and J. Papavassiliou, Phys. Rev. D 66 (2002) 111901; J. Phys. G 30 (2004) 203.
- [12] H. M. Georgi, S. L. Glashow, M. E. Machacek and D. V. Nanopoulos, Phys. Rev. Lett. 40 (1978) 692; T. Inami, T. Kubota and Y. Okada, Z. Phys. C 18 (1983) 69; A. Djouadi, M. Spira and P. M. Zerwas, Phys. Lett. B 264 (1991) 440; M. Spira, A. Djouadi, D. Graudenz and P. M. Zerwas, Nucl. Phys. B 453 (1995) 17 [arXiv:hep-ph/9504378].

- [13] A. D edes and S. M oretti, *Phys. Rev. Lett.* 84 (2000) 22; *Nucl. Phys. B* 576 (2000) 29; S. Y . Choi and J. S. Lee, *Phys. Rev. D* 61 (2000) 115002; S. Y . Choi, K . Hagiwara and J. S. Lee, *Phys. Lett. B* 529 (2002) 212; A. A rhib, D . K . Ghosh and O . C . Kong, *Phys. Lett. B* 537 (2002) 217; E . Christova, H . Eberl, W . M a jorotto and S. K ram l, *Nucl. Phys. B* 639 (2002) 263; *JHEP* 0212 (2002) 021; W . K hater and P. O sland, *Nucl. Phys. B* 661 (2003) 209.
- [14] J. F. G union and H . E . Haber, *Phys. Rev. D* 48 (1993) 5109; D . L. B orden, D . A . Bauer and D . O . Caldwell, *Phys. Rev. D* 48 (1993) 4018; B . G rzadkowski and J. F. G union, *Phys. Lett. B* 294 (1992) 361 [[arXiv:hep-ph/9206262](#)]; M . K ram er, J. H . K uhn, M . L. Stong and P. M . Zerwas, *Z. Phys. C* 64 (1994) 21 [[arXiv:hep-ph/9404280](#)]; G . J. G ounaris and G . P. T sirigoti, *Phys. Rev. D* 56 (1997) 3030 [Erratum -*ibid.* D 58 (1998) 059901] [[arXiv:hep-ph/9703446](#)]; I. F. G inzburg, G . L. K otkin, S. L. Pan l, V. G . Serbo and V . I. Tehov, *Nucl. Instrum . Meth. A* 219 (1984) 5; B . Badelek et al. [ECFA/DESY Photon Collider Working Group], *Int. J. Mod. Phys. A* 19 (2004) 5097 [[arXiv:hep-ex/0108012](#)].
- [15] S. Y . Choi and J. S. Lee, *Phys. Rev. D* 62 (2000) 036005; E . A sakawa, S. Y . Choi, K . Hagiwara and J. S. Lee, *Phys. Rev. D* 62 (2000) 115005; J. S. Lee, [hep-ph/0106327](#); S. Y . Choi, B . C . Chung, P. K o and J. S. Lee, *Phys. Rev. D* 66 (2002) 016009; R . M . Godbole, S. D . R indani and R . K . Singh, *Phys. Rev. D* 67 (2003) 095009; E . A sakawa and K . Hagiwara, *Eur. Phys. J. C* 31 (2003) 351 [[arXiv:hep-ph/0305323](#)]; S. Y . Choi, J. K alinowski, Y . Liao and P. M . Zerwas, *Eur. Phys. J. C* 40 (2005) 555 [[arXiv:hep-ph/0407347](#)]; B . G rzadkowski, Z . H ioki, K . O hkum a and J. W udka, *JHEP* 0511 (2005) 029 [[arXiv:hep-ph/0508183](#)]; R . M . Godbole, S. K ram l, S. D . R indani and R . K . Singh, *Phys. Rev. D* 74 (2006) 095006 [Erratum -*ibid.* D 74 (2006) 119901] [[arXiv:hep-ph/0609113](#)].
- [16] J. R . Ellis, J. S. Lee and A . Pilaftsis, *Nucl. Phys. B* 718 (2005) 247 [[arXiv:hep-ph/0411379](#)]; J. S. Lee, *Mod. Phys. Lett. A* 22 (2007) 1191 [[arXiv:0705.1089](#) [[hep-ph](#)]].
- [17] M . Spira, A . D juadi, D . G raudenz and P. M . Zerwas, *Phys. Lett. B* 318 (1993) 347; M . Spira, A . D juadi, D . G raudenz and P. M . Zerwas in Ref. [12]; M . K ram er, E . Laenen and M . Spira, *Nucl. Phys. B* 511 (1998) 523 [[arXiv:hep-ph/9611272](#)]; A . D juadi, M . Spira and P. M . Zerwas, *Phys. Lett. B* 264 (1991) 440; S. D awson, *Nucl. Phys. B* 359 (1991) 283; D . G raudenz, M . Spira and P. M . Zerwas, *Phys. Rev. Lett.* 70 (1993) 1372; R . P. K au man and W . Scha er, *Phys. Rev. D* 49 (1994) 551 [[arXiv:hep-ph/9305279](#)]; S. D awson and R . K au man, *Phys. Rev. D* 49 (1994) 2298 [[arXiv:hep-ph/9310281](#)].

- [18] S. Dawson, A. Djuradi and M. Spira, *Phys. Rev. Lett.* 77 (1996) 16 [arXiv:hep-ph/9603423]; M. Muhlleitner and M. Spira, *Nucl. Phys. B* 790 (2008) 1 [arXiv:hep-ph/0612254].
- [19] J. Guasch, W. Hollik and S. Penaranda, *Phys. Lett. B* 515 (2001) 367 [arXiv:hep-ph/0106027].
- [20] W. Y. Keung and W. J. Marciano, *Phys. Rev. D* 30 (1984) 248; A. Djuradi, J. Kalinowski and P. M. Zerwas, *Z. Phys. C* 70 (1996) 435 [arXiv:hep-ph/9511342]; S. Moretti and W. J. Stirling, *Phys. Lett. B* 347 (1995) 291 [Erratum -*ibid.* B 366 (1996) 451] [arXiv:hep-ph/9412209]; R. Decker, M. Nowakowski and A. Pilaftsis, *Z. Phys. C* 57 (1993) 339 [arXiv:hep-ph/9301283].
- [21] I. B. Khriplovich and S. K. Lamoreaux, *CP Violation Without Strangeness* (Springer, New York, 1997).
- [22] For a recent review, see, M. Pospelov and A. Ritz, *Annals Phys.* 318 (2005) 119.
- [23] For two-loop Higgs-mediated contributions to EDMs in the CP-violating MSSM, see D. Chang, W. Y. Keung and A. Pilaftsis, *Phys. Rev. Lett.* 82 (1999) 900; A. Pilaftsis, *Nucl. Phys. B* 644 (2002) 263; D. A. Demir, O. Lebedev, K. A. Olive, M. Pospelov and A. Ritz, *Nucl. Phys. B* 680 (2004) 339; K. A. Olive, M. Pospelov, A. Ritz and Y. Santoso, *Phys. Rev. D* 72 (2005) 075001 [arXiv:hep-ph/0506106].
- [24] J. R. Ellis, J. S. Lee and A. Pilaftsis, *Phys. Rev. D* 72 (2005) 095006 [arXiv:hep-ph/0507046].
- [25] J. S. Lee and S. Scopel, *Phys. Rev. D* 75 (2007) 075001 [arXiv:hep-ph/0701221]; J. S. Lee, arXiv:0706.2222 [hep-ph].
- [26] B. C. Regan, E. D. Commins, C. J. Schmidt and D. DeMille, *Phys. Rev. Lett.* 88 (2002) 071805.
- [27] G. Abbiendi et al. [OPAL Collaboration], *Eur. Phys. J. C* 37 (2004) 49 [arXiv:hep-ex/0406057]; OPAL Physics Note PN505, <http://opal.web.cern.ch/Opal/pubs/physnote/html/pn505.html>; A. Heister et al. [ALEPH Collaboration], *Phys. Lett. B* 526 (2002) 191 [arXiv:hep-ex/0201014]; J. Abdallah et al. [DELPHI Collaboration], *Eur. Phys. J. C* 32 (2004) 145 [arXiv:hep-ex/0303013]; P. Achard et al. [L3 Collaboration], *Phys. Lett. B* 545 (2002) 30 [arXiv:hep-ex/0208042]; S. Schael et al. [ALEPH Collaboration], *Eur. Phys. J. C* 47 (2006) 547 [arXiv:hep-ex/0602042].
- [28] P. Franzini et al., *Phys. Rev. D* 35, 2883 (1987).

- [29] W . M . Yao et al. [Particle Data Group], *J. Phys. G* 33 (2006) 1.
- [30] A . D edes and A . P ilafts is, *Phys. Rev. D* 67 (2003) 015012 [arX iv:hep-ph/0209306].
- [31] T . A altonen et al. [CDF Collaboration], *Phys. Rev. Lett.* 100 (2008) 101802 [arX iv:0712.1708 [hep-ex]].
- [32] E . Barberio et al. [Heavy Flavor Averaging Group (HFAG) Collaboration], arX iv:0704.3575 [hep-ex].
- [33] M . S . Carena, D . G arcia, U . N ierste and C . E . M . W agner, *Phys. Lett. B* 499, 141 (2001) [arX iv:hep-ph/0010003]. G . D egrassi, P . G ambino and G . F . G iudice, *JHEP* 0012, 009 (2000) [arX iv:hep-ph/0009337].
- [34] F . Borzum ati, J . S . Lee and W . Y . Song, *Phys. Lett. B* 595 (2004) 347. [arX iv:hep-ph/0401024].
- [35] K . Ikado et al., *Phys. Rev. Lett.* 97 (2006) 251802. [arX iv:hep-ex/0604018]; B . A ubert [The BABAR Collaboration], arX iv:0708.2260 [hep-ex].
- [36] B . A ubert et al. [BABAR Collaboration], *Phys. Rev. Lett.* 96 (2006) 241802 [arX iv:hep-ex/0511015].
- [37] H . G . Evans [CDF Collaboration], arX iv:0705.4598 [hep-ex].
- [38] M . M isiak et al., *Phys. Rev. Lett.* 98 (2007) 022002, [arX iv:hep-ph/0609232].

Table 1: The contents of the array RAUX_H. In RAUX_H(22) and RAUX_H(23), the notation h_f^0 is for the Yukawa couplings without including the threshold corrections. The notations which are not explained in the text follow the conventions of CPsuperH [3] and Refs. [5{7].

RAUX_H(1)	m_b^{pole}	RAUX_H(26)	$\hbar_t(Q_{tb})j$	RAUX_H(120)	d^H	10^{24} ecm
RAUX_H(2)	$m_b(m_b^{\text{pole}})$	RAUX_H(27)	$\hbar_b(m_t^{\text{pole}})j$	
RAUX_H(3)	$s(m_b^{\text{pole}})$	RAUX_H(28)	$\hbar_b(Q_b)j$	
RAUX_H(4)	m_c^{pole}	RAUX_H(29)	$\hbar_b(Q_{tb})j$	
RAUX_H(5)	$m_c(m_c^{\text{pole}})$	RAUX_H(30)	M_A^2	
RAUX_H(6)	$s(m_c^{\text{pole}})$	RAUX_H(31)	$\langle e_{H+H}^b (M_H^{\text{pole}2}) \rangle$	
...	...	RAUX_H(32)	$4v^2(m_t^{\text{pole}})=2$	
...	...	RAUX_H(33)	$4(m_t^{\text{pole}})$	RAUX_H(130)	$B(B_s!)$	10^7
...	...	RAUX_H(34)	$1(m_t^{\text{pole}})$	RAUX_H(131)	$B(B_d!)$	10^7
RAUX_H(10)	M_H^{pole} or M_H^e :	RAUX_H(35)	$2(m_t^{\text{pole}})$	RAUX_H(132)	$M_{B_d}^{\text{SU SY}}$	ps^1
RAUX_H(11)	Q_t^2	RAUX_H(36)	$34(m_t^{\text{pole}})$	RAUX_H(133)	$M_{B_s}^{\text{SU SY}}$	ps^1
RAUX_H(12)	Q_b^2	RAUX_H(134)	R_B	
RAUX_H(13)	Q_{tb}^2	RAUX_H(135)	$B(B! X_s)$	10^4
RAUX_H(14)	$v_1(m_t^{\text{pole}})$	RAUX_H(101)	$\frac{p}{s}$	RAUX_H(136)	$A_{CP}(B! X_s)\%$	
RAUX_H(15)	$v_1(Q_t)$	
RAUX_H(16)	$v_1(Q_b)$	
RAUX_H(17)	$v_1(Q_{tb})$	RAUX_H(111)	d_{T1}^H	
RAUX_H(18)	$v_2(m_t^{\text{pole}})$	RAUX_H(112)	$(d_{T1}^H)^e$			10^{24} ecm
RAUX_H(19)	$v_2(Q_t)$	RAUX_H(113)	$(d_{T1}^H)^{C_s}$			10^{24} ecm
RAUX_H(20)	$v_2(Q_b)$	RAUX_H(114)	d_e^H			10^{26} ecm
RAUX_H(21)	$v_2(Q_{tb})$	RAUX_H(115)	$(d_e^H)^t$			10^{26} ecm
RAUX_H(22)	$\hbar_t^0(m_t^{\text{pole}})j$	RAUX_H(116)	$(d_e^H)^b$			10^{26} ecm
RAUX_H(23)	$\hbar_b^0(m_t^{\text{pole}})j$	RAUX_H(117)	$(d_e^H)^t$			10^{26} ecm
RAUX_H(24)	$\hbar_t(m_t^{\text{pole}})j$	RAUX_H(118)	$(d_e^H)^b$			10^{26} ecm
RAUX_H(25)	$\hbar_t(Q_t)j$	RAUX_H(119)	$(d_e^H)^{\sim}$			10^{26} ecm

Table 2: The contents of the array CAUX_H. The notations which are not explained in the text follow the CPsuperH [3] convention.

CAUX_H(1)	$h_t = h_{tj}$	CAUX_H(112)	$D_{4;1}^{H^0}(\hat{s})$	CAUX_H(140)	$S_1^g(\frac{p_-}{\hat{s}})$
CAUX_H(2)	$h_b = h_{bj}$	CAUX_H(113)	$D_{4;2}^{H^0}(\hat{s})$	CAUX_H(141)	$P_1^g(\frac{p_-}{\hat{s}})$
...	...	CAUX_H(114)	$D_{4;3}^{H^0}(\hat{s})$	CAUX_H(142)	$S_2^g(\frac{p_-}{\hat{s}})$
...	...	CAUX_H(115)	$D_{4;4}^{H^0}(\hat{s})$	CAUX_H(143)	$P_2^g(\frac{p_-}{\hat{s}})$
...	...	CAUX_H(116)	$D_H^{H^0}(\hat{s})$	CAUX_H(144)	$S_3^g(\frac{p_-}{\hat{s}})$
...	...	CAUX_H(117)	$D_H^{H^0}(\hat{s})$	CAUX_H(145)	$P_3^g(\frac{p_-}{\hat{s}})$
CAUX_H(100)	$D_{1;1}^{H^0}(\hat{s})$	CAUX_H(118)	$D_G^{H^0}(\hat{s})$
CAUX_H(101)	$D_{1;2}^{H^0}(\hat{s})$	CAUX_H(119)	$D_G^{H^0}(\hat{s})$
CAUX_H(102)	$D_{1;3}^{H^0}(\hat{s})$	CAUX_H(150)	$h_b^0 j H_e^{B=2} \mathcal{B}_d^0 i_{SU SY}$
CAUX_H(103)	$D_{1;4}^{H^0}(\hat{s})$	CAUX_H(151)	$h_b^0 j H_e^{B=2} \mathcal{B}_s^0 i_{SU SY}$
CAUX_H(104)	$D_{2;1}^{H^0}(\hat{s})$
CAUX_H(105)	$D_{2;2}^{H^0}(\hat{s})$	CAUX_H(130)	$S_1(\frac{p_-}{\hat{s}})$
CAUX_H(106)	$D_{2;3}^{H^0}(\hat{s})$	CAUX_H(131)	$P_1(\frac{p_-}{\hat{s}})$
CAUX_H(107)	$D_{2;4}^{H^0}(\hat{s})$	CAUX_H(132)	$S_2(\frac{p_-}{\hat{s}})$
CAUX_H(108)	$D_{3;1}^{H^0}(\hat{s})$	CAUX_H(133)	$P_2(\frac{p_-}{\hat{s}})$
CAUX_H(109)	$D_{3;2}^{H^0}(\hat{s})$	CAUX_H(134)	$S_3(\frac{p_-}{\hat{s}})$
CAUX_H(110)	$D_{3;3}^{H^0}(\hat{s})$	CAUX_H(135)	$P_3(\frac{p_-}{\hat{s}})$
CAUX_H(111)	$D_{3;4}^{H^0}(\hat{s})$

Table 3: The contents of the extended SMPARA_H(IP).

IP	Parameter	IP	Parameter	IP	Parameter	IP	Parameter
1	$m_{em}^1(M_Z)$	6	m	11	$m_u(m_t^{pole})$	16	A
2	$m_s(M_Z)$	7	m	12	$m_c(m_t^{pole})$	17	
3	M_Z	8	$m_d(m_t^{pole})$	13	m_t^{pole}	18	
4	$\sin^2 w$	9	$m_s(m_t^{pole})$	14	w	19	
5	m_e	10	$m_b(m_t^{pole})$	15	z	20	
							...




Figure 5: The branching ratio $B(B_s \rightarrow \mu\mu) \times 10^7$ as a function of Φ_3 for four values of $\tan\beta$: $\tan\beta = 10$ (upper left), 20 (upper right), 30 (lower left), and 40 (lower right). The CPX scenario is taken with $M_{SUSY} = 0.5 \text{ TeV}$ and $M_H = 200 \text{ GeV}$ in the convention $\arg(\mu) = 0$. In each frame, the lower three lines are for the case $\rho = 1$ and the upper lines for $\rho = 10$ where the solid, dashed, and dash-dotted lines are for $\arg(\mu) = 0, 90, \text{ and } 180$, respectively. The current 95% experimental upper bound, 5.8×10^{-8} [31], is also shown as a horizontal line in each frame.




Figure 6: The branching ratio $B(B_s \rightarrow \mu\mu) \times 10^7$ as a function of the common hierarchy factor $\rho = \frac{v_c}{v} = \frac{v_D}{v}$ for four values of $\tan\beta$: $\tan\beta = 10$ (upper left), 20 (upper right), 30 (lower left), and 40 (lower right). The CPX scenario is taken with $M_{\text{SUSY}} = 0.5$ TeV and $M_H = 200$ GeV in the convention $\arg(\mu) = 0$. In each frame, the solid line is for $(\Phi_3; \Phi_A) = (0; 180)$ and the dashed one for $(90; 90)$. The current 95% experimental upper bound, 5.8×10^{-8} [31], is also shown as a horizontal line in each frame.




Figure 7: The branching ratio $\mathcal{B}^{\mu\mu} = B(B_s \rightarrow \mu\mu) \times 10^7$ in the $(\tan\beta; M_{H_1})$ plane. The CPX scenario is taken with $\mu_A = \mu_3 = 90$ and $M_{\text{SUSY}} = 0.5$ TeV for two values of the common hierarchy factor: $\rho = 1$ (left) and 10 (right). The unshaded region is not theoretically allowed. The different shaded regions correspond to different ranges of $\mathcal{B}^{\mu\mu}$, as shown: specifically, $\mathcal{B}^{\mu\mu} < 0.58$ in the lowest (blue) low- $\tan\beta$ region, consistent with the current upper limit at 95 % C.L.




Figure 8: The branching ratio $\mathcal{B}_s^{\text{sg}} = B(B \rightarrow X_s \gamma) \times 10^4$ in the $(\tan\beta; M_{H_1})$ plane. The same CPX scenario with $\mu_A = \mu_3 = 90$ is taken as in Fig. 7. The different shaded regions correspond to different ranges of $\mathcal{B}_s^{\text{sg}}$, as shown: specifically, $3.03 < \mathcal{B}_s^{\text{sg}} \leq 4.07$ in the uppermost (blue) high- $\tan\beta$ region, consistent with the current experimentally allowed 2-region, $3.03 < \mathcal{B}_s^{\text{sg}} \leq 4.07$ [32].




Figure 9: The ratio R_{BTV} in the $(\tan \beta; M_{H_1})$ plane. The same CPX scenario with $\alpha_A = \alpha_3 = 90^\circ$ is taken as in Fig. 7 for two values of ρ : $\rho = 1$ (left) and $\rho = 10$ (right). The different shaded regions correspond to the regions allowed at the 1- and 2- levels by the recent BELLE and BABAR results: $R_B^{EXP} = 1:0.38$ [5,35]. In the right frame, specifically, the 2- excluded regions are shown as $R_{BTV} > 1.76$ (in the high-tan region) and $R_{BTV} \leq 0.24$ (in the middle-tan region).




Figure 10: The experimental constraints from $B(B_s \rightarrow s\gamma)$ (95%), $B(B \rightarrow X_s)$ (2%), and R_B (1) in the $(\tan \beta; M_{H_1})$ plane for two values of ρ . The same CPX scenario with $\alpha_A = \alpha_3 = 90^\circ$ is taken as in Fig. 7.




Figure 11: The region allowed experimentally by the measurement $\mathbf{B}(\mathbf{B}_d \rightarrow \tau^+ \tau^-) < 4.1 \times 10^{-3}$ (90%) [36] (upper frames) and the regions where the SUSY contribution is smaller than the measured values of $B_s^0 - B_s^0$ mass difference [37] (middle frames) and $B_d^0 - B_d^0$ mass difference [29] (lower frames), in the $(\tan \beta; M_{H_1})$ plane. The left three frames are for $\rho = 1$ and the right ones for $\rho = 10$. The same CPX scenario with $\alpha_A = \alpha_3 = 90^\circ$ is taken as in Fig. 7.

Table 4: The contents of the extended SSPARA_H(IP).

IP	Parameter	IP	Parameter	IP	Parameter	IP	Parameter
1	\tan	8	\mathcal{M}_2	15	m_{E_3}	22	\mathcal{Q}
2	M_H^{pole}	9	\mathcal{M}_{3j}	16	\mathcal{A}_{tj}	23	\mathcal{U}
3	jj	10	\mathcal{M}_3	17	\mathcal{A}_t	24	\mathcal{D}
4		11	$m_{\mathcal{Q}_3}$	18	\mathcal{A}_{bj}	25	\mathcal{L}
5	\mathcal{M}_{1j}	12	$m_{\mathcal{U}_3}$	19	\mathcal{A}_b	26	\mathcal{E}
6	\mathcal{M}_1	13	$m_{\mathcal{D}_3}$	20	\mathcal{A}_j	27	...
7	\mathcal{M}_{2j}	14	$m_{\mathcal{L}_3}$	21	\mathcal{A}	28	...