

(Total) Vector Domination for Graphs with Bounded Branchwidth

Toshimasa Ishii¹, Hirotaka Ono², and Yushi Uno³

¹ Graduate School of Economics and Business Administration, Hokkaido University,
Sapporo 060-0809, Japan

² Department of Economic Engineering, Faculty of Economics, Kyushu University,
Fukuoka 812-8581, Japan

³ Department of Mathematics and Information Sciences, Graduate School of Science,
Osaka Prefecture University, Sakai 599-8531, Japan

Abstract. Given a graph $G = (V, E)$ of order n and an n -dimensional non-negative vector $d = (d(1), d(2), \dots, d(n))$, called demand vector, the vector domination (resp., total vector domination) is the problem of finding a minimum $S \subseteq V$ such that every vertex v in $V \setminus S$ (resp., in V) has at least $d(v)$ neighbors in S . The (total) vector domination is a generalization of many dominating set type problems, e.g., the dominating set problem, the k -tuple dominating set problem (this k is different from the solution size), and so on, and its approximability and inapproximability have been studied under this general framework. In this paper, we show that a (total) vector domination of graphs with bounded branchwidth can be solved in polynomial time. This implies that the problem is polynomially solvable also for graphs with bounded treewidth. Consequently, the (total) vector domination problem for a planar graph is subexponential fixed-parameter tractable with respect to k , where k is the size of solution.

1 Introduction

Given a graph $G = (V, E)$ of order n and an n -dimensional non-negative vector $d = (d(1), d(2), \dots, d(n))$, called *demand vector*, the *vector domination* (resp., *total vector domination*) is the problem of finding a minimum $S \subseteq V$ such that every vertex v in $V \setminus S$ (resp., in V) has at least $d(v)$ neighbors in S . These problems were introduced by [19], and they contain many existing problems, such as the minimum dominating set and the k -tuple dominating set problem (this k is different from the solution size) [20,21], and so on. Indeed, by setting $d = (1, \dots, 1)$, the vector domination becomes the minimum dominating set forms, and by setting $d = (k, \dots, k)$, the total vector dominating set becomes the k -tuple dominating set. If in the definition of total vector domination, we replace open neighborhoods with closed ones, we get the *multiple domination*. In this paper, we sometimes refer to these problems just as *domination problems*. Table 1 of [9] summarizes how related problems are represented in the scheme of domination problems. Many variants of the basic concepts of domination and their applications have appeared in [21,22].

Since the vector or multiple domination includes the setting of the ordinary dominating set problem, it is obviously NP-hard, and further it is NP-hard to approximate within $(c \log n)$ -factor, where c is a positive constant, e.g., 0.2267 [1,24]. As for the approximability, since the domination problems are special cases of a set-cover type integer problem, it is known that the polynomial-time greedy algorithm achieves an $O(\log n)$ -approximation factor [15]; it is already optimal in terms of order. We can see further analyses of the approximability and inapproximability in [8,9].

In this paper, we focus on another aspect of designing algorithms for domination problems, that is, the polynomial-time solvability of the domination problems for graphs of bounded treewidth or branchwidth. In [3], it is shown that the vector domination problem is $W[1]$ -hard with respect to treewidth. This result and Courcelle’s meta-theorem about MSOL [10] imply that the vector domination is unlikely expressible in MSOL; it is not obvious to obtain a polynomial time algorithm.

In this paper, we present a polynomial-time algorithm for the domination problems of graphs with bounded branchwidth. The branchwidth is a measure of the “global connectivity” of a graph, and is known to be a counterpart of treewidth. It is known that

$$\max\{bw(G), 2\} \leq tw(G) + 1 \leq \max\{3bw(G)/2, 2\},$$

where $bw(G)$ and $tw(G)$ denote the branchwidth and treewidth of graph G , respectively [26]. Due to the linear relation of these two measures, polynomial-time solvability of a problem for graphs with bounded treewidth implies polynomial-time solvability of a problem for graphs with bounded branchwidth, and vice versa. Hence, our results imply that the domination problems (i.e., vector domination, total vector domination and multiple domination) can be solved in polynomial time for graphs with bounded treewidth; the polynomial-time solvability for all the problems (except the dominating set problem) in Table 1 of [9] is newly shown. Also, they answer the question by [8,9] about the complexity status of the domination problems of graphs with bounded treewidth.

Furthermore, by using the polynomial-time algorithms for graphs of bounded treewidth, we can show that these problems for a planar graph are subexponential fixed-parameter tractable with respect to the size k of the solution, that is, there is an algorithm whose running time is $2^{O(\sqrt{k} \log k)} n^{O(1)}$. To our best knowledge, these are the first fixed-parameter algorithms for the total vector domination and multiple domination, whereas the vector domination for planar graphs has been shown to be FPT [25]. For the latter case, our algorithm greatly improves the running time.

Note that the polynomial-time solvability of the vector domination problem for graphs of bounded treewidth has been independently shown very recently [7]. They considered a further generalization of the vector domination problem, and gave a polynomial-time algorithm for graphs of bounded clique-width. Since $cw(G) \leq 2^{tw(G)+1} + 1$ holds where $cw(G)$ denotes the clique-width of graph G ([11]), their polynomial-time algorithm implies the polynomial-time solvability

of the vector domination problem for graphs of bounded treewidth and bounded branchwidth.

1.1 Related Work

The dominating set problem itself is one of the most fundamental graph optimization problems, and it has been intensively and extensively studied from many points of view. In the sense that the vector or multiple domination contains the setting of not only the ordinary dominating set problem but also many variants, there are an enormous number of related studies. Here we pick some representatives up.

As a research of the domination problems from the viewpoint of the algorithm design, Cicalese, Milanic and Vaccaro gave detailed analyses of the approximability and inapproximability [8,9]. They also provided some exact polynomial-time algorithms for special classes of graphs, such as complete graphs, trees, P_4 -free graphs, and threshold graphs.

For graphs with bounded treewidth (or branchwidth), the ordinary domination problems can be solved in polynomial time. As for the fixed-parameter tractability, it is known that even the ordinary dominating set problem is W[2]-complete with respect to solution size k ; it is unlikely to be fixed-parameter tractable [16]. In contrast, it can be solved in $O(2^{15.13\sqrt{k}} + n^3)$ time for planar graphs, that is, it is subexponential fixed-parameter tractable [18]. The subexponent part comes from the inequality $bw(G) \leq 12\sqrt{k} + 9$, where k is the size of a dominating set of G . Behind the inequality, there is a unified property of parameters, called *bidimensionality* [14]. Namely, the subexponential fixed-parameter algorithm of the dominating set for planar graphs (more precisely, H -minor-free graphs [13]) is based on the bidimensionality.

A maximization version of the ordinary dominating set is also considered. *Partial Dominating Set* is the problem of maximizing the number of vertices to be dominated by using a given number k of vertices. In [2], it was shown that partial dominating set problem is FPT with respect to k for H -minor-free graphs. Later, [17] gives a subexponential FPT with respect to k for apex-minor-free graphs, also a super class of planar graphs. Although partial dominating set is an example of problems to which the bidimensionality theory cannot be applied, they develop a technique to reduce an input graph so that its treewidth becomes $O(\sqrt{k})$.

For the vector domination, a polynomial-time algorithm for graphs of bounded treewidth has been proposed very recently [7], as mentioned before. In [25], it is shown that the vector domination for ρ -degenerated graphs can be solved in $k^{O(\rho k^2)} n^{O(1)}$ time, if $d(v) > 0$ holds for $\forall v \in V$ (positive constraint). Since any planar graph is 5-degenerated, the vector domination for planar graphs is fixed-parameter tractable with respect to solution size, under the positive constraint. Furthermore, the case where $d(v)$ could be 0 for some v can be easily reduced to the positive case by using the transformation discussed in [3], with increasing the degeneracy only 1. It follows that the vector domination for planar graphs

is FPT with respect to solution size k . However, for the total vector domination and multiple domination, neither polynomial time algorithm for graphs of bounded treewidth nor fixed-parameter algorithm for planar graphs has been known.

Other than these, several generalized versions of the dominating set problem are also studied. (k, r) -center problem is the one that asks the existence of set S of k vertices satisfying that for every vertex $v \in V$ there exists a vertex $u \in S$ such that the distance between u and v is at most r ; $(k, 1)$ -center corresponds to the ordinary dominating set. The (k, r) -center for planar graphs is shown to be fixed-parameter tractable with respect to k and r [12]. For $\sigma, \rho \subseteq \{0, 1, 2, \dots\}$ and a positive integer k , $\exists[\sigma, \rho]$ -dominating set is the problem that asks the existence of set S of k vertices satisfying that $|N(v) \cap S| \in \sigma$ holds for $\forall v \in S$ and $|N(v) \cap S| \in \rho$ for $\forall v \in V \setminus S$, where $N(v)$ denotes the open neighborhood of v . If $\sigma = \{0, 1, \dots\}$ and $\rho = \{1, 2, \dots\}$, $\exists[\sigma, \rho]$ -dominating set is the ordinary dominating set problem, and if $\sigma = \{0\}$ and $\rho = \{0, 1, 2, \dots\}$, it is the independent set. In [6], the parameterized complexity of $\exists[\sigma, \rho]$ -dominating set with respect to treewidth is also considered.

1.2 Our Results

Our results are summarized as follows:

- We present a polynomial-time algorithm for the vector domination of graph $G = (V, E)$ with bounded branchwidth. The running time is roughly $O(n^{6bw(G)+2})$.
- We present polynomial-time algorithms for the total vector domination and multiple domination of graph G with bounded branchwidth. The running time is roughly $O(2^{9bw(G)/2} n^{6bw(G)+2})$.
- Let G be a planar graph. Then, we can check in $O(n^4 + \min\{k, d^*\}^{40\sqrt{k}+34}n)$ time whether G has a vector dominating set with cardinality at most k or not, where $d^* = \max\{d(v) \mid v \in V\}$.
- Let G be a planar graph. Then, we can check in $O(n^4 + 2^{30\sqrt{k}+51/2} \min\{k, d^*\}^{40\sqrt{k}+34}n)$ time whether G has a total vector dominating set and a multiple dominating set with cardinality at most k or not.

It should be noted that it is actually possible to design directly polynomial time algorithms for graphs with bounded treewidth, but they are slower than the ones for graphs with bounded branchwidth; this is the reason why we adopt the branchwidth instead of the treewidth.

As far as the authors know, the second and fourth results give the first polynomial time algorithms and the first fixed-parameter algorithm for the total vector domination and multiple domination of graphs with bounded branchwidth (or treewidth) and planar graphs, respectively. As for the vector domination, we give an $O(n^{6bw(G)+2})$ -time algorithm, whose running time is $O(n^{6(tw(G)+1)+2})$ in terms of the treewidth, whereas the recent paper [7] gives an $O(cw(G)|\sigma|(n+1)^{5cw(G)})$ -time algorithm, where $|\sigma|$ is the encoding length of k -expression used in the algorithm, and is bounded by a polynomial in the input size for fixed k .

Since $cw(G) \leq 2^{tw(G)+1} + 1$ holds, this is an $O(2^{tw(G)+1}|\sigma|(n+1)^{5 \cdot 2^{tw(G)+1}})$ -time algorithm.

Also, the third result shows that the vector domination of planar graphs is subexponential FPT with respect to k , and it greatly improves the running time of existing $k^{O(k^2)}n^{O(1)}$ -time algorithm ([25]). It was shown in [5] that for the ordinary dominating set problem (equivalently, the vector domination (or multiple domination) with $d = (1, 1, \dots, 1)$) in planar graphs, there is no $2^{o(\sqrt{k})}n^{O(1)}$ -time algorithm unless the Exponential Time Hypothesis (i.e., the assumption that there is no $2^{o(n)}$ -time algorithm for n -variable 3SAT [23]) fails. Hence, in this sense, our algorithm in third result (or the fourth results for the multiple domination) is optimal if d^* is a constant.

The third and fourth results give subexponential fixed-parameter algorithms of the domination problems for planar graphs. It should be noted that the domination problems themselves do not have the bidimensionality, mentioned in the previous subsection, due to the existence of the vertices with demand 0. Instead, by reducing irrelevant vertices, we obtain a similar inequality about the branchwidth and the solution size of the domination problems, which leads to the subexponential fixed-parameter algorithms.

The remainder of the paper is organized as follows. In Section 2, we introduce some basic notations and then explain the branch decomposition. Section 3 is the main part of the paper, and presents our dynamic programming based algorithms for the considered problems. Section 4 explains how we extend the algorithms of Section 3 to fixed-parameter algorithms for planar graphs.

2 Preliminaries

A graph G is an ordered pair of its vertex set $V(G)$ and edge set $E(G)$ and is denoted by $G = (V(G), E(G))$. Let $n = |V(G)|$ and $m = |E(G)|$. We assume throughout this paper that all graphs are undirected, and simple, unless otherwise stated. Therefore, an edge $e \in E(G)$ is an unordered pair of vertices u and v , and we often denote it by $e = (u, v)$. Two vertices u and v are *adjacent* if $(u, v) \in E(G)$. For a graph G , the (*open*) *neighborhood* of a vertex $v \in V(G)$ is the set $N_G(v) = \{u \in V(G) \mid (u, v) \in E(G)\}$, and the *closed neighborhood* of v is the set $N_G[v] = N_G(v) \cup \{v\}$.

For a graph $G = (V, E)$, let $d = (d(v) \mid v \in V)$ be an n -dimensional non-negative vector. Then, we call a set $S \subseteq V$ of vertices a *d -vector dominating set* (resp., *d -total vector dominating set*) if $|N_G(v) \cap S| \geq d(v)$ holds for every vertex $v \in V - S$ (resp., $v \in V$). We call a set $S \subseteq V$ of vertices a *d -multiple dominating set* if $|N_G[v] \cap S| \geq d(v)$ holds for every vertex $v \in V$. We may drop d in these notations if there are no confusions.

2.1 Branch decomposition

A *branch decomposition* of a graph $G = (V, E)$ is defined as a pair $(T = (V_T, E_T), \tau)$ such that (a) T is a tree with $|E|$ leaves in which every non-leaf

node has degree 3, and (b) τ is a bijection from E to the set of leaves of T . Throughout the paper, we shall use *node* to denote an element in V_T for distinguishing it from an element in V .

For an edge f in T , let T_f and $T - T_f$ be two trees obtained from T by removing f , and E_f and $E - E_f$ be two sets of edges in E such that $e \in E_f$ if and only if $\tau(e)$ is included in T_f . The *order function* $w : E(T) \rightarrow 2^V$ is defined as follows: for an edge f in T , a vertex $v \in V$ belongs to $w(f)$ if and only if there exist an edge in E_f and an edge in $E - E_f$ which are both incident to v . The *width* of a branch decomposition (T, τ) is $\max\{|w(f)| \mid f \in E_T\}$, and the *branchwidth* of G , denoted by $bw(G)$, is the minimum width over all branch decompositions of G .

In general, computing the branchwidth of a given graph is NP-hard [28]. On the other hand, Bodlaender and Thilikos [4] gave a linear time algorithm which checks in linear time whether the branchwidth of a given graph is at most k or not, and if so, outputs a branch decomposition of minimum width, for any fixed k . Also, as shown in the following lemma, it is known that for planar graphs, it can be done in polynomial time for any given k , where a graph is called *planar* if it can be drawn in the plane without generating a crossing by two edges.

Lemma 1. ([28]) *Let G be a planar graph. Then, it can be checked in $O(n^2)$ time whether $bw(G) \leq k$ or not for a given integer k . Also, we can construct a branch decomposition of G with width $bw(G)$ in $O(n^4)$ time. \square*

Here, we introduce the following basic properties about branch decompositions, which will be utilized in the subsequent sections.

Lemma 2. *Let (T, τ) be a branch decomposition of G .*

(i) *For a tree T , let x be a non-leaf node and $f_i = (x, x_i)$, $i = 1, 2, 3$, be an edge incident to x (note that the degree of x is three). Then, $w(f_i) - w(f_j) - w(f_k) = \emptyset$ for every $\{i, j, k\} = \{1, 2, 3\}$. Hence, $w(f_i) \subseteq w(f_j) \cup w(f_k)$.*

(ii) *Let f be an edge of T , V_1 be the set of all end-vertices of edges in E_f , and V_2 be the set of all end-vertices of edges in $E - E_f$. Then, $(V_1 - w(f)) \cap (V_2 - w(f)) = \emptyset$ holds. Also, there is no edge in E connecting a vertex in $V_1 - w(f)$ and a vertex in $V_2 - w(f)$.*

Proof. (i) Without loss of generality, assume that $E_{f_1} \cap E_{f_2} = \emptyset$, $E_{f_2} \cap E_{f_3} = \emptyset$, $E_{f_3} \cap E_{f_1} = \emptyset$, and $E_{f_1} \cup E_{f_2} \cup E_{f_3} = E$. Let $v \in w(f_1)$ be a vertex. From the definition of $w(f_1)$, there exist two edges $e \in E_{f_1}$ and $e' \in E - E_{f_1}$ such that both of e and e' are incident to v . If $e' \in E_{f_2}$ (resp., $e' \in E_{f_3}$), then $v \in w(f_2)$ (resp., $v \in w(f_3)$) also holds. Thus, we can observe that there is no vertex in $w(f_1) - w(f_2) - w(f_3)$ for every $\{i, j, k\} = \{1, 2, 3\}$.

(ii) Assume by contradiction that there exists a vertex $v \in (V_1 - w(f)) \cap (V_2 - w(f))$. From definition of V_1 and V_2 , then there exists an edge $e_1 \in E_f$ and an edge $e_2 \in E - E_f$ such that both of e_1 and e_2 are incident to v . From the existence of e_1 and e_2 and the definition of $w(f)$, it follows that $w(f)$ also contains v , which contradicts $v \notin w(f)$.

Assume by contradiction that there exists an edge $e = (u_1, u_2) \in E$ such that $u_1 \in V_1 - w(f)$ and $u_2 \in V_2 - w(f)$. If we assume that $e \in E_f$ without loss of generality, then $u_2 \in V_1 - w(f)$ also holds, which contradicts $(V_1 - w(f)) \cap (V_2 - w(f)) = \emptyset$. \square

3 Domination problems in graphs of bounded branchwidth

In this section, we propose dynamic programming algorithms for the vector domination problem, the total vector domination problem, and the multiple domination problem, by utilizing a branch decomposition of a given graph. The techniques are based on the one developed by Fomin and Thilikos for solving the dominating set problem with bounded branchwidth [18]. Throughout this section, for a given graph $G = (V, E)$, the demand of each vertex $v \in V$ is denoted by $d(v)$, and let $d^* = \max\{d(v) \mid v \in V\}$.

3.1 Vector domination

In this subsection, we consider the vector domination problem, and show the following theorem.

Theorem 1. *If a branch decomposition of G with width b is given, a minimum vector dominating set in G can be found in $O((d^* + 2)^b \{(d^* + 1)^2 + 1\}^{b/2} m)$ time.* \square

Due to the assumption of the above theorem, we need to consider how we obtain a branch decomposition of G for the completeness of an algorithm of the vector domination problem. For a branch decomposition, there exists an $O(2^{b \lg 27} n^2)$ -time algorithm that given a graph G , reports $bw(G) \geq b$, or outputs a branch decomposition of G with width at most $3b$ [27,13]. Thus, the time to find a branch decomposition with width at most $3bw(G)$ is $O(\log bw(G) 2^{3bw(G) \lg 27} n^2)$ (smaller than the time complexity below), and we have the following corollary.

Corollary 1. *A minimum vector dominating set in G can be found in $O((d^* + 2)^{3bw(G)} \{(d^* + 1)^2 + 1\}^{3bw(G)/2} n^2)$ time.* \square

Below, for proving this theorem, we will give a dynamic programming algorithm for finding a minimum vector dominating set in G in $O((d^* + 2)^b \{(d^* + 1)^2 + 1\}^{b/2} m)$ time, based on a branch decomposition of G .

Let (T', τ) be a branch decomposition of $G = (V, E)$ with b , and $w' : E(T') \rightarrow 2^V$ be the corresponding order function. Let T be the tree from T' by inserting two nodes r_1 and r_2 , deleting one arbitrarily chosen edge $(x_1, x_2) \in E(T')$, adding three new edges (r_1, r_2) , (x_1, r_2) , and (x_2, r_2) ; namely, $T = (V(T') \cup \{r_1, r_2\}, E(T') \cup \{(r_1, r_2), (x_1, r_2), (x_2, r_2)\} - \{(x_1, x_2)\})$. Here, we regard T with a rooted tree by choosing r_1 as a root. Let $w(f) = w'(f)$ for every $f \in E(T) \cap E(T')$, $w(x_1, r_2) = w(x_2, r_2) = w'(x_1, x_2)$, and $w(r_1, r_2) = \emptyset$.

Let $f = (y_1, y_2) \in E$ be an edge in T such that y_1 is the parent of y_2 . Let $T(y_2)$ be the subtree of T rooted at y_2 , $E_f = \{e \in E \mid \tau(e) \in V(T(y_2))\}$, and G_f be the subgraph of G induced by E_f . Note that $w(f) \subseteq V(G_f)$ holds, since each vertex in $w(f)$ is an end-vertex of some edge in E_f by definition of the order function w . In the following, each vertex $v \in w(f)$ will be assigned one of the following $d(v) + 2$ colors

$$\{\top, 0, 1, 2, \dots, d(v)\}.$$

The meaning of the color of a vertex v is as follows: for a vertex set (possibly, a vector dominating set) D ,

- \top means that $v \in D$.
- $i \in \{0, 1, \dots, d(v)\}$ means that $v \notin D$ and $|N_{G_f}(v) \cap D| \geq d(v) - i$.

Notice that a vertex colored by $i > 0$ may need to be dominated by some vertices in $V - V(G_f)$ for the feasibility. Given a coloring $c \in \{\top, 0, 1, 2, \dots, d^*\}^{|w(f)|}$, let $D_f(c) \subseteq V(G_f)$ be a vertex set with the minimum cardinality satisfying the following (1)–(3), where $c(v)$ denotes the color assigned to a vertex $v \in V$:

$$c(v) = \top \text{ if and only if } v \in D_f(c) \cap w(f). \quad (1)$$

$$\text{If } c(v) = i, \text{ then } v \in w(f) - D_f(c) \text{ and } |N_{G_f}(v) \cap D_f(c)| \geq d(v) - i. \quad (2)$$

$$|N_{G_f}(v) \cap D_f(c)| \geq d(v) \text{ holds for every vertex } v \in V(G_f) - w(f) - D_f(c). \quad (3)$$

Intuitively, $D_f(c)$ is a minimum vector dominating set in G_f under the assumption that the color for every vertex in $w(f)$ is restricted to c . Note that a vertex in $w(f)$ is allowed not to meet its demand in G_f , because it can be dominated by some vertices in $V - V(G_f)$. Also note that every vertex in $V(G_f) - w(f)$ is not adjacent to any vertex in $V - V(G_f)$ by Lemma 2(ii), and it needs to be dominated by vertices only in $V(G_f)$ for the feasibility. We define $A_f(c)$ as $A_f(c) = |D_f(c)|$ if $D_f(c)$ exists and $A_f(c) = \infty$ otherwise.

Our dynamic programming algorithm proceeds bottom-up in T , while computing $A_f(c)$ for all $c \in \{\top, 0, 1, 2, \dots, d^*\}^{|w(f)|}$ for each edge f in T . We remark that $A_{(r_1, r_2)}(c)$ is the cardinality of a minimum vector dominating set, because $w(r_1, r_2) = \emptyset$ and $G_{(r_1, r_2)} = G$. The algorithm consists of two types of procedures: one is for leaf edges and the other is for non-leaf edges, where a *leaf edge* denotes an edge incident to a leaf of T .

Procedure for leaf edges: In the first step of the algorithm, we compute $A_f(c)$ for each edge f incident to a leaf of T . Then, for all colorings $c \in \{\top, 0, 1, 2, \dots, d^*\}^{|w(f)|}$, let $A_f(c)$ be the number of vertices colored by \top if G_f and c satisfy (1) – (3), and $A_f(c) = \infty$ otherwise.

For a fixed c , we need to check if (1) – (3) hold. This can be done in $O(|w(f)|)$ time. Hence, this step takes $O((d^* + 2)^{|w(f)|} |w(f)|)$ time.

Procedure for non-leaf edges: After the above initialization step, we visit non-leaf edges of T from leaves to the root of T . Let $f = (y_1, y_2)$ be a non-leaf edge of T such that y_1 is the parent of y_2 , y_3 and y_4 are the children of y_2 , and

$f_1 = (y_2, y_3)$ and $f_2 = (y_2, y_4)$. Now we have already obtained $A_{f_j}(c')$ for all $c' \in \{\top, 0, 1, 2, \dots, d^*\}^{|w(f_j)|}$, $j = 1, 2$. By Lemma 2(i), we have $w(f) \subseteq w(f_1) \cup w(f_2)$, $w(f_1) \subseteq w(f_2) \cup w(f)$, and $w(f_2) \subseteq w(f) \cup w(f_1)$; let $X_1 = w(f) - w(f_2)$, $X_2 = w(f) - w(f_1)$, $X_3 = w(f) \cap w(f_1) \cap w(f_2)$, and $X_4 = w(f_1) - w(f)$ ($= w(f_2) - w(f)$).

We say that a coloring $c \in \{\top, 0, 1, 2, \dots, d^*\}^{|w(f)|}$ of $w(f)$ is *formed* from a coloring c_1 of $w(f_1)$ and a coloring c_2 of $w(f_2)$ if the following (P1)–(P5) hold.

(P1) For every $v \in X_1 \cup X_2 \cup X_3$ with $c(v) = \top$,

- (a) If $v \in X_1 \cup X_3$, then $c_1(v) = \top$.
- (b) If $v \in X_2 \cup X_3$, then $c_2(v) = \top$.

(P2) For every $v \in X_4$, $c_1(v) = \top$ if and only if $c_2(v) = \top$.

(P3) For every $v \in X_j - D_{c_1, c_2}$ where $\{j, j'\} = \{1, 2\}$ and $D_{c_1, c_2} = \{v \in X_1 \cup X_2 \cup X_3 \cup X_4 \mid c_1(v) = \top \text{ or } c_2(v) = \top\}$,

If $c(v) = i$, then $c_j(v) = \min\{d(v), i + |D_{c_1, c_2} \cap N_{G_f}(v) \cap X_{j'}|\}$.
(Intuitively, if $v \in X_j - D_{c_1, c_2}$ needs to be dominated by at least $d(v) - i$ vertices in G_f , then at least $\max\{0, d(v) - i - |D_{c_1, c_2} \cap N_{G_f}(v) \cap X_{j'}|\}$ vertices from $V(G_{f_j})$ are necessary.)

(P4) For every $v \in X_3 - D_{c_1, c_2}$,

If $c(v) = i$, then $c_1(v) = \min\{d(v), i + |D_{c_1, c_2} \cap N_{G_f}(v) \cap X_2| + i_1\}$ and $c_2(v) = \min\{d(v), i + |D_{c_1, c_2} \cap N_{G_f}(v) \cap X_1| + i_2\}$ for some nonnegative integers i_1, i_2 with $i_1 + i_2 = \max\{0, d(v) - i - |D_{c_1, c_2} \cap N_{G_f}(v)|\}$.
(Intuitively, if $v \in X_3 - D_{c_1, c_2}$ needs to be dominated by at least $d(v) - i$ vertices in G_f , then at least $\max\{0, d(v) - i - |D_{c_1, c_2} \cap N_{G_f}(v)|\}$ vertices from $(V(G_{f_1}) - w(f_1)) \cup (V(G_{f_2}) - w(f_2))$ are necessary for dominating v . If i_1 (resp., i_2) vertices among those vertices belong to $V(G_{f_2}) - w(f_2)$ (resp., $V(G_{f_1}) - w(f_1)$), then at least $\max\{0, d(v) - i - |D_{c_1, c_2} \cap N_{G_f}(v) \cap X_{j'}| - i_j\}$ vertices from $V(G_{f_j})$ are necessary for $\{j, j'\} = \{1, 2\}$.)

(P5) For every $v \in X_4 - D_{c_1, c_2}$,

$c_1(v) = \min\{d(v), |D_{c_1, c_2} \cap N_{G_f}(v) \cap X_2| + i_1\}$ and $c_2(v) = \min\{d(v), |D_{c_1, c_2} \cap N_{G_f}(v) \cap X_1| + i_2\}$ for some nonnegative integers i_1, i_2 with $i_1 + i_2 = \max\{0, d(v) - |D_{c_1, c_2} \cap N_{G_f}(v)|\}$.
(This case can be treated in a similar way to (P4).)

As we will show in Lemmas 3 and 4, there exist a coloring c_1 of $w(f_1)$ and a coloring c_2 of $w(f_2)$ forming c such that $D_{f_1}(c_1) \cup D_{f_2}(c_2)$ satisfies (1)–(3) and $|D_{f_1}(c_1) \cup D_{f_2}(c_2)| = A_f(c)$. Namely, we have

$$A_f(c) = \min\{|A_{f_1}(c_1)| + |A_{f_2}(c_2)| - |D_{c_1, c_2} \cap (X_3 \cup X_4)| \mid c_1, c_2 \text{ forms } c\}.$$

Thus, for all colorings $c \in \{\top, 0, 1, 2, \dots, d^*\}^{|w(f)|}$, we can compute $A_f(c)$ from the information of f_1 and f_2 . By repeating these procedure bottom-up in T , we can find a minimum vector dominating set in G .

Here, for a fixed c , we analyze the time complexity for computing $A_f(c)$. Let $D_c = \{v \in w(f) \mid c(v) = \top\}$, $x_j = |X_j|$ for $j = 1, 2, 3, 4$, $z_3 = |X_3 - D_c|$, and z_4 be the number of vertices in X_4 not colored by \top . The number of pairs of a coloring c_1 of $w(f_1)$ and a coloring c_2 of $w(f_2)$ forming c is at most

$$(d^* + 1)^{z_3} \sum_{z_4=0}^{x_4} \binom{x_4}{z_4} (d^* + 1)^{z_4} (d^* + 1)^{z_4}$$

since the number of pairs (i_1, i_2) in (P4) or (P5) is at most $d^* + 1$ for each vertex in $X_3 - D_c$ or each vertex in X_4 not colored by \top .

Hence, for an edge f , the number of pairs forming c is at most

$$\begin{aligned} & (d^* + 2)^{x_1+x_2} \sum_{z_3=0}^{x_3} \binom{x_3}{z_3} (d^* + 1)^{z_3} (d^* + 1)^{z_3} \sum_{z_4=0}^{x_4} \binom{x_4}{z_4} (d^* + 1)^{z_4} (d^* + 1)^{z_4} \\ & = (d^* + 2)^{x_1+x_2} \{(d^* + 1)^2 + 1\}^{x_3+x_4} \end{aligned}$$

in total. Now we have $x_1 + x_2 + x_3 \leq b$, $x_1 + x_3 + x_4 \leq b$, and $x_2 + x_3 + x_4 \leq b$ (recall that b is the width of (T', τ)). By considering a linear programming problem which maximizes $(x_1 + x_2) \log(d^* + 2) + (x_3 + x_4) \log\{(d^* + 1)^2 + 1\}$ subject to these inequalities, we can observe that $(d^* + 2)^{x_1+x_2} \{(d^* + 1)^2 + 1\}^{x_3+x_4}$ attains the maximum when $x_1 = x_2 = x_4 = b/2$ and $x_3 = 0$. Thus, it takes in $O((d^* + 2)^b \{(d^* + 1)^2 + 1\}^{b/2})$ time to compute $A_f(c)$ for all colorings c of $w(f)$.

Since $|E(T)| = O(m)$ and the initialization step takes $O((d^* + 2)^b m)$ time in total, we can obtain $A_{(r_1, r_2)}(c)$ in $O(((d^* + 2)^b \{(d^* + 1)^2 + 1\}^{b/2} m)$ time.

Lemma 3. *Let $c \in \{\top, 0, 1, 2, \dots, d^*\}^{|w(f)|}$ be a coloring of $w(f)$. If a coloring c_1 of $w(f_1)$ and a coloring c_2 of $w(f_2)$ forms c , then $D_{f_1}(c_1) \cup D_{f_2}(c_2)$ satisfies (1)–(3) for f .*

Proof. We denote $D_{f_1}(c_1) \cup D_{f_2}(c_2)$ by D' , and $D' \cap (X_1 \cup X_2 \cup X_3 \cup X_4)$ by D'_{c_1, c_2} . Clearly, (1) holds, since $v \in D' \cap w(f)$ if and only if $c(v) = \top$ by the above (P1).

We next show that D' satisfies (2). Let v be a vertex in $X_1 - D' = X_1 - D'_{c_1, c_2}$. From the above (P3), we have $|N_{G_{f_1}}(v) \cap D'| \geq d(v) - i - |D'_{c_1, c_2} \cap N_{G_f}(v) \cap X_2|$. It follows that $|N_{G_f}(v) \cap D'| \geq |N_{G_{f_1}}(v) \cap D'| + |D'_{c_1, c_2} \cap N_{G_f}(v) \cap X_2| \geq d(v) - i$. Also, the case of $v \in X_2 - D'$ can be treated similarly.

Let v be a vertex in $X_3 - D' = X_3 - D'_{c_1, c_2}$. Since $|N_{G_f}(v) \cap D'| \geq |N_{G_f}(v) \cap D'_{c_1, c_2}|$ clearly holds, then we have only to consider the case of $|N_{G_f}(v) \cap D'_{c_1, c_2}| < d(v) - i$. From (P4), we have $|N_{G_{f_1}}(v) \cap D'| \geq \max\{0, d(v) - i - |D'_{c_1, c_2} \cap N_{G_f}(v) \cap X_2| - i_1\}$ and $|N_{G_{f_2}}(v) \cap D'| \geq \max\{0, d(v) - i - |D'_{c_1, c_2} \cap N_{G_f}(v) \cap X_1| - i_2\}$ where $i_1 + i_2 = d(v) - i - |D'_{c_1, c_2} \cap N_{G_f}(v)|$ (note that $i_1 + i_2 > 0$ from the assumption of this case). Notice that $(V(G_{f_1}) - w(f_1)) \cap (V(G_{f_2}) - w(f_2)) = \emptyset$ by Lemma 2(ii). It follows that $|N_{G_f}(v) \cap D'| \geq |N_{G_{f_1}}(v) \cap D'| + |N_{G_{f_2}}(v) \cap D'| - |N_{G_f}(v) \cap D'_{c_1, c_2} \cap (X_3 \cup X_4)| \geq 2(d(v) - i) - |N_{G_f}(v) \cap D'_{c_1, c_2}| - i_1 - i_2 = d(v) - i$.

We finally show that D' satisfies (3). Let v be a vertex in $X_4 - D'$. Since $|N_{G_f}(v) \cap D'| \geq |N_{G_f}(v) \cap D'_{c_1, c_2}|$ clearly holds, then we have only to consider the case of $|N_{G_f}(v) \cap D'_{c_1, c_2}| < d(v)$. From (P5), we have $|N_{G_{f_1}}(v) \cap D'| \geq$

$\max\{0, d(v) - |D'_{c_1, c_2} \cap N_{G_f}(v) \cap X_2| - i_1\}$ and $|N_{G_{f_2}}(v) \cap D'| \geq \max\{0, d(v) - |D'_{c_1, c_2} \cap N_{G_f}(v) \cap X_1| - i_2\}$ where $i_1 + i_2 = d(v) - |D'_{c_1, c_2} \cap N_{G_f}(v)| > 0$. Hence, we have $|N_{G_f}(v) \cap D'| \geq |N_{G_{f_1}}(v) \cap D'| + |N_{G_{f_2}}(v) \cap D'| - |N_{G_f}(v) \cap D'_{c_1, c_2} \cap (X_3 \cup X_4)| = 2d(v) - |N_{G_f}(v) \cap D'_{c_1, c_2}| - i_1 - i_2 = d(v)$. Also, it follows from the definition of $D_{f_j}(c_j)$ that $v \in V(G_{f_j}) - w(f_j)$ satisfies (3) for $j = 1, 2$. \square

Lemma 4. *Let $c \in \{\top, 0, 1, 2, \dots, d^*\}^{|w(f)|}$ be a coloring of $w(f)$. There exist a coloring c_1 of $w(f_1)$ and a coloring c_2 of $w(f_2)$ forming c such that $|D_{f_1}(c_1) \cup D_{f_2}(c_2)| \leq A_f(c)$.*

Proof. For each vertex $v \in w(f_j)$, $j = 1, 2$, let

$$c_j(v) = \begin{cases} \top & \text{if } v \in D_f(c), \\ \min\{d(v), c(v) + |N_{G_f}(v) \cap D_f(c) - V(G_{f_j})|\} & \text{if } v \in X_j - D_f(c), \\ \max\{0, d(v) - |N_{G_{f_j}}(v) \cap D_f(c)|\} & \text{if } v \in X_3 \cup X_4 - D_f(c). \end{cases}$$

For $v \in X_j - D_f(c)$, we have $|N_{G_f}(v) \cap D_f(c)| = |N_{G_{f_j}}(v) \cap D_f(c)| + |N_{G_f}(v) \cap D_f(c) - V(G_{f_j})| \geq d(v) - c(v)$, since $D_f(c)$ satisfies (2). Hence, $|N_{G_{f_j}}(v) \cap D_f(c)| \geq \max\{0, d(v) - c(v) - |N_{G_f}(v) \cap D_f(c) - V(G_{f_j})|\} = d(v) - c_j(v)$ for all $v \in w(f_j) - D_f(c)$. It follows from that the minimality of $A_{f_j}(c_j)$ implies that $|D_f(c) \cap V(G_{f_j})| \geq A_{f_j}(c_j)$; hence, $A_f(c) \geq |D_{f_1}(c_1) \cup D_{f_2}(c_2)|$. On the other hand, c_1 and c_2 does not necessarily form c . Below, we show that there exist a coloring c'_1 of $w(f_1)$ and a coloring c'_2 of $w(f_2)$ forming c such that $c'_j(v) \geq c_j(v)$ for every $v \in w(f_j) - D_f(c)$ for $j = 1, 2$. Note that $D_{f_j}(c_j)$ satisfies (1)–(3) also for c'_j , since $|N_{G_{f_j}}(v) \cap D_{f_j}(c)| \geq d(v) - c_j(v) \geq d(v) - c'_j(v)$ for every $v \in w(f_j) - D_f(c)$. Hence, from the minimality of $|D_{f_j}(c'_j)|$, we have $A_f(c) \geq |D_{f_1}(c_1) \cup D_{f_2}(c_2)| \geq |D_{f_1}(c'_1) \cup D_{f_2}(c'_2)|$, which proves this lemma.

We can construct such c'_1, c'_2 as follows. First let $c'_j(v) = c_j(v)$ for all $v \in X_1 \cup X_2 \cup D_f(c)$; c'_1 and c'_2 satisfy (P1) and (P2) in the definition of a coloring c formed by c_1 and c_2 . By Lemma 2(ii), every $v \in X_j$ satisfies $N_{G_f}(v) \cap D_f(c) - V(G_{f_j}) = N_{G_f}(v) \cap D_f(c) \cap X_{j'}$ for $\{j, j'\} = \{1, 2\}$. Hence, $c'_j(v) (= c_j(v))$ for $v \in X_j - D_f(c)$, $j = 1, 2$ satisfies (P3).

Let $v \in X_3 - D_f(c)$. Since $D_f(c)$ satisfies (2), we have $|N_{G_f}(v) \cap D_f(c)| \geq d(v) - c(v)$. Now from construction of c_1 and c_2 , the value i'_1 (resp., i'_2) corresponding to i_1 (resp., i_2) in (P4) in the definition of c formed by c_1 and c_2 is $\max\{0, d(v) - |N_{G_{f_1}}(v) \cap D_f(c)| - c(v) - |N_{G_f}(v) \cap X_2 \cap D_f(c)|\}$ (resp., $\max\{0, d(v) - |N_{G_{f_2}}(v) \cap D_f(c)| - c(v) - |N_{G_f}(v) \cap X_1 \cap D_f(c)|\}$). It follows that $i'_1 + i'_2 \leq \max\{0, d(v) - c(v) - |N_{G_f}(v) \cap D_f(c) \cap (X_1 \cup X_2 \cup X_3 \cup X_4)|\}$ (note that the final inequality follows from $|N_{G_f}(v) \cap D_f(c)| \geq d(v) - c(v)$).

Let $v \in X_4 - D_f(c)$. Since $D_f(c)$ satisfies (2), we have $|N_{G_f}(v) \cap D_f(c)| \geq d(v)$. From construction of c_1 and c_2 , the value i'_1 (resp., i'_2) corresponding to i_1 (resp., i_2) in (P5) in the definition of c formed by c_1 and c_2 is $\max\{0, d(v) - |N_{G_{f_1}}(v) \cap D_f(c)| - |N_{G_f}(v) \cap X_2 \cap D_f(c)|\}$ (resp., $\max\{d(v) - |N_{G_{f_2}}(v) \cap D_f(c)| - |N_{G_f}(v) \cap X_1 \cap D_f(c)|\}$). It follows that $i'_1 + i'_2 \leq \max\{0, d(v) - |N_{G_f}(v) \cap D_f(c) \cap (X_1 \cup X_2 \cup X_3 \cup X_4)|\}$.

Consequently, we can construct a coloring c'_1 of $w(f_1)$ and a coloring c'_2 of $w(f_2)$ forming c such that $c'_j(v) \geq c_j(v)$ for every $v \in X_3 \cup X_4 - D_f(c)$

and $c'_j(v) = c_j(v)$ for every $v \in D_f(c) \cup X_1 \cup X_2$ for $j = 1, 2$ by increasing i'_1 or i'_2 for each vertex $v \in X_3 \cup X_4 - D_f(c)$ so that $i'_1 + i'_2$ becomes equal to $\max\{0, d(v) - c(v) - |N_{G_f}(v) \cap D_f(c) \cap (X_1 \cup X_2 \cup X_3 \cup X_4)|\}$ (resp., $\max\{0, d(v) - |N_{G_f}(v) \cap D_f(c) \cap (X_1 \cup X_2 \cup X_3 \cup X_4)|\}$) if $v \in X_3$ (resp., $v \in X_4$). \square

Summarizing the arguments given so far, we have shown Theorem 1.

3.2 Total vector domination and multiple domination

We consider the total vector domination problem. The difference between the total vector domination and the vector domination is that each vertex selected as a member in a dominating set needs to be dominated or not. Hence, we will modify the following parts (I)–(III) in the algorithm for vector domination given in the previous subsection so that each vertex selected as a member in a dominating set also satisfies its demand.

(I) Color assignments: Let $f \in E(T)$ be an edge in a branch decomposition T of G . We will assign to each vertex $v \in w(f)$ an ordered pair (ℓ, i) of colors, $\ell \in \{\top, \perp\}$, $i \in \{0, 1, \dots, d(v)\}$, where \top means that v belongs to the dominating set, \perp means that v does not belong to the dominating set, and i means that v is dominated by at least $d(v) - i$ vertices in G_f .

(II) Conditions for $D_f(c)$: For a coloring $c \in (\{\top, \perp\} \times \{0, 1, 2, \dots, d^*\})^{|w(f)|}$, we modify (1)–(3) as follows, where let $c(v) = (c^1(v), c^2(v))$:

$$c^1(v) = \top \text{ if and only if } v \in D_f(c) \cap w(f).$$

$$\text{If } c^2(v) = i, \text{ then } |N_{G_f}(v) \cap D_f(c)| \geq d(v) - i.$$

$$|N_{G_f}(v) \cap D_f(c)| \geq d(v) \text{ holds for every vertex } v \in V(G_f) - w(f).$$

(III) Definition of a coloring c formed by c_1 and c_2 : For a coloring $c \in (\{\top, \perp\} \times \{0, 1, 2, \dots, d^*\})^{|w(f)|}$, we modify (P1)–(P5) as follows:

(P1') For every $v \in X_1 \cup X_2 \cup X_3$ with $c^1(v) = \top$ (resp., $c^1(v) = \perp$),

(a) If $v \in X_1 \cup X_3$, then $c_1^1(v) = \top$ (resp., $c_1^1(v) = \perp$).

(b) If $v \in X_2 \cup X_3$, then $c_2^1(v) = \top$ (resp., $c_2^1(v) = \perp$).

(P2') For every $v \in X_4$, $c_1^1(v) = \top$ (resp., $c_1^1(v) = \perp$) if and only if $c_2^1(v) = \top$ (resp., $c_2^1(v) = \perp$).

(P3') For every $v \in X_j$ where $\{j, j'\} = \{1, 2\}$ and $D_{c_1, c_2} = \{v \in X_1 \cup X_2 \cup X_3 \cup X_4 \mid c_1^1(v) = \top \text{ or } c_2^1(v) = \top\}$,

$$\text{If } c^2(v) = i, \text{ then } c_j^2(v) = \min\{d(v), i + |D_{c_1, c_2} \cap N_{G_f}(v) \cap X_{j'}|\}.$$

(P4') For every $v \in X_3$,

$$\text{If } c^2(v) = i, \text{ then } c_1^2(v) = \min\{d(v), i + |D_{c_1, c_2} \cap N_{G_f}(v) \cap X_2| + i_1\} \text{ and } c_2^2(v) = \min\{d(v), i + |D_{c_1, c_2} \cap N_{G_f}(v) \cap X_1| + i_2\} \text{ for some nonnegative integers } i_1, i_2 \text{ with } i_1 + i_2 = \max\{0, d(v) - i - |D_{c_1, c_2} \cap N_{G_f}(v) \cap (X_1 \cup X_2 \cup X_3 \cup X_4)|\}.$$

(P5') For every $v \in X_4$,

$$c_1^2(v) = \min\{d(v), |D_{c_1, c_2} \cap N_{G_f}(v) \cap X_2| + i_1\} \text{ and } c_2^2(v) = \min\{d(v), |D_{c_1, c_2} \cap N_{G_f}(v) \cap X_1| + i_2\} \text{ for some nonnegative integers } i_1, i_2 \text{ with } i_1 + i_2 = \max\{0, d(v) - |D_{c_1, c_2} \cap N_{G_f}(v) \cap (X_1 \cup X_2 \cup X_3 \cup X_4)|\}.$$

We analyze the time complexity of this modified algorithm. Similarly to the case of the vector domination, the total running time is dominated by total complexity for computing $A_f(c)$ for non-leaf edges f .

Let f be a non-leaf edge of T and $x_i, i = 1, 2, 3, 4$ and z_4 be defined as the previous subsection. The number of pairs of a coloring c_1 of $w(f_1)$ and a coloring c_2 of $w(f_2)$ forming c is at most

$$(d^* + 1)^{x_3} \sum_{z_4=0}^{x_4} \binom{x_4}{z_4} (d^* + 1)^{x_4} (d^* + 1)^{x_4}$$

since the number of pairs (i_1, i_2) in (P4') or (P5') is at most $d^* + 1$ for each vertex in $X_3 \cup X_4$. Hence, for an edge f , the number of pairs forming c is at most

$$\{2(d^* + 1)\}^{x_1+x_2} \sum_{z_3=0}^{x_3} \binom{x_3}{z_3} (d^* + 1)^{x_3} (d^* + 1)^{x_3} \sum_{z_4=0}^{x_4} \binom{x_4}{z_4} (d^* + 1)^{x_4} (d^* + 1)^{x_4} \\ = \{2(d^* + 1)\}^{x_1+x_2} \{2(d^* + 1)^2\}^{x_3+x_4}$$

in total. Since $x_1 + x_2 + x_3 \leq b$, $x_1 + x_3 + x_4 \leq b$, and $x_2 + x_3 + x_4 \leq b$, it follows that $(x_1 + x_2) \log(2d^* + 2) + (x_3 + x_4) \log\{2(d^* + 1)^2\}$ attains the maximum when $x_1 = x_2 = x_4 = b/2$ and $x_3 = 0$. Thus, it takes in $O(2^{3b/2}(d^* + 1)^{2b})$ time to compute $A_f(c)$ for all colorings c of $w(f)$. Namely, we obtain the following theorem.

Theorem 2. *If a branch decomposition of G with width b is given, a minimum total vector dominating set in G can be found in $O(2^{3b/2}(d^* + 1)^{2b}m)$ time. \square*

Also, by replacing $N_G()$ with $N_G[]$ in the modification for total vector domination, we can obtain the following theorem for the multiple domination problems.

Theorem 3. *If a branch decomposition of G with width b is given, a minimum multiple dominating set in G can be found in $O(2^{3b/2}(d^* + 1)^{2b}m)$ time. \square*

4 Subexponential fixed parameter algorithm for planar graphs

We consider the problem of checking whether a given graph G has a d -vector dominating set with cardinality at most k . As mentioned in Subsection 1.1, if G is ρ -degenerated, then the problem can be solved in $k^{O(\rho k^2)}n^{O(1)}$ time. Since a planar graph is 5-degenerated, it follows that the problem with a planar graph can be solved in $k^{O(k^2)}n^{O(1)}$ time. In this section, we give a subexponential fixed-parameter algorithm, parameterized by k , for a planar graph; namely, we will show the following theorem.

Theorem 4. *If G is a planar graph, then we can check in $O(n^4 + (\min\{d^*, k\} + 2)^{b^*} \{(\min\{d^*, k\} + 1)^2 + 1\}^{b^*/2} n)$ time whether G has a d -vector dominating set with cardinality at most k or not, where $b^* = \min\{12\sqrt{k+z} + 9, 20\sqrt{k} + 17\}$ and $z = |\{v \in V \mid d(v) = 0\}|$. \square*

This time complexity is roughly $O(n^4 + 2^{O(\sqrt{k} \log k)} n)$, which is subexponential with respect to k ; this improves the running time of the previous fixed-parameter algorithm.

Let $V_0 = \{v \in V \mid d(v) = 0\}$ and $z = |V_0|$. In [18, Lemma 2.2], it was shown that if a planar graph G' has an ordinary dominating set (i.e., a $(1, 1, \dots, 1)$ -vector dominating set) with cardinality at most k , then $bw(G') \leq 12\sqrt{k} + 9$. This bound is based on the *bidimensionality* [14], and was used to design the subexponential fixed-parameter algorithm with respect to k for the ordinary dominating set problem. In the case of our domination problems, however, it is difficult to say that they have the bidimensionality, due to the existence of V_0 vertices. Instead, we give a similar bound on the branchwidth not w.r.t k but w.r.t $k+z$ as follows: For any (total, multiple) d -vector dominating set D of G ($|D| \leq k$), $D \cup V_0$ is an ordinary dominating set of G , and this yields $bw(G) \leq 12\sqrt{k+z} + 9$.

Actually, it is also possible to exclude z from the parameters, though the coefficient of the exponent becomes larger. To this end, we use the notion of $(k, 2)$ -center. Recall that a (k, r) -center of G' is a set W of vertices of G' with size k such that any vertex in G' is within distance r from a vertex of W . For a (k, r) -center, a similar bound on the branchwidth is known: if a planar graph G' has a (k, r) -center, then $bw(G') \leq 4(2r+1)\sqrt{k} + 8r + 1$ ([12, Theorem 3.2]). Here, we use this bound. We can assume that for $v \in V_0$, $N_G(v) \not\subseteq V_0$ holds, because $v \in V_0$ satisfying $N_G(v) \subseteq V_0$ is never selected as a member of any optimal solution; it is *irrelevant*, and we can remove it. That is, every vertex in V_0 has at least one neighbor from $V - V_0$. Then, for any (total, multiple) d -vector dominating set D of G ($|D| \leq k$), D is a $(k, 2)$ -center of G . This is because any vertex in $V - V_0$ is adjacent to a vertex in D and any vertex in V_0 is adjacent to a vertex in $V - V_0$. Thus, we have $bw(G) \leq 20\sqrt{k} + 17$.

In summary, we have the following lemma.

Lemma 5. *Assume that G is a planar graph without irrelevant vertices, i.e., $N_G(v) \not\subseteq V_0$ holds for each $v \in V_0$. Then, if G has a (total, multiple) vector dominating set with cardinality at most k , then we have $bw(G) \leq \min\{12\sqrt{k+z} + 9, 20\sqrt{k} + 17\}$. \square*

Combining this lemma with the algorithm in Subsection 3.1, we can check whether a given graph has a vector dominating set with cardinality at most k according to the following steps 1 and 2:

Step 1: Let $b^* = \min\{12\sqrt{k+z}+9, 20\sqrt{k}+17\}$. Check whether the branchwidth of G is at most b^* . If so, then go to Step 2, and otherwise halt after outputting ‘NO’.

Step 2: Construct a branch decomposition with width at most b^* , and apply the dynamic programming algorithm in Subsection 3.1 to find a minimum vector dominating set for G .

By Lemma 1, Theorem 1, and the fact that any planar graph G' satisfies $|E(G')| = O(|V(G')|)$, it follows that the running time of this procedure is $O(n^4 + (d^* + 2)^{b^*} \{(d^* + 1)^2 + 1\}^{b^*/2} n)$. Hence, in the case of $d^* \leq k$, Theorem 4 has been proved.

The case of $d^* > k$ can be reduced to the case of $d^* \leq k$ by the following standard kernelization method, which proves Theorem 4. Assume that $d^* > k$. Let $V_{\max}(d)$ be the set of vertices v with $d(v) = d^*$. For the feasibility, we need to select each vertex $v \in V_{\max}(d)$ as a member in a vector dominating set. Hence, if $|V_{\max}(d)| > k$, then it turns out that G has no vector dominating set with cardinality at most k . Assume that $|V_{\max}(d)| \leq k$. Then, it is not difficult to see that we can reduce an instance $I(G, d, k)$ with G , d , and k to an instance $I(G', d', k')$ such that $G' = G - V_{\max}(d)$ (i.e., G' is the graph obtained from G by deleting $V_{\max}(d)$), $d'(v) = \max\{0, d(v) - |N_G(v) \cap V_{\max}(d)|\}$ for all vertices $v \in V(G')$, and $k' = \max\{0, k - |V_{\max}(d)|\}$. Based on this observation, we can reduce $I(G, d, k)$ to an instance $I(G'', d'', k'')$ with $\max\{d''(v) \mid v \in V(G'')\} \leq k'' \leq k$ or output ‘YES’ or ‘NO’ in the following manner:

(a) After setting $G' := G$, $d' := d$, and $k' := k$, repeat the procedures (b1)–(b3) while $k' < d'^*$ ($= \max\{d'(v) \mid v \in V(G')\}$).

(b1) If $k' < |V_{\max}(d')|$, then halt after outputting ‘NO.’

(b2) If $k' \geq |V_{\max}(d')|$ and $V(G') = V_{\max}(d')$, then halt after outputting ‘YES.’

(b3) Otherwise after setting $G'' := G' - V_{\max}(d')$, $d''(v) := \max\{0, d'(v) - |N_{G'}(v) \cap V_{\max}(d')|\}$ for each $v \in V(G'')$, and $k'' := \max\{0, k' - |V_{\max}(d')|\}$, redefine G'' , d'' , and k'' as G' , d' , and k' , respectively.

Next, we consider the total vector domination problem and the multiple domination problem. For these problems, since all vertices $v \in V$ need to be dominated by $d(v)$ vertices, the condition that $d^* \leq k$ is necessary for the feasibility. Similarly, we have the following theorem by Theorems 2 and 3.

Theorem 5. *Assume that a given graph G is planar, and let $b^* = \min\{12\sqrt{k+z}+9, 20\sqrt{k}+17\}$.*

(i) *We can check in $O(n^4 + 2^{3b^*/2}(\min\{d^*, k\} + 2)^{2b^*} n)$ time whether G has a total vector dominating set with cardinality at most k or not.*

(ii) *We can check in $O(n^4 + 2^{3b^*/2}(\min\{d^*, k\} + 2)^{2b^*} n)$ time whether G has a multiple dominating set with cardinality at most k or not. \square*

Before concluding this section, we mention that the above result can be extended to apex-minor-free graphs, a superclass of planar graphs. For apex-minor-free graphs, the following lemma is known.

Lemma 6. ([17, Lemma 2]) *Let G be an apex-minor-free graph. If G has a (k, r) -center, then the treewidth of G is $O(r\sqrt{k})$.*

From this lemma, the linear relation of treewidth and branchwidth, and the $2^{O(bw(G))}n^2$ -time algorithm for computing a branch decomposition with width $O(bw(G))$ (mentioned after Theorem 1), we obtain the following corollary.

Corollary 2. *Let G be an apex-minor-free graph. We can check in $2^{O(\sqrt{k} \log k)}n^{O(1)}$ time whether G has a (total, multiple) vector dominating set with cardinality at most k or not.*

References

1. N. Alon, D. Moshkovitz, and S. Safra. Algorithmic construction of sets for k -restrictions. *ACM Transactions on Algorithms (TALG)*, 2(2):153–177, 2006.
2. O. Amini, F. V. Fomin, and S. Saurabh. Implicit branching and parameterized partial cover problems. *Journal of Computer and System Sciences*, 77(6):1159–1171, 2011.
3. N. Betzler, R. Bredereck, R. Niedermeier, and J. Uhlmann. On bounded-degree vertex deletion parameterized by treewidth. *Discrete Applied Mathematics*, 160(1):53–60, 2012.
4. H. L. Bodlaender and D. M. Thilikos. Constructive linear time algorithms for branchwidth. In *Automata, Languages and Programming*, volume 1256 of *Lecture Notes in Computer Science*, pages 627–637. Springer, 1997.
5. L. Cai and D. Juedes. On the existence of subexponential parameterized algorithms. *Journal of Computer and System Sciences*, 67(4):789–807, 2003.
6. M. Chapelle. Parameterized complexity of generalized domination problems on bounded tree-width graphs. *arXiv preprint arXiv:1004.2642*, 2010.
7. F. Cicalese, G. Cordasco, L. Gargano, M. Milanić, and U. Vaccaro. Latency-bounded target set selection in social networks. In *The Nature of Computation. Logic, Algorithms, Applications*, pages 65–77. Springer, 2013.
8. F. Cicalese, M. Milanić, and U. Vaccaro. Hardness, approximability, and exact algorithms for vector domination and total vector domination in graphs. In *Fundamentals of Computation Theory*, volume 6914 of *Lecture Notes in Computer Science*, pages 288–297. Springer Berlin Heidelberg, 2011.
9. F. Cicalese, M. Milanić, and U. Vaccaro. On the approximability and exact algorithms for vector domination and related problems in graphs. *Discrete Applied Mathematics*, 161(6):750 – 767, 2013.
10. B. Courcelle. The monadic second-order logic of graphs. I. recognizable sets of finite graphs. *Information and computation*, 85(1):12–75, 1990.
11. B. Courcelle and S. Olariu. Upper bounds to the clique width of graphs. *Discrete Applied Mathematics*, 101(1):77–114, 2000.

12. E. D. Demaine, F. V. Fomin, M. Hajiaghayi, and D. M. Thilikos. Fixed-parameter algorithms for (k, r) -center in planar graphs and map graphs. *ACM Transactions on Algorithms (TALG)*, 1(1):33–47, 2005.
13. E. D. Demaine, F. V. Fomin, M. Hajiaghayi, and D. M. Thilikos. Subexponential parameterized algorithms on bounded-genus graphs and H -minor-free graphs. *Journal of the ACM (JACM)*, 52(6):866–893, 2005.
14. E. D. Demaine and M. Hajiaghayi. The bidimensionality theory and its algorithmic applications. *The Computer Journal*, 51(3):292–302, 2008.
15. G. Dobson. Worst-case analysis of greedy heuristics for integer programming with nonnegative data. *Mathematics of Operations Research*, 7(4):515–531, 1982.
16. R. G. Downey and M. R. Fellows. *Fixed-parameter tractability and completeness*. Cornell University, Mathematical Sciences Institute, 1992.
17. F. V. Fomin, D. Lokshtanov, V. Raman, and S. Saurabh. Subexponential algorithms for partial cover problems. *Information Processing Letters*, 111(16):814–818, 2011.
18. F. V. Fomin and D. M. Thilikos. Dominating sets in planar graphs: branch-width and exponential speed-up. *SIAM Journal on Computing*, 36(2):281–309, 2006.
19. J. Harant, A. Pruchnewski, and M. Voigt. On dominating sets and independent sets of graphs. *Combinatorics, Probability and Computing*, 8:547–553, 10 1999.
20. F. Harary and T. W. Haynes. Double domination in graphs. *Ars Combinatoria*, 55:201–214, 2000.
21. T. W. Haynes, S. T. Hedetniemi, and P. J. Slater. *Domination in graphs: advanced topics*, volume 40. Marcel Dekker New York, 1998.
22. T. W. Haynes, S. T. Hedetniemi, and P. J. Slater. *Fundamentals of domination in graphs*. Marcel Dekker, 1998.
23. R. Impagliazzo, R. Paturi, and F. Zane. Which problems have strongly exponential complexity? *Journal of Computer and System Sciences*, 63(4):512–530, 2001.
24. C. Lund and M. Yannakakis. On the hardness of approximating minimization problems. *Journal of the ACM (JACM)*, 41(5):960–981, 1994.
25. V. Raman, S. Saurabh, and S. Srihari. Parameterized algorithms for generalized domination. In *Combinatorial Optimization and Applications*, pages 116–126. Springer, 2008.
26. N. Robertson and P. D. Seymour. Graph minors. X. obstructions to tree-decomposition. *Journal of Combinatorial Theory, Series B*, 52(2):153–190, 1991.
27. N. Robertson and P. D. Seymour. Graph minors. XIII. the disjoint paths problem. *Journal of Combinatorial Theory, Series B*, 63(1):65–110, 1995.
28. P. D. Seymour and R. Thomas. Call routing and the ratcatcher. *Combinatorica*, 14(2):217–241, 1994.