

Preprint No. M 07/03

**Edge colouring by total labellings**

Brandt, Stephan; Rautenbach, Dieter;  
Stiebitz, Michael

Juni 2007

**Impressum:**

Hrsg.: Leiter des Instituts für Mathematik  
Weimarer Straße 25  
98693 Ilmenau  
Tel.: +49 3677 69 3621  
Fax: +49 3677 69 3270  
<http://www.tu-ilmenau.de/ifm/>

ISSN xxxx-xxxx

ilmedia

# Edge colouring by total labellings

Stephan Brandt, Dieter Rautenbach, and Michael Stiebitz \*

June 22, 2007

## Abstract

We introduce the concept of an edge-colouring total  $k$ -labelling. This is a labelling of the vertices and the edges of a graph  $G$  with labels  $1, 2, \dots, k$  such that the weights of the edges define a proper edge colouring of  $G$ . Here the weight of an edge is the sum of its label and the labels of its two endvertices. We define  $\chi'_t(G)$  to be the smallest integer  $k$  for which  $G$  has an edge-colouring total  $k$ -labelling. This parameter has natural upper and lower bounds in terms of the maximum degree  $\Delta$  of  $G$ :  $\lceil(\Delta + 1)/2\rceil \leq \chi'_t(G) \leq \Delta + 1$ . We improve the upper bound by 1 for every graph and prove a general upper bound of  $\chi'_t(G) \leq \Delta/2 + \mathcal{O}(\sqrt{\Delta \log \Delta})$ . Moreover, we investigate some special classes of graphs.

**Keywords** Edge colouring; total labelling; irregularity strength; discrepancy

**MSC Classification** 05C15; 05C78; 05D40

## 1 Introduction

For a graph  $G = (V(G), E(G))$  an *edge-colouring total  $k$ -labelling* is a function  $f : V(G) \cup E(G) \rightarrow \{1, 2, \dots, k\}$  such that the weights of the edges defined by

$$w(uv) := f(u) + f(uv) + f(v)$$

form a proper edge colouring. The smallest integer  $k$  for which there exists an edge-colouring total  $k$ -labelling is denoted by  $\chi'_t(G)$ .

A related concept which has recently received a lot of attention was proposed by Karoński, Łuczak and Thomason [16]. They conjectured that the edges of every graph  $G$  with no  $K_2$  component can be labeled with labels  $1, 2, 3$  such that the sums of the edge labels incident to the vertices of  $G$  define a proper vertex colouring. Addario-Berry, Dalal and Reed [2] recently proved that the labels  $1, 2, \dots, 16$  are always sufficient, i.e. every

---

\*Technische Universität Ilmenau, Fak. Mathematik & Naturwissenschaften, TU Ilmenau, Postfach 100565, 98684 Ilmenau, Germany, e-mail: {stephan.brandt, dieter.rautenbach,michael.stiebitz}@tu-ilmenau.de

graph with no  $K_2$  component has a *vertex-colouring edge 16-labelling* (cf. also [1, 3]). A total version of vertex-colouring labellings was discussed by Przybyło and Woźniak who proved [19] by similar methods as in [2] that every graph has a *vertex-colouring total 11-labelling* and conjecture that 2 labels are enough.

The vertex-colouring edge labellings can be considered a relaxation of the well-known *irregularity strength* of graphs [10, 4, 18, 14] where the label sums for all vertices are required to be different. Similarly, the edge-colouring total labellings which we study here can be considered a relaxation of *edge-irregular total labellings* introduced by Bača, Jendrol', Miller, and Ryan [6], where the weights of all edges are required to be different. The *total edge irregularity strength*  $\text{tes}(G)$  is defined as the smallest integer  $k$  for which a graph  $G$  has an edge-irregular total  $k$ -labelling. A simple lower bound is

$$\text{tes}(G) \geq \max \left\{ \left\lceil \frac{|E(G)| + 2}{3} \right\rceil, \left\lceil \frac{\Delta(G) + 1}{2} \right\rceil \right\}.$$

and Ivančo and Jendrol' [15] conjectured that this bound is attained for all graphs except  $K_5$ . Brandt, Miškuf, and Rautenbach [8, 9] recently proved that this is true for graphs whose size is at least 111000 times their maximum degree.

Let us return to the edge-colouring total  $k$ -labellings and the corresponding graph parameter  $\chi'_t(G)$  which has natural upper and lower bounds in terms of the maximum degree  $\Delta$  of  $G$ . Obviously,

$$\chi'_t(G) \leq \Delta + 1$$

by Vizing's Theorem [20], since a proper edge colouring together with a constant labelling of the vertices defines an edge-colouring total labelling of  $G$ . Furthermore, since the possible weights of the edges incident with a vertex  $v$  of maximum degree  $\Delta$  in an edge-colouring  $k$ -labelling  $f$  are  $f(v) + \{2, 3, \dots, 2k\}$ , we get a lower bound of

$$\chi'_t(G) \geq \left\lceil \frac{\Delta + 1}{2} \right\rceil.$$

The following is our main result whose proof we postpone to Section 3.

**Theorem 1.1** *If  $G$  is a graph of maximum degree  $\Delta$ , then*

$$\chi'_t(G) \leq \left\lfloor \frac{1}{2} \left( \Delta + \left\lfloor \sqrt{2\Delta(1 + \ln(2\Delta^2 - 2\Delta + 2))} \right\rfloor \right) \right\rfloor + 1 = \Delta/2 + \mathcal{O}(\sqrt{\Delta \log \Delta}).$$

Before we proceed to Section 2 where we study  $\chi'_t(G)$  for some special graphs, we show how to reduce the upper bound by one for every graph and relate  $\chi'_t(G)$  to the chromatic index. The next result already illustrates our general approach which is to combine edge colouring methods with suitable partitions of the vertex set.

**Theorem 1.2** *If  $G$  is a graph of maximum degree  $\Delta$ , then  $\chi'_t(G) \leq \Delta$*

**Proof:** Let  $c : E(G) \rightarrow \{1, 2, \dots, \Delta + 1\}$  be a proper edge colouring of  $G$  which exists by Vizing's Theorem [20]. Since the subgraph containing the edges coloured  $\Delta$  and  $\Delta + 1$  consists of paths and even cycles, it is bipartite. Fix a bipartition  $A \cup B$  of  $V(G)$  such that all edges with colours  $\Delta$  and  $\Delta + 1$  have one endvertex in  $A$  and the other endvertex in  $B$ .

Assign to all vertices of  $A$  the label 1 and to all vertices of  $B$  the label  $\Delta$ . Assign label  $c(e)$  to all edges between vertices of  $A$  and label  $c(e) + 1$  to all edges between vertices of  $B$ . Finally, determine the labels of the edges in the bipartite graph spanned by the edges between  $A$  and  $B$  by a proper  $\Delta$ -edge colouring  $c'$ .

The edges joining vertices of  $A$  receive weights between 3 and  $(\Delta - 1) + 1 + 1 = \Delta + 1$ , the edges joining  $A$  to  $B$  receive weights between  $\Delta + 2$  and  $2\Delta + 1$ , and the edges joining vertices of  $B$  receive weights between  $2\Delta + 2$  and  $3\Delta$ . Since these weights form proper edge colourings inside and between the sets, they form a proper edge colouring of the entire graph.  $\square$

The upper bound  $\chi'_t(G) \leq \Delta$  can only be tight for small values of  $\Delta$ . From Theorem 1.1 follows that for  $\Delta \geq 19$  we have  $\chi'_t(G) < \Delta$ , and, in fact, with a more refined reasoning along the same lines the threshold can be reduced to  $\Delta \geq 14$ . We are not aware of any graph with  $\Delta > 3$  and  $\chi'_t(G) = \Delta$ .

Next we show that an edge-colouring total  $k$ -labelling gives rise to a proper edge colouring with  $2k - 1$  colours. Conversely, this means that for every type II graph (i.e.  $\chi'(G) = \Delta(G) + 1$ ) we have  $\chi'_t(G) > \frac{\Delta(G)+1}{2}$ .

**Lemma 1.3** *If  $\chi'_t(G) = k$  for a graph  $G$ , then  $\chi'(G) \leq 2k - 1$ .*

**Proof:** Consider an edge-colouring total  $k$ -labelling  $f$  of  $G$ . Note that for  $l \leq k + 1$  the edges of weights  $l$  and  $l + 2k - 1$  cannot have a common endvertex and therefore form a matching. Thus we can decompose the edge set into  $2k - 1$  matchings:  $k - 1$  matchings with the edges of weight  $l$  and  $l + 2k - 1$  for  $3 \leq l \leq k + 1$ , and  $k$  matchings with the edges of weight  $l$  for  $k + 2 \leq l \leq 2k + 1$ .  $\square$

## 2 Special classes of graphs

If  $G$  is a graph of maximum degree  $\Delta = 1$ , then  $\chi'_t(G) = 1$ . If  $\Delta = 2$ , then  $\chi'_t(G) = 2$  by Theorem 1.2. Similarly, if  $\Delta(G) = 3$ , then  $2 \leq \chi'_t(G) \leq 3$ . In our first result we characterize cubic graphs with  $\chi'_t(G) = 2$ .

**Theorem 2.1** *A cubic graph  $G$  satisfies  $\chi'_t(G) = 2$  if and only if its vertex set can be partitioned into two parts  $A$  and  $B$  that induce perfect matchings.*

**Proof:** Let  $f$  be an edge-colouring total 2-labelling of a cubic graph  $G$ . For every vertex  $v \in V(G)$  the three edges incident with  $v$  must receive the weights 3, 4, 5, if  $f(v) = 1$ , and the weights 4, 5, 6, if  $f(v) = 2$ . The edges of weight 3 and weight 6 join two vertices with the same label.

If  $f(v) = 1$ , then the other endvertex of the edge of weight 5 incident with  $v$  has label 2. So there are at least as many vertices with label 2 as with label 1. Conversely, for  $f(v) = 2$  the edge of weight 4 incident to  $v$  has its other endvertex labelled 1. So there are at least as many vertices labelled 1 as with label 2. Together, there are equally many vertices labelled 1 and 2 and the edges of weights 4 and 5 form a 2-regular graph joining vertices of label 1 to vertices of label 2. Therefore,  $G$  has the indicated structure.

Conversely, if  $G$  has the indicated structure, then  $|A| = |B|$ . We assign label 1 to the vertices and edges in  $A$  and label 2 to the vertices and edges in  $B$ . Labelling the edges of the 2-regular bipartite graph between  $A$  and  $B$  by 1 and 2 according to a proper 2-edge colouring results in an edge-colouring total 2-labelling.  $\square$

It is an easy observation that the lower bound is tight for forests.

**Theorem 2.2** *If  $F$  is a forest of maximum degree  $\Delta$ , then  $\chi'_t(F) = \lceil \frac{\Delta+1}{2} \rceil$ .*

**Proof:** We prove the stronger statement that an edge-colouring total labelling exists using only two vertex labels 1 and  $k = \lceil \frac{\Delta+1}{2} \rceil$ . Obviously, it suffices to prove the statement for the tree components.

We proceed by induction on the number of vertices  $n$ . The statement is true for  $n \leq 2$  so assume  $n \geq 3$ . Let  $vw$  be an edge such that  $v$  has degree at least 2 and all neighbours of  $v$  except possibly  $w$  are leaves. Note that such an edge  $vw$  exists. Delete all neighbours of  $v$  except  $w$  to obtain a tree  $T'$ , which by induction has the required total labelling. Now label the deleted vertices with 1 and  $k$  such that at most  $\frac{d(v)+1}{2}$  of the neighbours of  $v$  (including the already labelled vertex  $w$ ) have the same label. Now the remaining edges can be easily labelled such that all edges incident with  $v$  have different weights.  $\square$

Next, we consider edge-colouring total labellings of complete graphs. In a graph  $G$  with a given edge colouring a *rainbow (perfect) matching* is a (perfect) matching, where all edges are of different colour. We need a lemma on rainbow matchings in the proof of our next result.

**Lemma 2.3** *(a) Every complete bipartite graph  $K_{k,k}$  has a proper  $k$ -edge colouring with a rainbow perfect matching if  $k$  is odd, and a rainbow matching of cardinality  $k - 1$  if  $k$  is even.*

*(b) Every complete graph  $K_{2k}$  of even order has a proper  $(2k - 1)$ -edge colouring with a rainbow perfect matching unless  $k = 2$ .*

**Proof:**

- (a) Let  $u_1, \dots, u_k$  and  $w_1, \dots, w_k$  be the vertices on both sides of the bipartition. Define a proper edge colouring of  $G$  by assigning the colour  $\ell \in \{1, \dots, k\}$  to the edge  $u_i w_j$ , if  $j - i \equiv \ell \pmod k$ . Now let  $a$  and  $b$  be the largest even and odd integer  $< \frac{k}{2} + 1$ , respectively. Choose a matching  $M$  consisting of the edges  $u_i w_{a+1-i}$  for  $1 \leq i \leq a$  and  $u_{a+i} w_{a+b+1-i}$  for  $1 \leq i \leq b$ . This is a rainbow matching of cardinality  $a + b = k - 1$ , if  $k$  is even and a rainbow perfect matching, if  $k$  is odd.

- (b) Let  $u_0, u_1, \dots, u_{2k-1}$  be the vertices of  $K_{2k}$ . First assume that  $k$  is odd. Take as the first colour class of edges the perfect matching  $M_0$  consisting of the edges  $u_i u_{2k-i-1}$  for  $0 \leq i \leq k-1$ . The remaining colour classes are obtained as follows: Embed the vertices of  $K_{2k}$  in the plane such that  $u_1, u_2, \dots, u_{2k-1}$  form the vertices of a regular  $(2k-1)$ -gon with center  $u_0$ . Rotating  $M_0$  by an angle of  $\frac{2\pi}{2k-1}$  a total number of  $2k-2$  times defines  $2k-2$  further perfect matchings (cf. Figure 1). Since the geometric lengths of all edges in one matching are different, this defines a proper edge-colouring of  $K_{2k}$  for which the matching  $u_0 u_{2k-1}, u_1 u_2, \dots, u_{2k-3} u_{2k-2}$  is a rainbow perfect matching.

Next, assume that  $k$  is even. Here we choose as the first colour class of edges the perfect matching  $M_0$  consisting of the edges  $u_i u_{2k-i-1}$  for  $0 \leq i < \frac{k}{4}$ ,  $u_i u_{2k-i-2}$  for  $\frac{k}{4} \leq i < \frac{3}{4}k-1$ ,  $u_i u_{2k-i-3}$  for  $\frac{3}{4}k-1 \leq i < k-1$ , and the additional edge  $u_i u_{i+\frac{k}{2}}$  for  $i = \lfloor \frac{5}{4}k \rfloor - 1$ . Again, the remaining  $2k-2$  colour classes are obtained by embedding the vertices of  $K_{2k}$  and rotating  $M_0$  as before (cf. Figure 1). Again the matching  $u_0 u_{2k-1}, u_1 u_2, \dots, u_{2k-3} u_{2k-2}$  is a rainbow perfect matching.

□




Figure 1

In a graph with a total labelling we denote the set of vertices with label  $i$  by  $V_i$ .

**Theorem 2.4** *If  $n \not\equiv 2 \pmod{4}$ , then  $\chi'_t(K_n) = \lceil \frac{n}{2} \rceil$  and if  $n \equiv 2 \pmod{4}$ , then  $\chi'_t(K_n) \leq \frac{n}{2} + 1$ .*

*Proof:* In view of the lower bound it suffices to describe suitable labellings of the complete graph.

First assume that  $n \equiv 0 \pmod{4}$ . Label half the vertices by 1 and the other half by  $k = \frac{n}{2}$ . Determine a proper edge colouring of the edges in  $V_1$  with labels  $1, \dots, k-1$ , a proper edge colouring of the edges in  $V_k$  with labels  $2, \dots, k$ , and a proper edge colouring of the edges joining  $V_1$  to  $V_k$  with labels  $1, \dots, k$ . It is now easy to verify, that this is an edge-colouring total  $k$ -labelling. Note that this also implies the result for  $n \equiv 3 \pmod{4}$  and  $n \equiv 2 \pmod{4}$  by considering edge-colouring total labellings of complete graphs of order  $n+1$  and  $n+2$ , respectively.

Therefore, only the case  $n \equiv 1 \pmod{4}$  remains.

Label  $\frac{n-1}{2} = k-1$  vertices by label 1 and  $k$ , respectively, and the remaining vertex  $v$  by  $\frac{k+1}{2}$ . Let  $u_{\frac{k+1}{2}+2}, \dots, u_{\frac{k+1}{2}+k}$  denote the vertices of  $V_1$  and  $w_{\frac{k+1}{2}+k+2}, \dots, w_{\frac{k+1}{2}+2k}$  be the vertices of  $V_k$ . Label the edges from  $v$  to  $u_i$  with  $i - \frac{k+1}{2} - 1$  and the edges  $v$  to  $w_j$  with  $j - \frac{k+1}{2} - k$ . Note that each vertex  $u_i$  and  $w_j$  is joined to  $v$  by an edge of weight  $i$  and  $j$ , respectively.

It remains to show that we can find an edge labelling of the edges not incident with  $v$ , such that the labels form a proper edge colouring of the remaining graph and the weight of each edge is different from the indices of its endvertices. The edges inside  $V_1$  will obtain the weights  $3, \dots, k+1$ , inside  $V_k$  the weights  $2k+2, \dots, 3k$ , and the edges between  $V_1$  and  $V_k$  will obtain weights  $k+2, \dots, 2k+1$ .

By Lemma 2.3 (b) we know that the complete graph  $K_{k-1}$  induced by  $V_1$  has a proper  $(k-2)$ -edge colouring  $c$  which has a rainbow perfect matching. Let  $\{2, \dots, k-1\}$  be the colours of the colouring and let  $\{\frac{k-1}{2}+1, \dots, k-1\}$  be the colours occurring in the rainbow perfect matching  $M$ . Assign the indices in such a way that the vertex  $u_i$  of index  $i$  is incident with the edge of colour  $i-2$  in the rainbow matching for  $\frac{k+1}{2}+2 \leq i \leq k+1$ . Finally, recolour the edges of the rainbow perfect matching  $M$  with colour 1 and take the colours of this new colouring  $c'$  as the labels of the edges inside  $V_1$ . Note that this edge labelling has the desired property that  $u_i$  is not joined to a vertex in  $V_1$  by an edge of weight  $i$ . Along the same line of argument we obtain a labelling of the edges inside  $V_k$  with labels  $\{2, \dots, k\}$  such that each vertex  $w_j$  is not joined to a vertex in  $V_k$  by an edge of weight  $j$ .

Finally, we need to label the edges in the bipartite graph spanned by the  $(V_1, V_k)$ -edges. This graph is isomorphic to  $K_{k-1, k-1}$ , where  $k-1 \equiv 0 \pmod{2}$ . By Lemma 2.3 (a) this graph has a proper  $(k-1)$ -edge colouring using the colours  $\{1, \dots, k-1\}$  with a rainbow matching  $M$  of cardinality  $k-2$  that avoids the colour  $\frac{k-1}{2}$ . Assign the indices in such a way that  $u_i$  is incident with the edge of  $M$  of weight  $i-k-1$  for  $k \leq i \leq \frac{k+1}{2}+k$ , and  $w_j$  is incident with the edge of  $M$  of weight  $j-k-1$  for  $\frac{k+1}{2}+k+2 \leq j \leq 2k$ . Moreover, let  $w_{2k+1}$  be the vertex in  $V_k$  that is not incident with an edge of  $M$ . Now recolour the edges of  $M$  with colour  $k$  to obtain a new colouring, which we use as the labelling of the  $(V_1, V_k)$  edges. By the construction it is easy to verify that the result is an edge-colouring total  $k$ -edge labelling.  $\square$

We conclude this section with some further results concerning the case  $n \equiv 2 \pmod{4}$  which might eventually allow to determine for which  $n \equiv 2 \pmod{4}$ ,  $\chi'_t(K_n) = \frac{\Delta+1}{2}$  holds, and for which  $\chi'_t(K_n) = \frac{\Delta+1}{2} + 1$ . We can show that the second equality holds for  $6 \leq n \leq 22$ . At the same time our result describes the distribution of the labels in some detail if the first equality holds.

**Lemma 2.5** *Let  $K_n$  be a complete graph with  $k = \chi'_t(K_n) = \frac{n}{2}$ . If  $V_i$  denotes the set of vertices labelled  $i$  in an edge-colouring total  $k$ -labelling of  $K_n$ , then the cardinality of each set  $V_i$  is even,  $|V_i| = |V_{k-i+1}|$  and  $|V_i| \leq |V_1| = |V_k|$  for  $i = 1, \dots, k$ . The edges of weight*

$k + 2$  have label 1 and the edges of weight  $2k + 1$  have label  $k$ . Moreover, if  $n \equiv 2 \pmod{4}$  then  $6 \leq |V_{\frac{k+1}{2}}| \equiv 2 \pmod{4}$ .

**Proof:** Since  $k = \frac{\Delta+1}{2}$ , each vertex  $v \in V_i$  is incident with an edge of weight  $i + \ell$  for  $2 \leq \ell \leq 2k$ . For  $2 \leq i \leq k$  the edges of weight  $i + 2$  form a matching between  $V_i$  and  $V_1$  and hence  $|V_i| \leq |V_1|$ . Similarly, each vertex  $v \in V_j$  is incident with an edge of weight  $j + 2k$  and for  $1 \leq j \leq k - 1$  these edges form a matching between  $V_j$  and  $V_k$ , implying  $|V_j| \leq |V_k|$ . Since the inequalities hold for  $i = k$  and  $j = 1$ , we obtain  $|V_1| = |V_k|$ .

Next we show that each of the sets  $V_i$  has even cardinality. This is true for  $V_1$ , since the edges of weight 3 form a perfect matching between the vertices in  $V_1$ . Now consider the vertex set  $U_i = V_1 \cup V_2 \cup \dots \cup V_i$ . Since the edges of weight  $i + 2$  form a perfect matching of  $U_i$ , and, by induction,  $U_{i-1}$  has even cardinality, the set  $V_i$  has even cardinality as well.

For  $i \leq \frac{k+1}{2}$  we prove by induction over  $i$  that the edges of weight  $2k + 1$  incident to a vertex in  $V_i$  have their other endvertex in  $V_{k-i+1}$ , and the edges of weight  $k + 2$  incident to a vertex in  $V_{k-i+1}$  have their other endvertex in  $V_i$ . In particular,  $|V_i| = |V_{k-i+1}|$  and the edges of weight  $k + 2$  and  $2k + 1$  have weight 1 and  $k$ , respectively.

The statement is true for  $i = 1$ , so assume that it is true for all indices  $< i$ . Let  $vw$  be the edge of weight  $2k + 1$  that is incident to  $v \in V_i$ . Since the label of  $vw$  is at most  $k$ , the vertex  $w$  has label  $s \geq k - i + 1$ . If  $s > k - i + 1$ , then by induction the other endvertex  $v$  of the edge of weight  $2k + 1$  incident to  $w$  has label  $t = k - s + 1 < i$ , contradicting  $v \in V_i$ . Analogously, for the vertices of  $V_{k-i+1}$  the other endvertex of the incident edge of weight  $k + 2$  lies in  $V_i$ . This completes the induction. If  $n \equiv 2 \pmod{4}$ , then  $6 \leq |V_{\frac{k+1}{2}}| \equiv 2 \pmod{4}$ , because of the parity conditions and since  $V_{\frac{k+1}{2}}$  has two disjoint perfect matchings consisting of the edges of weight  $k + 2$  and  $2k + 1$ .  $\square$

**Lemma 2.6** *Every edge-colouring total  $(2p + 1)$ -labelling of  $K_{4p+2}$  for  $p \geq 1$  uses at least 5 different vertex labels.*

**Proof:** For contradiction, we assume the existence of an edge-colouring total  $(2p + 1)$ -labelling using less than 5 different vertex labels. By Lemma 2.5, this implies that it has exactly 3 label classes  $V_1$ ,  $V_{p+1}$ , and  $V_{2p+1}$ . Moreover  $|V_{2p+1}| = |V_1| \geq |V_{p+1}| \geq 6$ . We know that all edges with weights  $3, \dots, p + 2$  have both endvertices in  $V_1$  and for each such weight value these edges form a perfect matching in  $V_1$ . Furthermore, all edges of weight  $p + 3, \dots, 2p + 2$  incident with a vertex in  $V_{p+1}$  have the other endvertex in  $V_1$ , and, finally, there is a perfect matching between  $V_{2p+1}$  and  $V_1$  of edges of weight  $2p + 3$ .

Let  $n_1$  be the number of vertices in  $V_1$  and  $n_{p+1}$  the number of vertices in  $V_{p+1}$ . Since we have  $n_1 = \frac{n-n_{p+1}}{2}$  for  $n = 4p + 2$ , there are exactly  $n_1/2$  edges in  $V_1$  of weight  $w$  for each  $3 \leq w \leq p + 2$  and  $\frac{n_1-n_{p+1}}{2}$  edges in  $V_1$  of weight  $w$  for each  $p + 3 \leq w \leq 2p + 2$  and no edges of weight  $\geq 2p + 3$ . Altogether, there are at most

$$p \left( \frac{n_1}{2} \right) + p \left( \frac{n_1 - n_{p+1}}{2} \right) = p \left( n_1 - \frac{n_{p+1}}{2} \right)$$



edges in  $V_1$ . Since  $2n_1 + n_{p+1} = n = 4p + 2$  we get  $p = \frac{1}{2}(n_1 + \frac{n_{p+1}}{2} - 1)$  and

$$\binom{n_1}{2} \leq \binom{n_1}{2} - \frac{1}{2}n_{p+1}^2 + \frac{1}{2}(n_1 - n_1 + 1)n_{p+1},$$

which is a contradiction since  $n_{p+1} \geq 6 > 1$ .  $\square$

### 3 The general upper bound

Our goal in this section is to prove Theorem 1.1. In order to clarify our approach, we present a number of intermediate results, some of which we think to be interesting on their own right. The first is a consequence of Vizing's Adjacency Lemma [21] (see also [13]). A graph  $G = (V, E)$  of maximum degree  $\Delta$  is called *critical* if  $\chi'(G) = \Delta + 1$  but  $\chi'(G - e) = \Delta$  for all  $e \in E$ .

**Lemma 3.1 (Vizing's Adjacency Lemma [21])** *Let  $G = (V, E)$  be a critical graph with maximum degree  $\Delta$  and  $\chi'(G) = \Delta + 1$ . If  $uv \in E$  then  $u$  is adjacent to at least  $\max\{2, \Delta - d_G(v) + 1\}$  many vertices of maximum degree.*

**Proposition 3.2** *Every graph  $G$  with maximum degree  $\Delta$  has a proper  $(\Delta + 1)$ -edge colouring such that no edge of colour  $\Delta + 1$  is incident with a vertex of degree less than  $\Delta$ .*

**Proof:** We apply induction on  $m := |E(G)|$ . If  $G$  has a proper  $\Delta$ -edge colouring, then the statement is vacuously true. Note that this already implies the result for  $m \leq 2$ . Therefore, we assume now that  $m \geq 3$  and that  $\chi'(G) = \Delta + 1$ .

It follows from Lemma 3.1 applied to a critical subgraph of  $G$  and a vertex  $u$  of maximum degree, that a neighbour  $w$  of  $u$  has maximum degree as well. By induction,  $G - uw$  has a proper  $(\Delta + 1)$ -edge colouring such that no edge of colour  $\Delta + 1$  is incident to a vertex of degree less than  $\Delta$ . Therefore, assigning the colour  $\Delta + 1$  to the edge  $uw$  yields the desired colouring.  $\square$

The construction in the next result relies on a suitable partition of the vertex set.

**Theorem 3.3** *If  $G$  is a graph of maximum degree  $\Delta$  whose vertex set has a partition  $V(G) = A \cup B$  such that every vertex has at most  $k - 1$  neighbours in  $A$  and at most  $k - 1$  neighbours in  $B$  for some  $k$  with  $k - 1 > \frac{\Delta}{2}$ , then  $\chi'_t(G) \leq k$ .*

**Proof:** Let  $V(G) = A \cup B$  be a partition as in the statement. Label the vertices of  $A$  with 1 and the vertices of  $B$  with  $k$ .

By Proposition 3.2,  $G[A]$  has a proper  $k$ -edge colouring that avoids colour  $k$  at the vertices of degree  $d_{G[A]}(v) < k - 1$ . Similarly,  $G[B]$  has a proper  $k$ -edge colouring that avoids colour 1 at the vertices of degree  $d_{G[B]}(v) < k - 1$ . We choose these edge colourings as the labellings of the edges in  $A$  and  $B$ , respectively. Let  $A'$  denote the set of vertices in

$A$  incident with an edge labelled  $k$  and let  $B'$  denote the set of vertices in  $B$  incident with an edge labelled 1.

It remains to label the edges between  $A$  and  $B$ . Let  $G(A, B)$  denote the bipartite spanning subgraph of  $G$  of maximum degree at most  $k - 1$  containing all edges between  $A$  and  $B$ . Considering a perfect matching in a bipartite  $(k - 1)$ -regular supergraph of  $G(A, B)$ , it follows that  $G(A, B)$  has a minimal matching  $M$  that saturates all vertices  $v$  with  $d_{G(A, B)}(v) = k - 1$ . Note that by the minimality requirement,  $M$  does not contain an  $(A', B')$ -edge, since for each vertex in  $u \in A' \cup B'$  we have  $d_{G(A, B)}(u) \leq \Delta(G) - (k - 1) < k - 1$ . We label the edges of  $M$  with one endvertex in  $A'$  with  $k$  and the remaining edges with 1. Now  $G(A, B) - M$  has maximum degree  $\leq k - 2$  and hence has a proper  $(k - 2)$ -edge colouring with colours  $2, 3, \dots, k - 1$  which we use as the labelling for the edges. It is easy to verify that the edge weights defined by this total  $k$ -labelling form a proper edge colouring of  $G$ .  $\square$

Our next goal is to find a partition as in Theorem 3.3 for some  $k$  close to  $\Delta/2$ . We do this using the probabilistic method via a discrepancy argument: For a graph  $G$  we consider the *discrepancy*  $\text{disc}(G)$  defined as follows:

$$\text{disc}(G) := \min_{g: V(G) \rightarrow \{-1, 1\}} \max_{u \in V(G)} \left| \sum_{v \in N_G(u)} g(v) \right|.$$

Note that  $\text{disc}(G)$  corresponds to the ordinary discrepancy of the hypergraph on the ground set  $V(G)$  whose hyperedges are the neighbourhoods of vertices in  $G$ .

Together with Theorem 3.3 we obtain.

**Corollary 3.4** *If  $G$  is a graph of maximum degree  $\Delta$ , then*

$$\chi_t(G) \leq \frac{\Delta + \text{disc}(G)}{2} + 1.$$

**Proof:** Let  $g : V(G) \rightarrow \{-1, 1\}$  be such that  $\text{disc}(G) = \max_{u \in V(G)} \left| \sum_{v \in N_G(u)} g(v) \right|$ . Let  $A = g^{-1}(1)$  and  $B = g^{-1}(-1)$ . For  $u \in V(G)$  let  $d_A(u) = |\{v \in N_G(u) \mid g(v) = 1\}|$  and  $d_B(u) = |\{v \in N_G(u) \mid g(v) = -1\}|$ . Since  $|d_A(u) - d_B(u)| \leq \text{disc}(G)$  and  $d_A(u) + d_B(u) \leq \Delta$ , we have  $\max\{d_A(u), d_B(u)\} \leq \frac{\Delta + \text{disc}(G)}{2}$  for every  $u \in V(G)$  and Theorem 3.3 implies the desired result.  $\square$

In order to bound the discrepancy we combine Chernoff's inequality with the Lovász Local Lemma.

**Lemma 3.5 (Chernoff's inequality [11], see also [5])** *Let  $X_1, \dots, X_n$  be mutually independent random variables with  $\mathbf{P}(X_i = 1) = \mathbf{P}(X_i = -1) = \frac{1}{2}$ . Then for  $S = X_1 + \dots + X_n$  and  $\delta > 0$  we get  $\mathbf{P}(|S| > \delta) < 2 \exp\left(-\frac{\delta^2}{2n}\right)$ .*

**Lemma 3.6 (Lovász Local Lemma [12], see also [5])** *Let  $A_1, A_2, \dots, A_n$  be events in an arbitrary probability space. Let  $\mathbf{P}(A_i) \leq p$  and let  $A_i$  be mutually independent of all but at most  $d \geq 2$  of the events  $A_j$  with  $j \neq i$  for each  $1 \leq i \leq n$ . If  $ep(d+1) \leq 1$ , then  $\mathbf{P}(\bigwedge_{i=1}^n \overline{A_i}) > 0$ , i.e. with positive probability none of the events  $A_i$  occurs.*

**Proposition 3.7** *If  $G$  is a graph of maximum degree  $\Delta$ , then*

$$\text{disc}(G) \leq \left\lfloor \sqrt{2\Delta(1 + \ln(2\Delta^2 - 2\Delta + 2))} \right\rfloor.$$

**Proof:** We consider a random function  $g : V(G) \rightarrow \{-1, 1\}$  where all values  $g(v)$  are 1 independently at random with probability  $1/2$ .

For some  $\delta > 0$  and  $u \in V(G)$  consider the event  $A_u$ :  $\left| \sum_{v \in N_G(u)} g(v) \right| > \delta$ . By Chernoff's inequality,

$$\mathbf{P}(A_u) \leq 2 \exp\left(\frac{-\delta^2}{2d_G(u)}\right) \leq 2 \exp\left(\frac{-\delta^2}{2\Delta}\right).$$

The events  $A_u$  and  $A_v$  are dependent only if there is a path of length exactly two between  $u$  and  $v$ . Therefore,  $A_u$  is independent of all but at most  $\Delta(\Delta - 1)$  many events  $A_v$  with  $v \neq u$ . For  $\delta := \sqrt{2\Delta(1 + \ln(2\Delta^2 - 2\Delta + 2))}$  we obtain

$$2 \exp\left(1 - \frac{\delta^2}{2\Delta}\right) (\Delta(\Delta - 1) + 1) = 1$$

and the Lovász Local Lemma implies the existence of a function  $g : V(G) \rightarrow \{-1, 1\}$  with  $\left| \sum_{v \in N_G(u)} g(v) \right| \leq \delta$  for all  $u \in V(G)$ .  $\square$

**Proof of Theorem 1.1:** The result follows immediately from Corollary 3.4 and Proposition 3.7.  $\square$

## 4 Concluding remarks

The upper bound  $\mathcal{O}(\sqrt{\Delta \log \Delta})$  for the discrepancy of a  $\Delta$ -regular graph is not far from being best possible. This is due to the fact, that there are graphs with discrepancy  $\Omega(\sqrt{\Delta})$ . The Paley graphs for example form an infinite sequence of graphs with  $\Delta = \frac{n-1}{2}$  and discrepancy  $\Omega(\sqrt{\Delta})$  by a result of Lovász and Sós (see [17, Theorem 4.5]). Conversely, the Beck-Fiala Conjecture (see [17]) says that the vertices of every hypergraph where each vertex belongs to at most  $\Delta$  hyperedges has discrepancy  $\mathcal{O}(\sqrt{\Delta})$ . If the Beck-Fiala Conjecture — or its restriction to  $\Delta$ -regular,  $\Delta$ -uniform hypergraphs — is true then we can improve the upper bound in Theorem 1.1 with the same reasoning to

$$\chi'_t(G) \leq \frac{\Delta + 1}{2} + \mathcal{O}(\sqrt{\Delta}). \tag{1}$$

On the other hand, if, like in most of our explicit labellings, the typical total  $k$ -labellings use on the vertices almost only the labels 1 and  $k$ , then the reduced upper bound in the formula above is tight in view of the Paley graphs.

So the main open question in this context might be the following:

**Problem 4.1** *Is there a constant  $K$  with*

$$\chi'_t(G) \leq \frac{\Delta + 1}{2} + K \quad (2)$$

*for all graphs  $G$  of maximum degree  $\Delta$ ?*

Surely there are further options except (1) and (2). Indications could be obtained from an answer to the question whether all graphs  $G$  have an edge-colouring total  $\chi'_t(G)$ -labelling with only few vertex labels.

In view of the graphs where we know the exact value of  $\chi'_t(G)$ , the constant  $K$  must be at least 1. With  $K = 1$  the bound (2) is attained with equality e.g. for cubic snarks and  $K_{4k+2}$  for  $1 \leq k \leq 5$ . For even  $\Delta$  we are not aware of any graph with  $\chi'_t(G) > \lceil \frac{\Delta+1}{2} \rceil$ . One first question in this direction is whether  $\chi'_t(G) = 3$  for all graphs with  $\Delta = 4$ . As a potential candidate for the general problem we checked the unique Paley graph on 17 vertices ( $\Delta = 8$ ), which is at the same time the  $(4, 4)$ -Ramsey graph, with a computer program, that came up with an edge-colouring total 5-labelling.

## Acknowledgement

We thank Friedrich Regen, Bonn, for writing this computer program.

## References

- [1] L. Addario-Berry, R.E.L. Aldred, K. Dalal, and B. Reed, Vertex colouring edge partitions, *J. Comb. Theory, Ser. B* **94** (2005), 237-244.
- [2] L. Addario-Berry, K. Dalal, and B. Reed, Degree constrained subgraphs, Proc. GRACO 2005, volume 19 of *Electron. Notes Discrete Math.*, Amsterdam (2005), 257-263 (electronic), Elsevier.
- [3] L. Addario-Berry, K. Dalal, C. McDiarmid, B. Reed, and A. Thomason, Vertex-Colouring Edge-Weightings, *Combinatorica* **27** (2007), 1-12.
- [4] M. Aigner and E. Triesch, Irregular assignments of trees and forests, *SIAM J. Discrete Math.* **3** (1990), 439-449.
- [5] N. Alon and J.H. Spencer, The probabilistic method (2nd ed.), Wiley, 2000.
- [6] M. Bača, S. Jendrol', M. Miller, and J. Ryan, On irregular total labellings, *Discrete Math.* **307** (2007), 1378-1388.

- [7] M. Bača, S. Jendrol', and M. Miller, On total edge irregular labelings of trees, *manuscript* 2006.
- [8] S. Brandt, J. Miškuf, and D. Rautenbach, On a Conjecture about Edge Irregular Total Labellings, *manuscript* 2006.
- [9] S. Brandt, J. Miškuf, and D. Rautenbach, Edge irregular total labellings for graphs of linear size, *manuscript* 2007.
- [10] G. Chartrand, M.S. Jacobson, J. Lehel, O.R. Oellermann, S. Ruiz, and F. Saba, Irregular networks, *Congr. Numerantium* **64** (1988), 197-210.
- [11] H. Chernoff, A measure of the asymptotic efficiency for tests of a hypothesis based on the sum of observations, *Ann. Math. Stat.* **23** (1952), 493-509.
- [12] P. Erdős and L. Lovász, Problems and results on 3-chromatic hypergraphs and some related questions, in: Infinite and finite sets (A. Hajnal ed.), North Holland, 1975, 609-628.
- [13] S. Fiorini and R. J. Wilson, Edge-colorings of graphs, in: Selected Topics in Graph Theory (L. W. Beineke, R. J. Wilson eds.), Academic Press, 1978, 103-126.
- [14] A. Frieze, R.J. Gould, M. Karoński, and F. Pfender, On graph irregularity strength, *J. Graph Theory* **41** (2002), 120-137.
- [15] J. Ivančo and S. Jendrol', Total edge irregularity strength of trees, *Discussiones Mathematicae Graph Theory* **26** (2006), 449-456.
- [16] M. Karoński, T. Łuczak and A. Thomason, Edge weights and vertex colours, *J. Comb. Theory, Ser. B* **91** (2004), 151-157 (2004).
- [17] J. Matoušek, Geometric Discrepancy, Springer, 1999.
- [18] T. Nierhoff, A tight bound on the irregularity strength of graphs, *SIAM J. Discrete Math.* **13** (1998), 313-323.
- [19] J. Przybyło and M. Woźniak, 1, 2 Conjecture, II, Preprint 2007.  
<http://www.ii.uj.edu.pl/preMD/MD26.pdf>
- [20] V.G. Vizing, On an estimate of the chromatic class of a p-graph (in russian), *Diskret. Analiz.* **3** (1964), 25-30.
- [21] V.G. Vizing, Critical graphs with given chromatic class (in russian), *Diskret. Analiz.* **5** (1965), 9-17.