

Selective Hypergraph Colourings

Yair Caro

Department of Mathematics
University of Haifa-Oranim
Israel

Josef Lauri

Department of Mathematics
University of Malta
Malta

Christina Zarb

Department of Mathematics
University of Malta
Malta

Abstract

We look at colourings of r -uniform hypergraphs, focusing our attention on unique colourability and gaps in the chromatic spectrum. The pattern of an edge E in an r -uniform hypergraph H whose vertices are coloured is the partition of r induced by the colour classes of the vertices in E . Let Q be a set of partitions of r . A Q -colouring of H is a colouring of its vertices such that only patterns appearing in Q are allowed. We first show that many known hypergraph colouring problems, including Ramsey theory, can be stated in the language of Q -colourings. Then, using as our main tools the notions of Q -colourings and Σ -hypergraphs, we define and prove a result on tight colourings, which is a strengthening of the notion of unique colourability. Σ -hypergraphs are a natural generalisation of σ -hypergraphs introduced by the first two authors in an earlier paper. We also show that there exist Σ -hypergraphs with arbitrarily large Q -chromatic number and chromatic number but with bounded clique number. Dvorak et al. have characterised those Q which can lead to a hypergraph with a gap in its Q -spectrum. We give a short direct proof of the necessity of their condition on Q . We also prove a partial converse for the special case of Σ -hypergraphs. Finally, we show that, for at least one family Q which is known to yield hypergraphs with gaps, there exist no Σ -hypergraphs with gaps in their Q -spectrum.

1 Introduction

Let $V = \{v_1, v_2, \dots, v_n\}$ be a finite set, and let $E = \{E_1, E_2, \dots, E_m\}$ be a family of subsets of V . The pair $H = (V, E)$ is called a *hypergraph* with vertex-set $V(H) = V$, and with edge-set $E(H) = E$. When all the subsets are of the same size r , we say that H is an *r -uniform hypergraph*.

Hypergraph vertex colourings is a widely studied topic, and several different types of colourings have been defined, such as classical colourings, where monochromatic edges are not allowed [2], Voloshin colourings [19], non monochromatic non-rainbow (NMNR) colourings [5] and constrained colouring [3, 7]. In

the latter three, one of the most interesting phenomenons studied is the existence of a *gap* in the *chromatic spectrum* of the hypergraph, which do not occur in classical colourings of neither graphs nor hypergraphs. The *chromatic spectrum* of a hypergraph H is the sequence, in increasing order, of all k such that H has a k -colouring which satisfies the type of colouring being considered. We say that the chromatic spectrum has a *gap* when there exist integers $k_1 < k_2 < k_3$ such that the hypergraph is k_1 - and k_3 -colourable but not k_2 -colourable.

In [5] and [7], the authors have studied the existence, or not, of gaps in the chromatic spectrum for NMNR colourings and constrained colourings in particular, using a construction defined in [5], the σ -hypergraph. A σ -hypergraph $H(n, r, q \mid \sigma)$, where σ is a partition of r , is an r -uniform hypergraph having nq vertices partitioned into n *classes* of q vertices each. If the classes are denoted by V_1, V_2, \dots, V_n , then a subset K of $V(H)$ of size r is an edge if the partition of r formed by the non-zero cardinalities $|K \cap V_i|$, $1 \leq i \leq n$, is σ . The non-empty intersections $K \cap V_i$ are called the parts of K . The number of parts of the partition σ is denoted by $s(\sigma)$, while the size of the largest and smallest parts of the partition σ are denoted by $\Delta = \Delta(\sigma)$ and $\delta = \delta(\sigma)$, respectively. Several interesting results emerged about the chromatic spectrum of σ -hypergraphs in conjunction with these two types of colourings

For the purpose of this paper we now extend this definition of a σ -hypergraph to the more general notion of a Σ -hypergraph. Let $P(r)$ denote the set of all partitions of r . A Σ -hypergraph $H(n, r, q \mid \Sigma)$, where $\Sigma \subseteq P(r)$, is an r -uniform hypergraph having nq vertices partitioned into n *classes* of q vertices each. If the classes are denoted by V_1, V_2, \dots, V_n , then a subset K of $V(H)$ of size r is an edge if the partition σ of r formed by the non-zero cardinalities $|K \cap V_i|$, $1 \leq i \leq n$, is in Σ . We call σ the *edge type* of the edge E , denoted by $\text{type}(E)$, and Σ is the set of allowable edge-types in $H(n, r, q \mid \Sigma)$.

In this paper we describe L -colourings and Q -colourings of r -uniform hypergraphs, which can encompass many different types of hypergraph colourings described in the literature. Such ideas, originated in Voloshin's seminal work in [13], were more explicit in [16], in the context of the colouring of Steiner systems for $r = 3$ and $r = 4$, and further studied in [11, 17], and studied in much more generality for oriented graphs in [14]. We define L - and Q -colourings as follows.

Let H be an r -uniform hypergraph, $r \geq 2$ and consider $E \in E(H)$. Then a colouring of the vertices of E induces a partition π of r whose parts are the numbers of vertices of each colour appearing in E . This partition is called the *colour pattern* of E and is written as $\text{pat}(E) = (n_1, n_2, \dots, n_k)$, where $n_1 \geq n_2 \geq \dots \geq n_k \geq 1$, where $\sum_{i=1}^k n_i = r$.

For any edge $E \in E(H)$, we assign $Q(E) \subseteq P(r)$. We say that a colouring of the vertices of H is an L -colouring, where $L = \{Q(E_i) : i = 1, \dots, |E(H)|\}$, if $\forall E_i \in E(H)$, $\text{pat}(E_i) \in Q(E_i)$. In the case when all the edges are assigned the same family of partitions Q , that is $Q(E_i) = Q$, $\forall E_i \in E(H)$, we call this a Q -colouring. We define $\Delta(Q) = \max\{\Delta(\pi) : \pi \in Q\}$ and let $s(Q) = \max\{s(\pi) : \pi \in Q\}$.

The Q -spectrum of H is the sequence, in increasing order, of all k such that H has a k - Q -colouring. The *lower chromatic number* χ_Q is defined as the least number k for which H has a k - Q -colouring. Similarly, the *upper chromatic*

number $\bar{\chi}_Q$ is the largest k for which H has a k - Q -colouring. Clearly, the first and last terms of this sequence are χ_Q and $\bar{\chi}_Q$ respectively. As described previously, we say that the Q -spectrum has a *gap*, or is *broken*, when there exist integers $k_1 < k_2 < k_3$ such that the hypergraph is k_1 - Q - and k_3 - Q -colourable but not k_2 - Q -colourable. We denote the Q -spectrum of a hypergraph H by $\text{Spec}_Q(H)$.

This paper is structured as follows. We first use the “language” of L - and Q -colourings to describe the different types of colourings covered by this concept. In Section 2 we introduce, motivated by the notion of uniquely-colourable graphs and hypergraphs [18], the concept of *tight Q -colourings* of hypergraphs and give some interesting results involving Σ -hypergraphs. In Section 3, we then consider the clique number of Σ -hypergraphs and present sparse r -uniform Σ -hypergraphs with bounded clique number and arbitrary large chromatic number, extending results from [6]. Finally, in Section 4 we concentrate on conditions on Q for the existence and non-existence of gaps in the Q -spectrum of r -uniform hypergraphs. These conditions have been completely characterised in [14] for oriented hypergraphs, with the results for non-oriented hypergraphs given as a corollary in Theorem 20 in [14]. Here we show that, in many cases, these conditions also guarantee the existence of a Σ -hypergraph with a broken Q -spectrum. But, we also show that Σ -hypergraphs cannot always provide examples with a broken Q -spectrum for all cases of Q for which it is known that a uniform hypergraph with a gap in its Q -spectrum does indeed exist.

Several types of colourings have been defined for hypergraphs. Here we look at the main types which have been studied, and describe them in terms of L - and Q -colourings. We also look at external colouring problems such as Ramsey Theory, and describe them also in terms of Q -colourings.

Two particularly important partitions of r will be used several times — the monochromatic partition (r) and the rainbow partition $(1, 1, \dots, 1)$. We use M and R to represent these partitions respectively.

1. For classical graph colourings, $Q = \{\pi \in P(r) : \pi = R\}$.
2. For, classical colourings of hypergraphs, $Q = \{\pi \in P(r) : \pi \neq M\}$. The chromatic number $\chi(H)$ is the smallest integer k for which a proper classical k -colouring of H exists. For any integer $p \geq \chi(H)$, there exists a proper classical p -colouring.
3. In Voloshin colourings of hypergraphs [19], or mixed hypergraphs, there exist two types of edges, D -edges and C -edges. A D -edge cannot be *monochromatic*, that is all vertices of the edge having the same colour, while a C -edge cannot be *polychromatic(rainbow)*, that is all vertices having a different colour. Hence for all D -edges, $Q_D = Q(E) = \{\pi \in P(r) : \pi \neq M\}$, while for all C -edges, $Q_C = Q(E) = \{\pi \in P(r) : \pi \neq R\}$, and $L = \{Q_C \cup Q_D\}$.
4. A non-monochromatic non-rainbow (NMNR) colouring, as discussed in [5], is a Q -colouring where $Q = \{\pi \in P(r) : \pi \notin \{M, R\}\}$. Such hypergraphs are often referred to as *bi-hypergraphs* and are a special type of Voloshin colourings.

5. An (α, β) -colouring of a hypergraph H , as described in [?], is the case where $Q = \{\pi \in P(r) : \pi = (n_1, n_2, \dots, n_k), \alpha \leq k \leq \beta\}$. This is based on the concept of *colour-bounded hypergraphs* first defined by Bujtas and Tuza in [3]. Observe that classical hypergraph colourings are $(2, r)$ -colourings, while NMNR-colourings are $(2, r - 1)$ -colourings.
6. Bujtas and Tuza defined another type of hypergraph colouring with further restrictions in [4]: a *stably bounded hypergraph* is a hypergraph together with four colour-bound functions which express restrictions on vertex colorings. Formally, an r -uniform stably bounded hypergraph is a six-tuple $H = (V(H), E(H), s, t, a, b)$, where s, t, a and b are positive integers, called *colour-bound functions*. We assume throughout that $1 \leq s \leq t \leq r$ and $1 \leq a \leq b \leq r$ hold. A proper vertex coloring of $H = (V(H), E(H), s, t, a, b)$ satisfies the following three conditions for every edge $E \in E(H)$.
 - The number of different colors assigned to the vertices of E is at least s and at most t .
 - There exists a colour assigned to at least a vertices of E .
 - Each color is assigned to at most b vertices of E .

Hence, such a colouring is a Q -colouring where

$$Q = \{\pi \in P(r) : \pi = (n_1, n_2, \dots, n_k), n_1 \geq n_2 \geq \dots \geq n_k, s \leq k \leq t, a \leq n_1 \leq b.\}$$

7. In [1], given a vertex coloring of a hypergraph, a vertex contained in an edge E is said to be uniquely colored in E , if its color is assigned to no other vertex in E . If every edge of a hypergraph contains a uniquely colored vertex, then the coloring is called *conflict-free*. In this case, $Q = \{\pi \in P(r) : \pi = (n_1, n_2, \dots, n_k), n_1 \geq n_2 \geq \dots \geq n_k = 1.\}$
8. We can also use the language of Q -colourings to express Ramsey's Theorem, which broadly states that one will find monochromatic cliques in any edge colouring of a sufficiently large complete graph. The theorem can also be extended to hypergraphs: for fixed integers r and k , and integers $n_1, \dots, n_k \geq r$, there exists an integer $R(r, n_1, \dots, n_k)$ such that if the edges of a complete r -uniform hypergraph on n vertices, $n \geq R(r, n_1, \dots, n_k)$ are coloured using k colours, then for some $1 \leq j \leq k$, there is a monochromatic complete r -uniform subhypergraph of order n_j and colour j .

Let us consider $K(n, r)$ to be a complete r -uniform hypergraph on n vertices. We define the hypergraph $H = H(n, r, p)$, where $p \geq r + 1$, such that the vertices of H are the edges of $K(n, r)$, so that $|V(H)| = \binom{n}{r}$. Now a collection of $\binom{p}{r}$ vertices in H is an edge if the corresponding edges in $K(n, r)$ are all taken from a set of p vertices in $K(n, r)$. So $|E(H)| = \binom{n}{p}$ and H is $\binom{p}{r}$ -uniform. We use this to express Ramsey's Theorem as follows:

Theorem 1.1. *For fixed k, r, p , there exists $N(k, r, p)$ such that for $n \geq N(k, r, p)$, if $Q \subseteq P(\binom{p}{r})$ and $M \notin Q$, then $H = H(n, r, p)$ is not k - Q -colourable.*

The definitions of L -colourings and Q -colourings encompasses all these different types of colourings as special cases, and allows for other types of colourings to be defined.

2 Tight colourings

We say that a hypergraph H is *tightly Q -colourable* if:

1. $\text{Spec}_Q(H) = \{k\}$
2. The k - Q -colouring is unique (up to exchanging colour classes)
3. The colour classes have equal size.
4. For any partition $\pi \in Q$, H is not $(Q \setminus \pi)$ -colourable.

Tight colouring is motivated by the concept of the so called uniquely colourable graphs [12]. While in a uniquely colourable graph, uniqueness is required only when the number of colours used is exactly $\chi(G)$ (but if we can use more than $\chi(G)$ colours, there is no restrictions on the colour class structure), in tight- Q colouring of hypergraphs, the restrictions are considerably stricter.

An important colouring that will be used frequently is the *canonical distinct monochromatic colouring* — in short CDMC — which is a colouring of the Σ -hypergraph $H(n, r, q | \Sigma)$ such that all vertices belonging to the same class are given the same colour (so each class is monochromatic), and vertices of distinct classes are given distinct colours (hence distinct). It is a simple but important fact that in a CDMC of $H(n, r, q | \Sigma)$, we have $\text{type}(E) = \text{pat}(E)$ for every edge $E \in E(H)$.

We are now ready to prove our main result on tight colouring.

Theorem 2.1. *Suppose $Q \subset P(r)$ such that $M, R \notin Q$. Then the Σ -hypergraph $H = H(2r, r, (r-1)^2 + 1 | \Sigma = Q)$ (on $(2r((r-1)^2 + 1))$ vertices) is tightly Q -colourable.*

Proof. Consider $H = H(2r, r, (r-1)^2 + 1 | \Sigma = Q)$. Clearly H is $2r$ - Q -colourable by the CMDC, with colour classes of equal size.

We first show that $\text{Spec}_Q(H) = \{2r\}$, and that H is uniquely $2r$ - Q -colourable.

Consider any k - Q -colouring of H . Let n_j be the number of distinct colours appearing in class V_j , for $j = 1 \dots 2r$. Let us consider those classes for which $n_j \geq r$. If there are at least r such classes, then we can choose any edge E of H with $\text{type}(E) = \sigma \in \Sigma$ to include r distinct colours, so that $\text{pat}(E) = R \notin Q$.

So there are at most $r-1$ such classes, and hence at least $r+1$ classes in which $n_j \leq r-1$. Let these classes be V_1 to V_{r+1} . Since $q = (r-1)^2 + 1$, it follows by the pigeon-hole principle that in each of the classes V_1 to V_{r+1} there

is a colour that appears at least r times. Let c_j be the colour which appears most frequently in class V_j for $j = 1 \dots r + 1$.

We show that the colours c_1, \dots, c_{r+1} must all be distinct and cannot appear in any other class. Suppose that colour c_j appears in another class among V_1 to V_{2r} , say in class V_t . Let σ be a partition such that $\Delta(\sigma) = \Delta(Q)$. Since $M \notin Q$, $\Delta(\sigma) = \Delta(Q) < r$. Let E be an edge with $\text{type}(E) = \sigma$. Let us choose the vertices of the Δ -part of E from the class V_j all of colour c_j , and we choose another part of E from class V_t to include the vertex with colour c_j , with the remaining parts chosen from any of the remaining classes. Then $\text{pat}(E)$ has $\Delta + 1$ vertices of the same colour, which is impossible by the maximality of $\Delta(Q)$. Hence c_1, \dots, c_{r+1} are all distinct and the colour c_j only appears in class V_j for $j = 1, \dots, r + 1$.

We now show that in fact, classes V_1 to V_{r+1} must be monochromatic of distinct colours. Suppose some class V_j , $1 \leq j \leq r + 1$, has two vertices x and y of distinct colours, one of which, say x , has colour c_j . Let σ be a partition such that $s(\sigma) = s(Q)$. Since $R \notin Q$, $s(\sigma) = s(Q) = s \leq r - 1$, and hence $\Delta(\sigma) \geq 2$. Also $s \geq 2$ since $M \notin Q$. We now take an edge E with $\text{type}(E) = \sigma$ such that the Δ part of E is chosen from V_j to include x and y . In the remaining classes from V_1 to V_{r+1} , there are at least another $r - 1$ distinct colours among the c_1 to c_{r+1} . So we can choose the remaining parts of σ with a distinct colour for each different to the colours of x and y . But then this edge includes $s(Q) + 1$ colours which is not possible by the maximality of $s(Q)$.

Finally, we show that all the class V_1 to V_{2r} must be monochromatic of distinct colours. Suppose V_j for $j > r + 1$, has two vertices, x and y of distinct colours. Observe that these colours cannot appear in classes V_1 to V_{r+1} as shown above. Let σ again be a partition such that $s(\sigma) = s(Q)$. Recall that, since $R \notin Q$, $s(\sigma) = s(Q) = s \leq r - 1$, and hence $\Delta(\sigma) \geq 2$, and also $s \geq 2$ since $M \notin Q$. We take an edge E with $\text{type}(E) = \sigma$ such that the Δ -part is taken from V_j to include x and y , and the remaining parts are chosen from the classes V_1 to V_{s-1} . Again E has a colour pattern with at least $s(Q) + 1$ distinct colours, a contradiction. Hence all classes must be monochromatic of distinct colour, and, up to permutation of classes, the CMDC is the only possible colouring of H .

We now show that for any partition $\pi \in Q$, H is not $(Q \setminus \pi)$ -colourable. Clearly if $|Q| = 1$, the result is trivial, so we may assume that $|Q| \geq 2$. Now consider $\pi \in Q$ and $Q^* = Q \setminus \pi$. Clearly the CMDC is not a valid $(2r)$ - Q^* -colouring of H , since if $E \in E(H)$ has $\text{type}(E) = \pi$, then also $\text{pat}(E) = \pi \in \Sigma = Q$, but $\text{pat}(E) = \pi \notin Q^*$.

So let us assume any Q^* -colouring of H which is not the CDMC. Once again let n_j be the number of colours appearing in V_j for $j = 1 \dots 2r$. If there are at least r classes for which $n_j \geq r$, then we can choose any edge E of H with $\text{type}(E) = \sigma \in \Sigma$ to include r distinct colours, so that $\text{pat}(E) = R \notin Q$ and hence not in Q^* . As before, there are at most $r - 1$ such classes, and hence at least $r + 1$ classes in which $n_j \leq r - 1$. Let these classes be V_1 to V_{r+1} . Since $q = (r - 1)^2 + 1$, it follows by the pigeon-hole principle that in each of the classes V_1 to V_{r+1} there is a colour that appears at least r times. Let c_j

be the colour which appears most frequently in class V_j for $j = 1 \dots r + 1$. We show that the colours c_1, \dots, c_{r+1} must all be distinct and cannot appear in any other class. Suppose that colour c_j appears in another class among V_1 to V_{2r} , say in class V_t . Let σ be a partition such that $\Delta(\sigma) = \Delta(Q)$ and observe that $\Delta(Q) \geq \Delta(Q^*)$. Also since $M \notin Q$, $\Delta(Q) = \Delta(\sigma) < r$.

Let E be an edge with $\text{type}(E) = \sigma$. We choose the Δ -part of E from the vertices in V_j of colour c_j , and another part from class V_t to include the vertex of colour c_j , with the rest of the parts chosen from the remaining classes. Then $\text{pat}(E)$ has at least $\Delta(Q) + 1 \geq \Delta(Q^*) + 1$ vertices of the same colour c_j , which is not valid in a Q^* -colouring of H . Hence c_1, \dots, c_{r+1} are all distinct and each colour c_j only appears in class V_j for $j = 1 \dots r + 1$. But now let us choose an edge E with $\text{type}(E) = \pi \in \Sigma$ from among the classes V_1, \dots, V_{r+1} . This edge has $\text{pat}(E) = \pi \notin Q^*$, making the Q^* -colouring of H invalid. Hence H is not $(Q \setminus \pi)$ -colourable. □

3 Bounded clique number and Unbounded Chromatic Number

We now consider the size of the largest clique, $\omega(H)$, for a Σ -hypergraph H . In general finding the size of the largest clique in a graph or hypergraph is NP -complete problem [9]. Here we will show that for Σ -hypergraphs $H(n, r, q | \Sigma)$, $\omega(H)$ depends only on some structural properties of Σ , and hence $\omega(H)$ can be computed in $O(1)$ time for fixed Σ .

We start with a definition. For a family $F \subseteq \Sigma$ of partitions of r and $k \geq r$, we say that F is k -full if the following conditions hold:

1. There exist positive integers b_1, b_2, \dots, b_t such that $b_1 + b_2 + \dots + b_t = k$.
2. For every $A = \{a_1, \dots, a_t\}$ such that $b_i \geq a_i \geq 0$ for $i = 1 \dots t$ and $\sum_{i=1}^t a_i = r$, there exists $\pi \in F$ such that the elements of π are $\text{sup}(A) = \{a_i : a_i > 0\}$, the *support* of A .

Theorem 3.1. *Given $H = H(n, r, q | \Sigma)$ with $n \geq s(\Sigma)$ and $q \geq \Delta(\Sigma)$, then $\omega(H) = \max\{k : \exists F \subseteq \Sigma \text{ where } F \text{ is } k\text{-full}\}$.*

Proof. Let K be a fixed maximal clique. Suppose $\{v_1, \dots, v_r\}$ is a set of r vertices in K . Since K is a clique, these vertices must form an edge E with $\text{type}(E) = \pi \in \Sigma$. Clearly $E = \bigcup \{E \cap V_i : i = 1 \dots n\}$.

Let $a_i = |E \cap V_i|$ and let $b_i = |K \cap V_i|$. We observe that $\sum_1^n b_i = |K|$ and $\sum_1^n a_i = r$, and $a_i \leq b_i$ for $i = 1 \dots n$. Hence we define $F = \{\pi : \exists E \subset K \text{ with } \text{type}(E) = \pi\}$, and we see that $F \subset \Sigma$ is k -full, proving that $\omega(H) \leq \max\{k : \exists F \subset \Sigma \text{ where } F \text{ is } k\text{-full}\}$.

Now suppose $F \subset \Sigma$ is k -full. Then there exist b_1, \dots, b_t such that $b_1 + b_2 + \dots + b_t = k$. Let us assume without loss of generality that $b_1 \geq b_2 \geq \dots \geq b_t$.

Let K be a set of vertices obtained by choosing b_1 vertices from V_i for $i = 1 \dots t$. We show that K is a clique.

Let $E = \{v_1, \dots, v_r\}$ be a set of vertices in K . Let $A = \{|E \cap V_i| = a_i, i = 1 \dots t\}$. We observe that $a_1 + a_2 + \dots + a_t = r$, and $a_i \leq b_i$.

Since F is k -full, there is a partition $\pi \in F$ whose parts are the elements of $\text{sup}(A)$. Hence E is an edge in K , and since E was arbitrarily chosen, K is a clique, proving that $\omega(H) \geq \max\{k : \exists F \subset \Sigma \text{ where } F \text{ is } k\text{-full}\}$. \square

The next corollary shows that unless M or R are in Σ , $H(n, r, q | \Sigma)$ has bounded clique size.

Corollary 3.2. *Consider $H = H(n, r, q | \Sigma)$ with $M, R \notin \Sigma$. Then $\omega(H) \leq (r - 1)^2$.*

Proof. Let K be a maximal clique in H . Suppose K intersects at least r classes V_i . Then if we take a single vertex from r different classes, this must form an edge E since K is a clique, but $\text{type}(E) = R \notin \Sigma$. Hence K intersects at most $r - 1$ classes.

Now consider the intersection of K with a single class V_i . Suppose $|K \cap V_i| \geq r$. Then if we choose r vertices from $K \cap V_i$, this must form an edge E since K is a clique, but $\text{type}(E) = M \notin \Sigma$. Hence the size of the intersection of K with any class must be less than r . Therefore $\omega(H) \leq (r - 1)^2$. \square

We now consider the Q -chromatic number $\chi_Q(H)$ and the chromatic number $\chi(H)$, and show that we can construct Σ -hypergraphs with arbitrarily large chromatic number, but with bounded clique number, a subject that got much attention [8, 10, 15].

Theorem 3.3. *Consider $H = H(n, r, q | \Sigma)$ with $M, R \notin \Sigma$ and $M \notin Q \subseteq P(r)$. Let t be an arbitrarily large integer. Then there exists a Σ -hypergraph H with $\chi_Q(H) \geq \chi(H) \geq t$ and $\omega(H) \leq (r - 1)^2$.*

Proof. Let $H = H(n, r, q | \Sigma)$. Let $n \geq (r - 2)(t - 1) + 1$ and $q = (r - 2)(t - 1) + 1$. By Theorem 3.2, $\omega(H) \leq (r - 1)^2$, and H is n - Q -colourable using the CDMC. This is a valid classical colouring since $M \notin \Sigma$, and hence $\chi_Q(H) \geq \chi(H)$.

We now show that H is not k -colourable, for $k < t$. Consider a colouring using k colours, where $1 < k < t$. Since $M, R \notin \Sigma$, it follows that $\Delta(\Sigma) \leq r - 1$ and $s(\Sigma) \leq r - 1$.

Since $q \geq (r - 2)(t - 1) + 1$ and we use at most $t - 1$ colours, in each class there must be a colour which appears at least $r - 1$ times. Since $n \geq (r - 2)(t - 1) + 1$ and we use at most $t - 1$ colours, it follows that one of the colours appears at least $r - 1$ times in $r - 1$ classes. Thus there must be an edge of H contained in these classes which is monochromatic, which is not a valid colouring, and hence not a valid Q -colouring of H . We conclude that $t \leq \chi(H) \leq \chi_Q(H) \leq n$, and $\omega(H) \leq (r - 1)^2$. \square

4 Q -colourings and gaps

We now turn to the conditions for the existence/non-existence of gaps in the Q -spectrum of a hypergraph H . We start with some definitions. These are

similar to the definitions given in [14], but in our case, they are restricted to non-oriented r -uniform hypergraphs. Our approach in this section incorporates and develops several arguments from [14], together with arguments specifically developed to treat Σ -hypergraphs.

Consider a partition $\sigma = (a_1, a_2, \dots, a_s)$ of r into s parts. We say that a partition π is *derived from σ by reduction* if it was formed by choosing a_i, a_j in σ and replacing them by $a_i + a_j$.

The set of partitions of r that are formed by repeatedly applying reduction starting from σ is denoted by $RD(\sigma)$. Similarly, for a set of partitions $Q \subseteq P(r)$, $RD(Q)$, the set of reduction-derived partitions is

$$\bigcup_{\sigma \in Q} RD(\sigma).$$

A set $Q \subseteq P(r)$ is called *reduction closed* if $RD(Q) = Q$, and $RD(Q)$ is called the *reduction closure of Q* . It is clear that $RD(RD(Q)) = RD(Q)$ and that M is always in $RD(Q)$.

As an example, consider $r = 6$ and $\sigma = (3, 1, 1, 1)$. The partitions derived from σ are $(4, 1, 1)$ and $(3, 2, 1)$ - applying reduction to these two partitions give the partitions $(5, 1)$ $(4, 2)$ and $(3, 3)$, which in turn give the partition $(6) = M$. Hence $RD((3, 1, 1, 1)) = \{(3, 1, 1, 1), (4, 1, 1), (3, 2, 1), (5, 1), (4, 2), (3, 3), M\}$.

Now consider a partition $\sigma = (a_1, a_2, \dots, a_s)$ of r into s parts. We say that a partition π is *derived from σ by expansion* if it was formed by choosing a_j in σ and replacing it by $a_j - 1$ and 1.

The set of partitions of r that are formed by repeatedly applying expansion starting from σ is denoted by $EX(\sigma)$. Similarly, for a set of partitions $Q \subseteq P(r)$, $EX(Q)$, the set of expansion derived partitions is

$$\bigcup_{\sigma \in Q} EX(\sigma).$$

A set $Q \subseteq P(r)$ is called *expansion closed* if $EX(Q) = Q$, and $EX(Q)$ is called the *expansion closure of Q* . It is clear that $EX(EX(Q)) = EX(Q)$ and that R is always in $EX(Q)$.

As an example, consider $r = 6$ and $\sigma = (3, 3)$. The partition derived from σ is $(3, 2, 1)$, and those derived from $(3, 2, 1)$ are $(2, 2, 1, 1)$ and $(3, 1, 1, 1)$. These in turn give $(2, 1, 1, 1, 1)$ and finally $(1, 1, 1, 1, 1, 1) = R$. Hence $EX((3, 3)) = \{(3, 3), (3, 2, 1), (2, 2, 1, 1), (3, 1, 1, 1), (2, 1, 1, 1, 1), R\}$.

A set Q of partitions of r is called *simply closed* if Q contains $EX(M) = \{(r), (r-1, 1), (r-2, 1, 1), \dots, (2, 1, 1, 1, \dots, 1), (1, 1, \dots, 1)\}$, the expansion closed family which is obtained starting with the partition M .

We call a set $Q \subseteq P(r)$ *robust* if it is reduction-closed, expansion-closed or simply-closed.

We can rephrase Theorem 20 in [14] using our terminology as follows:

Theorem 4.1. *There are no gaps in the Q -spectrum of any r -uniform hypergraph H if and only if $Q \subseteq P(r)$ is robust.*

We shall here prove the necessity of the condition.

Proof. Case 1 — Q is reduction-closed

Let H be an r -uniform hypergraph. Clearly, H is 1- Q -colourable since $M \in Q$. Now suppose k is the maximum number of colours for which H is k - Q -colourable. Consider a k - Q -colouring of H , hence each edge has $pat(E) \in Q$. Let us recolour the vertices coloured k with the colour $k - 1$. Let us consider the possible effects of this:

1. Consider the edges in which the colour k did not appear. Then the colouring pattern of these edges has not changed and hence is still a colour pattern in Q .
2. Consider edges in which the colour k appeared, but not the colour $k - 1$. If we change the colour k to $k - 1$, the colour pattern of the edges remain the same.
3. Consider edges in which the colour k appeared, as well as the colour $k - 1$. If there are a_i vertices coloured k and a_j vertices coloured $k - 1$, then the new edge has $a_i + a_j$ vertices coloured $k - 1$, and this colour pattern is in Q since Q is reduction closed.

So we have a valid $(k - 1)$ - Q -colouring of H . This process can be repeated until we end up with one colour, that is a monochromatic colouring. This is possible since Q is reduction closed. Therefore there are no gaps in the Q -spectrum of H .

Case 2 — Q is expansion-closed

Let H be an r -uniform hypergraph on n vertices. Clearly, H is n - Q -colourable since $R \in Q$. Now suppose the minimum colours for which a Q -colouring of H exists is $k < n$, and consider a k - Q -colouring of H . Let us choose a vertex v of colour x which appears at least twice (there must be such a colour since $k < n$). Let us change the colour of v to a new colour y . Hence there are now $k + 1$ colours on the vertices of H . Let us consider the possible effects of this:

1. For edges which do not include the vertex v , the colour pattern has not changed.
2. If the edge includes the vertex v , but no other vertex of colour x , the new edge has the same colour pattern.
3. If the edge includes the vertex v and other vertices of colour x , let the number of vertices of colour x (including v) be a_i . If we change the colour of v to colour y , we now have $(a_i - 1)$ vertices of colour x , and one vertex of colour y . Since Q is expansion-closed, this new colour pattern is in Q .

Hence we have a valid $(k + 1)$ - Q -colouring of H . Again, this process can be repeated until we reach n colours since Q is expansion-closed. Therefore there are no gaps in the Q -spectrum of H .

Case 3 — Q is simply-closed

Let H be an r -uniform hypergraph on n vertices. For each k , $1 \leq k \leq n$ we colour the vertices of H as follows: we take $k - 1$ vertices say v_1 to v_{k-1} and colour them with $k - 1$ distinct colors say $1, \dots, k - 1$. We colour the remaining vertices all with the same colour k . Any edge will have the colour pattern $(r - j, 1, 1, \dots, 1)$ for $0 \leq j \leq r$, so H is k - Q -colourable for $1 \leq k \leq n$. \square

Rather than dealing with any r -uniform hypergraph to prove the full converse (which can be done following the original proof modified to the Q -colouring language), we now prove a partial converse to Theorem 4.1 for Σ -hypergraphs, and then prove that hypergraphs of the form $H = H(n, 4, q | \Sigma)$ have no gap in their Q -spectrum where $Q = \{(3, 1)\}$ despite Q being non-robust.

Theorem 4.2. *If either $M \in Q$ or $R \in Q$, and Q is not a robust family, then there exists a Σ -hypergraph H with a gap in its Q -spectrum.*

We shall prove this theorem in a series of three lemmas.

Lemma 4.3. *Suppose $Q \subset P(r)$ and $M, R \in Q$, but Q is not simply closed. Then the Σ -hypergraph H on r^3 vertices (and a hypergraph K on r^2 vertices) has a gap in its Q -spectrum.*

Proof. Consider the complete r -uniform hypergraph K on r^2 vertices. It is 1- Q -colourable since $M \in Q$, and it is r^2 - Q -colourable since $R \in Q$. Now consider an r -colouring of K . Then at least one colour, say colour 1, appears at least r times, say on vertices v_1, \dots, v_r . Each other colour appears at least once, say colour j , for $2 \leq j \leq r$ appears on the vertex u_j . Then the edge containing k vertices chosen from v_1, \dots, v_r , and $r - k$ vertices chosen from u_2, \dots, u_r has colouring type $(r - k, 1, 1, \dots, 1)$, $\forall k = 0 \dots r$. But this means that $EX(M)$ must be contained in Q , a contradiction.

Now consider the Σ -hypergraphs $H = H(r^2, r, r \mid \Sigma = Q)$ and $H = H(r, r, r^2 \mid \Sigma = Q)$ — the hypergraphs are 1-colourable since $M \in Q$ and r^3 -colourable since $R \in Q$, but is not r -colourable since:

1. in $H = H(r^2, r, r \mid \Sigma = Q)$, the set of vertices obtained by choosing one vertex from each class has r^2 vertices and since $R \in \Sigma = Q$ it is a complete r -uniform hypergraph on r^2 vertices, so is not r - Q -colourable by the above argument.
2. in $H = H(r, r, r^2 \mid \Sigma = Q)$, each class V_i of H is a complete r -uniform hypergraph on r^2 vertices since $M \in Q$, and again, by the above argument is not r - Q -colourable.

\square

Lemma 4.4. *Let $Q \subset P(r)$ such that $R \in Q$, $M \notin Q$ and Q is not expansion-closed. Then the Σ -hypergraph $H = H(r, r, r^2 \mid \Sigma = Q)$ on r^3 vertices has a gap in its Q -spectrum.*

Proof. Consider $H = H(r, r, r^2 \mid \Sigma = Q)$. Then H is r - Q -colourable using the CDMC, and H is r^3 - Q -colourable since $R \in Q$. We claim that H is not $(r+1)$ - Q -colourable. Suppose that f is a proper $(r+1)$ - Q -colouring of H . We will show that this assumption contradicts the fact that Q is not expansion-closed.

Let f_i be the colour which appears most in class V_i — it is clear that since $q = r^2$ and we use at most $r+1$ colours in each class, f_i appears at least r times in class V_i , for $1 \leq i \leq r$. So we may assume that, without loss of generality, the first r vertices in V_i , $v_{i,j}$ for $j = 1 \dots r$, are such that $f(v_{i,j}) = f(i)$.

We first show that $f(i) \neq f(j)$ for $1 \leq i, j \leq r$. Suppose for contradiction that for some i, j , $f(i) = f(j)$. Recall that $\Delta(Q) = \max\{\Delta(\pi) : \pi \in Q\}$, and since $M \notin Q$, $\Delta(Q) < r$. Consider an edge $E \in E(H)$ with $\text{type}(E) = \sigma$ and $\Delta(\sigma) = \Delta(Q)$, such that the $\Delta(\sigma)$ vertices are chosen among the first r vertices in V_i , so that they all have colour $f(i)$, and such that another part of σ is chosen from the first r vertices of the class V_j . Since $f(i) = f(j)$, this would imply that edge E has at least $\Delta(Q) + 1$ vertices of the same colour, contradicting the maximality of $\Delta(Q)$.

We will now show that the assumption that f is a proper $(r+1)$ - Q -colouring of H and the fact that Q is not expansion-closed are contradictory. Since Q is not expansion-closed, there is a partition $\sigma = (a_1, a_2, \dots, a_s)$ such that for some $j \geq 2$, the derived partition $\sigma^* = (a_1, \dots, a_j - 1, \dots, a_s, 1)$ is not in Q .

Since there are r classes in H , we may assume that $f(i) = i$ for $i = 1 \dots r$ and that the colour $r+1$ appears in class V_1 , without loss of generality. Let us also assume that $v_{1,r+1}$ has colour $(r+1)$.

Let $E \in E(H)$ be an edge with $\text{type}(E) = \sigma$ such that the part a_j is taken from the first $a_j < r$ vertices of V_1 and the remaining parts from other classes, with all parts taken from the first r vertices of the respective class. Thus E is properly coloured. Now in part a_j let us replace the vertex $v_{1,1}$ with the vertex $v_{1,r+1}$ to create E^* which is still a valid edge. Clearly $\text{pat}(E^*) = \sigma^*$ which is not in Q , contradicting the assumption that f is a proper $(r+1)$ - Q -colouring of H . \square

Lemma 4.5. *Suppose $Q \subset P(r)$ such that $M \in Q$ but $R \notin Q$, and Q is not reduction-closed. Then the Σ -hypergraph $H = H(r^2, r, r \mid \Sigma = Q)$ on r^3 vertices has a gap in its Q -spectrum.*

Proof. Consider $H = H(r^2, r, r \mid \Sigma = Q)$. Clearly H is 1- Q -colourable since $M \in Q$, and H is r^2 - Q -colourable using the CDMC. We claim that H is not $(r^2 - 1)$ - Q -colourable. Suppose on the contrary that f is a proper $(r^2 - 1)$ - Q -colouring of H .

We first show that each class must be monochromatic. Assume without loss of generality that in class V_1 , $1 = f(v_{1,1}) \neq f(v_{1,2}) = 2$. Since we are using $r^2 - 1$ colours and each class has r vertices, there must be a vertex of another colour, say 3, not in V_1 . Let this vertex be $v_{2,1}$ in V_2 . Still V_1 and V_2 can have at most $2r$ colours, hence there is another vertex of colour say 4 not in $V_1 \cup V_2$. Let this vertex be $v_{1,3} \in V_3$. We can continue this process at least until class V_r , since $r - 1$ classes can have at most $r(r - 1) < r^2 - 1$ colours. Hence we conclude that we can assume the vertices $v_{1,1}, v_{1,2}, v_{2,1}, v_{3,1}, \dots, v_{r,1}$ are $r+1$

vertices of $r + 1$ distinct colours. Now let $\sigma \in \Sigma = Q$ be a partition such that $s(\sigma) = s(Q)$. Since $R \notin Q$, $s(Q) < r$ and hence $\Delta(\sigma) \geq 2$. We choose an edge E with $\text{type}(E) = \sigma$ such that the part $a_j \in \sigma$ for $j = 1 \dots s$ consists of the first a_j vertices of class V_j . Clearly the $\text{pat}(E) = \pi$ where $s(\pi) \geq s(\sigma) + 1$, hence $\pi \notin Q$. Thus all classes must be monochromatic.

Since there are r^2 classes and we are using $r^2 - 1$ colours, there are two classes, say V_1 and V_2 which are monochromatic of the same colour, while all other classes are monochromatic with pairwise distinct colours. Since Q is not reduction-closed, there is a partition $\pi = (a_1, \dots, a_s) \in Q$ such that for some $1 \leq i, j \leq s$, the partition π^* formed by replacing a_i and a_j by $a_i + a_j$ is not in Q .

Now consider the edge E^* with $\text{type}(E) = \pi \in \Sigma = Q$ such that the parts a_i and a_j are chosen from classes V_1 and V_2 , while the other parts are chosen from any of the remaining classes. Clearly $\text{pat}(E) = \pi^* \notin Q$, contradicting the assumption that f was a proper $(r^2 - 1)$ - Q -colouring. □

Combining these three lemmas, we have a proof of Theorem 4.2.

We now consider the case $Q = \{(3, 1)\}$. Clearly M and R are not in Q , and Q is not robust. By Theorem 4.1 there exists a hypergraph H that has a gap in its Q -spectrum.

We will show that this is not a Σ -hypergraph, hence only a partial converse to Theorem 4.1 can hold for Σ -hypergraphs. Yet we give a concrete example of a small 4-uniform hypergraph that has a gap in its Q -spectrum.

We have checked that

1. Theorem 4.1 holds for Σ -hypergraphs for $r = 3$ namely if Q is not robust then there is $H(n, 3, q|\Sigma)$ with a gap in its Q -spectrum.
2. Theorem 4.1 holds for Σ -hypergraphs for $r = 4$ unless $Q = \{(3, 1)\}$, namely if Q is not robust and not $\{(3, 1)\}$ then there is $H(n, 4, q|\Sigma)$ with a gap in its Q -spectrum.

The next Proposition covers the case $Q = \{(3, 1)\}$.

Proposition 4.6. *Consider $H(n, 4, q|\Sigma)$ where $\Sigma \subseteq P(4)$, and $Q = \{(3, 1)\}$. Then H does not have a gap in its Q -spectrum for $n \geq s(\Sigma)$, $q \geq \Delta(\Sigma)$.*

Proof. We first consider the case where Σ consists of a single partition of four (there are 5 such partitions). The conditions $n \geq s(\Sigma)$, $q \geq \Delta(\Sigma)$ are included so that all edge types in Σ can be realised.

1. Consider $H(n, 4, q|\sigma = (3, 1))$ where $n \geq 2$, $q \geq 3$. Then H is n - Q -colourable using the CDMC. In general, no class can contain 3 colours since in that case we can choose three vertices of distinct colours to give an edge containing three colours. So we may assume that each class contains at most two colours.

Consider a k -colouring, where $k < n$. Then at least one colour, say the colour 1, appears in two distinct classes, say V_1 and V_2 , without loss of

generality. If in one of these classes, say V_1 , there are three vertices of colour 1, then we can choose a monochromatic edge from V_1 and V_2 . If in say V_1 there are two vertices of the same colour but not colour 1, then we can choose these two vertices and the vertex of colour 1 from V_1 , plus the vertex of colour 1 in V_2 , to give an edge with $pat(E) = (2, 2)$. So there can only be one vertex of another colour, say 2, in V_1 . But then all vertices in V_2 must be of colour 1, because otherwise, if in V_2 there is a vertex of colour 2, we can choose an edge which includes two vertices of colour 1 and two vertices of colour 2, and if there is a vertex of another colour other than 1 and 2 in V_2 , then we can choose an edge to include three colours. But if all the vertices in V_2 are of colour 1, and there is a vertex in V_1 of colour 1, we can choose a monochromatic edge. Hence it is not possible to colour H with less than n colours.

Now consider a k -colouring, where $k > n \geq 2$. Then at least one class contains two vertices of distinct colour, say in V_1 there is a vertex of colour 1 and a vertex of colour 2. But since there are at least three colours, there is a vertex of another colour, say colour 3 in some other class in H . But then we can choose an edge to include the two vertices of colours 1 and 2 in V_1 , and this vertex of colour 3, giving an edge which includes 3 colours. Hence a Q -colouring with more than n colours is not possible. Therefore $Spec_Q(H) = \{n\}$.

2. Consider $H(n, 4, q | \sigma = (2, 2))$ where $n \geq 2, q \geq 2$. If $n = q = 2$ we can colour one class monochromatically, say with colour 1, and in the other class, we colour a vertex with colour 1 and one with colour 2, and this is the only valid Q -colouring of this graph, and hence its Q -spectrum is not broken. If $q > 2$, then if V_1 is monochromatic of say colour 1, then the vertices in V_2 must be coloured in such a way that we always choose a vertex of colour 1 and one of another colour. But this is impossible if $q > 3$.

Hence consider $q = 2$ and $n > 2$. If we colour the first 2 classes as described above, then if in the third class, both vertices are of colour 1, we can choose a monochromatic edge from classes V_1 and V_3 , while if both vertices are coloured 2, we can choose an edge E from classes V_1 and V_3 such that $pat(E) = (2, 2)$. If both colours 1 and 2 appear in V_3 , then we can choose an edge with colour pattern $(2, 2)$ from classes V_2 and V_3 . Finally, if a third colour appears in V_3 , then we can choose an edge which includes three colours from V_2 and V_3 . So again H is not colourable, and hence its Q -spectrum is empty.

3. Consider $H(n, 4, q | \sigma = (2, 1, 1))$ where $n \geq 3, q \geq 2$. Consider two vertices of the same colour, say colour 1, in a class, say V_1 . Hence without loss of generality, we may assume in V_2 there is a vertex of colour 1 and in V_3 a vertex of colour 2. This forces V_2 to be monochromatic of colour 1, otherwise the two vertices from V_1 and the vertex from V_3 together with a vertex from V_2 that is not of colour 1 will give either an edge E with $pat(E) = (2, 2)$ or $pat(E) = (2, 1, 1)$.

If V_3 is not monochromatic of colour 2 then either there is a vertex of colour 1 in V_3 and we have a monochromatic edge choosing the two vertices from V_1 , any vertex from V_2 and this vertex from V_3 , or there is an edge that includes three colours choosing the two vertices from V_3 and any vertex of V_2 and a vertex of colour 1 from V_1 . Hence V_3 must be monochromatic of colour 2 and choosing any two vertices from V_3 , one from V_2 and a vertex of colour 1 from V_1 , we get an edge with color pattern $(2, 2)$.

So assume that no two vertices in V_1 have the same colour. Then if we choose two vertices from this class, say one of colour 1 and one of colour 2 without loss of generality, then the remaining classes must be monochromatic, either all of colour 1 or all of colour 2. Let us say they are all of colour 1. But then we can choose a monochromatic edge by choosing a vertex of colour 1 from V_1 , a vertex from V_2 and two vertices from V_3 , all of colour 1. Hence H is not Q -colourable.

4. Consider $H(n, 4, q | \sigma = (1, 1, 1, 1))$ where $n \geq 4$, $q \geq 1$. If $n = 4$ and $q \geq 1$, then without loss of generality we can colour V_1, V_2 and V_3 monochromatically using colour 1, while the vertices in V_4 can be coloured using colours different from 1, that is using one colour, say colour 2, up to q colours. Hence $H(4, 4, 1 | \sigma = (1, 1, 1, 1))$ has Q -spectrum $\{2, 3, \dots, q + 1\}$. Now consider a k -colouring of H for $k > q + 1$. Let us choose an edge $E = \{v_1, v_2, v_3, v_4\}$ such that v_i is chosen from V_i for $i = 1 \dots 4$, so that $pat(E) = (3, 1)$, and without loss of generality, we assume that v_1, v_2 and v_3 are the same colour, say colour 1, and v_4 is of a different colour. Clearly, all vertices in V_1, V_2 and V_3 must be of colour 1, otherwise we can choose an edge with colour pattern $(2, 2)$ or $(2, 1, 1)$ which would make the colouring invalid. Hence only class V_4 can contain vertices of different colours, and since there are q vertices in a class, the maximum number of colours which can be used in a valid Q -colouring is $q + 1$.

If $n > 4$, then we must choose a vertex from each class such that three have the same colour, and one has different colour and if $q > 1$, this is clearly not possible.

5. Consider $H(n, 4, q | \sigma = (4))$ where $n \geq 1$, $q \geq 4$. Then we must choose four vertices from one class such that three are of the same colour, and one is of a different colour. If $q > 4$, this is not possible. For $q = 4$, we can colour each class to include 3 vertices of one colour, and one vertex of a different colour. This means that the Q -spectrum of such a graph is $\{2, 3, \dots, 2n\}$, and it contains no gaps.

Hence if $n \geq s(\sigma)$, $q \geq \Delta(\sigma)$, and $H(n, 4, q | \sigma)$ where $\sigma \in P(4)$, there is no gap in the Q -spectrum of H , for $Q = \{(3, 1)\}$.

Having done the above five cases we use the following simple observation. Let $\Sigma = \Sigma_1 \cup \Sigma_2$ where $\Sigma_1 \cap \Sigma_2 = \emptyset$ then

$$Spec_Q(H(n, r, q | \Sigma)) \subseteq Spec_Q(H(n, r, q | \Sigma_1)) \cap Spec_Q(H(n, r, q | \Sigma_2)),$$

and hence one can easily verify that for any Σ consisting of at least two partitions of 4, $\text{Spec}_Q(H(n, r, q|\Sigma))$ is either empty or contains exactly one value, hence it is not broken. \square

Although no Σ -hypergraph has a gap in its $(3, 1)$ -spectrum, we now give an example of a 4-uniform hypergraph with a broken $(3, 1)$ -spectrum.

Consider the 4-uniform hypergraph H having sixteen vertices. Suppose they are grouped into cells each containing two vertices, and the cells are arranged in a 4×2 grid. The cell in row i column j will be denoted by C_{ij} , $1 \leq i \leq 4$ and $1 \leq j \leq 2$. The vertices in cells C_{i1} and C_{i2} are said to be in the i^{th} row and the vertices in cells C_{1j} to C_{4j} are said to be in column j . All 4-subsets K of four vertices form an edge if the sizes of the non-empty intersections of K with the rows, and also the sizes of the non-empty intersections of K with the columns form the partition $(3, 1)$. Figure 1 shows an example with two edges in this hypergraph.




Figure 1: Two edges are shown with the vertices in one edge marked with \diamond while in the other edge the vertices are marked as \times

We show that H is 2- Q -colourable, 4- Q -colourable for $Q = \{(3, 1)\}$, but not 3-colourable. Consider a 2-colouring using colours 1 and 2 such that the vertices in column 1 receive colour 1, and the vertices in column 2 receive colour 2. It is easy to see that an edge $E \in E(H)$ is such that $\text{pat}(E) = (3, 1)$. Consider a 4-colouring of H such that the vertices in row i receive colour i for $i = 1 \dots 4$. Again it is easy to see that $\text{pat}(E) = (3, 1), \forall E \in E(H)$.

We now show that H is not 3- Q -colourable. Suppose H is coloured using three colours 1, 2 and 3. Suppose first that the vertices in a cell receive different

colours. We may assume without loss of generality that the vertices in C_{11} receive colours 1 and 2. There must be a vertex v which receives colour 3. If v is in column 1, then we can choose an edge which contains the three vertices of different colours, and hence the coloring is not valid. Therefore all vertices in column 1 must have colour 1 or 2.

Suppose v appears in column 2. If v is in row 1, we can choose an edge which includes the vertices in row 1 coloured 1,2 and 3, which is again invalid. Therefore we can assume v is not in row 1, and we assume without loss of generality that v is in C_{22} . SO we can choose an edge K as follows: K contains v and the two vertices in C_{21} . If these two vertices have colours 1 and 2 (recall that they cannot have colour 3), then already the colouring is invalid. If they are both of the same colour, say colour 1, we can choose the final vertex from C_{11} with column 2, (and similarly if both have colour 2, we can choose a vertex of colour 1 from C_{11}), again showing that the colouring is not valid.

So we may conclude that the vertices in one cell must have the same colour. Without loss of generality, suppose the vertices in C_{11} receive colour 1. If there is another vertex of colour 1 in the first column, say in row 2, then the vertices in C_{12} must receive another colour, say colour 2, otherwise we can choose a monochromatic edge. But there must be a vertex v coloured 3. If v is in the first column, then letting edge K contain v , the two vertices in C_{11} and a vertex coloured 2 from C_{12} gives an edge with three colours. Similarly, if v is in column 2 we have the same situation. Hence colour 1 cannot appear on the other cells in column 1. By the same argument, no colour can appear in two cells in the same column. But then for cells C_{21} , C_{31} and C_{41} only colours 2 and 3 are available (recall that the vertices in a cell must receive the same colour), and therefore the vertices in at least two cells must all receive the same colour, giving us the same contradiction.

Hence there is no valid 3- Q -colouring of H , and $\text{Spec}_Q(H)$ contains a gap.

5 Conclusion

In this paper we have tried to unify and clarify a number of hypergraph colouring problems using the language of Q -colourings and Σ -hypergraphs. We have defined tight colourings, which is a generalisation of unique colourability, and we have shown that for all Q not containing the partitions M and R there exist Σ -hypergraphs which are tightly Q -colourable. We have also shown that finding the clique number of Σ -hypergraphs, which is, in general, an NP -complete problem, can be done in $O(1)$ time and that there exist Σ -hypergraphs with arbitrarily high chromatic, and hence Q -chromatic, number but bounded clique number.

Finally we have considered the result in [14], which characterises those Q for which there exist hypergraphs with a gap in their Q -spectrum. In [14], this characterisation is given for oriented hypergraphs and the result for hypergraphs is given as a corollary. Here we give a short direct proof of the necessity of their condition for Σ -hypergraphs using the language of Q -colourings. We also prove a partial converse of their result for Σ -hypergraphs and show that, for

$Q = \{(3, 1)\}$, which is known to give hypergraphs with gaps, there is no Σ -hypergraph with a gap in its Q -spectrum. However, we present a hypergraph with a gap in its Q -spectrum for $Q = \{(3, 1)\}$. This hypergraph can be viewed as a further generalisation of σ -hypergraphs, which we intend to study further.

References

- [1] M. Axenovich and J. Rollin. Brooks Type Results for Conflict-Free Colorings and $\{a, b\}$ -factors in graphs. *ArXiv e-prints*, October 2014.
- [2] C. Berge and E. Minieka. *Graphs and hypergraphs*, volume 7. North-Holland publishing company Amsterdam, 1973.
- [3] C. Bujtás and Z. Tuza. Color-bounded hypergraphs, I: General results. *Discrete Mathematics*, 309(15):4890–4902, 2009.
- [4] C. Bujtás and Z. Tuza. Color-bounded hypergraphs, IV: Stable colorings of hypertrees. *Discrete Mathematics*, 310(9):1463–1474, 2010.
- [5] Y. Caro and J. Lauri. Non-monochromatic non-rainbow colourings of σ -hypergraphs. *Discrete Mathematics*, 318(0):96 – 104, 2014.
- [6] Y. Caro, J. Lauri, and C. Zarb. $(2, 2)$ -colourings and clique-free σ -hypergraphs. *Discrete Applied Mathematics*, 185(0):38 – 43, 2015.
- [7] Y. Caro, J. Lauri, and C. Zarb. Constrained colouring and σ -hypergraphs. *Discussiones Mathematicae Graph Theory*, 35(1):171–189, 2015.
- [8] P. Erdos and L. Lovász. Problems and results on 3-chromatic hypergraphs and some related questions. *Infinite and finite sets*, 10:609–627, 1975.
- [9] M.R. Garey and D.S. Johnson. *Computers and intractability*, volume 29. wh freeman, 2002.
- [10] H. Gebauer. On the construction of 3-chromatic hypergraphs with few edges. *Journal of Combinatorial Theory, Series A*, 120(7):1483 – 1490, 2013.
- [11] T. Griggs, G. Lo Faro, and G. Quattrocchi. On some colouring of 4-cycle systems with specified block colour patterns. *Discrete Mathematics*, 308(2):465–478, 2008.
- [12] J.L. Gross and J. Yellen. *Handbook of graph theory*. CRC press, 2004.
- [13] T. Jiang, D. Mubayi, Z. Tuza, V. Voloshin, and D.B. West. The chromatic spectrum of mixed hypergraphs. *Graphs and Combinatorics*, 18(2):309–318, 2002.
- [14] Z. Kára, J. Dvůrák, D. Pangrác, and O. Král. Pattern hypergraphs. *The Electronic Journal of Combinatorics*, 17(R15):1, 2010.

- [15] A. V. Kostochka and V Rödl. Constructions of sparse uniform hypergraphs with high chromatic number. *Random Structures & Algorithms*, 36(1):46–56, 2010.
- [16] S. Milici, A. Rosa, and V. Voloshin. Colouring steiner systems with specified block colour patterns. *Discrete Mathematics*, 240(13):145 – 160, 2001.
- [17] G. Quattrocchi. Colouring 4-cycle systems with specified block colour patterns: the case of embedding \tilde{p} 3-designs. *Journal of Combinatorics*, 8(1):R24–R24, 2001.
- [18] Z. Tuza, V. Voloshin, and H. Zhou. Uniquely colorable mixed hypergraphs. *Discrete Mathematics*, 248(13):221 – 236, 2002.
- [19] V. I. Voloshin. *Coloring mixed hypergraphs: theory, algorithms and applications*, volume 17 of *Fields Institute Monograph*. American Mathematical Society, 2002.