Low c-differential and c-boomerang uniformity of the swapped inverse function

Pantelimon Stănică Applied Mathematics Department, Naval Postgraduate School, Monterey, USA. E-mail: pstanica@nps.edu

September 29, 2020

Abstract

Modifying the binary inverse function in a variety of ways, like swapping two output points has been known to produce a 4-differential uniform permutation function. Recently, in [18] it was shown that this swapped version of the inverse function has boomerang uniformity exactly 10, if $n \equiv 0 \pmod{6}$, 8, if $n \equiv 3 \pmod{6}$, and 6, if $n \not\equiv 0 \pmod{3}$. Based upon the c-differential notion we defined in [14] and c-boomerang uniformity from [27], in this paper we characterize the c-differential and c-boomerang uniformity for the (0,1)-swapped inverse function in characteristic 2: we show that for all $c \neq 1$, the c-differential uniformity is upper bounded by 4 and the c-boomerang uniformity by 5 with both bounds being attained for $n \geq 4$.

Keywords: Boolean, *p*-ary functions, *c*-differentials, differential uniformity, boomerang uniformity, perfect and almost perfect *c*-nonlinearity **MSC 2000**: 06E30, 11T06, 94A60, 94D10.

1 Introduction and basic definitions

The boomerang attack on block ciphers developed by Wagner [32] was quickly picked by several works [3, 16, 2, 17] in the applied realm, but it took almost ten years until Cid et al. [12] came up with a theoretical tool called the Boomerang Connectivity Table (BCT), and further Boura and Canteaut [5] defined the boomerang uniformity to theoretical investigate

the functions' resistance to the boomerang attack. Using a multiplier differential we introduced in [14], we extended the notion of the c-Boomerang Connectivity Table (c-BCT) and c-boomerang uniformity in [27].

Here, we continue the work by investigating the well-known swapped inverse, proposed in [31], which was the subject of many papers since then (see [9, 24, 25, 28, 33], to cite just a few papers; a generalization allowing the modification of any two output values is done in [15]).

As customary, we let:

- 1. m, n are positive integers and p a prime number, #S denotes the cardinality of a set S and \bar{z} is the complex conjugate;
- 2. \mathbb{F}_{p^n} is the finite field with p^n elements, and $\mathbb{F}_{p^n}^* = \mathbb{F}_{p^n} \setminus \{0\}$ is the multiplicative group (for $a \neq 0$, we often write $\frac{1}{a}$ to mean the inverse of a in the multiplicative group);
- 3. We let \mathbb{F}_p^n be the *n*-dimensional vector space over \mathbb{F}_p . Next, $f: \mathbb{F}_{p^n}$ (or \mathbb{F}_p^n) to \mathbb{F}_p is a *p-ary function* on *n* variables. Any map $F: \mathbb{F}_{p^n} \to \mathbb{F}_{p^m}$ (or, $\mathbb{F}_p^n \to \mathbb{F}_p^m$) is called a *vectorial p-ary function*, or (n, m)-function.
- 4. When m=n, F can be uniquely represented as a univariate polynomial over \mathbb{F}_{p^n} (using some identification, via a basis, of the finite field with the vector space) of the form $F(x) = \sum_{i=0}^{p^n-1} a_i x^i$, $a_i \in \mathbb{F}_{p^n}$, whose algebraic degree is then the largest Hamming weight of the exponents i with $a_i \neq 0$.
- 5. Given a p-ary function f, the derivative of f with respect to $a \in \mathbb{F}_{p^n}$ is the p-ary function $D_a f(x) = f(x+a) f(x)$, for all $x \in \mathbb{F}_{p^n}$, which can be naturally extended to vectorial p-ary functions.
- 6. While the next concept can be defined for general (n,m)-functions, in this paper we consider m=n. For an (n,n)-function F, and $a,b\in\mathbb{F}_{p^n}$, we let $\Delta_F(a,b)=\#\{x\in\mathbb{F}_{p^n}:F(x+a)-F(x)=b\}$. We call the quantity $\delta_F=\max\{\Delta_F(a,b):a,b\in\mathbb{F}_{p^n},a\neq 0\}$ the differential uniformity of F. If $\delta_F=\delta$, then we say that F is differentially δ -uniform.
- 7. If $\delta = 1$, then F is called a *perfect nonlinear* (PN) function, or *planar* function. If $\delta = 2$, then F is called an *almost perfect nonlinear* (APN) function. It is well known that PN functions do not exist if p = 2.

The reader can consult [8, 10, 11, 13, 21, 29] for more on Boolean and p-ary functions.

The initial concept of boomerang uniformity was defined for permutations (of course, proper S-boxes) in the following way. Let F be a permutation on \mathbb{F}_{2^n} and $(a,b) \in \mathbb{F}_{2^n} \times \mathbb{F}_{2^n}$. One defines the entries of the Boomerang Connectivity Table (BCT) by

$$\mathcal{B}_F(a,b) = \#\{x \in \mathbb{F}_{2^n} | F^{-1}(F(x)+b) + F^{-1}(F(x+a)+b) = a\},\$$

where F^{-1} is the compositional inverse of F. The boomerang uniformity of F is defined as

$$\beta_F = \max_{a,b \in \mathbb{F}_{n^n} *} \mathcal{B}_F(a,b).$$

We also say that F is a β_F -uniform BCT function. Surely, $\Delta_F(a,b) = 0, 2^n$ and $\mathcal{B}_F(a,b) = 2^n$ whenever ab = 0. We know that $\delta_F = \delta_{F^{-1}}, \, \beta_F = \beta_{F^{-1}},$ and for permutations, $\beta_F \geq \delta_F$ and they are equal for APN permutations. Further, from [18] we know that for quadratic permutations, $\delta_F \leq \beta_F \leq \delta_F(\delta_F - 1)$.

We mention here that this concept was quickly picked up and studied in [5, 6, 9, 18, 19, 22, 30], to mention just a few recent papers.

Li et al. [18] (see also [22]) observed that

$$\mathcal{B}_{F}(a,b) = \# \left\{ (x,y) \in \mathbb{F}_{p^{n}} \times \mathbb{F}_{p^{n}} \mid {F(x)+F(y)=b \atop F(x+a)+F(y+a)=b} \right\},$$

$$\stackrel{y=x+\gamma}{=} \# \left\{ (x,\gamma) \in \mathbb{F}_{p^{n}} \times \mathbb{F}_{p^{n}} \mid {F(x+\gamma)+F(x)=b \atop F(x+\gamma+a)+F(x+a)=b} \right\}$$

$$= \sum_{\gamma \in \mathbb{F}_{p^{n}}} \# \left\{ x \in \mathbb{F}_{p^{n}} \mid D_{\gamma}F(x) = b \text{ and } D_{\gamma}F(x+a) = b \right\},$$

which allowed the concept to be extended to non-permutations, since it avoids the inverse of F.

We extended this notion recently in [27] to c-boomerang connectivity table, based upon our prior [14] c-differential concept (see [26, 34] for more work related to it).

For a p-ary (n,m)-function $F: \mathbb{F}_{p^n} \to \mathbb{F}_{p^m}$, and $c \in \mathbb{F}_{p^m}$, the (multiplicative) c-derivative of F with respect to $a \in \mathbb{F}_{p^n}$ is the function

$$_{c}D_{a}F(x) = F(x+a) - cF(x)$$
, for all $x \in \mathbb{F}_{p^{n}}$.

For an (n,n)-function F, and $a,b \in \mathbb{F}_{p^n}$, we let the entries of the c-Difference Distribution Table (c-DDT) be defined by ${}_c\Delta_F(a,b) = \#\{x \in \mathbb{F}_{p^n} : F(x+a) - cF(x) = b\}$. We call the quantity

$$\delta_{F,c} = \max \{ c \Delta_F(a,b) \mid a,b \in \mathbb{F}_{p^n}, \text{ and } a \neq 0 \text{ if } c = 1 \}$$

the c-differential uniformity of F. If $\delta_{F,c}=\delta$, then we say that F is differentially (c,δ) -uniform (or that F has c-uniformity δ , or for short, F is δ -uniform c-DDT). If $\delta=1$, then F is called a perfect c-nonlinear (PcN) function (certainly, for c=1, they only exist for odd characteristic p; however, as proven in [14], there exist PcN functions for p=2, for all $c\neq 1$). If $\delta=2$, then F is called an almost perfect c-nonlinear (APcN) function. When we need to specify the constant c for which the function is PcN or APcN, then we may use the notation c-PN, or c-APN. It is easy to see that if F is an (n,n)-function, that is, $F: \mathbb{F}_{p^n} \to \mathbb{F}_{p^n}$, then F is PcN if and only if cD_aF is a permutation polynomial.

Further, for an (n, n)-function $F, c \neq 0$, and $(a, b) \in \mathbb{F}_{p^n} \times \mathbb{F}_{p^n}$, we define the c-Boomerang Connectivity Table (c-BCT) entry at (a, b) to be

$$_{c}\mathcal{B}_{F}(a,b) = \# \left\{ x \in \mathbb{F}_{p^{n}} \mid F^{-1}(c^{-1}F(x+a)+b) - F^{-1}(cF(x)+b) = a \right\}.$$

Further, the *c-boomerang uniformity* of F is defined by

$$\beta_{F,c} = \max_{a,b \in \mathbb{F}_{n^n} *} {}_{c} \mathcal{B}_F(a,b).$$

If $\beta_{F,c} = \beta$, we also say that F is a β -uniform c-BCT function. We showed in [27] that we can avoid inverses, thus allowing the definition to be extended to all (n, m)-function, not only permutations. Precisely, the entries of the c-Boomerang Connectivity Table at $(a, b) \in \mathbb{F}_{p^n} \times \mathbb{F}_{p^n}$ can be given by

$${}_{c}\mathcal{B}_{F}(a,b) = \# \left\{ (x,\gamma) \in \mathbb{F}_{p^{n}} \times \mathbb{F}_{p^{n}} \middle| \begin{array}{c} F(x+\gamma) - cF(x) = b \\ F(x+\gamma+a) - c^{-1}F(x+a) = b \end{array} \right\}$$
$$= \sum_{\gamma \in \mathbb{F}_{p^{n}}} \# \left\{ x \in \mathbb{F}_{p^{n}} \middle| \begin{array}{c} cD_{\gamma}F(x) = b \text{ and } c^{-1}D_{\gamma}F(x+a) = b \\ \text{the } c\text{-boomerang system} \end{array} \right\}.$$

Besides various connections between c-DDT and c-BCT and characterizations via Walsh transforms, we investigated some of the known perfect nonlinear and the inverse function in all characteristics, in [27]. For example, we showed that in general, if $F(x) = x^{2^n-2}$ on \mathbb{F}_{2^n} , $_c\mathcal{B}_F(a,b) \leq 3$, and if $F(x) = x^{p^n-2}$ on \mathbb{F}_{p^n} (p odd), $_c\mathcal{B}_F(a,b) \leq 4$ and gave complete conditions when the upper bound happens. It is the purpose of this paper to investigate both the c-DDT and c-BCT for the (0,1)-output swapped inverse function in the binary case (which is a 4-uniform DDT permutation function occurring in many papers [20, 18, 23, 24, 25], to cite just a few works).

The rest of the paper is organized as follows. Section 2 and 3 deal with c-differential uniformity, respectively, c-boomerang uniformity of the swapped function. Section 4 concludes the paper.

2 The c-differential uniformity of the swapped inverse function

We will be using throughout Hilbert's Theorem 90 (see [7]), which states that if $\mathbb{F} \hookrightarrow \mathbb{K}$ is a cyclic Galois extension and σ is a generator of the Galois group $\operatorname{Gal}(\mathbb{K}/\mathbb{F})$, then for $x \in \mathbb{K}$, the relative trace $\operatorname{Tr}_{\mathbb{K}/\mathbb{F}}(x) = 0$ if and only if $x = \sigma(y) - y$, for some $y \in \mathbb{K}$. We also need the following two lemmas.

Lemma 1. Let n be a positive integer. We have:

- (i) The equation $x^2 + ax + b = 0$, with $a, b \in \mathbb{F}_{2^n}$, $a \neq 0$, has two solutions in \mathbb{F}_{2^n} if $\operatorname{Tr}\left(\frac{b}{a^2}\right) = 0$, and zero solutions otherwise (see [1]).
- (ii) The equation $x^2 + ax + b = 0$, with $a, b \in \mathbb{F}_{p^n}$, p odd, has (two, respectively, one) solutions in \mathbb{F}_{p^n} if and only if the discriminant a^2-4b is a (nonzero, respectively, zero) square in \mathbb{F}_{p^n} .

Lemma 2 ([14]). Let p, k, n be integers greater than or equal to 1 (we take $k \le n$, though the result can be shown in general). Then

$$\gcd(2^k + 1, 2^n - 1) = \frac{2^{\gcd(2k,n)} - 1}{2^{\gcd(k,n)} - 1}, \text{ and if } p > 2, \text{ then,}$$

$$\gcd(p^k + 1, p^n - 1) = 2, \text{ if } \frac{n}{\gcd(n,k)} \text{ is odd,}$$

$$\gcd(p^k + 1, p^n - 1) = p^{\gcd(k,n)} + 1, \text{ if } \frac{n}{\gcd(n,k)} \text{ is even.}$$

Consequently, if either n is odd, or $n \equiv 2 \pmod{4}$ and k is even, then $\gcd(2^k+1,2^n-1)=1$ and $\gcd(p^k+1,p^n-1)=2$, if p>2.

Given $F: \mathbb{F}_{p^n} \to \mathbb{F}_{p^n}$, and two points $x_0 \neq x_1$ in \mathbb{F}_{p^n} , we let $G_{x_0x_1}$ be the $\{x_0, x_1\}$ -swapping of F defined by

$$G_{x_0x_1}(x) = F(x) + ((x+x_0)^{p^n-1} + (x+x_1)^{p^n-1})(y_0 + y_1),$$

where $y_0 = F(x_0), y_1 = F(x_1)$. We will sometimes denote $G_{x_0x_1}$ simply by G if there is no danger of confusion. In this paper we consider the (0,1)-swapping of the inverse only, so, $G(x) = x^{p^n-2} + x^{p^n-1} + (x+1)^{p^n-1}$.

We now deal with the binary swapping of the inverse function, and its c-differential uniformity. There is no need to consider c=0 for the c-differential uniformity, since we already know that G is a permutation. Thus, in this paper we consider $c \neq 0,1$. We will be using the notation $i \mapsto c\Delta_G(a,b)$ to mean a contribution of i will be added to $c\Delta_G(a,b)$.

Theorem 3. Let $n \geq 2$ be a positive integer, $0, 1 \neq c \in \mathbb{F}_{2^n}$ and $F: \mathbb{F}_{2^n} \to \mathbb{F}_{2^n}$ be the inverse function defined by $F(x) = x^{2^n-2}$ and G be its (0,1)-swapping. If n=2, then ${}_c\Delta_G(a,b) \leq 1$; if n=3, ${}_c\Delta_G(a,b) \leq 3$. If $n \geq 4$, and for any $a,b \in \mathbb{F}_{p^n}$, the c-DDT entries satisfy ${}_c\Delta_G(a,b) \leq 4$ (all [1,2,3,4] c-DDT entries occur). Furthermore, ${}_c\Delta_G(a,b) = 4$ (so, the c-differential uniformity of G is $\delta_{F,c}=4$) if and only if any of the conditions happen:

- (i) For $a \in \mathbb{F}_{2^n}^*$ with $\text{Tr}\left(\frac{a}{a+1}\right) = 0$, $b = \frac{1}{a+1}$ and $c = \frac{1}{a^2+a}$, then $c\Delta_G(a, \frac{1}{a+1}) = 4$;
- (ii) For $a \in \mathbb{F}_{2^n}^*$ with $\operatorname{Tr}\left(\frac{a}{(a+1)^2}\right) = 0$, $b = \frac{1}{a^2}$, and $c = \frac{a+1}{a^2}$, then $c\Delta_G(a, \frac{1}{a^2}) = 4$.

Proof. We display first the c-differential equation

$$(x+a)^{2^{n}-2} + (x+a)^{2^{n}-1} + (x+a+1)^{2^{n}-1} + c\left(x^{2^{n}-2} + x^{p^{n}-1} + (x+1)^{2^{n}-1}\right) = b.$$
(1)

We ran a simple SageMath code and found easily the claims about n=2,3,4, so we now assume $n\geq 5$.

For $a, b \in \mathbb{F}_{2^n}$, we consider the *c*-differential equation (1). If a = 0, then (1) becomes

$$b = x^{2^{n}-2} + x^{2^{n}-1} + (x+1)^{2^{n}-1} + c\left(x^{2^{n}-2} + x^{2^{n}-1} + (x+1)^{2^{n}-1}\right)$$

= $(1+c)G(x)$.

Since G is also a permutation, then there exists a unique solution x, regardless of what $c \neq 1, b \in \mathbb{F}_{2^n}$. Thus, $_c\Delta_G(0,b) = 1$. From here on forward, we will assume that $a \neq 0$.

Case 1. Let a = 1, b = 0. Then (1) becomes

$$(x+1)^{2^{n}-2} + (x+1)^{2^{n}-1} + x^{2^{n}-1} + c\left(x^{2^{n}-2} + x^{2^{n}-1} + (x+1)^{2^{n}-1}\right) = 0.$$

Surely, x = 0 is a solution if and only if 1 + 1 + 0 + c(0 + 0 + 1) = 0, that is, c = 0, a contradiction. If x = 1, then 0 + 0 + 1 + c(1 + 1 + 0) = b, so b = 1, a contradiction. If $x \neq 0, 1$, then multiplying the displayed equation by x(x + 1), renders x + c(x + 1) = 0, and so, $x = \frac{c}{c+1}$, which implies $1 \mapsto_c \Delta_G(1, 0)$.

Case 2. Let a = 1, b = 1. If x = 0 in (1), then we must have 1 + 1 + 0 + c(0 + 0 + 1) = 1, so c = 1, a contradiction. If x = 1 in (1), then we must

have 0 + 0 + 1 + c(1 + 1 + 0) = 1, which is always true. If $x \neq 0, 1$, then multiplying the displayed equation by x(x + 1), we get

$$x + c(x + 1) = x(x + 1)$$
, that is, $x^2 + cx + c = 0$.

By Lemma 1, since $c \neq 0$, two solutions (surely, x = 1 cannot be one such solution) exist if and only if $\text{Tr}(c^2) = \text{Tr}(c) = 0$, therefore, we have $3 \mapsto {}_{c}\Delta_{G}(1,1)$ (including the prior x = 1), under this condition, and a contribution of 2, otherwise.

Case 3. Let $a=1, b \neq 0, 1$. If x=0, then we must have 1+1+0+c(0+0+1)=b, that is, b=c. Thus, $1 \mapsto {}_{c}\Delta_{G}(1,c)$. If x=1, then we must have 0+0+1+c(1+1+0)=b, and so, b=1, a contradiction. If $x \neq 0, 1$, multiplying Equation (1) by x(x+1) gives us

$$x + c(x+1) = bx(x+1)$$
, that is, $x^2 + \frac{b+c+1}{b}x + \frac{c}{b} = 0$,

which, by Lemma 1, under $b+c+1\neq 0$, has two distinct solutions if and only if $\operatorname{Tr}\left(\frac{bc}{(b+c+1)^2}\right)=0$. Thus, if b=c, and $\operatorname{Tr}(c)=0$, then $3\mapsto_c\Delta_G(1,c)$, and if b=c, $\operatorname{Tr}(c)=1$, then $1\mapsto_c\Delta_G(1,c)$. If $b\neq c$, and $\operatorname{Tr}\left(\frac{bc}{(b+c+1)^2}\right)=0$, then $2\mapsto_c\Delta_G(1,b)$.

Case 4. Let $a \neq 0, 1$. If x = 0, then we must have $\frac{1}{a} + 1 + 1 + c(0 + 0 + 1) = b$, so $b = c + \frac{1}{a}$. Thus, $1 \mapsto_{c} \Delta_{G}(a, c + \frac{1}{a})$. If x = 1, then $\frac{1}{a+1} + 1 + 1 + c(1+1+0) = b$, so $b = \frac{1}{a+1}$. Hence, $1 \mapsto_{c} \Delta_{G}(a, \frac{1}{a+1})$. If x = a, then $0 + 0 + 1 + c(\frac{1}{a} + 1 + 1) = b$, so $b = 1 + \frac{c}{a}$. Therefore, $1 \mapsto_{c} \Delta_{G}(a, 1 + \frac{c}{a})$. If x = a + 1, then $1 + 1 + 0 + c(\frac{1}{a+1} + 1 + 1) = b$, so $b = \frac{c}{a+1}$. Thus, $1 \mapsto_{c} \Delta_{G}(a, \frac{c}{a+1})$. Assuming $x \neq 0, 1, a, a + 1$, multiplying Equation (1) by x(x + a) implies

$$x + c(x + a) = bx(x + a)$$
, that is, $x^2 + \frac{ab + c + 1}{b}x + \frac{ac}{b} = 0$.

By Lemma 1, under $ab+c+1\neq 0$, this last equation has two distinct solutions if and only if $\operatorname{Tr}\left(\frac{abc}{(ab+c+1)^2}\right)=0$. We now summarize the largest contributions to $_c\Delta_G(a,b)$: if b=c+1

We now summarize the largest contributions to $_c\Delta_G(a,b)$: if $b=c+\frac{1}{a}=\frac{1}{a+1}$ (so, $c=\frac{1}{a^2+a}$; observe that $Tr(\frac{1}{c})=0$) and $Tr\left(\frac{abc}{(ab+c+1)^2}\right)=Tr\left(\frac{a^2}{(a+1)^2}\right)=Tr\left(\frac{a}{a+1}\right)=0$, then $4\mapsto {}_c\Delta_G(a,\frac{1}{a+1})$; if $b=c+\frac{1}{a}=\frac{c}{a+1}$ (so, $c=\frac{a+1}{a^2}$ and $b=\frac{1}{a^2}$; observe that Tr(c)=0) and $Tr\left(\frac{a}{(a+1)^2}\right)=0$,

then $4 \mapsto_c \Delta_G(a, \frac{1}{a^2})$. All values [1, 2, 3, 4] as c-DDT entries will occur, and we argue that next. For the values 1, 2, 3, that is obvious from our case splitting. The value 4 will occur if we can find parameters satisfying the previous conditions, and we achieve that in the following way. We can take $\beta \in \mathbb{F}_{2^n}^*$ such that $\text{Tr}(\beta) = \text{Tr}(1)$ and set $a := \frac{1}{\beta} + 1$, therefore, $\text{Tr}\left(\frac{a}{a+1}\right) = \text{Tr}(1) + \text{Tr}\left(\frac{1}{a+1}\right) = \text{Tr}(1) + \text{Tr}(\beta) = 0$. Thus, the second condition is satisfied, and therefore $\delta_{F,c} \leq 4$ with the upper bound being attained.

3 The c-boomerang uniformity of the swapped inverse function

We now investigate the c-BCT for the same swapped inverse function. It was shown in [18] that if c = 1, then the boomerang uniformity of G_{01} is 10, if $n \equiv 0 \pmod{6}$, 8, if $n \equiv 3 \pmod{6}$, and 6, if $n \not\equiv 0 \pmod{3}$. We will be using the notation $i \mapsto {}_{c}\mathcal{B}_{G}(a,b)$ to mean a contribution of i will be added to ${}_{c}\mathcal{B}_{G}(a,b)$.

We will be using throughout the c-boomerang system for the (0,1)swapping of the binary inverse function:

$$(x+\gamma)^{p^{n}-2} + (x+\gamma)^{p^{n}-1} + (x+\gamma+1)^{p^{n}-1} - c\left(x^{p^{n}-2} + x^{p^{n}-1} + (x+1)^{p^{n}-1}\right) = b,$$

$$(x+\gamma+a)^{p^{n}-2} + (x+\gamma+a)^{p^{n}-1} + (x+\gamma+a+1)^{p^{n}-1} - c^{-1}\left((x+a)^{p^{n}-2} + (x+a)^{p^{n}-1} + (x+a+1)^{p^{n}-1}\right) = b.$$
(2)

Theorem 4. Let $n \geq 2$ be a positive integer, $0, 1 \neq c \in \mathbb{F}_{2^n}$ and $F : \mathbb{F}_{2^n} \to \mathbb{F}_{2^n}$ be the inverse function defined by $F(x) = x^{2^n-2}$ and G be its (0,1)-swapping. If n = 2, $_c\mathcal{B}_G(a,b) = 1$; if n = 3, for $a,b \in \mathbb{F}_{2^n}$, $_c\mathcal{B}_G(a,b) \leq 4$ (all [1,2,3,4] entries occur). If $n \geq 4$, and for any $a,b \in \mathbb{F}_{2^n}$, the c-BCT entries satisfy $_c\mathcal{B}_G(a,b) \leq 5$. The upper bound is attained (at least) for $n \equiv 0 \pmod{3}$, $n \equiv 0 \pmod{4}$, $n \equiv 0 \pmod{5}$.

Proof. The instances n=2,3 were treated with a SageMath program. We now assume that $n \geq 4$. We do not need to consider the cases of a=0, since, then ${}_{c}\mathcal{B}_{G}(0,b) = {}_{c}\Delta_{G}(0,b)$, which was considered earlier.

Let a = 1. The c-boomerang system (2) becomes

$$(x+\gamma)^{2^{n}-2} + (x+\gamma)^{2^{n}-1} + (x+\gamma+1)^{2^{n}-1} + c\left(x^{p^{n}-2} + x^{p^{n}-1} + (x+1)^{p^{n}-1}\right) = b,$$

$$(x+\gamma+1)^{2^{n}-2} + (x+\gamma+1)^{2^{n}-1} + (x+\gamma)^{2^{n}-1} + \frac{1}{c}\left((x+1)^{p^{n}-2} + x^{p^{n}-1} + (x+1)^{p^{n}-1}\right) = b.$$
(3)

If x = 0, then $b = \gamma^{2^n - 2} + \gamma^{2^n - 1} + (\gamma + 1)^{2^n - 1} + c = (\gamma + 1)^{2^n - 2} + \gamma^{2^n - 1} + (\gamma + 1)^{2^n - 1}$. If $\gamma = 0, 1$, then c = 1, a contradiction. If $\gamma \neq 0, 1$, then $b = \frac{c}{c + 1}$ and $1 \mapsto {}_{c}\mathcal{B}_{G}\left(1, \frac{c}{c + 1}\right)$.

Next, if x = 1, then $b = (\gamma + 1)^{2^n - 2} + (\gamma + 1)^{2^n - 1} + \gamma^{2^n - 1} = \gamma^{2^n - 2} + \gamma^{2^n - 1} + (\gamma + 1)^{2^n - 1} + \frac{1}{c}$. If $\gamma = 0, 1$ we get a contradiction. If $\gamma \neq 0, 1$, then $b = \frac{1}{\gamma + 1} = \frac{1}{\gamma} + \frac{1}{c}$, rendering $b = \frac{1}{c + 1}$ and so, $1 \mapsto_c \mathcal{B}_G\left(1, \frac{1}{c + 1}\right)$.

If $x = \gamma$, then $b = 1 + \frac{c}{x} = \frac{1}{c(x+1)}$, so $cb^2 + (c^2 + c + 1)b + 1 = 0$. There exists a solution $b = c^{1/2}$ if $c^2 + c + 1 = 0$ and so, $1 \mapsto_c \mathcal{B}_G(1, c^{1/2})$, or two solutions b if $c^2 + c + 1 \neq 0$ and $\operatorname{Tr}\left(\frac{c}{(c^2 + c + 1)^2}\right) = 0$, and therefore $1 \mapsto_c \mathcal{B}_G(1, b)$, for any such b.

Similarly, if $x = \gamma + 1$, the obtained equation is $cb^2 + (c^2 + c + 1)b + c^2 = 0$. If $c^2 + c + 1 = 0$, then $b = c^{1/2}$ and $1 \mapsto_c \mathcal{B}_G(1, c^{1/2})$, and if $c^2 + c + 1 \neq 0$ and $\operatorname{Tr}\left(\frac{c^3}{(c^2 + c + 1)^2}\right) = 0$, and therefore $1 \mapsto_c \mathcal{B}_G(1, b)$, for any such b.

If $x \neq 0, 1, \gamma, \gamma + 1$, the system is then $b = \frac{1}{x+\gamma} + \frac{c}{x} = \frac{1}{x+\gamma+1} + \frac{1}{c(x+1)}$, implying that $b^2cx^2 + (1+c+b)(1+c+bc)x + c(1+c+bc) = 0$. If b = c+1, or $b = \frac{c+1}{c}$, there is a unique solution b, and if $c+1 \neq b \neq \frac{c+1}{c}$ and $\operatorname{Tr}\left(\frac{b^2c^2}{(1+c+bc)(1+c+b)^2}\right)$, then $1 \mapsto {}_c\mathcal{B}_G(1,b)$, for any such b.

We quickly see that the maximum claimed c-BCT entry cannot be achieved when a = 1.

From here on forward, we take $a \neq 0, 1$. We will consider the cases of (x, γ) being one of:

$$(0,0), (1,1), (0,1), (1,0), (0,a)_{a\neq0,1}, (a,0)_{a\neq0,1}, (0,a+1)_{a\neq0,1}, (a+1,0)_{a\neq0,1}, (1,a)_{a\neq0,1}, (a,1)_{a\neq0,1}, (1,a+1)_{a\neq0,1}, (a+1,1)_{a\neq0,1}, (a+1,1)_{a\neq0,1}, (a+1,a+1)_{a\neq0,1}, (a+1,a+1)_{a\neq0,1}, (a+1,a)_{a\neq0,1}, (a+1,a)_{a\neq0,1$$

Case 1. Let $x = \gamma = 0$. The c-boomerang system becomes (recall that $a \neq 0, 1$), $b = c + 1 = \frac{1}{a} + \frac{1}{ac} = \frac{c+1}{ac}$. Thus, $a = \frac{1}{c}, b = c+1$, so $1 \mapsto {}_{c}\mathcal{B}_{G}\left(\frac{1}{c}, c+1\right)$.

Case 2. Let $x = \gamma = 1$. The c-boomerang system becomes transforms into $b = 1 = \frac{1}{a} + \frac{1}{c(a+1)}$, implying a(a+1)c + c(a+1) + a = 0, that is,

$$a^2c + a + c = 0, (4)$$

which, by Lemma 1 has solutions a if and only $\operatorname{Tr}(c^2) = \operatorname{Tr}(c) = 0$ (using Hilbert's Theorem 90), so $1 \mapsto_{c} \mathcal{B}_{G}(a, 1)$, for such an a.

Case 3. Let $(x, \gamma) = (0, 1)$. Then, we must have $b = c = \frac{1}{a+1} + \frac{1}{ac}$, so,

$$a^{2}c^{2} + a(c^{2} + c + 1) + 1 = 0. (5)$$

If $c^2 + c + 1 = 0$, then $a = \frac{1}{c}$, so, $1 \mapsto_c \mathcal{B}_G\left(\frac{1}{c}, c\right)$, and if $c^2 + c + 1 \neq 0$ and $\operatorname{Tr}\left(\frac{c}{c^2 + c + 1}\right) = 0$, then $1 \mapsto_c \mathcal{B}_G\left(a, c\right)$.

Case 4. Let $(x, \gamma) = (1, 0)$. Then, we must have $b = 0 = \frac{1}{a+1} + \frac{1}{c(a+1)}$, so c = 1, which is not allowed.

Case 5. Let $(x, \gamma) = (0, a)$, $a \neq 0, 1$. We must have $b = \frac{1}{a} + c = 1 + \frac{1}{ca}$, so $a = \frac{1}{c}, b = 0$ and $1 \mapsto {}_c\mathcal{B}_G\left(\frac{1}{c}, 0\right)$.

Case 6. Let $(x, \gamma) = (a, 0)$, $a \neq 0, 1$. Then, $b = \frac{1}{a} + \frac{c}{a} = 1 + \frac{1}{c}$, so $a = c, b = 1 + \frac{1}{c}$, and $1 \mapsto_{c} \mathcal{B}_{G}\left(c, 1 + \frac{1}{c}\right)$.

Case 7. Let $(x, \gamma) = (0, a+1)$, $a \neq 0, 1$. Then, $b = \frac{1}{a+1} + c = \frac{1}{ac}$, so Equation (5) holds. If $c^2 + c + 1 = 0$, then $a = \frac{1}{c}, b = 1$, so, $1 \mapsto_c \mathcal{B}_G(\frac{1}{c}, 1)$, and if $c^2 + c + 1 \neq 0$ and $\text{Tr}\left(\frac{c}{c^2 + c + 1}\right) = 0$, then $1 \mapsto_c \mathcal{B}_G(a, \frac{1}{ac})$.

Case 8. Let $(x, \gamma) = (a + 1, 0)$, $a \neq 0, 1$. The c-boomerang system is then $b = \frac{1}{a+1} + \frac{c}{a+1} = 0$, so c = 1, a contradiction.

Case 9. Let $(x, \gamma) = (1, a), a \neq 0, 1$. We must have $b = \frac{1}{a+1} = \frac{1}{c(a+1)}$, so c = 1, a contradiction.

Case 10. Let $(x, \gamma) = (a, 1), a \neq 0, 1$. Then, $b = \frac{1}{a+1} + \frac{c}{a} = \frac{1}{c}$, so

$$a^2 + (c^2 + c + 1)a + c^2 = 0.$$

If $c^2 + c + 1 = 0$, $1 \mapsto_c \mathcal{B}_G\left(c, \frac{1}{c}\right)$, and if $c^2 + c + 1 \neq 0$ and $\operatorname{Tr}\left(\frac{c}{c^2 + c + 1}\right) = 0$, then $1 \mapsto_c \mathcal{B}_G\left(a, \frac{1}{c}\right)$.

Case 11. Let $(x, \gamma) = (1, a+1)$, $a \neq 0, 1$. Thus, $b = \frac{1}{a} = 1 + \frac{1}{c(a+1)}$, implying Equation (4). Thus, if $\operatorname{Tr}(c) = 0$, then $1 \mapsto_{c} \mathcal{B}_{G}\left(a, \frac{1}{a}\right)$.

Case 12. Let $(x, \gamma) = (a + 1, 1)$, $a \neq 0, 1$. Then, $b = \frac{1}{a} + \frac{c}{a+1} = 1$, so $a^2 + ac + 1 = 0$.

If $\operatorname{Tr}\left(\frac{1}{c}\right) = 0$, then $1 \mapsto {}_{c}\mathcal{B}_{G}\left(a, 1\right)$.

Case 13. Let $(x, \gamma) = (a, a), a \neq 0, 1$. Then, we must have $b = 1 + \frac{c}{a} = \frac{1}{a} + \frac{1}{c}$, so a = c, b = 0. Thus, $1 \mapsto_{c} \mathcal{B}_{G}(c, 0)$.

Case 14. Let $(x, \gamma) = (a + 1, a + 1), a \neq 0, 1$. Then, $b = 1 + \frac{c}{a+1} = \frac{1}{a}$, that is,

$$a^2 + ac + 1 = 0.$$

Therefore, if $\operatorname{Tr}\left(\frac{1}{c}\right) = 0$, $1 \mapsto {}_{c}\mathcal{B}_{G}\left(a, \frac{1}{a}\right)$.

Case 15. Let $(x, \gamma) = (a, a+1)$, $a \neq 0, 1$. Then, we must have $b = \frac{c}{a} = \frac{1}{a+1} + \frac{1}{c}$, and so, $a^2 + (c^2 + c + 1)a + c^2 = 0$. If $c^2 + c + 1 = 0$, $1 \mapsto_c \mathcal{B}_G(c, 1)$, and if $c^2 + c + 1 \neq 0$ and $\text{Tr}\left(\frac{c}{c^2 + c + 1}\right) = 0$, then $1 \mapsto_c \mathcal{B}_G\left(a, \frac{c}{a}\right)$.

Case 16. Let $(x, \gamma) = (a+1, a), a \neq 0, 1$. We must have $b = \frac{c}{a+1} = \frac{1}{a+1}$ that is, c = 1, a contradiction.

Case 17. Let $(x, \gamma) = (x, x)$, $x \neq 0, 1, a, a + 1$. Thus, $b = 1 + \frac{c}{x} = \frac{1}{a} + \frac{1}{c(x+a)}$, so

$$a^{2}cb^{2} + ba(c^{2} + c + 1 + ac) + ac + a + c^{2} = 0.$$
 (6)

Equation (6) will have one solution $b = \frac{1}{c^2+c+1}$ when $a = \frac{c^2+c+1}{c}$ (observe that $c^2+c+1 \neq 0$, since otherwise, c = 0), and so, $1 \mapsto {}_c\mathcal{B}_G\left(\frac{c^2+c+1}{c}, \frac{1}{c^2+c+1}\right)$,

and two b's if $\operatorname{Tr}\left(\frac{c(ac+a+c^2)}{(ac+c^2+c+1)^2}\right) = \operatorname{Tr}\left(\frac{c}{ac+c^2+c+1} + \frac{c^2}{(ac+c^2+c+1)^2} + \frac{c(a+1)}{(ac+c^2+c+1)^2}\right) = \operatorname{Tr}\left(\frac{c(a+1)}{(ac+c^2+c+1)^2}\right) = 0$, and, for such a, b, we have $1 \mapsto {}_{c}\mathcal{B}_{G}(a, b)$.

Case 18. Let $(x, \gamma) = (x, x+1), x \neq 0, 1, a, a+1$. Then, $b = \frac{c}{x} = \frac{1}{a+1} + \frac{1}{c(x+a)}$, so

$$a(a+1)cb^{2} + b(1+a+ac+c^{2}+ac^{2}) + c^{2} = 0,$$

which has a unique solution $b = \frac{c^2+c+1}{c+1}$ if $a = \frac{c^2+1}{c^2+c+1}$, and consequently, $1 \mapsto_c \mathcal{B}_G\left(\frac{c^2+1}{c^2+c+1}, \frac{c^2+c+1}{c+1}\right)$, or two solutions b if and only if $a \neq \frac{c^2+1}{c^2+c+1}$ and $\operatorname{Tr}\left(\frac{c^3a(a+1)}{(1+a+ac+c^2+ac^2)^2}\right) = 0$, when $1 \mapsto_c \mathcal{B}_G(a,b)$, for such a,b.

Case 19. Let $(x,\gamma)=(0,\gamma),\ \gamma\neq 0,1,a,a+1.$ The system becomes $b=\frac{1}{\gamma}+c=\frac{1}{\gamma+a}+\frac{1}{ca},$ implying

$$a^{2}cb^{2} + ba(1 + ac^{2}) + ac^{2} + ac + 1 = 0.$$
 (7)

If $a = \frac{1}{c^2}$, then b = c and $1 \mapsto_c \mathcal{B}_G\left(\frac{1}{c^2}, c\right)$. If $a \neq \frac{1}{c^2}$ and $\operatorname{Tr}\left(\frac{a^2c(ac^2 + ac + 1)}{(1 + ac^2)^2}\right) = \operatorname{Tr}\left(\frac{a^2c}{1 + ac^2} + \frac{a^4c^2}{(1 + ac^2)^2} + \frac{a^3(a + 1)c^2}{(1 + ac^2)^2}\right) = \operatorname{Tr}\left(\frac{a^3(a + 1)c^2}{(1 + ac^2)^2}\right) = 0$, then $1 \mapsto_c \mathcal{B}_G(a, b)$, for a, b satisfying Equation (7).

Case 20. Let $(x, \gamma) = (1, \gamma), \ \gamma \neq 0, 1, a, a + 1$. Thus, $b = \frac{1}{\gamma + 1} = \frac{1}{\gamma + a + 1} + \frac{1}{c(a+1)}$, so

$$a(a+1)cb^2 + ab + 1 = 0.$$

Thus, if $\operatorname{Tr}\left(\frac{(a+1)c}{a}\right) = 0$, then $1 \mapsto {}_{c}\mathcal{B}_{G}\left(a,b\right)$.

Case 21. Let $(x,\gamma)=(a,\gamma), \ \gamma\neq 0,1,a,a+1$. The system becomes $b=\frac{1}{\gamma+a}+\frac{c}{a}=\frac{1}{\gamma}+\frac{1}{c}$, so

$$a^{2}cb^{2} + ba(a+c^{2}) + c^{2} + ac + a = 0.$$
 (8)

If $a=c^2$, then $b=\frac{1}{c}$ and $1\mapsto {}_c\mathcal{B}_G\left(c^2,\frac{1}{c}\right)$, and if $a\neq c^2$ and $\operatorname{Tr}\left(\frac{c(c^2+ac+a)}{(a+c^2)^2}\right)=\operatorname{Tr}\left(\frac{c}{a+c^2}+\frac{a}{a+c^2}+\frac{a^2}{(a+c^2)^2}\right)=\operatorname{Tr}\left(\frac{c}{a+c^2}\right)=0$ (we used Hilbert's Theorem 90), then $1\mapsto {}_c\mathcal{B}_G\left(a,b\right)$, for a,b satisfying Equation (8).

Case 22. Let $(x, \gamma) = (a + 1, \gamma), \ \gamma \neq 0, 1, a, a + 1$. Then, $b = \frac{1}{\gamma + a + 1} + \frac{c}{a + 1} = \frac{1}{\gamma + 1}$, so

$$a(a+1)b^{2} + acb + c = 0. (9)$$

Thus, if $\operatorname{Tr}\left(\frac{a+1}{ac}\right) = 0$, then $1 \mapsto {}_{c}\mathcal{B}_{G}\left(a,b\right)$.

Case 23. Let $(x, \gamma) = (x, x+a), x \neq 0, 1, a, a+1, a \neq 0, 1$. Thus, $b = \frac{1}{a} + \frac{c}{x} = 1 + \frac{1}{c(x+a)}$, so

$$a^{2}cb^{2} + ba(c^{2} + c + 1 + ac) + 1 + ac + ac^{2} = 0.$$
 (10)

If $a = \frac{c^2 + c + 1}{c}$, then $b = \frac{c^2}{c^2 + c + 1}$, and $1 \mapsto_c \mathcal{B}_G\left(\frac{c^2 + c + 1}{c}, \frac{c^2}{c^2 + c + 1}\right)$. If $a \neq \frac{c^2 + c + 1}{c}$ and $\operatorname{Tr}\left(\frac{c(ac^2 + ac + 1)}{(c^2 + c + 1 + ac)^2}\right) = \operatorname{Tr}\left(\frac{c^2}{(ac + c^2 + c + 1)^2} + \frac{c}{ac + c^2 + c + 1} + \frac{(a + 1)c^3}{(ac + c^2 + c + 1)^2}\right) = \operatorname{Tr}\left(\frac{(a + 1)c^3}{(ac + c^2 + c + 1)^2}\right) = 0$, then $1 \mapsto_c \mathcal{B}_G(a, b)$.

Case 24. Let $(x,\gamma)=(x,x+a+1),\ x\neq 0,1,a,a+1,\ a\neq 1.$ Then, $b=\frac{1}{a+1}+\frac{c}{x}=\frac{1}{c(x+a)},$ so

$$b^{2}a(a+1)c + b(1+a+ac+c^{2}+ac^{2}) + 1 = 0.$$

If $a = \frac{c^2+1}{c^2+c+1}$, $b = \frac{c^2+c+1}{c(c+1)}$, and $1 \mapsto_c \mathcal{B}_G\left(\frac{c^2+1}{c^2+c+1}, \frac{c^2+c+1}{c(c+1)}\right)$, and if $a \neq \frac{c^2+1}{c^2+c+1}$ and $\operatorname{Tr}\left(\frac{a(a+1)c}{(1+a+ac+c^2+ac^2)^2}\right) = 0$, then $1 \mapsto_c \mathcal{B}_G(a,b)$.

Case 25. Let $x + \gamma \neq 0, 1, a, a + 1, x \neq 0, 1, a, a + 1$. Then, we must have $b = \frac{1}{x+\gamma} + \frac{c}{x} = \frac{1}{x+\gamma+a} + \frac{1}{c(x+a)}$, which implies

$$bx^{2} + cy + x(1+c+by) = 0, a+ac+a^{2}bc+bcx^{2} + (1+abc)y + x(1+c+bcy) = 0.$$

Adding to the second equation, a c multiple of the first gives us the system (the advantage is that it shows the linear relationship between x, y), and replacing y into the second displayed equation, we get

$$a + ac + a^{2}bc + (1 + c^{2})x + (1 + abc + c^{2})y = 0,$$

$$ab^{2}c x^{2} + (1 + ab + c)(1 + c + abc)x + ac(1 + c + abc) = 0.$$
(11)

If $b = \frac{c+1}{a}$, or $b = \frac{c+1}{ac}$, then the second equation in (11) has one solution x and if $\frac{c+1}{a} \neq b \neq \frac{c+1}{ac}$ and $\operatorname{Tr}\left(\frac{a^2b^2c^2}{(1+ab+c)^2(1+c+abc)}\right) = 0$ it has two solutions. In the first case, $1 \mapsto {}_c\mathcal{B}_G(a,b)$, and in the second case, $2 \mapsto {}_c\mathcal{B}_G(a,b)$ (for the same (a,b,c) tuple, each of the solutions x will render a solution y: that is the reason we wrote one of the equations as an affine plane).

Now, we need to see how the multiple (a, b) pairs (with or without conditions) overlap, as the contributions to the various c-BCT entries will be added. For easy referral, we shall combine them in a set where we put an asterisk if there are conditions on (a, b) (if the conditions are different, then we repeat that pair):

$$S = \left\{ \left(\frac{1}{c}, c+1 \right), (a,1)^*, \left(\frac{1}{c}, c \right)^*, (a,c)^*, (c,0), \left(a, \frac{1}{a} \right)^*, (c,1)^*, \left(a, \frac{c}{a} \right)^*, \right.$$

$$\left(\frac{1}{c}, 0 \right), \left(c, 1 + \frac{1}{c} \right), \left(\frac{1}{c}, 1 \right)^*, \left(a, \frac{1}{ac} \right)^*, (a,c)^*, \left(c, \frac{1}{c} \right)^*, \left(a, \frac{1}{c} \right)^*, \left(a, \frac{1}{a} \right)^*, \right.$$

$$\left(a, 1 \right)^*, \left(\frac{c^2 + c + 1}{c}, \frac{1}{c^2 + c + 1} \right), (a,b)^*, \left(\frac{c^2 + 1}{c^2 + c + 1}, \frac{c^2 + c + 1}{c + 1} \right), \left. (a,b)^*, \left(\frac{1}{c^2}, c \right), (a,b)^*, (a,b)^*, \left(\frac{c^2 + 1}{c^2 + c + 1}, \frac{c^2 + c + 1}{c^2 + c} \right), (a,b)^*, (a,b)^* \right\}.$$

$$\left(\frac{c^2 + c + 1}{c}, \frac{c^2}{c^2 + c + 1} \right), (a,b)^*, \left(\frac{c^2 + 1}{c^2 + c + 1}, \frac{c^2 + c + 1}{c^2 + c} \right), (a,b)^*, (a,b)^* \right\}.$$

We will prune the above set, concentrating on the maximal entry value ${}_{c}\mathcal{B}_{G}(a,b)$, which we claim to be 5. Going through every possibility (we shall go through similar arguments enabling us to motivate the maximum c-BCT entry below, so we skip the previous straightforward pruning argument, albeit tedious), we obtained that the only possible pairs that could possibly overlap five times (we write the constraints next to each of them) are in the

displayed list below (recall that p_{10} contributes twice to ${}_{c}\mathcal{B}_{G}(a,b)$). Since we shall refer to the respective pairs in the arguments below the list, by relabeling, we will call the pairs (a,b) satisfying the conditions from which E_{i} are derived, as coming from Case C_{i} . Also, if the pairs p_{i}, p_{j} match the conditions from cases C_{i}, C_{j} , we write $p_{i} = p_{j}$. We bring up an issue the reader should be aware of, which we alluded to previously: if the Case C_{i} holds, then $E_{i} = 0$; however, if $E_{i} = 0$, the pair (a, b) may or may not be p_{i} , so one has to continuously check. We find that to be the difficulty in the analysis below, along with the many cases.

$$C_0: p_0 = \left(a, \frac{1}{ac}\right): E_0 = a^2c^2 + (c^2 + c + 1)a + 1 = 0;$$

$$C_1: p_1 = \left(a, \frac{c}{a}\right): E_1 = a^2 + (c^2 + c + 1)a + c^2 = 0;$$

$$C_2: p_2 = (a, b): E_2 = a^2cb^2 + a(c^2 + c + 1 + ac)b + ac + a + c^2 = 0;$$

$$C_3: p_3 = (a, b): E_3 = a(a + 1)cb^2 + b(1 + a + ac + c^2 + ac^2) + c^2 = 0;$$

$$C_4: p_4 = (a, b): E_4 = a^2cb^2 + a(1 + ac^2)b + ac^2 + ac + 1 = 0;$$

$$C_5: p_5 = (a, b): E_5 = a(a + 1)cb^2 + ab + 1 = 0;$$

$$C_6: p_6 = (a, b): E_6 = a^2cb^2 + ba(a + c^2) + c^2 + ac + a = 0;$$

$$C_7: p_7 = (a, b): E_7 = a(a + 1)b^2 + acb + c = 0;$$

$$C_8: p_8 = (a, b): E_8 = a^2cb^2 + a(1 + c + c^2 + ac)b + 1 + ac + ac^2 = 0;$$

$$C_9: p_9 = (a, b): E_9 = a(a + 1)cb^2 + b(1 + a + ac + c^2 + ac^2) + 1 = 0;$$

$$C_{10}: p_{10} = (a, b): E_{10} = ab^2cx^2 + (1 + ab + c)(1 + c + abc)x + ac(1 + c + abc) = 0,$$
as well as $E_{10}' = a + ac + a^2bc + (1 + c^2)x + (1 + abc + c^2)y = 0;$

We make the following important observation: there is a duality between all of the E_i , $0 \le i \le 9$, expressions. By duality we mean that one expression is obtained from the other by replacing c by $\frac{1}{c}$. Precisely, $E_0 \longleftrightarrow E_1$, $E_2 \longleftrightarrow E_8$, $E_3 \longleftrightarrow E_9$, $E_4 \longleftrightarrow E_6$, $E_5 \longleftrightarrow E_7$.

And here is a simple argument that shows that in each dual pair, if one case occurs in a solution set, then the other cannot, which shows that the c-boomerang uniformity is upper bounded by 5 (we exclude two cases out of the possible 6 (recall that C_{10} contributes 2 to $_c\mathcal{B}_G(a,b)$) by showing that $\{p_0,p_1\}\cap\{p_5,p_7\}$, as well as $\{p_4,p_6\}\cap\{p_2,p_8\}$, if nonempty, cannot be part of quintuple solution sets).

If $p_0 = p_1$, then $b = \frac{c}{a} = \frac{1}{ac}$, implying c = 1, which is impossible. If $p_2 = p_8$, then $1 + a^2b^2c + a(b + b^2c) = 0$, $a^2b^2 + c + a(b^2 + bc) = 0$, If $p_3 = p_9$,

then adding $E_3+E_9=0$ we get c^2+1 , which is impossible. If $p_4=p_6$, then adding $(1+bc)E_4+c(b+c)E_6=0$, renders $b=\frac{bc+c^2+1}{b^2c+bc^2+bc+b+c}$, which when used in $bE_4+cE_6=0$, finds $b=\frac{c^2+c+1}{c}=0$ and a=1, which, when plugged back into p_4,p_6 imply $\frac{c\gamma_4+c+\gamma_4}{c\gamma_4}=\frac{c\gamma_4+c+\gamma_4}{\gamma_4+1}$ and $\frac{c\gamma_6+\gamma_6+1}{\gamma_6}=\frac{c\gamma_6+\gamma_6+1}{c(\gamma_6+1)}$, which gives c=1, an impossibility. If $p_5=p_7$, then adding $E_5+cE_7=0$, we get $b=\frac{1}{a}$, which, when replaced back, gives c(1+1/a)=0=1+1/a, and so, a=1. However, a=1,b=1, contradicts the conditions imposed in C_5,C_7 (Cases 20 and 21), since then $\gamma=0$. Further, if $p_0=p_5$, then $b=\frac{1}{ac}$, which used in E_5 gives $a=\frac{1}{c},b=1$ (and $c^2+c+1=0$). These values used in E_2,E_3,E_4,E_6,E_8,E_9 gives us

$$c + c^2, 1 + c, 1 + c, \frac{1}{c}, \frac{1+c}{c}, c + c^2,$$

and none can be zero. If $p_0 = p_7$, $b = \frac{1}{ac}$, then $a = \frac{1+c^2}{c}$, $b = \frac{1}{c^2+1}$ (and $c^5 + c^4 + c^3 + c + 1 = 0$). These values used in $E_2, E_3, E_4, E_6, E_8, E_9$ gives us

$$c, c^{2}(c^{2}+c+1), 1+c+c^{3}, \frac{1}{c^{2}}, c(c^{3}+1), \frac{c(c^{2}+c+1)}{(c+1)^{2}}.$$

If $c^2+c+1=0$, then $c^5+c^4+c^3+c+1=0$ implies c=1. If $c^3+c+1=0$, then c=0,1, which is not allowed. Next, we show that p_1 cannot be part of a quintuple solution along with any of the $\{p_5,p_7\}$. First, we assume that $p_1=p_5$ and so, $b=\frac{c}{a}$ and $a=\frac{c^3}{c^3+c+1}$. When plugged back into $E_1=0$, it implies $c^5+c^4+c^2+c+1=0$. Now, these values used in E_2,E_3,E_4,E_6,E_8,E_9 gives us

$$\frac{c+1}{c^5+1}, \frac{1+c}{c^2}, \frac{c}{1+c+c^3}, \frac{c^2(c^3+1)}{1+c+c^3}, c, \frac{(1+c)(1+c^2+c^3)}{c^2}.$$

Surely, if $c^3 + 1 = 0$, then $c^5 + c^4 + c^3 + c + 1 = c^4(c+1) \neq 0$, and therefore, we cannot make up a quintuple solution. If $p_1 = p_7$, then $b = \frac{c}{a}$, so a = c, and b = 1. Replacing these values into $E_2, E_3, E_4, E_6, E_8, E_9$ gives us

but none of these values can be zero.

We move on to the second dual pairs $\{p_4, p_6\} \cap \{p_2, p_8\}$. If $p_2 = p_4$, then adding $E_2 + E_4$ we get $b = \frac{1+c}{ac}$, which when put back into $E_2 = 0$, say, gives us c = 0, which is not allowed. If $p_2 = p_6$, then adding $E_2 + E_6$ implies ab(a+1)(c+1) = 0. If a = 0, then $E_2 = c^2$, which is not permitted. If

b=0, then $E_2=a+ac+c^2=0$, so $a=\frac{c^2}{c+1}$. Putting these values into $\{E_1,E_3,E_5,E_7,E_9\}$ renders

$$\left\{\frac{c^5}{(c+1)^2}, c^2, 1, c, 1\right\},\,$$

and none of these values can be zero, so no "winning" quintuple. If a=1, then $b=\frac{c^2+c+1}{c}$, which when used into $\{E_1,E_3,E_5,E_7,E_9\}$ gives

$$\left\{c, c+1, \frac{(c+1)^2}{c}, (c+1)^2, c(c+1)\right\}$$

which cannot be zero. If $p_4 = p_8$, adding $E_4 + E_8 = 0$, we get abc(a+1)(c+1). If a = 0, then $E_4 = 1$, which is not possible. If b = 0, then $a = \frac{1}{c^2+c}$, which, when used in $\{E_1, E_3, E_5, E_7, E_9\}$ gives

$$\left\{ \frac{c^6 + c + 1}{c^2(c+1)^2}, c^2, 1, c, 1 \right\},\,$$

so no possible quintuple. If a=1, then $b=\frac{c^2+c+1}{c}$, and $\{E_1,E_3,E_5,E_7,E_9\}$ become

$$\left\{c^2, \frac{c^4+c+1}{c}, 1, c, \frac{c^4+c^3+1}{c}\right\}.$$

We could possibly get a solution quintuple if $1+c+c^4=0$ and $1+c^3+c^4=0$, but that is impossible. If $p_6=p_8$, adding $E_6+E_8=0$, we get $b=\frac{1+c}{a}$, but with this value both E_6, E_8 become equal to c and consequently not zero.

The above discussion shows that ${}_{c}\mathcal{B}_{G}(a,b) \leq 5$. Since the rest of the proof (which is long enough, as it is) deals with finding appropriate cases where the upper bound is attained we moved it to Appendix A. The theorem is shown.

We make the following conjecture, based upon our extensive computations.

Conjecture 5. For all $n \ge 4$, then there exists $c \ne 0, 1$ such that $\beta_{F,c} = 5$.

4 Concluding remarks

In this paper we investigate the c-differential and c-boomerang uniformity of the swapped inverse function. We show that its c-differential uniformity is less than or equal to 4 and its c-boomerang uniformity is upper bounded

by 5. As we saw, investigating questions on c-differential and c-boomerang uniformities tends to be quite difficult, but it sure is interesting to find other classical PN/APN functions and study their properties through the new differential. It will also be worthwhile to check into the general p-ary versions of the results from this paper.

References

- [1] E. R. Berlekamp, H. Rumsey, G. Solomon, On the solutions of algebraic equations over finite fields, Information and Control 10 (1967), 553–564.
- [2] E. Biham, O. Dunkelman, N. Keller, New results on boomerang and rectangle attacks, In: Daemen J., Rijmen V. (eds.), Fast Software Encryption, FSE 2002, LNCS 2365, 2002, pp. 1–16, Springer, Berlin, Heidelberg.
- [3] A. Biryukov, D. Khovratovich, Related-key cryptanalysis of the full AES-192 and AES-256, In: Matsui M. (ed.), Adv. in Crypt. ASI-ACRYPT 2009, LNCS 5912, 2009, pp. 1–18, Springer, Berlin, Heidelberg.
- [4] A. W. Bluher, $On x^{q+1} + ax + b$, Finite Fields Appl. 10 (3) (2004), 285–305.
- [5] C. Boura, A. Canteaut, On the boomerang uniformity of cryptographic Sboxes, IACR Trans. Symmetric Cryptol. 3 (2018), 290–310.
- [6] C. Boura, L. Perrin, S. Tian, Boomerang Uniformity of Popular S-box Constructions, Workshop on Cryptography and Coding 2019, Paper 15, https://www.lebesgue.fr/sites/default/files/proceedings_WCC/WCC_2019_paper_15.pdf
- [7] N. Bourbaki, Elements of Mathematics, Algebra II (translated by P. M. Cohn and J. Howie), Springer, Berlin, 1990.
- [8] L. Budaghyan, Construction and Analysis of Cryptographic Functions, Springer-Verlag, 2014.
- [9] M. Calderini, I.Villa, On the Boomerang Uniformity of some Permutation Polynomials, https://eprint.iacr.org/2019/881.pdf.
- [10] C. Carlet, Boolean functions for cryptography and error correcting codes, In: Y. Crama, P. Hammer (eds.), Boolean Methods and Models, Cambridge Univ. Press, Cambridge, pp. 257–397, 2010.

- [11] C. Carlet, Vectorial Boolean Functions for Cryptography, In: Y. Crama, P. Hammer (eds.), Boolean Methods and Models, Cambridge Univ. Press, Cambridge, pp. 398–472, 2010.
- [12] C. Cid, T. Huang, T. Peyrin, Y. Sasaki, L. Song, Boomerang Connectivity Table: A new cryptanalysis tool, In: J. B. Nielsen and V. Rijmen (eds.), Adv. in Crypt. EUROCRYPT '18, pp. 683–714, Cham, 2018. Springer.
- [13] T. W. Cusick, P. Stănică, Cryptographic Boolean Functions and Applications (Ed. 2), Academic Press, San Diego, CA, 2017.
- [14] P. Ellingsen, P. Felke, C. Riera P. Stănică, A. Tkachenko, C-differentials, multiplicative uniformity and (almost) perfect cnonlinearity, to appear in IEEE Trans. Inf. Theory, 2020, https://doi.org/10.1109/TIT.2020.2971988.
- [15] N.S. Kaleyski, Changing APN functions at two points, Cryptogr. Commun. 11.6 (2019), 1165–1184.
- [16] J. Kelsey, T. Kohno and B. Schneier, Amplified boomerang attacks against reduced-round MARS and Serpent, In: Goos G., Hartmanis J., van Leeuwen J., Schneier B. (eds.), Fast Software Encryption, FSE 2000. LNCS 1978. Springer, Berlin, Heidelberg.
- [17] J. Kim, S. Hong, B. Preneel, E. Biham, O. Dunkelman, Related-Key Boomerang and Rectangle Attacks: Theory and Experimental Analysis, IEEE Trans. Inf. Theory 58(7) (2012), 4948–4966.
- [18] K. Li, L. Qu, B. Sun, C. Li, New results about the boomerang uniformity of permutation polynomials, IEEE Trans. Inf. Theory 65(11) (2019), 7542–7553.
- [19] N. Li, Z. Hu, M. Xiong, X. Zeng, 4-uniform BCT permutations from generalized butterfly structure, https://arxiv.org/abs/2001.00464.
- [20] Y. Li, M. Wang, Y. Yu, Constructing Differentially 4-Uniform Permutations Over $GF(2^{2k})$ From the Inverse Function Revisited, https://eprint.iacr.org/2013/731.
- [21] S. Mesnager, Bent functions: fundamentals and results, Springer Verlag, 2016.

- [22] S. Mesnager, C. Tang, M. Xiong, On the boomerang uniformity of quadratic permutations, https://eprint.iacr.org/2019/277.pdf.
- [23] J. Peng, C. H. Tan, New differentially 4-uniform permutations by modifying the inverse function on subfields, Cryptogr. Commun. 9 (2017), 363–378.
- [24] L. Qu, Y. Tan, C. H. Tan, C. Li, Constructing differentially 4-uniform permutations over $\mathbb{F}_{2^{2k}}$ via the switching method, IEEE Trans. Inf. Theory 59:4 (2013), 4675–4686.
- [25] L. Qu, Y. Tan, C. Li, and G. Gong, More constructions of differentially 4-uniform permutations on $\mathbb{F}_{2^{2k}}$, Des., Codes Cryptogr. 78 (2016), 391–408.
- [26] C. Riera, P. Stănică, *Investigations on c-(almost) perfect nonlinear functions*, https://arxiv.org/abs/2004.02245.
- [27] P. Stănică, Investigations on c-boomerang uniformity and perfect non-linearity, https://arxiv.org/abs/2004.11859, 2020.
- [28] D. Tang, C. Carlet, X. Tang, Differentially 4-uniform bijections by permuting the inverse function, Des. Codes. Cryptogr. 77 (2014), 117–141.
- [29] N. Tokareva, Bent Functions, Results and Applications to Cryptography, Academic Press, San Diego, CA, 2015.
- [30] Z. Tu, N. Li, X. Zeng, J. Zhou, A class of quadrinomial permutation with boomerang uniformity four, IEEE Trans. Inf. Theory, https://doi.org/10.1109/TIT.2020.2969578.
- [31] Y. Yu, M. Wang, Y. Li, Constructing differentially 4 uniform permutations from known ones, Chin. J. Electron. 22(3) (2013), 495–499.
- [32] D. Wagner, The boomerang attack, In: L. R. Knudsen (ed.), Fast Soft-ware Encryption, FSE '99, LNCS 1636, pages 156–170, 1999, Springer, Heidelberg.
- [33] Z. Zha, L. Hu, S. Sun, Constructing new differentially 4-uniform permutations from the inverse function, Finite Fields Appl. 25 (2014), 64–78.
- [34] H. Yan, S. Mesnager, Z. Zhou, Power Functions over Finite Fields with Low c-Differential Uniformity, https://arxiv.org/pdf/2003.13019.pdf.

Appendices

A Completing the proof of Theorem 4

Here, we finish the proof of Theorem 4. We cannot find a simple argument to show that 5 is always attained for all dimensions, but we can find some values for c such that ${}_c\mathcal{B}_G(a,b)=5$, namely, for $n\equiv 0\pmod 3$, $n\equiv 0\pmod 4$, $n\equiv 0\pmod 5$. We start by showing how we "guessed" a quintuple of cases that provide a solution set for the c-boomerang system. Recall that if the pair p_i happens then E_i will hold, but not necessarily the other way around.

If $p_0 = p_2$ (no need to consider $p_0 = p_1$ as they are dual), then $b = \frac{1}{ac}$, and so, $a = \frac{c^2 + c + 1}{c}$, $b = \frac{1}{c^2 + c + 1}$, along with $c^5 + c^4 + c^3 + c^2 + 1 = 0$. Using these values into $\{E_1, E_3, E_8, E_9\}$, we get

$$\left\{\frac{c^5+c^3+c^2+c+1}{c^2}, \frac{(c+1)^4}{c^2+c+1}, \frac{(c+1)^4}{c}, \frac{c(c+1)^2}{c^2+c+1}\right\},\,$$

and none of these values can be zero (if $c\neq 0,1$ and $c^5+c^4+c^3+c^2+1=0$). If $p_0=p_3$, then $b=\frac{1}{ac}$, and so, $a=\frac{c}{c^2+c+1},b=\frac{c^2+c+1}{c^2}$, along with $c^5+c^3+c^2+c+1=0$, which used in $E_1,E_2,E_4,E_5,E_6,E_7,E_8$ gives us

$$\left\{ \frac{c\left(c^5 + c^4 + c^3 + c^2 + 1\right)}{\left(c^2 + c + 1\right)^2}, \frac{(c+1)^4}{c^2 + c + 1}, \frac{(c+1)^3}{c^2 + c + 1}, \frac{c+1}{c^2}, \frac{(c+1)^3}{c}, \frac{(c+1)\left(c^3 + c + 1\right)}{c^3}, 0 \right\}.$$

We see that the only possibility is for $p_0 = p_3 = p_8$, and we can find x, y values such that $p_0 = p_3 = p_8 = p_{10}$ (under $c^5 + c^3 + c^2 + c + 1 = 0$, $n \equiv 0 \pmod{5}$) in the following way. The solutions x, y for C_{10} can be found by taking $b = \frac{1}{ac}$ and arriving at $a^2c^3 + ac(1+c+c^2)x + x^2 = 0$, $ac + (1+c^2)x + c^2y = 0$, for $E_{10} = E'_{10} = 0$. If $\text{Tr}\left(\frac{c}{c^2+c+1}\right) = 0$, then there is a solution x of the first equation, and consequently for y, using the second linear equation. We argue now why the trace is 0. We simply use the equation $c^5 + c^4 + c^3 + c^2 + 1 = 0$ and write it as $\frac{c^2}{(c^2+c+1)^2} = \frac{1}{c^3+c} = \frac{1}{c} + \frac{1}{c+1} + \frac{1}{(c+1)^2} = \frac{1}{c}$. Now, using the reciprocal polynomial of $c^5 + c^4 + c^3 + c^2 + 1$, we find that $\text{Tr}\left(\frac{1}{c}\right) = 0$. Thus, $c\mathcal{B}_G(a,b) = 5$ for $n \equiv 0 \pmod{5}$.

As an example for this argument, we take n = 5. The solutions for the c-boomerang system and the parameters (c, a, b, x, y) of the previous

argument (α is a primitive root, corresponding to the primitive polynomial $x^5 + x^2 + 1$) are (we used SageMath to find these solutions, in the finite field F_{25} given by the primitive polynomial $x^5 + x^2 + 1$:

 $(\alpha^4 + \alpha^2, \alpha^4, \alpha^3 + \alpha^2, 0, \alpha^4 + 1)$: solution from the case C_0 $(\alpha^4 + \alpha^2, \alpha^4, \alpha^3 + \alpha^2, \alpha^3 + \alpha^2 + \alpha, \alpha^4 + 1)$; solution from the case C_{10} $(\alpha^4 + \alpha^2, \alpha^4, \alpha^3 + \alpha^2, \alpha^4 + \alpha + 1, \alpha^3 + \alpha^2 + 1)$; solution from the case C_{10} $(\alpha^4 + \alpha^2, \alpha^4, \alpha^3 + \alpha^2, \alpha + 1, \alpha)$; solution from the case C_3 $(\alpha^4 + \alpha^2, \alpha^4, \alpha^3 + \alpha^2, \alpha^3 + \alpha + 1, \alpha^4 + \alpha^3 + \alpha + 1)$; solution from the case C_8 .

If $p_0 = p_4$, then $a = \frac{1}{c^2}$, b = c, which when used in $\{E_6, E_8, E_9\}$ finds the values

$$\left\{\frac{(c+1)^5}{c^3}, \frac{(c+1)^3}{c^2}, c(c+1)^2\right\}$$

and none of these will work. If $p_0 = p_6$, then $1 + c^2 + c^3 + a(1 + c + c^2) = 0$. So, if $c^2+c+1=0$, then $c^3+c^2+1=0$, and so c=0,1, which is not allowed. Thus, $a=\frac{c^3+c^2+1}{c^2+c+1}$, $b=\frac{c^2+c+1}{c(c^3+c^2+1)}$ and $E_0=0$ implies $c^5+c^4+c^2+c+1=0$ (thus, $n \equiv 0 \pmod{5}$, which used in $\{E_8, E_9\}$ gives us

$$\left\{ (c+1)^3, \frac{c^2(c+1)^2}{c^3+c^2+1} \right\},\,$$

and surely, none will work. If $p_0 = p_7$, then $a = \frac{1}{c^3 + c^2 + 1}$ ($c^3 + c^2 + 1 \neq 0$, since $E_4 = 0$ is equivalent to $1 + a(c^3 + c^2 + 1) = 0$), $b = \frac{c^3 + c^2 + 1}{c}$ and $c^5 + c^4 + c^3 + c + 1 = 0$. The values used in $\{E_8, E_9\}$ gives us

$$\left\{ \frac{(c+1)\left(c^3+c+1\right)}{c^3+c^2+1}, c^3(c+1) \right\},\,$$

and none of these values can be zero.

If $p_0 = p_8$, then $a^2c^2 + a(c^2 + c + 1) + 1 = 0$, $c + a(c^2 + c + 1) = 0$, $b = \frac{1}{ac}$. If $c^2 + c + 1 = 0$, then $a = \frac{1}{c}$, b = 1. When used in $E_9 = 0$, we get c(c + 1) = 0, which is not allowed. Thus $c^2 + c + 1 \neq 0$. Then $a = \frac{c}{c^2 + c + 1}$, $b = \frac{c^2 + c + 1}{c^2}$ and $c^5 + c^3 + c^2 + c + 1 = 0$ (thus, $n \equiv 0 \pmod{4}$), which put into $E_9 = 0$, gives us $(c+1)^2 = 0$. There is no need to check for further equalities involving p_0 . If $p_1 = p_2$, then $b = \frac{c}{a}$ and $E_1 = 0$, $E_2 = 0$ become $a^2 + c^2 + a(1+c+c^2) = 0$, $c + a(1+c+c^2) = 0$. Thus, $a = \frac{c}{c^2+c+1}$, $b = c^2+c+1$, and $E_1 = 0$ is transformed into $c^5 + c^4 + c^3 + c^2 + 1 = 0$ (which can always happen for some c if and only if $n \equiv 0 \pmod{5}$: that is easy to see since the underlying

irreducible polynomial $x^5 + x^4 + x^3 + x^2 + 1$ is a divisor of $x^{2^n - 1} + 1$ if and only if $n \equiv 0 \pmod{5}$. The values $a = \frac{c}{c^2 + c + 1}$, $b = c^2 + c + 1$ put back into E_3, E_4, E_6, E_8, E_9 (we use here the fact that p_0 and p_1, p_5, p_7 cannot be part of a winning "quintuple", via the prior discussion) renders

$$\left\{(c+1)^2,(c+1)^3,\frac{c(c+1)^3}{c^2+c+1},\frac{(c+1)^4}{c^2+c+1},0\right\}.$$

Thus $p_1 = p_2 = p_9$ and we can find (x, y), as before, such that $p_1 = p_2 = p_9 = p_{10}$, under $n \equiv 0 \pmod{5}$.

If $p_1 = p_3$, then a = 1, b = c, which when used in E_4, E_6, E_8, E_9 gives c = 1. If $p_1 = p_4$, then $a = \frac{c^3 + c + 1}{c(c^2 + c + 1)}$ (it is easy to see that $c^2 + c + 1 \neq 0$) and $b = \frac{c^2(c^2 + c + 1)}{c^3 + c + 1}$ and $c^5 + c^4 + c^3 + c + 1 = 0$. Used in E_6, E_9 renders c = 1, which is not allowed. If $p_1 = p_5$, then $a = \frac{c^3}{c^3 + c + 1}, b = \frac{c^3 + c + 1}{c^2}$ and $c^5 + c^4 + c^2 + c + 1 = 0$. Further, using these values in $\{E_6, E_8, E_9\}$, gives

$$\left\{ \frac{c^2(c+1)\left(c^2+c+1\right)}{c^3+c+1}, \frac{(c+1)\left(c^4+c^3+c^2+c+1\right)}{c^3+c+1}, \frac{(c+1)\left(c^3+c^2+1\right)}{c^2} \right\},\,$$

but these values cannot be zero under our conditions. If $p_1 = p_6$, then $a = c^2, b = \frac{1}{c}$, but these values, in E_7, E_8, E_9 will not work. If $p_1 = p_7$, then a = c, b = 1, which also imply that $E_8 = c + 1, E_9 = c(c + 1)$, and consequently, nonzero. If $p_1 = p_8$, then $a = \frac{c^2 + c + 1}{c}, b = \frac{c^2}{c^2 + c + 1}$, which is not going to vanish E_9 .

We now assume that $p_2=p_3$. Then, by adding $(a+1)E_2$ and aE_3 , we get $a+a^2+ac+a^2c+c^2+b(ac+a^2c+a^3c)=0$. If $1+a+a^2=0$, then we need $a+a^2+ac+a^2c+c^2=0$, that is, $c^2+c+1=0$. Thus, either a=c or $a=\frac{1}{c}$. If a=c, and $c^2+c+1=0$ (thus, $n\equiv 0 \mod 2$), then $E_2=E_3=0$, gives $1+c^2+bc^3+b^2c^3=1+c+bc+b^2(c^2+c^3)=0$. If further $p_2=p_3=p_4$, then adding $E_2+E_4=0$ gives b=1, which when used in E_2 implies c=1. If $p_2=p_3=p_5$, then, adding $E_2=0$, $E_5=0$ renders c=0, which is impossible. If $p_2=p_3=p_6$, then adding E_2 , E_6 gives $b=\frac{1}{c^2}$, which, when used in either E_2 , E_6 gives c=1. If c=1 if c=1

assumption $a = \frac{1}{c}$ and $c^2 + c + 1 = 0$. We now assume that $a^2 + a + 1 \neq 0$, and so, $b = \frac{a+a^2+ac+a^2c+c^2}{ac(1+a+a^2)}$. If $p_2 = p_3 = p_4$, then a = 1, which when used back into E_1 gives c = 0, 1. There is nothing tricky about the rest of the cases containing p_2 , so we move on. Similarly, for $p_3 = p_4$, $p_3 = p_5$.

cases containing p_2 , so we move on. Similarly, for $p_3 = p_4$, $p_3 = p_5$. We assume now $p_3 = p_6$. Adding $aE_3 + (a+1)E_6$ gives $b = \frac{a+a^2+ac+a^2c+c^2}{a(1+a^2+ac)}$. Continuing to add the other possible cases, p_7 , etc., and based upon observations, we can take a = c+1, $b = \frac{c^2+c+1}{c+1}$, where $1+c+c^2+c^3+c^4=0$ (which can always happen for some c when $n \equiv 0 \pmod{4}$), which will give us five solutions with $p_3 = p_6 = p_7 = p_{10}$, and so ${}_c\mathcal{B}_G(a,b) = 5$ for $n \equiv 0 \pmod{4}$. In a similar fashion, taking a = b = c+1, and c such that $c^3+c^2+1=0$ (thus, $n \equiv 0 \pmod{3}$), we see that $p_3 = p_7 = p_8 = p_{10}$, so ${}_c\mathcal{B}_G(a,b) = 5$ for $n \equiv 0 \pmod{3}$. With the same approach, with $a = \frac{c+1}{c}$, $b = \frac{c^2+c+1}{c(c+1)}$, and $c^4+c^3+c^2+c+1=0$ (so, $n \equiv 0 \pmod{4}$), then $p_4 = p_5 = p_9 = p_{10}$, so ${}_c\mathcal{B}_G(a,b) = 5$ in this case. The last claims of Theorem 4 on the bound being attained are shown.

B Some computational data

Below, we will compute the maximum c-boomerang uniformity of the (0,1)swapped binary inverse function in the fields \mathbb{F}_{2^2} , \mathbb{F}_{2^3} , \mathbb{F}_{2^4} , \mathbb{F}_{2^5} , determined
by the primitive polynomials X^2+X+1 , X^3+X+1 , X^4+X+1 , respectively, X^5+X^2+1 , each of some primitive root α .

The values of (c, a, b, x, y) achieving the (maximum) c-boomerang uniformity of 1 for n = 2 are:

```
(\alpha, \alpha, 0, \alpha, 0), (\alpha, \alpha, \alpha, \alpha, \alpha), (\alpha, \alpha, \alpha + 1, \alpha, \alpha + 1), (\alpha, \alpha, 1, \alpha, 1), (\alpha, \alpha + 1, 0, 0, \alpha + 1), \\ (\alpha, \alpha + 1, \alpha, 0, 1), (\alpha, \alpha + 1, \alpha + 1, 0, 0), (\alpha, \alpha + 1, 1, 0, \alpha), (\alpha, 1, 0, \alpha + 1, \alpha), \\ (\alpha, 1, \alpha, \alpha + 1, 0), (\alpha, 1, \alpha + 1, \alpha + 1, 1), (\alpha, 1, 1, \alpha + 1, \alpha + 1), (\alpha + 1, \alpha, 0, 0, \alpha), \\ (\alpha + 1, \alpha, \alpha, 0, 0), (\alpha + 1, \alpha, \alpha + 1, 0, 1), (\alpha + 1, \alpha, 1, 0, \alpha + 1), (\alpha + 1, \alpha + 1, 0, \alpha + 1, 0), \\ (\alpha + 1, \alpha + 1, \alpha, \alpha + 1, \alpha), (\alpha + 1, \alpha + 1, \alpha + 1, \alpha + 1), (\alpha + 1, \alpha + 1, 1, \alpha + 1, 1), \\ (\alpha + 1, 1, 0, \alpha, \alpha + 1), (\alpha + 1, 1, \alpha, \alpha, 1), (\alpha + 1, 1, \alpha + 1, \alpha, 0), (\alpha + 1, 1, 1, \alpha, \alpha).
```

The values of (c, a, b, x, y) achieving the (maximum) c-boomerang uniformity of 4 for n=3 are:

```
for (c, a, b) = (\alpha, \alpha^2 + 1, 0), the solutions (x, y) are : (0, \alpha^2 + 1), (\alpha, 0), (\alpha^2 + \alpha, \alpha + 1), (\alpha^2 + 1, \alpha^2 + \alpha + 1), for (c, a, b) = (\alpha^2, \alpha^2 + \alpha + 1, 0), the solutions (x, y) are : (0, \alpha^2 + \alpha + 1), (\alpha, \alpha^2 + 1), (\alpha^2, 0), (\alpha^2 + \alpha + 1, \alpha + 1); for (c, a, b) = (\alpha + 1, \alpha + 1, 0), the solutions (x, y) are :
```

```
(0, \alpha^2 + \alpha), (\alpha + 1, 0), (\alpha^2 + \alpha + 1, \alpha^2), (\alpha^2 + 1, \alpha + 1)
for (c, a, b) = (\alpha^2 + \alpha, \alpha + 1, 0), the solutions (x, y) are :
(0, \alpha + 1), (\alpha^2, \alpha^2 + \alpha + 1), (\alpha + 1, \alpha^2 + 1), (\alpha^2 + \alpha, 0);
for (c, a, b) = (\alpha^2 + \alpha + 1, \alpha^2 + \alpha + 1, 0), the solutions (x, y) are :
(0, \alpha^2), (\alpha + 1, \alpha^2 + \alpha + 1), (\alpha^2 + \alpha + 1, 0), (\alpha^2 + 1, \alpha);
for (c, a, b) = (\alpha^2 + 1, \alpha^2 + 1, 0), the solutions (x, y) are :
(0, \alpha), (\alpha + 1, \alpha^2 + \alpha), (\alpha^2 + \alpha + 1, \alpha^2 + 1), (\alpha^2 + 1, 0).
```

The values of (c, a, b, x, y) achieving the (maximum) c-boomerang uniformity of 5 for n=4 are:

```
for (c, a, b) = (\alpha^3, \alpha^3 + \alpha^2 + \alpha, \alpha^3 + \alpha^2), the solutions (x, y) are:
(0, \alpha^3 + \alpha^2 + 1), (\alpha, \alpha^3 + \alpha^2 + \alpha + 1), (\alpha^2, \alpha + 1), (\alpha^3 + \alpha^2 + 1, \alpha^3 + \alpha^2), (1, \alpha^3 + \alpha);
for (c, a, b) = (\alpha^3, \alpha^3 + 1, \alpha^3 + \alpha), the solutions (x, y) are:
(\alpha, \alpha + 1), (\alpha^3, \alpha^2 + 1), (\alpha^3 + \alpha^2, \alpha^3 + \alpha + 1), (\alpha^3 + \alpha, 1), (\alpha^3 + 1, \alpha):
for (c, a, b) = (\alpha^3 + \alpha^2, \alpha^3 + \alpha + 1, \alpha^3 + \alpha^2 + \alpha + 1), the solutions (x, y) are:
(0, \alpha^3 + \alpha^2 + \alpha), (\alpha^2, \alpha^3 + \alpha), (\alpha + 1, \alpha^2 + 1), (\alpha^3 + \alpha^2 + \alpha, \alpha^3 + \alpha^2 + \alpha + 1), (1, \alpha^3);
for (c, a, b) = (\alpha^3 + \alpha^2, \alpha^3 + \alpha^2 + 1, \alpha^3), the solutions (x, y) are:
(\alpha^2, \alpha^2 + 1), (\alpha^3, 1), (\alpha^3 + \alpha^2, \alpha), (\alpha^3 + \alpha^2 + \alpha + 1, \alpha^3 + 1), (\alpha^3 + \alpha^2 + 1, \alpha^2);
for (c, a, b) = (\alpha^3 + \alpha, \alpha^3 + \alpha + 1, \alpha^3 + \alpha^2 + \alpha + 1), the solutions (x, y) are:
(\alpha^3, \alpha^3 + \alpha^2 + \alpha), (\alpha^3 + \alpha + 1, \alpha^2 + 1), (\alpha^2 + 1, \alpha^2), (\alpha^3 + \alpha, \alpha + 1), (\alpha^3 + \alpha^2 + \alpha + 1, 1);
for (c, a, b) = (\alpha^3 + \alpha, \alpha^3 + \alpha^2 + 1, \alpha^3), the solutions (x, y) are :
(0, \alpha^3 + 1), (\alpha, \alpha^2), (\alpha^2 + 1, \alpha^3 + \alpha^2), (\alpha^3 + 1, \alpha^3), (1, \alpha^3 + \alpha^2 + \alpha + 1);
for (c, a, b) = (\alpha^3 + \alpha^2 + \alpha + 1, \alpha^3 + \alpha^2 + \alpha, \alpha^3 + \alpha^2), the solutions (x, y) are:
(\alpha + 1, \alpha), (\alpha^3 + \alpha^2, 1), (\alpha^3 + \alpha, \alpha^3 + \alpha^2 + 1), (\alpha^3 + \alpha^2 + \alpha, \alpha + 1), (\alpha^3 + \alpha^2 + \alpha + 1, \alpha^2);
for (c, a, b) = (\alpha^3 + \alpha^2 + \alpha + 1, \alpha^3 + 1, \alpha^3 + \alpha), the solutions (x, y) are:
(0, \alpha^3 + \alpha + 1), (\alpha^3 + \alpha, \alpha + 1, \alpha^3), (\alpha^3 + \alpha + 1, \alpha^3 + \alpha), (\alpha^2 + 1, \alpha), (1, \alpha^3 + \alpha^2).
```

The values of (c, a, b, x, y) achieving the (maximum) c-boomerang uniformity of 5 for n = 5 are:

```
for (c,a,b)=(\alpha^3,\alpha^4+\alpha^3+\alpha+1,\alpha+1), the solutions (x,y) are : (\alpha^2,0),(\alpha^3,\alpha^4+\alpha),(\alpha^3+\alpha,\alpha^4+\alpha^3+\alpha),(\alpha^4+1,\alpha^4+\alpha+1),(\alpha^4+\alpha^3+\alpha+1,1); for (c,a,b)=(\alpha^3+\alpha,\alpha,\alpha^2+1), the solutions (x,y) are : (\alpha,1),(\alpha^4,0),(\alpha^3+\alpha,\alpha^3+1),(\alpha^3+\alpha^2+\alpha,\alpha+1),(\alpha^3+\alpha^2,\alpha^3); for (c,a,b)=(\alpha^4+\alpha^2,\alpha^4,\alpha^3+\alpha^2), the solutions (x,y) are : (0,\alpha^4+1),(\alpha^3+\alpha^2+\alpha,\alpha^4+\alpha^3+\alpha^2+\alpha+1),(\alpha^4+\alpha+1,\alpha^4+\alpha^3+\alpha^2+\alpha),(\alpha+1,1),(\alpha^3+\alpha+1,\alpha^4); for (c,a,b)=(\alpha^3+\alpha^2+\alpha,\alpha^2,\alpha^4+1), the solutions (x,y) are :
```

```
(\alpha^2,1), (\alpha^3+\alpha^2+1,0), (\alpha^4+\alpha^3+\alpha,\alpha^3+\alpha), (\alpha^3+\alpha^2+\alpha,\alpha^3+\alpha+1), (\alpha^4+\alpha^3+\alpha^2+\alpha,\alpha^2+1); for (c,a,b)=(\alpha^4+\alpha^3+\alpha^2+1,\alpha^3+\alpha^2+1,\alpha^4+\alpha^3+\alpha), the solutions (x,y) are : (0,\alpha^3+\alpha^2), (\alpha^3,\alpha^4+\alpha+1), (\alpha^2+1,1), (\alpha^3+\alpha^2+\alpha+1,\alpha^3+\alpha^2+1), (\alpha^4+\alpha^3+\alpha^2+\alpha,\alpha^4+\alpha); for (c,a,b)=(\alpha^4+\alpha+1,\alpha^3+\alpha^2+1,\alpha^4+\alpha^3+\alpha), the solutions (x,y) are : (\alpha,0), (\alpha^3,\alpha^3+\alpha^2), (\alpha^2+1,\alpha^4+\alpha^3+\alpha^2+\alpha), (\alpha^3+\alpha^2+1,1), (\alpha^4+\alpha+1,\alpha^4+\alpha^3+\alpha^2+\alpha+1); for (c,a,b)=(\alpha^2+\alpha,\alpha^2,\alpha^4+1), the solutions (x,y) are : (0,\alpha^2+1), (\alpha^3+\alpha,\alpha^3+\alpha^2+\alpha+1), (\alpha^4+\alpha^3+\alpha,1), (\alpha^4+\alpha^3+\alpha^2+\alpha,\alpha^3+\alpha^2+\alpha), (\alpha^3+1,\alpha^2); for (c,a,b)=(\alpha^4+\alpha^3+\alpha^2+\alpha,\alpha^4,\alpha^3+\alpha^2), the solutions (x,y) are : (\alpha^4,1), (\alpha^4+\alpha^3+\alpha+1,0), (\alpha^4+\alpha+1,\alpha^4+1), (\alpha+1,\alpha^3+\alpha^2+\alpha), (\alpha^4+\alpha^3+\alpha^2+\alpha,\alpha^3+\alpha^2+\alpha+1); for (c,a,b)=(\alpha^4+\alpha^3+1,\alpha,\alpha^2+1), the solutions (x,y) are : (0,\alpha+1), (\alpha^3,\alpha^3+\alpha+1), (\alpha^3+\alpha^2+\alpha,\alpha^3+\alpha), (\alpha^3+\alpha^2,1), (\alpha^4+\alpha,\alpha); for (c,a,b)=(\alpha^4+\alpha^3+\alpha+1,\alpha,\alpha^2+\alpha), (\alpha^3+\alpha^2+\alpha), (\alpha^3+\alpha^2+\alpha), (\alpha^4+\alpha,\alpha); for (c,a,b)=(\alpha^4+\alpha^3+\alpha+1,\alpha,\alpha^2+\alpha), (\alpha^3+\alpha^2+\alpha), (\alpha^4+\alpha,\alpha); for (c,a,b)=(\alpha^4+\alpha^3+\alpha+1,\alpha,\alpha^2+\alpha), (\alpha^3+\alpha^2+\alpha), (\alpha^4+\alpha,\alpha); for (c,a,b)=(\alpha^4+\alpha^3+\alpha,\alpha,\alpha^3+\alpha+\alpha), (\alpha^3+\alpha^2+\alpha,\alpha^3+\alpha+\alpha), (\alpha^4+\alpha,\alpha); for (c,a,b)=(\alpha^4+\alpha^2+\alpha,\alpha^4+\alpha^3+\alpha+1,\alpha+1), (\alpha^4+\alpha^3+\alpha^2+\alpha+1), (\alpha^4+\alpha,\alpha); for (c,a,b)=(\alpha^4+\alpha^2+\alpha,\alpha^4+\alpha^3+\alpha+1,\alpha+1), (\alpha^4+\alpha^3+\alpha^2+\alpha+1), (\alpha^4+\alpha,\alpha); for (c,a,b)=(\alpha^4+\alpha^3+\alpha,\alpha^3+\alpha^2+\alpha,\alpha^3+\alpha+1), (\alpha^3+\alpha^2+\alpha,\alpha^3+\alpha^2+\alpha+1), (\alpha^4+\alpha,\alpha); for (c,a,b)=(\alpha^4+\alpha^2+\alpha,\alpha^4+\alpha^3+\alpha+1,\alpha+1), (\alpha^4+\alpha^3+\alpha^2+\alpha+1), (\alpha^4+\alpha,\alpha+1), (\alpha^4+
```