On the chromatic numbers of small-dimensional Euclidean spaces

Danila Cherkashin^{*}, Anatoly Kulikov[†], Andrei Raigorodskii[‡]

Abstract This paper is devoted to the study of the graph sequence $G_n = (V_n, E_n)$, where V_n is the set of all vectors $v \in \mathbb{R}^n$ with coordinates in $\{-1, 0, 1\}$ such that $|v| = \sqrt{3}$ and E_n consists of all pairs of vertices with scalar product 1. We find the exact value of the independence number of G_n . As a corollary we get new lower bounds on $\chi(\mathbb{R}^n)$ and $\chi(\mathbb{Q}^n)$ for small values of n. **1. Introduction** Let \mathbb{R}^n be the standard Euclidean space, where the distance between any two points x, y is denoted by |x - y|. Let V be an arbitrary point set in \mathbb{R}^n . Let a > 0 be a real number. By a distance graph with set of vertices V, we mean the graph G = (V, E) whose set of edges E contains all pairs of points from Vthat are at the distance a apart: $E = \{\{x, y\} : |x - y| = a\}.$ Distance graphs are among the most studied objects of combinatorial geometry. First of all, they are at the ground of the classical Hadwiger-Nelson problem, which was proposed around 1950 (see [9], [22]) and consists in determining the chromatic number of the space: $\chi(\mathbb{R}^n) = \min\{\chi : \mathbb{R}^n = V_1 \sqcup \ldots \sqcup V_{\chi}, \forall i \forall \mathbf{x}, \mathbf{y} \in V_i \ |\mathbf{x} - \mathbf{y}| \neq 1\},$ i.e., the minimum number of colors needed to color all the points in \mathbb{R}^n so that any two points at the distance 1 receive different colors. In other words, it is the chromatic number of the unit distance graph whose vertex set coincides with \mathbb{R}^n . Due to the extreme popularity of the subject, colorings of unit distance graphs are very deeply explored. Let us just refer the reader to several books and survey articles [1, 2, 5, 11, 18, 19, 21, 23]. In particular, the best known lower bounds for the chromatic numbers in dimensions ≤ 12 are given below:

the best known lower bounds for the chromatic numbers in dimensions ≤ 12 are given below:

$$\chi(\mathbb{R}^2) \ge 4 \ [18], \ \chi(\mathbb{R}^3) \ge 6 \ [17], \ \chi(\mathbb{R}^4) \ge 9 \ [8], \ \chi(\mathbb{R}^5) \ge 9 \ [4], \ \chi(\mathbb{R}^6) \ge 11 \ [6], \ \chi(\mathbb{R}^7) \ge 15 \ [18], \\ \chi(\mathbb{R}^8) \ge 16 \ [15], \ \chi(\mathbb{R}^9) \ge 21 \ [14], \ \chi(\mathbb{R}^{10}) \ge 23 \ [14], \ \chi(\mathbb{R}^{11}) \ge 25 \ [12], \ \chi(\mathbb{R}^{12}) \ge 27 \ [13].$$

Recently further improvements were announced:

 $\chi(\mathbb{R}^8) \ge 19 \ [10], \ \chi(\mathbb{R}^{10}) \ge 26 \ [7], \ [10], \ \chi(\mathbb{R}^{11}) \ge 32 \ [10], \ \chi(\mathbb{R}^{12}) \ge 36 \ [7].$

^{*}Saint Petersburg State University, Faculty of Mathematics and Mechanics; Moscow Institute of Physics and Technology, Laboratory of Advanced Combinatorics and Network Applications.

[†]Saint Petersburg State University, Faculty of Physics; The University of Queensland, School of Mathematics and Physics.

[‡]Moscow Institute of Physics and Technology, Faculty of Innovations and High Technology, Department of Discrete Mathematics and Laboratory of Advanced Combinatorics and Network Applications; Moscow State University, Mechanics and Mathematics Faculty, Department of Mathematical Statistics and Random Processes; Buryat State University, Institute of Mathematics and Informatics.

These improvements are essentially based on computer calculations.

In growing dimensions, the following bounds are the best known:

[20]
$$(1.239...+o(1))^n \leq \chi(\mathbb{R}^n) \leq (3+o(1))^n$$
 [15].

In this paper, we consider a special sequence of graphs defined in the following way.

Let V_n be the set of all vectors v from \mathbb{R}^n with coordinates in $\{-1, 0, 1\}$ and $|v| = \sqrt{3}$. The set V_n can be considered as the set of vertices of a graph $G_n = (V_n, E_n)$, where an edge connects two vertices if and only if the corresponding vectors have scalar product 1. Note that G_1 and G_2 are empty and G_3 is just a cube.

Recall that an *independent set* in a graph is any set of its vertices which are pairwise non-adjacent and the *independence number* of G denoted by $\alpha(G)$ is the size of a maximum independent set in the graph G.

Theorem 1. For $n \ge 1$, let c(n) denote the following constant:

$$c(n) = \begin{cases} 0 & if \ n \equiv 0 \\ 1 & if \ n \equiv 1 \\ 2 & if \ n \equiv 2 \ or \ 3 \end{cases} \pmod{4}.$$

Then, the independence number of G_n is given by the formula

$$\alpha(G_n) = \max\{6n - 28, 4n - 4c(n)\}.$$

Actually, the result of Theorem 1 is a far-reaching generalization of a much simpler lemma proved by Zs. Nagy (see [16]) in 1972 and used not only in combinatorial geometry, but also in Ramsey theory. In this lemma, $G'_n = (V'_n, E'_n)$, where V'_n is the set of all vectors v, $|v| = \sqrt{3}$, with coordinates in $\{0, 1\}$ and again an edge connects two vertices if and only if the corresponding vectors have scalar product 1. Lemma states that in this case $\alpha(G'_n) = n - c(n)$.

The proof of Theorem 1 is given in the following parts: some examples showing the lower bound in Theorem 1 and some preliminaries are given in Section 2; the upper bound is proved in Section 3 for n > 13 and in Section 4 for other values of n. Note that, roughly speaking, the quantity 13 is a threshold where the bound 6n - 28 starts dominating the bound 4n.

As a corollary of Theorem 1 we get the following bounds for the chromatic numbers of Euclidean spaces.

Theorem 2. Let c(n) be the constant defined in Theorem 1. Then, for all $n \ge 3$, we have

$$\chi(\mathbb{R}^n) \geqslant \chi(\mathbb{Q}^n) \geqslant \chi(G_n) \geqslant \frac{|V_n|}{\alpha(G_n)} = \frac{8C_n^3}{\max\{6n - 28, 4n - c(n)\}}.$$

Asymptotically, the bound in this theorem is $\frac{2}{9}n^2(1+o(1))$, which is a weak result. On the other hand, for small values of n, the theorem gives the best known bounds, namely:

$$\chi(\mathbb{R}^9) \ge \chi(\mathbb{Q}^9) \ge 21,$$

$$\chi(\mathbb{R}^{10}) \ge \chi(\mathbb{Q}^{10}) \ge 30,$$

$$\chi(\mathbb{R}^{11}) \ge \chi(\mathbb{Q}^{11}) \ge 35,$$

$$\chi(\mathbb{R}^{12}) \ge \chi(\mathbb{Q}^{12}) \ge 37.$$

Actually, we will show in Section 5 the following stronger result for n = 9. **Proposition 1.** The inequalities hold

$$\chi(\mathbb{R}^9) \geqslant \chi(\mathbb{Q}^9) \geqslant 22.$$

2. Lower bounds in Theorem 1 and some preliminaries

2.1. Auxiliary definitions

Consider the graph G_n . Any of its vertices has three non-zero coordinates and n-3 coordinates equal to 0. We call *base* the set of non-zero coordinates of a vertex. To make our exposition more concise, we will use the word "place" instead of the word "coordinate" or instead the expression "coordinate position". For example, it will be convenient to say (a bit informally) "vertex v intersects place x", if the vector vfrom \mathbb{R}^n corresponding to this vertex has nonzero value of the coordinate v_x . For the same reasons, we introduce the notion of a *signplace*: it is a coordinate with a fixed sign (plus or minus). In particular, from now on, we can say (again, a bit informally) "vertex v intersects signplace x^+ (x^-)", if it has the value of the coordinate v_x equal to +1 (-1). Finally, we define the *degree* of a place (signplace) in a set W of vertices of G_n as the number of vertices from W intersecting this place (signplace).

2.2. Constructions of independent sets in G_n

It suffices to show that $\alpha(G_n) \ge 4n - 4c(n)$ and that $\alpha(G_n) \ge 6n - 28$.

The first construction is as follows. Consider the first 4 places. Take all the 4 bases that can be taken on these places. For each base, consider 4 variants:

$$1, 1, 1; 1, -1, -1; -1, 1, -1; -1, -1, 1.$$

Clearly any two vectors with these bases have scalar product different from 1. We call this construction (and its natural analogs) quad.

Take [n/4] consecutive quads. If the remainder still consists of 3 places, then add 4 more bases. Eventually, we get exactly 4n - 4c(n) vectors that form an independent set in G_n .

Now, let us make the second construction. Take the following vectors:

$$1, -1, 0, 1, 0, \dots, 0, 0, 0, 0; \quad 1, -1, 0, 0, 1, 0, \dots, 0, 0, 0; \quad \dots; \quad 1, -1, 0, 0, 0, \dots, 0, 1, 0, 0, 0; \\0, 1, -1, 1, 0, \dots, 0, 0, 0, 0; \quad 0, 1, -1, 0, 1, 0, \dots, 0, 0, 0, 0; \quad \dots; \quad 0, 1, -1, 0, 0, \dots, 0, 1, 0, 0, 0; \\-1, 0, 1, 1, 0, \dots, 0, 0, 0, 0; \quad -1, 0, 1, 0, 1, 0, \dots, 0, 0, 0, 0; \quad \dots; \quad -1, 0, 1, 0, 0, \dots, 0, 1, 0, 0, 0.$$

In each line, we have a set of vectors, which is a particular case of what we will call *snake* in Section 3 and later. In every snake, we have n-6 vertices. Thus, the total amount of vertices here is 3n-18. Obviously, the union of these snakes is an independent set in G_n . Moreover, we can add to it 4 more vectors, which have a common base — the three first places: say,

$$1, 1, 1, 0, \dots, 0; \quad 1, -1, -1, 0, \dots, 0; \quad -1, 1, -1, 0, \dots, 0; \quad -1, -1, 1, 0, \dots, 0.$$

The whole construction is a particular case of a *cobra* discussed later in more details. Here the cobra contains 3n - 14 vertices.

Of course, we can take one more cobra, whose "head" is on the three last places and whose "tail" consists of minus ones instead of ones. Eventually, we get exactly 6n - 28 vertices forming an independent set in G_n .

The lower bound is proven.

It is worth noting that in Section 3 we will make a rather subtle analysis of possible independent sets in G_n . One would be able to derive from this analysis a complete description of examples giving the lower bound in Theorem 1. However, we will not present such description explicitly in this paper.

It is also worth noting that in the above example having 6n - 28 vertices and avoiding the scalar product 1, the scalar product -3 is also absent. Moreover, one can exclude 6 vertices from that example so that the scalar product -2 disappears as well.

2.3. Basic lemma

Let A be an arbitrary independent set of the maximum size in G_n . We already know that $|A| \ge \max\{6n - 28, 4n - 4c(n)\}$. Assume that we exclude some signplaces and all the vertices from the graph G_n intersecting them. Then we get a new graph G' with a possibly smaller independent set A'. Denote by a(A') the maximum degree of a signplace in the set A'. Denote by m(A') the number of signplaces in A'.

The following lemma is an important ingredient in the proof of the upper bound.

Lemma 1. Assume that we exclude k signplaces. Assume that the number of vertices excluded from A does not exceed 2k. Then we have either $a(A') \ge 5$ or m(A') < 14.

Proof of the lemma. By pigeon-hole principle $a(A') \ge 3(|A| - 2k)/(2n - k)$. If $|A| \ge 4n$, then $a(A') \ge 6$ and we are done. The inequality $|A| \ge 4n$ is true for n = 8, 12 and $n \ge 14$. Thus, it remains to consider only n = 7, 9, 10, 11, 13. If n = 7, then $|A| \ge \max\{14, 20\} = 20$. If k = 1, then $3(|A| - 2k)/(2n - k) \ge 54/13$, i.e., $a(A') \ge 5$. If $k \ge 2$, then it may happen that $3(|A| - 2k)/(2n - k) \le 4$. But in this case, $m(A') = 2n - k \le 14 - 2 < 14$. The same argument works for the 4 other values of n. The proof is complete.

3. Proof of Theorem 1 in the case $n \ge 14$

3.1. Starting the proof

Let A be an arbitrary independent set of the maximum size in G_n . Assume that we have already excluded several signplaces with the corresponding vertices (see Section 2.3). By Lemma 1 either $a(A') \ge 5$ or $m(A') \le 13$. The second case will be considered in Section 3.4. So we assume that $a(A') \ge 5$.

Consider a signplace with the maximum degree. Call it x_1^+ (each time when we choose a sign we can choose plus without loss of generality) and consider the set of vertices intersecting it (denote it by $N_{x_1^+}$). Note that no base can contain more than two vertices from $N_{x_1^+}$. Thus, we have at least three different bases. Also it is clear that any two bases containing vertices from $N_{x_1^+}$ intersect in exactly two signplaces. There are two different possibilities.

- 1) Among the bases, we have $\{x_1, x_2, x_3\}, \{x_1, x_2, x_4\}, \{x_1, x_3, x_4\}$. This case will be referred to as "quad" (cf. Section 2.2).
- 2) All the bases contain both x_1 and x_2 . This case will be referred to as "snake" (cf. Section 2.2).

The formal definition of a quad will be given in the next section, where we will analize Case 1. The same is for a snake in Section 3.3. In Section 3.4, we will complete the proof.

3.2. The first case — "quad"

We know that $a(A') \ge 5$. At the same time, $a(A') \le 6$, since otherwise the vertices from $N_{x_1^+}$ use at least 4 bases and therefore there is a base among $\{x_1, x_2, x_3\}, \{x_1, x_2, x_4\}, \{x_1, x_3, x_4\}$ such that it intersects the fourth base only on x_1^+ , which is impossible. Put a = a(A').

Thus, we have exactly three bases containing the vertices from $N_{x_1^+}$. Two of them (without loss of generality $\{x_1, x_2, x_3\}, \{x_1, x_2, x_4\}$) contain exactly 2 vertices from $N_{x_1^+}$ each, and the third one contains at least 1 vertex. Since $\{x_1^+, x_2, x_3\}$ contains two vertices, it intersects all the four signplaces in x_2, x_3 ; the same holds for $\{x_1^+, x_2, x_4\}$, which means that all the six signplaces of x_2, x_3, x_4 are necessarily intersected.

Consider the set U of all vertices intersecting $\{x_1, x_2, x_3, x_4\}$. There could be the following possibilities.

- Some vertices from U intersect x_1 . There are at most 2a such vertices.
- Some vertices from U lie on the base $\{x_2, x_3, x_4\}$. There are at most 4 such vertices.
- Some vertices from U intersect $\{x_2, x_3, x_4\}$ in one place and are not counted above. Actually, there are no such vertices because for every signplace in $\{x_2, x_3, x_4\}$ a vertex with a base in $\{x_1, x_2, x_3, x_4\}$ exists (do not forget that $\{x_1, x_2, x_3\}, \{x_1, x_2, x_4\}$ contain exactly 2 vertices each, and the third base contains at least 1 vertex).
- Some vertices from U intersect $\{x_2, x_3, x_4\}$ in two places and are not counted above. Again, there are no such vertices. Indeed, assume that some vertex (call it v) intersects $\{x_2, x_3, x_4\}$ in $\{x_i, x_j\}$. Then $\{x_1, x_2, x_3\}$ or $\{x_1, x_2, x_4\}$ intersects $\{x_i, x_j\}$ in exactly one place. This is impossible, since we know that two vertices from $N_{x_1^+}$ lie on $\{x_1, x_2, x_3\}$ and two vertices from $N_{x_1^+}$ lie on $\{x_1, x_2, x_4\}$.

Summarizing, we have at most $2a + 4 \leq 16$ vertices intersecting 8 signplaces. We call any of the corresponding constructions *quad*.

Now we may assume that A was transformed into A' in the following way (more details will be given in Section 3.4).

- First, all the signplaces of degree less than 3 have been deleted one by one. Note that by Lemma 1 during this process either $a \ge 5$ or $m \le 13$.
- Second, all the quads have been deleted one by one. Note that again by Lemma 1 during this process either $a \ge 5$ or $m \le 13$ (everytime the number of excluded signplaces is 8 and the number of excluded vertices is at most 16).
- Third, once again, all the signplaces of degree less than 3 have been deleted one by one. Obviously, there are no new quads and still by Lemma 1 $a \ge 5$ or $m \le 13$.

As before, we assume that $a \ge 5$ (since the case $m \le 13$ is considered in Section 3.4), and so we are prepared to the next case, in which we have $a(A') \ge 5$, there are no quads, and every signplace has degree at least 3.

3.3. The second case — "snake"

We start with a formal definition of a snake.

Definition 1. Snake is a set of vertices intersecting a signplace and a place and containing at least 5 vertices. Head of a snake is a couple of places, which intersect every vertex, and tail of a snake is the set of the remaining signplaces in each vertex. Size of a snake is the number of its vertices.

Clearly in the current case we have a snake of size $a \ge 5$ in A'. Let it be based on $\{x_1^+, x_2\}$ (with the head being $\{x_1, x_2\}$). Note that the size of its tail is equal to a, since vertices cannot intersect on tail.

Our aim is to prove that we can exclude some t signplaces with at most 3t - 14 vertices. Moreover, we will show that there is a special construction ("cobra", cf. Section 2.2), which has exactly 3t - 14 vertices on t signplaces and which is the only such construction up to the graph symmetries.

We have an alternative.

- 1) We can exclude 4 + a signplaces ($\{x_1, x_2\}$ with all possible signs and a signplaces of the tail) and 3a 2 vertices.
- 2) We have at least 3a 1 vertices intersecting the signplaces mentioned in the previous point.

In the first case, our aim is realized, since we can put t = 4 + a and get 3t - 14 = 3a - 2. In the second case, the analysis will be much longer.

Let us consider the second case of the alternative. Each vertex intersecting the tail of the snake that we analyze should intersect the head as well, and each of the a initial vertices intersects the head on two signplaces. Hence the sum of the degrees of the head signplaces is at least 4a - 1. But there is no signplace with degree exceeding a, so the degrees of the signplaces in the head are either

$$a, a, a, a$$
 or $a, a, a, a - 1$.

Anyway we have a place with two signplaces of degree exactly a. Without loss of generality, this place is x_1 . Since all quads are already excluded, we have two snakes with signplaces on x_1 : one signplace is x_1^+ and the second one is x_1^- . Consider their heads. They could both lie on $\{x_1, x_2\}$, or they could lie on $\{x_1, x_2\}$ and $\{x_1, x_3\}$ respectively.

In the first case, all the four signplaces of the head have degree a solely due to 2a vertices from the snakes. In addition, there are vertices intersecting the tail (since the degree of each signplace is at least 3 and two snakes could provide only two vertices on a signplace). Each vertex intersecting the tail should intersect the head as well, so the degree of some signplace in the head exceeds a, which contradicts the assumption that a is the maximum value of the degree.

We are left with the second case: there are two snakes of size a with heads on $\{x_1, x_2\}$ and $\{x_1, x_3\}$.

Let Q be the set of vertices lying fully on base $\{x_1, x_2, x_3\}$. Denote by B the set of signplaces in the intersection of the tails. Let C_1 and C_2 be the sets of the remaining signplaces in the corresponding tails. Let q, b, c_1, c_2 be the sizes of the corresponding sets. We have already described all the vertices intersecting x_1 , since the maximum degree is equal to a. Consider the sum of the degrees of the signplaces on x_2 and x_3 . Since the degree of each signplace is at least 3, we have a new vertex for each signplace from the intersection of the tails. Each vertex of this type should intersect both heads, and it cannot contain x_1 . Therefore, it contains both x_2 and x_3 and adds 2 to our sum. We have at least two vertices intersecting each signplace of the symmetric difference of the tails. Each vertex of this type should intersect the head of a corresponding snake and could intersect two signplaces of its tail. In total, these vertices add at least

$$2(c_1 + c_2)/2 = c_1 + c_2$$

to the sum. Each of the 2*a* initial vertices intersects $\{x_2, x_3\}$. Each vertex from *Q* adds yet another 1 to the sum, since it intersects $\{x_2, x_3\}$ on two places. Again in total, the sum of the degrees of the four signplaces on $\{x_2, x_3\}$ is at least

$$2b + c_1 + c_2 + 2a + q.$$

On the other hand, since the degree of each signplace is at most a, this sum does not exceed 4a. So we have

$$2b + c_1 + c_2 + q \leqslant 2a.$$

Each vertex from Q is in one snake. Consequently, $q = q_1 + q_2$ (q_i is the number of vertices lying in a corresponding snake),

$$b + q_1 + c_1 = a, \quad b + q_2 + c_2 = a,$$

and the inequality always turns to equality!

Thus, there is a set of

$$t := 6 + b + c_1 + c_2$$

signplaces intersected by

$$q + 3b + 3c_1 + 3c_2 \leq 3(b + c_1 + c_2) + 4 = 3t - 14$$

vertices.

The second case of the alternative is complete, and our aim is attained. However, we will also prove below an upper bound on t.

Suppose that the number of vertices is exactly 3t - 14. It means that all the intermediate inequalities turned to equalities. The last inequality turns to equality only when q = 4. One can see that any vertex intersecting a signplace from C_1 or C_2 should intersect 2 vertices of the tail, so it intersects $\{x_1, x_2, x_3\}$ only on 1 vertex, which contradicts q = 4. Hence $C_1 = C_2 = \emptyset$ and b = a - 2. For every signplace from $x^+ \in B$ there is a vertex intersecting x^+ and lying on base $\{x, x_2, x_3\}$. It turns out that there is a third snake on $\{x_2, x_3\}$. We call *cobra* the union of such 3 snakes. Finally, one can see that there is no place xsuch that x^+, x^- lie in B, otherwise there is an edge between two vertices on the base $\{x_1, x_2, x_3\}$.

As a result, t does not exceed n+3, and the tail of a corresponding snake can not contain two signplaces on the same place.

Summing up the above, if we have no quad, then there is a cobra, which consists of three snakes with a common tail and pairwise intersecting heads. It has 3t - 14 vertices on t signplaces, $8 \le t \le n + 3$.

3.4. Finishing the proof

In the previous sections we have shown that there are the following options.

- To exclude a signplace and at most 2 vertices intersecting it.
- To exclude 8 signplaces and at most 16 vertices intersecting it.
- To exclude t signplaces with at most 3t 14 vertices $(8 \le t \le n + 3)$.
- To get $m(A') \leq 13$.

Clearly the two first options yield at most 2m vertices on m signplaces. The same is true for the fourth option: we show it in Appendix using computer simulations.

According to this, only the following cases could occur.

- 1) There is no cobra. Then the number of vertices does not exceed $4n \leq 6n 28$.
- 2) There is one cobra and $t \leq n$. Then the number of vertices does not exceed

$$3t - 14 + 2(2n - t) \leq 5n - 14 \leq 6n - 28.$$

- 3) There is one cobra and t = n+1. We are left to prove that n-1 signplaces on n-3 places can contain at most 2n-3 vertices. Suppose the contrary, then, by pigeon-hole principle, there is a signplace of degree at least 3(2n-2)/(n-1) = 6. Using the same arguments as in Sections 3.1–3.3 we get a quad or a cobra, but both constructions contain 3 places with 2 signplaces, which contradicts our assumptions.
- 4) There is one cobra and t = n + 2. We are left to prove that n 2 signplaces on n 3 places can contain at most 2n 6 vertices. Then the number of vertices does not exceed

$$3(n+2) - 14 + 2n - 6 \leq 5n - 14 \leq 6n - 28$$

Again, suppose the contrary, so there is a singplace of degree at least 3(2n-5)/(n-2). For n < 4 the claim is obvious, and for $n \ge 4$ we have 3(2n-5)/(n-2) > 4. Using the same arguments as in Sections 3.1–3.3 we get a quad or a cobra, but both constructions contain 3 places with 2 signplaces. Thus, we get a contradiction.

5) There is one cobra and t = n+3. All other signplaces lie on distinct places, and we can apply Nagy's lemma (see Section 1 and [16]) to get an upper bound n-3 for the number of vertices. Then the number of vertices does not exceed

 $3(n+3) - 14 + n - 3 = 4n - 8 < 5n - 14 \le 6n - 28.$

6) There are two or more cobras. Then the bound 6n - 28 for the number of vertices is straightforward.

The proof of Theorem 1 for $n \ge 14$ is complete.

4. Proof of Theorem 1 in the case n < 14

We will deal with small values of n one by one. We have already proven the lower bounds in Section 2.2. So we have to prove only the upper bounds. Let us start from simple cases.

- In the case $n \leq 6$ the answer is easily found on a computer.
- The case n = 7 was considered by Cibulka in [6].
- In the case n = 9 we have to show that $\alpha(G_9) \leq 32$, and this will follow from the case n = 10 below, since $\alpha(G_n)$ is non-decreasing.

In the remaining cases the main lines of the proofs will be the same as in Sections 3.1–3.3, but we will make appropriate changes in the ends of the proofs.

4.1. The cases n = 8 and n = 12

We state that $\alpha(G_8) \leq 32$ and $\alpha(G_{12}) \leq 48$. Suppose the contrary. It means that $\alpha(G_n) > 4n$ for some $n \in \{8, 12\}$. Hence, by Section 3.4 there is a cobra on t signplaces with $t \ge 15$, since otherwise $3t - 14 \le 2t$ and thus in all the possibilities from Section 3.4 we have at most 2m vertices on m signplaces. But a cobra has at most n + 3 signplaces, so the only option is n = 12, t = 15. Let us exclude the cobra and note that all the 9 remaining signplaces lie on different places. Now we can apply Nagy's lemma (see Section 1 and [16]) to get in this case an upper bound 3t - 14 + 8 = 39 < 48. We get a contradiction and the claim is proven.

4.2. The case n = 10

We are going to prove that $\alpha(G_{10}) \leq 32$. Suppose we have an independent set A whose size is at least 32. Let us follow the proof of the case $n \geq 14$. If there is a quad, we exclude it and refer to the Appendix, where the results of some computer calculations are given, in particular, for m = 12, l = 6 (m is the number of signplaces, and l is the number of places). Thus, we get the needed bound by 16 + 16 = 32.

Then, we exclude some k signplaces whose degrees are at most 2. Suppose that the maximum degree in the remaining graph is at most 4. By Lemma 1 we get the bound $k \ge 8$. Then $|A| \le 2k + 4(2n - k)/3$. Hence $k \ge 10$. If among the excluded signplaces there were 6 signplaces on 3 places, then we can apply the bound for G_7 : $|A| \le \alpha(G_7) + 2 \cdot 6 = 32$. Otherwise there are at least n - 2 = 8 different places (and thus k is at most 12) in the remaining graph, and so one can apply computer calculations (see Appendix) for $m = 2n - k \le 10$, $l \ge 8$ and get the bounds by $2 \cdot 10 + 12 = 32$ or $2 \cdot 11 + 10 = 32$ when k = 10 or 11, respectively. For the case k = 12 Nagy's lemma gives us the inequality $|A| \le 2 \cdot 12 + 8 = 32$.

If a cobra exists, it cannot be intersected by any vertex (including the already deleted ones). The cobra uses at least 8 signplaces and leaves at least n-3=7 places, which are not fully used (i.e., at least one of

the signplaces on a place is free). Let t be the number of signplaces occupied by the cobra and m be equal to 2n - t. Then a graph G'_{10} , which is obtained by excluding the signplaces intersected by the cobra, has m signplaces lying on $l \ge 7$ different places.

Now we only have to sum up the number of vertices from the cobra and the number of vertices on the remaining signplaces (the last number is given in Appendix). So let us consider all the cases with t varying from 8 to n + 3 = 13:

- t = 8, m = 12, 3t 14 + 22 = 32;
- t = 9, m = 11, 3t 14 + 19 = 32;
- t = 10, m = 10, 3t 14 + 16 = 32;
- t = 11, m = 9, 3t 14 + 13 = 32, since we have at least 7 different places;
- t = 12, m = 8, 3t 14 + 10 = 32, since we have at least 7 different places;
- t = 13, m = 7, 3t 14 + 5 = 30, since all the places are distinct.

4.3. The case n = 11

We are going to prove that $\alpha(G_{11}) \leq 38$. Assume the contrary. If we have a quad, then after its exclusion we get exactly G_7 , but $\alpha(G_7) = 20$ and 20 + 16 < 38.

Now exclude one by one all the vertices with degree at most 2. Denote the number of excluded vertices by k. Following the proof of Lemma 1 we see that if k < 13, then a > 4, which means that a cobra exists. First, let $k \ge 13$. If among the excluded signplaces there where 8 signplaces on 4 places, then we can once again apply the bound for G_7 . Otherwise there are at least n - 3 = 8 different places in the remaining graph, so we can refer to Appendix with $m = 2n - k \le 9$, $l \ge 8$ and get at most 38 vertices in A.

If a cobra exists, it cannot be intersected by any vertex (including the already deleted ones). The cobra uses at least 8 signplaces and leaves at least n-3=8 places, which are not fully used. Let t be the number of signplaces occupied by the cobra and m be equal to 2n-t. Then a graph G'_{11} , which is obtained by excluding the signplaces intersected by the cobra, has m signplaces lying on $l \ge 8$ different places.

If $t \leq 8$, then we use the fact that all the signplaces, which are not in a cobra, have degrees at most two and get the declared bound: $3t - 14 + 2(22 - t) = t + 30 \leq 38$.

Now it remains to consider the cases with t from 9 to n + 3 = 14 and use the results from Appendix:

- t = 9, m = 13, 3t 14 + 25 = 38;
- t = 10, m = 12, 3t 14 + 22 = 38;
- t = 11, m = 11, 3t 14 + 19 = 38;
- t = 12, m = 10, 3t 14 + 16 = 38;
- t = 13, m = 9, 3t 14 + 13 = 38, since we have at least 8 different places;
- t = 14, m = 8, 3t 14 + 10 = 38, since all the places are distinct.

4.4. The case n = 13

We are going to prove that $\alpha(G_{13}) \leq 50$. Assume the contrary. If we have a quad, then after its exclusion we get exactly G_9 , but $\alpha(G_9) = 32$ and 32 + 16 < 50.

Now exclude one by one all the vertices with degree at most 2. Denote the number of excluded vertices by k. Following the proof of Lemma 1 we see that if k < 23, then a > 4, which means that a cobra exists. The case $k \ge 23$ is obvious.

Thus, a cobra exists, and it cannot be intersected by any vertex (including the already deleted ones). Define t as in the pevious cases. If t < 13, then we use the fact that all the signplaces, which are not in a cobra, have degrees at most two and get the declared bound: 3t - 14 + 2(26 - t) = t + 38 < 50.

Now it remains to consider the cases with t from 13 to n + 3 = 16 and use the results from Appendix:

- t = 13, m = 13, 3t 14 + 25 = 50;
- t = 14, m = 12, 3t 14 + 22 = 50;
- t = 15, m = 11, 3t 14 + 19 = 50;
- t = 16, m = 10, 3t 14 + 16 = 50.

The proof is complete.

5. Proof of Proposition 1

Suppose that $\chi(G_9) = 21$. Clearly $\frac{|V(G_9)|}{\alpha(G_9)} = 21$, and therefore every independent set has size 32. Revising the proof of Theorem 1 we see that the only way to reach 32 vertices in an independent set is by taking a couple of full quads. Thus, we have a collection of 21 pairs of full quads (denote it by A); this collection covers each base exactly two times, since every full quad has exactly 4 vertices on every covered base. Note that every pair of quads does not cover exactly one place, so one can split A into nine disjoint parts:

$$A = A_1 \sqcup \ldots \sqcup A_9.$$

Let S_1 be the set of all bases such that each of them does not contain the first place. Obviously $|S_1| = C_8^3 = 56$. Consider a pair of quads $p \in A$. Note that p covers 8 bases from S_1 , if $p \in A_1$, and 5 bases from S_1 otherwise. Denote the cardinalities of A_1 and $A \setminus A_1$ by a and b respectively. Every set in S_1 is covered twice, and therefore we have $2|S_1| = 112 = 8a + 5b$. Hence there are the following possibilities: (a = 14, b = 0), (a = 9, b = 8) and (a = 4, b = 16). But a + b = |A| = 21, so we get a contradiction. Proposition 1 is proved.

Appendix. Computer calculations

Let F be a subgraph of G_n . Denote the number of signplaces and places intersecting the vertices of F by m and l respectively.

We use the standard Bron–Kerbosch algorithm (see [3]) and get the following results:

- if m = 13, then $\alpha(F) \leq 25$;
- if m = 13 and l = 7, then $\alpha(F) \leq 18$;
- if m = 12, then $\alpha(F) \leq 22$;

- if m = 12 and l = 6, then $\alpha(F) \leq 16$;
- if m = 11, then $\alpha(F) \leq 19$;
- if m = 10, then $\alpha(F) \leq 16$;
- if m = 10 and $l \ge 8$, then $\alpha(F) \le 12$;
- if m = 9, then $\alpha(F) \leq 16$;
- if m = 9 and $l \ge 7$, then $\alpha(F) \le 13$;
- if m = 9 and $l \ge 8$, then $\alpha(F) \le 10$;
- if m = 8, then $\alpha(F) \leq 16$;
- if m = 8 and $l \ge 7$, then $\alpha(F) \le 10$;
- if m = 7, then $\alpha(F) \leq 10$;
- if m = l = 7, then $\alpha(F) \leq 5$;
- if m = 6, then $\alpha(F) \leq 7$;
- if m = 5, then $\alpha(F) \leq 5$;
- if m = 4, then $\alpha(F) \leq 4$;
- if m = 3, then $\alpha(F) \leq 1$.

Acknowledgements. Danila Cherkashin was supported by the grant of the Russian Scientific Foundation (grant N16-11-10039). All coauthors are grateful to Vyacheslav Sokolov for computer calculations.

References

- [1] P.K. Agarwal, J. Pach, Combinatorial geometry, John Wiley and Sons Inc., New York, 1995.
- [2] P. Brass, W. Moser, J. Pach, Research problems in discrete geometry, Springer, 2005.
- [3] C. Bron, J. Kerbosch, J. Algorithm 457: finding all cliques of an undirected graph, Communications of the ACM, 16 (1973), N9, 575 - 577.
- [4] K. Cantwell, Finite Euclidean Ramsey theory, J. Combin. Theory A, 73 (1996), N2, 273 285.
- [5] K.B. Chilakamarri, The unit-distance graph problem: A brief survey and some new results, Bull. Inst. Comb. Appl., 8 (1993), 39 - 60.
- [6] J. Cibulka, On the chromatic number of real and rational spaces, Geombinatorics, 2008.
- [7] G. Exoo, D. Ismailescu, On the Chromatic Number of \mathbb{R}^n for Small Values of n, http://arxiv.org/pdf/1408.2002v1.pdf.
- [8] G. Exoo, D. Ismailescu, M. Lim, On the Chromatic Number of ℝ⁴, Discrete Comput. Geom. 52 (2014), N2, 416 - 423.

- [9] H. Hadwiger, Ein Überdeckungssatz f
 ür den Euklidischen Raum, Portugaliae Math., 4 (1944), 140 -144.
- [10] M. Kahle, B. Taha, New lower bounds on $\chi(\mathbb{R}^d)$, $d = 8, \ldots, 12$, http://arxiv.org/pdf/1409.1278v1.pdf.
- [11] V. Klee, S. Wagon, Old and new unsolved problems in plane geometry and number theory, Math. Association of America, 1991.
- [12] A. Kupavskiy, On the chromatic number of \mathbb{R}^n with an arbitrary norm, Discrete Math., 311 (2011), N6, 437 440.
- [13] A.B. Kupavskiy, On coloring spheres embedded into \mathbb{R}^n , Sb. Math., 202 (2011), N6, 83 110.
- [14] A.B. Kupavskiy, A.M. Raigorodskii, On the chromatic number of \mathbb{R}^9 , J. of Math. Sci., 163 (2009), N6, 720 731.
- [15] D.G. Larman, C.A. Rogers, The realization of distances within sets in Euclidean space, Mathematika, 19 (1972), 1 - 24.
- [16] Z. Nagy, A certain constructive estimate of the Ramsey number, Matematikai Lapok, 23 (1972), N 301-302, 26.
- [17] O. Nechushtan, Note on the space chromatic number, Discrete Math., 256 (2002), 499 507.
- [18] A.M. Raigorodskii, Borsuk's problem and the chromatic numbers of metric spaces, Russian Math. Surveys, 56 (2001), N1, 103 - 139.
- [19] A.M. Raigorodskii, *Cliques and cycles in distance graphs and graphs of diameters*, "Discrete Geometry and Algebraic Combinatorics", AMS, Contemporary Mathematics, 625 (2014), 93 109.
- [20] A.M. Raigorodskii, On the chromatic number of a space, Russian Math. Surveys, 55 (2000), N2, 351 352.
- [21] A.M. Raigorodskii, Coloring Distance Graphs and Graphs of Diameters, Thirty Essays on Geometric Graph Theory, J. Pach ed., Springer, 2013, 429 - 460.
- [22] A. Soifer, The Mathematical Coloring Book, Springer, 2009.
- [23] L.A. Székely, Erdős on unit distances and the Szemerédi Trotter theorems, Paul Erdős and his Mathematics, Bolyai Series Budapest, J. Bolyai Math. Soc., Springer, 11 (2002), 649 - 666.