WEAVING SCHAUDER FRAMES

PETER G. CASAZZA, DANIEL FREEMAN, AND RICHARD G. LYNCH

ABSTRACT. We extend the concept of weaving Hilbert space frames to the Banach space setting. Similar to frames in a Hilbert space, we show that for any two approximate Schauder frames for a Banach space, every weaving is an approximate Schauder frame if and only if there is a uniform constant $C \geq 1$ such that every weaving is a C-approximate Schauder frame. We also study weaving Schauder bases, where it is necessary to introduce two notions of weaving. On one hand, we can ask if two Schauder bases are woven when considered as Schauder frames with their biorthogonal functionals, and alternatively, we can ask if each weaving of two Schauder bases remains a Schauder basis. We will prove that these two notions coincide when all weavings are unconditional, but otherwise they can be different. Lastly, we prove two perturbation theorems for approximate Schauder frames.

1. Introduction

The concept of weaving Hilbert space frames was first introduced in [BCGLL]. Two frames $\{x_j^0\}_{j\in J}$ and $\{x_j^1\}_{j\in J}$ for a Hilbert space H are called woven if there exist constants $0 < A \le B$ so that the weaving $\{x_j^0\}_{j\in \sigma} \cup \{x_j^1\}_{j\in \sigma^c}$ is a frame with bounds A, B for every choice of partition $\sigma \subset J$.

Weaving frames is motivated by distributed signal processing. We can think of each $j \in J$ as a sensor or node, and for each one we measure a signal with either x_j^0 or x_j^1 . Can a signal be recovered robustly regardless of how the measurements are taken at each node? Another way to think of this is if $(x_j^0)_{j\in J}$ and $(x_j^1)_{j\in J}$ are two sensor networks then we are interested in whenever parts of one network can be used to replace parts of the other. In [BCGLL], [CL], and [DV] the sensors are modeled by inner product with a vector in a frame or Riesz basis for a Hilbert space. In this paper, we extend the concept of weaving frames to the Banach space setting. We will model sensors by evaluation by linear functionals associated with an approximate Schauder frame or Schauder basis. We will be specifically considering weaving two bases or frames together, but we note that all of our results can be easily extended to weaving any finite number of bases or frames.

We begin by recalling the definition of a frame for a Hilbert space.

Definition 1.1. A set of vectors $(x_j)_{j\in J}$ in a Hilbert space H is called a *frame* for H if there exists positive constants A and B (called the lower and upper *frame bounds* respectively) such

²⁰⁰⁰ Mathematics Subject Classification. 46B20, 42C15.

The first and third authors were supported by NSF DMS 1307685; NSF ATD 1321779. The second author was supported by grant 353293 from the Simon's foundation.

that

$$A||x||^2 \le \sum_{i \in J} |\langle x, x_i \rangle|^2 \le B||x||^2 \quad \text{for all } x \in H.$$
 (1)

Given a set of vectors $(x_j)_{j\in J}$ in a Hilbert space H, we define an operator $S: H \to H$ (called the *frame operator*) by $S(x) := \sum_{j\in J} \langle x, x_i \rangle x_i$, for all $x\in H$, assuming that the series converges. The frame operator can be used to give a simple characterization of frames. Indeed, $(x_j)_{j\in J}$ is a frame if and only if the frame operator is well defined, bounded, and has bounded inverse. Furthermore, suppose that $(x_j)_{j\in J}$ is a frame with frame operator S, optimal lower frame bound S, and optimal upper frame bound S. Then |S| = B and $|S^{-1}| = A^{-1}$.

The characterization of a frame in terms of the frame operator is a useful perspective when considering how to generalize frame theory to Banach spaces. Frames have been generalized to Banach spaces in multiple ways. One way is to generalize the frame inequality (1) as in [G, FG], but the following definition of a Schauder frame and approximate Schauder frame instead generalizes the definition of the frame operator. Schauder frames were first used in [CDOSZ] with the goal of creating a procedure to represent vectors using quantized coefficients and are a generalization of framings of Banach spaces which were introduced in [CHL]. Approximate Schauder frames were defined in [FOSZ] and were used in the construction of a Schauder frame for $L_p(\mathbb{R}^d)$ with p > 2 formed by translations of a single function.

Definition 1.2. Let X be a Banach space with dual space X^* . A sequence $(x_i, f_i)_{i=1}^{\infty}$ in $X \times X^*$ is called an approximate Schauder frame for X if the operator $S: X \to X$ (called the frame operator) defined by $S(x) := \sum_{i=1}^{\infty} f_i(x)x_i$ is well defined and is bounded with bounded inverse. We say that $(x_i, f_i)_{i=1}^{\infty}$ is a C-approximate Schauder frame if $||S||, ||S^{-1}|| \le C$. We say that $(x_i, f_i)_{i=1}^{\infty}$ is a Schauder frame if the frame operator equals the identity operator.

An approximate Schauder frame is called unconditional if the series $\sum f_{\pi(i)}(x)x_{\pi(i)}$ converges to S(x) for every permutation π of \mathbb{N} . For $C_u \geq 1$, an unconditional approximate Schauder frame is called C_u -unconditional if for all $x \in X$ and all choices of $\varepsilon_i \in \{-1, 1\}$, $\|\sum_{i \in \mathbb{N}} \varepsilon_i f_i(x) x_i\| \leq C_u \|S(x)\|$. For $C_s \geq 1$, an unconditional approximate Schauder frame is C_s -suppression unconditional if for all $x \in X$ and all $\Gamma \subseteq \mathbb{N}$, $\|\sum_{i \in \Gamma} f_i(x) x_i\| \leq C_s \|S(x)\|$. As is the case for unconditional Schauder bases, we have that an approximate Schauder frame is unconditional if and only if it suppression unconditional and $C_s \leq C_u \leq 2C_s$.

Note that if $(x_i, f_i)_{i=1}^{\infty}$ is an approximate Schauder frame for a Banach space X with frame operator S then $(x_i, (S^{-1})^* f_i)_{i=1}^{\infty}$ is a Schauder frame for X. This is a generalization of the often used result that if $(x_i)_{i=1}^{\infty}$ is a frame for a Hilbert space H with frame operator S then $(S^{-1/2}x_i)_{i=1}^{\infty}$ is a Parseval frame for H.

Our goal is to extend the notion of weaving frames in a Hilbert space to weaving frames and bases for Banach spaces and to prove the analogous results in this setting. This is a different generalization of weaving frames than given in [DV], where they instead consider weaving together infinitely many Hilbert space frames.

Since ordering plays a crucial role, we index the weavings with elements from $\{0,1\}^{\mathbb{N}}$, the set of functions from \mathbb{N} to $\{0,1\}$. Note that $\{0,1\}^{\mathbb{N}}$ is a compact Hausdorff space in the product topology. We will repeatedly make use of this fact.

Definition 1.3. [BCGLL, Definition 3] Let $(x_j^0)_{j=1}^{\infty}$ and $(x_j^1)_{j=1}^{\infty}$ be two frames for a Hilbert space H. We say that $(x_j^0)_{j=1}^{\infty}$ and $(x_j^1)_{j=1}^{\infty}$ are woven if there are positive constants A and B such that for every function $\sigma \in \{0,1\}^{\mathbb{N}}$ we have that $(x_j^{\sigma(j)})_{j=1}^{\infty}$ is a frame for H with lower frame bound A and upper frame bound B.

We give the following natural generalization of weaving frames for Hilbert spaces to weaving approximate Schauder frames for Banach spaces.

Definition 1.4. Let $(x_j^0, f_j^0)_{j=1}^{\infty}$ and $(x_j^1, f_j^1)_{j=1}^{\infty}$ be two approximate Schauder frames for a Banach space X. A sequence $(x_j^{\sigma(j)}, f_j^{\sigma(j)})_{j=1}^{\infty}$ with $\sigma \in \{0, 1\}^{\mathbb{N}}$ is called a weaving of $(x_j^0, f_j^0)_{j=1}^{\infty}$ and $(x_j^1, f_j^1)_{j=1}^{\infty}$. Given $C \geq 1$, we say that $(x_j^0, f_j^0)_{j=1}^{\infty}$ and $(x_j^1, f_j^1)_{j=1}^{\infty}$ are C-woven if every weaving is a C-approximate Schauder frame. We say that $(x_j^0, f_j^0)_{j=1}^{\infty}$ and $(x_j^1, f_j^1)_{j=1}^{\infty}$ are woven if they are C-woven for some $C \geq 1$.

Note that this is a true generalization of the definition of woven frames for Hilbert spaces as two frames $(x_j^0)_{j=1}^{\infty}$ and $(x_j^1)_{j=1}^{\infty}$ of a Hilbert space H are woven if and only if the approximate Schauder frames $(x_j^0, x_j^0)_{j=1}^{\infty}$ and $(x_j^1, x_j^1)_{j=1}^{\infty}$ are woven (where we identify a vector $y \in H$ with the linear functional $x \mapsto \langle x, y \rangle$).

A Riesz basis for a Hilbert space H is any sequence of vectors that is image of some orthonormal basis under an invertible operator (this is equivalent to being a semi-normalized unconditional basis for a Hilbert space). In [BCGLL], it is proven that if $(x_j^0)_{j=1}^{\infty}$ and $(x_j^1)_{j=1}^{\infty}$ are two Riesz bases for a Hilbert space H then $(x_j^{\sigma(j)})_{j=1}^{\infty}$ is a Riesz basis for H for all $\sigma \in \{0,1\}^{\mathbb{N}}$. Thus, there is no need to distinguish between two Riesz bases having all weavings be Riesz bases and two Riesz bases having all weavings be frames. However, we will show that these two notions are not always the same for Schauder bases and thus we will require a second definition of weaving in this context.

A sequence $(x_j)_{j=1}^{\infty}$ in a Banach space X is a Schauder basis for X if for every $x \in X$ there exists a unique sequence of scalars $(a_j)_{j=1}^{\infty}$ such that $x = \sum_{j=1}^{\infty} a_j x_j$, and a basis is called unconditional if the series converges in every order for every $x \in X$. The biorthogonal functionals of $(x_j)_{j=1}^{\infty}$ is the sequence $(x_j^*)_{j=1}^{\infty}$ in X^* such that $x_j^*(x) = a_j$ for all $j \in \mathbb{N}$. A sequence $(x_j)_{j=1}^{\infty}$ in a Banach space X is called a basic sequence if it is a basis of its closed span $[x_j]_{j=1}^{\infty}$. The basis constant of a basic sequence $(x_j)_{j=1}^{\infty}$ is the least constant $C \geq 1$ such that $\|\sum_{i=1}^n a_i x_i\| \leq C \|\sum_{i=1}^N a_i x_i\|$ for all choices of scalars $(a_i)_{i=1}^N$ and all $n \leq N$.

Definition 1.5. Let $(x_j^0)_{j=1}^{\infty}$ and $(x_j^1)_{j=1}^{\infty}$ be two Schauder bases for a Banach space X. A sequence $(x_j^{\sigma(j)})_{j=1}^{\infty}$ with $\sigma \in \{0,1\}^{\mathbb{N}}$ is called a weaving of $(x_j^0)_{j=1}^{\infty}$ and $(x_j^1)_{j=1}^{\infty}$. We say that $(x_j^0)_{j=1}^{\infty}$ and $(x_j^1)_{j=1}^{\infty}$ are woven if every weaving is a Schauder basis for X.

Let $(x_j^0)_{j=1}^\infty$ and $(x_j^1)_{j=1}^\infty$ be two Schauder bases for a Banach space X with biorthogonal functionals $(x_j^{0*})_{j=1}^{\infty}$ and $(x_j^{1*})_{j=1}^{\infty}$. This implies that $(x_j^0, x_j^{0*})_{j=1}^{\infty}$ and $(x_j^1, x_j^{1*})_{j=1}^{\infty}$ are both Schauder frames for X. We can now consider whether or not the two bases $(x_j^0)_{j=1}^{\infty}$ and $(x_j^1)_{j=1}^{\infty}$ are woven and whether or not the two Schauder frames $(x_j^0, x_j^{0*})_{j=1}^{\infty}$ and $(x_j^1, x_j^{1*})_{j=1}^{\infty}$ are woven. In Example 3.2, we show that it is possible for two Schauder bases to be woven but for the corresponding Schauder frames to not be woven. The key idea for why this is possible is that when we weave approximate Schauder frames we are weaving both two sequences of vectors and two sequences of functionals, but if we weave two Schauder bases and obtain a basis the new corresponding biorthogonal functionals can be completely unrelated to the biorthogonal functionals of the two bases that we started with. This gives us more flexibility when weaving Schauder bases, but it is actually a problem when we consider our motivation as determining when different sensor networks can be combined. In that context, the given linear functionals are fundamental components of the sensors and thus we cannot change them based on what weaving we choose to use. For this reason, it is important for applications for two Schauder bases to be woven both as bases and as approximate Schauder frames. We prove that this is the case when all weavings are unconditional. In particular, in Theorem 4.5 we prove a number of equivalent properties, including that if $(x_i^0)_{i=1}^{\infty}$ and $(x_i^1)_{i=1}^{\infty}$ are two unconditional Schauder bases then every weaving of $(x_j^0)_{j=1}^{\infty}$ and $(x_j^1)_{j=1}^{\infty}$ is an unconditional basis if and only if every weaving of $(x_j^0, x_j^{0*})_{j=1}^{\infty}$ and $(x_i^1, x_i^{1*})_{i=1}^{\infty}$ is an unconditional approximate Schauder frame. In Theorem 4.5 we prove as well that if $(x_j^0)_{j=1}^\infty$ and $(x_j^1)_{j=1}^\infty$ are two unconditional Schauder bases of a Banach space X then every weaving of $(x_i^0)_{i=1}^{\infty}$ and $(x_i^1)_{i=1}^{\infty}$ is an unconditional basis of X if and only if every weaving is an unconditional basic sequence. Surprisingly, we show that this is not always the case if some weavings are conditional, and in Example 3.5 we give two unconditional Schauder bases for ℓ_1 such that every weaving is a basic sequence, but there exists a weaving which is not a Schauder basis for all of ℓ_1 .

Lastly, in Section 5 we prove two perturbation theorems for approximate Schauder frames and prove that these result in woven approximate Schauder frames. For a thorough approach to the basics of frame theory, see [CK, CL2, C]. For background on Banach space theory see [FHHMPZ] and [LT].

2. Uniform constants for woven approximate Schauder frames

One of the major results in [BCGLL] is a proof that if $(x_j^0)_{j=1}^{\infty}$ and $(x_j^1)_{j=1}^{\infty}$ are two frames for a Hilbert space H, then every weaving is a frame for H if and only if there is a uniform constant $C \geq 1$ such that $(x_j^0)_{j=1}^{\infty}$ and $(x_j^1)_{j=1}^{\infty}$ are C-woven. We shall prove in Theorem 2.4 that this same uniformity result holds more generally for approximate Schauder frames.

Let $(x_j^0, f_j^0)_{j=1}^{\infty}$ and $(x_j^1, f_j^1)_{j=1}^{\infty}$ be two approximate Schauder frames of X such that every weaving is an approximate Schauder frame. For $I \subseteq \mathbb{N}$ and $\sigma \in \{0, 1\}^{\mathbb{N}}$, we define a linear operator $P_{\sigma,I}: X \to X$ by $P_{\sigma,I}(x) = \sum_{j \in I} f_j^{\sigma(j)}(x) x_j^{\sigma(j)}$. Note that this series will converge for every interval $I \subseteq \mathbb{N}$, but that the series may not converge if the approximate Schauder frame is conditional and I is not an interval. In this notation, we have that $P_{\sigma,\mathbb{N}}$ is the frame

operator of the weaving $(x_j^{\sigma(j)}, f_j^{\sigma(j)})_{j=1}^{\infty}$. Thus, to prove that two approximate Schauder frames $(x_j^0, f_j^0)_{j=1}^{\infty}$ and $(x_j^1, f_j^1)_{j=1}^{\infty}$ are \tilde{C} -woven, we need to prove that $||P_{\sigma,\mathbb{N}}|| \leq C$ and $||P_{\sigma,\mathbb{N}}^{-1}|| \leq C$ for all $\sigma \in \{0,1\}^{\mathbb{N}}$.

Lemma 2.1. Let $(x_j^0, f_j^0)_{j=1}^{\infty}$ and $(x_j^1, f_j^1)_{j=1}^{\infty}$ be two approximate Schauder frames for X such that every weaving is an approximate Schauder frame for X. If $x \in X$ and $\varepsilon > 0$, then there exists $N \in \mathbb{N}$ such that $||P_{\sigma,[m,n]}x|| < \varepsilon$ for all $\sigma \in \{0,1\}^{\mathbb{N}}$ and $n \geq m \geq N$.

Proof. Let $x \in X$ and $\varepsilon > 0$. Assume the contrary that for all $N \in \mathbb{N}$, we may choose $\sigma \in \{0,1\}^{\mathbb{N}}$ and $n \geq m \geq N$ such that $\|P_{\sigma,[m,n]}x\| \geq \varepsilon$. We may then inductively choose $(\sigma_j)_{j=1}^{\infty}$ and $m_1 \leq n_1 < m_2 \leq n_2 < \dots$ such that $\|P_{\sigma_j,[m_j,n_j]}x\| \geq \varepsilon$. As, $([m_j,n_j])_{j=1}^{\infty}$ is a sequence of disjoint intervals, we may choose $\sigma \in \{0,1\}^{\mathbb{N}}$ such that $\sigma|_{[m_j,n_j]} = \sigma_j|_{[m_j,n_j]}$ for all $j \in \mathbb{N}$. Thus, we have that $\|P_{\sigma,[m_j,n_j]}x\| \geq \varepsilon$ for all $j \in \mathbb{N}$. Since we assumed that $(x_j^{\sigma(j)}, f_j^{\sigma(j)})$ is an approximate Schauder frame, this contradicts the fact that $P_{\sigma,\mathbb{N}}x$ converges.

Lemma 2.2. Let $(x_j^0, f_j^0)_{j=1}^\infty$ and $(x_j^1, f_j^1)_{j=1}^\infty$ be two approximate Schauder frames for X such that every weaving is an approximate Schauder frame for X. There exists a constant D > 0such that $||P_{\sigma,[m,n]}|| \leq D$ for all $\sigma \in \{0,1\}^{\mathbb{N}}$ and $m,n \in \mathbb{N}$.

Proof. Let $x \in X$. By Lemma 2.1 there exists $N \in \mathbb{N}$ such that $\|P_{\sigma,[n,\infty)}x\| < 1$ for all $\sigma \in \{0,1\}^{\mathbb{N}}$ and all $n \geq N$. We have that

$$\sup_{\sigma \in \{0,1\}^{\mathbb{N}}, n \le N} \|P_{\sigma,[n,\infty)}x\| \le \sup_{\sigma \in \{0,1\}^{\mathbb{N}}, n \le N} \|P_{\sigma,[n,N)}x\| + \sup_{\sigma \in \{0,1\}^{\mathbb{N}}} \|P_{\sigma,[N,\infty)}x\|.$$

Thus, we have

$$\sup_{\sigma \in \{0,1\}^{\mathbb{N}}, n \le N} \|P_{\sigma,[n,\infty)}x\| < \infty$$

as $\sup_{\sigma} \|P_{\sigma,[N,\infty)}x\| \le 1$ and the set $\{P_{\sigma,[n,N)}x\}_{\sigma\in\{0,1\}^{\mathbb{N}},n\le N}$ is finite. Hence, there exists a constant K > 0 such that $\|P_{\sigma,[n,\infty)}x\| \leq K$ for all $\sigma \in \{0,1\}^{\mathbb{N}}$ and all $n \in \mathbb{N}$. Therefore, for all $m, n \in \mathbb{N}$ with $m \leq n$ we have that

$$||P_{\sigma,[m,n]}x|| = ||P_{\sigma,[m,\infty)}x - P_{\sigma,[n+1,\infty)}x|| \le 2K.$$

Hence, $\{P_{\sigma,[m,n]}\}_{\sigma\in\{0,1\}^{\mathbb{N}},m\leq n}$ is a set of bounded operators such that for all $x\in X$, $\{P_{\sigma,[m,n]}x\}$ is uniformly bounded in norm. By the uniform boundedness principle, the set of operators $\{P_{\sigma,[m,n]}\}$ is uniformly bounded in norm.

Lemma 2.3. Let $(x_j^0, f_j^0)_{j=1}^{\infty}$ and $(x_j^1, f_j^1)_{j=1}^{\infty}$ be two approximate Schauder frames of X such that every weaving is an approximate Schauder frame of X. For all $x \in X$ there exists $\delta > 0$ such that for all $\sigma \in \{0,1\}^{\mathbb{N}}$ we have that $||P_{\sigma,\mathbb{N}}x|| \geq \delta ||x||$

Proof. Let $x \in X$ such that $x \neq 0$. By Lemma 2.1, for all $n \in \mathbb{N}$ there exists $M_n \in \mathbb{N}$ such that $\|P_{\gamma,[M_n,\infty)}x\|<\frac{1}{n}$ for all $\gamma\in\{0,1\}^{\mathbb{N}}$. Assume for the sake of contradiction that for all $n\in\mathbb{N}$ there exists $\sigma_n \in \{0,1\}^{\mathbb{N}}$ such that $\|P_{\sigma_n,\mathbb{N}}x\| < 1/n$. After passing to a subsequence, we may assume that $(\sigma_n)_{n=1}^{\infty}$ converges to some $\sigma \in \{0,1\}^{\mathbb{N}}$ and $\sigma_n|_{[1,M_n)} = \sigma|_{[1,M_n)}$ for all $n \in \mathbb{N}$. For each $n \in \mathbb{N}$, we have the following upperbound on the norm of $P_{\sigma,\mathbb{N}}x$.

$$||P_{\sigma,\mathbb{N}}x|| = ||P_{\sigma_n,\mathbb{N}}x - P_{\sigma_n,[M_n,\infty)}x + P_{\sigma,[M_n,\infty)}x||$$

$$\leq ||P_{\sigma_n,\mathbb{N}}x|| + ||P_{\sigma_n,[M_n,\infty)}x|| + ||P_{\sigma,[M_n,\infty)}x||$$

$$< 1/n + 1/n + 1/n$$

Thus, $||P_{\sigma,\mathbb{N}}x|| < 3/n$ for all $n \in \mathbb{N}$ and hence $P_{\sigma,\mathbb{N}}x = 0$. This contradicts that $(x_j^{\sigma(j)}, f_j^{\sigma(j)})_{j=1}^{\infty}$ is an approximate Schauder frame.

Theorem 2.4. Given two approximate Schauder frames for a Banach space X, they are woven if and only if every weaving is an approximate Schauder frame. That is, if every weaving is an approximate Schauder frame then there is a uniform constant $C \geq 1$ such that every weaving is a C-approximate Schauder frame. Furthermore, if every weaving is an unconditional approximate Schauder frame then there exists a constant $D \geq 1$ such that every weaving is a D-unconditional approximate Schauder frame.

Proof. Let $(x_j^0, f_j^0)_{j=1}^\infty$ and $(x_j^1, f_j^1)_{j=1}^\infty$ be two approximate Schauder frames of X such that every weaving is an approximate Schauder frame. Let $y \in X$. By Lemma 2.3, there exists $\delta > 0$ such that $\|y\| = \|P_{\sigma,\mathbb{N}}(P_{\sigma,\mathbb{N}}^{-1}y)\| \geq \delta \|P_{\sigma,\mathbb{N}}^{-1}y\|$ for all $\sigma \in \{0,1\}^{\mathbb{N}}$. Thus, $\|P_{\sigma,\mathbb{N}}^{-1}y\| \leq (1/\delta)\|y\|$ for all $\sigma \in \{0,1\}^{\mathbb{N}}$. We have that $\{P_{\sigma,\mathbb{N}}^{-1}\}_{\sigma \in \{0,1\}^{\mathbb{N}}}$ is a collection of bounded operators such that $\sup_{\sigma \in \{0,1\}^{\mathbb{N}}} \|P_{\sigma,\mathbb{N}}^{-1}\| < \infty$ by the uniform boundedness principle. Also, $\sup_{\sigma \in \{0,1\}^{\mathbb{N}}} \|P_{\sigma,\mathbb{N}}^{-1}\| < \infty$ by Lemma 2.2. Thus, there is a uniform constant $C \geq 1$ such that $\|P_{\sigma,\mathbb{N}}\| \leq C$ and $\|P_{\sigma,\mathbb{N}}^{-1}\| \leq C$ for all $\sigma \in \{0,1\}^{\mathbb{N}}$ and hence $(x_j^0, f_j^0)_{j=1}^\infty$ and $(x_j^1, f_j^1)_{j=1}^\infty$ are C-woven approximate Schauder frames of X.

We now assume that every weaving of $(x_j^0, f_j^0)_{j=1}^\infty$ and $(x_j^1, f_j^1)_{j=1}^\infty$ is an unconditional approximate Schauder frame. We have that there exists $C \geq 1$ such that every weaving of $(x_j^0, f_j^0)_{j=1}^\infty$ and $(x_j^1, f_j^1)_{j=1}^\infty$ is a C-approximate Schauder frame. For the sake of contradiction we assume that for all $n \in \mathbb{N}$, there exists $\sigma_n \in \{0, 1\}^\mathbb{N}$ such that the approximate Schauder $(x_j^{\sigma_n(j)}, f_j^{\sigma_n(j)})_{j=1}^\infty$ is not n-suppression unconditional. After passing to a subsequence, we may assume that there exists $\sigma \in \{0, 1\}^\mathbb{N}$ such that $\sigma_n \to \sigma$ in the product topology of $\{0, 1\}^\mathbb{N}$ and that $\sigma_n|_{[1,n]} = \sigma|_{[1,n]}$ for all $n \in \mathbb{N}$. We choose a subsequence $(m_n)_{n=1}^\infty$ of \mathbb{N} such that there exists a finite set $\Gamma_n \subseteq [1, m_n]$ and $x_n \in X$ with $||x_n|| = 1$ so that $||P_{\sigma_n,\Gamma_n}x_n|| \geq n||P_{\sigma_n,\mathbb{N}}x_n|| \geq nC^{-1}$.

We inductively define a sequence of natural numbers $(M_n)_{n=1}^{\infty}$ by $M_1 = m_1$ and for $n \in \mathbb{N}$ we let $M_{n+1} = m_{M_n}$. Let $\gamma \in \{0,1\}^{\mathbb{N}}$ be such that $\gamma|_{[1,m_1]} = \sigma_1|_{[1,m_1]}$ and for $n \in \mathbb{N}$, $\gamma|_{(M_n,M_{n+1}]} = \sigma_{M_n}|_{(M_n,M_{n+1}]}$. Let $D \geq 1$ be the suppression unconditionality constant of the weaving $(x_j^{\sigma(j)}, f_j^{\sigma(j)})_{j=1}^{\infty}$. For $n \in \mathbb{N}$, we have that

$$||P_{\gamma,\Gamma_{M_n}\cap(M_n,M_{n+1}]}|| = ||P_{\sigma_{M_n},\Gamma_{M_n}\cap(M_n,M_{n+1}]}||$$

$$\geq ||P_{\sigma_{M_n},\Gamma_{M_n}}|| - ||P_{\sigma_{M_n},\Gamma_{M_n}\cap[1,M_n]}||$$

$$= ||P_{\sigma_{M_n},\Gamma_{M_n}}|| - ||P_{\sigma,\Gamma_{M_n}\cap[1,M_n]}|| \geq M_n C^{-1} - DC.$$

Thus, we have that $\sup_{n\in\mathbb{N}} \|P_{\gamma,\Gamma_{M_n}\cap(M_n,M_{n+1}]}\| = \infty$ and hence $(x_j^{\gamma(j)},f_j^{\gamma(j)})_{j=1}^{\infty}$ is not unconditional.

3. Weaving bases

We now consider the case of weaving Schauder bases instead of approximate Schauder frames. Recall that if $(x_j)_{j=1}^{\infty}$ is a Schauder basic sequence then we say that it is C-basic for some $C \geq 1$ if $\|\sum_{j=1}^N a_j x_j\| \leq C \|\sum_{j=1}^\infty a_j x_j\|$ for all $N \in \mathbb{N}$ and all sequences of scalars $(a_j)_{j=1}^{\infty}$ such that $\sum_{j=1}^\infty a_i x_i$ converges. We say that an unconditional Schauder basic sequence is C-suppression unconditional if $\|\sum_{j\in E} a_j x_j\| \leq C \|\sum_{j=1}^\infty a_j x_j\|$ for all $E \subseteq \mathbb{N}$ and all sequences of scalars $(a_j)_{j=1}^\infty$ such that $\sum_{j=1}^\infty a_j x_j$ converges.

In Section 2 we proved that if every weaving of two approximate Schauder frames is an approximate Schauder frame then there is a uniform constant $C \geq 1$ such that the two approximate Schauder frames are C-woven. We now prove that this same uniformity theorem holds for weaving Schauder bases. The proof will follow the same concatonation argument as in the furthermore part of the proof of Theorem 2.4.

Proposition 3.1. Let $(x_i^0)_{i=1}^{\infty}$ and $(x_i^1)_{i=1}^{\infty}$ be two Schauder basic sequences in a Banach space X. If $(x_i^{\sigma(i)})_{i=1}^{\infty}$ is a basic sequence for all $\sigma \in \{0,1\}^{\mathbb{N}}$, then there is a uniform constant $C \geq 1$ such that for all $\sigma \in \{0,1\}^{\mathbb{N}}$, $(x_i^{\sigma(i)})_{i=1}^{\infty}$ is C-basic. Likewise, if $(x_j^{\sigma(j)})_{j=1}^{\infty}$ is an unconditional basic sequence for all $\sigma \in \{0,1\}^{\mathbb{N}}$, then there is a uniform constant $D \geq 1$ such that for all $\sigma \in \{0,1\}^{\mathbb{N}}$, $(x_i^{\sigma(j)})_{j=1}^{\infty}$ is D-unconditional.

Proof. Assume by way of contradiction that $(x_i^{\sigma(i)})_{i=1}^{\infty}$ is a basic sequence for all $\sigma \in \{0,1\}^{\mathbb{N}}$, but that for all $n \in \mathbb{N}$, there exists $\sigma_n \in \{0,1\}^{\mathbb{N}}$ such that $(x_i^{\sigma_n(i)})_{i=1}^{\infty}$ is not n-basic. After passing to a subsequence, we may assume that there exists $\sigma \in \{0,1\}^{\mathbb{N}}$ such that $\sigma_n \to \sigma$ in the product topology of $\{0,1\}^{\mathbb{N}}$. Let $C \geq 1$ be such that $(x_i^{\sigma(i)})_{i=1}^{\infty}$ is C-basic. After passing to a further subsequence, we may assume that $\sigma_n|_{[1,n]} = \sigma|_{[1,n]}$ for all $n \in \mathbb{N}$. Thus, for all $n \in \mathbb{N}$, $(x_i^{\sigma_n(i)})_{i=1}^n$ is C-basic, and hence $(x_i^{\sigma_n(i)})_{i=n+1}^{\infty}$ is not (n-C)-basic. Let $(m_n)_{n=1}^{\infty}$ be an increasing subsequence of natural numbers such that for all $n \in \mathbb{N}$, $(x_i^{\sigma_n(i)})_{i=n+1}^{m_n}$ is not (n-C)-basic. We can now create $\gamma \in \{0,1\}^{\mathbb{N}}$ by $\gamma|_{[1,m_1]} = \sigma_1|_{[1,m_1]}$, $\gamma|_{(m_1,m_{m_1}]} = \sigma_{m_1}|_{(m_1,m_{m_1}]}$, and so on. We have that $(x_i^{\gamma(i)})_{i=1}^{\infty}$ is not basic because it is formed by concatonating finite length basic sequences whose basis constants go to infinity, which is a contradiction.

A similar argument shows that if for all $\sigma \in \{0,1\}^{\mathbb{N}}$, $(x_i^{\sigma(i)})_{i=1}^{\infty}$ is an unconditional basic sequence then there is a universal constant $D \geq 1$ such that for all $\sigma \in \{0,1\}^{\mathbb{N}}$, $(x_i^{\sigma(i)})_{i=1}^{\infty}$ is D-unconditional.

Suppose that $(x_j^0)_{j=1}^\infty$ and $(x_j^1)_{j=1}^\infty$ are two Schauder bases for a Banach space X with biorthogonal functionals $(x_i^{0*})_{j=1}^\infty$ and $(x_i^{1*})_{j=1}^\infty$. We can consider the problem of whether or not $(x_i^0)_{j=1}^\infty$

and $(x_j^1)_{j=1}^\infty$ are woven Schauder bases, and we can consider the problem of whether or not $(x_j^0, x_j^{0*})_{j=1}^\infty$ and $(x_j^1, x_j^{1*})_{j=1}^\infty$ are woven approximate Schauder frames. The first natural question to ask is if these two notions are equivalent. In other words, are $(x_i^0)_{j=1}^\infty$ and $(x_j^1)_{j=1}^\infty$ woven Schauder bases if and only if $(x_j^0, x_j^{0*})_{j=1}^\infty$ and $(x_j^1, x_j^{1*})_{j=1}^\infty$ are woven approximate Schauder frames? Surprisingly, the answer is no.

Example 3.2. We have the following examples in c_0 and ℓ_1 .

- (1) Let $(e_j)_{j=1}^{\infty}$ be the standard unit vector basis for c_0 with biorthogonal functionals $(e_j^*)_{j=1}^{\infty}$, and let $(s_j)_{j=1}^{\infty}$ be the summing basis for c_0 with biorthogonal functionals $(s_j^*)_{j=1}^{\infty}$. That is, $s_j = \sum_{i=1}^{j} e_i$ for all $j \in \mathbb{N}$. Then, $(s_j)_{j=1}^{\infty}$ is a conditional Schauder basis for c_0 which is woven with the unconditional Schauder basis $(e_j)_{j=1}^{\infty}$, but $(s_j, s_j^*)_{j=1}^{\infty}$ and $(e_j, e_j^*)_{j=1}^{\infty}$ are not woven approximate Schauder frames.
- (2) Let $(e_j)_{j=1}^{\infty}$ be the standard unit vector basis for ℓ_1 with biorthogonal functionals $(e_j^*)_{j=1}^{\infty}$. Let $x_1 = e_1$ and $x_n = e_n - e_{n-1}$ for n > 1. Then, $(x_j)_{j=1}^{\infty}$ is a conditional Schauder basis for ℓ_1 which is woven with the unconditional Schauder basis $(e_j)_{j=1}^{\infty}$, but $(x_j, x_j^*)_{j=1}^{\infty}$ and $(e_j, e_j^*)_{j=1}^{\infty}$ are not woven approximate Schauder frames, where $(x_j^*)_{j=1}^{\infty}$ is the sequence of biorthogonal functionals to $(x_j)_{j=1}^{\infty}$.

Example 3.2 shows that it is possible for a conditional Schauder basis to be woven with an unconditional Schauder basis. The following proposition shows that this is impossible for woven approximate Schauder frames.

Proposition 3.3. Let $(x_j^0, f_j^0)_{j=1}^\infty$ and $(x_j^1, f_j^1)_{j=1}^\infty$ be C-approximate Schauder frames for a Banach space X. Suppose that $(x_j^0, f_j^0)_{j=1}^\infty$ is K-suppression unconditional and that $(x_j^1, f_j^1)_{j=1}^\infty$ is not D-suppression unconditional. Then $(x_j^0, f_j^0)_{j=1}^\infty$ and $(x_j^1, f_j^1)_{j=1}^\infty$ are not $(DC^{-1} - KC)$ -woven.

Proof. Let S_0 be the frame operator for $(x_j^0, f_j^0)_{j=1}^{\infty}$ and let S_1 be the frame operator for $(x_j^1, f_j^1)_{j=1}^{\infty}$. Let $x \in X$ and $A \subset \mathbb{N}$ such that $\|\sum_{j \in A} f_j^1(x) x_j^1\| > D \|S_1 x\| \ge DC^{-1} \|x\|$. Let $\sigma \in \{0, 1\}^{\mathbb{N}}$ be the characteristic function of A. Then

$$\left\| \sum_{j \in \mathbb{N}} f_j^{\sigma(j)}(x) x_j^{\sigma(j)} \right\| \ge \left\| \sum_{j \in A} f_j^1(x) x_j^1 \right\| - \left\| \sum_{j \notin A} f_j^0(x) x_j^0 \right\|$$

$$> D \|S_1 x\| - K \|S_0 x\|$$

$$\ge D C^{-1} \|x\| - K C \|x\|.$$

Thus the frame operator of $(x_j^{\sigma(j)}, f_j^{\sigma(j)})_{j=1}^{\infty}$ has norm greater than $(DC^{-1} - KC)$ and hence $(x_j^0, f_j^0)_{j=1}^{\infty}$ and $(x_j^1, f_j^1)_{j=1}^{\infty}$ are not $(DC^{-1} - KC)$ -woven.

We conclude the section with two examples that again demonstrate the significance of conditional compared to unconditional convergence. The next construction shows that it is possible for two woven unconditional Schauder bases to have some conditional weavings.

Example 3.4. Let $(e_j)_{j=1}^{\infty}$ be the standard unit vector basis for ℓ_1 . Define $(x_i^0)_{i=1}^{\infty}$ by $x_{2n-1}^0 = e_n$ and $x_{2n}^0 = e_{2n} - e_{2n-1}$ for all $n \in \mathbb{N}$ and define $(x_i^1)_{i=1}^{\infty}$ by $x_1^1 = e_1$, $x_2^1 = e_2$, and $x_{2n-1}^1 = e_{2n-1} - e_{2n-2}$ and $x_{2n}^1 = e_{2n}$ for all $n \geq 2$. Then $(x_i^0)_{i=1}^{\infty}$ and $(x_i^1)_{i=1}^{\infty}$ are woven Schauder bases, each of which are unconditional. However, the alternating weaving given by $(x_1^1, x_2^0, x_3^1, x_4^0, ...) = (e_1, e_2 - e_1, e_3 - e_2, e_4 - e_3, ...)$ is the same conditional basis used in Example 3.2.

The following construction shows it is possible for all the weavings of two unconditional Schauder bases to be Schauder basic sequences, but there exists a weaving which does not have dense span. In [BCGLL], it is proven that this is impossible for Riesz bases for Hilbert spaces, and in Theorem 4.5 we prove that this is impossible if all weavings are unconditional.

Example 3.5. Let $(e_j)_{j=1}^{\infty}$ be the standard unit vector basis for ℓ_1 . Define $(x_i^0)_{i=1}^{\infty}$ by $x_{2n-1}^0 = e_{2n-1} + e_{2n}$ and $x_{2n}^0 = e_{2n-1} - e_{2n}$ for all $n \in \mathbb{N}$ and define $(x_i^1)_{i=1}^{\infty}$ by $x_1^1 = e_1$ and $x_{2n}^1 = e_{2n} + e_{2n+1}$ and $x_{2n+1}^1 = e_{2n} - e_{2n+1}$ for all $n \in \mathbb{N}$. Then $(x_i^0)_{i=1}^{\infty}$ and $(x_i^1)_{i=1}^{\infty}$ are both unconditional Schauder bases of ℓ_1 and every weaving is a Schauder basic sequence. However, the alternating weaving $(x_1^0, x_2^1, x_3^0, x_4^1, \ldots) = (e_1 + e_2, e_2 + e_3, e_3 + e_4, e_4 + e_5, \ldots)$ does not contain e_1 in its closed span and thus is not a basis.

4. Weaving unconditional Schauder bases

In this section, we consider weaving unconditional bases. First, we show that two woven unconditional bases are necessarily equivalent. We conclude the section by proving that two unconditional bases that are woven and have all unconditional weavings is equivalent to five other characterizations, including in particular that all weavings are unconditional basic sequences and all weavings of the corresponding approximate Schauder frames are unconditional approximate Schauder frames. Recall that in Section 3 we gave counterexamples to these equivalences when it is not the case that all weavings are unconditional. We first cite the following theorem, which can be found in [LT] along with a thorough introduction to Banach lattices.

Theorem 4.1. Let $(x_i)_{i=1}^{\infty}$ be an unconditional basis for a Banach lattice X. There exists a constant $D \ge 1$ such that for every $n \in \mathbb{N}$ and sequence of scalars $(a_i)_{i=1}^n$, we have that

$$D^{-1} \left\| \left(\sum_{i=1}^{n} |a_i x_i|^2 \right)^{1/2} \right\| \le \left\| \sum_{i=1}^{n} a_i x_i \right\| \le D \left\| \left(\sum_{i=1}^{n} |a_i x_i|^2 \right)^{1/2} \right\|.$$

When we apply Theorem 4.1, the Banach lattice structure for X will be defined by a different 1-unconditional basis $(u_i)_{i=1}^{\infty}$ with biorthogonal functionals $(u_i^*)_{i=1}^{\infty}$. In this case, if $(x_i)_{i=1}^{\infty}$ is a sequence in X, and $(a_i)_{i=1}^n$ is a finite sequence of scalars then the vector $(\sum_{i=1}^n |a_i x_i|^2)^{1/2}$ is given by

$$\left(\sum_{i=1}^{n} |a_i x_i|^2\right)^{1/2} = \sum_{j=1}^{\infty} \left(\sum_{i=1}^{n} |a_i u_j^*(x_i)|^2\right)^{1/2} u_j.$$
 (2)

We have previously given an example of a conditional Schauder basis for ℓ_1 such that every weaving with the unit vector basis for ℓ_1 is a Schauder basis. In particular, this gives an

example of two non-equivalent Schauder bases for a Banach space X such that every weaving is a Schauder basis for X. Our next result shows that this is impossible if both bases are unconditional. Recall that two Schauder basic sequences $(x_j)_{j=1}^{\infty}$ and $(y_j)_{j=1}^{\infty}$ are said to be equivalent if there are constants c, C > 0 such that $c \| \sum a_j x_j \| \le \| \sum a_j y_j \| \le C \| \sum a_j x_j \|$ for all $(a_j)_{j=1}^{\infty} \in c_{00}$.

Theorem 4.2. Let $(x_i^0)_{i=1}^{\infty}$ and $(x_i^1)_{i=1}^{\infty}$ be two semi-normalized unconditional bases for a Banach space X such that for all $\sigma \in \{0,1\}^{\mathbb{N}}$, $(x_i^{\sigma(i)})_{i=1}^{\infty}$ is a basic sequence. Then, $(x_i^0)_{i=1}^{\infty}$ and $(x_i^1)_{i=1}^{\infty}$ are equivalent.

Proof. After renorming, we may assume that $(x_i^1)_{i=1}^{\infty}$ is 1-unconditional. Let X have the Banach lattice structure defined by the 1-unconditional basis $(x_i^1)_{i=1}^{\infty}$. Let D be the constant given in Theorem 4.1. As every weaving of $(x_i^0)_{i=1}^{\infty}$ and $(x_i^1)_{i=1}^{\infty}$ is a basic sequence, there is a constant C > 0 such that $|x_i^{1*}(x_i^0)| \geq C$ for all $i \in \mathbb{N}$. For any $(a_i)_{i=1}^{\infty} \in c_{00}$, we have the following estimate:

$$\left\| \sum_{i=1}^{\infty} a_i x_i^0 \right\| \ge D^{-1} \left\| \left(\sum_{i=1}^{\infty} |a_i x_i^0|^2 \right)^{1/2} \right\| \quad \text{by Theorem 4.1,}$$

$$= D^{-1} \left\| \sum_{j=1}^{\infty} \left(\sum_{i=1}^{\infty} |a_i x_j^{1*}(x_i^0)|^2 \right)^{1/2} x_j^1 \right\| \quad \text{by (2),}$$

$$\ge CD^{-1} \left\| \sum_{j=1}^{\infty} a_j x_j^1 \right\| \quad \text{as } (x_j^1) \text{ is 1-unconditional and } |x_j^{1*}(x_j^0)| \ge C \text{ for all } j \in \mathbb{N}.$$

Therefore, we have that the basis $(x_i^0)_{i=1}^{\infty}$ dominates the basis $(x^1)_{i=1}^{\infty}$, and by the same argument $(x_i^1)_{i=1}^{\infty}$ dominates $(x^0)_{i=1}^{\infty}$. Hence, $(x_i^0)_{i=1}^{\infty}$ and $(x^1)_{i=1}^{\infty}$ are equivalent.

Note that in proving Theorem 4.2 it was only required that there exists C > 0 such that $|x_j^{0*}(x_j^1)| \ge C$ and $|x_j^{1*}(x_j^0)| \ge C$ for $j \in \mathbb{N}$ which is a much weaker property than $(x_i^0)_{i=1}^{\infty}$ and $(x_i^1)_{i=1}^{\infty}$ are woven.

The following lemmas will be used in characterizing when every weaving of two unconditional Schauder bases is also an unconditional Schauder basis.

Lemma 4.3. Suppose X is a Banach space having subspaces Y and Z. Let $P: X \to Y$ be a projection from X onto Y and assume $P|_Z: Z \to Y$ is invertible. Then $Q = (P|_Z)^{-1}P$ is a projection onto Z.

Proof. The fact that Q is onto Z is clear. We check Q is a projection:

$$Q^2 = (P|_Z)^{-1} (P(P|_Z)^{-1} P) = (P|_Z)^{-1} P = Q$$

since if $x \in X$, then $Px \in Y$ and so $P(P|_Z)^{-1}Px = Px$.

For a Banach space X with subspaces $X_0, X_1 \subseteq X$, the distance between X_0 and X_1 measures how close a point in the sphere of one space can be to a point in the other space. That is,

$$d(X_0, X_1) = \inf\{\|x - y\| : x \in S_{X_i}, y \in X_{1-j} \text{ and } j \in \{0, 1\}.\}$$

Lemma 4.4. Suppose X is a Banach space and $X = X_1 \oplus X_2 = Y_1 \oplus Y_2$ for some subspaces X_1, X_2, Y_1, Y_2 of X. Assume P is a projection onto X_1 with (I - P) a projection onto X_2 , and Q is a projection on Y_1 with (I-Q) projecting onto Y_2 . Assume $Q|_{X_1}:X_1\to Y_1$ and $(I-P)|_{Y_2}: Y_2 \to X_2 \text{ are invertible and } d(X_1,Y_2) > 0. \text{ Then } X = X_1 \oplus Y_2.$

Proof. By Lemma 4.3, we have that $(Q|_{X_1})^{-1}Q$ is a projection onto X_1 and $((I-P)|_{Y_2})^{-1}(I-P)$ is a projection onto Y_2 . Furthermore, both

$$(Q|_{X_1})^{-1}Q((I-P)|_{Y_2})^{-1}(I-P)=0$$
 and $((I-P)|_{Y_2})^{-1}(I-P)(Q|_{X_1})^{-1}Q=0.$

Thus, the operator $R: X \to X$ defined by

$$R := (Q|_{X_1})^{-1}Q + ((I - P)|_{Y_2})^{-1}(I - P)$$

is a projection from X onto $X_1 \oplus Y_2$. We will show that the projection R is one-to-one (and hence the identity map) to prove that $X = X_1 \oplus Y_2$. Let $x \in X$ be so that Rx = 0. Then

$$||Rx|| = ||(Q|_{X_1})^{-1}Qx + ((I-P)|_{Y_2})^{-1}(I-P)x|| = 0$$

and since $d(X_1, Y_2) > 0$, it follows that

$$(Q|_{X_1})^{-1}Qx = 0$$
 and $((I-P)|_{Y_2})^{-1}(I-P)x = 0.$

Thus, Qx = 0 = (I - P)x so that x = Px = (I - Q)x. However, since $(Q|_{X_1})^{-1}Q$ is a projection onto X_1 , we have $Px = (Q|_{X_1})^{-1}QPx = (Q|_{X_1})^{-1}Qx = 0$ and therefore x = 0.

Given a sequence $(x_i)_{i\in\mathbb{N}}$ in a Banach space X and a set $\Gamma\subseteq\mathbb{N}$, we let $[x_i]_{i\in\Gamma}$ denote the subspace $\overline{\operatorname{span}}\{x_i\}_{i\in\Gamma}$.

Theorem 4.5. Let $(x_i^0)_{i=1}^{\infty}$ and $(x_i^1)_{i=1}^{\infty}$ be two C-unconditional bases for a Banach space X with biorthogonal functionals $(x_i^{0*})_{i=1}^{\infty}$ and $(x_i^{1*})_{i=1}^{\infty}$, respectively. The following are equivalent:

- (i) For every $\sigma \in \{0,1\}^{\mathbb{N}}$, $(x_i^{\sigma(i)})_{i=1}^{\infty}$ is an unconditional basis for X.
- (ii) There is a K > 0 so that for any $\sigma \in \{0,1\}^{\mathbb{N}}$, $(x_i^{\sigma(i)})_{i=1}^{\infty}$ is a K-unconditional basis for X.
- (iii) For every $\sigma \in \{0,1\}^{\mathbb{N}}$, $(x_i^{\sigma(i)}, x_i^{\sigma(i)*})_{i=1}^{\infty}$ is an unconditional approximate Schauder frame. (iv) For every $\sigma \in \{0,1\}^{\mathbb{N}}$, $(x_i^{\sigma(i)})_{i=1}^{\infty}$ is an unconditional basic sequence.
- (v) There exists a D > 0 so that for every $\sigma \in \{0,1\}^{\mathbb{N}}$,

$$d([x_i^0]_{i \in \sigma^{-1}(0)}, [x_i^1]_{i \in \sigma^{-1}(1)}) \ge D^{-1}.$$

(vi) There exists E > 0 such that for any $\sigma \in \{0,1\}^{\mathbb{N}}$, if P is the basis projection of $(x_i^0)_{i=1}^{\infty}$ onto $[x_i^0]_{i\in\sigma^{-1}(0)}$ and Q is the basis projection of $(x_i^1)_{i=1}^{\infty}$ onto $[x_i^1]_{i\in\sigma^{-1}(0)}$ then

$$P|_{[x_i^1]_{i\in\sigma^{-1}(0)}}:[x_i^1]_{i\in\sigma^{-1}(0)}\rightarrow [x_i^0]_{i\in\sigma^{-1}(0)}\quad and\quad Q|_{[x_i^0]_{i\in\sigma^{-1}(0)}}:[x_i^0]_{i\in\sigma^{-1}(0)}\rightarrow [x_i^1]_{i\in\sigma^{-1}(0)}$$

are invertible with inverses having norm at most E.

Proof. Note that (i) \Rightarrow (ii) follows from Proposition 3.1, and that (ii) \Rightarrow (iv) is immediate. We will prove (iv) \Rightarrow (v) \Rightarrow (vi) \Rightarrow (i) and (iii) \Rightarrow (vi) \Rightarrow (iii).

The case (iv) \Rightarrow (v) is well known, but we include a short proof here for completeness. By Proposition 3.1 there exists K > 0 such that $(x_i^{\sigma(i)})_{i \in \mathbb{N}}$ is K-suppression unconditional for all $\sigma \in \{0,1\}^{\mathbb{N}}$. Let $\sigma \in \{0,1\}^{\mathbb{N}}$, $x \in [x_i^0]_{i \in \sigma^{-1}(0)}$, and $y \in [x_i^1]_{i \in \sigma^{-1}(1)}$. We have that $||x-y|| \ge \frac{1}{K}||x||$, and hence $d([x_i^0]_{i \in \sigma^{-1}(0)}, [x_i^1]_{i \in \sigma^{-1}(1)}) \ge \frac{1}{K}$.

For (v) \Rightarrow (vi), we assume that (v) holds for some D > 0. Let $\sigma \in \{0,1\}^{\mathbb{N}}$. For $\Gamma \subseteq \mathbb{N}$ we let P_{Γ} be the basis projection of $(x_i^0)_{i=1}^{\infty}$ onto $[x_i^0]_{i\in\Gamma}$. Note that this implies that $I - P_{\Gamma} = P_{\Gamma^c}$ is the basis projection of $(x_i^0)_{i=1}^{\infty}$ onto $[x_i^0]_{i\in\Gamma^c}$. Let $x \in [x_i^1]_{i\in\sigma^{-1}(0)}$ with ||x|| = 1. Then

$$||P_{\sigma^{-1}(0)}(x)|| = ||x - (x - P_{\sigma^{-1}(0)}(x))|| \ge d\left([x_i^1]_{i \in \sigma^{-1}(0)}, [x_i^0]_{i \in \sigma^{-1}(1)}\right) \ge D^{-1}.$$
 (3)

Thus, we have that $P_{\sigma^{-1}(0)}$ is an isomorphism of $[x_i^1]_{i\in\sigma^{-1}(0)}$ onto a closed subspace of $[x_i^0]_{\sigma^{-1}(0)}$ with inverse having norm at most E:=D. It remains to prove that $P_{\sigma^{-1}(0)}\left([x_i^1]_{\sigma^{-1}(0)}\right)$ is dense in $[x_i^0]_{i\in\sigma^{-1}(0)}$.

Let $y \in \operatorname{span}(x_i^0)_{i \in \sigma^{-1}(0)}$ with $\|y\| = 1$. As $(x_i^0)_{i=1}^\infty$ and $(x_i^1)_{i=1}^\infty$ are unconditional bases there exists $N \in \mathbb{N}$ such that $y \in \operatorname{span}_{1 \leq i \leq N} x_i^0$ and for all $\Gamma \subset (N, \infty)$ we have that $\|\sum_{i \in \Gamma} x_i^{1*}(y) x_i^1\| < \varepsilon$. For $n \geq N$, we let $\sigma_n \in \{0,1\}^{\mathbb{N}}$ be defined by $\sigma_n|_{[1,n)} = \sigma|_{[1,n)}$ and $\sigma_n(i) = 1$ for $i \geq n$. A simple induction argument gives that $(x_i^{\sigma_n(i)})_{i=1}^\infty$ is an unconditional basis for X as we have changed only finitely many of the coordinates of $(x_i^1)_{i=1}^\infty$ and $d([x_i^0]_{i \in \sigma_n^{-1}(0)}, [x_i^1]_{i \in \sigma_n^{-1}(1)}) > 0$. Furthermore, $P_{\sigma_n^{-1}(0)}$ is an isomorphism of $[x_i^1]_{\sigma_n^{-1}(0)}$ onto $[x_i^0]_{\sigma_n^{-1}(0)}$ because these spaces are finite dimensional and have the same dimension. Hence, for each $n \geq N$ there exists unique $z_n \in \operatorname{span}(x_i^1)_{i \in \sigma_n^{-1}(0)}$ such that $y = P_{\sigma_n^{-1}(0)}(z_n) = \sum_{i \in \sigma_n^{-1}(0)} x_i^{0*}(z_n) x_i^0$ and $\|z_n\| \leq D$ by (3). Therefore,

$$P_{\sigma^{-1}(0)}z_n - y = \sum_{i \in \sigma^{-1}(0)} x_i^{0*}(z_n)x_i^0 - \sum_{i \in \sigma_n^{-1}(0)} x_i^{0*}(z_n)x_i^0 = \sum_{i \in \sigma^{-1}(0)\cap(n,\infty)} x_i^{0*}(z_n)x_i^0.$$
 (4)

If $(P_{\sigma^{-1}(0)}z_n)_{n=1}^{\infty}$ has a sequence of convex combinations which converges to y then we can conclude that $y \in P_{\sigma^{-1}(0)}([x_i^1]_{i \in \sigma^{-1}(0)})$ as the set is closed and convex. We assume by contradiction that this is not the case. Since any convex combination of y is itself, there exists F > 0 such that for all $n \geq N$,

$$\left\| \sum_{i \in \sigma^{-1}(0) \cap (n,\infty)} \sum_{j=n}^{\infty} x_i^{0*}(a_j z_j) x_i^0 \right\| > 2F \text{ for all } (a_j)_{j=n}^{\infty} \in [0,1]^{\mathbb{N}} \text{ with } \sum_{j=n}^{\infty} a_j = 1.$$
 (5)

Let $\varepsilon_n \searrow 0$ such that $\sum_{n=1}^{\infty} n\varepsilon_n < 1$ and $\sum_{n=1}^{\infty} \varepsilon_n < F$. We inductively choose a subsequence $(M_n)_{n=-1}^{\infty}$ of $\mathbb N$ with $M_{-1}=1$ and $M_0=N+1$ and sequences $(y_n)_{n=1}^{\infty}, (w_n)_{n=0}^{\infty} \subseteq X$ such that

- (a) $\left\| \sum_{i \in \sigma^{-1}(0) \cap [M_{n+1}, \infty)} x_i^{0*}(z_{M_j}) x_i^0 \right\| < \varepsilon_{n+1} \text{ for all } 1 \le j \le n \text{ and } n \in \mathbb{N}.$
- (b) $y_n := \sum_{i \in \sigma^{-1}(0) \cap [M_n, M_{n+1})} x_i^{0*}(z_{M_n}) x_i^0$ for all $n \in \mathbb{N}$ and $\|\sum_{j \in \mathbb{N}} a_j y_j\| > F$ for all $(a_j)_{j \in \mathbb{N}} \in [0, 1]^{\mathbb{N}}$ with $\sum_{j \in \mathbb{N}} a_j = 1$.

(c)
$$w_n := \lim_{i \to \infty} \sum_{i \in \sigma^{-1}(0) \cap [M_{n-1}, M_n)} x_i^{1*}(z_{M_i}) x_i^1$$
 for all $n \ge 0$.

(d)
$$\|w_n - \sum_{i \in \sigma^{-1}(0) \cap [M_{n-1}, M_n)} x_i^{1*}(z_{M_j}) x_i^1 \| < \varepsilon_j/j \text{ for all } j > n \ge 0.$$

Note that (a) can be achieved because the series converges and we are only working with finitely many z_j at each step. We have that (b) follows from (5), (a), and $\sum_{n=1}^{\infty} \varepsilon_n < F$. The definition of w_n in (c) can be made because the sum is over the finite interval $[M_{n-1}, M_n)$ and the sequence $(z_{M_j})_{j=n+1}^{\infty}$ is bounded, so we may pass to a subsequence of $(M_j)_{j\in\mathbb{N}}$ in which the limit exists. Finally, since the limit exists by (c), we can pass to yet another subsequence of $(M_j)_{j\in\mathbb{N}}$ to obtain (d).

For all $n \in \mathbb{N}$, we set $v_n := \sum_{i \in \sigma^{-1}(0) \cap [M_{n-1}, M_n)} x_i^{1*}(z_{M_n}) x_i^1$ and obtain the following:

(e)
$$||z_{M_n} - (v_n + \sum_{j=0}^{n-1} w_j)|| < \varepsilon_n \text{ for all } n \in \mathbb{N}.$$

(f)
$$||P_{\sigma^{-1}(0)}z_{M_n} - (y + y_n)|| < \varepsilon_{n+1} \text{ for all } n \in \mathbb{N}.$$

We have that (e) follows from (d) and the fact that $z_{M_n} \in [x_j^1]_{j=1}^{M_n-1}$. We have that (f) follows from (a), (b), and (3).

Let $M \in \mathbb{N}$ and $\Omega = \sigma^{-1}(0) \cap \bigcup_{n=1}^{2M} [M_{2n-1}, M_{2n})$. Then

$$\frac{1}{2M} \sum_{n=1}^{2M} (-1)^n v_{2n} \in [x_i^1]_{i \in \Omega} \quad \text{and} \quad \frac{1}{2M} \sum_{n=1}^{2M} (-1)^n (P_{\sigma^{-1}(1)} z_{M_{2n}} + y_{2n}) \in [x_i^0]_{i \in \mathbb{N} \setminus \Omega}. \tag{6}$$

Since $(y_n)_{n=N}^{\infty}$ is a block sequence of $(x_i^0)_{i\in\sigma^{-1}(0)}$, it is also C-unconditional and therefore by (b) we have that

$$\left\| \frac{1}{2M} \sum_{n=1}^{2M} (-1)^n \left(P_{\sigma^{-1}(1)} z_{M_{2n}} + y_{2n} \right) \right\| \ge C^{-1} \left\| \frac{1}{2M} \sum_{n=1}^{2M} y_{2n} \right\| > C^{-1} F.$$
 (7)

As $(w_i)_{i=0}^{\infty}$ is a block sequence of $(x_i^1)_{i=1}^{\infty}$, we have for all $n \in \mathbb{N}$ and $\Gamma \subseteq \{0, 1, ..., n\}$ that

$$\left\| \sum_{i \in \Gamma} w_i \right\| \le C \left\| \sum_{i=1}^n w_i \right\| = C \left\| \lim_{j \to \infty} \sum_{i \in \sigma^{-1}(0) \cap [1, M_n)} x_i^{1*}(z_{M_j}) x_i^1 \right\| \le \limsup_{j \to \infty} C^2 \|z_{M_j}\| \le C^2 D. \quad (8)$$

Thus, we have the following estimate:

$$\left\| \frac{1}{2M} \sum_{n=1}^{2M} (-1)^n \left(P_{\sigma^{-1}(1)} z_{M_{2n}} + y_{2n} \right) - \frac{1}{2M} \sum_{n=1}^{2M} (-1)^n v_{2n} \right\|$$

$$= \frac{1}{2M} \left\| \sum_{n=1}^{2M} (-1)^n \left(P_{\sigma^{-1}(1)} z_{M_{2n}} + y_{2n} - v_{2n} \right) \right\|$$

$$\leq \frac{1}{2M} \left\| \sum_{n=1}^{2M} (-1)^n \left(y + y_{2n} + P_{\sigma^{-1}(1)} z_{M_{2n}} - \left(v_{2n} + \sum_{j=0}^{2n-1} w_j \right) \right) \right\| + \frac{1}{2M} \left\| \sum_{j=1}^{M} w_{4j-2} + w_{4j-1} \right\|$$
13

$$\leq \frac{1}{2M} \sum_{n=1}^{2M} \left\| y + y_{2n} + P_{\sigma^{-1}(1)} z_{M_{2n}} - (v_{2n} + \sum_{j=0}^{2n-1} w_j) \right\| + \frac{C^2 D}{2M} \quad \text{by (8)}$$

$$< \frac{1}{2M} \sum_{n=1}^{2M} \left\| P_{\sigma^{-1}(0)} z_{M_{2n}} + P_{\sigma^{-1}(1)} z_{M_{2n}} - z_{M_{2n}} \right\| + \frac{1}{2M} \sum_{n=1}^{2M} (\varepsilon_{2n} + \varepsilon_{2n+1}) + \frac{C^2 D}{2M} \quad \text{by (e) and (f)}$$

$$< \frac{1}{2M} + \frac{C^2 D}{2M}.$$

The above estimate together with (6) and (7) gives that $d([x_i^0]_{i\in\mathbb{N}\setminus\Omega}, [x_i^1]_{i\in\Omega}) < CF^{-1}(1 + C^2D)(2M)^{-1}$. This contradicts (v) by choosing M large enough. An identical proof shows the claim for Q.

For (vi) \Rightarrow (i), we assume that (vi) holds for some E > 0. Let $\sigma \in \{0,1\}^{\mathbb{N}}$. For $j \in \sigma^{-1}(0)$ we let $f_j := \left(Q|_{[x_i^0]_{i \in \sigma^{-1}(0)}}^{-1}Q\right)^*x_j^{0*}$ and for $j \in \sigma^{-1}(1)$ we let $f_j := \left((I-P)|_{[x_i^1]_{i \in \sigma^{-1}(1)}}^{-1}(I-P)\right)^*x_j^{1*}$.

We first prove that $(x_i^{\sigma(i)}, f_i)_{i=1}^{\infty}$ is a biorthogonal system. Let $j, k \in \sigma^{-1}(0)$, then

$$f_j(x_k^0) = \left(\left(Q|_{[x_i^0]_{i \in \sigma^{-1}(0)}}^{-1} Q \right)^* x_j^{0*} \right) (x_k^0) = x_j^{0*} \left(\left(Q|_{[x_i^0]_{i \in \sigma^{-1}(0)}}^{-1} Q \right) x_k^0 \right) = x_j^{0*} (x_k^0) = \delta_{j,k}$$

For $j \in \sigma^{-1}(0)$ and $k \in \sigma^{-1}(1)$,

$$f_j(x_k^1) = \left(\left(Q|_{[x_i^0]_{i \in \sigma^{-1}(0)}}^{-1} Q \right)^* x_j^{0*} \right) (x_k^1) = x_j^{0*} \left(\left(Q|_{[x_i^0]_{i \in \sigma^{-1}(0)}}^{-1} Q \right) x_k^1 \right) = x_j^{0*}(0) = 0$$

Thus, if $j \in \sigma^{-1}(0)$ and $k \in \mathbb{N}$, then $f_j(x_k^{\sigma(k)}) = \delta_{j,k}$. Likewise, if $j \in \sigma^{-1}(1)$ and $k \in \mathbb{N}$, then $f_j(x_k^{\sigma(k)}) = \delta_{j,k}$. Thus, $(f_i)_{i=1}^{\infty}$ are biorthogonal functionals to $(x_i^{\sigma(i)})_{i=1}^{\infty}$. Let $\Gamma \subseteq \mathbb{N}$ and $x \in X$.

$$\left\| \sum_{i \in \Gamma} f_{i}(x) x_{i}^{\sigma(i)} \right\| \leq \left\| \sum_{i \in \Gamma \cap \sigma^{-1}(0)} (Q|_{[x_{i}^{0}]_{i \in \sigma^{-1}(0)}}^{-1} Q)^{*} x_{i}^{0*}(x) x_{i}^{0} \right\| + \left\| \sum_{i \in \Gamma \cap \sigma^{-1}(1)} ((I - P)|_{[x_{i}^{0}]_{i \in \sigma^{-1}(0)}}^{-1} (I - P))^{*} x_{i}^{1*}(x) x_{i}^{1} \right\|$$

$$\leq C \left\| \sum_{i \in \mathbb{N}} x_{i}^{0*} (Q|_{[x_{i}^{0}]_{i \in \sigma^{-1}(0)}}^{-1} Qx) x_{i}^{0} \right\| + C \left\| \sum_{i \in \mathbb{N}} x_{i}^{1*} ((I - P)|_{[x_{i}^{0}]_{i \in \sigma^{-1}(0)}}^{-1} (I - P)x) x_{i}^{1} \right\|$$

$$= C \left\| Q|_{[x_{i}^{0}]_{i \in \sigma^{-1}(0)}}^{-1} Qx \right\| + C \left\| (I - P)|_{[x_{i}^{0}]_{i \in \sigma^{-1}(0)}}^{-1} (I - P)x \right\|$$

$$\leq C \left\| Q|_{[x_{i}^{0}]_{i \in \sigma^{-1}(0)}}^{-1} Q \right\| \|x\| + C \left\| (I - P)|_{[x_{i}^{0}]_{i \in \sigma^{-1}(0)}}^{-1} (I - P) \right\| \|x\|$$

$$\leq 2CE \|x\|$$

Thus, $(x^{\sigma(i)})_{i=1}^{\infty}$ is an unconditional basic sequence. By Lemma 4.4, we have that $[x_i^{\sigma(i)}]_{i\in\mathbb{N}}=X$ and hence $(x^{\sigma(i)})_{i=1}^{\infty}$ is an unconditional basis of X.

For (vi) \Rightarrow (iii), we assume that (vi) holds for some E > 0. Let $\sigma \in \{0,1\}^{\mathbb{N}}$. We let P be the basis projection of $(x_i^0)_{i=1}^{\infty}$ onto $X_1 := [x_i^0]_{i \in \sigma^{-1}(0)}$ and Q be the basis projection of $(x_i^1)_{i=1}^{\infty}$ onto $Y_1 := [x_i^1]_{i \in \sigma^{-1}(0)}$. Note that (I - P) is the basis projection of $(x_i^0)_{i=1}^{\infty}$ onto $X_2 := [x_i^0]_{i \in \sigma^{-1}(1)}$ and (I - Q) is the basis projection of $(x_i^1)_{i=1}^{\infty}$ onto $Y_2 := [x_i^1]_{i \in \sigma^{-1}(1)}$. We have that $P|_{Y_1}, Q|_{X_1}$,

 $(I-P)|_{Y_2}$, and $(I-Q)|_{Y_1}$ are all invertible with bounded inverses. We let S be the frame operator of $(x_i^{\sigma(i)}, x_i^{\sigma(i)*})_{i=1}^{\infty}$. Then for all $x \in X$, we have that

$$S(x) = \sum_{i=1}^{\infty} x_i^{\sigma(i)*}(x) x_i^{\sigma(i)} = \sum_{i \in \sigma^{-1}(0)} x_i^0(x) x_i^0 + \sum_{i \in \sigma^{-1}(1)} x_i^1(x) x_i^1 = P(x) + (I - Q)(x).$$

Thus, S=P+(I-Q) is the frame operator of $(x_i^{\sigma(i)},x_i^{\sigma(i)*})_{i=1}^{\infty}$, and hence is well defined and bounded. We consider the operator $T=P|_{Y_1}^{-1}Q|_{X_1}^{-1}Q+(I-Q)|_{X_2}^{-1}(I-P)|_{Y_2}^{-1}(I-P)$ and will prove that T is the inverse of S and hence $(x_i^{\sigma(i)},x_i^{\sigma(i)*})_{i=1}^{\infty}$ would be an approximate Schauder frame. We have that

$$ST = (P + (I - Q))P|_{Y_1}^{-1}Q|_{X_1}^{-1}Q + (P + (I - Q))(I - Q)|_{X_2}^{-1}(I - P)|_{Y_2}^{-1}(I - P)$$

$$= PP|_{Y_1}^{-1}Q|_{X_1}^{-1}Q + (I - Q)(I - Q)|_{X_2}^{-1}(I - P)|_{Y_2}^{-1}(I - P)$$

$$= Q|_{X_1}^{-1}Q + (I - P)|_{Y_2}^{-1}(I - P) = I.$$

where the second equation follows from $(I-Q)P|_{Y_1}^{-1}=0=P(I-Q)|_{X_2}^{-1}$ and the last equation by Lemma 4.4. Thus, ST=I. To show TS=I, we have

$$TS = P|_{Y_1}^{-1}Q_{X_1}^{-1}Q(P + (I - Q)) + (I - Q)|_{X_2}^{-1}(I - P)|_{Y_2}^{-1}(I - P)(P + (I - Q))$$

$$= P|_{Y_1}^{-1}Q_{X_1}^{-1}QP + (I - Q)|_{X_2}^{-1}(I - P)|_{Y_2}^{-1}(I - P)(I - Q)$$

$$= P^{-1}|_{Y_1}P + (I - Q)|_{X_2}^{-1}(I - Q) = I$$

where the last equation follows from Lemma 4.4. Hence, we have that T is the inverse of S and $(x_i^{\sigma(i)}, x_i^{\sigma(i)*})_{i=1}^{\infty}$ is an approximate Schauder frame. We now check that $(x_i^{\sigma(i)}, x_i^{\sigma(i)*})_{i=1}^{\infty}$ is an unconditional approximate Schauder frame. Let $x \in X$ and $\Gamma \subseteq \mathbb{N}$. We have that

$$\left\| \sum_{i \in \Gamma} x_i^{\sigma(i)*}(x) x_i^{\sigma(i)} \right\| \le \left\| \sum_{i \in \Gamma \cap \sigma^{-1}(0)} x_i^{0*}(x) x_i^{0} \right\| + \left\| \sum_{i \in \Gamma \cap \sigma^{-1}(1)} x_i^{1*}(x) x_i^{1} \right\|$$

$$\le C \left\| \sum_{i \in \mathbb{N}} x_i^{0*}(x) x_i^{0} \right\| + C \left\| \sum_{i \in \mathbb{N}} x_i^{1*}(x) x_i^{1} \right\|$$

$$= 2C \|x\|.$$

Thus, $(x_i^{\sigma(i)}, x_i^{\sigma(i)*})_{i=1}^{\infty}$ is a 2C-unconditional approximate Schauder frame.

For (iii) \Rightarrow (vi), we assume that for every $\phi \in \{0,1\}^{\mathbb{N}}$, we have that $(x_i^{\phi(i)}, x_i^{\phi(i)*})_{i=1}^{\infty}$ is an unconditional approximate Schauder frame. By Theorem 2.4 there exists $E \geq 1$ such that for every $\phi \in \{0,1\}^{\mathbb{N}}$, we have that $(x_i^{\phi(i)}, x_i^{\phi(i)*})_{i=1}^{\infty}$ is a E-unconditional approximate Schauder frame whose frame operator S satisfies ||S||, $||S^{-1}|| \leq E$. Let $\sigma \in \{0,1\}^{\mathbb{N}}$ and $y \in [x_i^0]_{i \in \sigma^{-1}(0)}$. We let P be the basis projection of $(x_i^0)_{i=1}^{\infty}$ onto $[x_i^0]_{i \in \sigma^{-1}(0)}$ and Q be the basis projection of $(x_i^1)_{i=1}^{\infty}$ onto $[x_i^1]_{i \in \sigma^{-1}(0)}$. Thus, the frame operator of $(x_i^{\sigma(i)}, x_i^{\sigma(i)*})_{i=1}^{\infty}$ is given by S = P + (I - Q).

There exists unique $x \in X$ with $||x|| \le E||y||$ such that S(x) = y. We have that

$$y = S(x) = (P + (I - Q))(Q + (I - Q))x = PQx + P(I - Q)x + (I - Q)x$$

For the sake of contradiction, we assume that $(I-Q)x \neq 0$. As, $y \in [x_i^0]_{i \in \sigma^{-1}(0)}$, we have that (I-P)y = 0. Hence, 0 = (I-P)y = (I-P)(I-Q)x. Thus, 0 = (Q+(I-P))(I-Q)x and hence the operator (Q+(I-P)) is not invertible. This is a contradiction because (Q+(I-P)) is the frame operator of $(x_i^{1-\sigma(i)}, x_i^{(1-\sigma(i))*})_{i=1}^{\infty}$. Thus, (I-Q)x = 0 and $x = Q(x) \in [x_i^1]_{i \in \sigma^{-1}(0)}$. This implies $P|_{[x_i^1]_{i \in \sigma^{-1}(0)}} : [x_i^1]_{i \in \sigma^{-1}(0)} \to [x_i^0]_{i \in \sigma^{-1}(0)}$ is invertible and $\|P|_{[x_i^0]_{i \in \sigma^{-1}(0)}}^{-1}\| \leq E$. Likewise, $Q|_{[x_i^0]_{i \in \sigma^{-1}(0)}} : [x_i^0]_{i \in \sigma^{-1}(0)} \to [x_i^1]_{i \in \sigma^{-1}(0)}$ is invertible and $\|Q|_{[x_i^0]_{i \in \sigma^{-1}(0)}}^{-1}\| \leq E$.

5. Perturbations

Perturbation theorems are powerful tools in both creating new coordinate systems and in showing that coordinate systems are resilient to error. We first recall the following classical perturbation theorem for Schauder bases which is often referred to as the small perturbation lemma (see Theorem 6.18 in [FHHMPZ]).

Theorem 5.1. Let $(x_j^0)_{j=1}^{\infty}$ be a Schauder basis for a Banach space X with biorthogonal functionals $(x_j^{0*})_{j=1}^{\infty}$. If $(x_j^1)_{j=1}^{\infty}$ is a sequence in X such that $\sum_{j=1}^{\infty} \|x_j^0 - x_j^1\| \|x_j^{0*}\| < 1$ then $(x_j^1)_{j=1}^{\infty}$ is a Schauder basis for X and is equivalent to $(x_j^0)_{j=1}^{\infty}$.

Note that if $(x_j^1)_{j=1}^{\infty}$ is a perturbation of $(x_j^0)_{j=1}^{\infty}$ in the sense of Theorem 5.1, then every weaving is also a perturbation of $(x_j^0)_{j=1}^{\infty}$ and hence $(x_j^1)_{j=1}^{\infty}$ and $(x_j^0)_{j=1}^{\infty}$ are woven. In other words, a perturbation of a Schauder basis is a Schauder basis which is woven with it. Perturbation theorems have been considered for Hilbert space frames and Schauder frames as well [BCGLL, CLZ]. However, we note that perturbation theorems are particularly natural in the context of approximate Schauder frames because when a Schauder frame is perturbed, one might expect the frame operator to be perturbed as well. Hence, the perturbation of a Schauder frame would be expected to be an approximate Schauder frame, and indeed the name "approximate Schauder frame" was chosen for this very reason. We prove two perturbation theorems for approximate Schauder frames and prove they result in woven approximate Schauder frames.

Theorem 5.2. Suppose $(x_i, f_i)_{i=1}^{\infty}$ is a C-suppression unconditional approximate Schauder frame for X with frame operator S. If T is a bounded operator satisfying

$$||Id - T|| < C^{-1},$$

then $(Tx_i, f_i)_{i=1}^{\infty}$ is an unconditional approximate Schauder frame for X and is woven with $(x_i, f_i)_{i=1}^{\infty}$.

Proof. Set $(x_i^0, f_i^0)_{i=1}^{\infty} = (x_i, f_i)_{i=1}^{\infty}$ and $(x_i^1, f_i^1)_{i=1}^{\infty} = (Tx_i, f_i)_{i=1}^{\infty}$ for ease of notation. Let $x \in X$ and $\sigma \in \{0, 1\}^{\mathbb{N}}$. We will first prove that the series $\sum_{i=1}^{\infty} f_i^{\sigma(i)}(x) x_i^{\sigma(i)}$ is unconditionally Cauchy and hence the frame operator of $(x_i^{\sigma(i)}, f_i^{\sigma(i)})_{i=1}^{\infty}$ is well defined. Let $\varepsilon > 0$ and choose $N \in \mathbb{N}$ so

that for all $J \subseteq \mathbb{N}$ with $\min(J) \geq N$, we have that $(1 + ||T||) ||\sum_{i \in J} f_i(x) x_i|| < \varepsilon$. For a fixed $J \subseteq \mathbb{N}$ with $\min(J) \geq N$, we have that

$$\left\| \sum_{i \in J} f_i^{\sigma(i)}(x) x_i^{\sigma(i)} \right\| = \left\| \sum_{i \in J \cap \sigma^{-1}(0)} f_i(x) x_i + \sum_{i \in J \cap \sigma^{-1}(1)} f_i(x) T(x_i) \right\|$$

$$\leq \left\| \sum_{i \in J \cap \sigma^{-1}(0)} f_i(x) x_i \right\| + \left\| \sum_{i \in J \cap \sigma^{-1}(1)} f_i(x) x_i \right\| \|T\|$$

$$< \varepsilon.$$

Hence, the frame operator of $(x_i^{\sigma(i)}, f_i^{\sigma(i)})_{i=1}^{\infty}$ is well defined and converges unconditionally.

Let $\sigma \in \{0,1\}^{\mathbb{N}}$ and let S_{σ} be the frame operator of $(x_i^{\sigma(i)}, f_i^{\sigma(i)})_{i=1}^{\infty}$. We will prove that $||Id - S_{\sigma}S^{-1}|| < 1$, hence showing that S_{σ} is bounded with bounded inverse. If $x \in X$ and $y = S^{-1}x$, then

$$\begin{aligned} \|(Id - S_{\sigma}S^{-1})x\| &= \|Sy - S_{\sigma}y\| \\ &= \left\| \sum_{i=1}^{\infty} f_{i}^{0}(y)x_{i}^{0} - \sum_{i=1}^{\infty} f_{i}^{\sigma(i)}(y)x_{i}^{\sigma(i)} \right\| \\ &= \left\| \sum_{i \in \sigma^{-1}(1)} \left[f_{i}(y)x_{i} - f_{i}(y)T(x_{i}) \right] \right\| \\ &\leq \left\| \sum_{i \in \sigma^{-1}(1)} f_{i}(y)x_{i} \right\| \|Id - T\| \\ &\leq C\|Sy\| \|Id - T\| \\ &= C\|x\| \|Id - T\|. \end{aligned}$$

Thus, $||Id - S_{\sigma}S^{-1}|| \leq C||Id - T|| < 1$, and hence S_{σ} is bounded with bounded inverse.

Note that Theorem 5.2 is only stated for the case that $(x_i, f_i)_{i=1}^{\infty}$ is an unconditional approximate Schauder frame. It is natural to ask if the same conclusion holds for conditional approximate Schauder frames as well. This is not possible because if $(x_i, f_i)_{i=1}^{\infty}$ is a conditional approximate Schauder frame and α is any scalar other than 1 then $(x_i, f_i)_{i=1}^{\infty}$ is not woven with $(\alpha x_i, f_i)_{i=1}^{\infty}$. However, the following perturbation result holds for both unconditional and conditional approximate Schauder frames.

Theorem 5.3. If $(x_i^0, f_i^0)_{i=1}^{\infty}$ is an approximate Schauder frame for X with frame operator S and $(x_i^1, f_i^1)_{i=1}^{\infty}$ is a sequence in $X \times X^*$ satisfying

$$\sum_{i=1}^{\infty} (\|f_i^0 - f_i^1\| \|x_i^0\| + \|x_i^0 - x_i^1\| \|f_i^1\|) < \|S^{-1}\|^{-1}$$

then $(x_i^1, f_i^1)_{i=1}^{\infty}$ is an approximate Schauder frame for X that is woven with $(x_i^0, f_i^0)_{i=1}^{\infty}$.

Proof. Let $\sigma \in \{0,1\}^{\mathbb{N}}$ and $x \in X$. We first show that the series $\sum_{i=1}^{\infty} f_i^{\sigma(i)}(x) x_i^{\sigma(i)}$ is Cauchy and hence the frame operator of $(x_i^{\sigma(i)}, f_i^{\sigma(i)})_{i=1}^{\infty}$ is well defined. Let $\varepsilon > 0$ and choose $N \in \mathbb{N}$ so that for all $m > n \ge N$,

$$\left\| \sum_{i=n}^{m} f_i^0(x) x_i^0 \right\| + \sum_{i=n}^{m} (\|f_i^0 - f_i^1\| \|x_i^0\| + \|x_i^0 - x_i^1\| \|f_i^1\|) \|x\| < \epsilon.$$

Thus.

$$\left\| \sum_{i=n}^{m} f_{i}^{\sigma(i)}(x) x_{i}^{\sigma(i)} \right\| \leq \left\| \sum_{i=n}^{m} f_{i}^{0}(x) x_{i}^{0} \right\| + \left\| \sum_{i \in [n,m] \cap \sigma^{-1}(1)} \left[f_{i}^{0}(x) x_{i}^{0} - f_{i}^{1}(x) x_{i}^{1} \right] \right\|$$

$$\leq \left\| \sum_{i=n}^{m} f_{i}^{0}(x) x_{i}^{0} \right\| + \sum_{i=n}^{m} \left[\left\| f_{i}^{0}(y) x_{i}^{0} - f_{i}^{1}(y) x_{i}^{0} \right\| + \left\| f_{i}^{1}(x) x_{i}^{0} - f_{i}^{1}(x) x_{i}^{1} \right\| \right]$$

$$\leq \left\| \sum_{i=n}^{m} f_{i}^{0}(x) x_{i}^{0} \right\| + \sum_{i=n}^{m} (\left\| f_{i}^{0} - f_{i}^{1} \right\| \left\| x_{i}^{0} \right\| + \left\| x_{i}^{0} - x_{i}^{1} \right\| \left\| f_{i}^{1} \right\|) \|x\|$$

$$\leq \epsilon.$$

Hence, the frame operator of $(x_i^{\sigma(i)}, f_i^{\sigma(i)})_{i=1}^{\infty}$ is well defined. Let $\sigma \in \{0,1\}^{\mathbb{N}}$ and let T be the frame operator of $(x_i^{\sigma(i)}, f_i^{\sigma(i)})_{i=1}^{\infty}$. We will prove that $\|Id - TS^{-1}\| < 1$, hence showing that T is bounded with bounded inverse. If $x \in X$ and $y = S^{-1}x$, then

$$\begin{aligned} & \left\| (Id - TS^{-1})x \right\| = \left\| Sy - Ty \right\| \\ & = \left\| \sum_{i=1}^{\infty} f_i^0(y) x_i^0 - \sum_{i=1}^{\infty} f_i^{\sigma(i)}(y) x_i^{\sigma(i)} \right\| \\ & = \left\| \sum_{i \in \sigma^{-1}(1)} \left[f_i^0(y) x_i^0 - f_i^1(y) x_i^1 \right] \right\| \\ & \leq \sum_{i=1}^{\infty} \left\| f_i^0(y) x_i^0 - f_i^1(y) x_i^1 \right\| \\ & \leq \sum_{i=1}^{\infty} \left\| f_i^0(y) x_i^0 - f_i^1(y) x_i^0 \right\| + \left\| f_i^1(y) x_i^0 - f_i^1(y) x_i^1 \right\| \\ & \leq \left(\sum_{i=1}^{\infty} \left\| f_i^0 - f_i^1 \right\| \left\| x_i^0 \right\| + \left\| x_i^0 - x_i^1 \right\| \left\| f_i^1 \right\| \right) \left\| y \right\| \end{aligned}$$

$$\leq \left(\sum_{i=1}^{\infty} \|f_i^0 - f_i^1\| \|x_i^0\| + \|x_i^0 - x_i^1\| \|f_i^1\| \right) \|S^{-1}\| \|x\|.$$

Thus, $||Id - TS^{-1}|| \leq (\sum_{i=1}^{\infty} ||f_i^0 - f_i^1|| ||x_i^0|| + ||x_i^0 - x_i^1|| ||f_i^1||) ||S^{-1}|| < 1$ proving that T is bounded with bounded inverse.

References

- [BCGLL] T. Bemrose, P. G. Casazza, K. Gröchenig, M. C. Lammers, and R. G. Lynch, Weaving frames, submitted, see arXiv:1503.03947.
- [CDOSZ] P. G. Casazza, S. J. Dilworth, E. Odell, Th. Schlumprecht, and A. Zsak, Coefficient Quantization for Frames in Banach Spaces, J. Math. Anal. Appl. 348 (2008), 66–86.
- [CHL] P. G. Casazza, D. Han, and D. Larson, Frames for Banach spaces, Cont. Math., 247 (1999), 149-182.
- [CK] P.G. CASAZZA AND G. KUTYNIOK, Finite frames: Theory and applications, Springer (2013).
- [CL] P. G. Casazza and R. G. Lynch, Weaving properties of Hilbert space frames, Proceedings of SampTA (2015).
- [CL2] P. G. Casazza and R. G. Lynch, A brief introduction to Hilbert space frame theory and its applications, submitted.
- [CLZ] D.Y. Chen, L. Li, and B.T. Zheng, Perturbations of frames, Acta Math. Sinica, English Series, 30 (2014), no. 7, 1089–1108.
- [C] O. Christensen, An Introduction to Frames and Riesz Bases, Birkhäuser, Boston (2003).
- [DV] Deepshikha and L.K. Vashisht, On weaving frames, preprint.
- [FHHMPZ] M. Fabian, P. Habala, P. Hjek, V. Montesinos Santaluca, J. Pelant, and V. Zizler, Functional Analysis and Infinite-Dimensional Geometry, CMS Books Math./Ouvrages Math. SMC, vol. 8 Springer, New York (2001).
- [FOSZ] D. Freeman, E. Odell, Th. Schlumprecht, and A. Zsak, Unconditional structures of translates for $L_p(\mathbb{R}^d)$. Israel J. Math., 203 (2014), no. 1, 189–209.
- [FG] H. G. Feichtinger and K. Gröchenig, A unified approach to atomic decompositions via integrable group representations, In: Proc. Conf. Function Spaces and Applications, Lecture Notes Math. 1302. BerlinHeidelberg New York: Springer (1988), 52-73.
- [G] K. Gröchenig, Describing functions: Atomic decompositions versus frames, Monatshefte fur Math. 112 (1991), 1–41.
- [LT] J. Lindenstrauss and L. Tzafriri, Classical Banach spaces I and II, Springer, New York (1977).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA, MO 65211 *E-mail address*: casazzap@mail.missouri.edu

Department of Mathematics and Computer Science, Saint Louis University, St Louis, MO 63103

E-mail address: dfreema7@slu.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA, MO 65211

E-mail address: rglz82@mail.missouri.edu