arXiv:1609.08073v1 [math.PR] 26 Sep 2016

ON NON-POLYNOMIAL LOWER ERROR BOUNDS FOR ADAPTIVE STRONG APPROXIMATION OF SDES

LARISA YAROSLAVTSEVA

ABSTRACT. Recently, it has been shown in [9] that there exists a system of stochastic differential equations (SDE) on the time interval [0, T] with infinitely often differentiable and bounded coefficients such that the Euler scheme with equidistant time steps converges to the solution of this SDE at the final time in the strong sense but with no polynomial rate. Even worse, in [20] it has been shown that for any sequence $(a_n)_{n\in\mathbb{N}} \subset (0,\infty)$, which may converge to zero arbitrary slowly, there exists an SDE on [0,T] with infinitely often differentiable and bounded coefficients such that no approximation of the solution of this SDE at the final time based on n evaluations of the driving Brownian motion at fixed time points can achieve a smaller absolute mean error than the given number a_n . In the present article we generalize the latter result to the case when the approximations may choose the location as well as the number of the Brownian motion observed so far.

1. INTRODUCTION

Let $d, m \in \mathbb{N}, T \in (0, \infty)$, consider a *d*-dimensional system of autonomous stochastic differential equations (SDE)

(1)
$$dX(t) = \mu(X(t)) dt + \sigma(X(t)) dW(t), \quad t \in [0, T]$$
$$X(0) = x_0$$

with a deterministic initial value $x_0 \in \mathbb{R}^d$, a drift coefficient $\mu \colon \mathbb{R}^d \to \mathbb{R}^d$, a diffusion coefficient $\sigma \colon \mathbb{R}^d \to \mathbb{R}^{d \times m}$ and an *m*-dimensional driving Brownian motion W, and assume that (1) has a unique strong solution $(X(t))_{t \in [0,T]}$. Our computational task is to approximate X(T) by means of methods that use finitely many evaluations of the driving Brownian motion W. In particular we are interested in the following question: under which assumptions on the coefficients μ and σ exists a method of the latter type, which converges to X(T) in absolute mean with a polynomial rate?

It is well-known that if the coefficients μ and σ are globally Lipschitz continuous then the classical Euler scheme achieves the rate of convergence 1/2, see [26]. Moreover, the recent literature on numerical approximation of SDEs contains a number of results on approximation schemes that are specifically designed for non-Lipschitz coefficients and achieve polynomial convergence rates for suitable classes of such SDEs, see e.g. [16, 12, 18, 25, 38, 35, 37, 3, 21, 4] for SDEs with globally monotone coefficients and see e.g. [2, 8, 5, 1, 32, 17, 19, 23, 24, 33, 11] for SDEs with possibly non-monotone coefficients.

On the other hand, it has recently been shown in [20] that for any sequence $(a_n)_{n \in \mathbb{N}} \subset (0, \infty)$, which may converge to zero arbitrary slowly, there exists an SDE (1) with d = 4 and m = 1 and

with infinitely often differentiable and bounded coefficients μ and σ such that no approximation of X(T) based on finitely many evaluations of the driving Brownian motion W converges in absolute mean faster than the given sequence $(a_n)_{n \in \mathbb{N}}$. More formally,

(2)
$$\inf_{\substack{s_1,\ldots,s_n \in [0,T] \\ \text{measurable}}} \inf_{\substack{u \colon \mathbb{R}^n \to \mathbb{R}^4 \\ \text{measurable}}} \mathbb{E} \left\| X(T) - u \big(W(s_1),\ldots,W(s_n) \big) \right\| \ge a_n.$$

In particular, there exists an SDE (1) with infinitely often differentiable and bounded coefficients μ and σ such that its solution at the final time can not be approximated with a polynomial rate of convergence based on finitely many evaluations of the driving Brownian motion W. We add that the latter statement in the special case when the approximation is given by the Euler scheme with equidistant time steps has first been shown in [9].

Note that the time points $s_1, \ldots, s_n \in [0, T]$ that are used by an approximation $u(W(s_1), \ldots, W(s_n))$ in (2) are fixed, and therefore this negative result does not cover approximations that may choose the number as well as the location of the evaluation sites of the driving Brownian motion W in an adaptive way, e.g. numerical schemes that adjust the actual step size according to a criterion that is based on the values of the driving Brownian motion W observed so far, see e.g. [6, 29, 30, 27, 34, 22, 13, 14] and the references therein. See Section 4 for the formal definition of that type of approximations. It is well-known that for SDEs (1) with (essentially) globally Lipschitz continuous coefficients μ and σ adaptive approximations can not achieve a better rate of convergence compared to what is best possible for non-adaptive ones, which at the same time coincides with the best possible rate of convergence that can be achieved by any approximation based on $W(\frac{T}{n}), W(\frac{2T}{n}), \ldots, W(T)$, see [29, 30]. However, as has recently turned out, this is not necessarily the case anymore if the coefficients μ and σ are not both globally Lipschitz continuous. In [10] it has been shown that for the one-dimensional squared Bessel process, which is the solution of the SDE (1) with $d = m = \mu = 1$ and $\sigma(x) = 2\sqrt{|x|}$ for $x \in \mathbb{R}$ the following holds: the best possible rate of convergence that can be achieved by any approximation based on $W(\frac{T}{n}), W(\frac{2T}{n}), \ldots, W(T)$ equals 1/2, i.e. there exist $c_1, c_2 > 0$ such that

$$c_1 \cdot n^{-1/2} \le \inf_{\substack{u \colon \mathbb{R}^n \to \mathbb{R} \\ \text{measurable}}} \mathbb{E} \left| X(T) - u \left(W(\frac{T}{n}), W(\frac{2T}{n}), \dots, W(T) \right) \right| \le c_2 \cdot n^{-1/2},$$

while the best possible rate of convergence that can be achieved by approximations based on n adaptively chosen evaluations of the driving Brownian motion W equals infinity. More formally, for every $\alpha > 0$ there exists c > 0 and a sequence of approximations \hat{X}_n based on n adaptively chosen evaluations of W such that

$$\mathbb{E}|X(T) - \widehat{X}_n| \le c \cdot n^{-\alpha}.$$

In view of the latter result one might hope that a non-polynomial lower error bound a_n in (2) could be overcome by using adaptive approximations, see also the discussion in [7, p. 2]. In the present article we prove that the pessimistic alternative is true. We show that for any sequence $(a_n)_{n \in \mathbb{N}} \subset (0, \infty)$, which may converge to zero arbitrary slowly, there exists an SDE (1) with d = 4 and m = 1 and with infinitely often differentiable and bounded coefficients μ and σ such that no approximation based on n adaptively chosen evaluations of the driving Brownian motion

W on average can achieve a smaller absolute mean error than the given number a_n , i.e.

$$\mathbb{E}\left\|X(T) - \widehat{X}_n\right\| \ge a_n$$

for any approximation \widehat{X}_n of the latter type. This fact is an immediate consequence of Corollary 2 in Section 5 together with an appropriate scaling argument. For the proof of the latter result we employ the same class of SDEs as in [20]. Thus, roughly speaking, these SDEs can not be solved approximately in the strong sense in a reasonable computational time by means of any kind of adaptive (or nonadaptive) method based on finitely many evaluations of the driving Brownian motion W.

We conjecture that a similar negative result does even if one allows for adaptive approximations based on finitely many evaluations of arbitrary linear continuous functionals of the driving Brownian motion W. However, in this case one can not employ the class of SDEs from [20] since for every such SDE its solution at the final time can be approximated with error zero based on the evaluation of only two linear continuous functionals of the driving Brownian motion W, see (4)

We add that negative results in the spirit of (2) for quadrature problems for marginal distributions of SDEs have recently been established in [31].

We briefly describe the content of the paper. In Section 2 we fix some notation. In Section 3 we briefly introduce the class of SDEs from [20], which is studied in this article as well. In Section 4 we formally define the class of adaptive approximations, which are analysed in this article. Our lower error bounds are stated in Section 5. The proof of the main result, Theorem 1, is carried out in Section 6.

2. NOTATION

Throughout this article the following notation is used. For a set A, a vector space V, a set $B \subseteq V$, and a function $f: A \to B$ we put $\operatorname{supp}(f) = \{x \in A: f(x) \neq 0\}$. For sets A, B, a function $f: A \to B$ and a subset $E \subseteq A$ we denote by $f|_E$ the restriction of f to E. Moreover, for $d \in \mathbb{N}$ and $v \in \mathbb{R}^d$ we write ||v|| for the Euclidean norm of v. For $n \in \mathbb{N}$ and $-\infty < a < b < \infty$ we denote by $\mathfrak{B}(\mathbb{R}^n)$ and $\mathfrak{B}(C([a, b]))$ the Borel σ -fields on \mathbb{R}^n and on C([a, b]), respectively, where the latter space is equipped with the supremum norm. For S being a finite product of the latter two spaces we denote by $\mathfrak{B}(S)$ the Borel σ -field on S generated by the respective product topology.

3. A family of SDEs with smooth and bounded coefficients

Throughout this article we study SDEs provided by the following setting.

Let $T \in (0, \infty)$, let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space with a normal filtration $(\mathcal{F}_t)_{t \in [0,T]}$, and let $W: [0,T] \times \Omega \to \mathbb{R}$ be a standard $(\mathcal{F}_t)_{t \in [0,T]}$ -Brownian motion on $(\Omega, \mathcal{F}, \mathbb{P})$.

Let $0 < \tau_1 < \tau_2 < T$ and let $f, g, h \in C^{\infty}(\mathbb{R})$ be bounded and satisfy $\operatorname{supp}(f) \subseteq (-\infty, \tau_1]$, $\inf_{t \in [0, \tau_1/2]} |f'(t)| > 0$, $\operatorname{supp}(g) \subseteq [\tau_1, \tau_2]$, $g \neq 0$, $\operatorname{supp}(h) \subseteq [\tau_2, \infty)$, and $\int_{\tau_2}^T h(t) dt \neq 0$.

For every $\psi \in C^{\infty}(\mathbb{R})$ let $\mu^{\psi} \colon \mathbb{R}^4 \to \mathbb{R}^4$ and $\sigma \colon \mathbb{R}^4 \to \mathbb{R}^4$ be given by

$$\mu^{\psi}(x) = (1, 0, 0, h(x_1) \cdot \cos(x_2 \,\psi(x_3))),$$

$$\sigma(x) = (0, f(x_1), g(x_1), 0)$$

and consider the following 4-dimensional system of SDEs

(3)
$$dX^{\psi}(t) = \mu^{\psi}(X^{\psi}(t)) dt + \sigma(X^{\psi}(t)) dW(t), \quad t \in [0, T], \\ X^{\psi}(0) = 0.$$

Remark 1. Note that for every $\psi \in C^{\infty}(\mathbb{R})$ the functions μ^{ψ} and σ are infinitely often differentiable and bounded.

Remark 2. It is easy to see that for every $\psi \in C^{\infty}(\mathbb{R})$ the SDE (3) has a unique strong solution given by

(4)

$$X_{1}^{\psi}(t) = t, \qquad X_{2}^{\psi}(t) = \int_{0}^{\min(t,\tau_{1})} f(s) \, dW(s),$$

$$X_{3}^{\psi}(t) = \mathbf{1}_{[\tau_{1},T]}(t) \cdot \int_{\min(t,\tau_{1})}^{\min(t,\tau_{2})} g(s) \, dW(s),$$

$$X_{4}^{\psi}(t) = \mathbf{1}_{[\tau_{2},T]}(t) \cdot \cos\left(X_{2}^{\psi}(\tau_{1}) \, \psi\left(X_{3}^{\psi}(\tau_{2})\right)\right) \cdot \int_{\tau_{2}}^{t} h(s) \, ds$$

for all $t \in [0, T]$.

4. Adaptive strong approximations

Let $\delta \in (0, T]$. We study general strong approximations of $X^{\psi}(T)$ based on $(W(t))_{t \in [\delta, T]}$ and on finitely many sequential evaluations of W in the interval $(0, \delta)$. Every such approximation $\widehat{X} : \Omega \to \mathbb{R}^4$ is defined by three sequences

$$\varphi = (\varphi_n)_{n \in \mathbb{N}}, \quad \chi = (\chi_n)_{n \in \mathbb{N}}, \quad \phi = (\phi_n)_{n \in \mathbb{N}}$$

of measurable mappings

(5)

$$\begin{aligned} \varphi_n \colon \mathbb{R}^{n-1} \times \mathbb{R}^{[\delta,T]} \to (0,\delta), \\ \chi_n \colon \mathbb{R}^n \times \mathbb{R}^{[\delta,T]} \to \{0,1\}, \\ \phi_n \colon \mathbb{R}^n \times \mathbb{R}^{[\delta,T]} \to \mathbb{R}^4. \end{aligned}$$

The sequence φ determines the evaluation sites of a trajectory of W in the interval $(0, \delta)$. The total number of evaluations is determined by the sequence χ of stopping rules. Finally, the sequence ϕ is used to obtain the approximation to $X^{\psi}(T)$ from the observed data.

More precisely, let $\omega \in \Omega$, let $w = W(\omega)$ be the corresponding trajectory of W and put $v = (w(t))_{t \in [\delta,T]}$. The sequential observation of w starts at the knot $\varphi_1(v)$. After n steps the available information is then given by $D_n(\omega) = (y_1, \ldots, y_n, v)$, where $y_1 = w(\varphi_1(v)), \ldots, y_n = w(\varphi_n(y_1, \ldots, y_{n-1}, v))$, and we decide whether we stop or further evaluate w according to the value of $\chi_n(D_n(\omega))$. The total number of observations of w in the interval $(0, \delta)$ is thus given by

(6)
$$\nu(\omega) = \min\{n \in \mathbb{N} : \chi_n(D_n(\omega)) = 1\}.$$

If $\nu(\omega) < \infty$, then the data $D_{\nu(\omega)}(\omega)$ is used to construct the estimate $\phi_{\nu(\omega)}(D_{\nu(\omega)}(\omega)) \in \mathbb{R}^4$.

For obvious reasons we require that $\nu < \infty$ P-a.s. Then the resulting approximation is given by

$$\widehat{X} = \phi_{\nu}(D_{\nu}).$$

Without loss of generality we assume that

(7)
$$\varphi_k(y_1,\ldots,y_{k-1},v) \neq \varphi_l(y_1,\ldots,y_{l-1},v)$$

for all $y \in \mathbb{R}^{\infty}$, all $k, l \in \mathbb{N}$ with $k \neq l$ and all $v \in C([\delta, T])$. We put

$$c(\widehat{X}) = \mathbb{E}\nu,$$

that is the expected number of evaluations of the driving Brownian motion W in the interval $(0, \delta)$. We denote by \mathcal{X}^{δ} the class of all methods of the above form and for $N \in \mathbb{N}$ we put

$$\mathcal{X}_N^{\delta} = \{ \widehat{X} \in \mathcal{X}^{\delta} : c(\widehat{X}) \le N \}.$$

Clearly, $\mathcal{X}_N^{\delta_1} \subseteq \mathcal{X}_N^{\delta_2}$ for all $0 < \delta_2 \leq \delta_1 \leq T$ and all $N \in \mathbb{N}$. Let us stress that the class \mathcal{X}_N^{δ} contains in particular all methods from the literature, which use a step size control based on N sequential evaluations of W on average, see e.g. [6, 29, 30,27, 34, 22, 13, 14] and the references therein. Moreover, \mathcal{X}_N^{δ} of course contains all nonadaptive approximations $u(W(s_1), \ldots, W(s_N), (W(t))_{t \in [\delta, T]})$ based on N evaluations of W at fixed time points $s_1, \ldots, s_N \in (0, \delta)$ and on $(W(t))_{t \in [\delta, T]}$, as studied in [20]. In the latter case one can take any sequences φ, χ and ϕ satisfying

$$\varphi_n = s_n \text{ for } n \leq N, \chi_1 = \dots \chi_{N-1} = 0, \chi_N = 1 \text{ and } \phi_N = u.$$

5. Main results

Assume the setting in Section 3 and put

(8)
$$\alpha = \inf_{t \in [0,\tau_1/2]} |f'(t)|^2, \quad \beta = \int_{\tau_1}^{\tau_2} g^2(t) dt, \quad \gamma = \int_{\tau_2}^T h(t) dt$$

as well as

$$c_1 = \frac{\gamma \exp\left(-\frac{\pi^2}{4} - \frac{1}{\beta}\right)}{8\pi\sqrt{2\pi\beta}}, \quad c_2 = \frac{\gamma \exp\left(-\frac{\pi^2}{4}\right)}{4\pi}.$$

Our main result is stated in Theorem 1. It provides a uniform lower bound for the mean absolute error of any strong approximation of $X^{\psi}(T)$ that is based on $(W(t))_{t \in [\delta,T]}$ and on N sequential evaluations of W in the interval $(0, \delta)$ on average in the case that ψ is positive, strictly increasing and satisfies $\lim_{x\to\infty} \psi(x) = \infty$ as well as $1 \in \psi(\mathbb{R})$. See Section 6 for the proof.

Theorem 1. Let $\delta \in (0,T]$ and let $\psi \in C^{\infty}(\mathbb{R})$ be positive, strictly increasing with $\lim_{x\to\infty} \psi(x) =$ ∞ and $1 \in \psi(\mathbb{R})$. Then for all $N \in \mathbb{N}$ and all $\widehat{X} \in \mathcal{X}_N^{\delta}$ we have

(9)
$$\mathbb{E}\|X^{\psi}(T) - \widehat{X}\| \ge c_1 \cdot \exp\left(-\frac{1}{\beta} \cdot \left(\psi^{-1}\left(\left(1 + \sqrt{\frac{96}{\alpha(\min(\delta,\tau_1/2))^3}}\right)N^3\right)\right)^2\right) - \frac{c_2}{N}.$$

As a consequence of Theorem 1 we obtain a non-polynomial decay of the smallest possible mean absolute error of strong approximation of $X^{\psi}(T)$ based on $(W(t))_{t \in [\delta,T]}$ and on N sequential evaluations of W in the interval $(0, \delta)$ on average if ψ additionally satisfies an exponential growth condition.

Corollary 1. Let $\delta \in (0, T]$ and let $\psi \in C^{\infty}(\mathbb{R})$ be positive, strictly increasing with $\lim_{x\to\infty} \psi(x) = \infty$ and $1 \in \psi(\mathbb{R})$. Moreover assume that for all $q \in (0, \infty)$

$$\lim_{x \to \infty} \psi(x) \cdot \exp(-qx^2) = \infty.$$

Then for all $q \in (0, \infty)$ we have

$$\lim_{N \to \infty} \left(N^q \cdot \inf_{\widehat{X} \in \mathcal{X}_N^{\delta}} \mathbb{E} \| X^{\psi}(T) - \widehat{X} \| \right) = \infty.$$

Proof. The assumptions on the function ψ ensure that for all $q \in (0, \infty)$

(10)
$$\lim_{N \to \infty} \left(N^q \cdot \exp\left(-\frac{1}{\beta} \cdot \left(\psi^{-1}\left(\left(1 + \sqrt{\frac{96}{\alpha(\min(\delta, \tau_1/2))^3}}\right)N^3\right)\right)^2\right) \right) = \infty,$$

see Lemma 4.5 in [20]. This in particular implies that there exists $N_0 \in \mathbb{N}$ such that for all $N \geq N_0$

$$\frac{c_1}{2} \cdot \exp\left(-\frac{1}{\beta} \cdot \left(\psi^{-1}\left(\left(1 + \sqrt{\frac{96}{\alpha(\min(\delta,\tau_1/2))^3}}\right)N^3\right)\right)^2\right) \ge \frac{c_2}{N}$$

Employing Theorem 1 we therefore conclude that for all $N \geq N_0$

$$\inf_{\widehat{X}\in\mathcal{X}_{N}^{\delta}}\mathbb{E}\|X^{\psi}(T)-\widehat{X}\| \geq \frac{c_{1}}{2} \cdot \exp\left(-\frac{1}{\beta} \cdot \left(\psi^{-1}\left(\left(1+\sqrt{\frac{96}{\alpha(\min(\delta,\tau_{1}/2))^{3}}}\right)N^{3}\right)\right)^{2}\right).$$

The latter estimate and (10) imply the statement of the corollary.

The following result shows that the smallest possible mean absolute error of strong approximation of $X^{\psi}(T)$ based $(W(t))_{t \in [\delta,T]}$ and on N sequential evaluations of W in the interval $(0, \delta_N)$ on average may converge to zero arbitrarily slow even then when the sequence $(\delta_N)_{N \in \mathbb{N}}$ tends to zero with any given speed.

Corollary 2. Let $(a_N)_{N \in \mathbb{N}} \subset (0, \infty)$ and $(\delta_N)_{N \in \mathbb{N}} \subset (0, T]$ satisfy $\lim_{N \to \infty} a_N = 0$ and $\lim_{N \to \infty} \delta_N = 0$. Then there exists $\kappa > 0$ and $\psi \in C^{\infty}(\mathbb{R})$ such that for all $N \in \mathbb{N}$ we have

$$\inf_{\widehat{X}\in\mathcal{X}_N^{\delta_N}} \mathbb{E} \|X^{\psi}(T) - \widehat{X}\| \ge \kappa \cdot a_N.$$

Proof. We proceed similar to the proof of Corollary 4.3 in [20]. Without loss of generality we may assume that the sequences $(a_N)_{N \in \mathbb{N}}$ and $(\delta_N)_{N \in \mathbb{N}}$ are strictly decreasing. Let

$$N_0 = \min\left\{N \in \mathbb{N} \colon a_N + \frac{c_2}{N} \le c_1\right\}$$

and for $N \geq N_0$ put

$$b_N = \sqrt{-\beta \ln\left(\frac{1}{c_1} \cdot \left(a_N + \frac{c_2}{N}\right)\right)}, \quad d_N = \left(1 + \sqrt{\frac{96}{\alpha(\min(\delta_N, \tau_1/2))^3}}\right) N^3.$$

Note that the sequences $(b_N)_{N\geq N_0}$ and $(d_N)_{N\geq N_0}$ are strictly increasing and satisfy

$$\lim_{N \to \infty} b_N = \lim_{N \to \infty} d_N = \infty.$$

Define a function $\psi \colon \mathbb{R} \to \mathbb{R}$ by

$$\psi(x) = \begin{cases} d_{N_0} \cdot \left(1 - \exp\left(\frac{1}{x - b_{N_0}}\right)\right), & \text{if } x < b_{N_0}, \\ \\ d_N, & \text{if } x = b_N \text{ and } N \ge N_0, \\ \\ d_{N-1} + \frac{d_N - d_{N-1}}{1 + \exp\left(\frac{1}{x - b_{N-1}} - \frac{1}{b_N - x}\right)}, & \text{if } x \in (b_{N-1}, b_N) \text{ and } N > N_0 \end{cases}$$

Then ψ is positive, strictly increasing, infinitely often differentiable and satisfies $\lim_{x\to\infty} \psi(x) = \infty$ as well as $1 \in \psi(\mathbb{R})$.

For $N \in \mathbb{N}$ put

$$\varepsilon_N = \inf_{\widehat{X} \in \mathcal{X}_N^{\delta_N}} \mathbb{E} \| X^{\psi}(T) - \widehat{X} \|.$$

Theorem 1 implies that for all $N \ge N_0$

$$\varepsilon_N \ge c_1 \cdot \exp\left(-\frac{1}{\beta} \cdot (\psi^{-1}(d_N))^2\right) - \frac{c_2}{N} = c_1 \cdot \exp\left(-\frac{1}{\beta} \cdot b_N^2\right) - \frac{c_2}{N} = a_N.$$

Since the sequence $(\varepsilon_N)_{N \in \mathbb{N}}$ is decreasing hence for all $N \in \{1, 2, \dots, N_0\}$

$$\varepsilon_N \ge \varepsilon_{N_0} \ge a_{N_0}$$

Using the assumption that the sequence $(a_N)_{N \in \mathbb{N}}$ is strictly decreasing we therefore conclude that for all $N \in \mathbb{N}$

$$\varepsilon_N \ge \min\{1, a_{N_0}/a_N\} \cdot a_N \ge \frac{a_{N_0}}{a_1} \cdot a_N$$

which completes the proof of the corollary with $\kappa = a_{N_0}/a_1$.

6. Proof of Theorem 1

Let $\delta \in (0,T]$ and let $\widehat{X} \in \mathcal{X}^{\delta}$ be given by sequences $\varphi = (\varphi_n)_{n \in \mathbb{N}}, \chi = (\chi_n)_{n \in \mathbb{N}}$ and $\phi = (\phi_n)_{n \in \mathbb{N}}$, see (5). Recall the definition (6) of ν . We first determine the regular conditional distribution $\mathbb{P}^{W|D_{\nu}}$.

For $n \in \mathbb{N}$ put

$$S_n = \{s \in (0, \delta)^n : |\{s_1, \dots, s_n\}| = n\}.$$

For $n \in \mathbb{N}$, $s \in S_n$, $y \in \mathbb{R}^n$ and $v \in C([\delta, T])$ define functions

$$m_{s,y,v} \colon [0,T] \to \mathbb{R} \text{ and } R_s \colon [0,T]^2 \to \mathbb{R}$$

as follows. If $s_1 < \ldots < s_n$ put $s_0 = y_0 = 0$, $s_{n+1} = \delta$ and $y_{n+1} = v(\delta)$ and let

$$m_{s,y,v}(t) = \begin{cases} \frac{s_k - t}{s_k - s_{k-1}} \cdot y_{k-1} + \frac{t - s_{k-1}}{s_k - s_{k-1}} \cdot y_k, & \text{if } t \in [s_{k-1}, s_k) \text{ for } k \in \{1, \dots, n+1\}, \\ v(t), & \text{if } t \in [\delta, T] \end{cases}$$

as well as

$$R_s(r,t) = \begin{cases} \frac{(s_k - \max(r,t)) \cdot (\min(r,t) - s_{k-1})}{s_k - s_{k-1}}, & \text{if } r, t \in [s_{k-1}, s_k) \text{ for } k \in \{1, \dots, n+1\}, \\ 0, & \text{otherwise}, \end{cases}$$

for $r, t \in [0, T]$. Otherwise put

$$m_{s,y,v} = m_{(s_{\pi(1)},\dots,s_{\pi(n)}),(y_{\pi(1)},\dots,y_{\pi(n)}),v}, \quad R_s = R_{(s_{\pi(1)},\dots,s_{\pi(n)})},$$

where π is the permutation of $\{1, \ldots, n\}$ such that $s_{\pi(1)} < \ldots < s_{\pi(n)}$.

For $n \in \mathbb{N}$, $y \in \mathbb{R}^n$, $v \in C([\delta, T])$ and $k = 1, \ldots, n$ put

(11)
$$s_k^{y,v} = \varphi_k(y_1, \dots, y_{k-1}, v).$$

Note that due to the assumption (7) we have $|\{s_1^{y,v}, \ldots, s_n^{y,v}\}| = n$. Let $Q_{y,v}$ denote the Gaussian measure on $\mathfrak{B}(C([0,T]))$ with mean $m(Q_{y,v}) = m_{s^{y,v},y,v}$ and covariance function $R(Q_{y,v}) = R_{s^{y,v}}$. Consider the measurable space

(12)
$$(\Omega_1, \mathcal{F}_1) = \left(\bigcup_{n=1}^{\infty} \mathbb{R}^n \times C([\delta, T]), \sigma\left(\bigcup_{n=1}^{\infty} \mathfrak{B}(\mathbb{R}^n \times C([\delta, T]))\right)\right)$$

It is easy to see that $D_{\nu}: \Omega \to \Omega_1$ is \mathcal{F} - \mathcal{F}_1 measurable. Define the mapping

$$K: \Omega_1 \times \mathfrak{B}(C([0,T])) \to [0,1]$$

by

$$K((y,v),A) = Q_{y,v}(A)$$

for all $y \in \bigcup_{n=1}^{\infty} \mathbb{R}^n, v \in C([\delta, T])$ and $A \in \mathfrak{B}(C([0, T]))$.

Lemma 1. K is a version of the regular conditional distribution $\mathbb{P}^{W|D_{\nu}}$.

In the case of $\delta = T$ the statement of Lemma 1 seems to be well-known, see, e.g., [15, 28, 29, 30], but a proof of it seems not to be available in the literature. If, additionally, ν is constant then Lemma 1 follows from Lemma 2.9.7 in [36, p. 474], but measurability issues have not been fully addressed in the proof of the latter result. For convenience of the reader we therefore provide a proof of Lemma 1 here.

Proof. Clearly, for all $(y, v) \in \Omega_1$ the mapping

$$\mathfrak{B}(C([0,T])) \ni A \mapsto K((y,v),A) \in [0,1]$$

is a probability measure on $\mathfrak{B}(C([0,T]))$.

Next, let $A \in \mathfrak{B}(C([0,T]))$. We show that the mapping

(13)
$$\Omega_1 \ni (y, v) \mapsto K((y, v), A) \in [0, 1]$$

is \mathcal{F}_1 - $\mathfrak{B}([0,1])$ measurable. For $n \in \mathbb{N}$, $s \in S_n$ and $u \in C([0,\delta])$ define a function

$$F_{s,u}\colon [0,T] \to \mathbb{R}$$

by

$$F_{s,u}(t) = \begin{cases} u(t) - m_{s,(u(s_1),\dots,u(s_n)),0}(t), & \text{if } t \in [0,\delta], \\ 0, & \text{if } t \in (\delta,T] \end{cases}$$

for $t \in [0,T]$. It is easy to see that for all $(y,v) \in \Omega_1$

$$Q_{y,v} = \mathbb{P}^{F_s y, v, (W(t))_{t \in [0,\delta]} + m_s y, v, y, v}$$

and therefore for all $(y, v) \in \Omega_1$

(14)
$$K((y,v),A) = \int_{C([0,\delta])} 1_A(F_{s^{y,v},u} + m_{s^{y,v},y,v}) \mathbb{P}^{(W(t))_{t \in [0,\delta]}}(du).$$

Clearly,

$$\{((y,v),u) \in \Omega_1 \times C([0,\delta]) : F_{s^{y,v},u} + m_{s^{y,v},y,v} \in A\} = \bigcup_{n=1}^{\infty} A_n,$$

where

$$A_n = \{ (y, v, u) \in \mathbb{R}^n \times C([\delta, T]) \times C([0, \delta]) : F_{s^{y, v}, u} + m_{s^{y, v}, y, v} \in A \}$$

for $n \in \mathbb{N}$. The measurability of the functions $\varphi_n, n \in \mathbb{N}$, imply that for every $n \in \mathbb{N}$ the mapping

$$\mathbb{R}^n \times C([\delta, T]) \ni (y, v) \mapsto s^{y, v} \in S_n$$

is $\mathfrak{B}(\mathbb{R}^n \times C([\delta, T])) - \mathfrak{B}(S_n)$ measurable. Thus, observing that for every $n \in \mathbb{N}$ the mappings

$$S_n \times C([0, \delta]) \ni (s, u) \mapsto F_{s, u} \in C([0, T])$$

and

$$S_n \times \mathbb{R}^n \times C([\delta, T]) \ni (s, y, v) \mapsto m_{s, y, v} \in C([0, T])$$

are continuous we conclude that for every $n \in \mathbb{N}$ the mapping

$$\mathbb{R}^n \times C([\delta, T]) \times C([0, \delta]) \ni (y, v, u) \mapsto F_{s^{y, v}, u} + m_{s^{y, v}, y, v} \in C([0, T])$$

is $\mathfrak{B}(\mathbb{R}^n \times C([\delta, T]) \times C([0, \delta]) - \mathfrak{B}(C([0, T]))$ measurable. Hence for every $n \in \mathbb{N}$

$$A_n \in \mathfrak{B}(\mathbb{R}^n \times C([\delta, T]) \times C([0, \delta])) \subset \mathcal{F}_1 \otimes \mathfrak{B}(C([0, \delta])),$$

which implies that the mapping

$$\Omega_1 \times C([0,\delta]) \ni ((y,v),u) \mapsto 1_A(F_{s^{y,v},u} + m_{s^{y,v},y,v}) \in \mathbb{R}$$

is $\mathcal{F}_1 \otimes \mathfrak{B}(C([0, \delta])) - \mathfrak{B}(R)$ measurable. Using (14) and employing Fubini's theorem we thus conclude that the mapping (13) is $\mathcal{F}_1 - \mathfrak{B}([0, 1])$ measurable.

Finally, let $A \in \mathfrak{B}(C([0,T]))$ and $E \in \mathcal{F}_1$. We show that

(15)
$$\mathbb{P}(\{W \in A\} \cap \{D_{\nu} \in E\}) = \int_{E} K((y,v),A) \ \mathbb{P}^{D_{\nu}}(d(y,v)).$$

We have

$$\{D_{\nu} \in E\} \cap \{\nu < \infty\} = \bigcup_{n=1}^{\infty} (\{D_{\nu} \in E\} \cap \{\nu = n\})$$
$$= \bigcup_{n=1}^{\infty} \left(\{D_{n} \in E\} \cap \{\chi_{n}(D_{n}) = 1\} \cap \bigcap_{k=1}^{n-1} \{\chi_{k}(D_{k}) = 0\}\right)$$
$$= \bigcup_{n=1}^{\infty} \{D_{n} \in E \cap C_{n}\},$$

where $C_n \in \mathfrak{B}(\mathbb{R}^n \times C([\delta, T]))$ is given by

$$C_n = \chi_n^{-1}(\{1\}) \cap \bigcap_{k=1}^{n-1} \{(y,v) \in \mathbb{R}^n \times C([\delta,T]) : (y_1,\dots,y_k,v) \in \chi_k^{-1}(\{0\})\}$$

for $n \in \mathbb{N}$. Since $\nu < \infty$ a.s. we thus obtain

$$\mathbb{P}(\{W \in A\} \cap \{D_{\nu} \in E\}) = \sum_{n=1}^{\infty} \mathbb{P}(\{W \in A\} \cap \{D_n \in E \cap C_n\})$$
$$= \sum_{n=1}^{\infty} \int_{E \cap C_n} \mathbb{P}^{W|D_n = (y,v)}(A) \mathbb{P}^{D_n}(d(y,v))$$

Similarly to the proof of Lemma 2.9.7 on page 474 in [36] one can show that for all $n \in \mathbb{N}$ and all $G \in \mathfrak{B}(\mathbb{R}^n \times C([\delta, T]))$

$$\int_G \mathbb{P}^{W|D_n=(y,v)}(A) \ \mathbb{P}^{D_n}(d(y,v)) = \int_G Q_{y,v}(A) \ \mathbb{P}^{D_n}(d(y,v)).$$

Hence

$$\mathbb{P}(\{W \in A\} \cap \{D_{\nu} \in E\}) = \sum_{n=1}^{\infty} \int_{E \cap C_n} Q_{y,v}(A) \mathbb{P}^{D_n}(d(y,v))$$
$$= \sum_{n=1}^{\infty} \int_{E \cap C_n} K((y,v),A) \mathbb{P}^{D_n}(d(y,v))$$

Since $\mathbb{P}^{D_n}(G \cap C_n) = \mathbb{P}^{D_\nu}(G)$ for all $n \in \mathbb{N}$ and all $G \in \mathfrak{B}(\mathbb{R}^n \times C([\delta, T]))$ we thus conclude

$$\mathbb{P}(\{W \in A\} \cap \{D_{\nu} \in E\}) = \sum_{n=1}^{\infty} \int_{E \cap (\mathbb{R}^n \times C([\delta, T]))} K((y, v), A) \ \mathbb{P}^{D_{\nu}}(d(y, v)),$$

which implies (15) and completes the proof of the lemma.

Next, assume that $\delta \in (0, \tau_1]$, let $n \in \mathbb{N}$, $y \in \mathbb{R}^n$ and $v \in C([\delta, T])$. Recall the definition (11) of the time points $s_1^{y,v}, \ldots, s_n^{y,v}$ and put

(16)
$$s_0^{y,v} = 0, \quad s_{n+1}^{y,v} = \delta.$$

Let π be the permutation of $\{0, \ldots, n+1\}$ such that $s_{\pi(0)}^{y,v} < \ldots < s_{\pi(n+1)}^{y,v}$. Let $i^* \in \{0, \ldots, n\}$ and put

(17)
$$t_0 = s_{\pi(i^*)}^{y,v}, \quad t_1 = s_{\pi(i^*+1)}^{y,v}.$$

Define mappings

$$W: C([0,T]) \to C([0,t_0] \cup [t_1,\tau_1]), \ B: C([0,T]) \to C([t_0,t_1])$$

by

(18)
$$W(w) = w|_{[0,t_0] \cup [t_1,\tau_1]}$$

for $w \in C([0,T])$ and

(19)
$$B(w)(t) = w(t) - \frac{t_1 - t}{t_1 - t_0} w(t_0) - \frac{t - t_0}{t_1 - t_0} w(t_1)$$

for $w \in C([0,T])$ and $t \in [t_0, t_1]$. In the following lemma we present properties of the measures $Q_{y,v}^{\widetilde{W}}, Q_{y,v}^B$ and $Q_{y,v}^{(\widetilde{W},B)}$, which will be used in the proof of Theorem 1.

Lemma 2. We have

- (i) Q_{y,v}^{W̃} is the Gaussian measure on 𝔅(C([0,t₀] ∪ [t₁,τ₁])) with mean m(Q_{y,v}^{W̃}) = m_{s^{y,v},y,v}|_{[0,t₀]∪[t₁,τ₁]} and covariance function R(Q_{y,v}^{W̃}) = R_{s^{y,v}}|_{([0,t₀]∪[t₁,τ₁])²}.
 (ii) Q_{y,v}^B is the Gaussian measure on 𝔅(C([t₀,t₁])) with mean m(Q_{y,v}^B) = 0 and covariance function

$$R(Q_{y,v}^B)(r,t) = \frac{(t_1 - \max(r,t)) \cdot (\min(r,t) - t_0)}{t_1 - t_0}, \quad r,t \in [t_0,t_1].$$

 $\begin{array}{ll} \text{(iii)} & Q^B_{y,v} = Q^{-B}_{y,v}, \\ \text{(iv)} & Q^{(\widetilde{W},B)}_{y,v} = Q^{\widetilde{W}}_{y,v} \times Q^B_{y,v}. \end{array}$

Proof. The property (i) is obvious. Since B is a linear continuous mapping hence $Q_{y,v}^B$ is a Gaussian measure. Next, observe that for all $t \in [t_0, t_1]$ we have

$$m(Q_{y,v})(t) = \frac{t_1 - t}{t_1 - t_0} y_{\pi(i^*)} + \frac{t - t_0}{t_1 - t_0} y_{\pi(i^*+1)}.$$

In particular, for all $t \in [t_0, t_1]$,

(20)
$$m(Q_{y,v})(t) - \frac{t_1 - t}{t_1 - t_0} m(Q_{y,v})(t_0) - \frac{t - t_0}{t_1 - t_0} m(Q_{y,v})(t_1) = 0.$$

Hence for all $t \in [t_0, t_1]$,

$$\begin{split} m(Q_{y,v}^B)(t) &= \int_{C([t_0,t_1])} b(t) Q_{y,v}^B(db) \\ &= \int_{C([0,T])} \left(w(t) - \frac{t_1 - t}{t_1 - t_0} w(t_0) - \frac{t - t_0}{t_1 - t_0} w(t_1) \right) Q_{y,v}(dw) \\ &= m(Q_{y,v})(t) - \frac{t_1 - t}{t_1 - t_0} m(Q_{y,v})(t_0) - \frac{t - t_0}{t_1 - t_0} m(Q_{y,v})(t_1) = 0 \end{split}$$

and for all $r, t \in [t_0, t_1]$,

$$\begin{aligned} R(Q_{y,v}^B)(r,t) &= \int_{C([t_0,t_1])} \left(b(r) - m(Q_{y,v}^B)(r) \right) \cdot \left(b(t) - m(Q_{y,v}^B)(t) \right) Q_{y,v}^B(db) \\ &= \int_{C([0,T])} \left(w(r) - \frac{t_1 - r}{t_1 - t_0} w(t_0) - \frac{r - t_0}{t_1 - t_0} w(t_1) \right) \\ &\quad \cdot \left(w(t) - \frac{t_1 - t}{t_1 - t_0} w(t_0) - \frac{t - t_0}{t_1 - t_0} w(t_1) \right) Q_{y,v}(dw) \end{aligned}$$

and hence

$$\begin{split} R(Q_{y,v}^B)(r,t) &= \int_{C([0,T])} \Big(w(r) - m(Q_{y,v})(r) - \frac{t_1 - r}{t_1 - t_0} (w(t_0) - m(Q_{y,v})(t_0)) \\ &\quad - \frac{r - t_0}{t_1 - t_0} (w(t_1) - m(Q_{y,v})(t_1)) \Big) \\ &\quad \cdot \left(w(t) - m(Q_{y,v})(t) - \frac{t_1 - t}{t_1 - t_0} (w(t_0) - m(Q_{y,v})(t_0)) \\ &\quad - \frac{t - t_0}{t_1 - t_0} (w(t_1) - m(Q_{y,v})(t_1)) \Big) Q_{y,v}(dw) \\ &= R(Q_{y,v})(r,t) - \frac{t_1 - t}{t_1 - t_0} R(Q_{y,v})(r,t_0) - \frac{t - t_0}{t_1 - t_0} R(Q_{y,v})(r,t_1) \\ &\quad - \frac{t_1 - r}{t_1 - t_0} \Big(R(Q_{y,v})(t_0,t) - \frac{t_1 - t}{t_1 - t_0} R(Q_{y,v})(t_0,t_0) - \frac{t - t_0}{t_1 - t_0} R(Q_{y,v})(t_1,t_1) \Big) \\ &\quad - \frac{r - t_0}{t_1 - t_0} \Big(R(Q_{y,v})(t_1,t) - \frac{t_1 - t}{t_1 - t_0} R(Q_{y,v})(t_1,t_0) - \frac{t - t_0}{t_1 - t_0} R(Q_{y,v})(t_1,t_1) \Big). \end{split}$$

Observing that $R(Q_{y,v})(t_0,t) = R(Q_{y,v})(t_1,t) = 0$ for all $t \in [t_0,t_1]$ we thus obtain

$$R(Q_{y,v}^B)(r,t) = R(Q_{y,v})(r,t)$$

for all $r, t \in [t_0, t_1]$, which completes the proof of (ii). The property (ii) implies that $Q_{y,v}^{-B}$ is the Gaussian measure on $\mathfrak{B}(C([t_0, t_1]))$ with mean $m(Q_{y,v}^{-B}) = 0$ and covariance function $R(Q_{y,v}^{-B}) = R(Q_{y,v}^{B})$, which yields the property (iii). Next, we prove (iv). Using the properties (i), (ii) as well as (20) and the fact that $R(Q_{y,v})(r,t) = 0$ for all $r \in [0, t_0] \cup [t_1, \tau_1]$ and $t \in [t_0, t_1]$ we obtain

$$\begin{split} &\int_{C([0,T])} \left(\widetilde{W}(w)(r) - m(Q_{y,v}^{\widetilde{W}})(r)\right) \cdot \left(B(w)(t) - m(Q_{y,v}^{B})(t)\right) Q_{y,v}(dw) \\ &= \int_{C([0,T])} \left(w(r) - m(Q_{y,v})(r)\right) \cdot \left(w(t) - \frac{t_1 - t}{t_1 - t_0} w(t_0) - \frac{t - t_0}{t_1 - t_0} w(t_1)\right) Q_{y,v}(dw) \\ &= \int_{C([0,T])} \left(w(r) - m(Q_{y,v})(r)\right) \cdot \left(w(t) - m(Q_{y,v})(t) - \frac{t_1 - t}{t_1 - t_0} (w(t_0) - m(Q_{y,v})(t_0)) \right. \\ &\qquad \left. - \frac{t - t_0}{t_1 - t_0} (w(t_1) - m(Q_{y,v})(t_1))\right) Q_{y,v}(dw) \\ &= R(Q_{y,v})(r,t) - \frac{t_1 - t}{t_1 - t_0} R(Q_{y,v})(r,t_0) - \frac{t - t_0}{t_1 - t_0} R(Q_{y,v})(r,t_1) = 0 \end{split}$$

for all $r \in [0, t_0] \cup [t_1, \tau_1]$ and $t \in [t_0, t_1]$, which means that $\widetilde{W}(r)$ and B(t) are uncorrelated. This and the fact that $Q_{y,v}^{\widetilde{W}}$ and $Q_{y,v}^B$ are Gaussian measures implies (iv).

In the proof of Theorem 1 we employ the following lower bound for the first absolute moment of the sine of a normally distributed random variable, which is a generalization of Lemma 4.2 from [20], where a centered normally distributed random variable has been considered.

Lemma 3. Let $a \in \mathbb{R}$, $\tau \in [1, \infty)$, and let $Y \colon \Omega \to \mathbb{R}$ be a $\mathcal{N}(a, \tau^2)$ -distributed random variable. Then

$$\mathbb{E}\left[|\sin(Y)|\right] \ge \frac{\exp\left(-\frac{\pi^2}{8}\right)}{\sqrt{8\pi}}.$$

The proof of Lemma 3 is a straighforward generalization of the proof of Lemma 4.2 from [20]. We now proceed with the proof of Theorem 1.

We first consider the case $\delta \in (0, \tau_1/2]$. Let $N \in \mathbb{N}$ be such that $\widehat{X} \in \mathcal{X}_N^{\delta}$. We may then assume that $\widehat{X} \in \mathcal{X}_{N+1}^{\delta}$ and that \widehat{X} uses the evaluation site $\delta/2$. Let $\psi \in C^{\infty}(\mathbb{R})$ be positive, strictly increasing with $\lim_{x\to\infty} \psi(x) = \infty$ and $1 \in \psi(\mathbb{R})$. Using Remark 2, the assumptions on the functions f and g and Itô's formula we obtain that \mathbb{P} -a.s.

(21)
$$X_2^{\psi}(\tau_1) = -\int_0^{\tau_1} f'(t)W(t)dt, \quad X_3^{\psi}(\tau_2) = -\int_{\tau_1}^{\tau_2} g'(t)W(t)dt.$$

Thus, \mathbb{P} -a.s.

(22)
$$X_4^{\psi}(T) = F(W)$$

where $F: C([0,T]) \to \mathbb{R}$ is given by

$$F(w) = \gamma \cdot \cos\left(\int_0^{\tau_1} f'(t)w(t)dt \cdot \psi\left(-\int_{\tau_1}^{\tau_2} g'(t)w(t)dt\right)\right)$$

and γ is defined in (8). Recall the definition (12) of the measurable space $(\Omega_1, \mathcal{F}_1)$ and define a function $G: \Omega_1 \to \mathbb{R}^4$ by

$$G(y,v) = \phi_n(y,v)$$

for $y \in \mathbb{R}^n$ and $v \in C([\delta, T])$. Due to the measurability of the functions ϕ_n , $n \in \mathbb{N}$, the function G is \mathcal{F}_1 - $\mathfrak{B}(\mathbb{R}^4)$ measurable. Moreover,

(23)
$$\widehat{X} = \phi_{\nu}(D_{\nu}) = G(D_{\nu})$$

Let $pr_4: \mathbb{R}^4 \to \mathbb{R}$ denote the projection to the fourth component. Due to (22) and (23) we have

(24)
$$\mathbb{E}|X_4^{\psi}(T) - \hat{X}_4| = \mathbb{E}|F(W) - \mathrm{pr}_4(G(D_{\nu}))| = \mathbb{E}\left(\mathbb{E}\left(|F(W) - \mathrm{pr}_4(G(D_{\nu}))| \mid D_{\nu}\right)\right).$$

Lemma 1 implies that for $\mathbb{P}^{D_{\nu}}$ -a.a. $(y, v) \in \Omega_1$

(25)
$$\mathbb{E}(|F(W) - \mathrm{pr}_4(G(D_\nu))| \mid D_\nu = (y, v)) = \int_{C([0,T])} |F(w) - \mathrm{pr}_4(G(y, v))| Q_{y,v}(dw).$$

Fix $n \in \mathbb{N}$, $y \in \mathbb{R}^n$ and $v \in C([\delta, T])$. We show that

(26)
$$\int_{C([0,T])} |F(w) - \operatorname{pr}_4(G(y,v))| Q_{y,v}(dw) \geq \frac{\gamma \exp\left(-\frac{\pi^2}{4}\right)}{8\pi} \cdot \mathbb{1}_{[1,\infty)} \left(\sqrt{\frac{\alpha\delta^3}{96n^3}} \cdot \psi\left(-\int_{\tau_1}^{\tau_2} g'(t) v(t) dt\right)\right),$$

where α is given by (8). Recall the definitions (11) and (16) of the time points $s_0^{y,v}, \ldots, s_{n+1}^{y,v}$, let π be the permutation of $\{0, \ldots, n+1\}$ such that $s_{\pi(0)}^{y,v} < \ldots < s_{\pi(n+1)}^{y,v}$, and let t_0 and t_1 be given by (17) with

(27)
$$i^* = \min\{i \in \{0, \dots, n\} : |s_{\pi(i+1)}^{y,v} - s_{\pi(i)}^{y,v}| \ge \delta/(n+1)\}.$$

Put

$$C_1 = C([0, t_0] \cup [t_1, \tau_1]), \quad C_2 = C([t_0, t_1]), \quad C_3 = C([\tau_1, \tau_2]),$$

let $\widetilde{W}: C([0,T]) \to C_1$ and $B: C([0,T]) \to C_2$ be given by (18) and (19). respectively, and define $\overline{W}: C([0,T]) \to C_3$ by

$$\overline{W}(w) = w|_{[\tau_1, \tau_2]}$$

for $w \in C([0,T])$. Moreover, define mappings $H_1: C_1 \to \mathbb{R}, H_2: C_2 \to \mathbb{R}, H_3: C_3 \to \mathbb{R}$ and $J: C_1 \times C_2 \times C_3 \to \mathbb{R}$ by

$$H_{1}(\widetilde{w}) = \int_{0}^{t_{0}} f'(t) \,\widetilde{w}(t) dt + \int_{t_{0}}^{t_{1}} f'(t) \left(\frac{t - t_{0}}{t_{1} - t_{0}} \widetilde{w}(t_{1}) + \frac{t_{1} - t}{t_{1} - t_{0}} \widetilde{w}(t_{0})\right) dt + \int_{t_{1}}^{\tau_{1}} f'(t) \,\widetilde{w}(t) dt,$$

$$H_{2}(b) = \int_{t_{0}}^{t_{1}} f'(t) \,b(t) dt, \quad H_{3}(\overline{w}) = -\int_{\tau_{1}}^{\tau_{2}} g'(t) \,\overline{w}(t) dt$$

as well as

$$J(\widetilde{w}, b, \overline{w}) = \gamma \cdot \cos\left(\left(H_1(\widetilde{w}) + H_2(b)\right) \cdot \psi(H_3(\overline{w}))\right)$$

for $\widetilde{w} \in C_1$, $b \in C_2$ and $\overline{w} \in C_3$. We then have

$$F = J(\overline{W}, B, \overline{W}).$$

Clearly, $Q_{y,v}^{\overline{W}}$ is the dirac measure concentrated at $v|_{[\tau_1,\tau_2]}$. Using Lemma 2(iii),(iv) and the triangle inequality we thus obtain

$$\begin{split} \int_{C([0,T])} |F(w) - \mathrm{pr}_4(G(y,v))| \, Q_{y,v}(dw) \\ &= \int_{C_1} \int_{C_2} |J(\widetilde{w}, b, v|_{[\tau_1, \tau_2]}) - \mathrm{pr}_4(G(y,v))| \, Q_{y,v}^B(db) \, Q_{y,v}^{\widetilde{W}}(d\widetilde{w}) \\ &= \int_{C_1} \int_{C_2} \frac{1}{2} \left(|J(\widetilde{w}, b, v|_{[\tau_1, \tau_2]}) - \mathrm{pr}_4(G(y,v))| \right) \\ &+ |J(\widetilde{w}, -b, v|_{[\tau_1, \tau_2]}) - \mathrm{pr}_4(G(y,v))| \right) \, Q_{y,v}^B(db) \, Q_{y,v}^{\widetilde{W}}(d\widetilde{w}) \\ &\geq \int_{C_1} \int_{C_2} \frac{1}{2} \left| J(\widetilde{w}, b, v|_{[\tau_1, \tau_2]}) - J(\widetilde{w}, -b, v|_{[\tau_1, \tau_2]}) \right| \, Q_{y,v}^B(db) \, Q_{y,v}^{\widetilde{W}}(d\widetilde{w}). \end{split}$$

Put $z = H_3(v|_{[\tau_1,\tau_2]})$. The fact that for all $x, y \in \mathbb{R}$

$$\cos(x) - \cos(y) = 2\sin(\frac{y-x}{2})\sin(\frac{y+x}{2})$$

thus implies

$$\begin{split} \int_{C([0,T])} |F(w) - \mathrm{pr}_4(G(y,v))| \, Q_{y,v}(dw) \\ &\geq \gamma \int_{C_1} |\sin\big(H_1(\widetilde{w}) \cdot \psi(z)\big)| \, Q_{y,v}^{\widetilde{W}}(d\widetilde{w}) \cdot \int_{C_2} |\sin\big(H_2(b) \cdot \psi(z)\big)| \, Q_{y,v}^B(db) \\ &= \gamma \int_{\mathbb{R}} |\sin(y \cdot \psi(z))| \, Q_{y,v}^{H_1(\widetilde{W})}(dy) \cdot \int_{\mathbb{R}} |\sin(y \cdot \psi(z))| \, Q_{y,v}^{H_2(B)}(dy). \end{split}$$

Due to Lemma 2(i),(ii), $Q_{y,v}^{\widetilde{W}}$ and $Q_{y,v}^B$ are Gaussian measures on $\mathfrak{B}(C_1)$ and $\mathfrak{B}(C_2)$, respectively. Since the mappings H_1 and H_2 are linear and continuous we conclude that $Q_{y,v}^{H_1(\widetilde{W})}$ and $Q_{y,v}^{H_2(B)}$ are Gaussian measures on $\mathfrak{B}(\mathbb{R})$. Let m_1, σ_1^2 and m_2, σ_2^2 denote the mean and the variance of $Q_{y,v}^{H_1(\widetilde{W})}$ and $Q_{y,v}^{H_2(B)}$, respectively. Applying Lemma 3 we obtain

$$\int_{\mathbb{R}} |\sin(y \cdot \psi(z))| \, Q_{y,v}^{H_1(\widetilde{W})}(dy) \ge \frac{\exp\left(-\frac{\pi^2}{8}\right)}{\sqrt{8\pi}} \cdot \mathbb{1}_{[1,\infty)}(\sigma_1 \cdot \psi(z))$$

as well as

$$\int_{\mathbb{R}} |\sin(y \cdot \psi(z))| Q_{y,v}^{H_2(B)}(dy) \ge \frac{\exp\left(-\frac{\pi^2}{8}\right)}{\sqrt{8\pi}} \cdot 1_{[1,\infty)}(\sigma_2 \cdot \psi(z)).$$

Hence

(28)
$$\int_{C([0,T])} |F(w) - \operatorname{pr}_4(G(y,v))| Q_{y,v}(dw) \ge \frac{\gamma \exp\left(-\frac{\pi^2}{4}\right)}{8\pi} \cdot \mathbb{1}_{[1,\infty)}\left(\min(\sigma_1, \sigma_2) \cdot \psi(z)\right).$$

Next we derive lower bounds for σ_1^2 and σ_2^2 . Due to Lemma 2(ii) we have

$$m_2 = \int_{C_2} H_2(b) Q_{y,v}^B(db) = \int_{t_0}^{t_1} f'(t) \cdot m(Q_{y,v}^B)(t) dt = 0$$

as well as

$$\sigma_2^2 = \int_{C_2} (H_2(b) - m_2)^2 Q_{y,v}^B(db) = \int_{t_0}^{t_1} \int_{t_0}^{t_1} f'(r) \cdot f'(t) \cdot R(Q_{y,v}^B)(r,t) dr dt$$
$$= \int_{t_0}^{t_1} \int_{t_0}^{t_1} f'(r) \cdot f'(t) \cdot \frac{(t_1 - \max(r,t)) \cdot (\min(r,t) - t_0)}{t_1 - t_0} dr dt.$$

It is easy to see that for all $a, b \in \mathbb{R}$ with a < b

(29)
$$\int_{a}^{b} \int_{a}^{b} \frac{(b - \max(r, t)) \cdot (\min(r, t) - a)}{b - a} dr dt = \frac{(b - a)^{3}}{12}$$

Moreover, the assumption $\inf_{t \in [0, \tau_1/2]} |f'(t)| > 0$ implies that for all $r, t \in [0, \tau_1/2]$

(30)
$$f'(r) \cdot f'(t) = |f'(r) \cdot f'(t)| \ge \alpha$$

Observing (27) we thus obtain

(31)
$$\sigma_2^2 \ge \frac{\alpha (t_1 - t_0)^3}{12} \ge \frac{\alpha \delta^3}{12(n+1)^3}.$$

Put $s_i = s_{\pi(i)}^{y,v}$ for $i = 0, \ldots, n+1$ as well as $s_{n+2} = \tau_1$. Clearly,

$$\begin{split} m_1 &= \int_{C_1} H_1(\widetilde{w}) \, Q_{y,v}^{\widetilde{W}}(d\widetilde{w}) = \sum_{i \in \{0,\dots,n+1\} \setminus \{i^*\}} \int_{s_i}^{s_{i+1}} f'(t) m(Q_{y,v}^{\widetilde{W}})(t) dt \\ &+ \int_{t_0}^{t_1} f'(t) \Big(\frac{t-t_0}{t_1-t_0} m(Q_{y,v}^{\widetilde{W}})(t_1) + \frac{t_1-t}{t_1-t_0} m(Q_{y,v}^{\widetilde{W}})(t_0) \Big) dt \end{split}$$

Hence

$$\begin{split} \sigma_{1}^{2} &= \int_{C_{1}} (H_{1}(\widetilde{w}) - m_{1})^{2} Q_{y,v}^{\widetilde{W}}(d\widetilde{w}) \\ &= \int_{C_{1}} \left(\sum_{i \in \{0, \dots, n+1\} \setminus \{i^{*}\}} \int_{s_{i}}^{s_{i+1}} f'(t) \cdot (\widetilde{w}(t) - m(Q_{y,v}^{\widetilde{W}})(t)) dt \\ &+ \int_{t_{0}}^{t_{1}} f'(t) \cdot \frac{t - t_{0}}{t_{1} - t_{0}} \cdot (\widetilde{w}(t_{1}) - m(Q_{y,v}^{\widetilde{W}})(t_{1})) dt \\ &+ \int_{t_{0}}^{t_{1}} f'(t) \cdot \frac{t_{1} - t}{t_{1} - t_{0}} \cdot (\widetilde{w}(t_{0}) - m(Q_{y,v}^{\widetilde{W}})(t_{0})) dt \right)^{2} Q_{y,v}^{\widetilde{W}}(d\widetilde{w}). \end{split}$$

Lemma 2(i) implies that $R(Q_{y,v}^{\widetilde{W}})(r,t) = 0$ for all $r \in [s_i, s_{i+1}]$ and $t \in [s_j, s_{j+1}]$ and all $i, j \in \{0, \ldots, n\}$ with $i \neq j$ as well as $R(Q_{y,v}^{\widetilde{W}})(r,t) = 0$ for all $r \in [0, \tau_1]$ and $t \in \{t_0, t_1\} \cup [s_{n+1}, s_{n+2}]$. Thus

$$\sigma_1^2 = \sum_{i \in \{0,\dots,n\} \setminus \{i^*\}} \int_{s_i}^{s_{i+1}} \int_{s_i}^{s_{i+1}} f'(r) \cdot f'(t) \cdot R(Q_{y,v}^{\widetilde{W}})(r,t) dr dt$$
$$= \sum_{i \in \{0,\dots,n\} \setminus \{i^*\}} \int_{s_i}^{s_{i+1}} \int_{s_i}^{s_{i+1}} f'(r) \cdot f'(t) \cdot \frac{(s_{i+1} - \max(r,t)) \cdot (\min(r,t) - s_i)}{s_{i+1} - s_i} dr dt.$$

Using (30), (29) and the Hölder inequality we therefore obtain

$$\sigma_1^2 \ge \sum_{i \in \{0, \dots, n\} \setminus \{i^*\}} \int_{s_i}^{s_{i+1}} \int_{s_i}^{s_{i+1}} |f'(r) \cdot f'(t)| \cdot \frac{(s_{i+1} - \max(r, t)) \cdot (\min(r, t) - s_i)}{s_{i+1} - s_i} dr dt$$
$$\ge \frac{\alpha}{12} \sum_{i \in \{0, \dots, n\} \setminus \{i^*\}} (s_{i+1} - s_i)^3$$
$$\ge \frac{\alpha}{12n^2} \cdot \left(\sum_{i \in \{0, \dots, n\} \setminus \{i^*\}} (s_{i+1} - s_i)\right)^3 = \frac{\alpha}{12n^2} \cdot (\delta - (t_1 - t_0))^3.$$

The assumption that \widehat{X} uses the evaluation site $\delta/2$ implies that $t_1 - t_0 \leq \delta/2$. Hence

(32)
$$\sigma_1^2 \ge \frac{\alpha \delta^3}{96n^2}.$$

The desired lower bound (26) follows from (28), (31) and (32).

We conclude from (24),(25) (26) and (21) that

(33)
$$\mathbb{E}|X_4^{\psi}(T) - \widehat{X}_4| \ge \frac{\gamma \exp\left(-\frac{\pi^2}{4}\right)}{8\pi} \cdot \mathbb{P}\left(\left\{\psi(X_3^{\psi}(\tau_2)) \ge \sqrt{\frac{96\nu^3}{\alpha\delta^3}}\right\}\right).$$

Put

$$A = \left\{ \psi(X_3^{\psi}(\tau_2)) \ge \left(1 + \sqrt{\frac{96}{\alpha\delta^3}}\right) N^3 \right\}, \quad B = \{\nu \le N^2\}.$$

Clearly,

(34)
$$\mathbb{P}\Big(\Big\{\psi(X_3^{\psi}(\tau_2)) \ge \sqrt{\frac{96\nu^3}{\alpha\delta^3}}\Big\}\Big) \ge \mathbb{P}\Big(\Big\{\psi(X_3^{\psi}(\tau_2)) \ge \sqrt{\frac{96\nu^3}{\alpha\delta^3}}\Big\} \cap \{\nu \le N^2\}\Big) \ge \mathbb{P}(A \cap B) \ge \mathbb{P}(A) - \mathbb{P}(B^c).$$

Note that $X_3^{\psi}(\tau_2) \sim \mathcal{N}(0,\beta)$, where β is given by (8). Moreover, the assumption that ψ is continuous with $\lim_{x\to\infty} \psi(x) = \infty$ and $1 \in \psi(\mathbb{R})$ ensures that

$$\left(1 + \sqrt{\frac{96}{\alpha\delta^3}}\right)N^3 \in \psi(\mathbb{R}).$$

Hence

(35)

$$\mathbb{P}(A) = \mathbb{P}\left(\left\{X_3^{\psi}(\tau_2) \ge \psi^{-1}\left(\left(1 + \sqrt{\frac{96}{\alpha\delta^3}}\right)N^3\right)\right\}\right)$$
$$\ge \frac{1}{\sqrt{2\pi\beta}} \int_{\psi^{-1}\left(\left(1 + \sqrt{\frac{96}{\alpha\delta^3}}\right)N^3\right)}^{1+\psi^{-1}\left(\left(1 + \sqrt{\frac{96}{\alpha\delta^3}}\right)N^3\right)} \exp\left(-\frac{x^2}{2\beta}\right) dx$$
$$\ge \frac{1}{\sqrt{2\pi\beta}} \cdot \exp\left(-\frac{1}{2\beta} \cdot \left(1 + \psi^{-1}\left(\left(1 + \sqrt{\frac{96}{\alpha\delta^3}}\right)N^3\right)\right)^2\right)$$
$$\ge \frac{\exp\left(-\frac{1}{\beta}\right)}{\sqrt{2\pi\beta}} \cdot \exp\left(-\frac{1}{\beta} \cdot \left(\psi^{-1}\left(\left(1 + \sqrt{\frac{96}{\alpha\delta^3}}\right)N^3\right)\right)^2\right).$$

By the Markov inequality and the fact that $\widehat{X} \in \mathcal{X}_{N+1}^{\delta}$,

(36)
$$\mathbb{P}(B^c) \le \frac{\mathbb{E}\nu}{N^2} \le \frac{N+1}{N^2} \le \frac{2}{N}.$$

Estimates (33), (34), (35) and (36) imply (9), which completes the proof of the theorem in the case $\delta \in (0, \tau_1/2]$.

If $\delta \in (\tau_1/2, T]$ then the lower bound (9) follows from the fact that $\mathcal{X}_N^{\delta} \subseteq \mathcal{X}_N^{\tau_1/2}$ and the lower bound (9) in the case $\delta = \tau_1/2$.

Acknowledgement

I am grateful to Thomas Müller-Gronbach for stimulating discussions on the topic of this article.

References

- ALFONSI, A. Strong order one convergence of a drift implicit Euler scheme: Application to the CIR process. Statist. Probab. Lett. 83, 2 (2013), 602–607.
- [2] BERKAOUI, A., BOSSY, M., AND DIOP, A. Euler scheme for SDEs with non-Lipschitz diffusion coefficient: strong convergence. ESAIM Probab. Stat. 12 (2008), 1–11 (electronic).
- [3] BEYN, W.-J., ISAAK, E., AND KRUSE, R. Stochastic C-stability and B-consistency of explicit and implicit Euler-type schemes. J. Sci. Comput. (2015), 1–33.
- [4] BEYN, W.-J., ISAAK, E., AND KRUSE, R. Stochastic C-stability and B-consistency of explicit and implicit Milstein-type schemes. arXiv:1512.06905 (2016), 32 pages.
- [5] DEREICH, S., NEUENKIRCH, A., AND SZPRUCH, L. An Euler-type method for the strong approximation of the Cox-Ingersoll-Ross process. Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci. 468, 2140 (2012), 1105–1115.
- [6] GAINES, J. G., AND LYONS, T. J. Variable step size control in the numerical solution of stochastic differential equations. SIAM J. Appl. Math. 57, 5 (1997), 1455–1484.
- [7] GOBET, E., AND STAZHYNSKI, U. Optimal discretization of stochastic integrals for degenerate semimartingale. https://hal-polytechnique.archives- ouvertes.fr/hal-01241190 (2016), 31 pages.
- [8] GYÖNGY, I., AND RÁSONYI, M. A note on Euler approximations for SDEs with Hölder continuous diffusion coefficients. *Stochastic Process. Appl.* 121, 10 (2011), 2189–2200.
- [9] HAIRER, M., HUTZENTHALER, M., AND JENTZEN, A. Loss of regularity for Kolmogorov equations. Ann. Probab. 43, 2 (2015), 468–527.
- [10] HEFTER, M., AND HERZWURM, A. Optimal strong approximation of the one-dimensional squared Bessel process. arXiv:1601.01455 (2016), 20 pages.
- [11] HEFTER, M., AND HERZWURM, A. Strong convergence rates for Cox-Ingersoll-Ross processes full parameter range. arXiv:1608.00410 (2016), 18 pages.
- [12] HIGHAM, D. J., MAO, X., AND STUART, A. M. Strong convergence of Euler-type methods for nonlinear stochastic differential equations. SIAM J. Numer. Anal. 40, 3 (2002), 1041–1063 (electronic).
- [13] HOEL, H., VON SCHWERIN, E., SZEPESSY, A., AND TEMPONE, R. Adaptive multilevel Monte Carlo simulation. In Numerical analysis of multiscale computations, vol. 82 of Lect. Notes Comput. Sci. Eng. Springer, Heidelberg, 2012, pp. 217–234.
- [14] HOEL, H., VON SCHWERIN, E., SZEPESSY, A., AND TEMPONE, R. Implementation and analysis of an adaptive multilevel Monte Carlo algorithm. *Monte Carlo Methods Appl. 20*, 1 (2014), 1–41.
- [15] HOFMANN, N., MÜLLER-GRONBACH, T., AND RITTER, K. The optimal discretization of stochastic differential equations. J. Complexity 17, 1 (2001), 117–153.
- [16] HU, Y. Semi-implicit Euler-Maruyama scheme for stiff stochastic equations. In Stochastic analysis and related topics, V (Silivri, 1994), vol. 38 of Progr. Probab. Birkhäuser Boston, Boston, MA, 1996, pp. 183–202.
- [17] HUTZENTHALER, M., AND JENTZEN, A. On a perturbation theory and on strong convergence rates for stochastic ordinary and partial differential equations with non-globally monotone coefficients. arXiv:1401.0295 (2014), 41 pages.
- [18] HUTZENTHALER, M., JENTZEN, A., AND KLOEDEN, P. E. Strong convergence of an explicit numerical method for SDEs with non-globally Lipschitz continuous coefficients. Ann. Appl. Probab. 22, 4 (2012), 1611–1641.
- [19] HUTZENTHALER, M., JENTZEN, A., AND NOLL, M. Strong convergence rates and temporal regularity for Cox-Ingersoll-Ross processes and Bessel processes with accessible boundaries. arXiv: (2014), 32 pages.
- [20] JENTZEN, A., MÜLLER-GRONBACH, T., AND YAROSLAVTSEVA, L. On stochastic differential equations with arbitrary slow convergence rates for strong approximation. *Commun. Math. Sci.* 14, 7 (2016), 1477–1500.
- [21] KUMAR, C., AND SABANIS, S. On Milstein approximations with varying coefficients: the case of super-linear diffusion coefficients. arXiv:1601.02695 (2016), 31 pages.
- [22] LAMBA, H., MATTINGLY, J. C., AND STUART, A. M. An adaptive Euler-Maruyama scheme for SDEs: convergence and stability. *IMA J. Numer. Anal.* 27, 3 (2007), 479–506.
- [23] LEOBACHER, G., AND SZÖLGYENYI, M. A strong order 1/2 method for multidimensional SDEs with discontinuous drift. arXiv:1512.02807 (2015), 26 pages.

- [24] LEOBACHER, G., AND SZÖLGYENYI, M. A numerical method for SDEs with discontinuous drift. BIT 56, 1 (2016), 151–162.
- [25] MAO, X., AND SZPRUCH, L. Strong convergence rates for backward Euler-Maruyama method for non-linear dissipative-type stochastic differential equations with super-linear diffusion coefficients. *Stochastics* 85, 1 (2013), 144–171.
- [26] MARUYAMA, G. Continuous Markov processes and stochastic equations. Rend. Circ. Mat. Palermo (2) 4 (1955), 48–90.
- [27] MOON, K.-S., SZEPESSY, A., TEMPONE, R., AND ZOURARIS, G. E. Convergence rates for adaptive weak approximation of stochastic differential equations. *Stoch. Anal. Appl.* 23, 3 (2005), 511–558.
- [28] MÜLLER-GRONBACH, T. The optimal uniform approximation of systems of stochastic differential equations. Ann. Appl. Probab. 12, 2 (2002), 664–690.
- [29] MÜLLER-GRONBACH, T. Strong approximation of systems of stochastic differential equations. Habilitation thesis, TU Darmstadt (2002), iv+161.
- [30] MÜLLER-GRONBACH, T. Optimal pointwise approximation of SDEs based on Brownian motion at discrete points. Ann. Appl. Probab. 14, 4 (2004), 1605–1642.
- [31] MÜLLER-GRONBACH, T., AND YAROSLAVTSEVA, L. On hard quadrature problems for marginal distributions of SDEs with bounded smooth coefficients. arXiv:1603.08686 (2016), 38 pages.
- [32] NEUENKIRCH, A., AND SZPRUCH, L. First order strong approximations of scalar SDEs defined in a domain. Numerische Mathematik (2014), 1–34.
- [33] NGO, H.-L., AND TAGUCHI, D. Strong rate of convergence for the Euler-Maruyama approximation of stochastic differential equations with irregular coefficients. *Math. Comp.* 85, 300 (2016), 1793–1819.
- [34] RÖMISCH, W., AND WINKLER, R. Stepsize control for mean-square numerical methods for stochastic differential equations with small noise. SIAM J. Sci. Comput. 28, 2 (2006), 604–625 (electronic).
- [35] SABANIS, S. A note on tamed Euler approximations. *Electron. Commun. Probab.* 18 (2013), 1–10.
- [36] TRAUB, J. F., WASILKOWSKI, G., AND WOŹNIAKOWSKI, H. Information-based complexity. Boston, MA: Academic Press, Inc., 1988.
- [37] TRETYAKOV, M., AND ZHANG, Z. A fundamental mean-square convergence theorem for SDEs with locally Lipschitz coefficients and its applications. SIAM J. Numer. Anal. 51, 6 (2013), 3135–3162.
- [38] WANG, X., AND GAN, S. The tamed Milstein method for commutative stochastic differential equations with non-globally Lipschitz continuous coefficients. J. Difference Equ. Appl. 19, 3 (2013), 466–490.

FAKULTÄT FÜR INFORMATIK UND MATHEMATIK, UNIVERSITÄT PASSAU, INNSTRASSE 33, 94032 PASSAU, GER-MANY

E-mail address: larisa.yaroslavtseva@uni-passau.de