FORBIDDING INDUCED EVEN CYCLES IN A GRAPH: TYPICAL STRUCTURE AND COUNTING

JAEHOON KIM, DANIELA KÜHN, DERYK OSTHUS, TIMOTHY TOWNSEND

ABSTRACT. We determine, for all $k \geq 6$, the typical structure of graphs that do not contain an induced 2k-cycle. This verifies a conjecture of Balogh and Butterfield. Surprisingly, the typical structure of such graphs is richer than that encountered in related results. The approach we take also yields an approximate result on the typical structure of graphs without an induced 8-cycle or without an induced 10-cycle.

1. Introduction

1.1. **Background.** The enumeration and description of the typical structure of graphs with given side constraints has become a successful and popular area at the interface of probabilistic, enumerative, and extremal combinatorics (see e.g. [7] for a survey of such work). For example, a by now classical result of Erdős, Kleitman and Rothschild [12] shows that almost all triangle-free graphs are bipartite (given a fixed graph H, a graph is called H-free if it does not contain H as a not necessarily induced subgraph). This result was generalised to K_k -free graphs by Kolaitis, Prömel and Rothschild [14]. There are now many precise results on the number and typical structure of H-free graphs and more generally graphs, hypergraphs and other combinatorial structures with a given (anti-)monotone property.

Given a fixed graph H, a graph is called induced-H-free if it does not contain H as an induced subgraph. Associated counting and structural questions are equally natural as in the non-induced case, but seem harder to solve. Thus much less is known about the typical structure and number of induced-H-free graphs than that of H-free graphs, though considerable work has been done in this area (see, e.g. [2, 4, 13, 18, 19, 20]). In particular, Prömel and Steger [20] obtained an asymptotic counting result for the number of induced-H-free graphs on n vertices, showing that the logarithm of this number is essentially determined by the so-called colouring number of H. This was generalised to arbitrary hereditary properties independently by Alekseev [1] as well as Bollobás and Thomason [8]. Recent exciting developments in [5, 21] have opened up the opportunity to replace counting results by more precise results which identify the typical asymptotic structure.

In this paper we determine the typical structure of induced- C_{2k} -free graphs (from which the corresponding asymptotic counting result follows immediately). The key difficulty we encounter is that the typical structure turns out to be more complex than encountered in previous results on forbidden induced subgraphs. This requires new ideas and a more intricate analysis when 'excluding' classes of graphs which might be candidates for typical induced- C_{2k} -free graphs.

1.2. Graphs with forbidden induced cycles. Given a class of graphs \mathcal{A} , we let \mathcal{A}_n denote the set of all graphs in \mathcal{A} that have precisely n vertices, and we say that almost all graphs in \mathcal{A} have property \mathcal{B} if

$$\lim_{n\to\infty}\frac{|\{G\in\mathcal{A}_n:G\text{ has property }\mathcal{B}\}|}{|\mathcal{A}_n|}=1.$$

Date: August 14, 2018.

The research leading to these results was partially supported by the European Research Council under the European Union's Seventh Framework Programme (FP/2007-2013) / ERC Grant Agreements no. 258345 (D. Kühn) and 306349 (J. Kim and D. Osthus).

Given graphs H_1, \ldots, H_m , we say G can be covered by H_1, \ldots, H_m if V(G) admits a partition $A_1 \cup \cdots \cup A_m = V(G)$ such that $G[A_i]$ is isomorphic to H_i for every $i \in \{1, \ldots, m\}$.

Prömel and Steger proved in [18] that almost all induced- C_4 -free graphs can be covered by a clique and an independent set, and in [17] characterised the structure of almost all induced- C_5 -free graphs too. More recently, Balogh and Butterfield [4] determined the typical structure of induced-H-free graphs for a wide class of graphs H. In particular they proved that almost all induced- C_7 -free graphs can be covered by either three cliques or two cliques and an independent set, and that for $k \geq 4$ almost all induced- C_{2k+1} -free graphs can be covered by k cliques. They also conjectured that for $k \geq 6$ almost all induced- C_{2k} -free graphs can be covered by k-2 cliques and a graph whose complement is a disjoint union of stars and triangles. Our main result completely verifies this conjecture.

Theorem 1.1. For $k \geq 6$, almost all induced- C_{2k} -free graphs can be covered by k-2 cliques and a graph whose complement is a disjoint union of stars and triangles.

Theorem 1.1 together with the discussed results in [4, 12, 17, 18] implies that the typical structure of induced- C_k -free graphs is determined for every $k \in \mathbb{N}$ apart from $k \in \{6, 8, 10\}$. For the cases k = 8 and k = 10 the methods we use to prove Theorem 1.1 allow us to also prove an approximate result on the typical structure of induced- C_k -free graphs. In order to state this result we require the following definitions.

Given $\eta > 0$ and graphs G and G' on the same vertex set, we say G' is η -close to G if G' can be made into G by changing (i.e. adding or deleting) at most $\eta |G|^2$ edges. We say a graph G is a sun if either G consists of a single vertex or V(G) can be partitioned into sets A, B such that $E(G) = \{uv : |\{u,v\} \cap B| \leq 1\}$. We call A the body of the sun and B the side of the sun. Note that all stars and cliques (including triangles) are suns, and that we consider a single vertex to be both a star of order one and a clique of order one.

Theorem 1.2.

- (i) For every $\eta > 0$, almost all induced C_{10} -free graphs are η -close to graphs that can be covered by three cliques and a graph whose complement is a disjoint union of cliques.
- (ii) For every $\eta > 0$, almost all induced C_8 -free graphs are η -close to graphs that can be covered by two cliques and a graph whose complement is a disjoint union of suns.

We remark that in Theorems 1.1 and 1.2 we get exponential bounds on the proportion of induced- C_{2k} -free graphs that do not satisfy the relevant structural description. Our proofs also show that the k-2 cliques in the covering have size close to n/(k-1) in Theorem 1.1, with analogous bounds in Theorem 1.2. Theorem 1.1 also strengthens a result by Kang, McDiarmid, Reed and Scott [13] showing that almost all induced- C_{2k} -free graphs have a linear sized homogeneous set. (Their results were motivated by the Erdős-Hajnal conjecture, and actually apply to a large class of forbidden graphs H.)

It would of course be interesting to determine the typical structure of induced- C_6 -free graphs.

Question 1.3. What is the typical structure of induced- C_6 -free graphs?

It seems likely that almost all induced- C_6 -free graphs can be covered by one clique and one cograph, where a cograph is a graph not containing an induced copy of P_4 . Another natural question is that of the typical structure of induced-H-free graphs of a given density. In particular, an intriguing question is whether their typical structure exhibits a non-trivial 'phase transition' as found for triangle-free graphs [16] and more generally K_r -free graphs [6].

1.3. Overview of the paper. A key tool in our proofs is the recent hypergraph container approach, which was developed independently by Balogh, Morris and Samotij [5], and Saxton and Thomason [21]. Briefly, their result states that under suitable conditions on a uniform hypergraph

G, there is a small collection \mathcal{C} of small subsets (known as containers) of V(G) such that every independent set of vertices in G is a subset of some element of \mathcal{C} . The precise statement of the application used here is deferred until Section 3.

Given a graph G and a set $A \subseteq V(G)$, we denote by G[A] the graph induced on G by A, and we denote the complement of G by \overline{G} . For $k \in \mathbb{N}$ and a set V of vertices we define an *ordered* k-partition of V to be a k-partition of V such that one partition class is labelled and the rest are unlabelled. If Q is an ordered k-partition with labelled class Q_0 and unlabelled classes Q_1, \ldots, Q_{k-1} then we write $Q = (Q_0, \{Q_1, \ldots, Q_{k-1}\})$.

For $k \geq 4$, we say that a graph G is a k-template if V(G) has an ordered (k-1)-partition $Q = (Q_0, \{Q_1, \ldots, Q_{k-2}\})$ such that $G[Q_i]$ is a clique for all $i \in [k-2]$ and one of the following holds.

- k = 4 and $\overline{G}[Q_0]$ is a disjoint union of suns.
- k = 5 and $\overline{G}[Q_0]$ is a disjoint union of stars and cliques.
- $k \ge 6$ and $\overline{G}[Q_0]$ is a disjoint union of stars and triangles.

Clearly every k-template is induced- C_{2k} -free. If V(G) has such an ordered (k-1)-partition Q, we say that G is a k-template on Q, or G has ordered (k-1)-partition Q. If Q' is the (unordered) (k-1)-partition with the same partition classes as Q, we may also say that G is a k-template on Q'. Thus Theorem 1.1 can be reformulated as:

'For $k \geq 6$, almost all induced C_{2k} -free graphs are k-templates.'

Theorem 1.2 can be similarly reformulated in terms of 4- and 5-templates. As mentioned earlier, the main difficulty in proving Theorem 1.1 (compared to related results) is that typically $G[Q_0]$ is close to, but not quite, a complete graph. This makes it very difficult to rule out other similar classes of graphs as typical structures. To overcome this we use tools such as Ramsey's theorem to classify the graphs according to the neighbourhoods of certain vertices.

More precisely, our approach to proving our main result is as follows. Firstly, in Section 3 we use the hypergraph containers result discussed above to show that almost all induced- C_{2k} -free graphs are close to being a k-template, for every $k \geq 4$ (see Lemma 3.1). Note that Lemma 3.1 immediately implies Theorem 1.2.

In Section 4 we prove upper and lower bounds on the number of k-templates on n vertices (see Lemmas 4.4 and 4.6). In Section 5 we prove some preliminary results about graphs that are close to being a k-template.

In Section 6 we state a key result which is a version of Theorem 1.1 with respect to a given ordered (k-1)-partition (see Lemma 6.1) and use it together with Lemma 4.6 to derive Theorem 1.1. The remainder of the paper is devoted to proving Lemma 6.1 via an inductive argument, which we introduce at the end of Section 6. This argument involves partitioning the class of graphs considered in Lemma 6.1 into three 'bad' classes of graphs, and in each of Sections 7, 8 and 9 we use Lemma 4.4 and the results in Section 5 to prove an upper bound on the number of graphs in a different one of these classes (see Lemmas 7.3, 8.7 and 9.8). In particular, Lemmas 7.3 and 8.7 already show that almost all induced- C_{2k} -free graphs are 'extremely close' to being k-templates (see Proposition 9.1). Finally in Section 10 we use Lemmas 3.1, 7.3, 8.7 and 9.8 to complete the inductive argument set up in Section 6 and so prove Lemma 6.1. Before starting on any of this however, we lay out some notation and set out some useful tools in Section 2, below.

2. Notation and tools

Given a graph G, a vertex $x \in V(G)$, and an ordered (k-1)-partition $Q = (Q_0, \{Q_1, \dots, Q_{k-2}\})$ of V(G), we let $N(x), \overline{N}(x)$ denote the set of neighbours and non-neighbours of x in G, respectively. We also let $N_{Q_i}(x), \overline{N}_{Q_i}(x)$ denote the set of neighbours of x in Q_i and non-neighbours of x in Q_i ,

respectively. We sometimes use the notation $d_{G,Q}^i(x) = |N_{Q_i}(x)|$ and $\overline{d}_{G,Q}^i(x) = |\overline{N}_{Q_i}(x)|$ when we want to emphasise which graph we are working with. For a set A of vertices in G, we define

$$N(A) := \bigcap_{v \in A} N(v), \qquad \overline{N}(A) := \bigcap_{v \in A} \overline{N}(v),$$

$$N_{Q_i}(A) := \bigcap_{v \in A} N_{Q_i}(v), \qquad \overline{N}_{Q_i}(A) := \bigcap_{v \in A} \overline{N}_{Q_i}(v).$$

If it generates no ambiguity, we may write $N_i(x)$, $\overline{N}_i(x)$, $N_i(A)$, $\overline{N}_i(A)$ for $N_{Q_i}(x)$, $\overline{N}_{Q_i}(x)$, $N_{Q_i}(A)$, and $\overline{N}_{Q_i}(A)$ respectively. Given $A, B \subseteq V(G)$, we define

$$N^*(A, B) := N(A) \cap \overline{N}(B), \quad N_i^*(A, B) := N_i(A) \cap \overline{N}_i(B).$$

In the case when A and B both have size one, containing vertices a, b respectively, we may write $N^*(a,b)$ for $N^*(A,B)$ and $N_i^*(a,b)$ for $N_i^*(A,B)$.

We say that a partition of vertices is balanced if the sizes of any two partition classes differ by at most one. Given a (k-1)-partition Q of [n] with partition classes Q_0, \ldots, Q_{k-2} , and a graph G = (V, E) on vertex set [n], and an edge or non-edge e = uv with $u \in Q_i$ and $v \in Q_j$, we call ecrossing if $i \neq j$ and internal if i = j.

We denote a path on m vertices by P_m . Given a path $P = p_1 \dots p_m$ and a sequence A_1, \dots, A_m of sets of vertices, we say that P has type A_1, \ldots, A_m if $p_\ell \in A_\ell$ for every $\ell \in [m]$. We call a graph a linear forest if it is a forest such that all components are paths or isolated vertices.

Given $\ell, t \in \mathbb{N}$ we let $R_{\ell}(t)$ denote the ℓ -colour Ramsey number for monochromatic t-cliques, i.e. $R_{\ell}(t)$ is the smallest $N \in \mathbb{N}$ such that every ℓ -colouring of the edges of K_N yields a monochromatic copy of K_t .

We define

$$n_k := \left\lceil \frac{n}{k-1} \right\rceil.$$

In a number of our proofs we shall use the following Chernoff bound.

Lemma 2.1 (Chernoff bound). Let X have binomial distribution and let $0 < a \leq \mathbb{E}[X]$. Then

(i)
$$P(X > \mathbb{E}[X] + a) \le \exp\left(-\frac{a^2}{4\mathbb{E}[X]}\right)$$
.
(ii) $P(X < \mathbb{E}[X] - a) \le \exp\left(-\frac{a^2}{2\mathbb{E}[X]}\right)$.

(ii)
$$P(X < \mathbb{E}[X] - a) \le \exp\left(-\frac{a^2}{2\mathbb{E}[X]}\right)$$

Whenever this does not affect the argument, we assume all large numbers to be integers, so that we may sometimes omit floors and ceilings for the sake of clarity. In some proofs, given $a, b \in \mathbb{R}$ with 0 < a, b < 1, we will use the notation $a \ll b$ to mean that we can find an increasing function g for which all of the conditions in the proof are satisfied whenever $a \leq g(b)$. Throughout we write $\log x$ to mean $\log_2 x$.

We define $\xi(p) := -3p(\log p)/2$. The following bounds will prove useful to us. For $n \geq 1$ and $3\log n/n \le p \le 10^{-11}$,

(2.1)
$$\binom{n}{\leq pn} := \sum_{i=0}^{\lfloor pn \rfloor} \binom{n}{i} \leq pn \left(\frac{en}{pn}\right)^{pn} \leq 2^{\xi(p)n},$$

and

(2.2)
$$\xi(p) \le \frac{3}{2}p\left(\frac{1}{p}\right)^{1/8} \le p^{3/4}.$$

3. Approximate structure of typical induced- C_{2k} -free graphs

The main result of this section is Lemma 3.1, which approximately determines the typical structure of induced- C_{2k} -free graphs. As mentioned earlier, we make use of a 'container theorem' which reduces the proof of Lemma 3.1 to an extremal problem involving induced- C_{2k} -free graphs. More precisely, the argument is structured as follows.

We first introduce a number of tools (see Subsection 3.1): a 'Containers' theorem (Theorem 3.2), a Stability theorem (Theorem 3.3), and two Removal Lemmas (Theorem 3.4, Lemma 3.5). In Subsection 3.2 we use Theorem 3.3 to derive a Stability result involving induced- C_{2k} -free graphs (Lemma 3.7). Similarly we use Theorem 3.4 to derive another specialised version of the Removal Lemma (Lemma 3.9). In Subsection 3.3 we use Theorem 3.2 together with Lemmas 3.5, 3.7 and 3.9 to determine the approximate structure of typical induced- C_{2k} -free graphs.

We denote the number of (labelled) induced- C_{2k} -free graphs on n vertices by F(n,k).

Lemma 3.1. Let $k \geq 4$. For every $\eta > 0$ there exists $\varepsilon > 0$ such that the following holds for all sufficiently large n. All but at most $F(n,k)2^{-\varepsilon n^2}$ induced- C_{2k} -free graphs on n vertices can be made into a k-template by changing at most ηn^2 edges.

Note that Lemma 3.1 immediately implies Theorem 1.2.

3.1. Tools: containers, stability and removal lemmas. The key tool in this section is Theorem 3.2, which is an application of the more general theory of Hypergraph Containers developed in [5, 21]. We use the formulation of Theorem 1.5 in [21]. We require the following definitions in order to state it.

A 2-coloured multigraph G on vertex set [N] is a pair of edge sets $G_R, G_B \subseteq [N]^{(2)}$, which we call the red and blue edge sets respectively. If H is a fixed graph on vertex set [h], a copy of H in G is an injection $f:[h] \to [N]$ such that for every edge uv of H, $f(u)f(v) \in G_R$, and for every non-edge u'v' of H, $f(u')f(v') \in G_B$. We write $H \subseteq G$ if G contains a copy of G, and we say that G is G is G if there are no copies of G in G. We say that G is G if G is G if G is G if G is G if G if

Theorem 3.2. Let H be a fixed graph with h := |V(H)|. For every $\varepsilon > 0$, there exists c > 0 such that for all sufficiently large N, there exists a collection C of complete 2-coloured multigraphs on vertex set [N] with the following properties.

- (a) For every graph I on [N] that contains no induced copy of H, there exists $G \in \mathcal{C}$ such that $I \subseteq G$.
- (b) Every $G \in \mathcal{C}$ contains at most εN^h copies of H.
- (c) $\log |\mathcal{C}| \le cN^{2-(h-2)/(\binom{h}{2}-1)} \log N$.

Another tool that we will use is the following classical Stability theorem of Erdős and Simonovits (see e.g. [10, 11, 22]). By $T_k(n)$ we denote the $Tur\'an\ graph$, the largest complete k-partite graph on n vertices, and we define $t_k(n) := e(T_k(n))$. Given a family $\mathcal H$ of fixed graphs, we say a graph G is $\mathcal H$ -free if G does not contain any $H \in \mathcal H$ as a (not necessarily induced) subgraph, and we say G is $induced\mathcal H$ -free if G does not contain any G as an induced subgraph.

Theorem 3.3. Let $\mathcal{H} = \{H_1, \dots, H_\ell\}$ be a family of fixed graphs, and let $k := \min_{1 \leq i \leq \ell} \chi(H_i)$. For every $\delta > 0$ there exists $\varepsilon > 0$ such that the following holds for all sufficiently large n. If a graph G on n vertices is \mathcal{H} -free and $e(G) \geq t_{k-1}(n) - \varepsilon n^2$, then G can be obtained from $T_{k-1}(n)$ by changing at most δn^2 edges.

The final tools that we introduce in this subsection are the following two Removal Lemmas. The first is an extension of the Induced Removal Lemma to families of forbidden graphs, and is due to Alon and Shapira [3]. The original statement of this theorem also applies to infinite families

of forbidden graphs, but the version for finite families is sufficient for our purposes. The second is a version of the Removal Lemma applicable to complete 2-coloured multigraphs. The proof is similar to that of the standard Removal Lemma, so we omit it here; for details see [23]. For two sets A, B, we denote their symmetric difference by $A\triangle B$. For 2-coloured multigraphs G, G' on the same vertex set we define their distance by $\operatorname{dist}(G, G') := |G_R \triangle G'_R| + |G_B \triangle G'_B|$.

- **Theorem 3.4.** [3] For every finite family of fixed graphs \mathcal{H} and every $\delta > 0$, there exists $\varepsilon > 0$ such that the following holds for all sufficiently large n. If a graph G on n vertices contains at most εn^h induced copies of each graph $H \in \mathcal{H}$ on h vertices, then G can be made induced- \mathcal{H} -free by changing at most δn^2 edges.
- **Lemma 3.5.** For every fixed graph H on h vertices, and every $\delta > 0$, there exists $\varepsilon > 0$ such that the following holds for all sufficiently large n. If a complete 2-coloured multigraph G on vertex set [n] contains at most εn^h copies of H, then there exists a complete 2-coloured multigraph G' on vertex set [n] such that G' is H-free and $\operatorname{dist}(G, G') \leq \delta n^2$.
- 3.2. Stability and removal lemmas for even cycles. Suppose H is a complete 2-coloured multigraph on m vertices with $H_R \cap H_B = \emptyset$. If m = 3 and $|H_R| \le 1$ we call H a mostly blue triangle. For $k \in \{4,5,6\}$, if m = 4 and $|H_R| \ge 6 k$ and H^B contains a copy of P_4 then we call H a k-good tetrahedron. The following technical proposition will be useful in proving Lemmas 3.7 and 3.9.

Proposition 3.6. Let $k \geq 4$ and let G be a complete 2-coloured multigraph on 2k vertices. If G satisfies one of the following properties then G contains a copy of C_{2k} . Below, r_i always denotes a red edge.

- (E1) $G_R \triangle G_B$ is a set of at most k disjoint (red or blue) edges.
- (E2) $G_R \triangle G_B$ is the edge set of two disjoint copies of a blue K_k .
- (E3) $G_R \triangle G_B$ is the edge set of a union of disjoint graphs $K_3^1, K_3^2, r_1, \ldots, r_{k-3}$, where each K_3^i is a mostly blue triangle.
- (E4) $G_R \triangle G_B$ is the edge set of a union of disjoint graphs $K_4^1, r_1, \ldots, r_{k-2}$, where K_4^1 is a 4-good tetrahedron.
- (E5) $k \geq 5$ and $G_R \triangle G_B$ is the edge set of a union of disjoint graphs $K_4^1, r_1, \ldots, r_{k-2}$, where K_4^1 is a 5-good tetrahedron.
- (E6) $k \geq 6$ and $G_R \triangle G_B$ is the edge set of a union of disjoint graphs $K_4^1, r_1, \ldots, r_{k-2}$, where K_4^1 is a 6-good tetrahedron.

Proof. Let $V(G) = \{v_1, \ldots, v_{2k}\}$. Let $C = c_1 \ldots c_{2k}$ be a 2k-cycle. Note that if there exists a permutation σ of [2k] such that for every edge $c_i c_j \in E(C)$ we have $v_{\sigma(i)} v_{\sigma(j)} \in G_R$ and such that for every non-edge $c_{i'} c_{j'} \notin E(C)$ we have $v_{\sigma(i')} v_{\sigma(j')} \in G_B$, then $v_{\sigma(1)} \ldots v_{\sigma(2k)}$ is a copy of C_{2k} in G. We call such a permutation σ a covering permutation from C to G. For ease of reading, we will write a permutation σ on [2k] using the notation $\sigma = (\sigma(1), \ldots, \sigma(2k))$. If σ restricted to $\{m, m+1, \ldots, 2k\}$ is the identity permutation, we may simply write $\sigma = \{\sigma(1), \ldots, \sigma(m-1)\}$ instead. So for example if $\sigma = (1, 3, 4, 2)$ is a covering permutation from C to G, then $v_1 v_3 v_4 v_2 v_5 \ldots v_{2k}$ is a copy of C_{2k} in G.

We now show that each of the properties (E1),...,(E6) imply that there exists a covering permutation from C to G, and hence that G contains a copy of C_{2k} .

- (E1) There exists $b, r \in \mathbb{N} \cup \{0\}$ with $b+r \leq k$ such that, by relabelling vertices if necessary, $G_B \setminus G_R = \{v_1v_2, \dots, v_{2b-1}v_{2b}\}$ and $G_R \setminus G_B = \{v_{2b+1}v_{2b+2}, \dots, v_{2(b+r)-1}v_{2(b+r)}\}$. Depending on the value of b we find the following covering permutations σ from C to G, as required.
 - If b = 0 then σ is the identity permutation.
 - If b = 1 then $\sigma = (1, 3, 4, 2)$.
 - If $b \ge 2$ then $\sigma = (1, 3, \dots, 2b 1, 2, 4, \dots, 2b)$.

- (E2) Let $\{v_1,\ldots,v_k\},\{v_{k+1},\ldots,v_{2k}\}$ be the respective vertex sets of the two copies of a blue K_k in $G_R \triangle G_B$. Then $\sigma = (1, k+1, 2, k+2, \dots, k, 2k)$ is a covering permutation from C to G, as required.
- (E3) Let $V(K_3^1) = \{v_1, v_2, v_3\}, V(K_3^2) = \{v_4, v_5, v_6\}$ and $V(r_i) = \{v_{2i+5}, v_{2i+6}\}$ for every $i \in [k-3]$. Depending on the colour of the edges in K_3^1, K_3^2 we find the following covering permutations σ from C to G, as required.
 - If K_3^1, K_3^2 both contain no red edges, then $\sigma = (1, 4, 2, 5, 3, 6)$.
 - If K_3^1 contains exactly one red edge v_1v_2 and K_3^2 contains no red edges, then $\sigma =$
 - If K_3^1 contains exactly one red edge v_1v_2 and K_3^2 contains exactly one red edge v_5v_6 , then $\sigma = (1, 2, 4, 3, 5, 6).$
- (E4) Let $V(K_4^1) = \{v_1, v_2, v_3, v_4\}$ and $V(r_i) = \{v_{2i+3}, v_{2i+4}\}$ for every $i \in [k-2]$. Depending on the configuration of red edges in K_4^1 we find the following covering permutations σ from C to G, as required.
 - If K_4^1 contains exactly three red edges v_1v_2, v_2v_3, v_3v_4 , then σ is the identity permutation.

 - If K_4^1 contains exactly two red edges v_1v_2, v_2v_3 , then $\sigma=(1,2,3,5,6,4)$. If K_4^1 contains exactly two red edges v_1v_2, v_3v_4 , then $\sigma=(1,2,5,6,3,4)$.
- (E5) We may assume that K_4^1 contains exactly one red edge, since that is the only case not covered by (E4). Let $V(K_4^1) = \{v_1, v_2, v_3, v_4\}$ and $V(r_i) = \{v_{2i+3}, v_{2i+4}\}$ for every $i \in [k-2]$, and let v_1v_2 be the red edge in K_4^1 . Then $\sigma=(1,2,5,6,3,7,8,4)$ is a covering permutation from C to G, as required.
- (E6) We may assume that K_4^1 contains no red edges, since that is the only case not covered by (E5). Let $V(K_4^1) = \{v_1, v_2, v_3, v_4\}$ and $V(r_i) = \{v_{2i+3}, v_{2i+4}\}$ for every $i \in [k-2]$. Then $\sigma = (1, 5, 6, 2, 7, 8, 3, 9, 10, 4)$ is a covering permutation from C to G, as required.

We now use Theorem 3.3 and Proposition 3.6 to prove the following more specialised Stability result involving C_{2k} -free 2-coloured multigraphs.

Lemma 3.7. Let $k \geq 4$. For every $\delta > 0$ there exists $\varepsilon > 0$ such that the following holds for all sufficiently large n. If a complete 2-coloured multigraph G on vertex set [n] is C_{2k} -free and $|G_R \cap G_B| \ge t_{k-1}(n) - \varepsilon n^2$, then the graph $([n], G_R \cap G_B)$ can be obtained from $T_{k-1}(n)$ by changing at most δn^2 edges.

Proof. Choose $n_0 \in \mathbb{N}$ and $\varepsilon > 0$ such that $1/n_0 \ll \varepsilon \ll \delta$. Let $n \geq n_0$. Since G is C_{2k} -free, we know by Proposition 3.6 that no 2k vertices of G induce on G a 2-coloured multigraph G' that satisfies (E1). So, since G is complete, the graph $([n], G_R \cap G_B)$ must be $T_k(2k)$ -free. Note that $\chi(T_k(2k)) = k$. By Theorem 3.3, this together with the fact that $|G_R \cap G_B| \ge t_{k-1}(n) - \varepsilon n^2$ implies that the graph $([n], G_R \cap G_B)$ can be obtained from $T_{k-1}(n)$ by changing at most δn^2 edges.

The following proposition characterises the structure of graphs without k-good tetrahedrons. It will be useful in proving Lemma 3.9. The proof is fairly straightforward so we give only a sketch of it here.

Proposition 3.8. Let G be a 2-coloured multigraph with $G_R \cap G_B = \emptyset$.

- (i) If G does not contain a 6-good tetrahedron then G^B is a disjoint union of stars and triangles.
- (ii) If G does not contain a 5-good tetrahedron then G^B is a disjoint union of stars and cliques.
- (iii) If G does not contain a 4-good tetrahedron then G^B is a disjoint union of suns.

Proof. (i) follows immediately from the fact that if G is 6-good tetrahedron-free then G^B does not contain a P_4 .

To see (ii), note that if G is 5-good tetrahedron-free and P is a copy of P_4 in G^B , then $G^B[V(P)] = K_4$. So every component H of G^B is either a star or a triangle or contains a K_4 . But in the latter case it is easy to check that H is actually a clique.

It remains to prove (iii). If G is 4-good tetrahedron-free and P is a copy of P_4 in G^B , then $G^B[V(P)]$ is either a K_4 or a copy of the graph K_4^- obtained from K_4 by deleting one edge. So every component H of G^B is either a star or a clique or contains an induced copy of K_4^- . Using induction on |H|, it is not hard to show that in the latter case H must be a sun.

We now use Theorem 3.4 together with Propositions 3.6 and 3.8 to prove the following more specialised Removal Lemma involving even cycles.

Lemma 3.9. For every $k \geq 4$ and every $\delta > 0$ there exists $\varepsilon > 0$ such that the following holds for all sufficiently large n. Suppose G is a complete 2-coloured multigraph on n vertices such that $G_R \cap G_B = E(T_{k-1}(n))$. Let Q be the unique (k-1)-partition of the vertices of G such that no partition class induces an edge in $G_R \cap G_B$. Suppose further that G contains at most εn^{2k} copies of C_{2k} . Then there exists a k-template $T = (V(G), E^T)$ on Q such that $|G_R \triangle E^T| \leq \delta n^2$.

Proof. We first prove the lemma in the case $k \geq 6$. Choose $n_0 \in \mathbb{N}$ and $\varepsilon, \gamma > 0$ such that $1/n_0 \ll \varepsilon \ll \gamma \ll \delta, 1/k$. Let $n \geq n_0$ and let $Q = (Q_1, \ldots, Q_{k-1})$. Let $c := \varepsilon^{1/3}$.

We claim that for no two distinct $i, j \in [k-1]$ do $G[Q_i]$ and $G[Q_j]$ both contain at least cn^k copies of a blue K_k . Indeed, if they do then there are at least $c^2n^{2k} > \varepsilon n^{2k}$ sets of 2k vertices that each induce on G a 2-coloured multigraph G' that satisfies (E2). By Proposition 3.6 each such G' contains a copy of C_{2k} . This contradicts the assumption that G contains at most εn^{2k} copies of C_{2k} , which proves the claim.

Thus there exists $J \subseteq [k-1]$ with $|J| \le 1$ such that for all $i \in [k-1]$ with $i \notin J$, $G[Q_i]$ contains fewer than cn^k copies of a blue K_k . Together with Theorem 3.4 (applied to $G^B[Q_i]$) this implies that $G[Q_i]$ can be made free of blue cliques of size k by changing the colour of at most γn^2 edges inside Q_i . So by Turán's Theorem, for all $i \in [k-1]$ with $i \notin J$, $G[Q_i]$ must have at least

$$(k-1)\binom{n/(k-1)^2}{2} - 2\gamma n^2 \ge \frac{n^2}{4(k-1)^3}$$

red edges.

Claim 1: There is at most one index $i \in [k-1]$ such that $G[Q_i]$ contains at least cn^3 mostly blue triangles. Moreover, if there is such an index i then $J \subseteq \{i\}$, and if there is no such index then $J = \emptyset$.

Indeed, suppose for a contradiction that there exist distinct $i, j \in [k-1]$ such that Q_i, Q_j both contain at least cn^3 mostly blue triangles. Note that any class that contains at least cn^k copies of a blue K_k must contain at least cn^3 mostly blue triangles. So we may assume that $J \subseteq \{i, j\}$. Thus for every index $\ell \neq i, j$, $G[Q_\ell]$ contains at least $n^2/(4(k-1)^3)$ red edges. Thus there are at least $2\varepsilon n^{2k}$ sets of 2k vertices that each induce on G a 2-coloured multigraph G' that satisfies (E3). (To see this, note that to choose such a set of 2k vertices we may choose, for both indices i, j, the vertices of any one of the at least cn^3 mostly blue triangles in $G[Q_i]$, $G[Q_j]$ respectively, and then choose, for each index $\ell \neq i, j$, any one of the at least $n^2/(4(k-1)^3)$ red edges in Q_ℓ .) By Proposition 3.6 each such G' contains a copy of C_{2k} . This contradicts the assumption that G contains at most εn^{2k} copies of C_{2k} , which proves the claim.

Let J' consist of the index $j_0 \in [k-1]$ such that $G[Q_{j_0}]$ contains at least cn^3 mostly blue triangles, if such an index exists. Otherwise let $J' := \emptyset$. Thus $J \subseteq J'$. For all $i \in [k-1]$ with $i \notin J'$, Claim 1 together with Theorem 3.4 (applied to $G^B[Q_i]$) implies that $G[Q_i]$ can be made free of mostly blue triangles by changing the colour of at most γn^2 edges inside Q_i . This implies that the blue edges inside Q_i after such a change form a matching. Hence $G[Q_i]$ contains at most $2\gamma n^2$ blue edges.

If $J' = \emptyset$ then $G[Q_i]$ contains at most $2\gamma n^2$ blue edges for all $i \in [k-1]$, and hence $|G_B \setminus G_R| \leq \delta n^2$ (since $\gamma \ll \delta, 1/k$). In this case we are done by setting T to be K_n . Otherwise, $J' = \{j_0\}$ and it suffices to show that the blue edges in $G[Q_{j_0}]$ can be made into the edge set of a disjoint collection of stars and triangles by changing the colour of at most γn^2 edges inside Q_{j_0} , since then we are done by setting T to be K_n minus this disjoint collection of stars and triangles.

Claim 2(a): $G[Q_{j_0}]$ contains fewer than cn^4 6-good tetrahedrons.

Indeed, otherwise there are at least $\varepsilon^{1/2}n^{2k}$ sets of 2k vertices that each induce on G a 2-coloured multigraph G' that satisfies (E6). (To see this, note that to choose such a set of 2k vertices we may first choose the vertices of any one of the at least cn^4 6-good tetrahedrons, and then choose, for each other class Q_i , any one of the at least $n^2/(4(k-1)^3)$ red edges in Q_i .) By Proposition 3.6 each such G' contains a copy of C_{2k} . This contradicts the assumption that G contains at most εn^{2k} copies of C_{2k} , which proves the claim.

Claim 2(a) together with Theorem 3.4 (applied to $G^B[Q_{j_0}]$) implies that $G[Q_{j_0}]$ can be made free of 6-good tetrahedrons by changing the colour of at most γn^2 edges inside Q_{j_0} . Proposition 3.8(i) implies that after such a change, all blue edges inside Q_{j_0} form a disjoint collection of stars and triangles, as required. This completes the proof in the case $k \geq 6$.

For the case k = 5, the proof is almost identical to the case $k \geq 6$, except that instead of Claim 2(a) we prove the following weaker claim, which follows in a similar way.

Claim 2(b): $G[Q_{i_0}]$ contains fewer than cn^4 5-good tetrahedrons.

Claim 2(b) together with Theorem 3.4 (applied to $G^B[Q_{j_0}]$) implies that $G[Q_{j_0}]$ can be made free of 5-good tetrahedrons by changing the colour of at most γn^2 edges inside Q_{j_0} . Proposition 3.8(ii) implies that after such a change, all blue edges inside Q_{j_0} form a disjoint collection of stars and cliques. We are now done by setting T to be K_n minus this disjoint collection of stars and cliques.

For the case k = 4, the proof is again almost identical to the case $k \ge 6$, except that instead of Claim 2(a) we prove the following even weaker claim, which follows in a similar way.

Claim 2(c): $G[Q_{j_0}]$ contains fewer than cn^4 4-good tetrahedrons.

Claim 2(c) together with Theorem 3.4 (applied to $G^B[Q_{j_0}]$) implies that $G[Q_{j_0}]$ can be made free of 4-good tetrahedrons by changing the colour of at most γn^2 edges inside Q_{j_0} . Proposition 3.8(iii) implies that after such a change, all blue edges inside Q_{j_0} form a disjoint collection of suns. We are now done by setting T to be K_n minus this disjoint collection of suns.

3.3. Approximate structure of typical induced C_{2k} -free graphs. We are now in a position to prove the main result of this section.

Proof of Lemma 3.1. Choose $n_0 \in \mathbb{N}$ and $\varepsilon, \delta, \gamma, \beta > 0$ such that $1/n_0 \ll \varepsilon \ll \delta \ll \gamma \ll \beta \ll \eta, 1/k$. Let $\varepsilon' := 2\varepsilon$ and $n \geq n_0$. First we claim that $F(n,k) \geq 2^{t_{k-1}(n)}$. To see this, first note that any graph G that contains $\overline{T_{k-1}(n)}$ is induced- C_{2k} -free (since for any set of 2k vertices on G, 3 of them must form a triangle). Moreover, there are precisely $2^{t_{k-1}(n)}$ such graphs for any given labelling of the vertices, which proves the claim.

By Theorem 3.2 (with C_{2k} , n and ε' taking the roles of H, N and ε respectively) there is a collection \mathcal{C} of complete 2-coloured multigraphs on vertex set [n] satisfying properties (a)–(c). In particular, by (a), every induced- C_{2k} -free graph on vertex set [n] is contained in some $G \in \mathcal{C}$. Let \mathcal{C}_1 be the family of all those $G \in \mathcal{C}$ for which $|G_R \cap G_B| \geq t_{k-1}(n) - \varepsilon' n^2$. Then the number of (labelled) induced- C_{2k} -free graphs not contained in some $G \in \mathcal{C}_1$ is at most

$$|\mathcal{C}| 2^{t_{k-1}(n)-\varepsilon'n^2} \le 2^{-\varepsilon n^2} F(n,k),$$

because $|\mathcal{C}| \leq 2^{n^{2-\varepsilon'}}$, by (c), and $F(n,k) \geq 2^{t_{k-1}(n)}$. We claim that for every $G \in \mathcal{C}_1$ there exists a complete 2-coloured multigraph \tilde{G} and a k-template T on partition $Q = \{Q_0, Q_1, \dots, Q_{k-2}\}$ such that

$$\tilde{G}_R \cap Q_i^{(2)} = E(T[Q_i])$$
 and $\tilde{G}_R \cap \tilde{G}_B \cap Q_i^{(2)} = \emptyset$

for every $i \in \{0, 1, \dots, k-2\}$, and $\operatorname{dist}(G, \tilde{G}) \leq \eta n^2$. (Note that this claim implies that every induced- C_{2k} -free graph contained in G can be made into a k-template by changing a total of at most ηn^2 edges within the vertex classes Q_i .) Indeed, by (b), each $G \in \mathcal{C}_1$ contains at most $\varepsilon' n^{2k}$ copies of C_{2k} . Thus by Lemma 3.5 there exists a complete 2-coloured multigraph G' on the same vertex set that is C_{2k} -free, such that $\operatorname{dist}(G,G') \leq \delta n^2$. Then $|G'_R \cap G'_B| \geq t_{k-1}(n) - (\varepsilon' + \delta)n^2$. Thus by Lemma 3.7 there exists a complete 2-coloured multigraph G'' on the same vertex set, with $G''_R \cap G''_B = E(T_{k-1}(n))$ and such that $\operatorname{dist}(G',G'') \leq \gamma n^2$. Note that G'' can contain at most γn^{2k} copies of C_{2k} , since G' is C_{2k} -free. Let $Q = \{Q_0,Q_1,\dots,Q_{k-2}\}$ be the unique (k-1)-partition of V(G'') such that no partition class induces an edge in $G''_R \cap G''_B$. Thus by Lemma 3.9, there exists a k-template $T = (V(G), E^T)$ on Q such that $|G''_R \triangle E^T| \leq \beta n^2$. Define \tilde{G} to be the 2-coloured multigraph with $\tilde{G}_R \cap \tilde{G}_B = G''_R \cap G''_B$ and $\tilde{G}_R \cap Q_i = E(T[Q_i])$ for every $i \in \{0, 1, \dots, k-2\}$. Then $\operatorname{dist}(G, \tilde{G}) \leq (\delta + \gamma + \beta)n^2 \leq \eta n^2$, and \tilde{G} satisfies the required properties. This proves the claim and thus the lemma.

4. The number of k-templates

For $k \geq 4$ we denote the set of all k-templates on n vertices by T(n,k). Let $T_Q(n,k)$ denote the set of all k-templates on n vertices for which Q is an ordered (k-1)-partition. The aim of this section is to estimate $|T_Q(n,k)|$ and |T(n,k)| (see Lemmas 4.4 and 4.6 respectively). Before we start with this we need to introduce some more notation. A k-sun is defined as follows.

- If k = 4, a k-sun is any sun (as defined in Section 3.3).
- If k = 5, a k-sun is a star or a clique.
- If $k \ge 6$, a k-sun is a star or a triangle.

Note that the results of this section are only needed for Theorem 1.1 (and not Theorem 1.2) and so we would only need to consider the case $k \geq 6$. However, including the cases k = 4, 5 makes little difference to the proofs, and are also interesting in their own right, so we work with all $k \geq 4$ throughout this section.

Let $F_k(n)$ denote the set of all *n*-vertex graphs whose complement is a disjoint union of *k*-suns. Define $f_k(n) := |F_k(n)|$. A pair of vertices x, y is called a *twin pair* if $N(x) \setminus \{y\} = N(y) \setminus \{x\}$.

The following two lemmas give some estimates of the value of $f_k(n)$. Note that we do not make use of the upper bound in Lemma 4.1 anywhere in this paper, but we include it for its intrinsic interest. It would not be difficult to obtain more accurate bounds, though an asymptotic formula would probably require more work.

Lemma 4.1. For all $n \in \mathbb{N}$ and $k \geq 4$,

$$2^{n\log n - en\log\log n} \le f_k(n) \le 2^{n\log n - n\log\log n + n}.$$

Proof. Let P(n) denote the number of partitions of an n element set. It is well known (see e.g. [9]) that

$$2^{n\log n - en\log\log n} \le P(n) \le 2^{n\log n - n\log\log n}.$$

We will count the number $f_k(n)$ of graphs $G \in F_k(n)$. Note that $f_k(n) \ge P(n)$ follows by considering each partition class as the vertex set of a star in \overline{G} . This then immediately yields the lower bound in Lemma 4.1. Now note that if we choose a partition of [n] into the vertex sets of disjoint suns in \overline{G} (for which there are at most $2^{n \log n - n \log \log n}$ choices), and then for every vertex choose whether the vertex will be in the body of its sun or side of its sun (for which there are a total of 2^n choices),

we can generate every possible graph $G \in F_k(n)$ (note that some such graphs can be generated by multiple different choices). This yields the upper bound in Lemma 4.1.

Lemma 4.2. For $k \ge 4$ and $n > s \ge 10^7$,

$$s^{s/2} \le \frac{f_k(n)}{f_k(n-s)}$$
 and $\frac{f_k(n)}{f_k(n-1)} \le n^2$.

Proof. By Lemma 4.1, $f_k(n) \ge f_k(s) f_k(n-s) \ge 2^{s \log s - es \log \log s} f_k(n-s) \ge 2^{s \log s/2} f_k(n-s)$, which gives us the lower bound in the statement of the lemma.

For the upper bound, note that every graph in $F_k(n)$ has a twin pair. For any twin pair $i, j \in [n]$ the number of graphs in $F_k(n)$ for which i, j are twins is at most $2f_k(n-1)$, since every such graph can be obtained from a graph in $F_k(n-1)$ on vertex set $[n] \setminus \{i\}$ by adding the vertex i and choosing whether to add the edge ij (note that all other edges incident to i are prescribed, since i, j are twins). Thus

$$f_k(n) \le \sum_{0 < i \le n-1} \sum_{i < j \le n} 2f_k(n-1) \le n^2 f_k(n-1),$$

as required.

The following proposition can be proved by a simple but tedious calculation, which we omit here.

Proposition 4.3. Let $k, n \in \mathbb{N}$ with $n \ge k \ge 2$ and let 0 < s < n.

(i) Suppose G is a k-partite graph on n vertices in which some vertex class A satisfies $|A-n/k| \ge s$. Then

$$e(G) \le t_k(n) - s\left(\frac{s}{2} - k\right).$$

(ii)
$$t_{k-1}(n) \ge t_{k-1}(n-s) + sn(k-2)/(k-1) - s(k-2) - t_{k-1}(s)$$
.

Lemma 4.4. Let $k \geq 4$. There exists $n_0 \in \mathbb{N}$ such that for every $n \geq n_0$ and every ordered (k-1)-partition Q of [n], the number of k-templates on Q satisfies

$$|T_Q(n,k)| \le 2^{6(\log n)^2} 2^{t_{k-1}(n)} f_k(n_k),$$

where we recall that $n_k := \lceil n/(k-1) \rceil$.

Proof. Denote the classes of Q by $Q_0, Q_1, \ldots, Q_{k-2}$ and let $b := ||Q_0| - \lceil \frac{n}{k-1} \rceil|$. Then by Proposition 4.3(i) the number of k-templates on this partition is at most

$$f_k(|Q_0|)2^{\sum_{0\leq i< j\leq k-2}|Q_i||Q_j|} \leq f_k(n_k+b)2^{t_{k-1}(n)-b(b/2-(k-1))}.$$

Let $h(b) := f_k(n_k + b)2^{t_{k-1}(n) - b(b/2 - (k-1))}$. Then by Lemma 4.2,

$$\frac{h(b+1)}{h(b)} \le \left(\frac{n}{k-1} + b + 2\right)^2 2^{-((2b+1)/2 - (k-1))}.$$

Thus h(b) is a decreasing function for $b \ge 3 \log n$. This together with Lemma 4.2 gives us that the number of k-templates on Q is at most

$$h(b) \le f_k (n_k + 3\log n) 2^{t_{k-1}(n)} \le (n^2)^{3\log n} 2^{t_{k-1}(n)} f_k (n_k)$$

= $2^{6(\log n)^2} 2^{t_{k-1}(n)} f_k (n_k)$,

as required. \Box

We call a component of a graph *non-trivial* if it contains at least 2 vertices. The proof of Lemma 4.6 will make use of the following proposition.

Proposition 4.5. Let $k \geq 4$. There exists $n_0 \in \mathbb{N}$ such that the following holds for every $n \geq n_0$. Let Q be a balanced ordered (k-1)-partition of [n]. The proportion of k-templates G on Q that are such that $\overline{G}[Q_0]$ has at most one non-trivial component is at most 2^{-n} .

Proof. Since Q is balanced, the number of k-templates on Q is at least $2^{t_{k-1}(n)} f_k(\lfloor \frac{n}{k-1} \rfloor)$.

We can generate all possible edge sets for $G[Q_0]$ such that $\overline{G}[Q_0]$ has at most one non-trivial component in the following way. Note that for every such $G[Q_0]$, $\overline{G}[Q_0]$ contains at most one disjoint sun S of order at least two. For every vertex in Q_0 we choose whether it will belong to the body of S, the side of S, or neither (for which there are a total of at most 3^n choices). Hence the number of k-templates G on Q that are such that $\overline{G}[Q_0]$ has at most one non-trivial component is at most $3^n 2^{t_{k-1}(n)}$.

Since we have by Lemma 4.1 that $f_k(m) \geq 2^{m \log m - e m \log \log m}$ for all $m \in \mathbb{N}$, the result follows (with some room to spare).

The following trivial observation will be useful in the proof of Lemma 4.6.

(4.1) If a graph G is a disjoint union of suns then G contains no induced 4-cycles.

Lemma 4.6. For every $k \geq 4$ there exists $n_0 \in \mathbb{N}$ such that the following holds for all $n \geq n_0$, where we recall that $n_k := \lceil n/(k-1) \rceil$. The number of k-templates on vertex set [n] satisfies

$$|T(n,k)| \ge \frac{(k-1)^n}{2(k-2)!n^k} 2^{t_{k-1}(n)} f_k(n_k).$$

Proof. Choose n_0 such that $1/n_0 \ll 1/k$, and let $n \ge n_0$. Given a k-template G on vertex set [n] and an ordered (k-1)-partition $Q = (Q_0, \{Q_1, \ldots, Q_{k-2}\})$ of [n], we say that G is Q-compatible if G is a k-template on Q and the following hold:

(α) Whenever $\ell \leq 2k$ and $0 \leq i \leq k-2$ and $v_1, v_2, \ldots, v_\ell \in V(G) \setminus Q_i$, we have that

$$|\overline{N}_{Q_i}(\{v_1, v_2, \dots, v_\ell\})| \ge \frac{n}{2^{\ell+1}(k-1)}.$$

 (β) $\overline{G}[Q_0]$ has at least 2 non-trivial components.

Claim 1: Given a balanced ordered (k-1)-partition $Q = (Q_0, \{Q_1, \ldots, Q_{k-2}\})$ of [n], the number of k-templates G on vertex set [n] that are Q-compatible is at least $2^{t_{k-1}(n)-1}f_k(n_k)/n^2$. Indeed, consider a random graph G where for each potential crossing edge with respect to Q we choose the edge to be present or not, each with probability 1/2, independently; we let $G[Q_0]$ be one of the $f_k(|Q_0|)$ graphs in $F_k(|Q_0|)$, chosen uniformly at random; and we choose all edges to be present inside Q_i for every i > 0. So each k-template on Q is equally likely to be generated. Note that the number of potential crossing edges with respect to Q is $2^{t_{k-1}(n)}$. This together with Lemma 4.2 implies that the number of graphs in the probability space is at least $2^{t_{k-1}(n)}f_k(n_k)/n^2$. By Lemma 2.1(ii) and Proposition 4.5 respectively, we have that at least half of all graphs G in the probability space satisfy (α) and (β) , which proves the claim.

Claim 2: Given two balanced ordered (k-1)-partitions $Q=(Q_0,\{Q_1,\ldots,Q_{k-2}\})$ and $Q'=(Q'_0,\{Q'_1,\ldots,Q'_{k-2}\})$ of [n], and a k-template G on [n] that is both Q-compatible and Q'-compatible, there exist k vertices $u_0,v_0,v_1,\ldots,v_{k-2}\in [n]$ that are such that $G[\{u_0,v_0,v_1,\ldots,v_{k-2}\}]$ contains exactly one edge u_0v_0 and $u_0\in Q_0\cap Q'_0$ and $v_i\in Q_i\cap Q'_i$ for all $i\geq 0$.

To show this, we first choose a set of 2k vertices $U = \{u_{0,1}, w_{0,1}, u_{0,2}, w_{0,2}, u_1, w_1, \dots, u_{k-2}, w_{k-2}\}$ such that $u_{0,1}, w_{0,1}, u_{0,2}, w_{0,2} \in Q_0$ and $u_i, w_i \in Q_i$ for every i > 0 and

$$E(G[U]) = \{u_{0,1}u_{0,2}, u_{0,2}w_{0,1}, w_{0,1}w_{0,2}, w_{0,2}u_{0,1}, u_1w_1, \dots, u_{k-2}w_{k-2}\}.$$

This is possible since G satisfies (α) , (β) with respect to Q. Now if there exist distinct i, j > 0 such that $u_i, w_i, u_j, w_j \in Q'_0$ then $\overline{G}[Q'_0]$ contains the induced 4-cycle $u_i u_j w_i w_j$, which by (4.1) contradicts the fact that G is a k-template on Q'. So, by relabelling vertices if necessary, we may assume that $u_2, \ldots, u_{k-2} \notin Q'_0$. If $u_{0,1}, w_{0,1} \notin Q'_0$ then by the pigeon-hole principle there must exist i > 0 such that Q'_i contains at least 2 elements of $\{u_{0,1}, w_{0,1}, u_2, \ldots, u_{k-2}\}$, contradicting the assumption that $G[Q'_i]$ is a clique. So, by relabelling vertices if necessary, we may assume that $u_{0,1} \in Q'_0$, and similarly that $u_{0,2} \in Q'_0$. Now if $u_1, w_1 \in Q'_0$ then $\overline{G}[Q'_0]$ contains the induced 4-cycle $u_{0,1}u_1u_{0,2}w_1$, which by (4.1) contradicts the fact that G is a k-template on Q'. So, by relabelling vertices if necessary, we may assume that $u_1 \notin Q'_0$, and thus $u_1, \ldots, u_{k-2} \notin Q'_0$. Recall that for all i > 0, $G[Q'_i]$ is a clique, so Q'_i can contain at most one vertex in $\{u_1, \ldots, u_{k-2}\}$. Thus we may assume, by relabelling indices if necessary, that $u_{0,1}, u_{0,2} \in Q_0 \cap Q'_0$ and $u_i \in Q_i \cap Q'_i$ for every i > 0. So setting $u_0 := u_{0,1}, v_0 := u_{0,2}$ and $v_i := u_i$ for all i > 0 yields the required set of vertices.

Claim 3: If there exist balanced ordered (k-1)-partitions $Q = (Q_0, \{Q_1, \ldots, Q_{k-2}\})$ and $Q' = (Q'_0, \{Q'_1, \ldots, Q'_{k-2}\})$ of [n], and a k-template G on [n] that is both Q-compatible and Q'-compatible, then Q = Q'.

Consider any k vertices $u_0, v_0, \ldots, v_{k-2} \in V(G)$ that are such that $G[\{u_0, v_0, v_1, \ldots, v_{k-2}\}]$ contains exactly one edge u_0v_0 and $u_0 \in Q_0 \cap Q'_0$ and $v_i \in Q_i \cap Q'_i$ for all $i \geq 0$. Such vertices exist by Claim 2. For i > 0 define

$$\overline{N}_i := \overline{N}_{Q_i}(\{u_0, v_0, \dots, v_{k-2}\} \setminus \{v_i\}).$$

$$\overline{N}'_i := \overline{N}_{Q'_i}(\{u_0, v_0, \dots, v_{k-2}\} \setminus \{v_i\}).$$

Since both \overline{N}_i and \overline{N}_i' are subsets of the common non-neighbourhood of $\{u_0, v_0, v_1, \dots, v_{k-2}\} \setminus \{v_i\}$, neither can intersect Q_j or Q_j' for $j \notin \{0, i\}$. Note that all vertices in \overline{N}_i are adjacent. Thus $|\overline{N}_i \cap Q_0'| \leq 1$, since otherwise $\overline{G}[Q_0']$ contains an induced 4-cycle on u_0, v_0 together with 2 vertices from \overline{N}_i , which by (4.1) contradicts the fact that G is a k-template on Q'. Similarly, $|\overline{N}_i' \cap Q_0| \leq 1$. Define

$$\overline{N}_i^{\dagger} := (\overline{N}_i \cup \overline{N}_i') \setminus (Q_0 \cup Q_0').$$

Then $\overline{N}_i^{\dagger} \subseteq Q_i \cap Q_i'$.

Now we consider any vertex $w \in Q_0$. Since G satisfies (α) with respect to Q, we have that for every i > 0,

$$(4.2) |\overline{N}_{Q_i'}(w)| \ge |\overline{N}(w) \cap \overline{N}_i^{\dagger}| \ge |\overline{N}(w) \cap \overline{N}_i| - 1$$

$$= |\overline{N}_{Q_i}(\{u_0, v_0, \dots, v_{k-2}, w\} \setminus \{v_i\})| - 1 \ge \frac{n}{2^{k+1}(k-1)} - 1 \ge 1.$$

Thus w must belong to Q'_0 , since $G[Q'_i]$ is a clique for every i > 0. Hence $Q_0 \subseteq Q'_0$. In the same way we can show that $Q'_0 \subseteq Q_0$. Thus $Q_0 = Q'_0$.

Now we consider any vertex $w \in Q_j$, for j > 0. Since G satisfies (α) with respect to Q, we have (similarly to (4.2)) that for every $i \neq j$ with i > 0,

$$|\overline{N}_{Q_i'}(w)| \ge |\overline{N}(w) \cap \overline{N}_i^{\dagger}| \ge 1.$$

Thus $w \in Q'_0 \cup Q'_j$. Together with the fact that $Q_0 = Q'_0$ this implies that $w \in Q'_j$. Thus $Q_j \subseteq Q'_j$ for all i > 0.

Hence Q = Q', which proves the claim.

We now count the number of balanced ordered (k-1)-partitions. Since the vertex classes of a balanced ordered (k-1)-partition of [n] have sizes $\lceil \frac{n}{k-1} \rceil, \lceil \frac{n-1}{k-1} \rceil, \ldots, \lceil \frac{n-k+2}{k-1} \rceil$, the number of such (k-1)-partitions is

$$\frac{1}{(k-2)!} \binom{n}{\lceil \frac{n}{k-1} \rceil, \lceil \frac{n-1}{k-1} \rceil, \dots, \lceil \frac{n-k+2}{k-1} \rceil}.$$

This together with Claims 1 and 3 implies that

$$(4.3) |T(n,k)| \ge \frac{1}{2(k-2)!n^2} \binom{n}{\lceil \frac{n}{k-1} \rceil, \lceil \frac{n-1}{k-1} \rceil, \dots, \lceil \frac{n-k+2}{k-1} \rceil} 2^{t_{k-1}(n)} f_k(n_k).$$

Now note that if $a_1 + \cdots + a_{k-1} = n$, then $\binom{n}{a_1, a_2, \dots, a_{k-1}}$ is maximized by taking $a_j := \lceil \frac{n-j+1}{k-1} \rceil$ for every j. This implies that

$$(k-1)^n = \sum_{a_1 + \dots + a_{k-1} = n} \binom{n}{a_1, a_2, \dots, a_{k-1}} \le n^{k-2} \binom{n}{\lceil \frac{n}{k-1} \rceil, \lceil \frac{n-1}{k-1} \rceil, \dots, \lceil \frac{n-k+2}{k-1} \rceil},$$

which together with (4.3) implies the result.

5. Properties of Near-k-Templates

In this section we collect some properties of graphs which are close to being k-templates. In particular, when $k \geq 6$, this means we consider graphs G which have a vertex partition such that each vertex class induces on G an almost complete graph. (As in the previous section, we will need the results of this section for our main results only for the case $k \geq 6$, but we prove the results for all $k \geq 4$ since it makes little difference to the proofs.) More formally, given $k \geq 4$, a graph G on vertex set [n], and an ordered (k-1)-partition Q of [n] we define

$$h(Q,G) := \sum_{i=0}^{k-2} |E(\overline{G}[Q_i])|.$$

We say Q is an optimal ordered (k-1)-partition of G if h(Q,G) is the minimum value h(Q',G) takes over all partitions Q' of [n]. Note that if h(Q,G)=0 then G is a k-template on Q, and that the following also holds.

(5.1) If
$$k \ge 6$$
 then every k-template G' on Q satisfies $h(Q, G') \le n$.

Note that (5.1) does not hold for $k \in \{4,5\}$. We will require the following definitions in what follows.

- Recall that F(n,k) denotes the set of all labelled induced- C_{2k} -free graphs on vertex set [n].
- Given $n \in \mathbb{N}$, $k \geq 4$, and $\eta > 0$, we define $F(n, k, \eta) \subseteq F(n, k)$ to be the set of all graphs in F(n, k) such that $h(Q, G) \leq \eta n^2$ for some optimal ordered (k-1)-partition Q of G.
- Given further an ordered (k-1)-partition $Q=(Q_0,\{Q_1,\ldots,Q_{k-2}\})$ of [n] we define $F_Q(n,k)\subseteq F(n,k)$ to be the set of all graphs in F(n,k) for which Q is an optimal ordered (k-1)-partition
- Similarly we define $F_Q(n, k, \eta) \subseteq F(n, k, \eta)$ to be the set of all graphs in $F(n, k, \eta)$ for which Q is an optimal ordered (k-1)-partition.

Recall that, given a graph G on vertex set [n] and an index $i \in \{0, 1, ..., k-2\}$, we let $d_{G,Q}^i(x), \overline{d}_{G,Q}^i(x)$ denote the number of neighbours and non-neighbours of x in Q_i , respectively. The following proposition follows immediately from the definition of optimality.

Proposition 5.1. Let $k \geq 4$, let $\eta > 0$, let $Q = (Q_0, \{Q_1, \dots, Q_{k-2}\})$ be an ordered (k-1)-partition of [n], and let $G \in F_Q(n, k, \eta)$. For any two distinct indices $i, j \in \{0, 1, \dots, k-2\}$ every vertex $x \in Q_i$ satisfies $\overline{d}_{G,Q}^j(x) \geq \overline{d}_{G,Q}^i(x)$.

Next we show that for most graphs which are close to being k-templates, the bipartite graphs between the partition classes are quasirandom.

Given $k \geq 4$, $\nu = \nu(n) > 0$ and an ordered (k-1)-partition $Q = (Q_0, \{Q_1, \dots, Q_{k-2}\})$ of [n], we define the following properties that a graph on vertex set [n] may satisfy with respect to Q.

 $(\mathrm{F1})_{\nu} \ \ \mathrm{If} \ U_i \subseteq Q_i \ \mathrm{and} \ U_j \subseteq Q_j \ \mathrm{with} \ |U_i||U_j| \ge \nu^2 n^2 \ \mathrm{for \ distinct} \ 0 \le i,j \le k-2, \ \mathrm{then} \ \tfrac{1}{4} \le \tfrac{|e(U_i,U_j)|}{|U_i||U_j|} \le \tfrac{3}{4}.$ $(\mathrm{F2})_{\nu} \ ||Q_i| - \tfrac{n}{k-1}| \le \nu n \ \mathrm{for \ every} \ 0 \le i \le k-2.$

Given $\eta, \mu > 0$ we define $F_Q(n, k, \eta, \mu)$ to be the set of all graphs in $F_Q(n, k, \eta)$ that satisfy $(F1)_{\mu}$ and $(F2)_{\mu}$ with respect to Q.

Lemma 5.2. Let $n \ge k \ge 4$, let $0 < \eta < 1$, let $6k/n \le \nu = \nu(n) \le 1$, let $6\log n \le m = m(n) \le 1$ $10^{-11}n^2$, and let $Q=(Q_0,\{Q_1,\ldots,Q_{k-2}\})$ be an ordered (k-1)-partition of [n]. Then the following hold.

- (i) The number of graphs G in $F_Q(n,k,\eta)$ that fail to satisfy $(F1)_{\nu}$ with respect to Q and that have at most m internal non-edges is at most $2^{t_{k-1}(n)+\xi(m/n^2)n^2}2^{2n+1}\exp(-\nu^2n^2/32)$.
- (ii) The number of graphs G in $F_Q(n,k,\eta)$ that fail to satisfy $(F2)_{\nu}$ with respect to Q and that have at most m internal non-edges is at most $2^{t_{k-1}(n)+\xi(m/n^2)n^2}\exp(-\nu^2n^2/6)$.

Proof. For both (i) and (ii) we consider constructing such a graph G. By (2.1) there are at most $\binom{n^2}{\leq m} \leq 2^{\xi(m/n^2)n^2}$ choices for the internal edges of G.

We first prove (i). For a given choice of internal edges, consider the random graph H where for each possible crossing edge with respect to Q we choose the edge to be present or not, with probability 1/2, independently. Note that the total number of ways to choose the crossing edges is at most $2^{t_{k-1}(n)}$, and each possible configuration of crossing edges is equally likely. So an upper bound on the number of graphs $G \in F_Q(n, k, \eta)$ that fail to satisfy property $(F1)_{\nu}$ with respect to Q and that have at most m internal non-edges is

(5.2)
$$2^{t_{k-1}(n)+\xi(m/n^2)n^2}\mathbb{P}(H \text{ fails to satisfy } (F1)_{\nu} \text{ with respect to } Q).$$

Note that the number of choices for $U_i \subseteq Q_i, U_j \subseteq Q_j$ with $|U_i||U_j| \ge \nu^2 n^2$ is at most 2^{2n} and that $\mathbb{E}(e(U_i,U_i)) = |U_i||U_i|/2 \ge \nu^2 n^2/2$. Hence by Lemma 2.1,

$$\mathbb{P}(H \text{ fails to satisfy } (\mathrm{F1})_{\nu} \text{ with respect to } Q) \leq 2^{2n+1} \exp\left(-\frac{\nu^2 n^2}{32}\right).$$

This together with (5.2) yields the result.

We now prove (ii). If $||Q_i| - \frac{n}{k-1}| > \nu n$ for some $0 \le i \le k-2$, then by Proposition 4.3(i) the number of crossing edges in G is at most

$$t_{k-1}(n) - \frac{\nu^2 n^2}{3}.$$

We can conclude that the number of $G \in F_Q(n,k,\eta)$ that fail to satisfy $(F2)_{\nu}$ with respect to Q and that have at most m internal non-edges is at most

$$2^{\xi(m/n^2)n^2} 2^{t_{k-1}(n) - \frac{\nu^2 n^2}{3}} \le 2^{t_{k-1}(n) + \xi(m/n^2)n^2} \exp\left(-\frac{\nu^2 n^2}{6}\right),$$

as required.

We will apply the following special case of Lemma 5.2 in Section 10 in the proof of Lemma 6.1.

Corollary 5.3. Let $k \ge 4$ and let $0 < \eta, \mu < 10^{-11}$ be such that $\mu^2 > 24\xi(\eta)$. There exists an integer $n_0 = n_0(\mu, k)$ such that for all $n \ge n_0$ and every ordered (k-1)-partition Q of [n],

$$|F_O(n, k, \eta) \setminus F_O(n, k, \eta, \mu)| \le 2^{t_{k-1}(n) - \frac{\mu^2 n^2}{100}}.$$

Proof. We choose n_0 such that $1/n_0 \ll \eta, \mu, 1/k$. Applying Lemma 5.2 with $\mu, \eta n^2$ playing the roles of ν, m respectively yields that

$$|F_Q(n,k,\eta)\setminus F_Q(n,k,\eta,\mu)|\leq 2^{t_{k-1}(n)+\xi(\eta)n^2}2^{2n+1}\left(e^{-\frac{\mu^2n^2}{6}}+e^{-\frac{\mu^2n^2}{32}}\right)\leq 2^{t_{k-1}(n)-\frac{\mu^2n^2}{100}},$$
 as required.

The next proposition follows immediately from [4, Lemma 2.22]. We will use it to find induced copies of C_{2k} . (Usually T will be a suitable induced subgraph of C_{2k} and the A_i, B_i will be the intersection of (non-)neighbourhoods of vertices that we have already embedded.)

Proposition 5.4. Let $n_0, k \in \mathbb{N}$ and $\eta, \mu > 0$ be chosen such that $k \geq 4$ and $1/n_0 \ll \eta \ll \mu \ll 1/k$. Then the following holds for all $n \in \mathbb{N}$ with $n \geq n_0$. Let $Q = (Q_0, \{Q_1, \ldots, Q_{k-2}\})$ be an ordered (k-1)-partition of [n] and suppose $G \in F_Q(n, k, \eta, \mu)$. Let $I \subseteq \{0, 1, \ldots, k-2\}$. For every $i \in I$ let $A_i, B_i \subseteq Q_i$ be disjoint with $|A_i|, |B_i| \geq \mu^{1/2}n$. Let T be a 2|I|-vertex graph with a perfect matching whose edges are $v_i u_i$ for every $i \in I$. Then there exists an injection $f : V(T) \to V(G)$ such that $f(v_i) \in A_i, f(u_i) \in B_i$ for every $i \in I$, and f(V(T)) induces on G a copy of T.

Finally we show that if G is close to being a k-template then removing a small number of vertices from G does not alter its optimal ordered (k-1)-partition very much.

Given $m, n \in \mathbb{N}$ and an ordered (k-1)-partition Q of [n], we define $\mathcal{P}(Q, m)$ to be the collection of all ordered (k-1)-partitions of [n] that can be obtained from Q by moving at most m vertices between partition classes, and possibly choosing a different partition class to be the labelled one. Then it is easy to see that

$$|\mathcal{P}(Q,m)| \le k \binom{n}{m} k^m \le k \left(\frac{ekn}{m}\right)^m \le 2^{m \log(ek^2 n/m)}.$$

Given an ordered (k-1)-partition Q of [n] and a set $S \subseteq [n]$, let Q-S denote the ordered (k-1)-partition (possibly with some empty classes) obtained from Q by deleting all elements of S from their partition classes.

Lemma 5.5. Let $k \geq 4$, let $0 < \eta, \mu \leq 1/k^3$, let $0 < \nu = \nu(n) \leq 1/k^3$, and let $0 \leq m = m(n) \leq n^2$ with $\nu^2 > 4m/n^2$ for all $n \in \mathbb{N}$. There exists $n_0 \in \mathbb{N}$ such that the following holds for all $n \geq n_0$. Let $Q = (Q_0, \{Q_1, \dots, Q_{k-2}\})$ be an ordered (k-1)-partition of [n] and let $S \subseteq [n]$ with $|S| \leq n/k^2$. Then for every $G \in F_Q(n, k, \eta, \mu)$ that satisfies $(F1)_{\nu}$ with respect to Q and that has at most m internal non-edges, every optimal ordered (k-1)-partition of G - S is an element of $\mathcal{P}(Q - S, k^4 \nu^2 n)$.

Proof. Let $G \in F_Q(n, k, \eta, \mu)$ have at most m internal non-edges and satisfy $(F1)_{\nu}$ with respect to Q, and let $Q' = (Q'_0, \{Q'_1, \dots, Q'_{k-2}\})$ be an optimal ordered (k-1)-partition of G-S. By optimality of Q' it must be that G-S has at most m internal non-edges with respect to Q'.

For every $i \in \{0, 1, ..., k-2\}$, since $|Q_i - S| \ge n/(k-1) - \mu n - n/k^2 \ge n/k$, the pigeon-hole principle implies that there exists $j \in \{0, 1, ..., k-2\}$ such that $|Q_i \cap Q'_j| \ge n/k^2$. We define a function σ by setting $\sigma(i)$ to be an index in $\{0, 1, ..., k-2\}$ that satisfies $|Q_i \cap Q'_{\sigma(i)}| \ge n/k^2$, for every $i \in \{0, 1, ..., k-2\}$. Suppose for a contradiction that there exists $i' \in \{0, 1, ..., k-2\}$ with $i \ne i'$ such that $|Q_{i'} \cap Q'_{\sigma(i)}| \ge k^2 \nu^2 n$. Then since G satisfies $(F1)_{\nu}$ with respect to Q we have that the number of internal non-edges in G - S with respect to Q' is at least $|Q_i \cap Q'_{\sigma(i)}| |Q_{i'} \cap Q'_{\sigma(i)}|/4 \ge n$

 $\nu^2 n^2/4 > m$. This contradicts our previous observation that G-S has at most m internal non-edges with respect to Q'. Hence σ is a permutation on $\{0, 1, \ldots, k-2\}$. Moreover $|Q_i \cap Q'_j| < k^2 \nu^2 n$ for all $j \in \{0, 1, \ldots, k-2\}$ with $j \neq \sigma(i)$.

Let \mathcal{P} be the set of all ordered (k-1)-partitions of $[n] \setminus S$ for which such a permutation exists. So by the above we have that for every $G \in F_Q(n,k,\eta)$ that satisfies $(F1)_{\nu}$ with respect to Q and that has at most m internal non-edges, every optimal ordered (k-1)-partition of G-S is an element of \mathcal{P} . So it remains to show that $\mathcal{P} \subseteq \mathcal{P}(Q-S,k^4\nu^2n)$. This follows from the observation that every element of \mathcal{P} can be obtained by starting with the (labelled) (k-1)-partition $Q_0 \setminus S, Q_1 \setminus S, \ldots, Q_{k-2} \setminus S$, applying a permutation of $\{0,1,\ldots,k-2\}$ to the partition class labels, then for every ordered pair of partition classes moving at most $k^2\nu^2n$ elements from the first partition class to the second, and finally unlabelling all but one of the resulting partition classes.

The following is an immediate corollary of Lemma 5.5, applied with $\mu, \eta n^2$ playing the roles of ν, m , respectively.

Corollary 5.6. Let $k \geq 4$ and $0 < \eta, \mu < 1/k^3$ with $\mu^2 > 4\eta$. There exists $n_0 \in \mathbb{N}$ such that the following holds for all $n \geq n_0$. Let $Q = (Q_0, \{Q_1, \dots, Q_{k-2}\})$ be an ordered (k-1)-partition of [n] and let $S \subseteq [n]$ with $|S| \leq n/k^2$. Then for every $G \in F_Q(n, k, \eta, \mu)$, every optimal ordered (k-1)-partition of G - S is an element of $\mathcal{P}(Q - S, k^4\mu^2n)$.

6. Derivation of Theorem 1.1 from the main Lemma

The following lemma is the key result in our proof of Theorem 1.1. Together with Lemma 4.4 it implies that, for $k \geq 6$, almost all induced- C_{2k} -free graphs G with a given optimal ordered (k-1)-partition are k-templates. Recall that $n_k := \lceil n/(k-1) \rceil$, that $f_k(n)$ and $T_Q(n,k)$ were defined at the beginning of Section 4, and that $F_Q(n,k)$ was defined at the beginning of Section 5.

Lemma 6.1. For every $n, k \in \mathbb{N}$ with $k \geq 6$ there exists $C \in \mathbb{N}$ such that the following holds. For every ordered (k-1)-partition Q of [n],

$$|F_Q(n,k)| \le |T_Q(n,k)| + 5C2^{-n^{\frac{1}{2k^2}/3}} f_k(n_k) 2^{t_{k-1}(n)}$$

Lemma 6.1 will be proved in the remaining sections of this paper. We will now use it to derive Theorem 1.1.

Proof of Theorem 1.1. Let $n_0 \in \mathbb{N}$ be as in Lemma 4.6, let $C \in \mathbb{N}$ be as in Lemma 6.1, let $n_1 \in \mathbb{N}$ satisfy $1/n_1 \ll 1/k$, let $n \in \mathbb{N}$ with $n \ge \max\{n_0, n_1\}$, and let \mathcal{Q} be the set of all ordered (k-1)-partitions of [n]. Since $T(n,k) \subseteq F(n,k)$ and $T_{\mathcal{Q}}(n,k) \subseteq F_{\mathcal{Q}}(n,k)$ for every $\mathcal{Q} \in \mathcal{Q}$, Lemma 6.1 implies that

$$|F(n,k)| - |T(n,k)| = |F(n,k)\backslash T(n,k)| \le \sum_{Q\in\mathcal{Q}} |F_Q(n,k) \backslash T_Q(n,k)|$$

$$= \sum_{Q\in\mathcal{Q}} (|F_Q(n,k)| - |T_Q(n,k)|) \le 5C(k-1)^n 2^{-n^{\frac{1}{2k^2}/3}} f_k(n_k) 2^{t_{k-1}(n)}$$

$$\le C 2^{-n^{\frac{1}{2k^2}/4}} \frac{(k-1)^n}{2(k-2)!n^k} f_k(n_k) 2^{t_{k-1}(n)}.$$

This together with Lemma 4.6 implies that

$$|F(n,k)| - |T(n,k)| \le C2^{-n^{\frac{1}{2k^2}/4}} |T(n,k)| = o(|T(n,k)|),$$

where we use the little o notation with respect to n. So |F(n,k)| = (1+o(1))|T(n,k)|, as required.

Sections 7–10 are devoted to proving Lemma 6.1 by an inductive argument. For the remainder of the paper we fix constants $C, k, n_0 \in \mathbb{N}$ with $k \geq 6$ and $\varepsilon, \eta, \mu, \gamma, \beta, \alpha > 0$ such that

(6.1)
$$\frac{1}{C} \ll \frac{1}{n_0} \ll \varepsilon \ll \eta \ll \mu \ll \gamma \ll \beta \ll \alpha \ll \frac{1}{k}.$$

We also set $M := R_{2k-2}(\lceil \frac{1}{\gamma} \rceil) + 1$, fix an arbitrary integer $n \geq n_0$, and fix an arbitrary ordered (k-1)-partition $Q = (Q_0, \{Q_1, \dots, Q_{k-2}\})$ of [n].

We make the following inductive assumption in Sections 7, 8 and 9: for every $n' \leq n-1$, and every ordered (k-1)-partition $Q' = (Q'_0, \{Q'_1, \dots, Q'_{k-2}\})$ of [n'],

$$|F_{Q'}(n',k) \setminus T_Q(n',k)| \le 5C2^{-(n')^{\frac{1}{2k^2}}/3} f_k(n'_k) 2^{t_{k-1}(n')}.$$

Note that this together with Lemma 4.4 implies that

$$|F_{Q'}(n',k)| \le 6C2^{6(\log n')^2} f_k(n'_k) 2^{t_{k-1}(n')}.$$

We now give a number of definitions that will be used in the remaining sections. Given an index $i \in \{0, 1, \dots, k-2\}$, we call a vertex x of a graph G i-light if at least one of the following holds.

- (A1) $d_{G,Q}^i(x) \leq \alpha n$.
- (A2) $\overline{d}_{G,O}^{i}(x) \leq \alpha n$.
- (A3) There exists $z \in V(G)$ such that $|N_i^*(x,z)| + |N_i^*(z,x)| \leq \alpha n$.

(Intuitively, the neighbourhood in Q_i of an *i*-light vertex is 'atypical', and this is unlikely to happen.) Given $\psi > 0$ and an index $i \in \{0, 1, ..., k-2\}$, we call $\{x, y_1, y_2, y_3\} \subseteq V(G)$ a (k, x, i, ψ) configuration if it satisfies the following.

- (C1) $G[\{x, y_1, y_2, y_3\}]$ is a linear forest.
- (C2) $\overline{d}_{G,O}^{j}(x) \ge 13 \cdot 6^{k} \psi n \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\}.$
- (C3) There exists $i' \neq i$ such that $d_{G,Q}^j(x) \geq 13 \cdot 6^k \psi n$ for all $j \in \{0, 1, \dots, k-2\} \setminus \{i, i'\}$.
- (C4) $\min\{d_{G,O}^{i}(y_j), \overline{d}_{G,O}^{i}(y_j)\} \le \psi^2 n \text{ for all } j \in [3].$

(Intuitively, (C1)–(C3) of the definition of (k, x, i, ψ) -configurations are useful for 'building' induced copies of C_{2k} , so the existence of a (k, x, i, ψ) -configuration in an induced- C_{2k} -free graph G severely constrains the choices for the remaining edge set of G. The bounds arising from this are still not sufficiently strong though; we also need (C4), which gives further constraints on the choices for the remaining edge set of G.)

We partition $F_Q(n, k, \eta, \mu)$ into the sets T_Q, F_Q^1, F_Q^2, F_Q^3 defined as follows.

- (F0) $T_Q := T_Q(n,k) \cap F_Q(n,k,\eta,\mu)$. (F1) $F_Q^1 \subseteq F_Q(n,k,\eta,\mu) \setminus T_Q$ is the set of all remaining graphs G which satisfy one of the following.
 - (i) G contains a (k, x, i, ψ) -configuration for some $i \in \{0, 1, \dots, k-2\}$, some $x \in V(G)$ and some $\psi \in \{\beta^{1/2}, \beta^2\}.$
 - (ii) G contains a vertex x which is both i-light and j-light for some distinct indices $i, j \in$ $\{0, 1, \ldots, k-2\}.$
- (F2) $F_Q^2 \subseteq F_Q(n, k, \eta, \mu) \setminus (T_Q \cup F_Q^1)$ is the set of all remaining graphs that for some $i \in \{0, 1, \dots, k-1\}$ 2} contain a vertex $x \in Q_i$ that satisfies $\overline{d}_{G,Q}^i(x), d_{G,Q}^i(x) \ge \beta n$.
- (F3) $F_Q^3 := F_Q(n, k, \eta, \mu) \setminus (T_Q \cup F_Q^1 \cup F_Q^2)$ is the set of all remaining graphs.

Sections 7, 8 and 9 are devoted to proving upper bounds on $|F_Q^1|$, $|F_Q^2|$ and $|F_Q^3|$ respectively. As mentioned earlier, it turns out that F_Q^3 is the class of induced- C_{2k} -free graphs which are 'extremely close' to being k-templates (see Proposition 9.1). In Section 10 we will use these bounds to complete the proof of Lemma 6.1.

7. Estimation of $|F_Q^1|$

To estimate $|F_Q^1|$ we will bound the number of graphs satisfying (F1)(i) and (F1)(ii) separately. The main difficulty is in estimating those satisfying (F1)(i), i.e. the ones containing a (k, x, i, ψ) -configuration. The idea here is that a (k, x, i, ψ) -configuration has many potential extensions into an induced copy of C_{2k} . More precisely, given a (k, x, i, ψ) -configuration H we can find many disjoint 'skeleton' graphs L with the same number of components as H such that $H \cup L$ is a linear forest on 2k vertices (i.e. $H \cup L$ has a potential extension into an induced C_{2k}). Thus each skeleton induces a restriction on further edges that can be added. Since the skeletons are disjoint we obtain many edge restrictions in total, and thus a good bound on the number of graphs containing a (k, x, i, ψ) -configuration. The next two propositions are used to formalise the notion of extendibility into an induced C_{2k} . (Roughly, in these propositions one can consider L_1 as a (k, x, i, ψ) -configuration and L_2 as an associated skeleton.)

Proposition 7.1. Let $c \ge 1$ and let L_1, L_2 be disjoint linear forests, each with exactly c components, such that $|V(L_1)| + |V(L_2)| = 2k$. Then there exists a set E' of edges between $V(L_1)$ and $V(L_2)$ such that the graph $(V(L_1) \cup V(L_2), E' \cup E(L_1) \cup E(L_2))$ is isomorphic to C_{2k} .

The proof of Proposition 7.1 is trivial, and is omitted. Proposition 7.2 follows from an easy application of Proposition 7.1, and we give only a brief sketch of the proof.

Proposition 7.2. Let $c \geq 1$ and let L_1, L_2 be linear forests that satisfy the following.

- $V(L_1) \cap V(L_2) = \{x\}.$
- $|V(L_1)|, |V(L_2)| > 1$.
- $d_{L_1}(x) + d_{L_2}(x) = 2$.
- L_1 and $L_2 \{x\}$ both have exactly c components.
- $|V(L_1) \cup V(L_2)| = 2k$.

Then there exists a set E' of edges between $V(L_1)\setminus\{x\}$ and $V(L_2)\setminus\{x\}$ such that the graph $(V(L_1)\cup V(L_2), E'\cup E(L_1)\cup E(L_2))$ is isomorphic to C_{2k} .

Proof. If $d_{L_1}(x) = 0$ we apply Proposition 7.1 to $L_1 - x, L_2$; if $d_{L_2}(x) = 0$ we apply Proposition 7.1 to $L_1, L_2 - x$. If $d_{L_1}(x) = d_{L_2}(x) = 1$ one can easily find E' directly.

Lemma 7.3. $|F_O^1| \le C2^{-\frac{\beta^2 n}{14^k}} f_k(n_k) 2^{t_{k-1}(n)}$.

Proof. Let $F_{Q,(i)}^1$ denote the set of all graphs in F_Q^1 that satisfy (F1)(i). Similarly let $F_{Q,(ii)}^1$ denote the set of all graphs in F_Q^1 that satisfy (F1)(ii). Clearly,

$$|F_Q^1| \le |F_{Q,(i)}^1| + |F_{Q,(ii)}^1|.$$

We will first estimate the number of graphs in $F_{Q,(i)}^1$. Any graph $G \in F_{Q,(i)}^1$ can be constructed as follows. We first choose $\psi \in \{\beta^2, \beta^{1/2}\}$, and then perform the following steps.

- We choose an index $i \in \{0, 1, ..., k-2\}$, a set of three (labelled) vertices $Y = \{y_1, y_2, y_3\}$ in [n], a vertex $x \in [n] \setminus Y$, and a set E of edges between these four vertices such that $Y \cup \{x\}$ spans a linear forest. Let b_1 denote the number of such choices. The choices in the next steps will be made such that $Y \cup \{x\}$ is a (k, x, i, ψ) -configuration in G.
- Next we choose the graph G' on vertex set $[n] \setminus Y$ such that $G[[n] \setminus Y] = G'$. Let b_2 denote the number of possibilities for G'.
- Next we choose the set E' of edges in G between Y and $Q_i \setminus (Y \cup \{x\})$ such that E' is compatible with our previous choices. Let b_3 denote the number of possibilities for E'.
- Finally we choose the set E'' of edges in G between Y and $[n]\setminus (Q_i \cup Y \cup \{x\})$ such that E'' is compatible with our previous choices. Let b_4 denote the number of possibilities for E''.

Hence,

$$|F_{Q,(i)}^1| \le 2 \max_{\psi \in \{\beta^2, \beta^{1/2}\}} \{b_1 \cdot b_2 \cdot b_3 \cdot b_4\}.$$

We then estimate the number of graphs in $F_{Q,(ii)}^1$. Any graph $G \in F_{Q,(ii)}^1$ can be constructed as follows.

- We first choose a single vertex x from [n] and distinct indices $i, j \in \{0, 1, ..., k-2\}$. Let c_1 denote the number of such choices. The choices in the next steps will be made such that x is both i-light and j-light in G.
- Next we choose the graph G' on vertex set $[n]\setminus\{x\}$ such that $G[[n]\setminus\{x\}]=G'$. Let c_2 denote the number of possibilities for G'.
- Next we choose the set E of edges in G between $\{x\}$ and $(Q_i \cup Q_j) \setminus \{x\}$ such that E is compatible with our previous choices. Let c_3 denote the number of possibilities for E.
- Finally we choose the set E' of edges in G between $\{x\}$ and $[n]\setminus (Q_i \cup Q_j \cup \{x\})$. Let c_4 denote the number of possibilities for E'.

Hence,

$$|F_{Q,(ii)}^1| \le c_1 \cdot c_2 \cdot c_3 \cdot c_4.$$

The following series of claims will give upper bounds for the quantities $b_1, \ldots, b_4, c_1, \ldots, c_4$. Claims 1 and 5 are trivial, while the proof of Claim 6 is almost identical to that of Claim 2; we give proofs of Claims 2,3,4,7 and 8.

Claim 1: $b_1 \leq 2^6 k n^4$.

Claim 2: $b_2 \leq C2^{\mu^{1/2}n} f_k(n_k) 2^{t_{k-1}(n-3)}$.

Indeed, note that for every graph $\tilde{G} \in F_{Q,(i)}^1$, Corollary 5.6 together with (5.3) implies that every optimal ordered (k-1)-partition of $\tilde{G}[[n]\backslash Y]$ is contained in some set \mathcal{P} of size at most $2^{\mu n}$. Since $G[[n]\backslash Y]$ is clearly induced- C_{2k} -free, this together with (6.2) implies that

$$b_2 \le \sum_{Q' \in \mathcal{P}} |F_{Q'}(n-3,k)| \le 6C 2^{\mu n} 2^{6(\log n)^2} f_k(\lceil (n-3)/(k-1) \rceil) 2^{t_{k-1}(n-3)}$$

$$\le C 2^{\mu^{1/2}n} f_k(n_k) 2^{t_{k-1}(n-3)},$$

as required.

Claim 3: $b_3 \le 2^{4\psi^{3/2}n}$.

Indeed, for every graph $\tilde{G} \in F_{Q,(i)}^1$ for which $\{x, y_1, y_2, y_3\}$ is a (k, x, i, ψ) -configuration we have that $\min\{d_{\tilde{G},Q}^i(y_j), \overline{d}_{\tilde{G},Q}^i(y_j)\} \leq \psi^2 n$ for all $j \in [3]$. So $b_3 \leq \prod_{j=1}^3 h(j)$ where h(j) denotes the number of possibilities for a set of edges between $\{y_j\}$ and $Q_i \setminus (Y \cup \{x\})$ such that either $d_{G,Q}^i(y_j) \leq \psi^2 n$ or $\overline{d}_{G,Q}^i(y_j) \leq \psi^2 n$. Note that by $(2.1), h(j) \leq 2\binom{n}{\leq \psi^2 n} \leq 2^{\xi(\psi^2)n+1}$. Hence,

$$b_3 \le \prod_{j=1}^3 h(j) \le (2^{\xi(\psi^2)n+1})^3 \stackrel{(2.2)}{\le} 2^{4\psi^{3/2}n},$$

as required.

Claim 4: $b_4 \le 2^{3(k-2)n/(k-1)} 2^{\mu^{1/2}n} 2^{-\psi n/11^k}$.

Indeed, first define L to be the graph on vertex set $Y \cup \{x\}$ that satisfies E(L) = E. We say an induced subgraph H of G' - x is an L-compatible skeleton if it satisfies the following.

• |V(H)| = 2k - 4.

- $G'[V(H) \cup \{x\}]$ is a linear forest.
- In G', x has $2 d_L(x)$ neighbours in V(H).
- L and H have the same number of components.

Given an L-compatible skeleton H, note that Proposition 7.2, applied with $L, G'[V(H) \cup \{x\}]$ playing the roles of L_1 , L_2 respectively, implies that there exists a set $E_{L,H}$ of possible edges between Y and V(H) such that $(Y \cup \{x\} \cup V(H), E \cup E(H) \cup E_{L,H})$ is isomorphic to C_{2k} .

We will show that there exist a large number of disjoint L-compatible skeletons in G'-x. Since there is a limited number of ways to choose edges between Y and each of these L-compatible skeletons so as not to create an induced copy of C_{2k} , this will imply the claim.

For every index $j \neq i$, let $N_j^1(x), N_j^2(x) \subseteq N_{Q_j}(x)$ be disjoint with $|N_j^1(x)|, |N_j^2(x)| \ge \lfloor \frac{1}{2} |N_{Q_j}(x)| \rfloor$. Similarly, let $\overline{N}_{j}^{1}(x)$, $\overline{N}_{j}^{2}(x) \subseteq \overline{N}_{Q_{j}}(x)$ be disjoint with $|\overline{N}_{j}^{1}(x)|$, $|\overline{N}_{j}^{2}(x)| \ge \lfloor \frac{1}{2} |\overline{N}_{Q_{j}}(x)| \rfloor$. Note that we may assume that there exists an index $i' \in \{0, 1, \dots, k-2\} \setminus \{i\}$ such that in

 $G', |\overline{N}_{Q_j}(x)| \ge 12 \cdot 6^k \psi n \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ and } |N_{Q_j}(x)| \ge 12 \cdot 6^k \psi n \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ and } |N_{Q_j}(x)| \ge 12 \cdot 6^k \psi n \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ and } |N_{Q_j}(x)| \ge 12 \cdot 6^k \psi n \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ and } |N_{Q_j}(x)| \ge 12 \cdot 6^k \psi n \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ and } |N_{Q_j}(x)| \ge 12 \cdot 6^k \psi n \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ and } |N_{Q_j}(x)| \ge 12 \cdot 6^k \psi n \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ and } |N_{Q_j}(x)| \ge 12 \cdot 6^k \psi n \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ and } |N_{Q_j}(x)| \ge 12 \cdot 6^k \psi n \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ and } |N_{Q_j}(x)| \ge 12 \cdot 6^k \psi n \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ and } |N_{Q_j}(x)| \ge 12 \cdot 6^k \psi n \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ and } |N_{Q_j}(x)| \ge 12 \cdot 6^k \psi n \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ and } |N_{Q_j}(x)| \ge 12 \cdot 6^k \psi n \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ and } |N_{Q_j}(x)| \ge 12 \cdot 6^k \psi n \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ and } |N_{Q_j}(x)| \ge 12 \cdot 6^k \psi n \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ and } |N_{Q_j}(x)| \ge 12 \cdot 6^k \psi n \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ and } |N_{Q_j}(x)| \ge 12 \cdot 6^k \psi n \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ and } |N_{Q_j}(x)| \ge 12 \cdot 6^k \psi n \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ and } |N_{Q_j}(x)| \ge 12 \cdot 6^k \psi n \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ for all } j \in \{0, 1, \dots, k-2\} \setminus \{i\} \text{ for all }$ $\{0,1,\ldots,k-2\}\setminus\{i,i'\}$, since otherwise $\{x,y_1,y_2,y_3\}$ cannot be a (k,x,i,ψ) -configuration. Define $\ell_1, \ldots, \ell_{k-2}$ such that $\{\ell_1, \ldots, \ell_{k-2}\} = \{0, 1, \ldots, k-2\} \setminus \{i\}$ and $\ell_{k-2} = i'$. Thus the following hold.

(a)
$$|N_{\ell_j}^1(x)|, |N_{\ell_j}^2(x)|, |\overline{N}_{\ell_j}^1(x)|, |\overline{N}_{\ell_j}^2(x)| \ge 6 \cdot 6^k \psi n \text{ for all } j \in \{1, \dots, k-3\}.$$

(b)
$$|\overline{N}_{\ell_{k-2}}^1(x)|, |\overline{N}_{\ell_{k-2}}^2(x)| \ge 6 \cdot 6^k \psi n.$$

We now show that G'-x contains at least $5\cdot 6^k \psi n$ disjoint L-compatible skeletons. Define t to be the number of components of L, and define $s := d_L(x)$. Then 1 < t < 4 and 0 < s < 2. Note that $t+s \ge 2$, since a 4-vertex linear forest with one component contains no isolated vertices. We consider two cases. In each case we will describe the length and type of t path components, P^1, \ldots, P^t , each with an even number of vertices. Proposition 5.4 (applied repeatedly) together with (a),(b) will then imply that G'-x contains at least $5 \cdot 6^k \psi n$ disjoint L-compatible skeletons, each consisting exactly of t components isomorphic to P^1, \ldots, P^t . (We can apply Proposition 5.4) here since in each case $P^1 \cup \cdots \cup P^t$ will contain a perfect matching.)

Case 1: s = 2.

• For
$$1 \le r \le t-1$$
, P^r is a K_2 of type $\overline{N}_{\ell_r}^1(x)$, $\overline{N}_{\ell_r}^2(x)$.

• For
$$1 \le r \le t-1$$
, P^r is a K_2 of type $\overline{N}_{\ell_r}^1(x), \overline{N}_{\ell_r}^2(x)$.
• P^t is a $P_{2k-2t-2}$ of type $\overline{N}_{\ell_t}^1(x), \overline{N}_{\ell_t}^2(x), \overline{N}_{\ell_{t+1}}^1(x), \overline{N}_{\ell_{t+1}}^2(x), \dots, \overline{N}_{\ell_{k-2}}^1(x), \overline{N}_{\ell_{k-2}}^2(x)$.

Case 2: Either s = 1 or s = 0, t > 1.

• For
$$1 \le r \le 1 - s$$
, P^r is a K_2 of type $N^1_{\ell_r}(x)$, $\overline{N}^1_{\ell_r}(x)$.

•
$$P^{2-s}$$
 is a $P_{2k-2t-2}$ of type $N^1_{\ell_{2-s}}(x), \overline{N}^2_{\ell_{2-s}}(x), \overline{N}^1_{\ell_{3-s}}(x), \overline{N}^2_{\ell_{3-s}}(x), \overline{N}^1_{\ell_{3-s}}(x), \overline{N}^1_{\ell_{k-t-s}}(x), \overline{N}^1_{\ell_{k-t-s}}(x), \overline{N}^1_{\ell_{k-t-s}}(x)$

• For
$$k-t-s+1 \le r \le k-2$$
, P^r is a K_2 of type $\overline{N}_{\ell_r}^1(x), \overline{N}_{\ell_r}^2(x)$.

Since $t+s\geq 2$, this covers all cases. Now fix a set SK of $5\cdot 6^k \psi n$ disjoint L-compatible skeletons in G'-x, and let $H \in SK$. Let h_H denote the number of possibilities for a set E^* of edges between Y and V(H). Note that such a set E^* cannot equal $E_{L,H}$, since G needs to be induced- C_{2k} -free. Thus $h_H \leq 2^{|Y||V(H)|} - 1 = 2^{6(k-2)} - 1$. Note that by (F2)_u the number of vertices outside Q_i that are not contained in some graph $H \in SK$ is at most $(k-2)n/(k-1) + \mu n - 10(k-2)6^k \psi n$. Hence,

$$b_{4} \leq 2^{3(k-2)n/(k-1)-30(k-2)6^{k}} \psi^{n+3\mu n} \prod_{H \in SK} h_{H}$$

$$\leq 2^{3(k-2)n/(k-1)-30(k-2)6^{k}} \psi^{n+3\mu n} \left(2^{6(k-2)} \left(1 - 2^{-6(k-2)} \right) \right)^{5 \cdot 6^{k}} \psi^{n}$$

$$\leq 2^{3(k-2)n/(k-1)} 2^{3\mu n} e^{-5 \cdot 6^{k}} \psi^{n/(2^{6(k-2)})} \leq 2^{3(k-2)n/(k-1)} 2^{\mu^{1/2} n} 2^{-\psi^{n/11^{k}}},$$

as required.

Claim 5: $c_1 \le k^2 n$.

Claim 6: $c_2 \leq C 2^{\mu^{1/2} n} f(n_k) 2^{t_{k-1}(n-1)}$.

Claim 7: $c_3 \le 2^{7\xi(\alpha)n}$.

Indeed, for every graph $\tilde{G} \in F_{Q,(ii)}^1$ for which x is both i-light and j-light, we have that, for every $\ell \in \{i,j\}$, either $\min\{|N_{Q_\ell}(x)|, |\overline{N}_{Q_\ell}(x)|\} \leq \alpha n$ or else there exists a vertex $z \neq x$ such that $|N_\ell^*(x,z)| + |N_\ell^*(z,x)| \leq \alpha n$.

For $\ell \in \{i, j\}$, let $h(\ell, 1)$ denote the number possibilities for a set of edges in G between $\{x\}$ and $Q_{\ell} \setminus \{x\}$ such that $\min\{|N_{Q_{\ell}}(x)|, |\overline{N}_{Q_{\ell}}(x)|\} \leq \alpha n$. Then $h(\ell, 1) \leq 2\binom{n}{\leq \alpha n} \leq 2^{\xi(\alpha)n+1}$. For $\ell \in \{i, j\}$, let $h(\ell, 2)$ denote the number possibilities for a set of edges between $\{x\}$ and $Q_{\ell} \setminus \{x\}$ such that there exists a vertex $z \neq x$ such that $|N_{\ell}^*(x, z)| + |N_{\ell}^*(z, x)| \leq \alpha n$. Then $h(\ell, 2) \leq n\binom{|N_{Q_{\ell}}(z)|}{\leq \alpha n}\binom{|\overline{N}_{Q_{\ell}}(z)|}{\leq \alpha n}$ $\leq 2^{3\xi(\alpha)n}$.

Hence

$$c_3 \le (h(i,1) + h(i,2))(h(j,1) + h(j,2)) \le (2^{\xi(\alpha)n+1} + 2^{3\xi(\alpha)n})^2 \le 2^{7\xi(\alpha)n}$$

as required.

Claim 8: $c_4 \leq 2^{(k-3)n/(k-1)}2^{2\mu n}$.

Indeed, since the number of possible edges between $\{x\}$ and $[n]\setminus (Q_i \cup Q_j \cup \{x\})$ is at most $(k-3)n/(k-1)+2\mu n$, we have that $c_4 \leq 2^{(k-3)n/(k-1)+2\mu n}$, as required.

Now (7.2) together with Claims 1–4 and Proposition 4.3(ii) implies that

$$(7.4) |F_{Q,(i)}^{1}|$$

$$\leq 2 \max_{\psi \in \{\beta^{2}, \beta^{1/2}\}} \left\{ 2^{6}kn^{4} \cdot C2^{\mu^{1/2}n} f_{k}(n_{k}) 2^{t_{k-1}(n-3)} \cdot 2^{4\psi^{3/2}n} \cdot 2^{\frac{3(k-2)n}{k-1}} 2^{\mu^{1/2}n} 2^{-\frac{\psi n}{11^{k}}} \right\}$$

$$\leq \max_{\psi \in \{\beta^{2}, \beta^{1/2}\}} \left\{ Cf_{k}(n_{k}) 2^{t_{k-1}(n-3) + \frac{3(k-2)n}{k-1}} 2^{-\frac{\psi n}{12^{k}}} \right\} \leq Cf_{k}(n_{k}) 2^{t_{k-1}(n)} 2^{-\frac{\beta^{2}n}{13^{k}}}.$$

Similarly, (7.3) together with Claims 5–8 and Proposition 4.3(ii) implies that

$$(7.5) |F_{Q,(ii)}^{1}| \leq k^{2}n \cdot C2^{\mu^{1/2}n} f_{k}(n_{k}) 2^{t_{k-1}(n-1)} \cdot 2^{7\xi(\alpha)n} \cdot 2^{\frac{(k-3)n}{k-1}} 2^{2\mu n}$$

$$\leq C2^{\mu^{1/3}n} f_{k}(n_{k}) 2^{t_{k-1}(n-1) + \frac{(k-2)n}{k-1}} 2^{-\frac{n}{k-1}} 2^{7\xi(\alpha)n} \leq Cf_{k}(n_{k}) 2^{t_{k-1}(n)} 2^{-\frac{n}{k}}.$$

Now (7.1) together with (7.4) and (7.5) implies that

$$|F_Q^1| \le Cf_k(n_k)2^{t_{k-1}(n)} \left(2^{-\frac{\beta^2 n}{13^k}} + 2^{-\frac{n}{k}}\right) \le Cf_k(n_k)2^{t_{k-1}(n)}2^{-\frac{\beta^2 n}{14^k}},$$

as required.

8. Estimation of $|F_Q^2|$

Given $G \in F_Q^2 \cup F_Q^3$ and $i \in \{0, 1, \dots, k-2\}$, let $A_G^i := \{x \in Q_i : \overline{d}_{G,Q}^i(x), d_{G,Q}^i(x) \geq \beta n\}$. The key result of this section (Lemma 8.5) states that A_G^i has bounded size. To prepare for this, we will classify the pairs of vertices in A_G^i according to their (non-)neighbourhood intersection pattern. The fact that $G \notin F_Q^1$ allows us to observe some restrictions on these patterns (see Propositions 8.3 and 8.4). In the proof of Lemma 8.5 we use a Ramsey argument to restrict our view to one abundant type of pattern. This quickly leads to a contradiction if $|A_G^i|$ is large. Using the fact that $G \notin F_Q^1$ we show that the remainder of each class (i.e. $G[Q_i \backslash A_G^i]$) induces a very simple structure

(Proposition 8.2). We translate this structural information into a sufficiently strong bound on the number of graphs in F_G^2 , in Lemma 8.7.

Let \mathcal{L} denote the collection of all 4-vertex linear forests. The following proposition is an analogue of Proposition 3.8(i) that can be applied to graphs rather than 2-coloured multigraphs. It follows immediately from Proposition 3.8(i).

Proposition 8.1. Let G be a graph such that for every $H \in \mathcal{L}$, G is induced H-free. Then \overline{G} is a disjoint union of stars and triangles.

Proposition 8.2. Let $G \in F_Q^2 \cup F_Q^3$ and $i \in \{0, 1, ..., k-2\}$. Then $\overline{G}[Q_i \setminus A_G^i]$ is a disjoint union of stars and triangles.

Proof. Suppose for a contradiction that $\overline{G}[Q_i \setminus A_G^i]$ is not a disjoint union of stars and triangles. Then Proposition 8.1 implies that $\overline{G}[Q_i \setminus A_G^i]$ contains an induced copy of a graph in \mathcal{L} , with vertex set $\{x, y_1, y_2, y_3\}$ say. We will show that $\{x, y_1, y_2, y_3\}$ is a $(k, x, i, \beta^{1/2})$ -configuration, which contradicts the fact that $G \notin F_Q^1$. Note that $G[\{x, y_1, y_2, y_3\}]$ is a linear forest, and so $\{x, y_1, y_2, y_3\}$ satisfies (C1). By the definition of A_G^i we have that $\min\{d_{G,Q}^i(y_j), \overline{d}_{G,Q}^i(y_j)\} \leq \beta n$ for all $j \in [3]$, and so $\{x, y_1, y_2, y_3\}$ satisfies (C4). Since $G \notin F_Q^1$, x is j-light for at most one index $j \in \{0, 1, \dots, k-2\}$. Since $x \in Q_i \setminus A_G^i$, x is i-light. Thus for every $j \in \{0, 1, \dots, k-2\}$ with $i \neq j$ we have that x is not j-light, and hence $d_{G,Q}^j(x), \overline{d}_{G,Q}^j(x) > \alpha n > 13 \cdot 6^k \cdot \beta^{1/2} n$, and so $\{x, y_1, y_2, y_3\}$ satisfies (C2) and (C3). Therefore $\{x, y_1, y_2, y_3\}$ is a $(k, x, i, \beta^{1/2})$ -configuration, as required.

The following definitions will be useful in order to show that $|A_G^i|$ is small. Suppose S is a star or triangle. If S is a star on at least three vertices, we call the unique vertex in S of degree greater than one the *centre* of S. Otherwise we call the vertex of S with the smallest label the centre of S. Let $G \in F_Q^2 \cup F_Q^3$ and $i, j \in \{0, 1, \ldots, k-2\}$ and let $x, y \in A_G^i$.

- We say x, y are j-irregular if $|\overline{N}_{j}(\{x, y\})| \leq \gamma n$.
- We say x, y are j-asymmetric if $|N_j^*(x, y)| + |N_j^*(y, x)| > 3\gamma n$ and either $|N_j^*(x, y)| \le \gamma n$ or $|N_j^*(y, x)| \le \gamma n$.
- We say x, y are j-identical if $|N_j^*(x, y)| + |N_j^*(y, x)| \le 3\gamma n$.

Roughly speaking, if one of the above holds then the neighbourhoods of x, y do not behave in a 'random' like way (thus constraining the number of possibilities for choosing the neighbourhoods). The following statement follows immediately from the above definitions and the fact that $\gamma \ll \alpha$.

(8.1) If
$$x, y$$
 are j -identical then x, y are both j -light.

Proposition 8.3. Let $G \in F_Q^2 \cup F_Q^3$ and $i \in \{0, 1, ..., k-2\}$ and let $x, y \in A_G^i$. Then x, y are j-identical for at most one index $j \in \{0, 1, ..., k-2\}$.

Proof. Suppose x, y are j-identical for some $j \in \{0, 1, \dots, k-2\}$ and suppose $j' \in \{0, 1, \dots, k-2\}$ with $j' \neq j$. It suffices to show that x, y are not j'-identical. Note that x is j-light by (8.1). Since $G \notin F_Q^1$, x is j''-light for at most one index $j'' \in \{0, 1, \dots, k-2\}$. Thus x is not j'-light, and hence by (8.1) x, y are not j'-identical, as required.

Proposition 8.4. Let $G \in F_Q^2 \cup F_Q^3$ and $i \in \{0, 1, ..., k-2\}$ and let $x, y \in A_G^i$. Then there exists an index $j \in \{0, 1, ..., k-2\}$ such that x, y are j-irregular or j-asymmetric (or both).

Proof. Suppose for a contradiction that for every index $\ell \in \{0, 1, \dots, k-2\}$, x, y are neither ℓ -irregular nor ℓ -asymmetric. Since, by Proposition 8.3, x, y are j-identical for at most one index j, and $k \geq 6$, we may assume without loss of generality that x, y are not ℓ -identical for $\ell \in \{1, 2, 3\}$. We consider the following two cases.

Case 1: x, y are adjacent.

In this case we define sets A_{ℓ} , B_{ℓ} for $\ell \in \{0, 1, \dots, k-2\}$ as follows. We will use these sets to extend x, y into an induced copy of C_{2k} .

- Let $A_1 := N_1^*(x, y)$ and $B_1 := \overline{N}_1(\{x, y\})$.
- Let $A_2 := N_2^*(y, x)$ and $B_2 := \overline{N}_2(\{x, y\})$.
- For every $\ell \in \{0, 1, \dots, k-2\} \setminus \{1, 2\}$, let $A_{\ell}, B_{\ell} \subseteq \overline{N}_{\ell}(\{x, y\})$ be disjoint and satisfy $|A_{\ell}|, |B_{\ell}| \ge ||\overline{N}_{\ell}(\{x, y\})|/2|$.

Since for every $\ell \in \{0, 1, \dots, k-2\}$ x, y are neither ℓ -irregular nor ℓ -asymmetric, and for every $\ell \in \{1, 2\}$ x, y are not ℓ -identical, we have that $|A_{\ell}|, |B_{\ell}| \geq \gamma n/3$ for every $\ell \in \{0, 1, \dots, k-2\}$. This together with Proposition 5.4 and the fact that $\mu \ll \gamma$ implies that there exists in G an induced copy of P_{2k-2} of type $A_1, B_1, A_0, B_0, A_3, B_3, \dots, A_{k-2}, B_{k-2}, B_2, A_2$. By the definition of the sets A_{ℓ}, B_{ℓ} , the vertices of this P_{2k-2} together with x, y induce on G a copy of C_{2k} . This contradicts the fact that $G \in F_Q(n, k)$.

Case 2: x, y are not adjacent.

In this case we define sets A_{ℓ} , B_{ℓ} for $\ell \in \{0, 1, ..., k-2\}$ as follows. Similarly to the previous case, we will find an induced C_{2k} which contains x, y together with exactly one vertex from each of these sets.

- Let $A_1 := N_1^*(x, y)$ and $B_1 := N_1^*(y, x)$.
- Let $A_2 := N_2^*(x, y)$ and $B_2 := \overline{N}_2(\{x, y\})$.
- Let $A_3 := N_3^*(y, x)$ and $B_3 := \overline{N}_3(\{x, y\})$.
- For every $\ell \in \{0, 1, \dots, k-2\} \setminus \{1, 2, 3\}$, let $A_{\ell}, B_{\ell} \subseteq \overline{N}_{\ell}(\{x, y\})$ be disjoint and satisfy $|A_{\ell}|, |B_{\ell}| \ge ||\overline{N}_{\ell}(\{x, y\})|/2|$.

Since for every $\ell \in \{0, 1, ..., k-2\}$ x, y are neither ℓ -irregular nor ℓ -asymmetric, and for every $\ell \in \{1, 2, 3\}$ x, y are not ℓ -identical, we have that $|A_{\ell}|, |B_{\ell}| \ge \gamma n/3$ for every $\ell \in \{0, 1, ..., k-2\}$. As before, this together with Proposition 5.4 implies that there exists in G an induced copy of the graph H that consists of the following two components:

- One P_{2k-4} of type $A_2, B_2, A_0, B_0, A_4, B_4, \dots, A_{k-2}, B_{k-2}, B_3, A_3$.
- One K_2 of type A_1, B_1 .

By the definition of the sets A_{ℓ} , B_{ℓ} , the vertices of H together with x, y induce on G a copy of C_{2k} . This contradicts the fact that $G \in F_O(n, k)$.

This covers all cases, and hence completes the proof.

Recall from Section 6 that $M := R_{2k-2}(\lceil \frac{1}{\gamma} \rceil) + 1$.

Lemma 8.5. Let $G \in F_Q^2 \cup F_Q^3$ and $i \in \{0, 1, ..., k-2\}$. Then $|A_G^i| < M$.

Proof. Suppose for a contradiction that $|A_G^i| \geq M$. Consider an auxiliary complete graph H_i with $V(H_i) = A_G^i$. We define a (2k-2)-edge-colouring \mathcal{C} of H_i with colours $\{a_0, b_0, a_1, b_1, \ldots, a_{k-2}, b_{k-2}\}$ as follows.

- For every $j \in \{0, 1, ..., k-2\}$, an edge $xy \in E(H)$ is coloured a_j if x, y are j-irregular and for every $j' \in \{0, 1, ..., k-2\}$ with j' < j, x, y are not j'-irregular.
- An edge $xy \in E(H)$ that was not coloured in the previous step is coloured b_j if x, y are j-asymmetric, and for every $j' \in \{0, 1, \dots, k-2\}$ with j' < j, x, y are not j'-asymmetric.

Note that by Proposition 8.4, every edge is coloured by a unique colour in \mathcal{C} .

Now since $M > R_{2k-2}(\lceil 1/\gamma \rceil)$, H_i contains a monochromatic clique of size at least $1/\gamma$. Let $X = \{x_1, x_2, \dots, x_{\lceil 1/\gamma \rceil}\}$ be the vertex set of such a monochromatic clique. We consider the following two cases.

Case 1: X has colour a_j for some $j \in \{0, 1, ..., k-2\}$.

In this case every pair of vertices in X is j-irregular, by definition of \mathcal{C} . Let $X' := \{x_1, x_2, \dots, x_{\lceil \beta/2\gamma \rceil}\}$ and suppose $z, z' \in X'$. By the definition of j-irregularity, $|\overline{N}_j(z) \cap \overline{N}_j(z')| \leq \gamma n$. Note also that $|\overline{N}_j(z)| \geq \beta n$ by Proposition 5.1 and the fact that $z \in A_G^i$. So by the inclusion-exclusion principle,

$$2n/(k-1) \ge n/(k-1) + \mu n \ge |Q_j| \ge \sum_{z \in X'} |\overline{N}_j(z)| - \sum_{\substack{z,z' \in X' \\ z \ne z'}} |\overline{N}_j(z) \cap \overline{N}_j(z')|$$

$$\geq \beta \lceil \beta/2\gamma \rceil n - \lceil \beta^2/(4\gamma^2) \rceil \gamma n \geq \beta^2 n/5\gamma > 2n/(k-1),$$

where the last inequality follows from the fact that $\gamma \ll \beta$. This is a contradiction.

Case 2: X has colour b_j for some $j \in \{0, 1, \dots, k-2\}$.

In this case every pair of vertices in X is j-asymmetric, by definition of C. Suppose $\ell, \ell' \in [[1/\gamma]]$ are distinct. By the definition of j-asymmetry, exactly one of the following holds.

- (a) $|N_j^*(x_\ell, x_{\ell'})| \le \gamma n$ and $|N_j^*(x_{\ell'}, x_\ell)| > 2\gamma n$. (b) $|N_j^*(x_{\ell'}, x_\ell)| \le \gamma n$ and $|N_j^*(x_\ell, x_{\ell'})| > 2\gamma n$.

Consider the auxiliary tournament T with V(T) = X and $E(T) = \{ \overrightarrow{x_\ell x_{\ell'}} : \ell, \ell' \text{ satisfy (a)} \}$. By Redei's theorem every tournament contains a directed Hamilton path. So, by relabelling the indices if necessary, we may assume that $\overline{x_\ell x_{\ell+1}} \in E(T)$ for every $\ell \in [\lceil 1/\gamma \rceil - 1]$. Thus for every $\ell \in [\lceil 1/\gamma \rceil - 1]$. $\lceil \lceil 1/\gamma \rceil - 1 \rceil$,

$$|\overline{N}_j(x_{\ell+1})| = |\left(\overline{N}_j(x_{\ell}) \setminus N_j^*(x_{\ell+1}, x_{\ell})\right) \cup N_j^*(x_{\ell}, x_{\ell+1})| \le |\overline{N}_j(x_{\ell})| - 2\gamma n + \gamma n \le |\overline{N}_j(x_{\ell})| - \gamma n.$$

Hence,

$$|\overline{N}_j(x_{\lceil 1/\gamma \rceil})| \le |\overline{N}_j(x_1)| - \left(\frac{1}{\gamma} - 1\right) \cdot \gamma n \le |Q_j| - (1 - \gamma)n < 0,$$

which is a contradiction.

This covers all cases, and hence completes the proof.

Suppose $G \in F_Q^2$ and $i \in \{0, 1, \dots, k-2\}$. By Proposition 8.2, $\overline{G}[Q_i \setminus A_G^i]$ is a disjoint union of stars and triangles. Let S be the set of components of $\overline{G}[Q_i \setminus A_G^i]$ with the largest number of vertices. Let S^{\diamond} be the component in S whose centre c has the smallest label. Define $Y_i = Y_i(G,Q)$ to be the set of all isolated vertices in $\overline{G}[Q_i \setminus A_G^i]$ together with all vertices in $V(S^{\diamond}) \setminus \{c\}$.

Lemma 8.6. Let $G \in F_Q^2$ and $i \in \{0, 1, ..., k-2\}$. Then $|Y_i| \ge 10n/\log n$.

Proof. Define $s := \lceil 10n/\log n \rceil$. Suppose for a contradiction that $|Y_i| < s$. Since $G \in F_Q^2$, there exists an index $i' \in \{0, 1, \dots, k-2\}$ such that $|A_G^{i'}| > 0$. Let $x \in A_G^{i'}$. The definition of $A_G^{i'}$ together with Proposition 5.1 implies that $|\overline{N}_{Q_j}(x)| \ge \beta n$ for every $j \in \{0, 1, \dots, k-2\}$. This together with Lemma 8.5 implies that $|\overline{N}_{Q_i}(x) \setminus A_G^i| \geq \beta n - M > 2s$. Also, since $|Y_i| < s$, at most s components in $\overline{G}[Q_i \setminus A_G^i]$ are isolated vertices and every component in $\overline{G}[Q_i \setminus A_G^i]$ has order at most s. Thus there are at least two non-trivial components S, S' of $\overline{G}[Q_i \setminus A_G^i]$ that each contain a non-neighbour of x.

Since S is a non-trivial component of $\overline{G}[Q_i \setminus A_G^i]$ there exist vertices $y, y' \in S$ such that $xy, yy' \notin S$ $E(G[Q_i])$. Let $y'' \in S'$ be such that $xy'' \notin E(G[Q_i])$. Since y'' belongs to a different component of $\overline{G}[Q_i \setminus A_G^i]$ to y and y', it follows that $yy'', y'y'' \in E(G[Q_i])$. Thus,

(8.2)
$$E(G[\{x, y, y', y''\}]) \in \{\{yy'', y'y''\}, \{xy', yy'', y'y''\}\}.$$

Claim: $\{x, y, y', y''\}$ is a (k, x, i, β^2) -configuration.

Indeed, by (8.2), $G[\{x,y,y',y''\}]$ is a linear forest and so $\{x,y,y',y''\}$ satisfies (C1). As observed above, $\overline{d}_{G,Q}^j(x) \geq \beta n > 13 \cdot 6^k \beta^2 n$ for every $j \in \{0,1,\ldots,k-2\}$, and so $\{x,y,y',y''\}$ satisfies (C2). Since $G \notin F_Q^1$, there do not exist distinct $j,j' \in \{0,1,\ldots,k-2\}$ such that x is both j-light and j'-light. So there exists $j \in \{0,1,\ldots,k-2\}$ such that for every $j' \in \{0,1,\ldots,k-2\}$ with $j' \neq j$, $d_{G,Q}^{j'}(x) > \alpha n > 13 \cdot 6^k \beta^2 n$, and so $\{x,y,y',y''\}$ satisfies (C3). Since S,S' each contain at most s vertices, y,y',y'' each have at most s non-neighbours in $G[Q_i \setminus A_G^i]$. This together with Lemma 8.5 implies that y,y',y'' each have at most $s+M \leq \beta^4 n$ non-neighbours in $G[Q_i]$, and so $\{x,y,y',y''\}$ satisfies (C4). Hence $\{x,y,y',y''\}$ is a (k,x,i,β^2) -configuration, as required.

The above claim contradicts the fact that $G \notin F_Q^1$, and hence completes the proof.

Lemma 8.6 guarantees a large set of vertices in each class Q_i (namely Y_i) with an extremely restricted (non-)neighbourhood. This is the key idea in our estimation of $|F_Q^2|$.

Lemma 8.7. $|F_Q^2| \le C2^{-n} f_k(n_k) 2^{t_{k-1}(n)}$.

Proof. Define $s := \lceil 10n/\log n \rceil$. Since by Lemma 8.6 $|Y_i(G,Q)| \ge s$ for every graph $G \in F_Q^2$, any graph $G \in F_Q^2$ can be constructed as follows.

- First we choose sets $S_{\ell} \subseteq Q_{\ell}$ such that $|S_{\ell}| = s$, for every $\ell \in \{0, 1, \dots, k-2\}$. Let b_1 denote the number of such choices.
- Next we choose the graph G' on $[n] \setminus \bigcup_{\ell \in \{0,1,\dots,k-2\}} S_{\ell}$ such that $G[[n] \setminus \bigcup_{\ell \in \{0,1,\dots,k-2\}} S_{\ell}] = G'$. Let b_2 denote the number of possibilities for G'.
- Next we choose the set E' of internal edges of G that are incident to at least one vertex in $\bigcup_{\ell \in \{0,1,\ldots,k-2\}} S_{\ell}$ in such a way that the resulting graph G will satisfy $S_{\ell} \subseteq Y_{\ell}(G,Q)$ for every $\ell \in \{0,1,\ldots,k-2\}$. Let b_3 denote the number of possibilities for E'.
- Finally we choose the set E'' of crossing edges of G that are incident to at least one vertex in $\bigcup_{\ell \in \{0,1,\ldots,k-2\}} S_{\ell}$. Let b_4 denote the number of possibilities for E''.

Hence,

$$(8.3) |F_Q^2| \le b_1 \cdot b_2 \cdot b_3 \cdot b_4.$$

The following series of claims will give upper bounds for the quantities b_1, \ldots, b_4 . The proof of Claim 2 is almost identical to that of Claim 2 in Lemma 7.3; we give proofs of the others.

Claim 1: $b_1 \leq 2^n$.

Indeed,

$$b_1 \le {n \choose \left\lceil \frac{10n}{\log n} \right\rceil}^{k-1} \le {\left(\left(\frac{e \log n}{10} \right)^{\frac{10n}{\log n}} \right)}^{k-1} \le 2^n,$$

as required.

Claim 2: $b_2 \leq C 2^{\mu^{1/2} n} f_k(\lceil n/(k-1) - s \rceil) 2^{t_{k-1}(n-(k-1)s)}$.

Claim 3: $b_3 \le 2^n$.

Indeed, for every graph $G^* \in F_Q^2$ for which $S_\ell \subseteq Y_\ell(G^*,Q)$ for every $\ell \in \{0,1,\ldots,k-2\}$, let $G_{B,\ell}^* := \overline{G^*}[Q_\ell \backslash A_{G^*}^\ell]$. Then each S_ℓ consists of isolated vertices in $G_{B,\ell}^*$ as well as non-centre vertices of a single component \tilde{C} of $G_{B,\ell}^*$. (Note that \tilde{C} is a star or triangle in $G_{B,\ell}^*$, with some centre $u \in Q_\ell \backslash (A_{G^*}^\ell \cup S_\ell)$.) By Lemma 8.5, we also have that $|A_{G^*}^\ell| \leq M$.

Hence $b_3 \leq \prod_{\ell=0}^{k-2} \prod_{j=1}^3 h(\ell,j)$, where the quantities $h(\ell,j)$ are defined as follows. Let $h(\ell,1)$ denote the number of ways to choose a set $\tilde{A}^{\ell} \subseteq Q_{\ell} \backslash S_{\ell}$ of size at most M. (In what follows \tilde{A}^{ℓ} will

play the role of A_G^i .) Then $h(\ell,1) \leq n^M$. Given such a set \tilde{A}^ℓ , let $h(\ell,2)$ denote the number of ways to choose \tilde{C} . Then $h(\ell,2) \leq n2^{|S_\ell|} + n^3 \leq n2^{s+1}$. (Indeed, if \tilde{C} is a star we have at most n choices for the centre u, and for every vertex $v \in S_\ell$ we can choose whether v is adjacent to u or not; if \tilde{C} is a triangle we have at most n^3 choices for its vertices.) Given a set \tilde{A}^ℓ as above, let $h(\ell,3)$ denote the number of possible sets of edges between S_ℓ and \tilde{A}^ℓ . Then $h(\ell,3) \leq 2^{|S_\ell|}|\tilde{A}^\ell| \leq 2^{sM}$. Hence

$$b_3 \le \prod_{\ell=0}^{k-2} \prod_{j=1}^3 h(\ell,j) \le (n^M \cdot n2^{s+1} \cdot 2^{sM})^{k-1} \le 2^n,$$

as required.

Claim 4: $b_4 \leq 2^{(k-2)sn - \binom{k-1}{2}s^2}$.

Indeed, note that, for a fixed index $\ell \in \{0, 1, \dots, k-2\}$, the number h_{ℓ} of possible crossing edges in G that are incident to a vertex in S_{ℓ} is at most $s(n-|Q_{\ell}|)$. Also, the number of possible crossing edges in G that are incident to two vertices in $\bigcup_{\ell \in \{0,1,\dots,k-2\}} S_{\ell}$ is exactly $\binom{k-1}{2} s^2$. Hence,

$$b_4 \le 2^{\sum_{\ell=0}^{k-2} h_\ell} 2^{-\binom{k-1}{2}s^2} \le 2^{(k-2)sn - \binom{k-1}{2}s^2},$$

as required.

Note that $t_{k-1}(s(k-1)) = {k-1 \choose 2}s^2$ and that by Lemma 4.2, $f_k(n_k) \ge s^{s/2}f_k(\lceil n/(k-1) - s \rceil) \ge 2^{4n}f_k(\lceil n/(k-1) - s \rceil)$. These observations together with (8.3), Claims 1–4 and Proposition 4.3(ii) imply that

$$|F_Q^2| \le 2^n \cdot C2^{\mu^{1/2}n} f_k(\lceil n/(k-1) - s \rceil) 2^{t_{k-1}(n-(k-1)s)} \cdot 2^n \cdot 2^{(k-2)sn - \binom{k-1}{2}s^2}$$

$$\le C2^{3n} 2^{-4n} f_k(n_k) 2^{t_{k-1}(n-(k-1)s) + (k-2)sn - s(k-1)(k-2) - t_{k-1}(s(k-1))}$$

$$\le C2^{-n} f_k(n_k) 2^{t_{k-1}(n)},$$

as required.

9. Estimation of $|F_Q^3|$

The information we have gained so far allows us to easily deduce that every $G \in F_Q^3$ is extremely close to being a k-template (see Proposition 9.1). One advantage of this is that it allows us to use more precise estimates when applying induction (see Corollary 9.2).

Proposition 9.1. Let $G \in F_Q^3$ and $i \in \{0, 1, ..., k-2\}$. Then the following hold.

- (i) $\overline{G}[Q_i]$ is a disjoint union of stars and triangles.
- (ii) G contains at most n internal non-edges.
- (iii) Every vertex $x \in Q_i$ satisfies $\overline{d}_{G,Q}^i(x) < \beta n$.

Proof.

- (i) Since $G \notin F_Q^2$, every vertex $x \in Q_i$ satisfies $\min\{d_{G,Q}^i(x), \overline{d}_{G,Q}^i(x)\} < \beta n$. Thus $A_G^i = \emptyset$, and so by Proposition 8.2, $\overline{G}[Q_i]$ is a disjoint union of stars and triangles.
- (ii) This follows immediately from (i).
- (iii) Let $x \in Q_i$. Let us first show that $d^i_{G,Q}(x) \ge \beta n$. Suppose not. Then $\overline{d}^i_{G,Q}(x) = |Q_i| d^i_{G,Q}(x) 1 > |Q_i| \beta n 1 \ge n/(k-1) \mu n \beta n 1$. Thus for every $j \in \{0, 1, \dots, k-2\}$

with $j \neq i$, Proposition 5.1 implies that

$$d_{G,Q}^{j}(x) = |Q_{j}| - \overline{d}_{G,Q}^{j}(x) \le |Q_{j}| - \overline{d}_{G,Q}^{i}(x) < \left(\frac{n}{k-1} + \mu n\right) - \left(\frac{n}{k-1} - \mu n - \beta n - 1\right)$$

$$= \beta n + 2\mu n + 1 < \alpha n,$$

where the last inequality follows from the fact that $\mu, \beta \ll \alpha$. Thus x is both i-light and j-light, which contradicts the fact that $G \notin F_Q^1$. Thus $d_{G,Q}^i(x) \geq \beta n$. This together with the fact (observed in the proof of (i), above) that $A_G^i = \emptyset$ implies that $\overline{d}_{G,Q}^i(x) < \beta n$, as required.

Recall the definition of property $(F1)_{\nu}$ in Section 5. We define $T_Q^*(n,k) \subseteq F_Q^3$ to be the set of all (labelled) graphs in F_Q^3 that satisfy property $(F1)_{(40n \log n)^{1/2}/n}$ with respect to Q. Proposition 9.1(ii) together with Lemma 5.2(i) applied with $(40n \log n)^{1/2}/n$, n playing the roles of ν , m respectively implies that

$$(9.1) |F_Q^3 \setminus T_Q^*(n,k)| \le 2^{t_{k-1}(n) - n\log n/5}.$$

So (9.1) allows us to restrict our attention to the class $T_Q^*(n,k)$. In particular, this allows us to apply property $(F1)_{\nu}$ to much smaller vertex sets than in the preceding sections. This in turn gives us a much better bound on the number of partitions that may arise after deleting a small number of vertices. More precisely, Lemma 5.5 applied with $(40n \log n)^{1/2}/n$, n playing the roles of ν , m respectively implies the following result. Recall that $\mathcal{P}(Q,s)$ was defined before (5.3).

Corollary 9.2. Let $S \subseteq [n]$ with $|S| \le n/k^2$. Then for every $G \in T_Q^*(n,k)$, every optimal ordered (k-1)-partition of G-S is an element of $\mathcal{P}(Q-S,40k^4\log n)$.

In order to estimate $|T_Q^*(n,k)|$ (and thus $|F_Q^3|$) we will further split $T_Q^*(n,k)$ into four classes $\mathcal{A}_1, \ldots, \mathcal{A}_4$. To define these classes we require some further notation. We say that G contains a (6,3)-forest with respect to Q if there exist distinct indices $i, j \in \{0,1,\ldots,k-2\}$ such that there exist six vertices in $Q_i \cup Q_j$ that induce on G a linear forest with at most three components. A (6,3)-forest has potential extensions into an induced C_{2k} , so its existence in every $G \in \mathcal{A}_3$ (see below) constrains the possible edge sets for G (and thus the number of choices for G). To obtain a significant constraint on the possible edge sets however, we first need to exclude the situations that arise in the classes \mathcal{A}_1 and \mathcal{A}_2 , described below. These involve the structure of the stars of the complement graph inside the vertex classes, so to describe these classes of graphs recall that the centres of stars and triangles were defined before Proposition 8.3. Given a graph $G \in F_Q^3$ and an index $i \in \{0, 1, \ldots, k-2\}$ we define the following sets.

- $C^{i}(G,Q)$ is the set of all centres of triangles and non-trivial stars in $\overline{G}[Q_{i}]$.
- $C_{high}^i(G,Q)$ is the set of all centres of stars in $\overline{G}[Q_i]$ of order at least $n^{1-\frac{1}{2k^2}}/200k^2$.
- $B_{high}^{i}(G,Q)$ is the set of all vertices in Q_i which have a non-neighbour in C_{high}^i .
- $C_{low}^i(G,Q)$ is the set of all centres of triangles and non-trivial stars in $\overline{G}[Q_i]$ of order less than $n^{1-\frac{1}{2k^2}}/200k^2$.
- $B_{low}^i(G,Q)$ is the set of all vertices in Q_i which have a non-neighbour in C_{low}^i .
- $C_0^i(G,Q)$ is the set of all isolated vertices in $\overline{G}[Q_i]$.

We may sometimes write C^i for $C^i(G,Q)$ when the graph G and ordered (k-1)-partition Q we consider are clear from the context (and similarly for $C^i_{high}, B^i_{high}, C^i_{low}, B^i_{low}, C^i_0$). Note that Proposition 9.1(i) implies that $C^i_{high}, B^i_{high}, C^i_{low}, B^i_{low}, C^i_0$ form a partition of Q_i . Given a subset $B \subseteq B^i_{low}$, we denote by C(B) the set of all vertices in C^i_{low} that have a non-neighbour in B.

We partition $T_Q^*(n,k)$ into the sets A_1, \ldots, A_4 defined as follows.




Figure 1: An illustration of \overline{G} for $G \in A_2$. Note that (9.2) implies that at most one of xy_1, xy_2 is an edge in \overline{G} .

- \mathcal{A}_1 is the set of all graphs $G \in T_Q^*(n,k)$ for which there exist distinct indices $i,j \in \{0,1,\ldots,k-2\}$ such that $|B_{low}^i| \geq n/2k^2$ and there exist distinct vertices $y_1,y_2,y_3 \in Q_j$ that satisfy $|\overline{N}(\{y_1,y_2,y_3\}) \cap B_{low}^i| \leq n/200k^2$.
 \mathcal{A}_2 is the set of all graphs $G \in T_Q^*(n,k) \setminus \mathcal{A}_1$ for which there exist distinct indices $i,j \in \mathcal{A}_2$
- \mathcal{A}_2 is the set of all graphs $G \in T_Q^*(n,k) \setminus \mathcal{A}_1$ for which there exist distinct indices $i,j \in \{0,1,\ldots,k-2\}$ such that $|B_{low}^i| \geq n/2k^2$ and there exist distinct vertices $y_1,y_2,y_3 \in Q_j$ with $y_1,y_2 \notin C^j(G,Q)$ that satisfy

(9.2)
$$C(\overline{N}(\{y_1, y_2, y_3\}) \cap B_{low}^i) \cap \overline{N}(\{y_1, y_2\}) = \emptyset.$$

(See Figure 1.)

- \mathcal{A}_3 is the set of all graphs $G \in T_Q^*(n,k) \setminus (\mathcal{A}_1 \cup \mathcal{A}_2)$ such that G contains a (6,3)-forest with respect to Q.
- $\mathcal{A}_4 := T_O^*(n,k) \setminus (\mathcal{A}_1 \cup \mathcal{A}_2 \cup \mathcal{A}_3)$ is the set of all remaining graphs.

We will estimate the sizes of A_1, \ldots, A_4 separately. Lemma 9.3 below gives a bound on $|A_1|$. The idea of the proof of Lemma 9.3 is that in this case the neighbourhoods of y_1, y_2, y_3 are 'atypical', and hence a Chernoff estimate (see Claim 4) shows that graphs in A_1 are rare.

Lemma 9.3.
$$|\mathcal{A}_1| \leq C2^{-n/150k^2} f_k(n_k) 2^{t_{k-1}(n)}$$
.

Proof. Any graph $G \in \mathcal{A}_1$ can be constructed as follows.

- First we choose distinct indices $i, j \in \{0, 1, \dots, k-2\}$, distinct vertices $y_1, y_2, y_3 \in Q_j$, and a set E of edges between y_1, y_2, y_3 . Let b_1 denote the number of such choices. The choices in the next steps will be made such that G satisfies $|B_{low}^i| \ge n/2k^2$ and $|\overline{N}(\{y_1, y_2, y_3\}) \cap B_{low}^i| \le n/200k^2$.
- Next we choose the graph G' on vertex set $[n] \setminus \{y_1, y_2, y_3\}$ such that $G[[n] \setminus \{y_1, y_2, y_3\}] = G'$. Let b_2 denote the number of possibilities for G'.
- Next we choose the set E' of edges in G between $\{y_1, y_2, y_3\}$ and $Q_j \setminus \{y_1, y_2, y_3\}$. Let b_3 denote the number of possibilities for E'.
- Finally we choose the set E'' of edges in G between $\{y_1, y_2, y_3\}$ and $[n] \setminus Q_j$ such that E'' is compatible with our previous choices. Let b_4 denote the number of possibilities for E''.

Hence,

$$(9.3) |\mathcal{A}_1| \leq b_1 \cdot b_2 \cdot b_3 \cdot b_4.$$

The following series of claims will give upper bounds for the quantities b_1, \ldots, b_4 . Claim 1 is trivial; we give proofs of the others.

Claim 1: $b_1 \le 2^3 k^2 n^3$.

Claim 2: $b_2 \le C2^{2(\log n)^3} f_k(n_k) 2^{t_{k-1}(n-3)}$.

Indeed, note that for every graph $\tilde{G} \in \mathcal{A}_1$, Corollary 9.2 together with (5.3) implies that every optimal ordered (k-1)-partition of $\tilde{G}[[n] \setminus \{y_1, y_2, y_3\}]$ is contained in some set \mathcal{P} of size at most $2^{(\log n)^3}$. Since $G[[n] \setminus \{y_1, y_2, y_3\}]$ is clearly induced- C_{2k} -free, this together with (6.2) implies that

$$b_2 \le \sum_{Q' \in \mathcal{P}} |F_{Q'}(n-3,k)| \le 6C2^{(\log n)^3} 2^{6(\log n)^2} f_k(\lceil (n-3)/(k-1)\rceil) 2^{t_{k-1}(n-3)}$$

$$\le C2^{2(\log n)^3} f_k(n_k) 2^{t_{k-1}(n-3)},$$

as required.

Claim 3: $b_3 \le 2^{3\xi(\beta)n}$.

Indeed, for every graph $\tilde{G} \in \mathcal{A}_1$ and every $\ell \in [3]$, Proposition 9.1(iii) implies that $\overline{d}_{\tilde{G},Q}^{j}(y_{\ell}) < \beta n$. Thus

$$b_3 \le \binom{n}{\le \beta n}^3 \le 2^{3\xi(\beta)n},$$

as required.

Claim 4: $b_4 \le 2^{3((k-2)n/(k-1)+\mu n)-n/128k^2}$.

Consider the graph obtained by starting with the graph $([n], E(G') \cup E')$ and adding edges between $\{y_1, y_2, y_3\}$ and $[n] \setminus Q_j$ randomly, independently, with probability 1/2. Note that the number of graphs that this process can generate is at most $2^{3((k-2)n/(k-1)+\mu n)}$, with each such graph equally likely to be generated. So an upper bound on b_4 is given by

$$b_4 \le 2^{3((k-2)n/(k-1)+\mu n)} \mathbb{P}\left(|\overline{N}(\{y_1, y_2, y_3\}) \cap B_{low}^i| \le \frac{n}{200k^2}\right).$$

Since G' was chosen such that $|B_{low}^i| \ge n/2k^2$, we have that $\mathbb{E}(|\overline{N}(\{y_1, y_2, y_3\}) \cap B_{low}^i|) \ge n/16k^2$. So Lemma 2.1(ii) implies that

$$\mathbb{P}\left(|\overline{N}(\{y_1, y_2, y_3\}) \cap B_{low}^i| \le \frac{n}{200k^2}\right) \le \exp\left(-\frac{n}{128k^2}\right) \le 2^{-\frac{n}{128k^2}}.$$

Hence $b_4 \le 2^{3((k-2)n/(k-1)+\mu n)-n/128k^2}$, as required.

Now (9.3) together with Claims 1-4 and Proposition 4.3(ii) implies that

$$\begin{split} |\mathcal{A}_1| &\leq 2^3 k^2 n^3 \cdot C 2^{2(\log n)^3} f_k(n_k) 2^{t_{k-1}(n-3)} \cdot 2^{3\xi(\beta)n} \cdot 2^{3((k-2)n/(k-1)+\mu n)-n/128k^2} \\ &\leq C 2^{-n/150k^2} f_k(n_k) 2^{t_{k-1}(n-3)+3(k-2)n/(k-1)-3(k-2)-3} \\ &\leq C 2^{-n/150k^2} f_k(n_k) 2^{t_{k-1}(n)}, \end{split}$$

as required.

Lemma 9.4. $|\mathcal{A}_2| \leq C 2^{-n^{1/2k^2}/3} f_k(n_k) 2^{t_{k-1}(n)}$.

Proof. Note that for every $G \in \mathcal{A}_2$ and every $s \in \{0, 1, \dots, k-2\}$ the definition of $C^s(G, Q)$ implies that $Q_s \setminus C^s(G, Q) \ge |Q_s|/2$. So any graph $G \in \mathcal{A}_2$ can be constructed as follows. We first choose $a \in \mathbb{N}$ such that $n/2k^2 \le a \le n$, and then perform the following steps.

• We choose distinct indices $i, j \in \{0, 1, \dots, k-2\}$, a set $W = \{y_1, y_2\} \cup \{w_\ell^s : \ell \in [2], s \in \{0, 1, \dots, k-2\} \setminus \{j\}\}$ of vertices satisfying $y_1, y_2 \in Q_j$ and $w_1^s, w_2^s \in Q_s$ for every $s \in \{0, 1, \dots, k-2\} \setminus \{j\}$, a vertex $y_3 \in Q_j \setminus W$, and a set E of edges between the vertices in $W \cup \{y_3\}$. Let b_1 denote the number of such choices. The choices in this step and the next steps will be made such that $y_1, y_2 \notin C^j(G, Q)$ and $w_1^s, w_2^s \notin C^s(G, Q)$ for every $s \in \{0, 1, \dots, k-2\} \setminus \{j\}$, and $|B_{low}^i| = a$ and $C(Y) \cap \overline{N}(\{y_1, y_2\}) = \emptyset$, where $Y := \overline{N}(\{y_1, y_2, y_3\}) \cap B_{low}^i(G, Q)$.

- Next we choose the graph G' on vertex set $[n]\backslash W$ such that $G[[n]\backslash W] = G'$. Let b_2 denote the number of possibilities for G'.
- Next we choose the set E' of internal edges in G with exactly one endpoint in W such that E' is compatible with our previous choices. Let b_3 denote the number of possibilities for E'.
- Next we choose the set E'' of crossing edges in G between W and $B_{low}^i \setminus W$ such that E'' is compatible with our previous choices. Let b_4 denote the number of possibilities for E''.
- Finally we choose the set E''' of crossing edges in G between W and $[n]\setminus (W \cup B^i_{low})$ such that E''' is compatible with our previous choices. Let b_5 denote the number of possibilities for E'''.

Hence

(9.4)
$$|\mathcal{A}_2| \le n \max_{n/2k^2 \le a \le n} \{b_1 \cdot b_2 \cdot b_3 \cdot b_4 \cdot b_5\}.$$

The main idea of the proof is that since Y is large for $G \in \mathcal{A}_2$, it follows that C(Y) is also large. So the assumption that every element of C(Y) has at least one neighbour in $\{y_1, y_2\}$ places a significant restriction on the number of choices for G. The role of the w_ℓ^s is to 'balance out' the vertex classes, i.e. in the proof of Claim 5 it will be useful that W contains two vertices from each vertex class.

The following series of claims will give upper bounds for the quantities b_1, \ldots, b_5 . Claims 1 and 4 are trivial, and the proof of Claim 2 proceeds in an almost identical way to that of Claim 2 in the proof of Lemma 9.3; we give proofs of Claims 3 and 5.

Claim 1: $b_1 \leq k^2 n^{2k-1} 2^{\binom{2k-1}{2}}$.

Claim 2: $b_2 \le C2^{2(\log n)^3} f_k(n_k) 2^{t_{k-1}(n-(2k-2))}$.

Claim 3: $b_3 \le n^{4(k-1)}$.

Indeed, for every graph $\tilde{G} \in \mathcal{A}_2$ such that $y_1, y_2 \notin C^j(\tilde{G}, Q)$ and $w_1^s, w_2^s \notin C^s(\tilde{G}, Q)$ for every $s \in \{0, 1, \dots, k-2\} \setminus \{j\}$, Proposition 9.1(i) implies that $\overline{d}_{\tilde{G},Q}^j(y_\ell), \overline{d}_{\tilde{G},Q}^s(w_\ell^s) \leq 2$ for every $\ell \in [2]$ and every $s \in \{0, 1, \dots, k-2\} \setminus \{j\}$. Thus

$$b_3 \le n^{2|W|} \le n^{4(k-1)},$$

as required.

Claim 4: $b_4 \le 2^{(2k-4)a}$.

Claim 5: $b_5 \le 2^{(2k-4)(n-a)} 2^{-2n^{1/2k^2}/5}$.

Indeed, suppose G satisfies $C(Y) \cap \overline{N}(\{y_1, y_2\}) = \emptyset$. Since we choose G such that $|B_{low}^i| = a \ge n/2k^2$, the fact that $G \notin A_1$ implies that $|Y| > n/200k^2$. Now the definitions of C_{low}^i , B_{low}^i imply that

$$|C(Y)| \ge \frac{200k^2|Y|}{n^{1-1/2k^2}} \ge n^{1/2k^2}.$$

So since in G every vertex in C(Y) must have at least one neighbour in $\{y_1, y_2\}$,

$$(9.5) b_5 \leq 2^{2\sum_{s\in\{0,1,\dots,k-2\}\setminus\{j\}}|[n]\setminus(Q_s\cup B_{low}^i)|}2^{2|[n]\setminus(Q_j\cup B_{low}^i\cup C(Y))|}3^{|C(Y)|}$$
$$\leq 2^{(2k-4)(n-a)}2^{-2n^{1/2}k^2/5},$$

as required. The second inequality of (9.5) is where it is important that W contains two vertices from each vertex class.

Now (9.4) together with Claims 1–5 and Proposition 4.3(ii) implies that

$$\begin{split} |\mathcal{A}_2| & \leq n \cdot k^2 n^{2k-1} 2^{\binom{2k-1}{2}} \cdot C 2^{2(\log n)^3} f_k(n_k) 2^{t_{k-1}(n-(2k-2))} \\ & \cdot n^{4(k-1)} \cdot \max_{n/2k^2 \leq a \leq n} \left\{ 2^{(2k-4)a} \cdot 2^{(2k-4)(n-a)} 2^{-2n^{1/2k^2}/5} \right\} \\ & \leq C 2^{-n^{1/2k^2}/3} f_k(n_k) \cdot 2^{t_{k-1}(n-(2k-2)) + (2k-2)(k-2)n/(k-1) - (2k-2)(k-2) - t_{k-1}(2k-2)} \\ & \leq C 2^{-n^{1/2k^2}/3} f_k(n_k) 2^{t_{k-1}(n)}, \end{split}$$

as required.

As mentioned earlier, a (6,3)-forest (with edge set E say) is a useful building block for constructing many induced copies of C_{2k} . More precisely, in Lemma 9.5 we will show that there are many 'E-compatible' linear forests H, which play a similar role to that of the skeletons in the proof of Lemma 7.3. Each such $E \cup E(H)$ gives us a non-trivial restriction on the remaining edge set, resulting in an adequate bound on $|\mathcal{A}_3|$.

Lemma 9.5.
$$|\mathcal{A}_3| \leq C 2^{-\frac{n}{2^{14k}}} f_k(n_k) 2^{t_{k-1}(n)}$$
.

Proof. Any graph $G \in \mathcal{A}_3$ can be constructed as follows.

- First we choose distinct indices $i, j \in \{0, 1, ..., k-2\}$, a set $X \subseteq Q_i \cup Q_j$ of six vertices, and a set E of edges between vertices in X such that the graph (X, E) is a linear forest with at most three components (so E will be the edge set of a (6,3)-forest in G). Let b_1 denote the number of such choices.
- Next we choose a graph G' on vertex set $[n]\backslash X$ such that $G[[n]\backslash X] = G'$. Let b_2 denote the number of possibilities for G'.
- Next we choose the set E' of internal edges in G with exactly one endpoint in X. Let b_3 denote the number of possibilities for E'.
- Finally we choose the set E'' of crossing edges in G between X and $[n]\backslash X$ such that E'' is compatible with our previous choices. Let b_4 denote the number of possibilities for E''.

Hence,

$$(9.6) |\mathcal{A}_3| \le b_1 \cdot b_2 \cdot b_3 \cdot b_4.$$

The following series of claims will give upper bounds for the quantities b_1, \ldots, b_4 . Claim 1 is trivial, and the proofs of Claims 2 and 3 follow in an almost identical way to those of Claims 2 and 3 in the proof of Lemma 9.3, so we give only a proof of Claim 4.

Claim 1: $b_1 < 2^{15}k^2n^6$.

Claim 2: $b_2 \le C2^{2(\log n)^3} f_k(n_k) 2^{t_{k-1}(n-6)}$.

Claim 3: $b_3 \le 2^{6\xi(\beta)n}$.

Claim 4:
$$b_4 \le 2^{\frac{6(k-2)n}{k-1}} 2^{\mu^{1/4}n} 2^{-\frac{n}{2^{13k}}}$$
.

Indeed, we define an E-compatible forest to be a linear forest H on 2k-6 vertices, with the same number of components as (X, E), such that $V(H) \cap Q_s$ induces a clique on two vertices for every $s \in \{0, 1, \ldots, k-2\} \setminus \{i, j\}$. Note that an E-compatible forest exists since $2k-6 \geq 2 \cdot 3$ and (X, E) has at most three components. Moreover, an E-compatible forest contains a perfect matching, so Proposition 5.4 implies that for every graph $\tilde{G} \in \mathcal{A}_3$, the number of disjoint E-compatible forests in \tilde{G} is at least

$$\left| \frac{n/(k-1) - \mu n - 2\mu^{1/2}n}{2} \right| \ge \frac{n}{2(k-1)} - 3\mu^{1/2}n.$$

Hence G' contains at least $n/2(k-1)-3\mu^{1/2}n$ disjoint E-compatible forests. Now fix a set CF of $n/2(k-1)-3\mu^{1/2}n$ disjoint E-compatible forests in G', and let $H\in CF$. Let h_H denote the number of possibilities for a set E^* of edges between X and V(H). By Proposition 7.1 there exists at least one set \tilde{E} of edges between X and V(H) such that the graph $(X\cup V(H), E\cup E(H)\cup \tilde{E})$ is isomorphic to C_{2k} . So since G must be induced- C_{2k} -free, we must have that $E^*\neq \tilde{E}$, and hence $h_H\leq 2^{|X||V(H)|}-1=2^{12(k-3)}-1$. Note that the number of vertices outside $Q_i\cup Q_j$ that are not contained in some graph $H\in CF$ is at most $(k-3)n/(k-1)+2\mu n-(2k-6)(n/2(k-1)-3\mu^{1/2}n)\leq 6k\mu^{1/2}n$. Hence,

$$b_{4} \leq 2^{6 \cdot \max\{|Q_{i}|,|Q_{j}|\}} 2^{6(6k\mu^{1/2}n)} \prod_{H \in CF} h_{H}$$

$$\leq 2^{6(n/(k-1)+\mu n)} 2^{6(6k\mu^{1/2}n)} \left(2^{12(k-3)} \left(1 - 2^{-12(k-3)}\right)\right)^{n/(2(k-1)) - 3\mu^{1/2}n}$$

$$\leq 2^{\frac{6(k-2)n}{k-1}} 2^{40k\mu^{1/2}n} e^{-\frac{n/(2(k-1))}{2^{12(k-3)}}} \leq 2^{\frac{6(k-2)n}{k-1}} 2^{\mu^{1/4}n} 2^{-\frac{n}{2^{13k}}},$$

as required.

Now (9.6) together with Claims 1–4 and Proposition 4.3(ii) implies that

$$\begin{aligned} |\mathcal{A}_{3}| &\leq 2^{15} k^{2} n^{6} \cdot C 2^{2(\log n)^{3}} f_{k}(n_{k}) 2^{t_{k-1}(n-6)} \cdot 2^{6\xi(\beta)n} \cdot 2^{\frac{6(k-2)n}{k-1}} 2^{\mu^{1/4}n} 2^{-\frac{n}{2^{13k}}} \\ &\leq C 2^{-\frac{n}{2^{14k}}} f_{k}(n_{k}) 2^{t_{k-1}(n-6)+6(k-2)n/(k-1)-6(k-2)-t_{k-1}(6)} \\ &\leq C 2^{-\frac{n}{2^{14k}}} f_{k}(n_{k}) 2^{t_{k-1}(n)}, \end{aligned}$$

as required.

The next proposition shows that for every $G \in \mathcal{A}_4$, the small stars and triangles in $\overline{G}[Q_0]$ do not cover too many vertices.

Proposition 9.6. For every $G \in \mathcal{A}_4$ and index $i \in \{0, 1, \dots, k-2\}$, $|B_{low}^i| < n/2k^2$.

Proof. Suppose for a contradiction that there exists a graph $G \in \mathcal{A}_4$ such that $|B^i_{low}| \geq n/2k^2$ for some index $i \in \{0, 1, \ldots, k-2\}$. Since $G \in \mathcal{A}_4 \subseteq F_Q^3$, G is not a k-template. This fact together with Proposition 9.1(i) implies that there exists an index $j \in \{0, 1, \ldots, k-2\} \setminus \{i\}$ and a non-edge y_1y_3 inside Q_j . At most one of y_1, y_3 can be in C^j (by definition of C^j), and so without loss of generality we assume that $y_1 \notin C^j$. So Proposition 9.1(i) implies that $\overline{d}_{G,Q}^j(y_1) \leq 2$. This together with the observation that $|Q_j \setminus C^j| \geq |Q_j|/2$ (by definition of C^j) implies that there exists a vertex $y_2 \in Q^j \setminus C^j$ that is a neighbour of y_1 .

Define $Y := \overline{N}(\{y_1, y_2, y_3\}) \cap B^i_{low}$. Since $|B^i_{low}| \ge n/2k^2$ and $G \notin \mathcal{A}_1$, $|Y| > n/200k^2$. Since $|B^i_{low}| \ge n/2k^2$ and $G \notin \mathcal{A}_2$, C(Y) contains a vertex $x_3 \in \overline{N}(\{y_1, y_2\})$. Since $x_3 \in C(Y)$ there exists a vertex $x_1 \in Y$ that is a non-neighbour of x_3 . By Proposition 9.1(iii), $\overline{d}^i_{G,Q}(x_1)$, $\overline{d}^i_{G,Q}(x_3) \le \beta n$. So since $|Y| > n/200k^2 \ge 2\beta n$, there exists a vertex $x_2 \in Y \cap N(\{x_1, x_3\})$.

Then $E(G[\{x_1, x_2, x_3, y_1, y_2, y_3\}]) = \{x_1x_2, x_2x_3, y_1y_2\} \cup E' \text{ with } E' \subseteq \{y_2y_3, y_3x_3\}$. Thus the set $\{x_1, x_2, x_3, y_1, y_2, y_3\} \subseteq Q_i \cup Q_j$ induces on G a linear forest with at most three components, and so G contains a (6,3)-forest with respect to G. This contradicts the fact that $G \notin A_3$, and hence completes the proof.

We now have sufficient information about the set \mathcal{A}_4 of remaining graphs to count them directly (i.e. \mathcal{A}_4 is the only class for which we do not use induction in our estimates). In particular, we now know that in \overline{G} every vertex class is the union of triangles and stars, where crucially the number of triangles and small stars is not too large (see Proposition 9.6). This allows us to show by a direct counting argument that $|\mathcal{A}_4|$ is negligible.

Lemma 9.7. $|\mathcal{A}_4| \leq 2^{-\frac{n \log n}{3k^2}} f_k(n_k) 2^{t_{k-1}(n)}$.

Proof. Any graph $G \in \mathcal{A}_4$ can be constructed as follows.

- First we choose a partition of Q_i into five sets, $C_h^i, B_h^i, C_\ell^i, B_\ell^i, C_z^i$, for every $i \in \{0, 1, \dots, k-1\}$ 2}. Let b_1 denote the number of such choices.
- Next we choose the set E of crossing edges in G with respect to Q. Let b_2 denote the number of possibilities for E.
- \bullet Finally we choose the set E' of internal edges in G with respect to Q such that G satisfies $C_h^i = C_{high}^i, B_h^i = B_{high}^i, C_\ell^i = C_{low}^i, B_\ell^i = B_{low}^i, \text{ and } C_z^i = C_0^i \text{ for every } i \in \{0, 1, \dots, k-2\}.$ Let b_3 denote the number of possibilities for E'.

Hence

$$(9.7) |\mathcal{A}_4| \le b_1 \cdot b_2 \cdot b_3.$$

The following series of claims will give upper bounds for the quantities b_1, b_2, b_3 . Claims 1 and 2 are trivial; we give only a proof of Claim 3.

Claim 1: $b_1 \leq 5^n$.

Claim 2: $b_2 \leq 2^{t_{k-1}(n)}$.

Claim 3:
$$b_3 \leq 2^{\frac{(k-1/2)n\log n}{k^2}}$$
.

Claim 3: $b_3 \leq 2^{\frac{(k-1/2)n\log n}{k^2}}$. For any given $i \in \{0, 1, \dots, k-2\}$ and any vertex $x \in B^i_{high}$, the number of possibilities for the unique non-neighbour of x in C^i_{high} (namely the centre of the star in \overline{G} containing x) is $|C^i_{high}|$. Now consider $x \in B_{low}^i$. Then x has a unique non-neighbour y in C_{low}^i , and has the possibility of either being part of a triangle in \overline{G} or a star in \overline{G} . Note also that $|B_{low}^i| < n/2k^2$ by Proposition 9.6, and that by definition of C_{high}^i ,

$$|C_{high}^i| \le \frac{200k^2n}{n^{1-1/2k^2}} \le 200k^2n^{1/2k^2}.$$

Hence,

$$b_{3} \leq \prod_{i=0}^{k-2} (2|C_{low}^{i}|)^{|B_{low}^{i}|} |C_{high}^{i}|^{|B_{high}^{i}|} \leq \prod_{i=0}^{k-2} n^{\frac{n}{2k^{2}}} (200k^{2})^{n} (n^{\frac{1}{2k^{2}}})^{n} = 2^{\frac{(k-1)n\log n}{k^{2}}} (200k^{2})^{n(k-1)}$$
$$< 2^{\frac{(k-1/2)n\log n}{k^{2}}}.$$

as required.

Now (9.7) together with Claims 1–3 and Lemma 4.1 implies that

$$|\mathcal{A}_4| \le 5^n \cdot 2^{t_{k-1}(n)} \cdot 2^{\frac{(k-1/2)n\log n}{k^2}}$$

$$\le 5^n 2^{-\frac{n\log n}{2k^2}} 2^{n_k \log n_k - en_k \log \log n_k} 2^{en_k \log \log n_k} 2^{t_{k-1}(n)} \le 2^{-\frac{n\log n}{3k^2}} f_k(n_k) 2^{t_{k-1}(n)},$$

as required.

Recall that $F_Q^3 = (F_Q^3 \setminus T_Q^*(n,k)) \cup A_1 \cup A_2 \cup A_3 \cup A_4$. The following bound on $|F_Q^3|$ follows immediately from this observation together with (9.1) and Lemmas 9.3, 9.4, 9.5 and 9.7.

Lemma 9.8.
$$|F_Q^3| \le 2C2^{-n^{\frac{1}{2k^2}}/3} f(n_k) 2^{t_{k-1}(n)}$$
.

10. Proof of Lemma 6.1

Proof of Lemma 6.1. Recall from Section 6 that we prove Lemma 6.1 by induction on n and that we choose constants satisfying (6.1). The fact that $1/C \ll 1/n_0, 1/k$ implies that the statement of Lemma 6.1 holds for all $n \le n_0$. So suppose that $n > n_0$ and that the statement holds for all n' < n. Then we obtain the bounds in Lemmas 7.3, 8.7 and 9.8. These bounds together with the fact that $F_Q(n, k, \eta, \mu) = T_Q \cup F_Q^1 \cup F_Q^2 \cup F_Q^3$ and $T_Q \subseteq T_Q(n, k)$ imply that

$$|F_Q(n,k,\eta,\mu) \setminus T_Q(n,k)| \le C \left(2^{-\beta^2 n/14^k} + 2^{-n} + 2 \cdot 2^{-n^{\frac{1}{2k^2}}/3} \right) f_k(n_k) 2^{t_{k-1}(n)}$$

$$\le 3C 2^{-n^{\frac{1}{2k^2}}/3} f_k(n_k) 2^{t_{k-1}(n)}.$$

This together with Corollary 5.3 implies that

$$|F_{Q}(n,k,\eta) \setminus T_{Q}(n,k)| \leq |F_{Q}(n,k,\eta) \setminus F_{Q}(n,k,\eta,\mu)| + |F_{Q}(n,k,\eta,\mu) \setminus T_{Q}(n,k)|$$

$$\leq \left(2^{-\frac{\mu^{2}n^{2}}{100}} + 3C2^{-n^{\frac{1}{2k^{2}}/3}} f_{k}(n_{k})\right) 2^{t_{k-1}(n)}$$

$$\leq 4C2^{-n^{\frac{1}{2k^{2}}/3}} f_{k}(n_{k}) 2^{t_{k-1}(n)}.$$

Note that Lemma 3.1 (applied with $\eta/2$ playing the role of η) together with (5.1) implies that

$$(10.2) |F(n,k) \setminus F(n,k,\eta)| \le 2^{-\varepsilon n^2} |F(n,k,\eta)|.$$

Let \mathcal{Q} denote the set of all ordered (k-1)-partitions of [n], and recall that our choice of $Q \in \mathcal{Q}$ was arbitrary. Now (10.1) together with (10.2) and Lemma 4.4 implies that

$$|F(n,k) \setminus F(n,k,\eta)| \le 2^{-\varepsilon n^2} \sum_{Q' \in \mathcal{Q}} \left(|F_{Q'}(n,k,\eta) \setminus T_{Q'}(n,k)| + |T_{Q'}(n,k)| \right)$$

$$\le 2^{-\varepsilon n^2} (k-1)^n \left(4C2^{-n^{\frac{1}{2k^2}/3}} + 2^{6(\log n)^2} \right) f_k(n_k) 2^{t_{k-1}(n)}$$

$$\le C2^{-\varepsilon n^2/2} f_k(n_k) 2^{t_{k-1}(n)}.$$

Now this together with (10.1) implies that

$$|F_{Q}(n,k)| \leq |F_{Q}(n,k,\eta)| + |F(n,k) \setminus F(n,k,\eta)|$$

$$\leq |T_{Q}(n,k)| + |F_{Q}(n,k,\eta) \setminus T_{Q}(n,k)| + |F(n,k) \setminus F(n,k,\eta)|$$

$$\leq |T_{Q}(n,k)| + \left(4 \cdot 2^{-n\frac{1}{2k^{2}}/3} + 2^{-\epsilon n^{2}/2}\right) C f_{k}(n_{k}) 2^{t_{k-1}(n)}$$

$$\leq |T_{Q}(n,k)| + 5C 2^{-n\frac{1}{2k^{2}}/3} f_{k}(n_{k}) 2^{t_{k-1}(n)},$$

which completes the inductive step, and hence the proof.

11. Acknowledgement

We are grateful to Mihyun Kang for helpful remarks on the number of k-templates.

References

- [1] V.E. Alekseev, Range of values of entropy of hereditary classes of graphs, Diskret. Mat. 1992, 148–157.
- [2] N. Alon, J. Balogh, B. Bollobás, R. Morris, The structure of almost all graphs in a hereditary property, J. Combin. Theory B 101 (2011), 85–110.
- [3] N. Alon, A. Shapira, A characterisation of the (natural) graph properties testable with one-sided error, in Proc. of FOCS 2005, 429-438, SIAM J. Comput. (Special Issue on FOCS 05) 37 (2008), 1703-1727.
- [4] J. Balogh, J. Butterfield, Excluding induced subgraphs: Critical graphs, Random Structures and Algorithms 38 (2011), 100–120.
- [5] J. Balogh, R. Morris, W. Samotij, Independent sets in hypergraphs, J. Amer. Math. Soc 28 (2015), 669-709.
- [6] J. Balogh, R. Morris, W. Samotij, L. Warnke, The typical structure of sparse K_{r+1} -free graphs, Trans. Amer. Math. Soc, to appear.
- [7] B. Bollobás, Hereditary and monotone properties of combinatorial structures, in A. Hilton, J. Talbot, Surveys in Combinatorics 2007 (Cambridge University Press, Cambridge, 2007), 1–40.
- [8] B. Bollobás, A. Thomason, Hereditary and monotone properties of graphs, *The mathematics of Paul Erdős II* 14 (1997), 70–78.
- [9] N.G. de Bruijn, Asymptotic methods in analysis, (Dover, New York, NY, 1958).
- [10] P. Erdős, On some new inequalities concerning extremal properties of graphs, in P. Erdős and G. Katona, eds., Theory of Graphs (Academic Press, New York, 1968), 77–81.
- [11] P. Erdős, Some recent results on extremal problems in graph theory, in P. Rosenstiehl, ed., *Theory of Graphs* (Gordon and Breach, New York; and Dunod, Paris, 1976), 117–123.
- [12] P. Erdős, D.J. Kleitman, B.L. Rothschild, Asymptotic enumeration of K_n-free graphs, in Colloquio Internazionale sulle Teorie Combinatorie (Rome, 1973), Vol. II, 19–27. Atti dei Convegni Lincei 17, Accad. Naz. Lincei, Rome, 1976.
- [13] R.J. Kang, C. McDiarmid, B. Reed, A. Scott, For most graphs H, most H-free graphs have a linear homogeneous set, Random Structures and Algorithms 45 (2014) 343–361.
- [14] Ph.G. Kolaitis, H.J. Prömel, B.L. Rothschild, $K_{\ell+1}$ -free graphs: asymptotic structure and a 0-1 law, *Trans. Amer. Math. Soc.* **303** (1987), 637–671.
- [15] D. Kühn, D. Osthus, T. Townsend, Y. Zhao, On the structure of oriented graphs and digraphs with forbidden tournaments or cycles, arXiv:1404.6178.
- [16] D. Osthus, H.J. Prömel, A. Taraz, For which densities are random triangle-free graphs almost surely bipartite?, *Combinatorica* **23** (2003), 105–150.
- [17] H.J. Prömel, A. Steger, Almost all Berge graphs are perfect, Combin. Prob. Comput. 1 (1992), 53–79.
- [18] H.J. Prömel, A. Steger, Excluding induced subgraphs: quadrilaterals, Random Structures and Algorithms 2 (1991), 55–71.
- [19] H.J. Prömel, A. Steger, Excluding induced subgraphs II: extremal graphs, Discrete Appl. Math. 44 (1993), 283–294.
- [20] H.J. Prömel, A. Steger, Excluding induced subgraphs III: a general asymptotic, Random Structures and Algorithms 3 (1992), 19–31.
- [21] D. Saxton, A. Thomason, Hypergraph containers, Inventiones mathematicae, to appear.
- [22] M. Simonovits, A method for solving extremal problems in graph theory, in P. Erdős and G. Katona, eds., Theory of Graphs (Academic Press, New York, 1968), 279–310.
- [23] T. Townsend, PhD Thesis, University of Birmingham, in preparation.

School of Mathematics

University of Birmingham

Edgbaston

Birmingham

B15 2TT

UK

E-mail addresses: {j.kim.3, d.kuhn, d.osthus, txt238}@bham.ac.uk