# Approximability of the Vertex Cover Problem in Power Law Graphs

Mikael Gast<sup>\*</sup> Mathias Hauptmann<sup>†</sup>

#### Abstract

In this paper we construct an approximation algorithm for the MINIMUM VERTEX COVER Problem (MIN-VC) with an expected approximation ratio of  $2 - \frac{\zeta(\beta) - 1 - \frac{1}{2\beta}}{2^{\beta}\zeta(\beta - 1)\zeta(\beta)}$  for random *Power Law Graphs* (PLG) in the  $(\alpha, \beta)$ -model of Aiello et. al.. We obtain this result by combining the Nemhauser and Trotter approach for MIN-VC with a new deterministic rounding procedure which achieves an approximation ratio of  $\frac{3}{2}$  on a subset of low degree vertices for which the expected contribution to the cost of the associated linear program is sufficiently large.

## 1 Introduction

In recent years topological analyses have been applied to a variety of real world graphs such as the World-Wide Web, the Internet, Collaboration and Social Networks, Protein Interaction Networks and other large-scale graphs of biological systems. Typical statistical parameters such as the *diameter*, robustness, clustering coefficient and degree distribution have been measured and compared to the expected values of these parameters in uniform random graph models such as the classical G(n, p)-Model due to Erdős and Rényi [ER60]. It turned out that the real world graphs are significantly different from the random models with respect to these statistical and topological properties. In subsequent studies the aim was to describe the properties of real world networks mathematically and to propose new models in order to meet these conditions.

As of 1999 Kumar et. al. [BKM<sup>+</sup>00, KRR<sup>+</sup>00], Kleinberg et. al. [KKR<sup>+</sup>99, KL01] and Faloutsos, Faloutsos and Faloutsos [FFF99, SFFF03] measured the degree sequence of the World-Wide Web and independently observed that it is well approximated by a power law distribution, i.e. the number of nodes  $y_i$  of a given degree i is proportional to  $i^{-\beta}$  where  $\beta > 0$ . This was later verified for a large number of existing real-world networks such as protein-protein interactions, gene regulatory networks, peer-to-peer networks, mobile call networks and social networks [JAB01, GBBK02, SMS<sup>+</sup>08, EKM<sup>+</sup>04].

In order to analyze these graphs, some research has been directed towards finding suitable models for describing structural properties quantitatively and qualitatively. A number of *Power Law Graph* (PLG) models have been proposed, such as the Barabási-Albert model of *Preferential Attachment* [BA99], the *Buckley-Osthus Model* [BO04], the *Cooper-Frieze Model* [CF03] and the *Copying Model* due to Kumar et. al. [KRR<sup>+</sup>00]. All these models describe a *random growth process* starting from a small seed graph and yielding – besides other features – a power law degree sequences in the limit.

A different approach is to take a power law degree sequence as input and to generate a graph instance with this distribution in a random fashion. Among the most widely known models of

<sup>\*</sup>Dept. of Computer Science, University of Bonn. e-mail: gast@cs.uni-bonn.de

<sup>&</sup>lt;sup>†</sup>Dept. of Computer Science, University of Bonn. e-mail: hauptman@cs.uni-bonn.de

this kind is the ACL-Model due to Aiello, Chung and Lu [ACL01]. Here, the number  $y_i$  of vertices with degree *i* is roughly given by  $y_i \approx e^{\alpha}/i^{\beta}$ , where  $e^{\alpha}$  is a normalization constant which determines the size of the graph. While this model is potentially less accurate than the detailed description of a growth process, it has the advantage of being robust and general, i.e., structural properties that are true in this model will be true for the majority of graphs with the given degree sequence.

All of the above models are well motivated and there exists a large body of literature on mathematical foundations and applications [BA99, ACL00, BR02, EKM<sup>+</sup>04, MPS06]. In this paper, we focus on the ACL-Model for random PLG which we will refer to as the  $(\alpha, \beta)$ -Model.

Apart from having certain structural properties, such as high clustering coefficient, smallworld characteristics and self similarity, there exists practical evidence that combinatorial optimization in PLG is easier than in general graphs [PL01, GMS03, EKM<sup>+</sup>04, KGS06]. Contrasting this Ferrante et. al. [FPP08] and Shen et. al. [SNT10] studied the approximation hardness of certain optimization problems in *combinatorial Power Law Graphs* and showed NP-hardness and APX-hardness of classical problems such as MINIMUM VERTEX COVER (MIN-VC), MAXI-MUM INDEPENDENT SET (MAX-IS) and MINIMUM DOMINATING SET (MIN-DS). In this paper we study the approximability of the MINIMUM VERTEX COVER problem in the random Power Law Graph model of Aiello et. al. [ACL01].

The MINIMUM VERTEX COVER is one of the most well-studied problems in combinatorial optimization. A vertex cover of a graph G = (V, E) is a set of vertices  $C \subseteq V$  such that each edge  $e = \{u, v\}$  of G has at least one endpoint in C. The MINIMUM VERTEX COVER problem (MIN-VC) is the the problem of finding a cover of minimum cardinality in a graph. The problem is known to be NP-complete due to Karp's original proof [Kar72] and APX-complete [PY91]. Moreover, it cannot be approximated within a factor of 1.3606 [DS05], unless P = NP, and is inapproximable within  $2 - \epsilon$  for any  $\epsilon > 0$  as long as the Unique Games Conjecture (UGC) holds true [KR08]. Here, we show that the MIN-VC problem can be approximated with an expected approximation ratio < 2 in random Power Law Graphs:

**Theorem 1.** There exists a polynomial time algorithm which approximates the MINIMUM VER-TEX COVER problem (MIN-VC) in random Power Law Graphs in the  $(\alpha, \beta)$ -Model for  $\beta > 2$ (where graphs are given instance by instance) with an expected approximation ratio of

$$\rho = 2 - \frac{\zeta(\beta) - 1 - \frac{1}{2^{\beta}}}{2^{\beta}\zeta(\beta - 1)\zeta(\beta)}.$$

We also give a refined analysis for the case  $\beta > 2.424$  and obtain the following improvement.

**Theorem 2.** For  $\beta > 2.424$ , the MINIMUM VERTEX COVER problem (MIN-VC) in the  $(\alpha, \beta)$ -Model can be approximated with expected asymptotic approximation ratio

$$\rho' = 2 - \frac{\left(\zeta(\beta) - 1 - \frac{1}{2^{\beta}}\right) \cdot \zeta(\beta - 1)}{\zeta(\beta - 1) \cdot \zeta(\beta)} \left[ 1 - \left(\frac{\zeta(\beta - 1) - \left(1 + \frac{1}{2^{\beta - 1}}\right)}{\zeta(\beta - 1)}\right)^3 \right].$$

In Figure 1 these two upper bounds  $\rho$  and  $\rho'$  are shown as functions of the parameter  $\beta$ .

The paper is organized as follows. In subsection 2.1 we describe the  $(\alpha, \beta)$ -model for Power Law Graphs, describe the random generation process and give a formal description of the model parameters. In subsection 2.2 we give some background on the MIN-VC problem and briefly describe the *half-integral solution* method proposed by Nemhauser and Trotter. Section 3 presents our new approximation algorithm for MIN-VC in Power Law Graphs. This algorithm basically consists of a deterministic rounding procedure on a half-integral solution for MIN-VC.




Figure 1: Comparison of first (---) and second (—) analysis in terms of functions of the parameter  $\beta$ , for  $\beta > 2$  and  $\beta > 2.424$ , respectively.

In Section 3.1 we show that this rounding procedure yields an approximation ratio of  $\frac{3}{2}$  in the subgraph induced by the low-degree vertices of the Power Law Graph and a 2-approximation in the residual graph. In Section 3.2 we construct upper and lower bounds on the expected size of the half-integral solution in the induced subgraph of low-degree vertices and finally prove our main theorems. We conclude the paper by giving a short summary and further research in Section 4.

## 2 Preliminaries


### **2.1** $(\alpha, \beta)$ -Power Law Graphs

In this section we describe the random PLG-model proposed by Aiello, Chung and Lu in [ACL01], which we will denote as  $M(\alpha, \beta)$ . This model considers a random graph with the following degree distribution depending on two given values  $\alpha$  and  $\beta$ : For each  $1 \leq i \leq \Delta = \left\lfloor e^{\frac{\alpha}{\beta}} \right\rfloor$  there are  $y_i$  vertices of degree i with

$$y_i = \begin{cases} \left\lfloor \frac{e^{\alpha}}{i^{\beta}} \right\rfloor & \text{if } i > 1 \text{ or } \sum_{i=1}^{\Delta} \left\lfloor \frac{e^{\alpha}}{i^{\beta}} \right\rfloor \text{ is even} \\ \left\lfloor e^{\alpha} \right\rfloor + 1 & \text{otherwise.} \end{cases}$$

Here, *i* and  $y_i$  satisfy  $\log y_i = \alpha - \beta \log i$ . The variable  $\alpha$  is the logarithm of the size of the graph and  $\beta$  is the log-log growth rate. Let  $\mathcal{G}_{(\alpha,\beta)}$  be the set of all undirected graphs with multi-edges and self-loops on  $n = \sum_{i=1}^{\Delta} y_i$  vertices which have  $y_i$  vertices of degree i  $(1 \leq i \leq \Delta)$ . Then  $M(\alpha, \beta)$  is the distribution on  $\mathcal{G}_{(\alpha,\beta)}$  obtained in the following way [ACL01]:

- 1. Generate a set L of d(v) distinct copies of each vertex v.
- 2. Generate a random matching on the elements of L.
- 3. For each pair of vertices u and v, the number of edges joining u and v in G is equal to the number of edges in the matching of L which join copies of u to copies of v.



As in [ACL01], in the following we will work with the real numbers  $\frac{e^{\alpha}}{i^{\beta}}$ ,  $e^{\frac{\alpha}{\beta}}$  instead of their integer counterparts. For  $\beta > 2$  the error is a lower order term (c.f. [ACL01], remark on page 6).

A graph  $G \in \mathcal{G}_{(\alpha,\beta)}$  has the following properties: The maximum degree of G is  $e^{\frac{\alpha}{\beta}}$ , and for  $\beta > 2$  the number of vertices is  $n = \sum_{i=1}^{e^{\alpha/\beta}} \frac{e^{\alpha}}{i^{\beta}} \approx \zeta(\beta)e^{\alpha}$  and the number of edges is  $m = \frac{1}{2}\sum_{i=1}^{e^{\alpha/\beta}} i \frac{e^{\alpha}}{i^{\beta}} \approx \frac{1}{2}\zeta(\beta-1)e^{\alpha}$  where the error terms are o(n) and o(m), respectively.

## 2.2 LP-Relaxation and Half-Integral Solution for Min-VC

In this section we give a brief outline of the Nemhauser-Trotter Theorem stated in [NT75] and show how this is used to approximate MIN-VC in a graph G = (V, E) as described by Hochbaum et. al. in [HMNT93].

Nemhauser and Trotter considered the following LP-relaxation, which applies to the more general *weighted* vertex cover problem:

 $\begin{array}{ll} \text{minimize} & \sum_{i=1}^{n} w_{i} x_{i},\\ \text{subject to} & x_{i} + x_{j} \geq 1, \quad \text{for each edge } \{v_{i}, v_{j}\} \in E,\\ & x_{i} \quad \geqslant 0, \quad \text{for each vertex } v_{i} \in V, \end{array}$ 

They show that there always exists an optimal solution x for this LP which is *half-integral*, i.e. for all  $i, x_i \in \{0, \frac{1}{2}, 1\}$ . Then they partition the set of vertices into subsets  $P, Q, R \subseteq V$ , such that  $v_i \in P$  if  $x_i = 1, v_i \in Q$  if  $x_i = \frac{1}{2}$  and  $v_i \in R$  if  $x_i = 0$  in this solution. They show that at least one optimal vertex cover in G contains the set P, that each vertex in R has all its neighbors in P and – moreover – that each cover in G has weight at least  $w(P) + \frac{1}{2}w(Q)$ . From this it follows that at least one optimal vertex cover in G consists of the set P and an optimal cover in the subgraph H[Q] induced by Q.

Hochbaum et. al. [HMNT93] showed that an integer solution y obtained by setting  $y_i = 1$  for all vertices  $v_i \in Q \cup P$  and  $y_i = 0$  for all  $v_i \in R$  is a 2-approximate solution for the MIN-VC problem in G. Our approximation algorithm for MIN-VC in random Power Law Graphs will make use of a half-integral solution x of the LP-relaxation along with the properties described in the Nemhauser-Trotter Theorem in order to achieve an approximation ratio strictly less than 2.

# **3** Approximation of Min-VC in $(\alpha, \beta)$ -PLG

In this section we present our main result, namely an approximation algorithm with expected approximation ratio  $2 - \frac{\zeta(\beta) - 1 - \frac{1}{2\beta}}{2^{\beta}\zeta(\beta-1)\zeta(\beta)}$  for the MIN-VC problem in  $(\alpha, \beta)$ -PLG for  $\beta > 2$ . Furthermore a refined analysis yields an improved *asymptotic* approximation ratio for the case  $\beta > 2.424$ .

Let us first give an outline of this algorithm. On instance  $G \in \mathcal{G}_{(\alpha,\beta)}$  the algorithm starts with a half-integral solution  $x: V \to \{0, \frac{1}{2}, 1\}$  of the associated LP and uses some deterministic rounding procedure to generate an integral solution  $y: V \to \{0, 1\}$ . We show that for the set  $V^* = \bigcup_{v:d(v)\in\{1,2\}} (\{v\} \cup \mathcal{N}(v))$  of degree-1 and degree-2 nodes and their neighbors in G, the rounding procedure satisfies  $y(V^*) \leq \frac{3}{2} \cdot x(V^*)$  and furthermore  $x(V^*)$  is sufficiently large (in expectation) with respect to  $M(\alpha, \beta)$ .

### 3.1 Approximation Algorithm

Now, we describe our deterministic rounding procedure (Algorithm 1) on G = (V, E) for  $G \in \mathcal{G}_{(\alpha,\beta)}$ . First, the algorithm processes all nodes of the subset  $V' = L \cup \mathcal{N}(L)$  where  $L = \left\{ v \in V | \left( d(v) = 2, x(v) = \frac{1}{2} \right) \lor (d(v) = 1) \right\}$  and provides a rounded integral solution y with  $y(V') \leq \frac{3}{2} \cdot x(V')$ . Furthermore we show that  $y(V^* \setminus V') \leq \frac{4}{3} \cdot x(V^* \setminus V')$  and  $y(V \setminus V^*) \leq 2 \cdot x(V \setminus V^*)$ .

An analysis of the algorithm is provided by the following Lemma 3 and Lemma 4.

**Lemma 3.** The assignment y generated by Algorithm 1 is an integer solution and satisfies y(u) = 1 for all  $u \in V'$  with  $d(u) \ge 3$ .

*Proof.* Any high-degree neighbor of degree-1 vertices is set to 1 in step (1) of the algorithm.

Since either step (3) or (4) is processing every single degree-2 vertex  $v \in V$  with  $x(v) = \frac{1}{2}$ , there are no leftover vertices  $v \in V'$  of degree 2 with fractional values.

Assume that there is a vertex  $u \in V'$ ,  $d(u) \ge 3$  and  $x(u) = y(u) = \frac{1}{2}$ . Then u has at least one degree 2 neighbor  $v_1$  with  $x(v_1) = \frac{1}{2}$ . Because of step (3) and (4) of the algorithm,  $v_1$  must have been processed by another degree 2 vertex  $v_2$ , setting  $y(v_1) = 1$ . This again introduces another neighbor w of  $v_2$  with y(w) = 1 and leads to the situation of a path  $uv_1v_2w$  described in step (2). In this case, the algorithm sets y(u) = 1 and thus we have a contradiction to the above assumption.

**Lemma 4.** The assignment y generated by Algorithm 1 satisfies  $y(V^*) \leq \frac{3}{2} \cdot x(V^*)$ .

*Proof.* The algorithm partitions the graph induced by  $V^*$  into edge-disjoint subgraphs, namely stars whose leaves are degree-1 vertices and paths of length  $\leq 4$  whose internal nodes are degree-2 vertices. We show that for each such subgraph  $P_i$ ,  $y(P_i) \leq \frac{3}{2} \cdot x(P_i)$  and furthermore y(v) = 1 for each  $v \in V^*$  which is contained in more than one such subgraph.

In step (1) of the algorithm all degree-1 vertices and their neighbors are processed.

In step (2) the subgraphs are unprocessed paths  $P_i$  of length 3. Since  $P_i = \underbrace{\bullet \quad v_1 \quad v_2 \quad w}_{v_1 \quad v_2 \quad w}$  contains two disjoint edges  $\{u, v_1\}, \{v_2, w\}, x(P_i) \ge 2$  and particularly  $x(v_2) + x(w) \ge 1$ . Therefore  $y(P_i) = 3 \le \underbrace{\frac{3}{2}} \cdot x(P_i)$  holds via mapping  $\overset{\geq 1/2}{\bullet} \underbrace{\overset{1/2}{\bullet}} \overset{\geq 1/2}{\bullet} \overset{\geq 0}{\bullet} \mapsto \underbrace{\overset{1}{\bullet}} \underbrace{\overset{1}{\bullet}} \overset{0}{\bullet} \overset{0}{\bullet}$ 

In step (3) all paths  $P_i = \underbrace{\bullet}_{u} \underbrace{\bullet}_{v} \underbrace{\bullet}_{w}$  are processed, where at least one of u, w is of degree  $\geq 3$ . In cases (3.1)-(3.4) the algorithm considers all possible combinations of some of these nodes being already processed.

In case (3.1) u is marked unprocessed, w is already processed and  $x(u) \ge \frac{1}{2}$ . The rounding algorithm sets y(v) = 0 and y(u) = 1, mapping  $\overset{\ge 1/2}{\bullet} \overset{1/2}{\bullet} \overset{1}{\bullet} \mapsto \overset{1}{\bullet} \overset{0}{\bullet} \overset{1}{\bullet} \overset{0}{\bullet} \overset{1}{\bullet}$ , again yielding a vertex cover  $y \upharpoonright P_i$  for  $P_i$  with  $y(P_i) \le x(P_i)$ .

In case (3.2) we have that both u, w are marked as unprocessed and since  $x(v) = \frac{1}{2}$  we have that  $x(u) \ge \frac{1}{2}$  and  $x(w) \ge \frac{1}{2}$ . The rounding algorithm sets y(v) = 0, y(u) = y(w) = 1, mapping  $\stackrel{\geq 1/2}{\bullet} \stackrel{1/2}{\bullet} \stackrel{1/2}{\bullet} \stackrel{0}{\bullet} \stackrel{1}{\bullet} \stackrel{0}{\bullet} \stackrel{1}{\bullet} \stackrel{1}{\bullet} \stackrel{1}{\bullet} \stackrel{0}{\bullet} \stackrel{1}{\bullet} \stackrel{1}{\bullet} \stackrel{1}{\bullet} \stackrel{0}{\bullet} \stackrel{1}{\bullet} \stackrel{1}{\bullet} \stackrel{1}{\bullet} \stackrel{1}{\bullet} \stackrel{0}{\bullet} \stackrel{1}{\bullet} \stackrel{1$ 

Algorithm 1: DETERMINISTIC ROUNDING

**Input**:  $G = (V, E), x : V \to \{0, \frac{1}{2}, 1\}.$ **Output**:  $y: V \to \{0, 1\}$ . forall the  $v \in V$  do y(v) := x(v);mark v as unprocessed; (0) compute G' = (V', E') induced by  $V' = L \cup \mathcal{N}(L)$  where  $L = \left\{ v \in V | \left( d(v) = 2, x(v) = \frac{1}{2} \right) \lor (d(v) = 1) \right\};$ (1) forall the  $v \in V$  with d(v) = 1 do let u be the neighbor of v in G; set y(v) = 0; set y(u) = 1; (2) forall the  $P = uv_1v_2w \subset G'$  unprocessed,  $d(u) \ge 3, d(v_1) = d(v_2) = 2$  do set  $y(u) = y(w) = y(v_1) = 1;$ set  $y(v_2) = 0;$ (3) forall the  $v \in V'$  unprocessed,  $d(v) = 2 \land \exists u \in \mathcal{N}(v), d(u) \ge 3$  do else if u unprocessed, w processed then (3.1)set y(v) = 0; set y(u) = 1; else if  $both \ u, w$  unprocessed then (3.2)set y(v) = 0; set y(u) = y(w) = 1; /\* with  $x(u) \ge \frac{1}{2}$  and  $x(w) \ge \frac{1}{2}$  \*/ if both u, w processed then (3.3) $| \quad \texttt{set } y(v) = 0;$ else if u processed, w unprocessed then (3.4)set y(v) = 0; set y(w) = 1; /\* y(u) = 1 already set and  $x(w) \ge \frac{1}{2}$  \*/ (4) forall the  $v \in V'$  unprocessed, d(v) = 2 do else if u unprocessed, w processed then (4.1)| set y(v) = 0; set y(u) = 1; else if both u, w unprocessed then (4.2)/\* with  $x(u) \ge \frac{1}{2}$  and  $x(w) \ge \frac{1}{2}$  \*/ set y(v) = 0; set y(u) = y(w) = 1;  ${\bf if} \ both \ u,w \ {\bf processed \ then} \\$ (4.3)set y(v) = 0;else if u processed, w unprocessed then (4.4)set y(v) = 0; set y(w) = 1; /\* y(u) = 1 already set and  $x(w) \ge \frac{1}{2}$  \*/ (5) for all the  $v \in V$  do if  $x(v) = \frac{1}{2}$  then  $/* y(v) = \min\{1, 2 \cdot x(v)\} */$ set y(v) = 1;

In case (3.3) both u, w are marked as processed and therefore y(u) = y(w) = 1, since u, w are adjacent to processed degree one or degree two vertices other than v. The algorithm sets y(v) = 0, mapping  $\begin{bmatrix} 1 & 1/2 & 1 \\ 0 & 0 & 0 \end{bmatrix} \mapsto \begin{bmatrix} 1 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}^1$ . Hence  $y \upharpoonright P_i$  is a vertex cover for  $P_i$  with  $y(P_i) \leq x(P_i)$ .

In case (3.4) u is already processed and w is still marked unprocessed. Since  $x(v) = \frac{1}{2}$  we have that  $x(w) \ge \frac{1}{2}$ . The rounding algorithm sets y(v) = 0 and y(w) = 1, mapping  $\begin{bmatrix} 1 & 1/2 \\ 0 & 0 \end{bmatrix} \xrightarrow{1} \begin{bmatrix} 1/2 \\ 0 & 0 \end{bmatrix} \xrightarrow{1} \begin{bmatrix} 0 \\ 0 & 0 \end{bmatrix} \xrightarrow{1} \begin{bmatrix} 1/2 \\ 0 & 0 \end{bmatrix} \xrightarrow{1} \begin{bmatrix} 0 \\ 0 & 0 \end{bmatrix} \xrightarrow{1} \begin{bmatrix} 1/2 \\ 0 & 0 \end{bmatrix} \xrightarrow{1} \begin{bmatrix} 0 \\ 0 & 0 \end{bmatrix} \xrightarrow$ 

Step (4) considers all remaining unprocessed vertices of degree 2. If v is such a vertex with neighborhood  $\mathcal{N}(v) = \{u, w\}$ , the sub-cases (4.1)-(4.4) are treated analogously to cases (3.1)-(3.4) and the mapping  $x \mapsto y$  achieves  $y(P_i) \leq \frac{4}{3} \cdot x(P_i)$  on the considered paths  $P_i$ .

After steps (0)-(4) of the algorithm there may still be some remaining high-degree vertices  $u \in V^*, d(u) \ge 3$  with  $x(u) = y(u) = \frac{1}{2}$ . These are treated separately (and rounded to y(u) = 1 together with all other vertices in  $V \setminus (V' \setminus V^*)$ ) in step (5) of the algorithm. We have to argue that  $y(V^*) \le \frac{3}{2} \cdot x(V^*)$  still holds true.

We consider first the case that  $u \in V', d(u) \ge 3$  and  $x(u) = y(u) = \frac{1}{2}$ . Then u has a neighbor v of degree  $\le 2$  with  $x(v) = \frac{1}{2}$  and y(v) = 1, and since  $y(u) = \frac{1}{2}$  we have d(v) = 2. Let  $v_2$  be the other neighbor of v, then  $d(v_2) = 1$  (since otherwise the second neighbor w of  $v_2$  would give rise to a path of length 3, containing also u and hence would have been processed in step (2)). But then locally on the set  $\{u, v, v_2\}$  we have the mapping  $\int_{-\infty}^{1/2} \int_{-\infty}^{1/2} \cdots \int_{-\infty}^{1/2} \int_{-\infty}^{0} \cdots \int_{-\infty}^$ 

Let us now assume  $u \in V^* \setminus V', d(u) \ge 3$  and  $x(u) = y(u) = \frac{1}{2}$ . Then every degree-2 neighbor v has  $x(v) \ne \frac{1}{2}$ , hence x(v) = 1, and therefore y(v) = 1. We show that  $v \notin V'$ , i.e. that v was not processed by the algorithm and can be treated as a part of a subgraph disjoint to G' in G. Let  $w \in \mathcal{N}(v)$  be the second neighbor of v besides u. Then x(w) = 0 since otherwise (in case  $x(w) \ge \frac{1}{2}$ ) we could decrease x(v) from 1 to  $\frac{1}{2}$  and still have a feasible half-integral solution, which would contradict the optimality of x. Therefore  $v, w \notin V'$ , which means that v, w are not processed by the algorithm. Rounding y(u) = 1, mapping  $\overset{1/2}{\bullet} \xrightarrow{1}{\bullet} \mapsto \overset{1}{\bullet} \xrightarrow{1}{\bullet}$ , yields a vertex cover  $y \upharpoonright \{u, v\}$  with  $y(\{u, v\}) \le \frac{4}{3} \cdot x(\{u, v\})$ .

We conclude that the assignment  $y: V \mapsto \{0, 1\}$  is a vertex cover of G with  $y(V^*) \leq \frac{3}{2} \cdot x(V^*)$ and  $y(V \setminus V^*) \leq 2 \cdot x(V \setminus V^*)$ .

## 3.2 Expected Approximation Ratio

The following lemma shows how to retrieve an expected approximation ratio for our algorithm for MIN-VC in G.

**Lemma 5.** If the rounding scheme  $x \mapsto y$  satisfies  $y(V^*) \leq \frac{3}{2} \cdot x(V^*)$  and  $y(V \setminus V^*) \leq 2 \cdot x(V \setminus V^*)$ then this gives an approximation ratio

$$\frac{y(V)}{\mathsf{OPT}} \leqslant \frac{y(V)}{x(V)} \leqslant \frac{x(V^*)}{x(V)} \cdot \frac{3}{2} + \frac{x(V \setminus V^*)}{x(V)} \cdot 2.$$

In order to apply Lemma 5 and to derive an expected approximation ratio for the algorithm, in the following we will give a lower bound on  $\mathbb{E}[x(V^*)]$  and an upper bound on x(V). The next lemma provides a lower bound on  $x(V^*)$  in terms of the number of high-degree vertices adjacent to degree-1 and degree-2 nodes. **Lemma 6.** Let  $G[V^*]$  be the subgraph of G induced by  $V^*$ . For every optimal half-integral solution x for the MIN-VC LP, the size of the half-integral solution restricted to  $V^*$  is lower-bounded by the size of the high-degree neighborhood of degree-1 and degree-2 vertices:

$$x(V^*) \ge \frac{1}{2} \cdot \left| \left\{ u \in V | d(u) \ge 3 \land \exists v \in \mathcal{N}(u), d(v) \in \{1, 2\} \right\} \right|$$

Proof. Let  $V^* = X \cup Y, X = \{v \in V | d(v) \in \{1,2\}\}$  and  $Y = \{u \in V | d(u) \ge 3 \land \exists v \in \mathcal{N}(u), d(v) \in \{1,2\}\}$ . Choose some arbitrary function  $f: Y \to E(X,Y)$  such that for every  $u \in Y, f(u) = \{u, v\}$  for some  $v \in X$  adjacent to u. f(Y) consists of pairwise disjoint paths  $Q_1, \ldots, Q_m$  of length  $\leq 2$ , such that each path contains one or two vertices from Y. This implies  $x(V^*) \ge m \ge \frac{|Y|}{2}$ .

#### **First Analysis**

We will now estimate the expected number of high-degree vertices adjacent to vertices of degree one or two, which – combined with the preceding Lemma 6 – gives a lower bound on  $\mathbb{E}[x(V^*)]$ . We prove the following theorem:

#### Theorem 7.

$$\mathbb{E}[x(V^*)] \ge \frac{1}{2} \cdot \mathbb{E}\left[\left|\left\{u \in V | d(u) \ge 3 \land \exists v \in \mathcal{N}(u), d(v) \in \{1, 2\}\right\}\right|\right]$$

$$= \frac{1}{2} \cdot \sum_{u:d(u)\ge 3} \eta(u) \tag{1}$$

$$e^{\alpha} \quad \zeta(\beta) - 1 - \frac{1}{2^{\beta}}$$

$$\geq \frac{1}{2^{\beta}} \cdot \frac{1}{\zeta(\beta-1)}$$
<sup>(2)</sup>

where  $\eta(u)$  is the probability that  $u \in V$  has a neighbor in the set of vertices of degree one or two.

In order to provide bounds on the probability  $\eta(u)$  for a vertex u of degree d of having a degree-1 or degree-2 neighbor, we consider how edges are generated in the random matching procedure of the distribution  $M(\alpha, \beta)$ : d(u) copies of u are randomly matched with the copies of the remaining vertices  $v \in V, v \neq u$ . We use the following lower bound on  $\eta(u)$ .

**Lemma 8.** For every u with  $d(u) \ge 3$ ,  $\eta(u) \ge \frac{1}{2^{\beta-1} \cdot \sum_{i=1}^{\Delta} \frac{1}{i^{\beta-1}}}$ .

Proof.

 $\eta(u) \ge \Pr(\text{the first copy of } u \text{ is neighbor of a degree-2-node})$ 

$$= \frac{2 \cdot \# \text{deg-2-nodes}}{(\sum_{v \in V} d(v)) - 1}$$
  
$$\geqslant \frac{2 \cdot \frac{e^{\alpha}}{2^{\beta}}}{\sum_{i=1}^{\Delta} i \cdot \frac{e^{\alpha}}{i^{\beta}}} = \frac{\frac{1}{2^{\beta-1}}}{\sum_{i=1}^{\Delta} \frac{1}{i^{\beta-1}}}, \text{ where } \Delta = e^{\frac{\alpha}{\beta}} \text{ is the maximum degree of } G.$$

In Equation 1 we substitute  $\eta(u)$  by the bound given in Lemma 8 and obtain:

$$\mathbb{E}[x(V^*)] \ge \frac{1}{2} \cdot \sum_{u:d(u)\ge 3} \eta(u)$$

$$= \frac{1}{2} \cdot \left(\sum_{i=1}^{\Delta} \frac{e^{\alpha}}{i^{\beta}} - e^{\alpha} - \frac{e^{\alpha}}{2^{\beta}}\right) \cdot \frac{1}{2^{\beta-1} \cdot \sum_{i=1}^{\Delta} \frac{1}{i^{\beta-1}}}$$

$$= \frac{e^{\alpha}}{2^{\beta}} \cdot \frac{\sum_{i=1}^{\Delta} \frac{1}{i^{\beta}-1} - \frac{1}{2^{\beta}}}{\sum_{i=1}^{\Delta} \frac{1}{i^{\beta-1}}}$$
(3)

We will now show that in Inequality 3 we can replace the terms  $\sum_{i=1}^{\Delta} \frac{1}{i^{\beta}}$  and  $\sum_{i=1}^{\Delta} \frac{1}{i^{\beta-1}}$  by  $\zeta(\beta)$  and  $\zeta(\beta-1)$ , respectively. We make use of the following lemma.

**Lemma 9.** For A, B, a, b > 0,  $\frac{A}{B} \ge \frac{A+a}{B+b} \iff \frac{A}{B} \ge \frac{a}{b}$ .

Therefore, in order to show

$$\mathbb{E}[x(V^*)] \ge \frac{e^{\alpha}}{2^{\beta}} \cdot \frac{\zeta(\beta) - 1 - \frac{1}{2^{\beta}}}{\zeta(\beta - 1)},$$

it is sufficient to show that there exists a  $\Delta_0$  such that for all  $\Delta \ge \Delta_0$  the following holds

$$\frac{\sum_{i=1}^{\Delta} \frac{1}{i^{\beta}} - 1 - \frac{1}{2^{\beta}}}{\sum_{i=1}^{\Delta} \frac{1}{i^{\beta-1}}} \ge \frac{\frac{1}{(\Delta+1)^{\beta}}}{\frac{1}{(\Delta+1)^{\beta-1}}} = \frac{1}{\Delta+1}$$

This is provided by the following lemma.

**Lemma 10.** There exists a  $\Delta_0 \ge 8$  such that for all  $\Delta \ge \Delta_0$ ,  $\frac{\sum_{i=1}^{\Delta} \frac{1}{i^{\beta}} - 1 - \frac{1}{2^{\beta}}}{\sum_{i=1}^{\Delta} \frac{1}{i^{\beta-1}}} \ge \frac{1}{\Delta+1}$ .

*Proof.* The above inequality is equivalent to

$$\sum_{i=1}^{\Delta} \frac{1}{i^{\beta}} - 1 - \frac{1}{2^{\beta}} \ge \sum_{i=1}^{\Delta} \frac{1}{\Delta + 1} \cdot \frac{1}{i^{\beta - 1}}$$

$$\iff \qquad \sum_{i=1}^{\Delta} \left( \frac{1}{i^{\beta}} - \frac{1}{\Delta + 1} \cdot \frac{1}{i^{\beta - 1}} \right) \ge 1 + \frac{1}{2^{\beta}}$$

$$\iff \qquad \sum_{i=1}^{\Delta} \frac{\Delta + 1 - i}{(\Delta + 1)i^{\beta}} \ge 1 + \frac{1}{2^{\beta}}$$
(4)

Suppose  $\Delta \ge 8$ , then the sum on the left-hand side of the Inequality 4 is bounded by the sum of the terms with indices i = 1, 2, 4, 8:

$$\sum_{i=1}^{\Delta} \frac{\Delta+1-i}{(\Delta+1)i^{\beta}} \ge \frac{\Delta}{\Delta+1} + \frac{\Delta-1}{(\Delta+1)2^{\beta}} + \frac{\Delta-3}{(\Delta+1)4^{\beta}} + \frac{\Delta-7}{(\Delta+1)8^{\beta}} = \frac{\Delta 8^{\beta} + (\Delta-1)4^{\beta} + (\Delta-3)2^{\beta} + \Delta-7}{(\Delta+1)8^{\beta}}.$$
 (5)

Using Inequality 5 and the fact that  $1 + \frac{1}{2^{\beta}} = \frac{(\Delta+1)8^{\beta} + (\Delta+1)4^{\beta}}{(\Delta+1)8^{\beta}}$ , in order to prove Inequality 4 it is sufficient to show the following:

$$\frac{\Delta 8^{\beta} + (\Delta - 1)4^{\beta} + (\Delta - 3)2^{\beta} + \Delta - 7}{(\Delta + 1)8^{\beta}} \stackrel{!}{\geqslant} \frac{(\Delta + 1)8^{\beta} + (\Delta + 1)4^{\beta}}{(\Delta + 1)8^{\beta}}$$
$$(\Delta - 3)2^{\beta} + \Delta - 7 \stackrel{!}{\geqslant} 8^{\beta} + 2 \cdot 4^{\beta}.$$

This is valid for  $\Delta \ge \frac{8^{\beta} + 2 \cdot 4^{\beta} + 6 \cdot 2^{\beta} + 7}{1 + 2^{\beta}}$ . Hence we choose  $\Delta_0 = \lceil * \rceil \frac{8^{\beta} + 2 \cdot 4^{\beta} + 6 \cdot 2^{\beta} + 7}{1 + 2^{\beta}}$ .

This completes the proof of Theorem 7. The next lemma provides an upper bound for x(V): Lemma 11.  $x(V) \leq \frac{1}{2}\zeta(\beta) e^{\alpha}$ 

*Proof.* In order to get an upper bound for x(V) we construct a feasible half-integral solution for G by setting  $x(v) = \frac{1}{2}$  for all  $v \in V$  where  $\frac{1}{2} \sum_{v \in V} \leq \frac{1}{2} \zeta(\beta) e^{\alpha}$ .

Now let us restate the main Theorem 1 and finish the proof.

**Theorem.** For  $\beta > 2$  the MINIMUM VERTEX COVER problem in  $(\alpha, \beta)$ -Power Law Graphs G can be approximated with expected approximation ratio  $\rho \leq 2 - \frac{\zeta(\beta) - 1 - \frac{1}{2\beta}}{2^{\beta}\zeta(\beta-1)\zeta(\beta)}$ .

*Proof.* Algorithm 1 achieves an approximation ratio of  $\frac{3}{2}$  for MIN-VC in the subgraph induced by  $V^*$  in G and a ratio of 2 in  $G[V \setminus V^*]$ , i.e.

$$\rho \leqslant \mathbb{E}\left[\frac{3}{2} \cdot \frac{x(V^*)}{x(V)} + 2 \cdot \frac{x(V) - x(V^*)}{x(V)}\right] = \mathbb{E}\left[2 - \frac{1}{2} \cdot \frac{x(V^*)}{x(V)}\right].$$

From Theorem 7 and Lemma 11 we have that  $\mathbb{E}[x(V^*)] \ge \frac{1}{2} \cdot \frac{\left(\zeta(\beta) - 1 - \frac{1}{2^{\beta}}\right)e^{\alpha}}{2^{\beta-1}\zeta(\beta-1)}$  and  $x(V) \le \frac{1}{2} \cdot \zeta(\beta) e^{\alpha}$ . This yields

$$\mathbb{E}\left[\frac{x(V^*)}{x(V)}\right] \geqslant \frac{\frac{1}{2} \cdot \frac{\left(\zeta(\beta) - 1 - \frac{1}{2^{\beta}}\right)e^{\alpha}}{2^{\beta - 1}\zeta(\beta - 1)}}{\frac{1}{2} \cdot \zeta(\beta)e^{\alpha}} = \frac{\zeta(\beta) - 1 - \frac{1}{2^{\beta}}}{2^{\beta - 1}\zeta(\beta - 1)\zeta(\beta)}$$

and

 $\Leftrightarrow$ 

$$\rho \leqslant 2 - \frac{1}{2} \cdot \frac{\zeta(\beta) - 1 - \frac{1}{2^{\beta}}}{2^{\beta - 1}\zeta(\beta - 1)\zeta(\beta)} = 2 - \frac{\zeta(\beta) - 1 - \frac{1}{2^{\beta}}}{2^{\beta}\zeta(\beta - 1)\zeta(\beta)}$$

#### **Refined Analysis for** $\beta > 2.424$

We will now refine the analysis of Algorithm 1 by giving a better estimate on the probability  $\eta(u, U)$  of a high-degree node u being adjacent to a vertex in the set U, i.e. a vertex of degree one or two. However, this analysis will only apply to the more restricted range of  $\beta > 2.424$ . Again, we will first obtain a bound on the expected approximation ratio of the algorithm in terms of the partial sums  $\sum_{i=1}^{\Delta} \frac{1}{i^{\beta}}$  and  $\sum_{i=1}^{\Delta} \frac{1}{i^{\beta}-1}$  and then show that these can be replaced by  $\zeta(\beta)$  and  $\zeta(\beta-1)$ , respectively.

**Lemma 12.** For every u with  $d(u) \ge 3$  and  $U \subseteq V$ ,

$$\eta(u,U) \geqslant \frac{\sum_{i=1}^{\Delta} \frac{e^{\alpha}}{i^{\beta-1}} - e^{\frac{\alpha}{\beta}} + 1}{\sum_{i=1}^{\Delta} \frac{e^{\alpha}}{i^{\beta-1}}} \left[ 1 - \left( \frac{\sum_{i=1}^{\Delta} \frac{e^{\alpha}}{i^{\beta-1}} - d(U) - 3 + 1}{\sum_{i=1}^{\Delta} \frac{e^{\alpha}}{i^{\beta-1}} - 3 + 1} \right)^3 \right].$$

*Proof.* For a given set U of vertices from G we let  $d(U) = \sum_{v \in U} d(v)$ . Furthermore let  $\eta(u, U)$  be the probability that u is connected to at least one node in U. We obtain

$$\eta(u, U) = \Pr(u \text{ matches to } U)$$
  
=  $\sum_{j=1}^{d(u)} \Pr(j\text{-th copy is first one matching to } U)$   
=  $\sum_{j=1}^{d(u)} \frac{d(u)}{\sum_{i=1} \frac{e^{\alpha}}{i^{\beta-1}} - (j-1) - 1} \prod_{k=1}^{j-1} \left(1 - \frac{d(U)}{\sum_{i=1} \frac{e^{\alpha}}{i^{\beta-1}} - 1 - (k-1)}\right)$ 

Now define  $N = \sum_{i=1} \frac{e^{\alpha}}{i^{\beta-1}}$ . We have:

$$\begin{split} \eta(u,U) &= \sum_{j=1}^{d(u)} \frac{d(U)}{N-j} \prod_{k=1}^{j-1} \frac{N-d(U)-k}{N-k} \\ &\geqslant \sum_{j=1}^{d(u)} \frac{d(U)}{N-j} \left( \frac{N-d(U)-d(u)+1}{N-d(u)+1} \right)^{j-1} \\ &\geqslant \sum_{j=1}^{d(u)} \frac{d(U)}{N} \left( \frac{N-d(U)-d(u)+1}{N-d(u)+1} \right)^{j-1} \\ &= \frac{d(U)}{N} \left[ \frac{1-\left(\frac{N-d(U)-d(u)+1}{N-d(u)+1}\right)^{d(u)}}{1-\frac{N-d(U)-d(u)+1}{N-d(u)+1}} \right] \\ &= \frac{d(U)}{N} \left[ 1-\left(\frac{N-d(U)-d(u)+1}{N-d(u)+1}\right)^{d(u)} \right] \cdot \frac{N-d(u)+1}{d(U)} \\ &= \frac{N-d(u)+1}{N} \left[ 1-\left(\frac{N-d(U)-d(u)+1}{N-d(u)+1}\right)^{d(u)} \right] \end{split}$$

Since the function  $\left(\frac{N-d(U)-d(u)+1}{N-d(u)+1}\right)^{d(u)}$  is monotone decreasing in d(u) it follows that:

$$\begin{split} \eta(u,U) &\ge \frac{N-\Delta+1}{N} \left[ 1 - \left(\frac{N-d(U)-3+1}{N-3+1}\right)^3 \right] \\ &= \frac{\sum_{i=1} \frac{e^{\alpha}}{i^{\beta-1}} - e^{\frac{\alpha}{\beta}} + 1}{\sum_{i=1} \frac{e^{\alpha}}{i^{\beta-1}}} \left[ 1 - \left(\frac{\sum_{i=1} \frac{e^{\alpha}}{i^{\beta-1}} - d(U) - 3 + 1}{\sum_{i=1} \frac{e^{\alpha}}{i^{\beta-1}} - 3 + 1} \right)^3 \right] \end{split}$$

Because of Equation 1 we have  $\mathbb{E}[x(V^*)] \ge \frac{1}{2} \cdot \sum_{u:d(u)\ge 3} \eta(u,U)$  and we obtain the following

approximation ratio:

$$\begin{split} \rho \leqslant \mathbb{E} \left[ 2 - \frac{1}{2} \cdot \frac{x(V^*)}{x(V)} \right] \\ \leqslant 2 - \frac{1}{2} \cdot \frac{\left( \sum_{i=1}^{\Delta} \frac{e^{\alpha}}{i^{\beta}} - e^{\alpha} - \frac{e^{\alpha}}{2^{\beta}} \right) \cdot \frac{\sum_{i=1}^{\Delta} \frac{e^{\alpha}}{i^{\beta-1}} - e^{\frac{\alpha}{\beta}} + 1}{\sum_{i=1}^{\Delta} \frac{e^{\alpha}}{i^{\beta-1}}} \left[ 1 - \left( \frac{\sum_{i=1}^{\Delta} \frac{e^{\alpha}}{i^{\beta-1}} - d(U) - 3 + 1}{\sum_{i=1}^{\Delta} \frac{e^{\alpha}}{i^{\beta-1}} - 3 + 1} \right)^{3} \right]}{\frac{1}{2} \sum_{i=1}^{\Delta} \frac{e^{\alpha}}{i^{\beta}}}{\frac{1}{2} \sum_{i=1}^{\Delta} \frac{e^{\alpha}}{i^{\beta}}} \\ = 2 - \underbrace{\left( \sum_{i=1}^{\Delta} \frac{1}{i^{\beta}} - 1 - \frac{1}{2^{\beta}} \right) \cdot \left( \sum_{i=1}^{\Delta} \frac{1}{i^{\beta-1}} - \frac{\alpha}{e^{\alpha}} + \frac{1}{e^{\alpha}} \right)}{\int \frac{\left( \sum_{i=1}^{\Delta} \frac{1}{i^{\beta-1}} - \frac{1}{e^{\alpha}} \right)}{F}} \left[ 1 - \underbrace{\left( \frac{\sum_{i=1}^{\Delta} \frac{1}{i^{\beta-1}} - \frac{d(v)}{e^{\alpha}} - \frac{2}{e^{\alpha}}}{\sum_{i=1}^{\Delta} \frac{1}{i^{\beta-1}} - \frac{2}{e^{\alpha}}} \right)^{3}}_{C} \right] \end{split}$$

$$(6)$$

Now we show that, in Inequality 6, we can replace the partial sums  $\sum_{i=1}^{\Delta} \frac{1}{i^{\beta}}$  and  $\sum_{i=1}^{\Delta} \frac{1}{i^{\beta-1}}$  by  $\zeta(\beta)$  and  $\zeta(\beta-1)$  respectively. First, we consider the term C where  $d(v) = e^{\alpha} \left(1 + \frac{1}{2^{\beta-1}}\right)$ , i.e. the number of copies of degree-1 and degree-2 vertices:

$$C = \frac{\sum_{i=1}^{\Delta} \frac{1}{i^{\beta-1}} - \frac{e^{\alpha} \left(1 + \frac{1}{2^{\beta-1}}\right)}{e^{\alpha}} - \frac{2}{e^{\alpha}}}{\sum_{i=1}^{\Delta} \frac{1}{i^{\beta-1}} - \frac{2}{e^{\alpha}}} = \frac{\sum_{i=1}^{\Delta} \frac{1}{i^{\beta-1}} - \left(1 + \frac{1}{2^{\beta-1}}\right) - \frac{2}{\Delta^{\beta}}}{\sum_{i=1}^{\Delta} \frac{1}{i^{\beta-1}} - \frac{2}{\Delta^{\beta}}}$$

We show that following inequality holds true:

We have

$$F = \frac{\left(\sum_{i=1}^{\Delta} \frac{1}{i^{\beta}} - 1 - \frac{1}{2^{\beta}}\right) \cdot \left(\sum_{i=1}^{\Delta} \frac{1}{i^{\beta-1}} - \frac{\Delta}{e^{\alpha}} + \frac{1}{e^{\alpha}}\right)}{\left(\sum_{i=1}^{\Delta} \frac{1}{i^{\beta-1}}\right) \cdot \left(\sum_{i=1}^{\Delta} \frac{1}{i^{\beta}}\right)}.$$

We let  $S_{\beta} = \sum_{i=1}^{\Delta} \frac{1}{i^{\beta}}$  and  $S_{\beta-1} = \sum_{i=1}^{\Delta} \frac{1}{i^{\beta-1}}$  and recall that  $e^{\alpha} = \Delta^{\beta}$ . According to Lemma 9 it remains to show the following inequality:

$$\frac{\left(S_{\beta} - 1 - \frac{1}{2^{\beta}}\right) \cdot \left(S_{\beta-1} - \frac{1}{\Delta^{\beta-1}} + \frac{1}{\Delta^{\beta}}\right)}{\left(S_{\beta-1}\right) \cdot \left(S_{\beta}\right)} \\
\geqslant \frac{1}{\frac{1}{\left(\Delta+1\right)^{\beta}} \left(S_{\beta-1}\right) + \frac{1}{\left(\Delta+1\right)^{\beta-1}} \left(S_{\beta} + \frac{1}{\left(\Delta+1\right)^{\beta}}\right)} \cdot \left[\frac{1}{\left(\Delta+1\right)^{\beta}} \left(S_{\beta-1} - \frac{1}{\Delta^{\beta-1}} + \frac{1}{\Delta^{\beta}}\right) \\
+ \left(S_{\beta} - 1 - \frac{1}{2^{\beta}} + \frac{1}{\left(\Delta+1\right)^{\beta}}\right) \cdot \left(-\frac{1}{\left(\Delta+1\right)^{\beta}} + \frac{1}{\left(\Delta+1\right)^{\beta-1}} + \frac{1}{\left(\Delta\right)^{\beta-1}} - \frac{1}{\left(\Delta\right)^{\beta}}\right)\right]$$

which is equivalent to

$$\begin{split} \left(S_{\beta}-1-\frac{1}{2^{\beta}}\right)\left(S_{\beta-1}-\frac{\Delta-1}{\Delta^{\beta}}\right)\left[\frac{1}{(\Delta+1)^{\beta}}S_{\beta-1}+\left(S_{\beta}+\frac{1}{(\Delta+1)^{\beta}}\right)\frac{1}{(\Delta+1)^{\beta-1}}\right]\\ \geqslant \left[\frac{1}{(\Delta+1)^{\beta}}\left(S_{\beta-1}+\frac{\Delta-1}{\Delta^{\beta}}\right)\cdot S_{\beta}\cdot S_{\beta-1}\right.\\ \left.+\left(S_{\beta}-1-\frac{1}{2^{\beta}}+\frac{1}{(\Delta+1)^{\beta}}\right)\cdot \left(\frac{\Delta}{(\Delta+1)^{\beta}}+\frac{\Delta-1}{\Delta^{\beta}}\right)\cdot S_{\beta}\cdot S_{\beta-1}\right]. \end{split}$$

We rearrange terms and get

$$S_{\beta-1}^{2}S_{\beta}\frac{1}{(\Delta+1)^{\beta}} + S_{\beta}^{2}S_{\beta-1}\left(\frac{\Delta}{(\Delta+1)^{\beta}} + \frac{\Delta-1}{\Delta^{\beta}}\right) + S_{\beta}S_{\beta-1}\left[-\frac{\Delta-1}{\Delta^{\beta}} \cdot \frac{1}{(\Delta+1)^{\beta}} + \left(\frac{\Delta}{(\Delta+1)^{\beta}} + \frac{\Delta-1}{\Delta^{\beta}}\right)\left(-1 - \frac{1}{2^{\beta}} + \frac{1}{(\Delta+1)^{\beta}}\right)\right] \\ \leqslant S_{\beta-1}^{2}S_{\beta}\frac{1}{(\Delta+1)^{\beta}} + S_{\beta-1}S_{\beta}^{2}\frac{1}{(\Delta+1)^{\beta-1}} + S_{\beta}S_{\beta-1}\left[\frac{1}{(\Delta+1)^{\beta}} \cdot \frac{1}{(\Delta+1)^{\beta-1}} - \frac{\Delta-1}{\Delta^{\beta}} - \left(1 + \frac{1}{2^{\beta}}\right)\frac{1}{(\Delta+1)^{\beta-1}}\right] \\ + S_{\beta}^{2}\frac{1}{(\Delta+1)^{\beta-1}} + S_{\beta-1}^{2}\left(-1 - \frac{1}{2^{\beta}}\right)\frac{1}{(\Delta+1)^{\beta}} + S_{\beta-1}\left[-\left(1 + \frac{1}{2^{\beta}}\right)\frac{1}{(\Delta+1)^{\beta}} \cdot \frac{1}{(\Delta+1)^{\beta-1}}\left(1 + \frac{1}{2^{\beta}}\right)\frac{\Delta-1}{\Delta^{\beta}} \cdot \frac{1}{(\Delta+1)^{\beta}}\right] \\ + S_{\beta}\left[\left(-1 - \frac{1}{2^{\beta}}\right)\frac{1 - \Delta}{\Delta^{\beta}} \cdot \frac{1}{(\Delta+1)^{\beta-1}} - \frac{\Delta-1}{\Delta^{\beta}} \cdot \frac{1}{(\Delta+1)^{\beta}} \cdot \frac{1}{(\Delta+1)^{\beta-1}}\right] \\ + \left(1 + \frac{1}{2^{\beta}}\right)\frac{\Delta-1}{\Delta^{\beta}} \cdot \frac{1}{(\Delta+1)^{\beta-1}}$$
(7)

The following lemma shows that in order to prove Inequality 7 it is sufficient to show the respective inequality given by the terms of slowest convergence as  $\Delta$  goes to infinity.

**Lemma 13.** Let  $f_{\beta}, g_{\beta}, F_{\beta}, G_{\beta}$  be functions of  $\Delta$  depending on the parameter  $\beta$  with  $|g_{\beta}|, |G_{\beta}| \leq c$  for a constant c depending only on  $\beta$ . Then  $f_{\beta}(\Delta) < F_{\beta}(\Delta)$  for almost all  $\Delta$  implies

$$f_{\beta}(\Delta) \cdot \frac{1}{\Delta^{\beta-1}} + g_{\beta}(\Delta) \cdot \frac{1}{\Delta^{\beta}} \leqslant F_{\beta}(\Delta) \cdot \frac{1}{\Delta^{\beta-1}} + G_{\beta}(\Delta) \cdot \frac{1}{\Delta^{\beta}}$$

for all but finitely many  $\Delta$ .

Hence it remains to show that

$$\begin{split} S_{\beta}^2 S_{\beta-1} \left( \frac{\Delta}{(\Delta+1)^{\beta}} + \frac{\Delta-1}{\Delta^{\beta}} \right) - S_{\beta} S_{\beta-1} \left( 1 + \frac{1}{2^{\beta}} \right) \left( \frac{\Delta}{(\Delta+1)^{\beta}} + \frac{\Delta-1}{\Delta^{\beta}} \right) \\ \leqslant S_{\beta} S_{\beta-1} (-1) \cdot \frac{\Delta-1}{\Delta^{\beta}} + S_{\beta-1} S_{\beta}^2 \frac{1}{(\Delta+1)^{\beta-1}} \end{split}$$

which holds true if and only if

$$S_{\beta}^{2}S_{\beta-1}\left(\frac{\Delta}{(\Delta+1)^{\beta}} + \frac{\Delta-1}{\Delta^{\beta}}\right)$$
  
$$\leqslant S_{\beta}S_{\beta-1}\left[\left(1 + \frac{1}{2^{\beta}}\right)\left(\frac{\Delta}{(\Delta+1)^{\beta}} + \frac{\Delta-1}{\Delta^{\beta}}\right) - \frac{\Delta-1}{\Delta^{\beta}}\right] + S_{\beta-1}S_{\beta}^{2}\frac{1}{(\Delta+1)^{\beta-1}}$$

which can be rewritten as

$$S_{\beta}^{2}S_{\beta-1}\left(\frac{1}{(\Delta+1)^{\beta}} + \frac{\Delta-1}{\Delta^{\beta}}\right) \leqslant S_{\beta}S_{\beta-1}\left[\left(1 + \frac{1}{2^{\beta}}\right)\frac{\Delta}{(\Delta+1)^{\beta}} + \frac{1}{2^{\beta}} \cdot \frac{\Delta-1}{\Delta^{\beta}}\right]$$

$$\iff \qquad S_{\beta}\left(\frac{\Delta-1}{\Delta^{\beta}}\right) \leqslant \left(1 + \frac{1}{2^{\beta}}\right)\frac{\Delta}{(\Delta+1)^{\beta}} + \frac{1}{2^{\beta}} \cdot \frac{\Delta-1}{\Delta^{\beta}}$$

$$\iff \qquad S_{\beta}(\Delta-1) \leqslant \left(1 + \frac{1}{2^{\beta}}\right)\frac{\Delta^{\beta+1}}{(\Delta+1)^{\beta}} + \frac{1}{2^{\beta}} \cdot (\Delta-1)$$

$$\iff \qquad \left(S_{\beta} - \frac{1}{2^{\beta}}\right)(\Delta-1) \leqslant \left(1 + \frac{1}{2^{\beta}}\right)\frac{\Delta^{\beta+1}}{(\Delta+1)^{\beta}}$$
(8)

Now Inequality 8 follows from the observation that for all  $\beta > 2.424$ ,  $S_{\beta} - \frac{1}{2^{\beta}} < 1 + \frac{1}{2^{\beta}}$ . Finally we have shown the following theorem.

**Theorem.** For all  $\beta > 2.424$  the MINIMUM VERTEX COVER problem on  $(\alpha, \beta)$ -Power Law Graphs G can be approximated with expected approximation ratio

$$\rho \leqslant 2 - \frac{\left(\zeta(\beta) - 1 - \frac{1}{2^{\beta}}\right) \cdot \left(\zeta(\beta - 1) - \frac{\Delta}{e^{\alpha}} + \frac{1}{e^{\alpha}}\right)}{\zeta(\beta - 1) \cdot \zeta(\beta)} \left[1 - \left(\frac{\zeta(\beta - 1) - \left(1 + \frac{1}{2^{\beta - 1}}\right) - \frac{2}{e^{\alpha}}}{\zeta(\beta - 1) - \frac{2}{e^{\alpha}}}\right)^3\right]$$

This converges to

$$\rho \leqslant 2 - \frac{\left(\zeta(\beta) - 1 - \frac{1}{2^{\beta}}\right) \cdot \zeta(\beta - 1)}{\zeta(\beta - 1) \cdot \zeta(\beta)} \left[ 1 - \left(\frac{\zeta(\beta - 1) - \left(1 + \frac{1}{2^{\beta - 1}}\right)}{\zeta(\beta - 1)}\right)^3 \right]$$

as  $\alpha \to \infty$ .

## 4 Conclusion

In Section 3 we presented a new approximation algorithm for MIN-VC in  $(\alpha, \beta)$ -PLG with expected approximation ratio of  $\rho \leq 2 - \frac{\zeta(\beta) - 1 - \frac{1}{2\beta}}{2^{\beta}\zeta(\beta - 1)\zeta(\beta)}$  in our first analysis of Section 3.2. Moreover, in our refined analysis we showed for  $\beta > 2.424$  an expected asymptotic approximation ratio of  $\rho' \leq 2 - \frac{(\zeta(\beta) - 1 - \frac{1}{2\beta}) \cdot \zeta(\beta - 1)}{\zeta(\beta - 1) \cdot \zeta(\beta)} \left[ 1 - \left(\frac{\zeta(\beta - 1) - (1 + \frac{1}{2\beta - 1})}{\zeta(\beta - 1)}\right)^3 \right]$ . The algorithm itself basically consists of a deterministic rounding proceedure on a half integral solution for MIN VC (a.f. Algorithm 1)

a deterministic rounding procedure on a half-integral solution for MIN-VC (c.f. Algorithm 1). We showed that this rounding procedure yields an approximation ratio of  $\frac{3}{2}$  in the subgraph induced by the low-degree vertices of the  $(\alpha, \beta)$ -PLG and a 2-approximation in the residual graph.

Further research will be directed towards extending the improved analysis also to the range  $\beta < 2.424$  and towards investigating the approximability of MIN-VC in other PLG-Models, e.g. the Preferential Attachment Model in [BA99].

## Acknowledgements

The first author is supported by the NRW State within the B-IT Research School. The authors would like to thank Marek Karpinski for helpful remarks and discussions.

# References

- [ACL00] William Aiello, Fan Chung, and Linyuan Lu. A random graph model for massive graphs. In Proceedings of the thirty-second annual ACM Symposium on Theory of Computing (STOC), pages 171–180. ACM, 2000.
- [ACL01] William Aiello, Fan Chung, and Linyuan Lu. A random graph model for power law graphs. *Experimental Mathematics*, 10(1):53–66, 2001.
- [BA99] Albert-László Barabási and Réka Albert. Emergence of scaling in random networks. Science, 286(5439):509, 1999.
- [BKM<sup>+</sup>00] Andrei Broder, Ravi Kumar, Farzin Maghoul, Prabhakar Raghavan, Sridhar Rajagopalan, Raymie Stata, Andrew S. Tomkins, and Janet Wiener. Graph structure in the Web. Computer networks, 33(1-6):309–320, June 2000.
- [BO04] Pierce G. Buckley and Deryk Osthus. Popularity based random graph models leading to a scale-free degree sequence. *Discrete Mathematics*, 282(1-3):53–68, 2004.
- [BR02] Béla Bollobás and Oliver M. Riordan. Mathematical results on scale-free random graphs. Handbook of graphs and networks, pages 1–34, 2002.
- [CF03] Colin Cooper and Alan Frieze. A general model of web graphs. *Random Structures & Algorithms*, 22(3):311–335, 2003.
- [DS05] Irit Dinur and Shmuel Safra. On the hardness of approximating minimum vertex cover. Annals of Mathematics, 162(1):439–485, 2005.
- [EKM<sup>+</sup>04] Stephen Eubank, V. S. Anil Kumar, Madhav V. Marathe, Aravind Srinivasan, and Nan Wang. Structural and algorithmic aspects of massive social networks. In Proceedings of the 15th annual ACM-SIAM Symposium on Discrete Algorithms, pages 718–727. SIAM, 2004.
- [ER60] Paul Erdős and Alfréd Rényi. On the evolution of random graphs. Publications of the Mathematical Institute of the Hungarian Academy of Sciences, 5:17–61, 1960.
- [FFF99] Michalis Faloutsos, Petros Faloutsos, and Christos Faloutsos. On power-law relationships of the internet topology. ACM SIGCOMM Computer Communication Review, 29(4):251–262, 1999.
- [FPP08] Alessandro Ferrante, Gopal Pandurangan, and Kihong Park. On the hardness of optimization in power-law graphs. *Theoretical Computer Science*, 393(1-3):220–230, March 2008.
- [GBBK02] Nabil Guelzim, Samuele Bottani, Paul Bourgine, and François Képès. Topological and causal structure of the yeast transcriptional regulatory network. *Nature Genetics*, 31(1):60–63, 2002.
- [GMS03] Christos Gkantsidis, Milena Mihail, and Amin Saberi. Conductance and congestion in power law graphs. SIGMETRICS Performance Evaluation Review, 31:148–159, 2003.
- [HMNT93] Dorit S. Hochbaum, Nimrod Megiddo, Joseph Naor, and Arie Tamir. Tight bounds and 2-approximation algorithms for integer programs with two variables per inequality. *Mathematical Programming*, 62(1):69–83, 1993.

- [JAB01] Mihajlo Jovanović, Fred S. Annexstein, and Kenneth A. Berman. Modeling peer-topeer network topologies through "small-world" models and power laws. In IX Telecommunications Forum, TELFOR, 2001.
- [Kar72] Richard M. Karp. Reducibility among combinatorial problems. Complexity of Computer Computations, 40(4):85–103, 1972.
- [KGS06] Mehmet Koyutürk, Ananth Grama, and Wojciech Szpankowski. Assessing significance of connectivity and conservation in protein interaction networks. *Journal of Computational Biology*, 14(6):747–64, 2006.
- [KKR<sup>+</sup>99] Jon M. Kleinberg, Ravi Kumar, Prabhakar Raghavan, Sridhar Rajagopalan, and Andrew S. Tomkins. The Web as a graph: measurements, models and methods. Proceedings of the 5th annual international Conference on Computing and Combinatorics, 1627:1–17, 1999.
- [KL01] Jon M. Kleinberg and Steve Lawrence. The structure of the Web. *Science*, 294(5548):1849, 2001.
- [KR08] Subhash Khot and Oded Regev. Vertex cover might be hard to approximate to within 2-ε. Journal of Computer and System Sciences, 74(3):335–349, May 2008.
- [KRR<sup>+00]</sup> Ravi Kumar, Prabhakar Raghavan, Sridhar Rajagopalan, D. Sivakumar, Andrew S. Tomkins, and Eli Upfal. Stochastic models for the web graph. In *Proceedings of the 41st* annual Symposium on Foundations of Computer Science, pages 57–65. IEEE, 2000.
- [MPS06] Milena Mihail, Christos H. Papadimitriou, and Amin Saberi. On certain connectivity properties of the internet topology. Journal of Computer and System Sciences, 72(2):239–251, 2006.
- [NT75] George L. Nemhauser and Leslie E. Trotter. Vertex packings: Structural properties and algorithms. *Mathematical Programming*, 8(1):232–248, December 1975.
- [PL01] Kihong Park and Heejo Lee. On the effectiveness of route-based packet filtering for distributed DoS attack prevention in power-law internets. ACM SIGCOMM Computer Communication Review, 31(4):15–26, October 2001.
- [PY91] Christos H. Papadimitriou and Mihalis Yannakakis. Optimization, approximation, and complexity classes. Journal of Computer and System Sciences, 43(3):425–440, December 1991.
- [SFFF03] Georgos Siganos, Michalis Faloutsos, Petros Faloutsos, and Christos Faloutsos. Power laws and the AS-level internet topology. *IEEE/ACM Transactions on Networking*, 11(4):514–524, 2003.
- [SMS<sup>+</sup>08] Mukund Seshadri, Sridhar Machiraju, Ashwin Sridharan, Jean Bolot, Christos Faloutsos, and Jure Leskovec. Mobile call graphs: beyond power-law and lognormal distributions. In Proceeding of the 14th ACM SIGKDD international conference on Knowledge discovery and data mining (KDD), pages 596–604, New York, New York, USA, August 2008. ACM Press.
- [SNT10] Yilin Shen, Dung T. Nguyen, and My T. Thai. On the hardness and inapproximability of optimization problems on power law graphs. *Combinatorial Optimization and Applications*, pages 197–211, 2010.