# Integration and Cell Decomposition in P-minimal Structures

Pablo Cubides Kovacsics \*1 and Eva Leenknegt † 2

<sup>1</sup>Laboratoire Paul Painlevé, Université de Lille 1, CNRS U.M.R. 8524, 59655 Villeneuve d'Ascq Cedex, France.

<sup>2</sup>Department of Mathematics, KULeuven, Celestijnenlaan 200B, 3001 Heverlee, Belgium.

July 8, 2021

#### Abstract

We show that the class of  $\mathcal{L}$ -constructible functions is closed under integration for any P-minimal expansion of a p-adic field  $(K, \mathcal{L})$ . This generalizes results previously known for semi-algebraic and sub-analytic structures. As part of the proof, we obtain a weak version of cell decomposition and function preparation for P-minimal structures, a result which is independent of the existence of Skolem functions. A direct corollary is that Denef's results on the rationality of Poincaré series hold in any P-minimal expansion of a p-adic field  $(K, \mathcal{L})$ .

## 1 Introduction

One of the main results of this paper is that, for arbitrary P-minimal structures over p-adic fields, the class of constructible functions is closed under integration. This generalizes a result which was previously known only for semi-algebraic and sub-analytic structures (and some intermediary cases).

As part of the proof, we obtain the second main result of this paper: a version of cell decomposition and function preparation for P-minimal structures. While our version is somewhat weaker than what was obtained in previous attempts by e.g. Mourgues [20], it does not depend on the existence of definable Skolem functions.

<sup>\*</sup>The research leading to these results has received funding from the European Research Council, ERC Grant nr. 615722, MOTMELSUM, 2014 - 2019.

<sup>&</sup>lt;sup>†</sup>During the realization of this project, the second author was a postdoctoral fellow of the Fund for Scientific Research - Flanders (Belgium) (F.W.O.).

In this introduction we give an informal motivation of our approach, discussing the historical connections between p-adic integration, rationality of Poincaré series, and cell decomposition. We will present exact statements of our main results in the next section.

The study of p-adic integrals was motivated by a number-theoretic question. It was conjectured by Borevich and Shafarevich that for  $f(x) \in \mathbb{Z}[x]$ , Poincaré series like e.g.  $P(T) := \sum_{m \in \mathbb{N}} N_m T^m$ , where

$$N_m := \#\{x \in (\mathbb{Z}/p^m\mathbb{Z})^n \mid f(x) \equiv 0 \mod p^m\},\$$

are rational functions of T. This was originally proven by Igusa [15–17]. Later, Denef [8] obtained a similar, more general result. He gave two proofs that were based on Macintyre's quantifier elimination for semi-algebraic sets [19], one using resolution of singularities, and one where he introduced cell decomposition techniques. We refer to [10] for a comparison of both approaches.

A first step towards a proof is the realization that the terms of a Poincaré series can be connected to the measure of certain semi-algebraic sets, and hence to p-adic integrals. For instance, one can check that

$$N_m = p^{nm} \cdot \mu(\{x \in \mathbb{Z}_p^n \mid \operatorname{ord} f(x) \geqslant m\}),$$

where  $\mu$  is the Haar measure normalized such that  $\mu(\mathbb{Z}_p) = 1$ , and ord denotes the valuation map ord :  $\mathbb{Q}_p \to \mathbb{Z} \cup \{\infty\}$ . To prove rationality, one needs to understand how the measure of this family of definable sets depends on m. (If the dependence is tame enough, identities like  $\sum_{n=0}^{\infty} x^n = \frac{1}{1-x}$  can be used to deduce rationality.) Hence the focus shifts to the computation of (families of) p-adic integrals.

In a p-adic integral, the integrand is usually the valuation 'ord f' or the p-adic norm  $|f| := p^{-\text{ord}f}$  of a semi-algebraic function f. More generally, one can consider so-called constructible functions, i.e.  $\mathbb{Q}$ -linear combinations of definable functions  $\alpha: \mathbb{Q}_p^r \to \mathbb{Z}$  and their induced functions  $p^{\alpha}$ . Note that it is natural here to work in a two-sorted structure  $(\mathbb{Q}_p, \mathbb{Z})$ , adding the value group  $\mathbb{Z} \cup \{\infty\}$  as a separate sort. The word definable should also be interpreted in that context, using a two-sorted language  $\mathcal{L}_{\text{ring},2} = (\mathcal{L}_{\text{ring}}, \mathcal{L}_{\text{Pres}}, \text{ord})$ , consisting of the ring language  $\mathcal{L}_{\text{ring}}$  for the main sort  $\mathbb{Q}_p$ , the Presburger language  $\mathcal{L}_{\text{Pres}} = (+, -, <, \equiv_n)$  for the value group sort  $\mathbb{Z} \cup \{\infty\}$ , and the valuation map ord :  $\mathbb{Q}_p \to \mathbb{Z} \cup \{\infty\}$  to connect the sorts.

Denef showed in [9] that given a semi-algebraic function f, the integral of |f| equals a constructible function, a result which was later generalized to the sub-analytic setting by Cluckers, Gordon and Halupczok [6]. We will discuss this in more detail in the next section. In section 4, we prove that this closure property holds in arbitrary P-minimal structures over p-adic fields. Corollary 4.6 then provides an example of how the rationality of Poincaré series can be deduced from this.

Let us now discuss the connection with cell decomposition techniques. The

general philosophy is to partition a definable set X in somewhat uniform parts, called cells. If X is the domain of a function f, an additional goal may be to prepare the function, i.e. choosing the partition in such a way that the function f, when restricted to each of the cells, has some additional nice properties. Cells are generally taken to be sets of the form

$$\left\{ (x,t) \in D \times T \, \middle| \, \begin{array}{c} \text{a condition of a fixed form describing} \\ t \text{ in terms of the other variables } x \end{array} \right\} \,,$$

where D is a definable set and T is one of the sorts. For instance, when working with semi-algebraic sets, this fixed form is a formula stating that x belongs to an interval-like set:

ord 
$$a_1(x) \square_1$$
 ord  $(t - c(x)) \square_2$  ord  $a_2(x)$  and  $t - c(x) \in \lambda P_n$ ,

where  $\Box_i$  may denote < or no condition,  $\lambda \in K$  and the functions  $a_i(x)$  are definable functions  $D \to K$ . We use the symbol  $P_n$  for the set of (non-zero) n-th powers. Note that if  $\lambda = 0$ , what we get is just the graph of the function c(x).

What Denef [8,12] showed is that given a semi-algebraic function  $f: X \subseteq K^{n+1} \to K$ , X can be decomposed into cells such that on each cell C,  $f_{|C}$  satisfies the following condition:

$$|f(x,t)| = |\lambda(t-c(x))|^{\frac{e}{n}}|h(x)|,$$

where h(x) and c(x) are definable functions and e, n are integer numbers. The version stated here is a reformulation by Cluckers [5], who also obtained an analogous result for subanalytic functions in [1].

This preparation result is particularly useful for integration purposes. Indeed, if the domain of a function  $f: X \to K$  can be partitioned into a finite number of sets  $\{(x,t) \in D_i \times T \mid \phi_i(x,t)\}$  on each of which f has the form prescribed above, then one gets that

$$\int_{X} |f(x,t)| |dx| |dt| = \sum_{i} \int_{D_{i}} |h(x)| \left[ \int_{\{t \in T | \phi_{i}(x,t)\}} |\lambda(t - c(x))|^{\frac{e}{n}} |dt| \right] |dx|.$$

Iteration of this theorem allows one to give very accurate descriptions of the value of the integral of f and its dependence on possible parameters. Similar strategies were applied for the subanalytic case, see e.g. [1,11].

When Haskell and Macpherson developed the notion of P-minimality [13], it was natural to ask how much of the above ideas could be generalized to that setting. One of the most notable results so far in this context is Mourgues' cell decomposition theorem [20]. She showed that in a P-minimal structure which admits definable Skolem functions, any definable set  $A \subseteq K^{r+1}$  can be partitioned in cells of the form

$$\{(x,t)\subseteq S\times K\mid \operatorname{ord} a(x) \square_1 \operatorname{ord} (t-c(x)) \square_2 \operatorname{ord} b(x); t-c(x)\in \lambda P_n\},\$$

where  $a, b, c: S \to K$  are definable functions,  $\square_i$  are either < or no condition,  $P_n$  denotes the set of n-th powers and  $\lambda \in K$ . Moreover, she showed that the existence of definable Skolem functions is a necessary condition for the existence of a decomposition using cells of this form. Note that is currently not known whether all P-minimal structures admit definable Skolem functions. Work by the second author on reducts of p-adically closed fields [18] seems to indicate that this may not be the case, and hence some caution is warranted when making this assumption.

One way to deal with this uncertainty is to replace P-minimality by a (possibly) more restrictive notion, explicitly adding the existence of Skolem function as a condition. An example of this approach is the recent attempt of Darnière (see [7] for his preprint), who suggests a notion of so-called P-optimal structures. Even with these stronger assumptions, it is still an open problem whether some version of the Denef-Cluckers preparation theorem holds for arbitrary P-minimal structures.

In this paper, we take an alternative approach: the results we present here do not rely on the existence of definable Skolem functions. Instead, we decided to shift the emphasis back to possible applications. While our versions of cell decomposition/preparation are certainly weaker than versions known for individual structures, they are still strong enough to prove the theorems that initially motivated their development. Theorem 4.1 is an illustration of this.

The rest of the paper is organized as follows. In section 2, we explain our results in more detail. The proof of the cell decomposition and preparation theorems will be given in section 3. In section 4 we show that constructible functions form a class that is stable under integration. This result will be used to derive the rationality of Poincaré series.

Our arguments use the fact that on every model M of Presburger arithmetic, there exists a definable total order  $\lhd$  of M, such that every definable subset of M has a  $\lhd$ -minimal element. We present this result in an appendix, as part of a more general framework. As a corollary, we obtain that Presburger arithmetic has elimination of imaginaries (a result already proven in [2]).

The authors would like to thank Raf Cluckers for for stimulating conversations during the preparation of this paper.

## 2 Overview of main results

In this section we state the main results of this paper. Proofs will be deferred to later sections.

We first fix some notations. Let K be a p-adically closed field (that is, elementarily equivalent to a p-adic field). We use the notation  $q_K$  for the number of elements of the residue field  $k_K$ ,  $\mathcal{O}_K$  for the valuation ring of K, and  $\pi_K$  for a uniformizing element. Write  $\mathrm{ac}_m: K \to (\mathcal{O}_K/\pi_K^m\mathcal{O}_K)^\times \cup \{0\}$  for the m-th

angular component map, which can be defined as

$$\operatorname{ac}_{m}(x) := \left\{ \begin{array}{ll} \tilde{x} \mod \pi_{K}^{m} & \text{if } 0 \neq x = \pi_{K}^{\operatorname{ord}x} \tilde{x}, \\ 0 & \text{if } x = 0. \end{array} \right.$$

In every expansion of a p-adically closed field K, such angular component maps exist and can be defined in a unique way, as was shown in Lemma 1.3 of [3]. For notational purposes, we will fix a definable set  $S \subseteq K^{m_0} \times \Gamma_K^{m_0}$  which we call a parameter set. Given a set  $X \subseteq S \times K$  and  $s \in S$ , we write

$$X_s := \{ x \in K \mid (s, x) \in X \}$$

to denote the fiber over s. Analogously, for a definable function  $f: X \to \Gamma_K$ , we use the notation  $f_s(\cdot)$  for the function  $f(s,\cdot): X_s \to \Gamma_K$ . Given two sets A and B, we write  $\Pi_A: A \times B \to A$  for the projection onto A, and  $\Pi_B: A \times B \to B$  for the projection onto B. For a positive integer  $n \geq 1$ ,  $A^{\leq n}$  denotes  $\bigcup_{i=1}^n A^i$ .

We will work with a two-sorted version of P-minimality, where we consider both the field sort and the value group sort  $\Gamma_K \cup \{\infty\}$  to be of equal importance. Let  $(K, \Gamma_K; \mathcal{L}_2)$  be a two-sorted structure, with language  $\mathcal{L}_2 = (\mathcal{L}, \mathcal{L}_{Pres}, \text{ord})$ . Here  $\mathcal{L}$ , the language for the K-sort, is assumed to be an expansion of the ring language  $\mathcal{L}_{\text{ring}}$ . For the value group sort  $\Gamma_K \cup \{+\infty\}$ , we use the language of Presburger arithmetic  $\mathcal{L}_{Pres} = (+, -, <, \equiv_n)$ . The sorts are connected through the valuation map ord :  $K \to \Gamma_K \cup \{+\infty\}$ . If the language  $\mathcal{L}_2$  is clear from the context, we will just write  $(K, \Gamma_K)$ . By a definable set we mean definable with parameters.

**Definition 2.1.** A two-sorted structure  $(K, \Gamma_K; \mathcal{L}_2)$  with  $\mathcal{L}_2 = (\mathcal{L}, \mathcal{L}_{\text{Pres}}, \text{ord})$  and  $\mathcal{L}_{\text{ring}} \subseteq \mathcal{L}$  is said to be P-minimal if the underlying structure  $(K, \mathcal{L})$  is P-minimal, that is, for every  $(K', \mathcal{L})$  elementarily equivalent to  $(K, \mathcal{L})$ , the  $\mathcal{L}$ -definable subsets of K' are  $\mathcal{L}_{\text{ring}}$ -definable.

This definition is motivated by the following observation, which was based on Wagner's results on definable functions in one variable on certain ordered abelian groups [21].

**Theorem 2.2** (Cluckers [2], Lemma 2 and Theorem 6). Let  $(K, \mathcal{L})$  be a P-minimal field.

For any  $\mathcal{L}$ -definable set  $X \subseteq (K^{\times})^m$ , the set

$$\operatorname{ord}(X) := \{ (\operatorname{ord} x_1, \dots, \operatorname{ord} x_m) \in \Gamma_K^m \mid (x_1, \dots, x_m) \in X \}$$

is  $\mathcal{L}_{Pres}$ -definable.

Let  $S \subseteq \Gamma_K^m$  be a Presburger-definable set. Then the set

$$\operatorname{ord}^{-1}(S) := \{ (x_1, \dots, x_m) \in (K^{\times})^m \mid \operatorname{ord} x \in S \}$$

is  $\mathcal{L}_{ring}$ -definable.

This theorem implies that, given a (mono-sorted) P-minimal structure  $(K, \mathcal{L})$ , the valuation map ord :  $K \to \Gamma_K \cup \{\infty\}$  induces a two-sorted structure  $(K, \Gamma_K)$  where every definable subset of  $\Gamma_K^m$  is  $\mathcal{L}_{\text{Pres}}$ -definable, and hence it is natural to take  $\mathcal{L}_{\text{Pres}}$  as the language for the value group sort.

Note that a two-sorted P-minimal structure  $(K, \Gamma_K)$  cannot have definable Skolem functions since any definable section of ord contradicts the assumption of P-minimality.

We will now explain our notion of cells and cell decomposition. We distinguish the following two kinds of cells:

**Definition 2.3** (Cells). Let  $(K, \Gamma_K)$  be an  $\mathcal{L}_2$ -structure.

• A subset  $C \subseteq S \times K$  is a K-cell if it is of the form

$$C = \left\{ (s,t) \in D \times K \middle| \begin{array}{l} \alpha(s) \square_1 \operatorname{ord}(t - c(s)) \square_2 \beta(s), \\ \operatorname{ord}(t - c(s)) \equiv k \mod n, \\ \operatorname{ac}_m (t - c(s)) = \xi \end{array} \right\},$$

where D, the base of the cell, is a definable subset of S, c is a definable function  $c: D \to K$ ,  $\alpha, \beta$  are  $\mathcal{L}_2$ -definable functions  $D \to \Gamma_K$ ,  $k, n, m \in \mathbb{N}$ ,  $\xi \in \operatorname{ac}_m(K)$  and the symbols  $\square_i$  may denote < or no condition. If  $\operatorname{ac}_m(t-c(s))=0$  (and hence t=c(s)), one should ignore the first two conditions.

• A subset  $B \subseteq S \times \Gamma_K$  is a  $\Gamma$ -cell if it is of the form

$$B = \left\{ (s, \gamma) \in D \times \Gamma_K \middle| \begin{array}{c} \alpha(s) \square_1 \ \gamma \square_2 \ \beta(s), \\ \gamma \equiv k \mod n \end{array} \right\},$$

where D, the base of the cell, is a definable subset of S,  $\alpha, \beta$  are definable functions  $D \to \Gamma_k$ ,  $k, n \in \mathbb{N}$  and the squares  $\square_i$  may denote < or no condition.

To each cell, one can associate a tuple  $(\Box, k, n, m, \xi)$ , respectively  $(\Box, k, n)$ , where  $\Box = (\Box_1, \Box_2) \in \{\emptyset, <\}^2, \xi \in \operatorname{ac}_m(K)$ , and (k, n, m) is a triple of nonnegative integers such that k < n. These tuples will be referred to as the *type* of the respective cells. We denote by  $P_K$  (resp.  $P_{\Gamma}$ ) the set of all possible types of K-cells (resp.  $\Gamma$ -cells).

We obtain cell decomposition results for definable sets both of the form  $X \subseteq S \times K$  and  $X \subseteq S \times \Gamma_K$ . Let us first consider the case where the last variable belongs to the  $\Gamma_K$ -sort. When X is a set  $X \subseteq S \times \Gamma_K$ , we obtain a partition into a finite union of  $\Gamma$ -cells. Moreover, for functions  $f: X \subseteq S \times \Gamma_K \to \Gamma_K$ , we describe explicitly how, on each cell in the decomposition of X, the value of f depends on the last variable.

**Proposition 2.4** (Function preparation). Let  $f: X \subseteq S \times \Gamma_K \to \Gamma_K$  be definable in a P-minimal structure  $(K, \Gamma_K)$ . There exists a finite partition of

X in  $\Gamma$ -cells C, such that on each cell C with type  $(\delta, k, n)$ , the function f has the form

$$f(x,\gamma) = a\left(\frac{\gamma - k}{n}\right) + \delta(x),$$

where  $a \in \mathbb{Z}, n, k \in \mathbb{N}$  and  $\delta$  is a definable function  $S \to \Gamma_K$ .

If X is a subset of  $S \times K$ , the statement of our cell decomposition result is more subtle. The main difference between classical cell decomposition and K-cell decomposition arises at the level of centers. In the classical definition, the centers appear as the images of definable functions from the parameter set S. Instead, a K-cell decomposition provides a partition of X into sets  $X_i$ , which are essentially a finite union of cells, together with a definable  $\Sigma_i$  containing all their possible tuples of centers. This type of decomposition is sufficiently strong for the computation of integrals: the Haar measure is translation invariant, and hence the centers are not of great importance here.

**Theorem-Definition 2.5** (K-cell decomposition). Let  $(K, \Gamma_K)$  be a P-minimal structure, and  $X \subseteq S \times K$  be a definable set. There exists a finite partition of X in sets  $X_i \subseteq S_i \times K$ . On each part  $X_i$ , there is an integer r and r associated K-cells  $C_j$  of the form

$$C_{j} := \left\{ (s, t) \in S_{i} \times K \middle| \begin{array}{l} \alpha_{j}(s) \square_{1, j} \text{ ord } t \square_{2, j} \beta_{j}(s) \land \\ \text{ord } t \equiv k_{j} \mod n \land \\ ac_{m}(t) = \xi_{j} \end{array} \right\},$$

where  $\alpha_j, \beta_j : S_i \to \Gamma_K$  are definable functions and  $(\Box_j, k_j, n, m, \xi_j) \in P_K$ . To each  $X_i$ , we associate a definable set  $\Sigma_i \subseteq S_i \times K^r$ , which has the following property. To any function

$$\sigma: S_i \to K^r: s \mapsto (\sigma_1(s), \dots, \sigma_r(s)),$$

whose graph is contained in  $\Sigma_i$ , we can associate a (bijective) translation  $T_{\sigma}$ :  $\sqcup_j C_j \to X_i$ , defined by

$$T_{\sigma}(s,t) = (s,t-\sigma_j(s))$$
 for all  $(s,t) \in C_j$ .

The tuple  $\{(X_i)_i, (\Sigma_i)_i, (C_{\delta_{ij}})_{i,j}\}$ , will be called a K-cell decomposition of X.

Notice that the cells  $C_j$  are not necessarily disjoint (in fact, some of them may even coincide.) What we obtain is a family of bijective translations  $T_{\sigma}$  between the disjoint union  $\sqcup_j C_j$  and one of the parts  $X_i$ . Also note that while the sets  $\Sigma_i$  are definable, we cannot assure that any of the individual curves  $\sigma$  contained in it will be definable. (If a definable  $\sigma$  exists and X only consists of K-variables, a cell decomposition with the functions  $\sigma(x)$  as centers will be very similar to what Mourgues obtained.)

In the second part of the paper (section 4), we discuss applications of the preparation and cell decomposition theorems. We will restrict our attention

to the case where K is a p-adic field. The results in other sections are valid for arbitrary p-adically closed fields.

Inspired by his rationality results, Denef decided to introduce the class of constructible functions:

**Definition 2.6.** Let X be an  $\mathcal{L}_2$ -definable set. Write  $\mathbb{A}_{q_K}$  for the ring

$$\mathbb{A}_{q_K} := \mathbb{Z}\left[q_K, q_K^{-1}, \left(\frac{1}{1 - q_K^{-i}}\right)_{i \in \mathbb{N}, i > 0}\right].$$

We say that a function  $f: X \to \mathbb{Q}$  is  $\mathcal{L}_2$ -constructible if it is contained in the  $\mathbb{A}_{q_K}$ -algebra generated by functions of the forms

$$\alpha: X \to \mathbb{Z}$$
 and  $X \to \mathbb{Z}: x \mapsto q_K^{\beta(x)}$ ,

where  $\alpha$  and  $\beta$  are  $\mathcal{L}_2$ -definable and  $\mathbb{Z}$ -valued.

When  $\mathcal{L}$  is  $\mathcal{L}_{\text{ring}}$ , the subanalytic language  $\mathcal{L}_{an}$  on K (see [11] for a definition), or some intermediary languages as in [4], the class of  $\mathcal{L}_2$ -constructible functions is known to be stable under integration (see [10], [1], and [6] for the most convenient dealing with integrability conditions). We show that (see Theorem 4.1), whenever  $(K, \mathbb{Z})$  is a P-minimal structure, the class of  $\mathcal{L}_2$ -constructible functions is stable under integration:

**Theorem.** Let K be a p-adic field,  $(K, \mathbb{Z})$  a P-minimal structure, and  $f: X \subseteq S \times K^m \to \mathbb{A}_{q_K}$  an  $\mathcal{L}_2$ -constructible function such that  $f(s, \cdot)$  is measurable and integrable on  $Y_s$  for all  $s \in S$ . There exists a constructible function  $g: S \to \mathbb{A}_{q_K}$ , such that

$$g(s) = \int_{X_s} f(s, x)|dx|,$$

for all  $s \in S$ .

We also extend the rationality results known so far only for the semi-algebraic [8] and subanalytic setting [11] (and thus also for any sublanguage), obtaining the following:

**Theorem.** Suppose that  $(K, \mathbb{Z})$  is P-minimal. Let X be a definable subset of  $\mathcal{O}_K^n \times \mathbb{N}$ , and let  $a_n$  be the Haar measure of  $X_n := \{x \in \mathcal{O}_K^n \mid (x, n) \in X\}$  for each  $n \geq 0$ . Then the series  $\sum_{i \geq 0} a_i T^i$  is rational.

Here we normalize the Haar measure on  $K^n$  so that  $\mathcal{O}_K^n$  has measure 1. For a more precise statement we refer to Corollary 4.6.

## 3 Cell decomposition and function preparation

In this section we will give a proof of the cell decomposition and preparation theorems. For the comfort of the reader, we will restate (an abbreviated version of) the theorems. To ease notation, we will assume that for any definable set  $X \subseteq S \times K$ , the projection onto the parameter set S is surjective, replacing S by  $\Pi_S(X)$  if necessary. The following notation will also be used in the proofs of this section.

**Definition 3.1.** A K-cell condition is a formula  $C_{\delta}(x, y, \alpha, \beta; s)$  of the form

$$C_{\delta}(x, y, \alpha, \beta; s) := \left(\begin{array}{c} \alpha(s) \ \Box_{1} \ \operatorname{ord}(x - y) \ \Box_{2} \ \beta(s) \ \land \\ \operatorname{ord}(x - y) \equiv k \mod n \ \land \\ \operatorname{ac}_{m}(x - y) = \xi \end{array}\right),$$

where  $(\Box, k, n, m, \xi) = \delta \in P_K$ , and  $\alpha, \beta$  are definable functions  $S \to \Gamma_K$ . When no s appears,  $\alpha, \beta$  are just elements of  $\Gamma_K$ .

**Theorem** (Theorem-Definition 2.5). Let  $X \subseteq S \times K$  be a set definable in a P-minimal structure  $(K, \Gamma_K)$ . There exists a K-cell decomposition  $\{(X_i)_i, (\Sigma_i)_i, (C_{\delta_{ij}})_{i,j}\}$  of X.

*Proof.* Fix a parameter  $s \in S$ . By the cell decomposition theorem for semi-algebraic sets, see e.g. [6, theorem 3.3.2], there exists a finite partition of  $X_s$  into K-cells

$$C_s = \left\{ t \in K \middle| \begin{array}{l} \alpha_s \square_1 \operatorname{ord}(t - c_s) \square_2 \beta_s, \\ \operatorname{ord}(t - c_s) \equiv k_s \mod n_s, \\ \operatorname{ac}_{m_s}(t - c_s) = \xi_{m_s,s} \end{array} \right\}, \tag{1}$$

where  $\alpha_s, \beta_s \in \Gamma_k, c_s \in K$  and  $(\Box, k_s, n_s, m_s, \xi_{m_s,s}) \in P_K$ . Note that the cell decomposition of  $X_s$  may contain multiple cells of the same type.

Claim 3.2. There is a natural number  $N \ge 1$  such that for every  $s \in S$ , the set  $X_s$  can be partioned as a union of at most N K-cells, and for each of these cells we can assure that  $n_s, m_s < N$ .

The claim will follow by a standard compactness argument. Recall that  $P_K$  consists of elements  $\delta = (\Box_{\delta}, k_{\delta}, n_{\delta}, m_{\delta}, \xi_{\delta})$ , encoding the type of a K-cell. For each positive integer N, put

$$P_{K,N} := \{ \delta \in P_K \mid n_{\delta} < N, m_{\delta} < N \},$$

and write

$$E_{K,N} := \bigsqcup_{i=1}^{N} P_{K,N},$$

for the disjoint union of N copies of  $P_{K,N}$ . Note that  $E_{K,N}$  is a finite set. For every  $J \subseteq E_{K,N}$ , fix an enumeration  $\{\delta_1,\ldots,\delta_{|J|}\}$  of J. Given  $y=(y_1,\ldots,y_{|J|}) \in K^{|J|}$ , and  $\alpha=(\alpha_1,\alpha_2)\in\Gamma_K^{2|J|}$  we will write

$$C_J(y,\alpha) := \begin{cases} \bigcup_{i=1}^{|J|} C_{\delta_i}(K, y_i, \alpha_{1i}, \alpha_{2i}) & \text{if the sets } C_{\delta_i}(K, y_i, \alpha_{1i}, \alpha_{2i}) \\ \emptyset & \text{otherwise,} \end{cases}$$

making use of K-cell conditions  $C_{\delta_i}$  as defined in Definition 3.1. Consider the set of formulas

$$\Sigma(x) := \left\{ \bigwedge_{J \subseteq E_{K,N}} \neg (\exists y \in K^{|J|}) (\exists \alpha \in \Gamma_K^{2|J|}) [X_x = C_J(y, \alpha)] \mid N \in \mathbb{N}^* \right\}.$$

Since each  $X_s$  can be partitioned in semi-algebraic cells as in (1),  $\Sigma(x)$  is inconsistent. Hence, by compactness there exists a finite subset  $\Sigma_0(x)$  which is inconsistent. Since  $\Sigma_0(x)$  is a finite subset of  $\Sigma(x)$ , one can find a positive integer  $N_0$  such that

$$\left[ \bigwedge_{J \subseteq E_{K,N_0}} \neg (\exists y \in K^{|J|}) (\exists \alpha \in \Gamma_K^{2|J|}) [X_x = C_J(y,\alpha)] \right] \models \Sigma_0(x).$$

This implies that there must exist N > 0 such that for every  $s \in S$ 

$$(K, \Gamma_K) \models \left[ \bigvee_{J \subseteq E_{K,N}} (\exists y \in K^{|J|}) (\exists \alpha \in \Gamma_K^{2|J|}) [X_s = C_J(y, \alpha)] \right],$$

which completes the claim.

Now choose an integer N satisfying the requirements of Claim 3.2. Let  $W_N$  denote the power set of  $E_{K,N}$ . Since  $W_N$  is finite, one can put a total ordering  $\leq$  on it. We will also put an alternative ordering  $\leq$  on the value group  $\Gamma_K$ , which is defined by :

$$x \triangleleft y \Leftrightarrow (0 \leqslant x < y) \lor (0 < x \leqslant -y) \lor (0 < -x < y) \lor (0 < -x < -y). \tag{2}$$

This produces a total ordering on  $\Gamma_K$  which can be extended to  $\Gamma_K^k$  lexicographically. We will also denote this extension by  $\triangleleft$ . The important property of the order  $\triangleleft$  is that every definable set of  $\Gamma_K$  has a  $\triangleleft$ -smallest element (for a proof of this, see the appendix, in particular A.7). Now consider the map

$$\tau:S\to W_N\times (\Gamma_K)^{\leq 2|W_N|}:s\mapsto (\tau_1(s),\tau_2(s)),$$

where  $\tau_1, \tau_2$  are defined as follows:

• put  $\tau_1(s) = J$ , if J is the  $\lessdot$ -smallest element of  $W_N$  such that

$$(K, \Gamma_K) \models \left[ (\exists y \in K^{|J|}) (\exists \alpha \in \Gamma_K^{2|J|}) [X_s = C_J(y, \alpha)] \right].$$

The claim ensures the existence of at least one such J in  $W_N$ .

• let  $\tau_2(s)$  be the  $\triangleleft$ -smallest tuple  $\alpha \in \Gamma_K^{2|\tau_1(s)|}$  such that

$$(K, \Gamma_K) \models (\exists y \in K^{|J|})[X_s = C_{\tau_1(s)}(y, \alpha)].$$

It is clear that the function  $\tau$  will be definable, using some fixed representation for the finite index set  $W_N$ . For each  $J \in \tau_1(S)$ , let  $S_J$  be the set  $\{s \in S \mid \tau_1(s) = J\}$ . These sets induce a partition of X into sets  $X_J := \{(s,x) \in X : s \in S_J\}$ . We show that this partition satisfies all conditions stated in the theorem. Fix  $\{\delta_1, \ldots, \delta_{|J|}\} = J \in \tau_1(S)$ . The integer r associated to  $X_J$  is precisely r := |J|. Let  $\Sigma_J := \bigcup_{s \in S_J} \Sigma_{J,s}$  be the set consisting of fibers

$$\Sigma_{J,s} := \{ y \in K^r \mid [X_s = C_J(y, \tau_2(s))] \}.$$

Note that these sets are non-empty by definition of  $\tau_1$ . Given any function  $\sigma: S_J \to K^r: s \mapsto (\sigma_1(s), \ldots, \sigma_r(s))$  whose graph is contained in  $\Sigma_J$ , one then has that  $X_s = C_J(\sigma(s), \tau_2(s))$  for all  $s \in S_J$ . For  $1 \leq j \leq r$  and  $i \in \{1, 2\}$ , define  $\alpha_{ij}: S_J \to \Gamma_K$  to be the  $ij^{\text{th}}$ -component in the tuple  $\tau_2(s)$ . We obtain that

$$X_J = \bigcup_{j=1}^r \{(s, x) \in S_J \times K \mid C_{\delta_j}(x, \sigma_j(s), \alpha_{1j}, \alpha_{2j}, s)\}.$$

Taking the r K-cells associated to  $X_J$  given by

$$C_i := \{(s, x) \in S_J \times K \mid C_{\delta_i}(x, 0, \alpha_{1i}, \alpha_{2i}, s)\}, 1 \le j \le r,$$

it is clear that the translation map  $T_{\sigma}$  stated in the theorem gives the required bijection.

We now present the preparation theorem for definable subsets of the form  $X \subseteq S \times \Gamma_K$ . The proof follows a similar scheme as the previous one.

**Proposition** (Proposition 2.4). Let  $(K, \Gamma_k, \mathcal{L}_2)$  be P-minimal. Let  $X \subseteq S \times \Gamma_K$  and  $f: X \to \Gamma_K$  a definable function. There exists a finite decomposition of X into  $\Gamma$ -cells C such that on each such cell C, there exists a constant  $a_C \in \mathbb{Z}$  and a definable function  $\delta: D \to \Gamma_K$ , such that for all  $(x, \gamma) \in C$ ,

$$f(x,\gamma) = a_C \left(\frac{\gamma - n_0}{n}\right) + \delta(x).$$

*Proof.* Since  $(K, \mathcal{L})$  is P-minimal, it follows from Theorem 2.2 that each of the fibers  $X_s$  is Presburger definable. Cluckers [2] obtained a cell decomposition theorem for Presburger structures. Applying this to the sets  $X_s$ , yields that each  $X_s$  can be partitioned into a finite union of cells of the form

$$C_s := \{ \gamma \in K \mid \alpha_s \square_1 \gamma \square_2 \beta_s \text{ and } \gamma \equiv n_0 \mod n_s \},$$

where  $\alpha_s, \beta_s \in \Gamma_K$  and  $n_s \in \mathbb{N}$  are constants depending on s. Also note that for any  $s \in S$ , the graph of the function  $f_s$  will be a Presburger set, by the assumption of P-minimality. Indeed, the related set

$$G_s := \{(x, y) \in (K^{\times})^2 \mid f_s(\operatorname{ord} x) = \operatorname{ord} y\}$$

is definable in a P-minimal structure, and hence by Theorem 2.2, the set

$$Graph(f_s) = \{(\operatorname{ord} x, \operatorname{ord} y) \in \Gamma^2 \mid (x, y) \in G_s\},\$$

is Presburger definable. This means that each  $f_s$  is a Presburger definable function, and hence must be piecewise linear (with coefficients in  $\mathbb{Q}$ ). In particular, the above partition can be taken such that on each  $C_s$ , there exist constants  $a_s \in \mathbb{Z}$ ,  $\delta_s \in \gamma_K$  such that for all  $\gamma \in C_s$ , we have that

$$f_s(\gamma) = a_s \left(\frac{\gamma - n_0}{n_s}\right) + \delta_s.$$

Claim 3.3. There is a natural number  $N \geq 1$  such that for every  $s \in S$ , the set  $X_s$  can be partioned as a union of at most N  $\Gamma$ -cells, and for each of these cells we can assure that  $n_s, |a_s| < N$ .

The claim follows by compactness and P-minimality using an analogous argument to the one presentend in Claim 3.2.

For an integer N satisfying the requirements of the claim, let

$$P_{\Gamma,N} := \{(\Box, k, n) \in P_{\Gamma} \mid n < N\}$$
 and  $E_{\Gamma,N} := \bigsqcup_{i=1}^{N} P_{\Gamma,N}$ .

Recall that  $P_{\Gamma}$  consists of elements  $\delta = (\Box_{\delta}, k_{\delta}, n_{\delta})$ , encoding the type of a  $\Gamma$ -cell. We will use the notation

$$C_{\delta}(\alpha, \beta) := \{ \gamma \in K \mid \alpha \square_{\delta, 1} \ \gamma \square_{\delta, 2} \ \beta \land \gamma \equiv k_{\delta} \mod n_{\delta} \}.$$

For every  $J \subseteq E_{\Gamma,N}$ , fix an enumeration  $\{\delta_1, \ldots, \delta_{|J|}\}$  of J. Given  $\alpha = (\alpha_1, \alpha_2) \in \Gamma_K^{2|J|}$ , we put

$$C_J(\alpha) := \begin{cases} \bigcup_{i=1}^{|J|} C_{\delta_i}(\alpha_{1i}, \alpha_{2i}) & \text{if the sets } C_{\delta_i}(\alpha_{1i}, \alpha_{2i}) \\ \emptyset & \text{otherwise.} \end{cases}$$

Let  $a \in \mathbb{Z}^{|J|}$  be such that  $|a_i| < N$  for all  $1 \le i \le |J|$ . Let x be a tuple of variables of the same length (and sorts) as elements in S, and  $\gamma$  a  $\Gamma_K$ -variable of length 1. The tuple  $\alpha = (\alpha_1, \alpha_2, \alpha_3)$  consists of (tuples of)  $\Gamma_K$ -variables:  $\alpha_1, \alpha_2$  and  $\alpha_3$  all have length |J|. We define the formula  $\phi_{J,a}(x,\alpha)$  as

$$\phi_{J,a}(x,\alpha) := \left( \begin{array}{c} X_x = C_J(\alpha_1, \alpha_2) \land \\ \bigwedge_{1 \le i \le |J|} (\forall \gamma) \left[ \gamma \in C_{\delta_i}(\alpha_{1i}, \alpha_{2i}) \to \left( f_x(\gamma) = a_i \left( \frac{\gamma - k_{\delta_i}}{n_{\delta_i}} \right) + \alpha_{3i} \right) \right] \end{array} \right).$$

Roughly, this formula states that the set  $X_x$  can be decomposed into finitely many disjoint  $\Gamma$ -cells, on each of which the function  $f_s$  satisfies the required preparation condition. Define the set  $W_N$  by

$$W_N := \{(J, a) : J \subseteq E_{\Gamma, N}, a \in \mathbb{Z}^{|J|}, |a_i| < N \text{ for all } 1 \le i \le |J|\}.$$

Since  $W_N$  is finite, one can put a total ordering  $\leq$  on it. As before we work with an alternative total ordering  $\leq$  on the value group  $\Gamma_K$  defined as in equation (2). We proceed as in the K-cell decomposition theorem and define a map

$$\sigma: S \to W_N \times (\Gamma_K)^{\leq 3|W_N|}: s \mapsto (\sigma_1(s), \sigma_2(s)),$$

where  $\sigma_1, \sigma_2$  are defined as follows:

• put  $\sigma_1(s) = (J, a)$ , if (J, a) is the  $\lessdot$ -smallest element of  $W_N$  such that

$$(K, \Gamma_K) \models (\exists \alpha \in \Gamma_K^{3|J|}) \phi_{J,a}(s, \alpha).$$

Claim 3.3 ensures the existence of at least one such (J, a) in  $W_N$ .

• let  $\sigma_2(s)$  be the  $\triangleleft$ -smallest tuple  $\alpha \in \Gamma_K^{3|\sigma_1(s)|}$  such that

$$(K, \Gamma_K) \models \phi_{\sigma_1(s)}(s, \alpha).$$

It is easy to see that the function  $\sigma$  will be definable, using some fixed representation of the finite index set  $W_N$ . We recover the  $\Gamma$ -cell decomposition for X and the linear functions satisfying the preparation condition in the following way. For each  $\lambda = (J, a) \in \sigma_1(S)$ , we define sets  $S_{\lambda}$  and  $X_{\lambda}$ , as

$$S_{\lambda} := \{ s \in S \mid \sigma_1(s) = \lambda \} \text{ and } X_{\lambda} := \{ (s, x) \in X : s \in S_{\lambda} \}.$$

This gives us a finite partition of X as  $X = \bigcup_{\lambda} X_{\lambda}$ . We will now partition the sets  $X_{\lambda}$  as a finite union of  $\Gamma$ -cells, on each of which f will have the required form.

For  $1 \leq j \leq |J|$  and  $i \in \{1, 2, 3\}$ , define  $\alpha_{\lambda ij} : S_{\lambda} \to \Gamma_K$  to be the  $ij^{\text{th}}$ -coordinate of  $\sigma_2(s)$ . Note that these functions are indeed definable, since  $\sigma_2$  is definable. The above construction now implies that

$$X_{\lambda} = \bigcup_{1 \leqslant j \leqslant |J|} C_{\lambda,j},$$

where  $C_{\lambda,j} := \{(s,\gamma) \in S_{\lambda} \times K \mid \alpha_{\lambda 1j} \square_{\delta_{j},1} \ \gamma \square_{\delta_{j},2} \ \alpha_{\lambda 2j} \land \gamma \equiv k_{\delta_{j}} \mod n_{\delta_{j}} \}$ . The formula  $\phi_{\lambda}(s,\alpha)$  then ensures that for all  $(s,\gamma) \in C_{\lambda,j}$ , it holds that

$$f_{\mid C_{\lambda,j}}(s,\gamma) = a_j \left(\frac{t - k_{\delta_j}}{n_{\delta_j}}\right) + \alpha_{\lambda 3j},$$

which completes the proof.

Remark. Proposition 2.4 can be used to translate theorems for parametrized Presburger definable sets  $X \subseteq S \times \Gamma_K^m$  to two-sorted P-minimal structures, in the following sense. The same theorems will hold in any two-sorted P-minimal structure, where the parameter set S can now be any  $\mathcal{L}_2$ -definable set containing variables in both K and  $\Gamma_K$ , and the involved Presburger-definable functions should be replaced by functions which are piecewise linear in the  $\Gamma_K$ -variables, in the sense of Proposition 2.4. The corollary stated below is an example of this.

Given a P-minimal structure  $(K, \Gamma_K, \mathcal{L}_2)$ , we call a definable function  $f: X \subseteq S \times \Gamma_K^m \to S \times \Gamma_K^l$  linear over S if there is a definable function  $g: S \to \Gamma_K^l$  and a linear definable function  $a: \Gamma_K^m \to \Gamma_K^l$  such that f(s,t) = (s,g(s)+a(t)) for all  $(s,t) \in X$ . We write H for the set  $H:=\{x \in \Gamma_K \mid x \geq 0\}$ .

**Corollary 3.4.** (Parametric rectilinearization) Let  $(K, \Gamma_K, \mathcal{L}_2)$  be a P-minimal structure and  $X \subseteq S \times \Gamma_K^m$  be a definable set. There exists a finite partition of X into definable sets such that the following holds.

For each part A, there is a set  $B \subseteq S \times \Gamma_K^m$  and a definable bijection  $\rho: A \to B$  which is linear over S such that, for each  $s \in S$ , the set  $B_s$  is a set of the form  $\Lambda_s \times H^l$  for a bounded subset  $\Lambda_s \subseteq H^{m-l}$ , depending on s (in a definable way), and for an integer  $l \geq 0$  only depending on A.

*Proof.* The proof is almost word for word the proof the same as the proof of the Parametric rectilinearization Theorem for Presburger definable sets (Theorem 3 in [2]). One just needs to replace every application of the Presburger function preparation theorem (Theorem 1 in [1]) by Proposition 2.4.

## 4 Integration and rationality

In this section, K denotes a p-adic field, so the value group  $\Gamma_K$  will just be  $\mathbb{Z}$ . Two types of integrals will appear. When integrating over (subsets of)  $K^m$ , the Haar measure  $\mu$  is used. When integrating over  $\mathbb{Z}^n$ , we use the counting measure. The notation  $\int_X |dx|$  will be used in both contexts, adapting the measure |dx| to the sort of the variables involved.

The results below are stated for an  $\mathcal{L}_2$ -constructible function  $f: X \subseteq S \times Y \to \mathbb{A}_{q_K}$ , where both S and Y are definable sets and S is considered a parameter set. Note that both S and Y may contain variables in both the K-sort and the  $\mathbb{Z}$ -sort, unless explicitly stated otherwise. Recall that the definition of constructible functions was given in Definition 2.6. For such a fuction f, we define its locus of integrability as the set

$$\operatorname{Int}(f,S) := \{ s \in S \mid f(s,\cdot) \text{ is measurable and integrable on } Y_s \}.$$

The main result of this section is the following theorem.

**Theorem 4.1.** Let K be a p-adic field and  $(K, \mathbb{Z}, \mathcal{L}_2)$  be a P-minimal structure. Let S be a definable set, and  $f: X \subseteq S \times Y \to \mathbb{A}_{q_K}$  an  $\mathcal{L}_2$ -constructible function such that Int(f, S) = S. There exists an  $\mathcal{L}_2$ -constructible function  $g: S \to \mathbb{A}_{q_K}$ , such that

$$g(s) = \int_{X_s} f(s, y)|dy|,$$

for all  $s \in S$ .

This is a partial generalization of results which were already proven for specific cases by Cluckers, Gordon and Halupczok in [6]. The generalization is

partial because their results do not require the assumption  $\operatorname{Int}(f,S)=S$ , instead relying on an interpolation lemma, replacing f by a function  $\tilde{f}$  that coincides with f on its locus of integrability, and for which  $\operatorname{Int}(\tilde{f},S)=S$ . If a similar interpolation lemma can be proven to hold in general P-minimal structures, the assumption that  $\operatorname{Int}(f,S)=S$ , can be removed from our result as well.

**Proposition 4.2.** Theorem 4.1 holds when  $X \subseteq \mathbb{Z}^r$ .

*Proof.* Note that in this case, X is  $\mathcal{L}_{Pres}$ -definable, by Theorem 2.2. Proofs can be found in [6], Theorem 2.1.6.

**Proposition 4.3.** Theorem 4.1 holds when  $\mathcal{L}_2 = \mathcal{L}_{rinq,2}$  or  $\mathcal{L}_{an,2}$ .

Proof. See [6], Theorem 3.1.1.

As a first step towards a general proof of Theorem 4.1, we show that it already holds when  $Y \subseteq \mathbb{Z}^r$ :

П

**Proposition 4.4.** Theorem 4.1 holds when  $Y \subseteq \mathbb{Z}^r$ .

*Proof.* This is essentially a consequence of Proposition 2.4 (see also the remark on page 13). In [6], this proposition was proven under the assumption that  $\mathcal{L}_2 = \mathcal{L}_{\text{ring},2}$  or  $\mathcal{L}_{\text{an},2}$ . Part (1) corresponds to Theorem 3.4.5 and part (2) to Theorem 3.1.1 in [6]. If one replaces their Parametric rectilinearization Theorem (Proposition 3.4.4 in [6]) by Corollary 3.4, the same proof also works for two-sorted *P*-minimal structures.

We will reduce the general case to Proposition 4.4 using the following observation on the measure of definable sets.

**Proposition 4.5.** Let  $(K, \Gamma_K, \mathcal{L}_2)$  be a P-minimal structure. Let  $X \subseteq S \times T$  be a definable set, where T is K or  $\mathbb{Z}$ . There exists a constructible function  $g: S \to \mathbb{A}_{q_K}$ , such that g uniformly measures the fibers  $X_s$ , that is,

$$g(s) = \int_{X_s} |dt|,$$

whenever  $X_s$  has finite measure. Moreover, the set

$$\tilde{S} := \{ s \in S \mid X_s \text{ has finite measure} \},$$

is definable.

*Proof.* When  $T = \mathbb{Z}$ , this is a consequence of Proposition 4.4, where f is the constant function f(x) = 1 for all  $x \in X$ . The fact that the set  $\tilde{S}$  is definable follows from the cell-decomposition part of Proposition 2.4. Indeed, a  $\Gamma$ -cell  $B \subseteq \mathbb{Z}$  (as in definition 2.3) has finite measure if and only if both  $\square_1$  and  $\square_2$  are < on such cell. This is a definable condition.

Let us now consider the case where T = K. By (the translation version of) the K-cell decomposition theorem (i.e., Theorem 2.5), we can partition X in

parts  $X_i \subseteq S_i \times K$ . On each of these parts, for any choice of a function  $\sigma$  with image contained in  $\Sigma$ , we have that

$$\int_{(X_i)_s} |dt| = \int_{T_\sigma(\sqcup(C_j)_s)} |dt| = \sum_j \left[ \int_{(C_j)_s} |dt| \right],$$

and hence it suffices to compute the integral  $\int_{(C_j)_s} |dt|$ . Assume that  $C_j$  is the zero-centered K -cell

$$C_{j} := \left\{ (s,t) \in S_{i} \times K \middle| \begin{array}{l} \alpha_{j}(s) \square_{j,1} \text{ ord } t \square_{j,2} \beta_{j}(s), \\ \text{ord } t \equiv k_{j} \mod N, \\ \text{ac}_{M}(x) = \xi_{j} \end{array} \right\}.$$

Computing the measures of these cells, we get that

$$\int_{(C_j)_s} |dt| = \sum_{\tau \in T_j} \mu \left( \xi_j \pi_K^{k_j + \tau N} (1 + \pi^M \mathcal{O}_K) \right),$$

$$= |\xi| q_K^{-(k_j + M)} \sum_{\tau \in T_j} (q_K^{-N})^{\tau},$$

where  $\pi_K$  is a uniformizing element for K and  $T_j$  is the set

$$T_j := \{ \tau \in \Gamma_K \mid \alpha_j(s) \square_{j,1} \ k_j + \tau N \square_{j,2} \ \beta_j(s) \}.$$

It is easy to see that  $(C_j)_s$  (and hence X) can only have finite measure if  $\square_{j,1}$  denotes < for  $j=1,\ldots,l$ . Since this is a property of the cell, this is a definable condition.

We get the following results for this sum. If we put  $\tilde{\alpha} := \lfloor \frac{\alpha - k}{N} \rfloor + 1$ , and  $\tilde{\beta} := \lceil \frac{\beta - k}{N} \rceil - 1$  (clearly these are still definable functions), then we get that

$$\sum_{\tau \in T_j} (q_K^{-N})^{\tau} = \begin{cases} \frac{q_K^{-N\tilde{\alpha}_j}}{1 - q_K^{-N}} & \text{if } \Box_{j,2} = \emptyset, \\ \frac{1}{1 - q_K^{-N}} (q_K^{-N\tilde{\alpha}_j} - q_K^{-N\tilde{\beta}_j}) & \text{if } \Box_{j,2} = <. \end{cases}$$
(3)

In both cases we obtain an  $\mathcal{L}_2$ -constructible function. Hence, we can conclude that  $\mu(X_s)$  is given by a constructible function as well.

We can now complete the proof of the main theorem:

Proof of Theorem 4.1. Since  $\operatorname{Int}(f,S)=S$ , by Fubini's theorem, the general result can be obtained by iteration and we may assume that either  $Y=\mathbb{Z}$ , or Y=K. The first case is already included in Proposition 4.4, so we only need to consider the case Y=K.

A general constructible function  $f: X \subseteq S \times K \to \mathbb{A}_{q_K}$  has the form

$$f(s,x) = \sum_{i=1}^{r} a_i q_K^{f_{i0}(s,x)} \prod_{j=1}^{r'} f_{ij}(s,x),$$

where the  $f_{ij}$  are definable functions  $X \to \mathbb{Z}$ , and  $a_i \in \mathbb{A}_{q_K}$ . Now put  $\gamma = (\gamma_{ij})_{i,j}$  and consider the set

$$G := \{ (s, \gamma, x) \in S \times \mathbb{Z}^{(r'+1)r} \times K \mid \gamma_{ij} = f_{ij}(s, x) \},$$

which is a permutated version of the combined graphs of the functions generating f. To ease notations, we will sometimes consider G as a subset of  $D \times K$ , where  $D = \prod_{S \times \Gamma_K^{(r'+1)r}}(G)$ . Let  $\mu$  denote the usual Haar measure. The integral of  $f_s$  can be written as a sum ranging over  $\mathrm{Im}(f_s)$ :

$$\int_{X_s} f(s, x) |dx| = \sum_{\delta \in \operatorname{Im}(f_s)} \delta \cdot \mu \{ x \in X_s \mid f_s(x) = \delta \},$$

and this sum can be expressed in terms of the variables  $\gamma$ , to obtain a sum

$$\sum_{\gamma \in D_s} \left[ \left( \sum_{i=1}^r a_i q_K^{\gamma_{i0}} \prod_{j=1}^{r'} \gamma_{ij} \right) \cdot \mu \left( \left\{ x \in X_s \mid \bigwedge_{ij} f_{ij}(s, x) = \gamma_{ij} \right\} \right) \right]$$

This reduces the integral to a sum

$$\int_{X_s} f(s,x)|dx| = \sum_{\gamma \in D_s} \left( \sum_{i=1}^r a_i q_K^{\gamma_{i0}} \prod_{j=1}^{r'} \gamma_{ij} \right) \cdot \mu(G_{s,\gamma}). \tag{4}$$

Applying Proposition 4.5, we know that  $\mu(G_{s,\gamma})$  is given by a constructible function, whenever  $G_{s,\gamma}$  has finite measure. Since this is a definable condition, we may as well assume that the measure of  $G_{s,\gamma}$  is finite for all  $s \in S$  and  $\gamma \in \text{Im}(f_s)$ . Hence, we can conclude that

$$\int_{X_s} f(s, x) |dx| = \sum_{\gamma \in D_s} h(s, \gamma), \tag{5}$$

for some constructible function  $h: D \to \mathbb{A}_{q_K}$ . Noticing that

$$\sum_{\gamma \in D_s} h(s,\gamma) = \int_{D_s} h(s,\gamma) |d\gamma|,$$

the result follows by Proposition 4.4 applied to the constructible function h.

As a consequence of Theorem 4.1, we obtain the following rationality result.

**Corollary 4.6.** Suppose that  $(K, \mathbb{Z})$  is P-minimal. Let X be a definable subset of  $\mathbb{N} \times D$ , where D is a compact subset of  $K^m$ . Then the series  $\sum_{n\geq 0} \mu(X_n)T^n$  is a rational function. More precisely,

$$\sum_{n>0} \mu(X_n) T^n = \frac{Q(T)}{\prod_{i=1}^r (1 - q_K^{-m_i} T_i^N)},$$

for certain integers  $m_i, r \in \mathbb{N}$ ,  $N_i > 0$  and  $Q(T) \in \mathbb{A}_{q_K}[T]$ .

*Proof.* Applying Theorem 4.1 to the set  $X \subseteq \mathbb{N} \times D$ , one can find a constructible function  $g: \mathbb{N} \to \mathbb{A}_{q_K}$ , such that

$$g(n) = \int_{X_n} |dx|.$$

This function must have the form

$$g(n) = \sum_{i=1}^{r} a_i q_K^{\alpha_i(n)} \prod_j \beta_{ij}(n),$$

where  $a_i \in \mathbb{A}_{q_K}$ , and the functions  $\alpha_i$  and  $\beta_{ij}$  are Presburger-definable functions  $\mathbb{N} \to \mathbb{Z}$ , and hence it is actually  $\mathcal{L}_{\text{ring},2}$ -constructible. Our claim now follows from Denef's rationality results in the semi-algebraic case, for which we refer to eg. [8, 9].

We finish by presenting as a conjecture a version of interpolation for P-minimal constructible functions:

**Conjecture 1.** (Interpolation) Let K be a p-adic field and  $(K, \mathbb{Z}, \mathcal{L}_2)$  be a P-minimal structure. For every  $\mathcal{L}_2$ -constructible function  $f: X \subseteq S \times Y \to \mathbb{A}_{q_K}$  there exists an  $\mathcal{L}_2$ -constructible function  $g: X \subseteq S \times Y \to \mathbb{A}_{q_K}$  such that Int(g, S) = S and f(s, y) = g(s, y) whenever  $s \in Int(f, S)$ .

Assuming this conjecture, Theorem 4.1 implies the full generalization of the stability result in [6].

Corollary 4.7. Let K be a p-adic field and  $(K, \mathbb{Z}, \mathcal{L}_2)$  be a P-minimal structure and suppose that the interpolation conjecture is true. Let S be a definable set, and  $f: X \subseteq S \times Y \to \mathbb{A}_{q_K}$  an  $\mathcal{L}_2$ -constructible function. There exists an  $\mathcal{L}_2$ -constructible function  $g: S \to \mathbb{A}_{q_K}$ , such that

$$g(s) = \int_{X_s} f(s, y)|dy|,$$

whenever  $s \in Int(f, S)$ .

## A Definably well-ordered structures

**Definition A.1.** A structure  $(M, \mathcal{L})$  is said to be *definably well-ordered* if there exists a definable linear order  $\triangleleft$  on M, such that every definable subset of M has a  $\triangleleft$ -minimal element.

**Lemma A.2.** Suppose that  $(M, \mathcal{L})$  is definably well-ordered, and that  $\triangleleft$  is defined by an  $\mathcal{L}(a)$ -formula. Then every structure  $(N, \mathcal{L}(a))$  which is elementarily equivalent to  $(M, \mathcal{L}(a))$ , is definably well-orderable.

*Proof.* Let  $\phi(x,y)$  be an  $\mathcal{L}(a)$ -formula with length(x)=1. By definition,

$$M \models (\forall y)[(\exists x)\phi(x,y) \to (\exists x)(\phi(x,y) \land (\forall z)[\phi(z,y) \to x \le z])].$$

Since N and M are elementarily equivalent as  $\mathcal{L}(a)$  structures, this implies that every definable subset of N has a  $\triangleleft$ -minimal element.  $\square$ 

The previous lemma shows that being a definably well-ordered structure is a property of Th(M,a), where a is a tuple of parameters used in a formula defining a linear order that satisfies the requirements of Definition A.1. We say that a theory T is definably well-orderable if it has some definably well-ordered model where the linear order is 0-definable. The following lemma shows the relation with cartesian powers:

#### Lemma A.3. The following are equivalent:

- 1.  $(M, \mathcal{L})$  is definably well-ordered;
- 2. There is an  $\mathcal{L}(M)$ -definable linear order  $\lhd$  on  $M^n$  such that any definable subset of  $M^n$  has a  $\lhd$ -minimal element.

*Proof.* That (1) implies (2) follows by equipping  $M^n$  with the lexicographic order induced by the definable linear order on M. For the converse, suppose that n > 1 and pick an element  $a \in M^{n-1}$ . Let  $\phi(x, y)$  be a formula defining  $x \triangleleft y$ . Then  $\phi(x_1, a; y_1, a)$  defines a well-order on M.

For a theory to have models which are definably well-ordered is a very strong property. As an example we show that such theories have definable choice, and thus eliminate imaginaries (for definitions we refer to [23] and [14]).

**Proposition A.4.** An definably well-ordered structure  $(M, \mathcal{L})$  has definable choice.

*Proof.* Let  $X \subseteq M^{m+n}$  be a definable set, and  $\triangleleft$  a fixed definable linear order on M, such that definable sets in any cartesian power of M have  $\triangleleft$ -minimal elements. Define  $f: \Pi_m(X) \to M^n$  to be the function sending x to the  $\triangleleft$ -least element in  $X_x$ . Clearly, if  $X_x = X_y$  then f(x) = f(y).

Corollary A.5. A definably well-ordered structure  $(M, \mathcal{L})$  has definable Skolem functions.

Corollary A.6. A definably well-ordered structure  $(M, \mathcal{L})$  has uniform elimination of imaginaries.

Notice that being a definably well-ordered structure is stronger than having definable choice. For instance, the real field has definable choice, yet by a result of Ramakrishnan in [22] every definable order embeds in  $(\mathbb{R}^n, <_{lex})$ . Therefore, no definable linear order has minimal elements for all definable subsets of the real line. Using this, one can show that no reduct of the real field is definably well-orderable.

Even though definably well-ordered structures have strong model-theoretic properties, they are not always model-theoretically tame. For instance, the theory of arithmetic is definably well-orderable and yet model-theoretically wild. The main example of a tame well-orderable theory is Presburger Arithmetic. That this theory is well-orderable, is a consequence of the following proposition:

**Proposition A.7.** Let  $\mathcal{L}$  be a language containing  $\{\leq,-\}$  (as  $\mathcal{L}_{Pres}$ ). Then  $Th(\mathbb{Z},\mathcal{L})$  is definably well-orderable.

*Proof.* Consider the following definable order

$$x \vartriangleleft y \Leftrightarrow \begin{cases} 0 \le x < y \\ 0 \le x < -y \\ 0 \le -x < y \\ 0 \le -x < -y \end{cases}$$

On  $\mathbb{Z}$  this defines the following well-order:

$$0 \triangleleft 1 \triangleleft -1 \triangleleft 2 \triangleleft -2 \triangleleft \cdots$$

Because of lemma A.2, this completes the proof.

Notice that the linear ordering defined in the previous proposition does not necessarily define a well-order on every  $\mathbb{Z}$ -group G. However, it does define a linear order such that for any definable subset  $A \subseteq G$ , A has a  $\triangleleft$ -minimal element. As a corollary we get a result from [2]

Corollary A.8. Presburger arithmetic has elimination of imaginaries.

## References

- [1] R. Cluckers, Analytic p-adic cell decomposition and integrals, Transactions of the American Mathematical Society **356** (2003), no. 4, 1489–1499.
- [2] R. Cluckers, Presburger sets and p-minimal fields, J. Symbolic Logic 68 (2003), no. 1, 153–162. MR 1959315 (2003m:03062)
- [3] R. Cluckers and E. Leenknegt, A version of p-adic minimality, J. Symbolic Logic 77 (2012), no. 2, 621–630.
- [4] R. Cluckers and L. Lipshitz, Fields with analytic structure, J. Eur. Math. Soc. (JEMS) 13 (2011), 1147–1223, math.LO/0610666.
- [5] Raf Cluckers, Classification of semi-algebraic p-adic sets up to semi-algebraic bijection, J. Reine Angew. Math. 540 (2001), 105–114. MR 1868600 (2002i:14052)
- [6] Raf Cluckers, Julia Gordon, and Immanuel Halupczok, Integrability of oscillatory functions on local fields: transfer principles, Duke Math. J. 163 (2014), no. 8, 1549–1600. MR 3210968

- [7] Luck Darniere, Cell decomposition and dimension theory on p-optimal fields, arXiv:1412.2571v1 (2014).
- [8] J. Denef, The rationality of the Poincaré series associated to the p-adic points on a variety, Invent. Math. 77 (1984), 1–23.
- [9] J. Denef, On the evaluation of certain p-adic integrals, Séminaire de théorie des nombres, Paris 1983–84, Progr. Math., vol. 59, Birkhäuser Boston, Boston, MA, 1985, pp. 25–47. MR 902824 (88j:11031)
- [10] \_\_\_\_\_\_, Arithmetic and geometric applications of quantifier elimination for valued fields, Model theory, algebra, and geometry, Math. Sci. Res. Inst. Publ., vol. 39, Cambridge Univ. Press, Cambridge, 2000, pp. 173–198. MR 1773707 (2001e:03063)
- [11] J. Denef and Lou van den Dries, p-adic and real subanalytic sets, Ann. of Math. (2) 128 (1988), no. 1, 79–138.
- [12] Jan Denef, p-adic semi-algebraic sets and cell decomposition, J. Reine Angew. Math. 369 (1986), 154–166. MR 850632 (88d:11030)
- [13] Deirdre Haskell and Dugald Macpherson, A version of o-minimality for the p-adics, J. Symbolic Logic 62 (1997), no. 4, 1075–1092. MR 1618009 (99j:03028)
- [14] Wilfrid Hodges, A shorter model theory, Cambridge University Press, New York, NY, USA, 1997.
- [15] Jun-ichi Igusa, Complex powers and asymptotic expansions. I. Functions of certain types, J. Reine Angew. Math. 268/269 (1974), 110–130, Collection of articles dedicated to Helmut Hasse on his seventy-fifth birthday, II. MR 0347753 (50 #254)
- [16] \_\_\_\_\_, Complex powers and asymptotic expansions. II. Asymptotic expansions, J. Reine Angew. Math. 278/279 (1975), 307–321. MR 0404215 (53 #8018)
- [17] \_\_\_\_\_\_, Forms of higher degree, Tata Institute of Fundamental Research Lectures on Mathematics and Physics, vol. 59, Tata Institute of Fundamental Research, Bombay; by the Narosa Publishing House, New Delhi, 1978. MR 546292 (80m:10020)
- [18] Eva Leenknegt, *Reducts of p-adically closed fields*, Accepted for publication in Archive for Mathematical Logic **20 pp.** (2013).
- [19] A. Macintyre, On definable subsets of p-adic fields, J. Symb. Logic 41 (1976), 605-610.
- [20] Marie-Hélène Mourgues, Cell decomposition for P-minimal fields, MLQ Math. Log. Q. 55 (2009), no. 5, 487–492. MR 2568759 (2011d:03054)

- [21] Françoise Point and Frank O. Wagner, Essentially periodic ordered groups, Ann. Pure Appl. Logic 105 (2000), no. 1-3, 261–291. MR 1786147 (2002a:03073)
- [22] Janak Ramakrishnan, Definable linear orders definably embed into lexicographic orders in o-minimal structures, Proc. Amer. Math. Soc. 141 (2013), no. 5, 1809–1819. MR 3020867
- [23] Lou van den Dries, *Tame topology and o-minimal structures*, London Mathematical Society Lecture Note Series, vol. 248, Cambridge University Press, Cambridge, 1998. MR 1633348 (99j:03001)