The Reducts of the Homogeneous Binary Branching C-relation^{*}

Manuel Bodirsky Institut für Algebra, TU Dresden, Germany manuel.bodirsky@tu-dresden.de

Peter Jonsson Department of Computer and System Science Linköpings Universitet, Linköping, Sweden peter.jonsson@liu.se

Trung Van Pham Institut für Computersprachen, Theory and Logic Group TU Wien, Austria pvtrung@logic.at

October 21, 2018

Abstract

Let $(\mathbb{L}; C)$ be the (up to isomorphism unique) countable homogeneous structure carrying a binary branching C-relation. We study the reducts of $(\mathbb{L}; C)$, i.e., the structures with domain \mathbb{L} that are first-order definable in $(\mathbb{L}; C)$. We show that up to existential interdefinability, there are finitely many such reducts. This implies that there are finitely many reducts up to first-order interdefinability, thus confirming a conjecture of Simon Thomas for the special case of $(\mathbb{L}; C)$. We also study the endomorphism monoids of such reducts and show that they fall into four categories.

Contents

1	Introduction	2
2	Results	3
3	Preliminaries	4
	3.1 Model theory	5
	3.2 The structure $(L; C)$: Fraïssé-amalgamation	5
	3.3 The structure $(L; C)$: an axiomatic approach $\ldots \ldots \ldots \ldots \ldots \ldots \ldots \ldots \ldots \ldots$	6
	3.4 The reduct $(\mathbb{L}; Q)$	9
	3.5 Convex orderings of C-relations	11

^{*}The first and third author have received funding from the European Research Council under the European Community's Seventh Framework Programme (FP7/2007-2013 Grant Agreement no. 257039). The third author has received funding from the project P27600 of the Austrian Science Fund (FWF) and the project "Models on graphs: enumerative combinatorics and algebraic structures" of the Vietnam National Foundation for Science and Technology Development (NAFOSTED). The second author is partially supported by the *Swedish Research Council* (VR) under Grant 621-2012-3239.

4	Aut	tomorphism groups of reducts	11
	4.1	Permutation group preliminaries	12
	4.2	The rerooting lemma	13
	4.3	Automorphism group classification	15
5	Ran	nsey theory for the C-relation	17
	5.1	Ramsey classes	17
	5.2	Canonical functions	18
	5.3	The ordering property	18
6	End	lomorphism monoids of reducts	19
6	End 6.1	lomorphism monoids of reducts Canonical behavior without constants	19 20
6	End 6.1 6.2	Iomorphism monoids of reducts Canonical behavior without constants Canonical behavior with one constant	19 20 24
6	End 6.1 6.2	lomorphism monoids of reducts Canonical behavior without constants Canonical behavior with one constant $6.2.1$	19 20 24 24
6	End 6.1 6.2	lomorphism monoids of reducts Canonical behavior without constants Canonical behavior with one constant $6.2.1$ The behavior cut _c $6.2.2$ The behavior rer _c	19 20 24 24 25
6	End 6.1 6.2	lomorphism monoids of reducts Canonical behavior without constants Canonical behavior with one constant $6.2.1$ The behavior cut _c $6.2.2$ The behavior rer _c $6.2.3$ The behavior rer _c	19 20 24 24 25 27
6	End 6.1 6.2	domorphism monoids of reducts Canonical behavior without constants Canonical behavior with one constant 6.2.1 The behavior cut_c 6.2.2 The behavior rer_c 6.2.3 The behavior rer_c 6.2.4 Classification of behaviors with one constant	 19 20 24 24 25 27 27
6	End 6.1 6.2	domorphism monoids of reducts Canonical behavior without constants Canonical behavior with one constant $6.2.1$ The behavior cut_c $6.2.2$ The behavior rer_c $6.2.3$ The behavior rer_c $6.2.4$ Classification of behaviors with one constant Canonical behavior with two constants	 19 20 24 24 25 27 27 29

1 Introduction

A structure Γ is called *homogeneous* (or sometimes *ultra-homogeneous* in order to distinguish it from other notions of homogeneity that are used in adjacent areas of mathematics) if every isomorphism between finite substructures of Γ can be extended to an automorphism of Γ . Many classical structures in mathematics are homogeneous such as (\mathbb{Q} ; <), the random graph, and the homogeneous universal poset.

C-relations are central for the structure theory of Jordan permutation groups [1-3, 36]. They also appear frequently in model theory. For instance, there is a substantial literature on *C*-minimal structures which are analogous to o-minimal structures but where a C-relation plays the role of the order in an o-minimal structure [30, 38]. In this article we study the universal homogeneous binary branching *C*-relation (\mathbb{L} ; *C*). This structure is one of the fundamental homogeneous structures [3,26,37] and can be defined in several different ways—we present two distinct definitions in Section 3. We mention that (\mathbb{L} ; *C*) is the up to isomorphism unique countable *C*-relation which is existential positive complete in the class of all *C*-relations – see [9] for the notion of existential positive completeness.

If Γ has a finite relational signature (as in the examples mentioned above), then homogeneity implies that Γ is ω -categorical, that is, every countable model of the first-order theory of Γ is isomorphic to Γ . A relational structure Δ is called a *reduct* of Γ if Δ and Γ have the same domain and every relation in Δ has a first-order definition (without parameters) in Γ . It is well known that reducts of ω -categorical structures are again ω -categorical [33]. Two reducts Δ_1 and Δ_2 are said to be first-order interdefinable if Δ_1 is first-order definable in Δ_2 , and vice versa. Existential and existential positive¹ interdefinability are defined analogously.

It turns out that several fundamental homogeneous structures with finite relational signatures have only finitely many reducts up to first-order interdefinability. This was shown for $(\mathbb{Q}; <)$ by Cameron [24] (and, independently and in somewhat different language, by Frasnay [29]), by Thomas for the the random graph [45], by Junker and Ziegler for the expansion of $(\mathbb{Q}; <)$ by a constant [34], by Pach, Pinsker, Pluhár, Pongrácz, and Szabó for the homogeneous universal poset [42], and by Bodirsky, Pinsker and Pongrácz for the random ordered graph [19]. Thomas has conjectured that all homogeneous structures with a finite relational signature have finitely many reducts [45]. In this paper, we study the reducts of $(\mathbb{L}; C)$ up to first-order, and even up

¹A first-order formula is *existential* if it is of the form $\exists x_1, \ldots, x_m \cdot \psi$ where ψ is quantifier-free, and *existential* positive if it is existential and does not contain the negation symbol \neg .

to existential and existential positive interdefinability. Our results for reducts up to first-order interdefinability confirm Thomas' conjecture for the case of $(\mathbb{L}; C)$.

Studying reducts of ω -categorical structures has an additional motivation coming from permutation group theory. We write S_{ω} for the group of all permutations on a countably infinite set. The group S_{ω} is naturally equipped with the topology of pointwise convergence. By the fundamental theorem of Engeler, Ryll-Nardzewski, and Svenonius, the reducts of an ω -categorical structure Γ are one-to-one correspondence with the *closed* subgroups of S_{ω} that contain the automorphism group of Γ . The automorphism groups of ω -categorical structures are important and well-studied groups in permutation group theory, and classifications of reducts up to first-order interdefinability shed light on their nature. Indeed, all the classification results mentioned above make extensive use of the group-theoretic perspective on reducts.

Let us also mention that reducts of $(\mathbb{L}; C)$ are used for modeling various computational problems studied in phylogenetic reconstruction [14, 22, 23, 31, 41, 44]. When Γ is such a structure with a finite relational signature, then the constraint satisfaction problem (CSP) for the template Γ is the problem to decide for a finite structure Δ with the same signature as Γ whether there exists a homomorphism from Δ to Γ or not. For example, the CSP for $(\mathbb{L}; C)$ itself has been called the rooted triple consistency problem and it is known to be solvable in polynomial time by a non-trivial algorithm [4, 14, 31]. Other phylogeny problems that can be modeled as CSPs for reducts of $(\mathbb{L}; C)$ are the NP-complete quartet consistency problem [44] and the NP-complete forbidden triples problem [22]. To classify the complexity of CSPs of reducts of an ω -categorical structure, a good understanding of the endomorphism monoids of these reducts is important; for example, such a strategy has been used successfully in [12, 18, 21]. In this paper, we show that the endomorphism monoids of (L;C) fall into four categories. In [10] the authors give a full complexity classification for CSPs for reducts of (L;C) and make essential use of this result.

2 Results

We show that there are only three reducts of $(\mathbb{L}; C)$ up to existential interdefinability (Corollary 3). In particular, there are only three reducts of $(\mathbb{L}; C)$ up to first-order interdefinability. The result concerning reducts up to first-order interdefinability can also be shown with a proof based on known results on Jordan permutation groups (Section 4). However, we do not know how to obtain our stronger statement concerning reducts up to existential interdefinability using Jordan group techniques.

Our proof of Corollary 3 uses Ramsey theory for studying endomorphism monoids of reducts of $(\mathbb{L}; C)$. More specifically, we use a Ramsey-type result for C-relations which is a special case of Miliken's theorem [39] (see [15] for a short proof). We use it to show that endomorphisms of reducts of $(\mathbb{L}; C)$ must behave *canonically* (in the sense of Bodirsky & Pinsker [16]) on large parts of the domain and this enables us to perform a combinatorial analysis of the endomorphism monoids. This approach provides additional insights which we describe next.

Assume that Γ is a homogeneous structure with a finite relational signature whose age² has the Ramsey property (all examples mentioned above are reducts of such a structure). Then, there is a general approach to analyzing reducts up to first-order interdefinability via the transformation monoids that contain Aut(Γ) instead of the closed permutation groups that contain Aut(Γ). This Ramsey-theoretic approach has been described in [16]. We write ω^{ω} for the transformation monoid of all unary functions on a countably infinite set. The monoid ω^{ω} is naturally equipped with the topology of pointwise convergence and the closed submonoids of ω^{ω} that contain Aut(Γ) are in one-to-one correspondence with the reducts of Γ considered up to existential positive interdefinability. We note that giving a complete description of the reducts up to existential positive interdefinability is usually difficult. For instance, already the structure (\mathbb{N} ;=) admits infinitely many such reducts [8]. However, it is often feasible to describe all reducts up to existential interdefinability; here, the Random Graph provides a good illustration [17]. In this paper, we show that it is feasible to describe all reducts of (\mathbb{L} ; C) up to existential positive interdefinability. In

²The age of a relational structure Γ is the set of finite structures that are isomorphic to some substructure of Γ .

particular, we show that the reducts of $(\mathbb{L}; C)$ fall into four categories. An important category is when a reduct Γ of $(\mathbb{L}; C)$ has the same endomorphisms as the reduct $(\mathbb{L}; Q)$. This reduct is a natural *D*-relation which is associated to $(\mathbb{L}; C)$ (see Subsection 3.4), and its known complexity allows us to derive the complexity of the CSP for a large class of the reducts of $(\mathbb{L}; C)$. Those four categories are stated in the following main result of our paper.

Theorem 1. Let Γ be a reduct of $(\mathbb{L}; C)$. Then one of the following holds.

- 1. Γ has the same endomorphisms as $(\mathbb{L}; C)$,
- 2. Γ has a constant endomorphism,
- 3. Γ is homomorphically equivalent to a reduct of $(\mathbb{L};=)$, or
- 4. Γ has the same endomorphisms as $(\mathbb{L}; Q)$.

We use this result to identify in Corollary 3 below the reducts of $(\mathbb{L}; C)$ up to existential interdefinability. The proof of Corollary 3 is based on a connection between existential and existential positive definability on the one hand, and the endomorphisms of Δ on the other hand.

Proposition 2 (Proposition 3.4.7 in [6]). For every ω -categorical structure Γ , it holds that

- a relation R has an existential positive definition in Γ if and only if R is preserved by the endomorphisms of Γ and
- a relation R has an existential definition in Γ if and only if R is preserved by the embeddings of Γ into Γ.

Corollary 3. Let Γ be a reduct of $(\mathbb{L}; C)$. Then Γ is existentially interdefinable with $(\mathbb{L}; C)$, with $(\mathbb{L}; Q)$, or with $(\mathbb{L}; =)$.

Our result has important consequences for the study of CSPs for reducts of $(\mathbb{L}; C)$. To see this, note that when two structures Γ and Δ are homomorphically equivalent, then they have the same CSP. Since the complexity of CSP(Γ) has been classified for all reducts Γ of $(\mathbb{L}; =)$ (see Bodirsky and Kára [11]) and since CSP(Γ) is trivial if Γ has a constant endomorphism, our result shows that we can focus on the case when Γ has the same endomorphisms as $(\mathbb{L}; C)$ or $(\mathbb{L}; Q)$. This kind of simplifying assumptions have proven to be extremely important in complexity classications of CSPs: examples include Bodirsky & Kára [13] and Bodirsky & Pinsker [18].

This article is organized as follows. The structure $(\mathbb{L}; C)$ is formally defined in Section 3. We then show (in Section 4) how to classify the reducts of $(\mathbb{L}; C)$ up to first-order interdefinability by using known results about Jordan permutation groups. For the stronger classification up to existential definability, we investigate transformation monoids. The Ramsey-theoretic approach works well for studying transformation monoids and will be described in Section 5. The main result is proved in Section 6.

3 Preliminaries

We will now present some important definitions and results. We begin, in Section 3.1, by providing a few preliminaries from model theory. Next, we define the universal homogeneous binary branching C-relation (\mathbb{L} ; C). There are several equivalent ways to do this and we consider two of them in Sections 3.2 and 3.3, respectively. The first approach is via Fraïssé-amalgamation and the second approach is an axiomatic approach based on Adeleke and Neumann [3]. In Section 3.4, we also give an axiomatic treatment of an interesting reduct of (\mathbb{L} ; C). In Section 3.5, we continue by introducing an ordered variant of the binary branching C-relation [25] which will be important in the later sections.

3.1 Model theory

We follow standard terminology as, for instance, used by Hodges [33]. Let τ be a relational signature (all signatures in this paper will be relational) and Γ a τ -structure. When $R \in \tau$, we write R^{Γ} for the relation denoted by R in Γ ; we simply write R instead of R^{Γ} when the reference to Γ is clear. Let Γ_1 and Γ_2 be two τ -structures with domains D_1 and D_2 , respectively, and let $f: D_1 \to D_2$ be a function. If $t = (t_1, \ldots, t_k) \in (D_1)^k$, then we write f(t) for $(f(t_1), \ldots, f(t_k))$, i.e. we extend single-argument functions pointwise to sequences of arguments. We say that f preserves R iff $f(t) \in R^{\Gamma_2}$ whenever $t \in R^{\Gamma_1}$. If $X \subseteq D_1$ and $R \in \tau$ is a k-ary relation, then we say that f preserves R on X if $f(t) \in R^{\Gamma_2}$ whenever $t \in R^{\Gamma_1} \cap X^k$. If f does not preserves R (on X), then we say that f violates R (on X).

A function $f: D_1 \to D_2$ is an *embedding* of Γ_1 into Γ_2 if f is injective and has the property that for all $R \in \tau$ (where R has arity k) and all $t \in (D_1)^k$, we have $f(t) \in R^{\Gamma_2}$ if and only if $t \in R^{\Gamma_1}$.

A substructure of a structure Γ is a structure Δ with domain $S = D_{\Delta} \subseteq D_{\Gamma}$ and $R^{\Delta} = R^{\Gamma} \cap S^{n}$ for each *n*-ary $R \in \tau$; we also write $\Gamma[S]$ for Δ . The *intersection* Δ of two τ -structures Γ, Γ' is the structure with domain $D_{\Gamma} \cap D_{\Gamma'}$ and relations $R^{\Delta} = R^{\Gamma} \cap R^{\Gamma'}$ for all $R \in \tau$; we also write $\Gamma \cap \Gamma'$ for Δ .

Let Γ_1, Γ_2 be τ -structures such that $\Delta = \Gamma_1 \cap \Gamma_2$ is a substructure of both Γ_1 and Γ_2 . A τ structure Δ' is an *amalgam of* Γ_1 and Γ_2 over Δ if for $i \in \{1, 2\}$ there are embeddings f_i of Γ_i to Δ' such that $f_1(a) = f_2(a)$ for all $a \in D_{\Delta}$. We assume that classes of structures are closed under isomorphism. A class \mathcal{A} of τ -structures has the *amalgamation property* if for all $\Delta, \Gamma_1, \Gamma_2 \in \mathcal{A}$ with $\Delta = \Gamma_1 \cap \Gamma_2$, there is a $\Delta' \in \mathcal{A}$ that is an amalgam of Γ_1 and Γ_2 over Δ . A class of finite τ -structures that has the amalgamation property, is closed under isomorphism and closed under taking substructures is called an *amalgamation class*.

A relational structure Γ is called *homogeneous* if all isomorphisms between finite substructures can be extended to automorphisms of Γ . A class \mathcal{K} of τ -structures has the joint embedding property if for any $\Gamma, \Gamma' \in \mathcal{K}$, there is $\Delta \in \mathcal{K}$ such that Γ and Γ' embed into Δ . An amalgamation class has the joint embedding property since it always contains an empty structure. The following basic result is known as Fraïssé's theorem.

Theorem 4 (see Theorem 6.1.2 in Hodges [33]). Let \mathcal{A} be an amalgamation class with countably many non-isomorphic members. Then there is a countable homogeneous τ -structure Γ such that \mathcal{A} is the class of structures that embeds into Γ . The structure Γ , which is unique up to isomorphism, is called the Fraïssé-limit of \mathcal{A} .

3.2 The structure (**L**; *C*): Fraïssé-amalgamation

We will now define the structure (L; C) as the Fraïssé-limit of an appropriate amalgamation class. We begin by giving some standard terminology concerning rooted trees. Throughout this article, a *tree* is a simple, undirected, acyclic, and connected graph. A *rooted tree* is a tree T together with a distinguished vertex r which is called the *root* of T. The vertices of T are denoted by V(T). The *leaves* L(T) of a rooted tree T are the vertices of degree one that are distinct from the root r. In this paper, a rooted tree is often drawn downward from the root.

For $u, v \in V(T)$, we say that u lies below v if the path from u to r passes through v. We say that u lies strictly below v if u lies below v and $u \neq v$. All trees in this article will be rooted and binary, i.e., all vertices except for the root have either degree 3 or 1, and the root has either degree 2 or 0. A subtree of T is a tree T' with $V(T') \subseteq V(T)$ and $L(T') \subseteq L(T)$. If the root of T' is different from the root of T, the subtree is called proper subtree. The youngest common ancestor (yca) of a non-empty finite set of vertices $S \subseteq V(T)$ is the (unique) node w that lies above all vertices in S and has maximal distance from r.

Definition 5. The leaf structure of a binary rooted tree T is the relational structure (L(T); C)where C(a, bc) holds in C if and only if $yca(\{b, c\})$ lies strictly below $yca(\{a, b, c\})$ in T. We call T the underlying tree of the leaf structure. We mention that the definition of C-relation on binary rooted trees can also be obtained from the relation | on trees with a distinguished leaf [25]. The slightly non-standard way of writing the arguments of the relation C has certain advantages that will be apparent in forthcoming sections.

Definition 6. For finite non-empty $S_1, S_2 \subseteq L(T)$, we write $S_1|S_2$ if neither of $yca(S_1)$ and $yca(S_2)$ lies below the other. For sequences of (not necessarily distinct) vertices x_1, \ldots, x_n and y_1, \ldots, y_m we write $x_1, \ldots, x_n | y_1, \ldots, y_m$ if $\{x_1, \ldots, x_n\} | \{y_1, \ldots, y_m\}$.

In particular, xy|z (which is the notation that is typically used in the literature on phylogeny problems) is equivalent to C(z, xy); it will be very convenient to have both notations available. Note that if xy|z then this includes the possibility that x = y; however, xy|z implies that $x \neq z$ and $y \neq z$. Hence, for every triple x, y, z of leaves in a rooted binary tree, we either have xy|z, yz|x, xz|y, or x = y = z. Also note that $x_1, \ldots, x_n|y_1, \ldots, y_m$ if and only if $x_ix_j|y_k$ and $x_i|y_ky_l$ for all $i, j \leq n$ and $k, l \leq m$. The following result is known but we have been unable to find an explicit proof in the literature. Hence, we give a proof for the convenience of the reader.

Proposition 7. The class C of all finite leaf structures is an amalgamation class.

Proof. Arbitrarily choose $B_1, B_2 \in \mathbb{C}$ such that $A = B_1 \cap B_2$ is a substructure of both B_1 and B_2 . We inductively assume that the statement has been shown for all triples (A, B'_1, B'_2) where $D(B'_1) \cup D(B'_2)$ is a proper subset of $D(B_1) \cup D(B_2)$.

Let T_1 be the rooted binary tree underlying B_1 and T_2 the rooted binary tree underlying B_2 . Let $B_1^1 \in \mathbb{C}$ be the substructure of B_1 induced by the vertices below the left child of T_1 and $B_1^2 \in \mathbb{C}$ be the substructure of B_1 induced by the vertices below the right child of T_1 . The structures B_2^1 and B_2^2 are defined analogously for B_2 .

First consider the case when there is a vertex u that lies in both B_1^1 and B_2^1 and a vertex v that lies in both B_2^1 and B_1^2 . We claim that in this case no vertex w from B_2^2 can lie inside B_1 . Assume the contrary and note that w is either in B_1^1 , in which case we have uw|v in B_1 , or in B_1^2 , in which case we have u|vw in B_1 . But since u, v, w are in A, this contradicts the fact that uv|w holds in B_2 . Let $C' \in \mathbb{C}$ be the amalgam of B_1 and B_2^1 over A (which exists by the inductive assumption) and let T' be its underlying tree. Consider a tree T with root r, T' as its left subtree, and the underlying tree of B_2^2 as its right subtree. It is straightforward to verify that the leaf structure of T is in \mathbb{C} and that it is an amalgam of B_1 and B_2 over A.

The above argument can also be applied to the cases where the role of B_1 and B_2 , or the role of B_1^1 with B_1^2 , or the role of B_2^1 with B_2^2 are exchanged. Hence, the only remaining essentially different case we have to consider is when $D(B_1^1) \cup D(B_2^1)$ and $D(B_1^2) \cup D(B_2^2)$ are disjoint. In this case, it is straightforward to first amalgamate B_1^1 with B_2^1 and B_1^2 with B_2^2 to obtain the amalgam of B_1 and B_2 ; the details are left to the reader.

We write $(\mathbb{L}; C)$ for the Fraïssé-limit of \mathcal{C} . Obvious reducts of $(\mathbb{L}; C)$ are $(\mathbb{L}; C)$ itself and $(\mathbb{L}; =)$. To define a third reduct, consider the 4-ary relation Q(xy, uv) with the following first-order definition over $(\mathbb{L}; C)$:

$$(xy|u \wedge xy|v) \lor (x|uv \wedge y|uv)$$

This relation is often referred to as the *quartet* relation [44].

3.3 The structure $(\mathbf{L}; C)$: an axiomatic approach

The structure $(\mathbb{L}; C)$ that we defined in Section 3.2 is an important example of a so-called *C*-relation. This concept was introduced by Adeleke & Neumann [3] and we closely follow their definitions in the following. A ternary relation $C \subseteq X^3$ is said to be a C-relation on X if the following conditions hold:

C1.
$$\forall a, b, c (C(a, bc) \Rightarrow C(a, cb))$$

C2. $\forall a, b, c (C(a, bc) \Rightarrow \neg C(b, ac))$

- C3. $\forall a, b, c, d (C(a, bc) \Rightarrow C(a, dc) \lor C(d, bc))$
- C4. $\forall a, b \ (a \neq b \Rightarrow C(a, b, b))$

A C-relation is called *proper* if it satisfies two further properties:

- C5. $\forall a, b \exists c (C(c, ab))$
- C6. $\forall a, b \ (a \neq b \Rightarrow \exists c (c \neq b \land C(a, bc)))$

These six axioms do not describe the Fraïssé-limit $(\mathbb{L}; C)$ up to isomorphism. To completely axiomatize the theory of $(\mathbb{L}; C)$, we need two more axioms.

C7. $\forall a, b, c \left(C(c, ab) \Rightarrow \exists e \left(C(c, eb) \land C(e, ab) \right) \right)$

C8. $\forall a, b, c ((a \neq b \lor a \neq c \lor b \neq c) \Rightarrow (C(a, bc) \lor C(b, ac) \lor C(c, ab)))$

C-relations that satisfy C7 are called *dense* and C-relations that satisfy C8 are called *binary* branching. Note that C1-C8 are satisfiable since $(\mathbb{L}; C)$ is a countable model of C1-C8.

We mention that the structure $(\mathbb{L}; C)$ is existential positive complete within the class of all Crelations, as defined in [9]: for every homomorphism h of $(\mathbb{L}; C)$ into another C-relation and every existential positive formula $\phi(x_1, \ldots, x_n)$ and all $p_1, \ldots, p_n \in \mathbb{L}$ such that $\phi(h(p_1), \ldots, h(p_n))$ holds we have that $\phi(p_1, \ldots, p_n)$ holds in $(\mathbb{L}; C)$, too. It is also easy to see that every existential positive complete C-relation must satisfy C7 and C8. These facts about existential positive completeness of $(\mathbb{L}; C)$ are not needed in the remainder of the article, but together with Lemma 11 below they demonstrate that the structure $(\mathbb{L}; C)$ can be seen as the (up to isomorphism unique) generic countable C-relation.

The satisfiability of C1-C8 can also be shown using the idea of constructing C-relations in [5, page 123]. Let \mathcal{F} be the set of functions $f: (0, \infty) \to \{0, 1\}$, where $(0, \infty)$ denotes the set of positive rational numbers with the standard topology, such that the following conditions hold.

- there exists $a \in (0, \infty)$ such that f(x) = 0 for every $x \in (0, a)$.
- f has finitely many points of discontinuity and for each point b of discontinuity, there exists $\epsilon \in (0, b)$ such that $f(x) \neq f(b)$ for every $x \in (b \epsilon, b)$.

For every $f, g \in \mathcal{F}$ such that $f \neq g$, let $\operatorname{pref}(f, g)$ denote the interval (0, c) such that f(x) = g(x)for every $x \in (0, c)$, and $f(c) \neq g(c)$. If f = g, let $\operatorname{pref}(f, g) := (0, \infty)$. Note that c is a point of discontinuity of either f or g. We define a relation C on \mathcal{F} by C(f, gh) if $\operatorname{pref}(f, h) \subsetneq \operatorname{pref}(g, h)$. We can easily to verify that $(\mathcal{F}; C)$ is a countable model of C1-C8.

We will now prove (in Lemma 11) that there is a unique countable model of C1-C8 up to isomorphism. It suffices to show that if Γ is a countable structure with signature $\{C\}$ satisfying C1-C8, then Γ is isomorphic to ($\mathbb{L}; C$). To do so, we need a number of observations (Lemmas 8, 9, and 10).

The following consequences of C1-C8 are used in the proofs without further notice.

Lemma 8 (C-consequences). Let C denote a C-relation. Then

- 1. $\forall x, y, z, t ((C(x, yz) \land C(x, yt)) \Rightarrow C(x, zt)),$
- 2. $\forall x, y, z, t ((C(x, zt) \land C(z, xy)) \Rightarrow (C(t, xy) \land C(y, zt))), and$
- 3. $\forall x, y, z, t \left((C(z, xy) \land C(y, xt)) \Rightarrow (C(z, yt) \land C(z, xt)) \right).$

Proof. We prove the first consequence. The others can be shown analogously. Assume to the contrary that C(x, zt) does not hold. By applying C3 to x, y, z, t, we get that C(t, yz) and C2 implies that C(z, yt) does not hold. By applying C3 to x, y, t, z, it follows that C(x, zt) holds and we have a contradiction.

For two subsets Y, Z of X, we write C(Y, Z) if

- 1. $C(y, z_1 z_2)$ for arbitrary $y \in Y$ and $z_1, z_2 \in Z$, and
- 2. $C(z, y_1y_2)$ for arbitrary $y_1, y_2 \in Y$ and $z \in Z$.

Lemma 9. Let C be a ternary relation on a countably infinite set X that satisfies C1-C8. Then for every finite subset Y of X of size at least 2 there are two non-empty subsets A, B of Y such that $A \cup B = Y$ and C(A, B).

Proof. We prove the lemma by induction on |Y|. Clearly, the claim holds if |Y| = 2 so we assume that the lemma holds when |Y| = k - 1 for some k > 2. Henceforth, assume |Y| = k. Arbitrarily choose $y \in Y$ and let $Y' = Y \setminus \{y\}$. By the induction hypothesis, there are two non-empty subsets A', B' of Y' such that $A' \cup B' = Y'$ and C(A', B'). Pick $a' \in A'$ and $b' \in B'$. One of the following holds.

- C(y, a'b'). Arbitrarily choose $c, d \in Y'$. We show that C(y, cd) holds. If $c, d \in A'$, then we have C(y, a'b'), C(b', a'c), and C(b', a'd). It follows immediately from Lemma 8 that C(y, cd). Analogously, C(y, cd) holds if $c, d \in B'$. It remains to consider the case $c \in A', d \in B'$. Here, we have C(y, a'b'), C(b', a'c) and C(a', b'd). Once again, it follows from Lemma 8 that C(y, cd) holds. By setting $A = \{y\}$ and $B = A' \cup B'$, the lemma of the lemma follows.
- C(b', ya'). We first show that for arbitrary $a'' \in A'$ and $b'' \in B'$, we have C(b'', a''y). This follows from Lemma 8 and the fact that C(b', ya'), C(b', a'a''), and C(a', b'b'') hold. We can now show that for arbitrary $b'', b''' \in B'$, we have that C(y, b''b'') holds. This follows from Lemma 8 and the fact that C(b', a'y), C(a', b'b''), and C(a', b'b'') hold. This implies that $C(A' \cup \{y\}, B')$ and we have proved the induction step by setting $A = A' \cup \{y\}$ and B = B'.
- C(a', yb'). This case can be proved analogously to the previous case: we get that A = A'and $B = B' \cup \{y\}$.

The case distinction is exhaustive because of C8.

We would like to point out an important property of maps that preserve C.

Lemma 10. Let $e: X \to \mathbb{L}$ for $X \subseteq \mathbb{L}$ be a function that preserves C. Then e is injective and preserves the relation $\{t \in \mathbb{L}^3 : t \notin C\}$.

Proof. Clearly, e preserves the binary relation $\{(x, y) \in \mathbb{L}^2 : x \neq y\} = \{(x, y) : \exists z.C(x, y, z)\}$, and so e is injective. Arbitrarily choose $u_1, u_2, u_3 \in \mathbb{L}$ such that $u_1|u_2u_3$ does not hold. If $|\{u_1, u_2, u_3\}| = 1$ then $e(u_1)|e(u_2)e(u_3)$ does not hold and there is nothing to show. If $|\{u_1, u_2, u_3\}| = 2$ then by C4 either $u_1 = u_2$ or $u_1 = u_3$, and $e(u_1)|e(u_2)e(u_3)$ does not hold. If $|\{u_1, u_2, u_3\}| = 3$ then by C6 we have either $u_2|u_1u_3$, or $u_3|u_1u_2$. It follows that $e(u_2)|e(u_1)e(u_3)$ or $e(u_3)|e(u_1)e(u_2)$. In both cases, $e(u_1)|e(u_2)e(u_3)$ does not hold by C2.

We will typically use the contrapositive version of Lemma 10 in the sequel. This allows to draw the conclusion a|bc under the assumption e(a)|e(b)e(c).

Lemma 11. Let Γ be a countable structure with signature $\{C\}$ that satisfies C1-C8. Then Γ is isomorphic to $(\mathbb{L}; C)$.

Proof. It is straightforward (albeit a bit tedious) to show that $(\mathbb{L}; C)$ satisfies C1-C8. It then remains to show that if Γ_1 and Γ_2 are two countably infinite $\{C\}$ -structures that satisfy C1-C8, then the two structures are isomorphic. Let X_1, X_2 denote the domains of Γ_1 and Γ_2 , respectively. This can be shown by a back-and-forth argument based on the following claim.

Claim: Let A be a non-empty finite subset of X_1 and let f denote a map from A to X_2 that preserves C. Then for every $a \in X_1$, the map f can be extended to a map g from $A \cup \{a\}$ to X_2 that preserves C.

It follows from Lemma 10 that f also preserves $\{(x, y, z) : \neg C(x, yz)\}$. We prove the claim by induction on |A|. Clearly, we are done if $a \in A$ or |A| = 1. Hence, assume that $a \notin A$ and $|A| \ge 2$. Let A_1, A_2 be subsets of A such that $A_1 \cup A_2 = A$ and $C(A_1, A_2)$, which exist due to Lemma 9. Note that $C(f(A_1), f(A_2))$ holds in $(X_2; C)$. Pick $a_1 \in A_1$ and $a_2 \in A_2$. We construct the map g in each of the following three cases.

• $C(a, a_1a_2)$. We claim that $C(\{a\}, A)$ holds. Arbitrarily choose $u, v \in A$. Then either $C(ua_1, a_2)$ or $C(a_1, a_2u)$ by the choice of a_1 and a_2 . Similarly, either $C(va_1, a_2)$ or $C(a_1, a_2v)$. So there are four cases to consider; we only treat the case $C(ua_1, a_2)$ and $C(a_1, va_2)$ since the other cases are similar or easier. Now, $C(ua_1, a_2)$ and $C(a_1a_2, a)$ imply that $C(a, ua_1)$ by item 3 of Lemma 8. Similarly, we have $C(a, va_2)$. Now $C(a_1a_2, a)$ and two applications of item 1 of Lemma 8 give C(uv, a), which proves the subclaim.

It follows from C5 that there exists an $a' \in X_2$ such that $C(a', f(a_1)f(a_2))$. Once again, we obtain $C(\{a'\}, f(A))$ as a consequence of Lemma 8. This implies that the map $g: A \cup \{a\} \to X_2$, defined by $g|_A = f$ and g(a) = a', preserves C.

• $C(a_2, aa_1)$. It follows from Lemma 8 that $C(\{a\} \cup A_1, A_2)$ holds. We consider the following cases.

 $|A_1| = 1$. There exists $a' \in X_2$ such that $C(f(a_2), a'f(a_1))$ by C6, and Lemma 8 implies that $\overline{C(\{f(a_1), a'\}, f(A_2))}$ holds. Since we also have $C(\{a, a_1\}, A_2)$, the map $g: A \cup \{a\} \to X_2$ defined by $g|_A = f$ and g(a) = a', preserves C.

 $|A_2| = 1$. This case can be treated analogously to the previous case.

 $|A_1| \ge 2$ and $|A_2| \ge 2$. Let B_1, B_2 be non-empty such that $C(B_1, B_2)$ and $A_1 = B_1 \cup B_2$. Arbitrarily choose $b_1 \in B_1, b_2 \in B_2$. The following cases are exhaustive by C8.

- $C(a, b_1b_2)$. It is a direct consequence of C7 that there exists an $a' \in X_2$ such that both $C(\{f(a_2)\}, \{f(b_1), f(b_2), a'\})$ and $C(a', f(b_1)f(b_2))$ hold. Furthermore, Lemma 8 implies that $C(\{a\}, A_1), C(\{a\} \cup A_1, A_2), C(\{a'\}, f(A_1)), \text{ and } C(\{a'\} \cup f(A_1), f(A_2)).$ Hence, the map $g: A \cup \{a\} \to X_2$, defined by $g|_A = f$ and g(a) = a', preserves C.
- $C(b_2, ab_1)$. By assumption we know that $|A_2| \ge 2$, and since $A_1 \cup A_2 = A$ it follows that $|A_1 \cup \{a\}| < |A|$. Hence, by the induction hypothesis there exists a map $h: A_1 \cup \{a\} \to X_2$ such that $h|_{A_1} = f|_{A_1}$ and h preserves C on $A_1 \cup \{a\}$. Since h preserves C on $A_1 \cup \{a\}$, we see that $C(h(b_2), h(a)h(b_1))$, and consequently that $C(f(b_2), h(a)f(b_1))$ holds. Since both $C(A_1, A_2)$ and $C(f(A_1), f(A_2))$ hold, it follows from Lemma 8 that $C(\{a\} \cup A_1, A_2)$ and $C(\{h(a)\} \cup f(A_1), f(A_2))$ hold. This implies that the map $g: A \cup \{a\} \to X_2$, defined by $g|_{A_1 \cup \{a\}} = h, g|_{A_2} = f|_{A_2}$, preserves C.
- $C(b_1, ab_2)$. The proof is analogous to the case above.
- $C(a_1, aa_2)$. The proof is analogous to the case when $C(a_2, aa_1)$.

The case distinction is exhaustive because of C8.

3.4 The reduct $(\mathbb{L}; Q)$

The reduct $(\mathbb{L}; Q)$ of $(\mathbb{L}; C)$ can be treated axiomatically, too. A 4-ary relation D is said to be a *D*-relation on X if the following conditions hold:

- D1. $\forall a, b, c, d \left(D(ab, cd) \Rightarrow D(ba, cd) \land D(ab, dc) \land D(cd, ab) \right)$
- D2. $\forall a, b, c, d (D(ab, cd) \Rightarrow \neg D(ac, bd))$
- D3. $\forall a, b, c, d, e \left(D(ab, cd) \Rightarrow D(eb, cd) \lor D(ab, ce) \right)$

D4. $\forall a, b, c ((a \neq c \land b \neq c) \Rightarrow D(ab, cc))$

A D-relation is called *proper* if it additionally satisfies the following condition:

D5. For pairwise distinct a, b, c there is $d \in X \setminus \{a, b, c\}$ with D(ab, cd).

As with $(\mathbb{L}; C)$, it is possible to axiomatize the theory of $(\mathbb{L}; Q)$ by adding finitely many axioms.

D6. $\forall a, b, c, d (D(ab, cd) \Rightarrow \exists e (D(eb, cd) \land D(ae, cd) \land D(ab, ed) \land D(ab, ce)))$

D7. $\forall a, b, c, d (|\{a, b, c, d\}| \ge 3 \Rightarrow (D(ab, cd) \lor D(ac, bd) \lor D(ad, bc)))$

D-relations satisfying D6 are called *dense*, and D-relations satisfying D7 are called *binary* branching.

We will continue by proving that if two countable structures with signature $\{D\}$ satisfy D1-D7, then they are isomorphic. For increased readability, we write D(xyz, uv) when $D(xy, uv) \land D(xz, uv) \land D(yz, uv)$, and we write D(xy, zuv) when $D(xy, zu) \land D(xy, zv) \land D(xy, uv)$. One may note, for instance, that D(xy, zuv) is equivalent to D(xy, uvz).

Lemma 12 (D-consequences). If D is a D-relation, then

- $\forall x, y, z, u, v((D(xy, zu) \land D(xy, zv)) \Rightarrow D(xy, uv)), and$
- $\forall x, y, z, u, v (D(xy, zu) \Rightarrow (D(xyv, zu) \lor D(xy, zuv))).$

Proof. We prove the first item. By applying D1 and D3 to $D(xy, zu) \wedge D(xy, zv)$, we get that

 $(D(yv, uz) \lor D(xy, uv)) \land (D(yu, vz) \lor D(xy, uv)).$

If D(xy, uv) does not hold, then $D(yv, uz) \wedge D(yu, vz)$ must hold. However, this immediately leads to a contradiction via D2: $D(yu, vz) \Rightarrow \neg D(yv, uz)$.

To prove the second item, assume that D(xy, zu) holds and arbitrarily choose v. By D3, we have $D(vy, zu) \lor D(xy, zv)$. Assume that D(vy, zu) holds; the other case can be proved in a similar way. By definition D(xyv, zu) if and only if $D(xy, zu) \land D(xv, zu) \land D(yv, zu)$. We know that D(xy, zu) holds and that D(vy, zu) implies D(yv, zu) via D1. It remains to show that D(xv, zu) holds, too. By once again applying D1, we see that $D(zu, yx) \land D(zu, yv)$. It follows that D(zu, xv) holds by the claim above and we conclude that D(xv, zu) holds by D1.

Lemma 13. Let $e: X \to \mathbb{L}$ for $X \subseteq \mathbb{L}$ be a function that preserves Q. Then e is injective and preserves the relation $\{q \in \mathbb{L}^4 : q \notin Q\}$.

Proof. The proof is very similar to the proof of Lemma 10 and is left to the reader.

We will typically use the contrapositive version of Lemma 13 in the sequel. This allows to draw the conclusion Q(ab, cd) under the assumption Q(e(a)e(b), e(c)e(d)).

Lemma 14. Let D be a 4-ary relation on a countably infinite set X that satisfies D1-D7. Then (X; D) is isomorphic to $(\mathbb{L}; Q)$, and homogeneous.

The proof of Lemma 14 is based on Lemma 11 and the idea of rerooting at a fixed leaf to create a C-relation from a D-relation. The idea of rerooting was already discussed in [25].

Proof. It is straightforward to verify that $(\mathbb{L}; Q)$ satisfies D1-D7. Let $(X_1; D)$ and $(X_2; D)$ be two countably infinite sets that satisfy D1-D7, let Y_1 be a finite subset of X_1 , and α an embedding of the structure induced by Y_1 in (X; D) into (X_2, D) . We will show that α can be extended to an isomorphism between $(X_1; D)$ and (X_2, D) . This can be applied to $(X_1; D) = (X_2; D) = (\mathbb{L}; Q)$ and hence also shows homogeneity of $(\mathbb{L}; Q)$.

Arbitrarily choose $c \in Y_1$. We define a relation C on $X'_1 := X_1 \setminus \{c\}$ as follows: for every $(x, y, z) \in (X'_1)^3$, let $(x, y, z) \in C$ if and only if D(cx, yz) holds. Similarly, we define a relation C on $X'_2 := X_2 \setminus \{\alpha(c)\}$ as follows: for every $(x, y, z) \in (X'_1)^3$, let $(x, y, z) \in C$ if and only if $D(\alpha(c)x, yz)$ holds.

One can verify that both (X'_1, C) and (X'_2, C) satisfies C1-C8. It follows from Lemma 11 that $(X'_1; C)$ and $(X'_2; C)$ are isomorphic to $(\mathbb{L}; C)$, and it follows from homogeneity of $(\mathbb{L}; C)$ that the restriction of α to $Y_1 \setminus \{c\}$ can be extended to an isomorphism α' between $(X'_1; C)$ and $(X'_2; C)$.

We conclude the proof by showing that the map $\beta: X_1 \to X_2$, defined by $\beta(c) := \alpha(c)$ and $\beta|_{X'_1} = \alpha'$, is an isomorphism between $(X_1; D)$ and $(X_2; D)$. Arbitrarily choose $x, y, u, v \in X_1$ satisfying D(xy, uv). By Lemma 13 it is sufficient to show that $D(\beta(x)\beta(y), \beta(u)\beta(v))$. Clearly, we are done if x, y, u, v are not pairwise distinct, or if $c \in \{x, y, u, v\}$, so assume otherwise. By Lemma 12 we have D(xyc, uv) or D(xy, cuv). In the former case, it follows from the definition of C on X'_1 and X'_2 that $D(\alpha(x)\alpha(c), \alpha(u)\alpha(v))$ and $D(\alpha(y)\alpha(c), \alpha(u)\alpha(v))$. Lemma 12 implies that $D(\alpha(x)\alpha(y), \alpha(u)\alpha(v))$, which is equivalent to $D(\beta(x)\beta(y), \beta(u)\beta(v))$. The case that D(xy, cuv) can be shown analogously to the previous case.

Corollary 15. There exists an operation $rer \in Aut(\mathbb{L}; Q)$ that violates C.

Proof. It follows from Lemma 14 that $\operatorname{Aut}(\mathbb{L}; Q)$ is 3-transitive. Since $\operatorname{Aut}(\mathbb{L}; C)$ is 2-transitive, but not 3-transitive, it follows that $\operatorname{Aut}(\mathbb{L}; C) \neq \operatorname{Aut}(\mathbb{L}; Q)$. Since $\operatorname{Aut}(\mathbb{L}; C) \subseteq \operatorname{Aut}(\mathbb{L}; Q)$, there is rer $\in \operatorname{Aut}(\mathbb{L}; Q)$ which violates C.

The name rer may seem puzzling at first sight: it is short-hand for *rerooting*. The choice of terminology will be clarified in the next section.

3.5 Convex orderings of C-relations

In the proof of our main result, it will be useful to work with an expansion $(L; C, \prec)$ of (L; C)by a certain linear order \prec on \mathbb{L} . We will next describe how this linear order is defined as a Fraïssé-limit. A linear order \prec on the elements of a leaf structure (L; C) is called *convex* if for all $x, y, z \in L$ with $x \prec y \prec z$ we have that either xy|z or that x|yz (but not xz|y). The concept of convex linear order was already discussed in [25] and in [32, page 162].

Proposition 16. Let (L(T); C) be the leaf structure of a finite binary rooted tree T and arbitrarily choose $a \in L(T)$. Then there exists a convex linear order \prec of L(T) whose maximal element is a. In particular, every leaf structure can be expanded to a convexly ordered leaf structure.

Proof. Perform a depth-first search of T, starting at the root, such that vertices that lie above a in T are explored latest possible during the search. Let \prec be the order on L(T) in which the vertices have been visited during the search. Clearly, \prec is convex and a is its largest element. \Box

Proposition 17. The class C' of all finite convexly ordered leaf structures is an amalgamation class and its Fraissé-limit is isomorphic to an expansion $(L; C, \prec)$ of (L; C) by a convex linear ordering \prec . The structure $(\mathbb{L}; C, \prec)$ is described uniquely up to isomorphism by the axioms C1-C8 and by the fact that \prec is a dense and unbounded linear order which is convex with respect to $(\mathbb{L}; C)$.

Proof. The proof that \mathcal{C}' is an amalgamation class is similar to the proof of Proposition 7. The Fraïssé-limit of \mathcal{C}' clearly satisfies C1-C8, it is equipped with a convex linear order, and all countable structures with these properties are in fact isomorphic; this can be shown by a back-and-forth argument. By Lemma 11, the structure obtained by forgetting the order is isomorphic to $(\mathbb{L}; C)$ and the statement follows.

By the classical result of Cantor [27], all countable dense unbounded linear orders are isomorphic to $(\mathbb{Q}; <)$, and hence Proposition 17 implies that $(\mathbb{L}; \prec)$ is isomorphic to $(\mathbb{Q}; <)$.

4 Automorphism groups of reducts

We will now show that the structure $(\mathbb{L}; C)$ has precisely three reducts up to first-order interdefinability. Our proof uses a result by Adeleke and Neumann [2] about primitive permutation groups with *primitive Jordan sets*. The link between reducts of $(\mathbb{L}; C)$ and permutation groups is given



Figure 1: Illustration of the 3 orbits of triples (a, b, c) with pairwise distinct entries of Aut(L; C).

by the theorem of Engeler, Ryll-Nardzewski, and Svenonius, which we briefly recall in Section 4.1. We continue in Section 4.2 by presenting some important lemmata about functions that preserve Q but violate C. With these results in place, we finally prove the main result of this section in Section 4.3.

4.1 Permutation group preliminaries

Our proof will utilize links between homogeneity, ω -categoricity, and permutation groups so we begin by discussing these central concepts. A structure Γ is ω -categorical if all countable structures that satisfy the same first-order sentences as Γ are isomorphic (see e.g. Cameron [26] or Hodges [33]). Homogeneous structures with finite relational signatures are ω -categorical, so the structure ($\mathbb{L}; C$) is ω -categorical. Moreover, all structures with a first-order definition in an ω categorical structure are ω -categorical (see again Hodges [33]). This implies, for instance, that ($\mathbb{L}; Q$) is ω -categorical.

The fundamental theorem by Engeler, Ryll-Nardzewski, and Svenonius is a characterization of ω -categoricity in terms of permutation groups. When G is a permutation group on a set X, then the *orbit* of a k-tuple t is the set $\{\alpha(t) \mid \alpha \in G\}$. We see that homogeneity of $(\mathbb{L}; C)$ implies that $\operatorname{Aut}(\mathbb{L}; C)$ has precisely three orbits of triples with pairwise distinct entries; an illustration of these orbits can be found in Figure 1. We now state the Engeler-Ryll-Nardzewski-Svenonius theorem and its proof can be found in, for instance, Hodges [33].

Theorem 18. A countable relational structure Γ is ω -categorical if and only if the automorphism group of Γ is oligomorphic, that is, if for each $k \geq 1$ there are finitely many orbits of k-tuples under Aut(Γ). A relation R has a first-order definition in an ω -categorical structure Γ if and only if R is preserved by all automorphisms of Γ .

This theorem implies that a structure $(\mathbb{L}; R_1, R_2, ...)$ is first-order definable in $(\mathbb{L}; C)$ if and only if its automorphism group contains the automorphisms of (L; C).

Automorphism groups G of relational structures carry a natural topology, namely the topology of *pointwise convergence*. Whenever we refer to topological properties of groups it will be with respect to this topology. To define this topology, we begin by giving the domain X of the relational structure the discrete topology. We then view G as a subspace of the Baire space X^X which carries the product topology; see e.g. Cameron [26]. A set of permutations is called *closed* if it is closed in the subspace Sym(X) of X^X , where Sym(X) is the set of all bijections from X to X. The *closure* of a set of permutations P is the smallest closed set of permutations that contains P and it will be denoted by \overline{P} . Note that \overline{P} equals the set of all permutations f such that for every finite subset A of the domain there is a $g \in P$ such that f(a) = g(a) for all $a \in A$.

We write $\langle P \rangle$ for the smallest permutation group that contains a given set of permutations P. Note that the smallest closed permutation group that contains a set of permutations P equals $\overline{\langle P \rangle}$. It is easy to see that a set of permutations G on a set X is a closed subgroup of the group of all permutations of X if and only if G is the automorphism group of a relational structure [26].

We need some terminology from permutation group theory and we mostly follow Bhattacharjee, Macpherson, Möller and Neumann [5]. A permutation group G on a set X is called

• *k*-transitive if for any two sequences a_1, \ldots, a_k and b_1, \ldots, b_k of k distinct points of X there exists g in G such that $g(a_i) = b_i$ for all $1 \le i \le k$,

- *transitive* if G is 1-transitive,
- *highly transitive* if it is *k*-transitive for all natural numbers *k*,
- primitive if it is transitive and all equivalence relations that are preserved by all operations in G are either the equivalence relation with one equivalence class or the equivalence relation with equivalence classes of size one.

The following simple fact illustrates the link between model theoretic and permutation group theoretic concepts.

Proposition 19. For an automorphism group G of a relational structure Γ with domain D, the following are equivalent.

- G is highly transitive.
- G equals the set of all permutations of D.
- Γ is a reduct of (D; =).

The pointwise stabilizer at $Y \subset X$ of a permutation group G on X is the permutation group on X consisting of all permutations $\alpha \in G$ such that $\alpha(y) = y$ for all $y \in Y$. A subset X' of X is said to be a *Jordan set* (for G in X) if |X'| > 1 and the pointwise stabilizer H of G at $X \setminus X'$ is transitive on X'.

If the group G is (k + 1)-transitive and X' is any co-finite subset with $|X \setminus X'| = k$, then X' is automatically a Jordan set. Such Jordan sets will be said to be *improper* while all other will be called *proper*. We say that the Jordan set X' is k-transitive if the pointwise stabilizer H is k-transitive on X'. The permutation group G on the set X is said to be a Jordan group if G is transitive on X and there exists a proper Jordan set for G in X. The main result that we will use in Section 4.3 is the following.

Theorem 20 (Note 7.1 in Adeleke and Neumann [2]). If G is primitive and has 2-transitive proper Jordan sets, then G is either highly transitive or it preserves a C- or D-relation on X.

Note that $\operatorname{Aut}(\mathbb{L}; C)$ is 2-transitive by homogeneity and that 2-transitivity implies primitivity. The following proposition shows that Theorem 20 applies in our setting.

Proposition 21. For two arbitrary distinct elements $a, b \in \mathbb{L}$, the set $S := \{x \in \mathbb{L} : ax | b\}$ is a 2-transitive proper primitive Jordan set of Aut(L; C).

Proof. The pointwise stabilizer of Aut(L; C) at $L \setminus S$ acts 2-transitively on S; this can be shown via a simple back-and-forth argument.

4.2 The rerooting lemma

We will now prove some fundamental lemmata concerning functions that preserve Q. They will be needed to prove Theorem 28 which is the main result of Section 4. They will also be used in subsequent sections: we emphasize that these results are not restricted to permutations. The most important lemma is the *rerooting lemma* (Lemma 26) about functions that preserve Q and violate C. The following notation will be convenient in the following.

Definition 22. We write $x_1 \ldots x_n : y_1 \ldots y_m$ if $Q(x_i x_j, y_k y_l)$ for all $i, j \leq n$ and $k, l \leq m$.

Lemma 23. Let $A_1, A_2 \subseteq \mathbb{L}$ be such that $A_1|A_2$ and let $f: A_1 \cup A_2 \to \mathbb{L}$ be a function that preserves Q and satisfies $f(A_1)|f(A_2)$. Then f also preserves C.

Proof. Since $A_1|A_2$, we have $A_1 \cup A_2 \ge 2$. Clearly, the claim of lemma holds if $|A_1 \cup A_2| = 2$. It remains to consider the case $|A_1 \cup A_2| \ge 3$. Let $a_1, a_2, a_3 \in A_1 \cup A_2$ be three distinct elements such that $a_1a_2|a_3$. We have to verify that $f(a_1)f(a_2)|f(a_3)$ and we do this by considering four different cases.

- $a_1, a_2 \in A_1$ and $a_3 \in A_2$. In this case, since $f(A_1)|f(A_2)$, we have in particular that $f(a_1)f(a_2)|f(a_3)$.
- $a_1, a_2 \in A_2$ and $a_3 \in A_1$. Analogous to the previous case.
- $a_1, a_2, a_3 \in A_1$. Let $b \in A_2$. Clearly $a_1a_2 : a_3b$, and $f(a_1)f(a_2) : f(a_3)f(b)$ since f preserves Q. Moreover, we have $f(a_1)f(a_2)f(a_3)|f(b)$, and thus $f(a_1)f(a_2)|f(a_3)$.
- $a_1, a_2, a_3 \in A_2$. Analogous to the previous case.

Since we have assumed that $A_1|A_2$, these cases are in fact exhaustive. One may, for instance, note that if $a_1, a_3 \in A_1$ and $a_2 \in A_2$, then $a_1a_3|a_2$ which immediately contradicts that $a_1a_2|a_3$.

Lemma 24. Let $A \subset \mathbb{L}$ be finite of size at least two and let $f: A \to \mathbb{L}$ be a function which preserves Q. Then there exists a non-empty $B \subsetneq A$ such that the following conditions hold:

- $f(B)|f(A \setminus B)$
- B|x for all $x \in A \setminus B$.

Proof. Let B_1, B_2 be non-empty such that $B_1 \cup B_2 = A$ and $f(B_1)|f(B_2)$. We see that B_1, B_2 is a partitioning of A since $f(B_1)|f(B_2)$ implies $B_1 \cap B_2 = \emptyset$. If $B_1|x$ for all $x \in B_2$, then we can choose $B = B_1$ and we are done. Otherwise there are $u, v \in B_1$ and $w \in B_2$ such that u|vw. We claim that in this case $x|B_2$ for all $x \in B_1$. Since f preserves Q on A and f(u)f(v) : f(w)f(x) holds for every $x \in B_2$, we have uv : wx by Lemma 13. Therefore u|wx and v|wx hold. This implies that $u|B_2$ holds. Let w', w'' be two arbitrary elements from B_2 and u' an arbitrary element from B_1 . We thus have f(w')f(w'') : f(u')f(u) and, once again by Lemma 13, we have uu' : w'w''. This implies u|w'w'' and consequently u'|w'w''. Hence, $u'|B_2$ for arbitrary $u' \in B_2$.

We will now introduce the idea of *c*-universality. This seemingly simple concept is highly important throughout the article and it will be encountered in several different contexts.

Definition 25. Arbitrarily choose $c \in \mathbb{L}$. A set $A \subseteq \mathbb{L} \setminus \{c\}$ is called c-universal if for every finite $U \subset \mathbb{L}$ and for every $u \in U$, there exists an $\alpha \in \operatorname{Aut}(\mathbb{L}; C)$ such that $\alpha(u) = c$ and $\alpha(U) \subseteq A \cup \{c\}$.

We continue by presenting the rerooting lemma which identifies permutations g of \mathbb{L} that preserve Q and can be used for generating all automorphisms of $(\mathbb{L}; Q)$ when combined with $\operatorname{Aut}(\mathbb{L}; C)$. The idea is based on the following observation: the finite substructures of $(\mathbb{L}; C)$ provide information about the root of the underlying tree whereas the finite substructures of $(\mathbb{L}; Q)$ only provide information about the underlying unrooted trees. Intuitively, we use the function g to change the position of the root in order to generate all automorphisms of $(\mathbb{L}; Q)$.

Lemma 26 (Rerooting Lemma). Arbitrarily choose $c \in \mathbb{L}$ and assume that $A \subseteq \mathbb{L} \setminus \{c\}$ is *c*-universal. If *g* is a permutation of \mathbb{L} that preserves *Q* on $A \cup \{c\}$ and satisfies g(A)|g(c), then

$$\operatorname{Aut}(\mathbb{L}; Q) \subseteq \overline{\langle \operatorname{Aut}(\mathbb{L}; C) \cup \{g\} \rangle}.$$

Proof. Arbitrarily choose $f \in \operatorname{Aut}(\mathbb{L}; Q)$ and let X be an arbitrary finite subset of \mathbb{L} . We have to show that $\langle \operatorname{Aut}(\mathbb{L}; C) \cup \{g\} \rangle$ contains an operation e such that e(x) = f(x) for all $x \in X$. This is trivial when |X| = 1 so we assume that $|X| \ge 2$. By Lemma 24, there exists a non-empty proper subset Y of X such that $f(Y)|f(X \setminus Y)$ and Y|x for all $x \in X \setminus Y$. By the homogeneity of $(\mathbb{L}; C)$, we can choose an element $c' \in \mathbb{L} \setminus X$ such that c'|Y and $(Y \cup \{c'\})|x$ for all $x \in X \setminus Y$. By c-universality, there exists an $\alpha \in \operatorname{Aut}(\mathbb{L}; C)$ such that $\alpha(X \cup \{c'\}) \subseteq A \cup \{c\}$ and $\alpha(c') = c$. Let $h := g \circ \alpha$. Note that h preserves Q on X and that h is a permutation. We continue by proving a particular property of h.

Claim. $h(Y)|h(X \setminus Y)$.

To prove this, we first show that $h(y_1)h(y_2)|h(y_3)$ for every $y_1, y_2 \in Y$ and $y_3 \in X \setminus Y$. We have $y_1y_2 : y_3c'$ by the choice of c' and this implies that $h(y_1)h(y_2) : h(y_3)h(c')$. Since $\alpha(X) \subseteq A$, it

follows from the definition of h that $h(y_1), h(y_2), h(y_3) \in g(A)$. Since g(c)|g(A) and $\alpha(y_i) \in A$ for every $i \in \{1, 2, 3\}$, we have $g(c)|h(y_1)h(y_2)h(y_3)$. Since h(c') = g(c) and $h(y_1)h(y_2) : h(y_3)h(c')$, it follows that $h(y_1)h(y_2)|h(y_3)$. In the same vein, we show that $h(y_1)|h(y_2)h(y_3)$ for every $y_1 \in Y$ and $y_2, y_3 \in X \setminus Y$. In this case, we have $y_1c' : y_2y_3$ by the choice of c' and this implies $h(y_1)h(c') :$ $h(y_2)h(y_3)$. Since h(c') = g(c) and $g(c)|h(y_1)h(y_2)h(y_3)$, we see that $h(y_1)|h(y_2)h(y_3)$.

Let $\beta: h(X) \to f(X)$ be defined by $\beta(x) = f(h^{-1}(x))$. Note that h^{-1} is well-defined since h is an injective function. Since both h and f preserve Q, we have that β preserve Q by Lemma 13.

Note that $\beta(h(Y))|\beta(h(X \setminus Y))$ since $\beta(h(x)) = f(x)$ and we have assumed that $f(Y)|f(X \setminus Y)$. Hence, the conditions of Lemma 23 apply to β for $A_1 := h(Y)$ and $A_2 := h(X \setminus Y)$ if we use the claim above. It follows that β preserves C. By the homogeneity of $(\mathbb{L}; C)$, there exists an $\gamma \in \operatorname{Aut}(\mathbb{L}; C)$ that extends β . Then $e := \gamma \circ h$ has the desired property. \Box

Observe the following important consequence of Lemma 26.

Corollary 27. Assume $f \in Aut(\mathbb{L}; Q)$ violates C. Then

$$\overline{\langle \operatorname{Aut}(\mathbb{L}; C) \cup \{f\} \rangle} = \operatorname{Aut}(\mathbb{L}; Q) \,.$$

Proof. The relation Q is first-order definable over $(\mathbb{L}; C)$ so $\operatorname{Aut}(\mathbb{L}; C) \subseteq \operatorname{Aut}(\mathbb{L}; Q)$. Furthermore, f preserves Q and it follows that

$$\overline{\langle \operatorname{Aut}(\mathbb{L}; C) \cup \{f\} \rangle} \subseteq \operatorname{Aut}(\mathbb{L}; Q)$$
.

For the converse, choose $f \in \operatorname{Aut}(\mathbb{L}; Q)$ such that there are $a_1, a_2, a_3 \in \mathbb{L}$ with $a_1|a_2a_3$ and $f(a_1)f(a_2)|f(a_3)$. Let $A = \{x \mid xa_1 : a_2a_3\}$. We will show that $f(A)|f(a_3)$. Let $x, y \in A$ be arbitrary. Since f preserves Q, we have $f(x)f(a_1) : f(a_2)f(a_3)$ and $f(y)f(a_1) : f(a_2)f(a_3)$. It follows from the condition $f(a_1)f(a_2)|f(a_3)$ that

$$f(x)f(a_1)|f(a_2) \wedge f(x)f(a_1)|f(a_3) \wedge f(y)f(a_1)|f(a_2) \wedge f(y)f(a_1)|f(a_3).$$

Since $f(x)f(a_1)|f(a_3) \wedge f(y)f(a_1)|f(a_3)$, we have $f(x)f(y)|f(a_3)$. Thus $f(A)|f(a_3)$. Clearly, A is a_3 -universal. Applying Lemma 26 to $c = a_3$ we have

$$\overline{\operatorname{Aut}(\mathbb{L};Q)\subseteq \langle \operatorname{Aut}(\mathbb{L};C)\cup \{f\}\rangle}$$

4.3 Automorphism group classification

We are now ready to prove the main result concerning automorphism groups of the reducts of $(\mathbb{L}; C)$.

Theorem 28. Let G be a closed permutation group on the set \mathbb{L} that contains $\operatorname{Aut}(\mathbb{L}; C)$. Then G is either $\operatorname{Aut}(\mathbb{L}; C)$, $\operatorname{Aut}(\mathbb{L}; Q)$, or $\operatorname{Aut}(\mathbb{L}; =)$.

Proof. Because G satisfies the conditions of Theorem 20, it is either highly transitive or it preserves a C- or D-relation. If G is highly transitive, then G equals $\operatorname{Aut}(\mathbb{L};=)$ by Proposition 19. Assume instead that G preserves a C-relation C'. We begin by making an observation.

Claim 0. All tuples $(o, p, q) \in C'$ with pairwise distinct entries satisfy o|pq.

Suppose for contradiction that p|oq. Then, (o, p, q) is in the same orbit as (q, p, o) in Aut $(\mathbb{L}; C)$ and therefore also in G. Since C' is preserved by G, we have C'(q, p, o) which contradicts C2. Similarly, it is impossible that q|op. Thus, the only remaining possibility is o|pq since C satisfies C8.

Arbitrarily choose $a, b, c \in \mathbb{L}$ such that a|bc and some $\alpha \in G$. If $|\{a, b, c\}| = 2$, then (by 2-transitivity of Aut($\mathbb{L}; C$)) we have that $(\alpha(a), \alpha(b), \alpha(c))$ is in the same orbit as (a, b, c) of Aut($\mathbb{L}; C$). Consequently, $(\alpha(a), \alpha(b), \alpha(c)) \in C$. Suppose instead that $|\{a, b, c\}| = 3$. Observe

that C' contains a triple with pairwise distinct entries. Arbitrarily choose two distinct elements $u, v \in \mathbb{L}$. Axiom C6 implies that there exists a $w \in \mathbb{L}$ such that C'(u, vw) and $w \neq v$. In fact, we also have $w \neq u$ since otherwise C'(u, vu) which is impossible due to C2 and C4. In particular, it follows that u|vw and therefore (u, v, w) is in the same orbit as (a, b, c) in $\operatorname{Aut}(\mathbb{L}; C)$. It follows that $(a, b, c) \in C'$. Since G preserves C' we have $C'(\alpha(a), \alpha(b)\alpha(c))$. By Claim 0, $\alpha(a)|\alpha(b)\alpha(c)$. We conclude that α preserves C and that $G = \operatorname{Aut}(\mathbb{L}; C)$.

Finally, we consider the case when G preserves a D-relation D. We begin by making three intermediate observations.

Claim 1. Every tuple $(a, b, c, d) \in D$ with pairwise distinct entries satisfies ab : cd.

Suppose for contradiction that ac: bd. Then either $ac|b \wedge ac|d$ or $a|bd \wedge c|bd$ by the definition of relation Q. In the first case, (a, b, c, d) is in the same orbit as (c, b, a, d) in $Aut(\mathbb{L}; C)$ so $(c, b, a, d) \in D$. Axiom D1 implies that $(a, d, b, c) \in D$ and this contradicts D2. If $a|bd \wedge c|bd$, then we can obtain a contradiction in a similar way. Finally, the case when ad: bc can be treated analogously. It follows that ab: cd since Q satisfies D7.

Claim 2. D contains a tuple (o, p, q, r) with pairwise distinct entries such that op|qr holds. Let $u, v, w \in \mathbb{L}$ be three distinct elements such that uv|w. There is an $x \in \mathbb{L} \setminus \{u, v, w\}$ such

that D(uv, wx) by D5. Claim 1 immediately implies that uv : wx. We consider the following cases.

- uv|wx. There is nothing to prove in this case.
- uvw|x. Choose $y \in \mathbb{L}$ be such that $y \neq w$ and uv|yw. It follows from D3 that D(uv, yw) or D(yv, wx). The second case is impossible since yv : wx does not hold. We see that $(u, v, y, w) \in D$ and we are done.
- uv|x and uvx|w. One may argue similarly as in the previous case by choosing $y \in \mathbb{L} \setminus \{u, v, w, x\}$ such that uv|yx and observe that $(u, v, x, w) \in D$ by D1.

Claim 3. D contains a tuple (a, b, c, d) with pairwise distinct entries such that $ab|c \wedge abc|d$. It follows from Claim 2 that there exists a tuple (o, p, q, r) with pairwise distinct entries such that op|qr holds. Choose $s \in \mathbb{L}$ such that opqr|s holds. Axiom D3 implies that D(sp, qr) or D(op, qs). We are done if the second case holds. If the first case holds, then we have D(qr, ps) by D1 and we are once again done.

Now, we show that every $f \in G$ preserves Q. Arbitrarily choose $a_1, a_2, a_3, a_4 \in \mathbb{L}$ such that $a_1a_2: a_3a_4$. We show that $(a_1, a_2, a_3, a_4) \in D$ (and, consequently, that $(f(a_1), f(a_2), f(a_3), f(a_4)) \in D$) by an exhaustive case analysis. Claim 2 implies that D contains a tuple (o, p, q, r) with pairwise distinct entries and op|qr. Consequently, D contains all tuples in the same orbit as (o, p, q, r) in Aut $(\mathbb{L}; C)$.

If a_1, a_2, a_3, a_4 are pairwise distinct and satisfy $a_1a_2|a_3a_4$, then $(a_1, a_2, a_3, a_4) \in D$ by Claim 1. Similarly, if a_1, a_2, a_3, a_4 are pairwise distinct and satisfy $a_1a_2|a_3$ and $a_1a_2a_3|a_4$, then Claim 3 implies that $(a_1, a_2, a_3, a_4) \in D$. If $a_2|a_3a_4$ and $a_1|a_2a_3a_4$, then $(a_1, a_2, a_3, a_4) \in D$ by D1. If $a_1a_2a_4|a_3$ and $a_1a_2|a_4$, or if $a_1|a_3a_4$ and $a_2|a_1a_3a_4$, then $(a_1, a_2, a_3, a_4) \in D$ by D1. If $a_3 = a_4$, $a_1 \neq a_3$, $a_2 \neq a_3$, then $(a_1, a_2, a_3, a_4) \in D$ by D4. The only remaining possibility to satisfy $a_1a_2: a_3a_4$ is that $a_1 = a_2, a_3 \neq a_1, a_4 \neq a_1$. In this case, $(a_1, a_2, a_3, a_4) \in D$ by D4 and D1. Hence, in all cases we have $(a_1, a_2, a_3, a_4) \in D$ and, consequently, $(f(a_1)f(a_2), f(a_3)f(a_4)) \in D$.

We can now conclude this part of the proof. If $f(a_1), f(a_2), f(a_3), f(a_4)$ are pairwise distinct, then $f(a_1)f(a_2): f(a_3)f(a_4)$ by Claim 1. Otherwise, one of the following cases hold:

- $f(a_1) = f(a_2), f(a_3) \neq f(a_1), \text{ and } f(a_4) \neq f(a_1),$
- $f(a_3) = f(a_4), f(a_1) \neq f(a_3)$, and $f(a_1) \neq f(a_4)$, or
- $f(a_1) = f(a_2)$ and $f(a_3) = f(a_4)$.

In all three cases, we have that $f(a_1)f(a_2) : f(a_3)f(a_4)$ and $G \subseteq \operatorname{Aut}(\mathbb{L}; Q)$. If $G = \operatorname{Aut}(\mathbb{L}; C)$, then we are done. Otherwise, pick one $f \in G \setminus \operatorname{Aut}(\mathbb{L}; C)$. Corollary 27 asserts that

$$\overline{\langle \operatorname{Aut}(\mathbb{L}; C) \cup \{f\} \rangle} = \operatorname{Aut}(\mathbb{L}; Q) \subseteq G$$

and it follows that $G = \operatorname{Aut}(\mathbb{L}; Q)$.

The following is an immediate consequence of Theorem 28 in combination with Theorem 18.

Corollary 29. Let Γ be a reduct of $(\mathbb{L}; C)$. Then Γ is first-order interdefinable with $(\mathbb{L}; C)$, $(\mathbb{L}; Q)$, or $(\mathbb{L}; =)$.

Proof. Since Γ is a reduct of $(\mathbb{L}; C)$, $\operatorname{Aut}(\Gamma)$ is a closed group that contains $\operatorname{Aut}(\mathbb{L}; C)$ and therefore equals $\operatorname{Aut}(\mathbb{L}; C)$, $\operatorname{Aut}(\mathbb{L}; Q)$, or $\operatorname{Aut}(\mathbb{L}; =)$ by Theorem 28. Theorem 18 implies that Γ is first-order interdefinable with $(\mathbb{L}; C)$, with $(\mathbb{L}; Q)$, or with $(\mathbb{L}; =)$.

Corollary 29 will be refined to a classification up to existential interdefinability in the forthcoming sections.

5 Ramsey theory for the C-relation

To analyze endomorphism monoids of reducts of $(\mathbb{L}; C)$, we apply Ramsey theory; a survey on this technique can be found in Bodirsky & Pinsker [16]. The basics of the Ramsey approach are presented in Section 5.1 and we introduce the important concepts of *canonicity* and the *ordering property* in Sections 5.2 and 5.3, respectively. We would like to mention that none of the results from the previous sections that use the theory of Jordan permutation groups is needed in the subsequent parts.

We will frequently use topological methods when studying transformation monoids. The definition of the topology of pointwise convergence for transformations monoids is analogous to the definition for groups: the closure \overline{F} of $F \subseteq \mathbb{L}^{\mathbb{L}}$ is the set of all functions $f \in \mathbb{L}^{\mathbb{L}}$ with the property that for every finite subset A of \mathbb{L} , there is a $g \in F$ such that f(a) = g(a) for all $a \in A$. A set of functions is closed if $F = \overline{F}$. We write $\langle F \rangle$ for the smallest transformation monoid that contains F. The smallest closed transformation monoid that contains a set of functions F equals $\overline{\langle F \rangle}$. The closed transformation monoids are precisely those that are endomorphism monoids of relational structures. We say that a function f is generated by a set of operations F is f is in the smallest closed monoid that contains F. A more detailed introduction to these concepts can be found in Bodirsky [6].

5.1 Ramsey classes

Let Γ, Δ be finite τ -structures. We write $\binom{\Delta}{\Gamma}$ for the set of all substructures of Δ that are isomorphic to Γ . When Γ, Δ, Θ are τ -structures, then we write $\Theta \to (\Delta)_r^{\Gamma}$ if for all functions $\chi: \binom{\Theta}{\Gamma} \to \{1, \ldots, r\}$ there exists $\Delta' \in \binom{\Theta}{\Delta}$ such that χ is constant on $\binom{\Delta'}{\Gamma}$.

Definition 30. A class of finite relational structures \mathbb{C} that is closed under isomorphisms and substructures is called Ramsey if for all $\Gamma, \Delta \in \mathbb{C}$ and arbitrary $k \geq 1$, there exists a $\Theta \in \mathbb{C}$ such that Δ embeds into Θ and $\Theta \to (\Delta)_k^{\Gamma}$.

A homogeneous structure Γ is called *Ramsey* if the class of all finite structures that embed into Γ is Ramsey. We refer the reader to Kechris, Pestov and Todorcevic [35] or Nešetřil [40] for more information about the links between Ramsey theory and homogeneous structures. An example of a Ramsey structure is (D; =)—the fact that the class of all finite structures that embed into (D; =) is Ramsey can be seen as a reformulation of Ramsey's classical result [43].

The Ramsey result that is relevant in our context (Theorem 31) is a consequence of a more powerful theorem due to Miliken [39]. The theorem in the form presented below and a direct proof of it can be found in Bodirsky & Piguet [15]. We mention that a weaker version of this theorem

(which was shown by the academic grand-father of the first author of this article [28]) has been known for a long time.

Theorem 31 (see Bodirsky [7] or Miliken [39]). The structure $(\mathbb{L}; C, \prec)$ is Ramsey.

We also need the following result.

Theorem 32 (see Bodirsky, Pinsker & Tsankov [20]). If Γ is homogeneous and Ramsey, then every expansion of Γ by finitely many constants is Ramsey, too.

5.2 Canonical functions

The typical usage of Ramsey theory in this article is for showing that the endomorphisms of Γ behave *canonically* on large parts of the domain; this will be formalized below. A wider introduction to canonical operations can be found in Bodirsky [6] and Bodirsky & Pinsker [16]. The definition of canonical functions given below is slightly different from the one given in [6] and [16]. It is easy to see that they are equivalent, though.

Definition 33. Let Γ, Δ be structures and let S be a subset of the domain D of Γ . A function $f: \Gamma \to \Delta$ is canonical on S as a function from Γ to Δ if for all $s_1, \ldots, s_n \in S$ and all $\alpha \in \operatorname{Aut}(\Gamma)$, there exists a $\beta \in \operatorname{Aut}(\Delta)$ such that $f(\alpha(s_i)) = \beta(f(s_i))$ for all $i \in \{1, \ldots, n\}$.

In Definition 33, we might omit the set S if S = D is clear from the context. Note that a function f from Γ to Δ is canonical if and only if for every $k \ge 1$ and every $t \in D^k$, the orbit of f(t) in Aut(Δ) only depends on the orbit of t in Aut(Γ).

Example 1. Write $x \succ y$ if $y \prec x$. The structure $(\mathbb{L}; C, \succ)$ is isomorphic to $(\mathbb{L}; C, \prec)$; let – be such an isomorphism. Note that – is canonical as a function from $(\mathbb{L}; C, \prec)$ to $(\mathbb{L}; C, \prec)$.

When Γ is Ramsey, then the following theorem allows us to work with canonical endomorphisms of Γ . It can be shown with the same proof as presented in Bodirsky, Pinsker, & Tsankov [20].

Theorem 34. Let Γ , Δ denote finite relational structures such that Γ is homogeneous and Ramsey while Δ is ω -categorical. Arbitrarily choose a function $f: \Gamma \to \Delta$. Then, there exists a function

 $g \in \overline{\{\alpha_1 f \alpha_2 : \alpha_1 \in \operatorname{Aut}(\Delta), \alpha_2 \in \operatorname{Aut}(\Gamma)\}}$

that is canonical as a function from Γ to Δ .

Note that expansions of homogeneous structures with constant symbols are again homogeneous. We obtain the following by combining the previous theorem and Theorem 32.

Corollary 35. Let Γ , Δ denote finite relational structures such that Γ is homogeneous and Ramsey while Δ is ω -categorical. Arbitrarily choose a function $f: \Gamma \to \Delta$ and elements c_1, \ldots, c_n of Γ . Then, there exists a function

$$g \in \overline{\{\alpha_1 f \alpha_2 : \alpha_1 \in \operatorname{Aut}(\Delta), \alpha_2 \in \operatorname{Aut}(\Gamma, c_1, \dots, c_n)\}}$$

that is canonical as a function from $(\Gamma, c_1, \ldots, c_n)$ to Δ .

5.3 The ordering property

Another important concept from Ramsey theory that we will exploit in the forthcoming proofs is the *ordering property*. We will next prove that the class of ordered leaf structure has this property.

Definition 36 (See Kechris, Pestov & Todorcevic [35] or Nešetřil [40]). Let \mathfrak{C}' be a class of finite structures over the signature $\tau \cup \{\prec\}$, where \prec denotes a linear order, and let \mathfrak{C} be the class of all τ -reducts of structures from \mathfrak{C}' . Then \mathfrak{C}' has the ordering property if for every $\Delta_1 \in \mathfrak{C}$ there exists a $\Delta_2 \in \mathfrak{C}$ such that for all expansions $\Delta'_1 \in \mathfrak{C}'$ of Δ_1 and $\Delta'_2 \in \mathfrak{C}'$ of Δ_2 there exists an embedding of Δ'_1 into Δ'_2 .

Proposition 37. Let Γ be a homogeneous relational τ -structure with domain D and suppose that Γ has an ω -categorical homogeneous expansion Γ' with signature $\tau \cup \{\prec\}$ where \prec denotes a linear order. Then, the following are equivalent.

- the class C' of finite structures that embed into Γ' has the ordering property and
- for every finite $X \subseteq D$ there exists a finite $Y \subseteq D$ such that for every $\beta \in \operatorname{Aut}(\Gamma)$ there exists an $\alpha \in \operatorname{Aut}(\Gamma')$ such that $\alpha(X) \subseteq \beta(Y)$.

Proof. First suppose that \mathcal{C}' has the ordering property and let $X \subseteq D$ be finite. Let Δ_1 be the structure induced by X in Γ . Then, there exists $\Delta_2 \in \mathcal{C}$ such that for for all expansions $\Delta'_1 \in \mathcal{C}'$ of Δ_1 and for all expansions $\Delta'_2 \in \mathcal{C}'$ of Δ_2 , there exists an embedding of Δ'_1 into Δ'_2 . Since every structure in \mathcal{C}' can be embedded into Γ' , we may assume that Δ'_2 is a substructure of Γ' with domain Y. Arbitrarily choose $\beta \in \operatorname{Aut}(\Gamma)$. Then, there exists an embedding from the structure induced by X in Γ' to the structure induced by $\beta(Y)$ in Γ' . By homogeneity of Γ' , this embedding can be extended to an automorphism α of Γ' which has the desired property.

For the converse direction, let Δ_1 be the τ -reduct of an arbitrary structure from \mathcal{C}' and let n denote the cardinality of Δ_1 . Since Γ' is ω -categorical, there is a finite number m of orbits of n-tuples. Hence, there exists a set Z of cardinality $n \cdot m$ such that for every embedding e of Δ_1 into Γ , there exists an automorphism α of Γ' such that the image of $\alpha \circ e$ is a subset of Z. By assumption, there exists a set $Y \subseteq D$ such that for every $\beta \in \operatorname{Aut}(\Gamma)$, there exists an $\alpha \in \operatorname{Aut}(\Gamma')$ and $\alpha(Z) \subseteq \beta(Y)$. Let Δ_2 be the structure induced by Y in Γ . Now, let $\Delta'_1 = (\Delta_1, \prec)$ and arbitrarily choose $\Delta'_2 = (\Delta_2, \prec) \in \mathcal{C}'$. By the choice of Z, there is an embedding f of Δ'_1 into the substructure of Γ' . By homogeneity of Γ , there is a $\beta \in \operatorname{Aut}(\Gamma)$ that maps Δ_2 to Δ'_2 . By the choice of Y, there exists an $\alpha \in \operatorname{Aut}(\Gamma')$ such that $\alpha(Z) \subseteq \beta(Y)$. Now, $\alpha \circ f$ is an embedding of Δ'_1 into Δ'_2 which concludes the proof.

Theorem 38. The class of all ordered leaf structures has the ordering property.

Proof. By Proposition 37, it is sufficient to show that for every finite $X \subseteq \mathbb{L}$, there exists a finite $Y \subseteq \mathbb{L}$ such that for every $\beta \in \operatorname{Aut}(\mathbb{L}; C)$ there exists an $\alpha \in \operatorname{Aut}(\mathbb{L}; C, \prec)$ satisfying $\alpha(X) \subseteq \beta(Y)$. Let X be an arbitrary finite subset of \mathbb{L} , let $Z = X \cup -X$ (where - is defined as in Example 1), and let Δ be the structure induced by Z in $(\mathbb{L}; C, \prec)$. Let Γ be the structure induced by a two-element subset of \mathbb{L} in $(\mathbb{L}; C, \prec)$. The exact choice is not important since all such structures are isomorphic. Since $(\mathbb{L}; C, \prec)$ is Ramsey by Theorem 31, there exists a leaf structure Θ such that $\Theta \to (\Delta)_2^{\Gamma}$. Let Y be the domain of Θ .

Now, choose some $\beta \in \operatorname{Aut}(\mathbb{L}; C)$ arbitrarily. Define the following 2-coloring of $\binom{\Theta}{\Gamma}$: suppose that $x, y \in Y$ satisfy $x \prec y$. Color the copy of Γ induced by $\{x, y\}$ red iff $\beta(x) \prec \beta(y)$ and blue otherwise. Then, there exists a copy Δ' of Δ in Θ such that all copies of Γ in Δ' have the same color. If the color is red, clearly there is an automorphism $\alpha \in \operatorname{Aut}(\mathbb{L}; C, \prec)$ such that $\alpha(X) \subseteq \beta(\Delta') \subseteq Y$. If the color is blue, then there is also an automorphism $\alpha \in \operatorname{Aut}(\mathbb{L}; C, \prec)$ such that $\alpha(X) \subseteq \beta(\Delta') \subseteq Y$ since Z also contains -X.

6 Endomorphism monoids of reducts

In this section we prove the remaining results that were stated in Section 2. We start with a description of the basic idea how to use the Ramsey theoretic tools introduced in the previous section. In our proof, we can exclusively focus on analyzing *injective* endomorphisms, because of a fundamental lemma which we describe next. Since $(\mathbb{L}; C)$ has a 2-transitive automorphism group, all reducts Γ of $(\mathbb{L}; C)$ also have a 2-transitive automorphism group. We can thus apply the following result.

Lemma 39 (see, e.g., Bodirsky [6]). Let Γ be a relational structure with a 2-transitive automorphism group. If Γ has a non-injective endomorphism, then it also has a constant endomorphism.

Let Γ be a reduct of $(\mathbb{L}; C)$. Suppose that Γ has an endomorphism e that does not preserve C, i.e., there is $(o, p, q) \in C$ such that $(e(o), e(p), e(q)) \notin C$. If e is not injective, then Γ also has a constant endomorphism by Lemma 39. In this case, the third item in Theorem 1 applies and we are done. So suppose in the following that e is injective. By Theorem 31, the structure $(\mathbb{L}; C, \prec)$ is Ramsey. Hence, Corollary 35 implies that $\{e\} \cup \operatorname{Aut}(\mathbb{L}; C)$ generates an injective function f that equals e on o, p, q and therefore still violates C, but is canonical as a function from $(\mathbb{L}; C, \prec, o, p, q)$ to $(\mathbb{L}; C, \prec)$.

As we have noted above, a canonical function f from Γ to Δ induces a function from the orbits of k-tuples in Aut(Γ) to the orbits of k-tuples in Aut(Δ); we will refer to those functions as the behavior of f. There are finitely many behaviors of canonical injections from $(\mathbb{L}; C, \prec, o, p, q)$ to $(\mathbb{L}; C, \prec)$: since the pre-image is homogeneous in a ternary language with three constants, their number is bounded by the number of functions from $\mathcal{O}_6 \to \mathcal{O}_3$, where \mathcal{O}_k denotes the set of orbits of k-tuples of distinct elements in $(\mathbb{L}; C, \prec)$. The function $s: k \mapsto |\mathcal{O}_k|$ is well-known in combinatorics (see Sloane's Integer Sequence A001813 and see [25] for various enumerative results for leaf structures on trees), and we have s(n) = (2n)!/n!. In particular, s(3) = 12 and s(6) = 30240. So the number of canonical behaviors of functions from $(\mathbb{L}; C, \prec, o, p, q)$ to $(\mathbb{L}; C, \prec)$ is bounded by 12^{30240} . For every function with one of those behaviors, we prove that Γ must be as described in item 3 and 4 of Theorem 1. Since 12^{30240} is a somewhat large number of cases, the way we treat these cases in the following is important. We then repeat the same strategy for the structure $(\mathbb{L}; Q)$ but here we have to expand with four constants, that is, we analyze canonical functions from $(\mathbb{L}; C, \prec, c_1, \ldots, c_4) \to (\mathbb{L}; C, \prec)$.

In the following, several arguments hold for the expansion of $(\mathbb{L}; C, \prec)$ by any finite number of constants $\bar{c} = (c_1, \ldots, c_n)$. The following equivalence relation plays an important role.

Definition 40. Let $\bar{c} = (c_1, \ldots, c_n) \in \mathbb{L}^n$. Then $E_{\bar{c}}$ denotes the equivalence relation defined on $\mathbb{L} \setminus \{c_1, \ldots, c_n\}$ by

$$E_{\bar{c}}(x,y) \Leftrightarrow \bigwedge_{i=1}^{n} xy | c_i$$

The equivalence classes of $E_{\bar{c}}$ are called cones (of $(\mathbb{L}; C, \bar{c})$). We write $S_a^{\bar{c}}$ for the cone that contains $a \in \mathbb{L} \setminus \{c_1, \ldots, c_n\}$.

Note that each cone induces in $(\mathbb{L}; C, \prec)$ a structure that is isomorphic to $(\mathbb{L}; C, \prec)$.

In Section 6.1, Section 6.2, and Section 6.3 we study the behavior of canonical functions with zero, one, and two constants, respectively. Finally, in Section 6.4, we put the pieces together and prove Theorem 1 and Corollary 3.

6.1 Canonical behavior without constants

In this section we analyze the behavior of canonical functions from $(\mathbb{L}; C, \prec)$ to $(\mathbb{L}; C, \prec)$. In particular, we discuss possible behaviors on cones (Corollary 45) and close with a useful lemma (Lemma 50) that shows that when a reduct Γ of $(\mathbb{L}; C)$ is preserved by functions with certain behaviors, then Γ is homomorphically equivalent to a reduct of $(\mathbb{L}; =)$.

Definition 41. Let $A \subseteq \mathbb{L}$ and $e \colon \mathbb{L} \to \mathbb{L}$ a function. Then we say that e has on A the behavior

- id if for all $x, y, z \in A$ with xy|z we have that e(x)e(y)|e(z).
- lin if for all $x, y, z \in A$ with $x \prec y \prec z$ we have that e(x)|e(y)e(z).
- nil if for all $x, y, z \in A$ with $x \prec y \prec z$ we have that e(x)e(y)|e(z).

In this case, we will also say that *e* behaves as id, lin, or nil on A, respectively. When f behaves as lin on $A = \mathbb{L}$, then we do not mention A and simply say that f behaves as lin; we make the analogous convention for all other behaviors that we define. We first prove that functions with behavior lin and nil really exist.

Lemma 42. There are functions from $\mathbb{L} \to \mathbb{L}$ which preserve \prec and have the behavior lin and nil.

Proof. A function f with behavior lin can be constructed as follows. Let v_1, v_2, \ldots be an enumeration of \mathbb{L} . Inductively suppose that there exists a function $f: \{v_1, \ldots, v_n\} \to \mathbb{L}$ such that for all $x, y, z \in \{v_1, \ldots, v_n\}$ with $x \prec y \prec z$ it holds that f(x)|f(y)f(z) and $f(x) \prec f(y) \prec f(z)$. This is clearly true for n = 1. We prove that f has an extension f' to v_{n+1} with the same property. Let w_1, \ldots, w_{n+1} be such that $\{w_1, \ldots, w_{n+1}\} = \{v_1, \ldots, v_{n+1}\}$ and $w_1 \prec \cdots \prec w_{n+1}$. We consider the following cases.

- $v_{n+1} = w_1$. There exists a $c \in \mathbb{L}$ such that $c|f(w_2)f(w_3)$ (see axiom C4). Note that if n = 1, then let c be such that $c \neq f(w_2)$. Pick c such that $c \prec f(w_2)$, and define $f'(v_{n+1}) = c$.
- $v_{n+1} = w_i$ for $i \in \{2, \ldots, n-1\}$. There exists a $c \in \mathbb{L}$ such that $f(w_{i-1})|cf(w_{i+1})$ and $c|f(w_{i+1})f(w_{i+2})$ (see Axiom C7). Pick c such that $f(w_{i-1}) \prec c \prec f(w_{i+1})$, and define $f'(v_{n+1}) = c$.
- $v_{n+1} = w_n$. There exists a $c \in \mathbb{L}$ such that $c \neq f(w_{n+1})$ and $f(w_{n-1})|f(w_{n+1})c$ (see Axiom C6). Pick c such that $f(w_{n-1}) \prec c \prec f(w_{n+1})$, and define $f'(v_{n+1}) = c$.
- $v_{n+1} = w_{n+1}$. There exists a $c \in \mathbb{L}$ such that $c \neq f(w_n)$ and $f(w_{n-1})|f(w_n)c$ (see Axiom C6). Pick c such that $f(w_n) \prec c$, and define $f'(v_{n+1}) = c$.

The function defined on all of \mathbb{L} in this way has the behavior lin. The existence of a function with behavior nil can be shown analogously.

The functions lin and nil constructed in Lemma 42 preserve the linear order \prec . In general, a function $f : \mathbb{L} \to \mathbb{L}$ with behavior lin or nil may not preserve \prec , however together with Aut($\mathbb{L}; C$) generates lin or nil, respective. In the following, we will use lin and nil also to denote the functions with behavior lin and nil that have been constructed in Lemma 42; whether we mean the behavior or the function lin and nil will always be clear from the context. As we see in the following proposition, the two functions are closely related.

Proposition 43. Aut($\mathbb{L}; C$) \cup {nil} generates lin, and Aut($\mathbb{L}; C$) \cup {lin} generates nil.

Proof. Let $n \geq 1$ and arbitrarily choose $t \in \mathbb{L}^n$. Then $-\operatorname{nil}(-t)$ and $\operatorname{lin}(t)$ induce isomorphic substructures in $(\mathbb{L}; C, \prec)$, and by the homogeneity of $(\mathbb{L}; C, \prec)$ there is an $\alpha \in \operatorname{Aut}(\mathbb{L}; C, \prec)$ such that $\alpha(-\operatorname{nil}(-t)) = \operatorname{lin}(t)$. It follows that $\operatorname{lin} \in \overline{\langle \operatorname{Aut}(\mathbb{L}; C) \cup \{\operatorname{nil}\} \rangle}$. The fact that $\operatorname{Aut}(\mathbb{L}; C) \cup \{\operatorname{lin}\}$ generates nil can be shown in the same way.

The following lemma classifies the behavior of canonical injective functions from $(\mathbb{L}; C, \prec)$ to $(\mathbb{L}; C, \prec)$ on sufficiently large subsets of \mathbb{L} .

Lemma 44. Let $S \subseteq \mathbb{L}$ be a set that contains four elements x, y, u, v such that xy|uv, and let $f: D \to D$ be injective and canonical on S as a function from $(\mathbb{L}; C, \prec)$ to $(\mathbb{L}; C, \prec)$. Then f behaves as id, lin, or nil on S.

Proof. Since f is canonical on S as a function from $(\mathbb{L}; C, \prec)$ to $(\mathbb{L}; C, \prec)$, it either preserves or reverses the order \prec on S. We focus on the case that f preserves \prec on S, since the order-reversing case is analogous. Without loss of generality, we assume that $x \prec y \prec u \prec v$. Since f preserves \prec , we have $f(x) \prec f(y) \prec f(u) \prec f(v)$. The following cases are exhaustive.

- f(x)f(y)|f(u) and f(y)f(u)|f(v). By canonicity, f behaves as nil on S.
- f(x)f(y)|f(u) and f(y)|f(u)f(v). By canonicity, f behaves as id on S.
- f(x)|f(y)f(u) and f(y)|f(u)f(v). By canonicity, f behaves as lin on S.
- f(x)|f(y)f(u) and f(y)f(u)|f(v). By canonicity, f(x)|f(y)f(v) and f(x)f(u)|f(v). It is easy to see that those conditions are impossible to satisfy over $(\mathbb{L}; C)$.



Figure 2: Illustration of the re-ordering of Lemma 47.

Corollary 45. Let $\bar{c} \in \mathbb{L}^n$ for $n \ge 0$, let S be a cone of $(\mathbb{L}; C, \bar{c})$, and let $f: \mathbb{L} \to \mathbb{L}$ be an injection that is canonical on S as a function from $(\mathbb{L}; C, \prec, \bar{c})$ to $(\mathbb{L}; C, \prec)$. Then f behaves as id, lin, or nil on S.

Proof. Note that f is on S canonical as a function from $(\mathbb{L}; C, \prec)$ to $(\mathbb{L}; C, \prec)$; also note that every cone contains elements x, y, u, v such that xy|uv. Hence, the statement follows from Lemma 44. \Box

We finally show that if Γ is preserved by lin, then Γ is homomorphically equivalent to a reduct of $(\mathbb{L}; =)$; this will be a consequence of the stronger Lemma 50 below.

Definition 46. For $a_1, a_2, \ldots, a_k \in \mathbb{L}$, $k \ge 2$, we write $\operatorname{Nil}(a_1, a_2, \ldots, a_k)$ if $a_1 \prec a_2 \prec \cdots \prec a_k$ and $a_1 a_2 \ldots a_{i-1} | a_i$ for all $i \in \{2, \ldots, k\}$.

Observe that for all $a_1, \ldots, a_k \in \mathbb{L}$ such that $a_1 \prec \cdots \prec a_k$ we have Nil(nil(a_1, \ldots, a_k)) (recall from Section 3.1 that we apply functions to tuples componentwise). Also observe that all k-tuples in Nil lie in the same orbit of k-tuples.

Lemma 47. Let $a_1, \ldots, a_k \in \mathbb{L}$ be such that $\operatorname{Nil}(a_1, \ldots, a_k)$. Then for every $p \in \{1, \ldots, k\}$ there is an $e \in \operatorname{Aut}(\mathbb{L}; C) \cup \{\operatorname{nil}\}$ such that $\operatorname{Nil}(e(a_p, a_1, \ldots, a_{p-1}, a_{p+1}, a_{p+2}, \ldots, a_k))$.

Proof. By Proposition 16 and the homogeneity of $(\mathbb{L}; C)$ there exists an $\alpha \in \operatorname{Aut}(\mathbb{L}; C)$ such that $\alpha(a_p) \prec \alpha(a_1) \prec \alpha(a_2) \prec \cdots \prec \alpha(a_{p-1}) \prec \alpha(a_{p+1}) \prec \cdots \prec \alpha(a_k)$; see Figure 2. Define $e := \operatorname{nil} \circ \alpha$. By the observation above we have $\operatorname{Nil}(e(a_p, a_1, a_2, \ldots, a_{p-1}, a_{p+1}, a_{p+2}, \ldots, a_k))$, as desired. \Box

Lemma 48. Let $a_1, \ldots, a_k \in \mathbb{L}$ be such that $\operatorname{Nil}(a_1, \ldots, a_k)$. Then for any $p \in \{1, 2, \ldots, k-1\}$, there is an $e \in \langle \operatorname{Aut}(\mathbb{L}; C) \cup \{\operatorname{nil}\} \rangle$ such that $e(a_p) = a_{p+1}$, $e(a_{p+1}) = a_p$, and $e(a_i) = a_i$ for every $i \in \{1, \ldots, k\} \setminus \{p, p+1\}$.

Proof. By the homogeneity of $(\mathbb{L}; C)$ there is an $\alpha \in \operatorname{Aut}(\mathbb{L}; C)$ such that $\alpha(a_{p+1}) \prec \alpha(a_p) \prec \alpha(a_{p-1}) \prec \cdots \prec \alpha(a_2) \prec \alpha(a_1) \prec \alpha(a_{p+2}) \prec \alpha(a_{p+3}) \prec \cdots \prec \alpha(a_k)$; see Figure 3. Let $z_i := \operatorname{nil}(\alpha(a_i))$ for $i \in \{1, 2, \ldots, k\}$. Clearly, we have $\operatorname{Nil}(z_{p+1}, z_p, z_{p-1}, \ldots, z_2, z_1, z_{p+2}, z_{p+3}, \ldots, z_k)$. Starting with the tuple $(z_{p+1}, z_p, z_{p-1}, \ldots, z_2, z_1, z_{p+2}, z_{p+3}, \ldots, z_k)$ we repeatedly apply Lemma 47 to the resulting tuple at the positions $p := 3, 4, \ldots, p-1$ in this order. In this way, we obtain in the first step an $e_1 \in M := \langle \operatorname{Aut}(\mathbb{L}; C) \cup \{\operatorname{nil}\} \rangle$ such that

Nil
$$(e_1(z_{p-1}, z_{p+1}, z_p, z_{p-2}, \dots, z_2, z_1, z_{p+2}, z_{p+3}, \dots, z_k))$$
.

In the second step, we obtain an $e_2 \in M$ such that

Nil
$$(e_2(z_{p-2}, z_{p-1}, z_{p+1}, z_p, z_{p-3}, \dots, z_2, z_1, z_{p+2}, z_{p+3}, \dots, z_k))$$
.



Figure 3: Illustration of the re-ordering of Lemma 48.

In the *i*-th step, we obtain an $e_i \in M$ such that

$$Nil(e_i(z_{p-i}, z_{p-i+1}, ..., z_{p-2}, z_{p-1}, z_{p+1}, z_p, z_{p-i-1}, ..., z_2, z_1, z_{p+2}, z_{p+3}, ..., z_k)).$$

For i = p - 1, we therefore obtain an $e' \in M$ such that

Nil
$$(e'(z_1, z_2, \ldots, z_{p-2}, z_{p-1}, z_{p+1}, z_p, z_{p+2}, \ldots, z_k))$$
.

Define $f := e' \circ \operatorname{nil} \circ \alpha$ and observe that $\operatorname{Nil}(f(a_1, a_2, \dots, a_{p-1}, a_{p+1}, a_p, a_{p+2}, \dots, a_k))$. Therefore, $f(a_1, \dots, a_{p-1}, a_{p+1}, a_p, a_{p+2}, \dots, a_k)$ and (a_1, a_2, \dots, a_k) are in the same orbit in $(\mathbb{L}; C)$, and there is $\gamma \in \operatorname{Aut}(\mathbb{L}; C)$ such that $\gamma(f(a_p, a_{p+1})) = (a_{p+1}, a_p)$ and $\gamma(f(a_i)) = a_i$ for all $i \in \{1, \dots, k\} \setminus \{p, p+1\}$. Then $e := \gamma \circ f \in M$ has the desired property.

We write S_k for the symmetric group on $\{1, \ldots, k\}$.

Lemma 49. Let $(x_1, x_2, \ldots, x_k), (y_1, \ldots, y_k) \in \text{Nil}$, and arbitrarily choose $\delta \in S_k$. Then there exists an $e \in M := \langle \text{Aut}(\mathbb{L}; C) \cup \{\text{nil}\} \rangle$ such that $e(x_i) = y_{\delta(i)}$.

Proof. By Lemma 48, for each $p \in \{1, 2, ..., k-1\}$ there is an $e_p \in M$ such that $e_p(x_p) = x_{p+1}$, $e_p(x_{p+1}) = x_p$, and $e_p(x_i) = x_i$ for all $i \in \{1, ..., k\} \setminus \{p, p+1\}$. Since S_k is generated by the transpositions $(1, 2), (2, 3), \ldots, (k-1, k)$, it follows that there exists $e' \in \langle \{e_p : 1 \le p \le k\} \rangle \subseteq M$ such that $e'(x_i) = x_{\delta(i)}$ for all $i \in \{1, ..., k\}$. By the homogeneity of $(\mathbb{L}; C)$, there exists an $\alpha \in \operatorname{Aut}(\mathbb{L}; C)$ such that $\alpha(x_i) = y_i$ for all $i \in \{1, ..., k\}$. Then $e := \alpha \circ e'$ satisfies $e(x_i) = y_{\delta(i)}$. \Box

We can finally prove the announced result.

Lemma 50. Let Γ be a reduct of $(\mathbb{L}; C)$. Let $\overline{c} \in \mathbb{L}^n$ and suppose that Γ has an endomorphism behaving as lin on a cone of $(\mathbb{L}; C, \overline{c})$. Then Γ is homomorphically equivalent to a reduct of $(\mathbb{L}; =)$.

Proof. Recall that each cone S of $(\mathbb{L}; C, \bar{c})$ induces in $(\mathbb{L}; C, \prec)$ a structure that is isomorphic to $(\mathbb{L}; C, \prec)$. Let e be an endomorphism of Γ that behaves as lin on S and arbitrarily choose a finite set $A \subseteq \mathbb{L}$. By the homogeneity of $(\mathbb{L}; C, \prec)$ there are automorphisms α, β of $(\mathbb{L}; C, \prec)$ such that $\lim(x) = \alpha(e(\beta(x)))$ for all $x \in A$. Since $\operatorname{End}(\Gamma)$ is closed we have that $\lim \in \operatorname{End}(\Gamma)$ and $\operatorname{nil} \in \operatorname{End}(\Gamma)$ by Proposition 43.

Our proof has two steps: we first prove that the structure Δ induced by $D := \operatorname{nil}(\mathbb{L})$ is isomorphic to a reduct of $(\mathbb{Q}; <)$ and then we prove in the next step that Δ is in fact isomorphic to a reduct of $(\mathbb{L}; =)$. Clearly, this implies the statement since Γ and Δ are homomorphically equivalent. Let τ be the signature of Γ . We first show that for every $R \in \tau$, the relation R^{Δ} has a first-order definition in $(D; \prec)$. The relation R^{Γ} has a first-order definition ϕ in $(\mathbb{L}; C)$. An atomic sub-formula C(x; yz) of ϕ holds in Δ if and only if $y \prec z \prec x, z \prec y \prec x$ or $y = z \land x \neq y$. Hence, if we replace in ϕ all occurrences of C(x; yz) by $y \prec z \prec x \lor z \prec y \prec x \lor (y = z \land x \neq y)$, we obtain a formula that defines R^{Δ} over $(D; \prec)$. Since $(\mathbb{L}; \prec)$ is isomorphic to $(\mathbb{Q}; <)$ and nil preserves \prec , it follows that $(D; \prec)$ is isomorphic to $(\mathbb{Q}; <)$, too. Hence, Δ is isomorphic to a reduct of $(\mathbb{Q}; <)$.

To show that Δ is isomorphic to a reduct of $(\mathbb{L};=)$, let X be a finite subset of D and α be a permutation of D. By Proposition 19 and the fact that $\operatorname{End}(\Delta)$ is a closed subset of D^D , it suffices to find an $e \in \operatorname{End}(\Delta)$ such that $e(x) = \alpha(x)$ for all $x \in X$. Since $X \subseteq \operatorname{nil}(\mathbb{L})$, the elements of X can be enumerated by x_1, \ldots, x_n such that $\operatorname{Nil}(x_1, \ldots, x_n)$. Since $\alpha(X) \subseteq D = \operatorname{nil}(\mathbb{L})$, there is a $\gamma \in S_n$ such that $\operatorname{Nil}(\alpha(x_{\gamma(1)}), \ldots, \alpha(x_{\gamma(n)}))$. We apply Lemma 49 to (x_1, \ldots, x_n) , $(y_1, \ldots, y_n) := (\alpha(x_{\gamma(1)}), \ldots, \alpha(x_{\gamma(n)}))$, and $\delta = \gamma^{-1}$, and obtain an $f \in \operatorname{End}(\Gamma)$ such that $f(x_i) =$ $y_{\delta(i)} = \alpha(x_{\gamma\gamma^{-1}(i)}) = \alpha(x_i)$ for all $i \in \{1, \ldots, k\}$. The restriction of $\operatorname{nil} \circ f$ to D is an endomorphism f' of Δ . Since nil preserves \prec and Δ is a reduct of $(D; \prec)$ which is isomorphic to $(\mathbb{Q}; <)$, we have that $(\alpha(x_1), \ldots, \alpha(x_n)) = (f(x_1), \ldots, f(x_n))$ and $(f'(x_1), \ldots, f'(x_n))$ lie in the same orbit in Δ . Hence, there exists an $\beta \in \operatorname{Aut}(\Delta)$ such that $\beta(f'(x_i)) = \alpha(x_i)$ for all $i \in \{1, \ldots, n\}$, and $\beta \circ f'$ is an endomorphism of Δ as required. \Box

6.2 Canonical behavior with one constant

In this section we study the behavior of canonical functions from $(\mathbb{L}; C, \prec, c_1)$ to $(\mathbb{L}; C, \prec)$. Some important behaviors are introduced in Definition 51. We then show in Sections 6.2.1, 6.2.2, and 6.2.3 that when a function f has some of those behaviors on a c-universal set, then $\{f\} \cup \operatorname{Aut}(\mathbb{L}; C)$ generates lin or $\operatorname{End}(\mathbb{L}; Q)$. Finally, Section 6.2.4 classifies behaviors of canonical functions from $(\mathbb{L}; C, \prec, c_1)$ to $(\mathbb{L}; C, \prec)$.

Definition 51. Let $c \in \mathbb{L}$ and $A \subseteq \mathbb{L} \setminus \{c\}$. Let $e: \mathbb{L} \to \mathbb{L}$ be a function such that

- 1. for any $a \in A$ we have that $e(c)|e(A \cap S_a^c)|$,
- 2. for any $a \in A$, e preserves C on $A \cap S_a^c$, and
- 3. for any $a, b \in A$ we have either $S_a^c = S_b^c$ or $e(A \cap S_a^c)|e(A \cap S_b^c)$.

Then we say that e has on A the behavior

 id_c iff for all $x, y, z \in A$ with x|yzc and y|zc we have that e(x)|e(y)e(z)e(c) and e(y)|e(z)e(c).

 cut_c iff for all $x, y, z \in A$ with x|yzc and y|zc we have that e(x)e(y)e(z)|e(c) and e(x)|e(y)e(z).

 rer_{c} iff for all $x, y, z \in A$ with x|yzc and y|zc we have that e(x)e(y)e(z)|e(c) and e(x)e(y)|e(z).

rer *c* iff for all $x, y \in A$ with x|yc we have that e(y)|e(x)e(c).

6.2.1 The behavior cut_c

Recall that a set $A \subseteq \mathbb{L} \setminus \{c\}$ is called *c*-universal if for any finite $U \subset \mathbb{L}$ and $u \in U$, there is $\alpha \in \operatorname{Aut}(\mathbb{L}; C)$ such that $\alpha(u) = c$ and $\alpha(U) \subseteq A \cup \{c\}$. In this section we prove that for all $c \in \mathbb{L}$, functions with behavior cut_c on a *c*-universal set together with $\operatorname{Aut}(\mathbb{L}; C)$ generate lin. This follows from the following more general fact.

Lemma 52 (Cut Lemma). Let $\operatorname{Aut}(\mathbb{L}; C) \subseteq M \subseteq \mathbb{L}^{\mathbb{L}}$ be such that for any finite $U \subset \mathbb{L}$ and $u \in U$, there exists $g \in M$ that behaves as cut_u on $U \setminus \{u\}$. Then M generates nil and lin.

Proof. By Proposition 43, it suffices to show that M generates nil. We show that for all k and all $x_1, \ldots, x_k \in \mathbb{L}$ there is an $f \in \langle M \rangle$ such that $f(x_i) = \operatorname{nil}(x_i)$ for all $i \leq k$. We prove this by induction on k. Clearly, the claim holds for k = 1 so assume that k > 1. Suppose without loss of generality that $x_1 \prec x_2 \prec \cdots \prec x_k$. We inductively assume that there is an $f' \in \langle M \rangle$ such that $f'(x_i) = \operatorname{nil}(x_i)$ for all i < k. Let $U := f'(\{x_1, \ldots, x_k\})$ and $u := f'(x_k)$. By assumption, there exists a $g \in M$ that behaves as cut_u on $U \setminus \{u\}$.

Set $f'' = g \circ f'$. We claim that $f''(x_1) \cdots f''(x_i) | f''(x_{i+1})$ for all $1 \leq i < k$. For i < k - 1, this follows from the inductive assumption that $f'(x_1) \cdots f'(x_i) | f'(x_{i+1})$, and the assumption that g behaves as cut_u on $U \setminus \{u\}$ and in particular preserves C on $U \setminus \{u\}$. For i = k - 1, note that that $g(u)|g(U \setminus \{u\})$ since g behaves as cut_u on $U \setminus \{u\}$, and $f''(x_1), \ldots, f''(x_i) \subseteq g(U \setminus \{u\})$. Therefore, $f''(x_1) \cdots f''(x_{k-1})|f''(x_k)$ which concludes the proof of the claim.

Since $\operatorname{nil}(x_1) \cdots \operatorname{nil}(x_i) | \operatorname{nil}(x_{i+1})$ for all $1 \leq i < k$, the homogeneity of $(\mathbb{L}; C)$ implies that there exists an $\alpha \in \operatorname{Aut}(\mathbb{L}; C) \subseteq M$ such that $\alpha(f''(x_i)) = \operatorname{nil}(x_i)$ for all $i \leq k$. Then $f := \alpha \circ f'' \in \langle M \rangle$ has the desired property which concludes the proof. \Box

Corollary 53. Let $c \in \mathbb{L}$, let $A \subseteq \mathbb{L} \setminus \{c\}$ be *c*-universal, and let *g* be a function that behaves as cut_c on *A*. Then $\{g\} \cup \operatorname{Aut}(\mathbb{L}; C)$ generates lin.

Proof. The *c*-universality of *A* implies that for every finite $U \subset \mathbb{L}$ and $u \in U$ there exists an $\alpha \in \operatorname{Aut}(\mathbb{L}; C)$ such that $\alpha(U \setminus \{u\}) \subseteq A$ and $\alpha(u) = c$. Then $g \circ \alpha$ behaves as cut_u on *U* and the statement follows from Lemma 52.

6.2.2 The behavior rer_c

We will next prove that for all $c \in \mathbb{L}$, functions with behavior rer_c on a c-universal set together with Aut($\mathbb{L}; C$) generate End($\mathbb{L}; Q$). We need the following lemmas in some later proofs.

Lemma 54. Arbitrarily choose $X \subseteq \mathbb{L}$ and $c \in X$. If $f : \mathbb{L} \to \mathbb{L}$ preserves Q on every 4-element subset of X that contains c, then f preserves Q on all of X.

Proof. Let $a_1, a_2, a_3, a_4 \in X$ be such that $a_1a_2 : a_3a_4$. We show $f(a_1)f(a_2) : f(a_3)f(a_4)$. It is easy to see that this holds if a_1, a_2, a_3, a_4 are not pairwise distinct and it holds by assumption when $c \in \{a_1, a_2, a_3, a_4\}$. Suppose that this is not the case. Then $a_1a_2c : a_3a_4$ or $a_1a_2 : ca_3a_4$. The latter case can be treated analogously to the former so we assume that $a_1a_2c : a_3a_4$. In particular, $a_1c : a_3a_4$ and $a_2c : a_3a_4$ so $f(a_1)f(c) : f(a_3)f(a_4)$ and $f(a_2)f(c) : f(a_3)f(a_4)$. It follows from Lemma 12 that $f(a_1)f(a_2) : f(a_3)f(a_4)$ as desired.

Lemma 55. Let $A \subseteq \mathbb{L} \setminus \{c\}$ and $f : \mathbb{L} \to \mathbb{L}$ be such that f behaves as rer_c on A. Then f preserves Q on $A \cup \{c\}$.

Proof. By Lemma 54 it suffices to show that f preserves Q on $\{x, y, z, c\}$, where x, y, z are pairwise distinct elements in A. The following cases are essential.

• xyz|c.

It follows from Item 1 and Item 2 of Definition 51 that f(x)f(y)f(z)|f(c) and f preserves C on $\{x, y, z\}$. This implies that (x, y, z, c) and (f(x), f(y), f(z), f(c)) are in the same orbit of Aut($\mathbb{L}; C$). Thus f preserves Q on $\{x, y, z, c\}$.

• xy|zc

Clearly, we have xy : zc. It follows from Item 1 of Definition of 51 that f(x)f(y)|f(c). It follows from Item 3 of Definition 51 that f(x)f(y)|f(z). It implies that f(x)f(y) : f(z)f(c). Thus f preseves Q on $\{x, y, z, c\}$.

• $x|yzc \wedge yz|c$

Clearly, we have xc : yz. It follows from Item 1 of Definition 51 that f(y)f(z)|f(c). It follows from Item 3 of Definition 51 that f(x)|f(y)f(z). It implies that f(y)f(z) : f(x)f(c). Thus f preserves Q on $\{x, y, z, c\}$.

• $x|yzc \wedge y|zc$

Clearly, we have xy : zc. By the definition of rer_c that $f(x)f(y)f(z)|f(c) \wedge f(x)f(y)|f(z)$. It implies that f(x)f(y) : f(z)f(c). Thus f preserves Q on $\{x, y, z, c\}$.

The other cases can be obtained from the above cases by exchanging the roles of x, y, z.

The following generation lemma is flexible and will be useful later.

Lemma 56 (Rerooting Lemma, general form). Let $\operatorname{Aut}(\mathbb{L}; C) \subseteq M \subseteq \mathbb{L}^{\mathbb{L}}$ be such that for any finite $U \subset \mathbb{L}$ and $u \in U$ there exists $g \in M$ that behaves as rer_u on $U \setminus \{u\}$. Then M generates $\operatorname{End}(\mathbb{L}; Q)$.

Proof. We follow almost literally the proof of Lemma 26 (the rerooting lemma). Arbitrarily choose $f \in \operatorname{End}(\mathbb{L}; Q)$ and let A be an arbitrary finite subset of \mathbb{L} . We have to show that $\langle M \rangle$ contains an operation e such that e(x) = f(x) for all $x \in A$. This is trivial when |A| = 1 so we assume that $|A| \geq 2$. By Lemma 24, there exists a non-empty proper subset B of A such that $f(B)|f(A \setminus B)$ and B|a for all $a \in A \setminus B$. By the homogeneity of $(\mathbb{L}; C)$ we can choose an element $c \in \mathbb{L} \setminus A$ such that c|B and $(B \cup \{c\})|a$ for all $a \in A \setminus B$. By assumption, there exists a $g \in M$ such that g behaves as rer_c on A. By Lemma 55, g preserves Q on A.

We claim that $g(B)|g(A \setminus B)$. First, we show that $g(b_1)g(b_2)|g(a)$ for every $b_1, b_2 \in B$ and $a \in A \setminus B$. By the choice of c we have $b_1b_2 : ac$. Since g preserves Q on A, we have $g(b_1)g(b_2) : g(a)g(c)$. Since $g(c)|g(b_1)g(b_2)g(a)$, we have $g(b_1)g(b_2)|g(a)$. Next, we show that $g(b)|g(a_1)g(a_2)$ for every $b \in B$ and $a_1, a_2 \in A \setminus B$. By the choice of c we have $bc : a_1a_2$. Since g preserves Q on A, we have $g(b)g(c) : g(a_1)g(a_2)$. Since $g(c)|g(b_2)g(a_2)$. Since $g(c)|g(b_2)g(a_2)$.

Let $\beta: g(A) \to f(A)$ be defined by $\beta(x) = f(g^{-1}(x))$ for all $x \in g(A)$. Since both g and f preserve Q, we have that β preserves Q by Lemma 13. Since $\beta(g(B))|\beta(g(A \setminus B))$, the conditions of Lemma 23 apply to β for $A_1 := g(B)$ and $A_2 := g(A \setminus B)$, and hence β preserves C. By the homogeneity of $(\mathbb{L}; C)$, there exists an $\gamma \in \operatorname{Aut}(\mathbb{L}; C) \subseteq M$ that extends β . Then $e := \gamma \circ g \in \langle M \rangle$ has the desired property.

Corollary 57. Let $c \in \mathbb{L}$, let $A \subseteq \mathbb{L} \setminus \{c\}$ be *c*-universal, and let *g* be a function that behaves as rer_c on *A*. Then $\{g\} \cup \operatorname{Aut}(\mathbb{L}; C)$ generates $\operatorname{End}(\mathbb{L}; Q)$.

Proof. We claim that the conditions of Lemma 56 apply to $M := \langle \{g\} \cup \operatorname{Aut}(\mathbb{L}; C) \rangle$. Let $U \subset \mathbb{L}$ be finite and arbitrarily choose $u \in U$. By *c*-universality of A there exists an $\alpha \in \operatorname{Aut}(\mathbb{L}; C)$ such that $\alpha(U) \subseteq A$ and $\alpha(u) = c$. Since g behaves as rer_c on $\alpha(U)$, the function $g \circ \alpha$ behaves as rer_u on U. By Lemma 56 $\langle \{g\} \cup \operatorname{Aut}(\mathbb{L}; C) \rangle$ generates $\operatorname{End}(\mathbb{L}; Q)$, and so does $\{g\} \cup \operatorname{Aut}(\mathbb{L}; C)$. \Box

By Lemma 55, a function with behavior rer_c on A preserves Q on A. We now characterize the situation where a function preserving Q behaves as rer_c.

Lemma 58. Let X be a subset of \mathbb{L} , arbitrarily choose $a \in X$, and let $f: \mathbb{L} \to \mathbb{L}$ be a function that preserves Q on X and has the property that $f(a)|f(X \setminus \{a\})$. Then f preserves C on $X \cap S_x^a$ for every $x \in X \setminus \{a\}$, and for any $x, y \in X \setminus \{a\}$ either $S_x^a = S_y^a$ or $f(X \cap S_x^a)|f(X \cap S_y^a)$.

Proof. Arbitrarily choose $x \in X \setminus \{a\}$ and pick pairwise distinct elements $u, v, w \in X \cap S_x^a$. By C8 we can assume that uv|w. Clearly, aw : uv and it follows that f(a)f(w) : f(u)f(v). Since f(a)|f(u)f(v)f(w) by the assumptions on f, we have f(w)|f(u)f(v). This concludes the proof of the first assertion of the lemma.

To show the remaining assertion, let $x, y \in X \setminus \{a\}$ and assume that $S_x^a \neq S_y^a$. We claim that $f(X \cap S_x^a)|f(y)$. It suffices to show that f(x')f(x)|f(y) for any $x' \in X \cap S_x^a$. Clearly, we have ay : xx' and, consequently, f(a)f(y) : f(x)f(x'). Since f(a)|f(x)f(x')f(y), it follows that f(x)f(x')|f(y). This concludes the proof of the claim. Similarly, it can be shown that $f(x)|f(X \cap S_y^a)$. To complete the argument, arbitrarily choose $x_1, x_2 \in X \cap S_x^a$ and $y_1, y_2 \in X \cap S_y^a$. The claim implies that $f(x_1)f(x)|f(y)$ and $f(x_2)f(x)|f(y)$. Similarly, $f(x)|f(y_1)f(y)$ and $f(x)|f(y_2)f(y)$ because $f(x)|f(X \cap S_y^a)$. This implies that $f(x_1)f(x_2)|f(y_1)f(y_2)$, and, consequently, $f(X \cap S_x^a)|f(X \cap S_y^a)$. \Box

Corollary 59. Let X be a subset of \mathbb{L} , arbitrarily choose $a \in X$, and let $f : \mathbb{L} \to \mathbb{L}$ be a function such that $f(a)|f(X \setminus \{a\})$. Then f behaves as rer_a on X if and only if f preserves Q on X.

Proof. By Lemma 55 if f behaves as rer_a on X then f preserves Q on X. Conversely, suppose that f preserves Q on X. By Lemma 58, it remains to show that for $x, y, z \in X \setminus \{a\}$ such that $x|yza \wedge y|za$ we have $f(a)|f(x)f(y)f(z) \wedge f(z)|f(x)f(y)$. Clearly, we have xy : za and f(x)f(y): f(z)f(a). Since f(a)|f(x)f(y)f(z), it follows that f(x)f(y)|f(z).

6.2.3 The behavior rer_c

We finally study functions with behavior \tilde{rer}_c on a *c*-universal set. We need the following lemma for the proof of the next important lemma.

Lemma 60. Let $c \in \mathbb{L}$ and $A \subseteq \mathbb{L} \setminus \{c\}$ be a *c*-universal set. Let $f : \mathbb{L} \to \mathbb{L}$ be a function that behaves as rer_c on A. Then f preserves Q on A.

Proof. Let x, y, z, t be elements in A such that xy : zt. It suffices to show that f(x)f(y) : f(z)f(t). Without loss of generality we assume that $xy|z \wedge xy|t$. The following cases are exhaustive.

• c|xy.

We will show that f(x)f(y)|f(z). If xyz|c holds, then by Item 2 of Definition 51 we have f(x)f(y)|f(z). If xy|zc holds, then by the definition of rer_c, we have f(z)|f(x)f(c) and f(z)|f(y)f(c). It follows that f(x)f(y)|f(z). If z|xyc holds, by Item 3 of Definition 51, we have f(x)f(y)|f(z). These cases are exhaustive, thus f(x)f(y)|f(z). By the same arguments we have f(x)f(y)|f(t). Thus f(x)f(y):f(z)f(t).

• $\neg c|xy$.

We consider the case cx|y. The case cy|x are proved similarly. Since xy|z and xy|t, it follows that cxy|z and cxy|t. By the definition of rer_c, we have $f(c)f(z)|f(x)\wedge f(c)f(z)|f(y)\wedge f(c)f(z)|f(y)$. It follows from $f(c)f(z)|f(x)\wedge f(c)f(t)|f(x)$ that f(z)f(t)|f(x). It follows from $f(c)f(z)|f(y)\wedge f(c)f(t)|f(y)$ that f(z)f(t)|f(y). It implies that $f(z)f(t)|f(x)\wedge f(z)f(t)|f(x)\wedge f(z)f(t)|f(y)$. Thus f(x)f(y):f(z)f(t).

Lemma 61. Let $c \in \mathbb{L}$, let $A \subseteq \mathbb{L} \setminus \{c\}$ be *c*-universal, and let $g: \mathbb{L} \to \mathbb{L}$ be a function that behaves as $\tilde{\operatorname{rer}}_c$ on A. Then $\{g\} \cup \operatorname{Aut}(\mathbb{L}; C)$ generates lin.

Proof. The proof has two steps. We first show that $\{g\} \cup \operatorname{Aut}(\mathbb{L}; C)$ generates $\operatorname{End}(\mathbb{L}; Q)$, and then prove that $\{g\} \cup \operatorname{End}(\mathbb{L}; Q)$ generates lin.

For the first step it suffices to show that $\langle \{g\} \cup \operatorname{Aut}(\mathbb{L}; C) \rangle$ satisfies the conditions of Lemma 56. Let U be a non-empty finite subset of \mathbb{L} and arbitrarily choose an element $u \in U$. Let $c' \in \mathbb{L}$ be such that $c' \neq u$ and for every $v \in U \setminus \{u\}$ we have uc'|v. Since A is c-universal, there is an $\alpha \in \operatorname{Aut}(\mathbb{L}; C)$ such that $\alpha(U) \subseteq A$ and $\alpha(c') = c$. Arbitrarily choose two members v_1, v_2 of $U \setminus \{u\}$. By the choice of c' we have that either $uc'|v_1 \wedge uc'v_1|v_2$, $uc'|v_2 \wedge uc'v_2|v_1$, or $v_1v_2|uc'$. Since α preserves C and g behaves as rer $_c$ on A, it follows that $g(\alpha(v_1))g(\alpha(v_2))|g(\alpha(u))$ in each of the three cases. This implies that $g(\alpha(U \setminus \{u\}))|g(\alpha(u))$. By Lemma 60, g preserves Q on A so $g \circ \alpha$ preserves Q on U. By Corollary 59 the function $g \circ \alpha$ behaves as rer $_u$ on U. This concludes the first step.

To show that $\{g\} \cup \operatorname{End}(\mathbb{L}; Q)$ generates lin, we use Lemma 52. Let $U \subseteq \mathbb{L}$ be finite and arbitrarily choose $u \in U$. Let $v \in \mathbb{L}$ be such that U|v. By the *c*-universality of A there is an $\alpha \in \operatorname{Aut}(\mathbb{L}; C)$ such that $\alpha(U \cup \{v\}) \subseteq A \cup \{c\}$ and $\alpha(u) = c$. Since for every $x \in U \setminus \{u\}$ we have $c\alpha(x)|\alpha(v)$, it follows that $g(c)g(\alpha(v))|g(\alpha(x))$ for every $x \in U \setminus \{u\}$. This property together with the homogeneity of $(\mathbb{L}; Q)$ allows us to choose $\beta \in \operatorname{Aut}(\mathbb{L}; Q)$ such that $\beta(g(c))\beta(g(\alpha(v)))|\beta(g(\alpha(U \setminus \{u\}))))$. Let $h = \beta \circ g \circ \alpha$. Since g behaves as $\tilde{\operatorname{rr}}_c$ on A, it follows from Lemma 60 that g preserves Qon A, therefore g preserves Q on $\alpha(\{v\} \cup U \setminus \{u\})$. This implies that h preserves Q on $\{v\} \cup U \setminus \{u\}$. Since $v|U \setminus \{u\}, h(v)|h(U \setminus \{u\})$, and h preserves Q on $\{v\} \cup U \setminus \{u\}$, it follows from Lemma 23 that h preserves C on $U \setminus \{u\}$. Since $h(u) = \beta(g(c))$, we have that $h(u)|h(U \setminus \{u\})$. It follows that h behaves as cut_u on U.

6.2.4 Classification of behaviors with one constant

The main result of this section is Lemma 64 below, which can be seen as a classification of canonical functions from $(\mathbb{L}; C, \prec, c)$ to $(\mathbb{L}; C, \prec)$. We first need two lemmata about *c*-universal sets.

Lemma 62. Choose $c \in \mathbb{L}$ and let $A \subseteq \mathbb{L} \setminus \{c\}$ be a *c*-universal set. Then for every finite subset X of \mathbb{L} there is an $\alpha \in \operatorname{Aut}(\mathbb{L}; C, \prec)$ such that $\alpha(X) \subseteq A$ and $\alpha(X)|c$.

Proof. Recall that the class of all ordered leaf structures has the ordering property (Theorem 38). By the formulation of the ordering property from Proposition 37, there exists a finite subset Y of \mathbb{L} such that for every $\beta \in \operatorname{Aut}(\mathbb{L}; C)$ there exists a $\alpha \in \operatorname{Aut}(\mathbb{L}; C, \prec)$ such that $\alpha(X) \subseteq \beta(Y)$. Let $y \in \mathbb{L}$ be such that y|Y holds. Since A is c-universal, there exists a $\gamma \in \operatorname{Aut}(\mathbb{L}; C)$ such that $\gamma(y) = c$ and $\gamma(Y) \subseteq A$. By the choice of Y there exists an $\alpha \in \operatorname{Aut}(\mathbb{L}; C, \prec)$ such that $\alpha(X) \subseteq \gamma(Y) \subseteq A$. Since $\gamma(X)|c$, we have $\alpha(X)|c$ which concludes the proof. \Box

Lemma 63. Let $c \in \mathbb{L}$, and let A be a subset of \mathbb{L} such that $A \subseteq \{x \in \mathbb{L} : x \prec c\}$ or $A \subseteq \{x \in \mathbb{L} : c \prec x\}$, and A is c-universal. Let $e : \mathbb{L} \to \mathbb{L}$ be an injective function that is canonical as a function from $(\mathbb{L}; C, \prec, c)$ to $(\mathbb{L}; C)$ and preserves C on $A \cap S_a^c$ for every $a \in A$. Then

- for every $a \in A$ we have $e(c)|e(A \cap S_a^c)$, and
- for every $a, b \in A$ we have either $S_a^c = S_b^c$ or $e(A \cap S_a^c) | e(A \cap S_b^c)$.

Proof. To prove the first assertion of the lemma, it suffices to prove that for arbitrary $x, y \in A$ satisfying xy|c we have e(x)e(y)|e(c). Assume for contradiction that there are $x_1, x_2 \in A$ such that $x_1x_2|c$ and $e(x_1)|e(x_2)e(c)$. Let $x_3 \in \mathbb{L}$ be such that $x_1 \neq x_3 \wedge x_1x_3|x_2$ holds. By Lemma 62 there is an $\alpha \in \operatorname{Aut}(\mathbb{L}; C, \prec)$ such that $\alpha(\{x_1, x_2, x_3\}) \subseteq A$ and $\alpha(\{x_1, x_2, x_3\})|c$. For $i \in \{1, 2, 3\}$ let $x'_i := \alpha(x_i)$. The pairs $(x_1, x_2), (x'_1, x'_2)$, and (x'_3, x'_2) are in the same orbit of $\operatorname{Aut}(\mathbb{L}; C, \prec, c)$: we have $x_1 \prec x_2$ if and only if $x'_1 \prec x'_2$ since α preserves \prec . Further, $x'_1 \prec x'_2$ if and only if $x'_3 \prec x'_2$ by convexity of \prec since $x_1x_3|x_2$ holds and α preserves C. Moreover, it holds that $x_1x_2|c$ by assumption, and $x'_1x'_2|c$ and $x'_3x'_2|c$ by the properties of α . In case that $A \subseteq \{x \in \mathbb{L} : x \prec c\}$ we have $x_1, x_2, x_3, x'_1, x'_2, x'_3 \prec c$, otherwise we have $c \prec x_1, x_2, x_3, x'_1, x'_2, x'_3$. By the homogeneity of $(\mathbb{L}; C, \prec, c)$.

By the canonicity of e, we have that $e(x'_1)|e(x'_2)e(c)$ and $e(x'_3)|e(x'_2)e(c)$. Since e preserves C on $S_x \cap A$, we have $e(x'_1)e(x'_3)|e(x'_2)$ which implies that $e(x'_1)e(x'_3)|e(x'_2)e(c)$. Since $x'_1x'_2x'_3|c$ and x'_1, x'_2, x'_3 are pairwise distinct, it follows that either (x'_1, x'_3) or (x'_3, x'_1) is in the same orbit as (x'_1, x'_2) in Aut($\mathbb{L}; C, \prec, c$). If (x'_1, x'_3) and (x'_1, x'_2) are in the same orbit in Aut($\mathbb{L}; C, \prec, c$), then by the canonicity of e, $(e(x'_1), e(x'_3), e(c))$ and $(e(x'_1), e(x'_2), e(c))$ are in the same orbit in Aut($\mathbb{L}; C$). It is impossible since $e(x'_1)e(x'_3)|e(x'_2)e(c)$. The case (x'_3, x'_1) and (x'_1, x'_2) are in the same orbit of Aut($\mathbb{L}; C, \prec, c$) is proved similarly. The first assertion of the lemma therefore follows.

It remains to show the second assertion of the lemma. We consider the case $A \subseteq \{x \in \mathbb{L} : x \prec c\}$. The case $A \subseteq \{x \in \mathbb{L} : c \prec x\}$ is argued similarly. We need the two following claims.

Claim 1. For any $x_1, x_2, x_3 \in A$ satisfying $x_1|x_2x_3c$ and $x_2x_3|c$ we have $e(x_1)|e(x_2)e(x_3)$.

Proof of Claim 1. For a contradiction we assume that $e(x_1)e(x_2)|e(x_3)$. Let $y_1, \ldots, y_5 \in A$ be pairwise distinct such that $y_1|y_2y_3y_4y_5c$, $y_2y_3y_4y_5|c$, and $y_2y_3|y_4y_5$. It follows from the convexity of \prec that $y_1 \prec y_i$ for $i \in \{2, \ldots, 5\}$. Since $y_2y_3|y_4y_5$, we have either $y_i \prec y_j$ for $i \in \{2, 3\}$ and $j \in \{4, 5\}$, or $y_i \prec y_j$ for $i \in \{4, 5\}$ and $j \in \{2, 3\}$. Without loss of generality we assume that $y_2 \prec y_3 \prec y_4 \prec y_5$. If $x_2 \prec x_3$, then the tuples (x_1, x_2, x_3) , (y_1, y_2, y_4) , and (y_1, y_4, y_5) are in the same orbit of Aut($\mathbb{L}; C, \prec, c$). Thus $e(y_1)e(y_2)|e(y_4)$ and $e(y_1)e(y_4)|e(y_5)$. Since e preserves C on $A \cap S_{y_2}^c$ and since $\{y_2, y_3, y_4, y_5\} \subseteq A \cap S_{y_2}^c$, we have $e(y_2)|e(y_4)e(y_5)$. These conditions are impossible to satisfy over $(\mathbb{L}; C)$. If $x_3 \prec x_2$ then we consider the three tuples (x_1, x_3, x_2) , (y_1, y_2, y_3) , and (y_1, y_3, y_4) that lie in the same orbit of Aut($\mathbb{L}; C, \prec, c$) and proceed analogously. \Box **Claim 2.** For any $x_1, x_2, x_3 \in A$ satisfying $x_1x_2|x_3c$ we have $e(x_1)e(x_2)|e(x_3)$.

This claim can be shown similarly as for the claim above by choosing five distinct elements y_1, \ldots, y_5 such that $y_1y_2y_3y_4|y_5c$ and $y_1y_2|y_3y_4$.

Let $a, b \in A$ such that $S_a^c \neq S_b^c$. This condition implies that ca|b or cb|a. We consider the case ca|b. The case cb|a is argued similarly. By the definition of cones, we have $S_a^c|S_b^c$, thus $A \cap S_a^c|A \cap S_b^c$. Let $x, y \in A \cap S_a^c$ and $z, t \in A \cap S_b^c$ be arbitrary. It follows from $ca|b \wedge xya|c \wedge ztb|c$ that $z|xyc \wedge xy|c \wedge t|xyc \wedge xy|c \wedge zt|xc \wedge zt|yc$. It follows from Claim 1 that $e(z)|e(x)e(y) \wedge e(t)|e(x)e(y)$, and it follows from Claim 2 that $e(z)e(t)|e(x) \wedge e(z)e(t)|e(y)$. Thus e(x)e(y)|e(z)e(t). The second assertion follows.

Lemma 64. Let $c \in \mathbb{L}$, and let A be a subset of \mathbb{L} such that $A \subseteq \{x \in \mathbb{L} : x \prec c\}$ or $A \subseteq \{x \in \mathbb{L} : c \prec x\}$, and A is c-universal. Let $e : \mathbb{L} \to \mathbb{L}$ be an injective function that is canonical on A as a function from $(\mathbb{L}; C, \prec, c)$ to $(\mathbb{L}; C, \prec)$. Then $\{e\} \cup \operatorname{Aut}(\mathbb{L}; C)$ generates lin or e behaves on A as id_c or rer_c .

Proof. We consider the case $A \subseteq \{x \in \mathbb{L} : x \prec c\}$. The case $A \subseteq \{x \in \mathbb{L} : c \prec x\}$ is argued similarly. The canonicity of e implies that either

- $e(x) \prec e(y)$ for all $x, y \in A$ such that xy|c and $x \prec y$, or
- $e(y) \prec e(x)$ for all $x, y \in A$ such that xy|c and $x \prec y$.

If the second case applies, we continue the proof with $-\circ e$ instead of e. Thus we assume in the following that the first case applies.

Since A is c-universal, there is an $a \in A$ such that $S_a^c \cap A$ contains four distinct elements x, y, u, v satisfying xy|uv. The function e is canonical on $S_a^c \cap A$ as a function from $(\mathbb{L}; C, \prec)$ to $(\mathbb{L}; C, \prec)$. Lemma 44 shows that e behaves as id, lin, or nil on $S_a^c \cap A$. The canonicity of e implies that e has the same behavior on all sets of the form $S_x^c \cap A$ for $x \in A$.

If e behaves as lin on all those sets, then we show that $\operatorname{Aut}(\mathbb{L}; C) \cup \{e\}$ generates lin. Let X be a finite subset of \mathbb{L} . By Lemma 62 there is an $\alpha \in \operatorname{Aut}(\mathbb{L}; C, \prec)$ such that $\alpha(X) \subseteq A$ and $\alpha(X)|c$. Since e behaves as lin on $\alpha(X)$ and α preserves \prec , the function $e \circ \alpha$ behaves as lin on X. This implies that $\{e\} \cup \operatorname{Aut}(\mathbb{L}; C)$ generates lin.

If e behaves as nil on all sets of the form $S_x^c \cap A$ for $x \in A$, then by the same argument it can be shown that $\{e\} \cup \operatorname{Aut}(\mathbb{L}; C)$ generates nil, and therefore lin by Proposition 43. We therefore assume in the following that e preserves C on each set of the form $S_x^c \cap A$, $x \in A$. Lemma 63 implies that the preconditions of the behaviors are satisfied (Definition 51). Let $u, v \in A$ be such that u|vc. Clearly, we have $u \prec v$. By C8, the following cases are exhaustive.

- 1. e(u)|e(v)e(c). Let $x, y, z \in A$ be such that x|yzc and y|zc. Clearly, we have $x \prec y \prec z \prec c$. Since (x, y) and (y, z) are in the same orbit of $\operatorname{Aut}(\mathbb{L}; C, \prec, c)$ as (u, v), we have e(x)|e(y)e(c) and e(y)|e(z)e(c) implying that e(x)|e(y)e(z)e(c). Thus, e behaves as id_c on A.
- 2. e(v)|e(u)e(c). Let $x, y, z \in A$ be such that x|yzc and y|zc. Since (x, y) and (y, z) are in the same orbit as (u, v) in $\operatorname{Aut}(\mathbb{L}; C, \prec, c)$, we have e(y)|e(x)e(c) and e(z)|e(y)e(c), and therefore e(z)|e(x)e(y)e(c). Thus, e behaves as rer_c on A, and $\{e\} \cup \operatorname{Aut}(\mathbb{L}; C)$ generates lin by Lemma 61.
- 3. e(u)e(v)|e(c). Canonicity of e on A implies that for any $x, y \in A$ we have e(x)e(y)|e(c) and thus e(A)|e(c). Let $a_1, a_2, a_3 \in A$ be three distinct elements such that $a_1|a_2a_3c$ and $a_2|a_3c$. By the convexity of \prec we have $a_1 \prec a_2 \prec a_3$ so (a_1, a_2) and (a_2, a_3) are in the same orbit of Aut($\mathbb{L}; C, \prec$) as (u, v). The canonicity of e implies that either $e(a_1) \prec e(a_2) \prec e(a_3)$ or $e(a_3) \prec e(a_2) \prec e(a_1)$ holds. It follows from the convexity of \prec that either $e(a_1)|e(a_2)e(a_3)$ or $e(a_1)e(a_2)|e(a_3)$ holds. If the first case holds then e behaves as cut_c , and if the second case holds then e behaves as rer_c on A.

We conclude that unless $\{e\} \cup \operatorname{Aut}(\mathbb{L}; C)$ generates lin, it behaves as id_c or rer_c on A.

6.3 Canonical behavior with two constants

In this section we analyze canonical functions from $(\mathbb{L}; C; \prec, c_1, c_2)$ to $(\mathbb{L}; C; \prec)$. For our purposes, it suffices to treat some special behaviors (Lemma 66); the motivation for those behaviors will become clear in the proof of Proposition 69. Then, we prove that certain behaviors of f imply that $\{f\} \cup \operatorname{Aut}(\mathbb{L}; C)$ generates lin (Lemma 67 and Lemma 68).

Definition 65. Let $c_1, c_2 \in \mathbb{L}$ be distinct. Then $A \subset \mathbb{L} \setminus \{c_1, c_2\}$ is called (c_1, c_2) -universal if for every finite $U \subset \mathbb{L}$ and $u_1, u_2 \in U$ there is an $\alpha \in \operatorname{Aut}(\mathbb{L}; C)$ such that $\alpha(U) \subseteq A \cup \{u_1, u_2\}$, $\alpha u_1 = c_1$, and $\alpha u_2 = c_2$.

Note that when A is (c_1, c_2) -universal then this implies in particular that $\{x \in A : xc_1|c_2\}$ and $\{x \in A : x|c_1c_2\}$ are c_1 -universal.

Lemma 66. Let $c_1, c_2 \in \mathbb{L}$ be distinct, and let A be (c_1, c_2) -universal such that all elements in $A_1 := \{x \in A : xc_1|c_2\}$ are in the same orbit in $(\mathbb{L}; C; \prec, c_1, c_2)$, and all elements in $A_2 := \{x \in A : x|c_1c_2\}$ are in the same orbit in $(\mathbb{L}; C; \prec, c_1, c_2)$. Let $f : \mathbb{L} \to \mathbb{L}$ be canonical on A as a function from $(\mathbb{L}; C; \prec, c_1, c_2)$ to $(\mathbb{L}; C; \prec)$. Then $\{f\} \cup \operatorname{Aut}(\mathbb{L}; C)$ generates lin or $\operatorname{End}(\mathbb{L}; Q)$, or f preserves C on $\{c_1\} \cup A_1 \cup A_2$.

Proof. It follows from the assumption on A_1 and A_2 that f is canonical on A_1 and A_2 as a function from $(\mathbb{L}; C, \prec, c_1)$ to $(\mathbb{L}; C, \prec)$. By Lemma 64, if f does not preserve C on $A_1 \cup \{c_1\}$ then $\{f\} \cup \operatorname{Aut}(\mathbb{L}; C)$ generates lin and we are done, or it behaves as rer_{c_1} on $A_1 \cup \{c_1\}$ in which case $\{f\} \cup \operatorname{Aut}(\mathbb{L}; C)$ generates $\operatorname{End}(\mathbb{L}; Q)$ by Corollary 57, and we are again done. The same argument applies if f does not preserve C on $A_2 \cup \{c_2\}$.

Thus, it remains to consider the case when f preserves C on both $A_1 \cup \{c_1\}$ and on $A_2 \cup \{c_1\}$. Let $a_1 \in A_1$ and $a_2 \in A_2$. We distinguish the following cases:

- $f(c_1)|f(a_1)f(a_2)$. It follows from the canonicity of f that $f(c_1)|f(a_1)f(x)$ for all $x \in A_2$ so $f(c_1)|f(A_2)$. This is impossible since f preserves C on $\{c_1\} \cup A_2$.
- $f(a_2)f(c_1)|f(a_1)$. It follows from the canonicity of f that $f(x)f(c_1)|f(a_1)$ for all $x \in A_2$, therefore $f(a_1)|f(A_2)$. This implies that f behaves as cut_{a_1} on A_2 . Since $c_1a_1|x$ for all $x \in A_2$ and A_2 is c_1 -universal, we have that A_2 is also a_1 -universal, and, by Corollary 53, $\{f\} \cup \operatorname{Aut}(\mathbb{L}; C)$ generates lin.
- $f(c_1)f(a_1)|f(a_2)$. It follows from the canonicity of f that f preserves C on $\{c_1\} \cup A_1 \cup A_2$, and we are done.

Since these three cases are exhaustive, the statement follows.

Lemma 67. Let $c_1, c_2 \in \mathbb{L}$ and $A \subseteq \mathbb{L} \setminus \{c_1, c_2\}$ be (c_1, c_2) -universal. Let $A_1 = \{x \in A : xc_1|c_2\}$, $A_2 = \{x \in A : c_1|xc_2\}$, and $A_3 = \{x \in A : c_1c_2|x\}$. Let $g \colon \mathbb{L} \to \mathbb{L}$ be an injection such that

- $g(A_1 \cup \{c_1\})|g(c_2),$
- g preserves C on $\{c_1\} \cup A_1$ and on $\{c_2\} \cup A_2 \cup A_3$, and
- $g(c_1)g(c_2)|g(x)$ for every $x \in A_2 \cup A_3$.

Then $\{g\} \cup \operatorname{Aut}(\mathbb{L}; C)$ generates nil.

Proof. We need to show that for all k and all $x_1, \ldots, x_k \in \mathbb{L}$ there is an $f \in M := \langle \operatorname{Aut}(\mathbb{L}; C) \cup \{g\} \rangle$ such that $f(x_j) = \operatorname{nil}(x_j)$ for all $j \leq k$. This is clearly true for $k \leq 2$. To prove it for $k \geq 3$, suppose without loss of generality that $x_1 \prec \cdots \prec x_k$. We first prove by induction on $i \in \{1, \ldots, k-1\}$ that there exists an $h \in M$ with the following properties.

- 1. $h(x_1) \cdots h(x_i) | h(x_j)$ for every $j \in \{i+1, \ldots, k\}$
- 2. h preserves C on $\{x_i, \ldots, x_k\}$
- 3. for $i \ge 2$ we additionally require that $h(x_1) \cdots h(x_j) | h(x_{j+1})$ for every $j \in \{1, \ldots, i-1\}$

For i = 1 the identity function has the properties that we require for $h \in M$. For $i \ge 2$, we inductively assume the existence of a function $h' \in M$ such that

• $h'(x_1)\cdots h'(x_{i-1})|h'(x_j)$ for every $j \in \{i,\ldots,k\}$,

- h' preserves C on $\{x_{i-1}, \ldots, x_k\}$, and
- if $i \ge 3$ we additionally have $h(x_1) \cdots h(x_j) | h(x_{j+1})$ for every $j \in \{1, \dots, i-2\}$.

By (c_1, c_2) -universality of A, there exists an $\alpha \in \operatorname{Aut}(\mathbb{L}; C)$ that maps $h'(x_1)$ to $c_1, h'(x_i)$ to c_2 , and such that $\alpha h'(\{x_1, \ldots, x_k\}) \subseteq A \cup \{c_1, c_2\}$.

Observation. $\alpha h'(\{x_1, ..., x_{i-1}\}) \subseteq \{c_1\} \cup A_1 \text{ and } \alpha h'(\{x_i, ..., x_k\}) \subseteq \{c_2\} \cup A_2 \cup A_3.$

Proof of the observation. The first property of h' implies that $h'(x_1)h'(x_{i-1})|h'(x_i)$. Therefore, $\alpha h'(x_1)\alpha h'(x_{i-1})|\alpha h'(x_i)$ and $c_1x|c_2$ for every $x \in \alpha h'(\{x_1,\ldots,x_{i-1}\})$ which concludes the proof of the first part of the observation.

To show the second part, arbitrarily choose $j \in \{i, \ldots, k\}$. If j = i then $\alpha h'(x_j) = \alpha h'(x_i) = c_2$ and there is nothing to show. Since $x_{i-1} \prec x_i \prec x_j$, we distinguish the cases that $x_{i-1}|x_ix_j$ and $x_{i-1}x_i|x_j$. By the inductive assumption, h' preserves C on $\{x_{i-1}, \ldots, x_k\}$ so we have

$$h'(x_{i-1})|h'(x_i)h'(x_j)$$
 or $h'(x_{i-1})h'(x_i)|h'(x_j)$.

First consider the case $h'(x_{i-1})|h'(x_i)h'(x_j)$. By the first property of h', we also have

$$h'(x_1)h'(x_{i-1})|h'(x_i)$$
 and $h'(x_1)h'(x_{i-1})|h'(x_i)h'(x_i)|$

Consequently, $\alpha h'(x_1)|\alpha h'(x_i)\alpha h'(x_j)$, and thus $c_1|c_2\alpha h'(x_j)$. Hence $\alpha h'(x_j) \in A_2$. Now consider the case $h'(x_{i-1})h'(x_i)|h'(x_j)$. Since $h'(x_1)h'(x_{i-1})|h'(x_j)$, we have $h'(x_1)h'(x_i)|h'(x_j)$. Thus $c_1c_2|\alpha h'(x_j)$, and $\alpha h'(x_j) \in A_3$.

We claim that $h := g \circ \alpha \circ h'$ satisfies the inductive claim so we have to verify the three properties from the inductive statement.

- Ad 1. By the observation above with the facts that $\alpha h'(x_i) = c_2$ and $\alpha h'$ is injective, it follows that $\alpha h'(\{x_{i+1}, \ldots, x_k\}) \subseteq A_2 \cup A_3$. Since $g(c_1)g(c_2)|g(x)$ for every $x \in A_2 \cup A_3$ and $g(A_1 \cup \{c_1\})|g(c_2)$, we have $g(\{c_1, c_2\} \cup A_1)|g(x)$ for every $x \in A_2 \cup A_3$. Therefore, $(g \circ \alpha \circ h')(\{x_1, \ldots, x_i\})|(g \circ \alpha \circ h')(x_j)$ for every $j \in \{i+1, \ldots, k\}$, or, equivalently, $h(x_1) \cdots h(x_i)|h(x_i)$, which is what we had to show.
- Ad 2. By the second property of h', the restriction of h' to $\{x_{i-1}, \ldots, x_k\}$ preserves C. Since $\alpha h'(\{x_i, \ldots, x_k\}) \subseteq \{c_2\} \cup A_2 \cup A_3$ and g preserves C over $\{c_2\} \cup A_2 \cup A_3$, the restriction of $h = g \circ \alpha \circ h'$ to $\{x_i, \ldots, x_k\}$ preserves C as well.
- Ad 3. We assume that $i \geq 3$ since otherwise there is nothing to show. Since g preserves C over $A_1 \cup \{c_1\}$ and $\alpha h'(\{x_1, \ldots, x_{i-1}) \subseteq A_1 \cup \{c_1\}$, the third property of h' implies that $g \circ \alpha \circ h'(\{x_1, \ldots, x_j\})|g \circ \alpha \circ h'(x_{j+1})$ for all $j \in \{1, \ldots, i-2\}$. Equivalently, $h(x_1) \cdots h(x_j)|h(x_{j+1})$ for all $j \in \{1, \ldots, i-2\}$. It remains to show that $h(x_{i-2})h(x_{i-1})|h(x_i)$. This follows directly from the fact that $g(A_1 \cup \{c_1\})|g(c_2)$ and $\alpha h'(x_i) = c_2$.

This concludes the induction. For i = k the third property of h implies that $h(x_1) \cdots h(x_j) | h(x_{j+1})$ for all $j \in \{1, \ldots, k-1\}$. This property and the homogeneity of $(\mathbb{L}; C)$ imply the existence of $\beta \in \operatorname{Aut}(\mathbb{L}; C)$ such that $\beta h(x) = \operatorname{nil}(x)$ for all $x \in X$, and hence $f := \beta \circ h \in M$ is a function with the desired properties. \Box

Lemma 68. Let $c_1, c_2 \in \mathbb{L}$ and $A \subseteq \mathbb{L} \setminus \{c_1, c_2\}$ be (c_1, c_2) -universal. Let $A_1 = \{x \in A : xc_1|c_2\}$, $A_2 = \{x \in A : c_1|xc_2\}$, and $A_3 = \{x \in A : c_1c_2|x\}$. Let $g \colon \mathbb{L} \to \mathbb{L}$ be an injection such that

- for all $a_1 \in A_1$, $a_2 \in A_2$ we either have $g(c_1)g(a_1)|g(a_2)$ or $g(c_1)g(a_2)|g(a_1)$;
- g preserves C on $\{c_1\} \cup A_1 \cup A_3$ and on $\{c_2\} \cup A_2 \cup A_3$, and
- $g(c_1)g(c_2)|g(x)$ for every $x \in A$.

Then $\{g\} \cup \operatorname{Aut}(\mathbb{L}; C)$ generates lin.

Proof. We first show that $\{g\} \cup \operatorname{Aut}(\mathbb{L}; C)$ generates a function f with the property that there are no $a, b, c, d \in \mathbb{L}$ such that f(a)f(b)|f(c)f(d). For this, it suffices by a standard application of König's tree lemma (see e.g. Section 3.1 in [6]) to show that for all finite $S = \{x_1, \ldots, x_k\} \subset \mathbb{L}$ there is an $h \in M := \langle \operatorname{Aut}(\mathbb{L}; C) \cup \{g\} \rangle$ such that there are no $a, b, c, d \in S$ with h(a)h(b)|h(c)h(d).

This is clearly true for $k \leq 1$. To prove it for $k \geq 2$, we prove by induction on $i \in \{1, \ldots, k-1\}$ that there exists an $h \in M$ with the following property.

 $P_{h,i}$ The equivalence relation \sim_h defined on $\{x_2, \ldots, x_k\}$ by $u \sim_h v$ iff $h(x_1)|h(u)h(v)$ has at least *i* equivalence classes.

For i = 1 the statement is trivial and we let $h \in M$ be the identity function. For $i \geq 2$, we inductively assume the existence of a function $h' \in M$ satisfying $P_{h',i-1}$. If $P_{h',i}$ holds, then there is nothing to be shown so we assume that there are distinct $p, q \leq k$ such that $h'(x_1)|h'(x_p)h'(x_q)$. By (c_1, c_2) -universality of A, there exists an $\alpha \in \operatorname{Aut}(\mathbb{L}; C)$ that maps $h'(x_1)$ to $c_1, h'(x_p)$ to c_2 , and such that $\alpha h(\{x_1, \ldots, x_k\}) \subseteq A \cup \{c_1, c_2\}$. We claim that $h := g \circ \alpha \circ h'$ satisfies $P_{h,i}$. To show that, we first prove that \sim_h has at least as many equivalence classes as $\sim_{h'}$.

Observe that when $r, s \in \{2, \ldots, k\}$ are such that $x_r \not\sim_{h'} x_s$, then $x_r \not\sim_h x_s$: when both $\alpha(h'(x_r)), \alpha(h'(x_s)) \in A_1 \cup A_3$ then this follows the assumption that g preserves C on $\{c_1\} \cup A_1 \cup A_3$; a similar argument applies when both $\alpha(h'(x_r)), \alpha(h'(x_s)) \in A_2 \cup A_3$. When $\alpha(h'(x_r)) \in A_1$ and $\alpha(h'(x_s)) \in A_2$ then either $g(c_1)g(\alpha(h'(x_r)))|g(\alpha(h'(x_s)))$ or $g(c_1)g(\alpha(h'(x_s)))|g(\alpha(h'(x_r)))|$, so $x_r \not\sim_h x_s$.

Next, consider the case that one of r, s, say r, equals p, that is, $\alpha(h'(x_r)) = c_2$. In this case we have by the third assumption on g in the statement of the lemma that $g(\alpha(h'(x_1)))g(\alpha(h'(x_r)))|g(x))$ for all $x \in A$, and in particular that $g(\alpha(h'(x_1)))g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g(\alpha(h'(x_r)))|g$

To see that \sim_h has *strictly* more equivalence classes than $\sim_{h'}$, observe that $x_p \sim_{h'} x_q$ but $x_p \nsim_h x_q$ as we have just seen. This concludes the inductive proof.

Note that $P_{h,k-1}$ implies that for all $p < q \le k$ we have $h(x_p)h(x_1)|h(x_q)$ or $h(x_q)h(x_1)|h(x_p)$, and in particular there cannot be $a, b, c, d \in S$ such that h(a)h(b)|h(c)h(d). This concludes our proof of the existence of f.

Since $(\mathbb{L}; C, \prec)$ is homogeneous, Ramsey, and ω -categorical, Theorem 34 asserts the existence of a function $f' \in \overline{\{\alpha_1 f \alpha_2 \mid \alpha_1, \alpha_2 \in \operatorname{Aut}(\mathbb{L}; C, \prec)\}}$ which is canonical as a function from $(\mathbb{L}; C, \prec)$ to $(\mathbb{L}; C)$. Clearly, f' also has the property that there are no $a, b, c, d \in \mathbb{L}$ such that f'(a)f'(b)|f'(c)f'(d), and the behavior of f' is either lin or nil by Lemma 44. In both cases, $\{f'\} \cup \operatorname{Aut}(\mathbb{L}; C)$ generates lin by Proposition 43.

6.4 Proof of the main result

In this section we finish the proof of Theorem 1 and Corollary 3. We begin by proving two auxiliary results in Propositions 69 and 71.

Proposition 69. Let Γ be a reduct of $(\mathbb{L}; C)$. Then one of the following applies.

- 1. $\operatorname{End}(\Gamma) = \operatorname{End}(\mathbb{L}; C);$
- 2. End(Γ) contains a constant operation;
- 3. End(Γ) contains lin;
- 4. End(Γ) contains End($\mathbb{L}; Q$).

Proof. If Γ has a non-injective endomorphism, then Γ also has a constant endomorphism by Lemma 39 and the second item of the statement of the proposition applies. Therefore we suppose in the following that all endomorphisms are injective. If all endomorphisms preserve C, then the first item applies and we are done. Hence, suppose that there is an injective endomorphism e that violates the rooted triple relation, that is, there are c_1, c_2, c_3 such that $c_1|c_2c_3$ and not $e(c_1)|e(c_2)e(c_3)$. Under this assumption, we claim that there are $d_1, d_2, d_3 \in \mathbb{L}$ such that $d_1|d_2d_3$ and $e(d_1)e(d_2)|e(d_3)$. By injectivity of *e*, we either have $e(c_1)e(c_3)|e(c_2)$ or $e(c_1)e(c_2)|e(c_3)$. In the first case, choose $(d_1, d_2, d_3) := (c_1, c_3, c_2)$ and in the second case choose $(d_1, d_2, d_3) := (c_1, c_2, c_3)$.

By convexity of \prec we have either $d_1 \prec d_2 \prec d_3$, $d_1 \prec d_3 \prec d_2$, $d_2 \prec d_3 \prec d_1$, or $d_3 \prec d_2 \prec d_1$. In each case, by the homogeneity of (L; C), there exists an $\alpha \in \operatorname{Aut}(\mathbb{L}; C)$ such that $\alpha d_1 \prec \alpha d_2 \prec \alpha d_3$. After replacing d_1, d_2, d_3 by $\alpha d_1, \alpha d_2, \alpha d_3$ and e by $x \mapsto e(\alpha^{-1}x)$, we still have $d_1|d_2d_3$ and $e(d_1)e(d_2)|e(d_3)$. So we assume in the following that $d_1 \prec d_2 \prec d_3$. There also exists $\beta \in \operatorname{Aut}(\mathbb{L}; C)$ such that $\beta(e(d_1)) \prec \beta(e(d_2)) \prec \beta(e(d_3))$. By replacing e with the function $x \mapsto \beta e(x)$, we may henceforth assume that $e(d_1) \prec e(d_2) \prec e(d_3)$.

Recall our strategy described at the beginning of this section: we explained that one can additionally assume (by Corollary 35) that e is canonical as a function from $(\mathbb{L}; C, \prec, d_1, d_2, d_3)$ to $(\mathbb{L}; C, \prec)$. Define

$$A_{1} := \{a : a \prec d_{1}, ad_{1}|d_{2}d_{3}\}$$
$$A_{2} := \{a : d_{1} \prec a \prec d_{2}, d_{1}|ad_{2}d_{3} \land a|d_{2}d_{3}\}$$
$$A_{3} := \{a : a \prec d_{1}, a|d_{1}d_{2}d_{3}\}$$

Note that $A := A_1 \cup A_2 \cup A_3$ is (d_1, d_2) -universal: for every finite $X \subseteq \mathbb{L}$ and arbitrary $x_1, x_2 \in X$ there exists an $\alpha \in Aut(\mathbb{L}; C)$ such that

- $\alpha x_1 = d_1$ and $\alpha x_2 = d_2$,
- $\{\alpha x : x \in X \setminus \{x_1, x_2\}, xx_1 | x_2\} \subseteq A_1,$
- $\{\alpha x : x \in X \setminus \{x_1, x_2\}, x_1 | x_2 x\} \subseteq A_2$, and
- $\{\alpha x : x \in X \setminus \{x_1, x_2\}, x | x_1 x_2\} \subseteq A_3.$

We observe that if A_i is d_j -universal, for $1 \leq i \leq 3$ and $1 \leq j \leq 3$, and e is canonical on A_i as a function from $(\mathbb{L}; C, \prec, d_j)$ to $(\mathbb{L}; C, \prec)$, then Lemma 64 implies that e behaves as id_{d_j} or rer_{d_j} on A_i unless $\{e\} \cup \mathrm{Aut}(\mathbb{L}; C)$ generates lin. If $\{e\} \cup \mathrm{Aut}(\mathbb{L}; C)$ generates lin then the third item of the statement of the proposition holds and we are done. If e behaves as rer_{d_j} on A_i then Corollary 57 implies that $\{e\} \cup \mathrm{Aut}(\mathbb{L}; C)$ generates $\mathrm{End}(\mathbb{L}; Q)$; in this case the fourth item of the statement holds. Therefore we assume in the following that e behaves as id_{d_j} on A_i . Note that this assumption implies that e preserves C on $\{d_j\} \cup A_i$.

Now, pick $r \in A_2$ arbitrarily. By the injectivity of e, the following cases are exhaustive.

- $e(d_1)e(d_2)|e(r)e(d_3)$. This is in contradiction with the assumption that e behaves as id_{d_2} on A_2 . To see this, choose an element $a \in A_2$ and note that $e(d_2)|e(a)e(d_3)$ by the canonicity of e on A_2 . This implies that $e(d_2)|e(A_2)$.
- $e(d_1)e(d_2)e(r)|e(d_3)$. This is in contradiction with the assumption that e behaves as id_{d_3} on A_2 . To see this, choose an element $a \in A_2$ and note that $e(d_2)e(a)|e(d_3)$ by the canonicity of e on A_2 . This implies that $e(d_3)|e(A_2)$.
- $e(d_1)e(d_2)e(d_3)|e(r)$. This is the remaining case that we will consider in the rest of the proof.

Lemma 66 applied to $f := e, c_1 := d_1, c_2 := d_2$, and A shows that e preserves C on $\{d_1\} \cup A_1 \cup A_3$, unless $\{e\} \cup \operatorname{Aut}(\mathbb{L}; C)$ generates lin or $\operatorname{End}(\mathbb{L}; Q)$. The same argument can be applied when we exchange d_2 with d_1 and A_2 with A_1 so we assume that e preserves C on $\{d_1\} \cup A_1 \cup A_3$ and on $\{d_2\} \cup A_2 \cup A_3$.

If there were a $u \in A_3$ such that $e(d_1)e(u)|e(d_3)$ or $e(d_1)|e(u)e(d_3)$ then e would not behave as id_{d_3} or id_{d_1} on A_3 since $e(d_3)|e(A_3)$ or $e(d_1)|e(A_3)$ by the canonicity of e, respectively. Hence, we have $e(d_1)e(d_2)e(d_3)|e(A_3)$. If there were a $u \in A_1$ such that $e(d_1)|e(u)e(d_2)$ then by the canonicity of e we would have $e(d_1)|e(A_1)$, and e would not behave as id_{d_1} on A_1 . Thus $e(u)e(d_1)|e(d_2)$ or $e(u)|e(d_1)e(d_2)$. This implies that either $e(u)e(d_1)|e(d_2)$ for all $u \in A_1$ or $e(u)|e(d_1)e(d_2)$ for all $u \in A_1$.

In the former case, we have $e(A_1 \cup \{d_1\})|e(d_2)$ and Lemma 67 applied to $c_1 = d_1$ and $c_2 = d_2$ shows that $\{e\} \cup \operatorname{Aut}(\mathbb{L}; C)$ generates nil, and therefore lin by Proposition 43.

In the latter case we show that the conditions in Lemma 68 are satisfied for $A, g := e, c_1 := d_1$, and $c_2 := d_2$. Clearly, the second and the third conditions are satisfied. It remains to show that the first condition is satisfied. Arbitrarily choose $a_1 \in A_1$ and $a_2 \in A_2$. If $e(d_1)|e(a_1)e(a_2)$ then for all $u \in A_1$ we have $e(d_1)|e(u)e(a_2)$ by the canonicity of e. This implies that $e(d_1)|e(A_1)$ which leads to a contradiction since e behaves as id_{d_1} on A_1 . Thus either $e(d_1)e(a_2)$ or $e(d_1)e(a_2)|e(a_1)|$ holds. Hence, Lemma 68 shows that $\{e\} \cup \operatorname{Aut}(\mathbb{L}; C)$ generates lin. \Box

Proposition 69 leaves us with the task of further analyzing the reducts of $(\mathbb{L}; Q)$. We first need the following lemma.

Lemma 70. Let $U \subset \mathbb{L}$ be finite and arbitrarily choose $c \in \mathbb{L} \setminus U$. Then there are $U_1, \ldots, U_k \subseteq U$ such that $U_1 \cup \cdots \cup U_k = U$ and $(\{c\} \cup \bigcup_{i=1}^{i-1} U_j) | U_i$ for all $i \leq k$.

Proof. By induction on the size of U. If $\{c\}|U$, then k := 1 and $U_1 := U$ satisfies the statement. Otherwise, $|U| \ge 2$, and by Lemma 9, there are two non-empty subsets V, W of U such that $V \cup W = U$ and V|W. We either have $(\{c\} \cup V)|W$ or $V|(W \cup \{c\})$. In the first case, we inductively have U_1, \ldots, U_{k-1} such that $U_1 \cup \cdots \cup U_{k-1} = V$ and $(\{c\} \cup \bigcup_{j=1}^{i-1} U_j)|U_i$ for all $i \le k-1$. Set $U_k := W$. Then $U_1, \ldots, U_{k-1}, U_k$ satisfy the requirements from the statement. The case when $V|(W \cup \{c\})$ can be shown analogously.

Proposition 71. Let Γ be a reduct of $(\mathbb{L}; Q)$. Then one of the following cases applies.

- 1. All endomorphisms of Γ preserve Q;
- 2. Γ has a constant endomorphism;
- 3. Γ is preserved by lin.

Proof. If all endomorphisms of Γ preserve Q, then we are in case one of the statement of the proposition; in the following we therefore assume that Γ has an endomorphism f that violates Q. We can then choose four elements $d_1, d_2, d_3, d_4 \in \mathbb{L}$ such that $d_1d_2 : d_3d_4$ and $f(d_1)f(d_3) : f(d_2)f(d_4)$. By the homogeneity of $(\mathbb{L}; Q)$ there are $\gamma, \delta \in \operatorname{Aut}(\mathbb{L}; Q)$ such that

- $\gamma(d_1) \prec \gamma(d_2) \prec \gamma(d_3) \prec \gamma(d_4),$
- $\gamma(d_1)\gamma(d_2)|\gamma(d_3)\gamma(d_4),$
- $\delta(f(d_1)) \prec \delta(f(d_3)) \prec \delta(f(d_2)) \prec \delta(f(d_4))$, and
- $\delta(f(d_1))\delta(f(d_3))|\delta(f(d_2))\delta(f(d_4)).$

(Here, the order \prec is still the order as defined in Section 3.5.) By replacing f by $\delta \circ f \circ \gamma^{-1}$, we can assume that $d_1 \prec d_2 \prec d_3 \prec d_4$, $d_1d_2|d_3d_4$, $f(d_1) \prec f(d_3) \prec f(d_2) \prec f(d_4)$, and $f(d_1)f(d_3)|f(d_2)f(d_4)$. Corollary 35 asserts the existence of a function

$$g \in \overline{\{\alpha_2 f \alpha_1 : \alpha_1 \in \operatorname{Aut}(\mathbb{L}; C, \prec, d_1, \dots, d_4), \alpha_2 \in \operatorname{Aut}(\mathbb{L}; C, \prec)\}}$$

which is canonical as a function from $(\mathbb{L}; C, \prec, d_1, \ldots, d_4)$ to $(\mathbb{L}; C, \prec)$. Note that there exists an $\alpha \in \operatorname{Aut}(\mathbb{L}; C, \prec)$ such that $g(d_i) = \alpha(f(d_i))$ for all $i \in \{1, \ldots, 4\}$, and, in particular, $g(d_1)g(d_2)g(d_2)g(d_4)$.

Let $S = \{x \in \mathbb{L} : d_1d_2|x \wedge d_1d_2x|d_3d_4 \wedge d_2 \prec x\}$. Note that S is d_1 -universal and d_2 -universal. By Lemma 64, either g behaves on S as id_{d_1} or rer_{d_1} , or $\{g\} \cup \mathrm{Aut}(\mathbb{L}; C)$ generates lin. Similarly, either g behaves on S as id_{d_2} or rer_{d_2} , or $\{g\} \cup \mathrm{Aut}(\mathbb{L}; C)$ generates lin. In the latter cases we are done, so assume that g behaves on S as id_{d_1} or rer_{d_1} , and as id_{d_2} or rer_{d_2} .

We then show that the conditions of Lemma 52 apply to $M := \langle \operatorname{Aut}(\mathbb{L}; Q) \cup \{g\} \rangle$. Let $U \subset \mathbb{L}$ be finite and arbitrarily choose $u \in U$. By Lemma 70 there exists a partition $U_1 \cup \cdots \cup U_k$ of $U \setminus \{u\}$ such that $(\{u\} \cup \bigcup_{i=1}^{i-1} U_j) | U_i$ for all $i \in \{1, \ldots, k\}$. By d_1 -universality of S, there are subsets



Figure 4: Illustration for the last proof step for Proposition 71.

 X_1, \ldots, X_k of S such that $X_i | (\bigcup_{j=i+1}^k X_j \cup \{d_1\})$ for all $i \in \{1, \ldots, k\}$, and $(X_i; C)$ is isomorphic to $(U_i; C)$. By the homogeneity of $(\mathbb{L}; Q)$ there is an $\alpha \in \operatorname{Aut}(\mathbb{L}; Q)$ such that $\alpha(u) = d_3$, $\alpha(U_i) = X_i$, and α preserves C on each U_i .

First consider the case that g behaves on S as rer_{d_1} . We claim that $g(d_3)|g(S)$ or $g(d_4)|g(S)$. Since $g(d_1)g(d_3)|g(d_2)g(d_4)$ and by the canonicity of g as a function from $(\mathbb{L}; C, \prec, d_1, \ldots, d_4)$ to $(\mathbb{L}; C, \prec)$, either

 $g(d_1)g(d_3)g(S)|g(d_2)g(d_4),g(d_1)g(d_3)|g(S)g(d_2)g(d_4), \text{ or } g(d_1)g(d_3)g(d_2)g(d_4)|g(S).$

In the first case $g(d_4)|g(S)$ holds while in the second and third case $g(d_3)|g(S)$ holds. We first consider the case $g(d_3)|g(S)$. Let $h := g \circ \alpha$. Clearly, h is in M. We will show that h behaves as cut_u on U. Since g behaves as rer_{d_1} on S and $X_i|(\bigcup_{j=i+1}^k X_j \cup \{d_1\})$ for all $i \in \{1, \ldots, k\}$, it follows from the definition of rer_{d_1} that $\bigcup_{j=1}^i g(X_j)|g(X_{i+1})$ for all $i \in \{1, \ldots, k-1\}$. It implies that $\bigcup_{j=1}^i h(U_j)|h(U_{i+1})$ for all $i \in \{1, \ldots, k-1\}$. Since α preserves C on each U_i and gpreserves C on each X_i , h preserves C on each U_i . Since $\bigcup_{j=1}^i U_j|U_{i+1}, \bigcup_{j=1}^i h(U_j)|h(U_{i+1})$ for all $i \in \{1, \ldots, k-1\}$ and h preserves C on each U_i , it follows that h preserves C on $\bigcup_{i=1}^k U_i$. Since $g(d_3)|g(S)$, we have that $h(u)|h(\bigcup_{i=1}^k U_i)$. Thus h behaves as cut_u on U. By Lemma 52, Γ is preserved by lin. The case $g(d_4)|g(S)$ can be treated similarly (by choosing $\alpha \in \operatorname{Aut}(\mathbb{L}; Q)$ such that $\alpha(u) = d_4$ instead of $\alpha(u) = d_3$).

Finally, we consider the case when g behaves on S as id_{d_1} . By the canonicity of g as a function from $(\mathbb{L}; C, \prec, d_1, \ldots, d_4)$ to $(\mathbb{L}; C, \prec)$ and since all the elements of S lie in the same orbit of $(\mathbb{L}; C, \prec, d_1, \ldots, d_4)$, either $g(d_1)|g(d_2)g(S)$, $g(S)g(d_1)|g(d_2)$, or $g(d_1)g(d_2)|g(x)$ for all $x \in S$. The first case is impossible because g behaves as id_{d_1} on S. The second case is impossible, too: to see this, pick $a, b, c \in S$ such that $d_1a|b$ and $d_1ab|c$. Since g behaves on S as id_{d_1} , we have $g(d_1)g(a)g(b)|g(c)$ and $g(d_1)g(a)|g(b)$, and $g(d_1)g(a)g(b)g(c)|g(d_2)$ by assumption. In case that g behaves as id_{d_2} we would have $g(d_2)g(a)|g(b)$ which is inconsistent with the above. In case that g behaves as rer_{d_2} we would have $g(d_2)g(a): g(b)g(c)$ which is inconsistent with the above, too.

In the third and last case, we first show that g does not behave as rer_{d_2} on S. Let $a, b, c \in S$ such that $d_1ab|c \wedge d_1a|b$. Since g behaves as id_{d_1} , we have $g(d_1)g(a)g(b)|g(c)$. By the assumption $g(d_1)g(d_2)|g(x)$ for all $x \in S$, we have $g(d_1)g(d_2)|g(c)$. It follows from $g(d_1)g(a)g(b)|g(c)$ that $g(d_2)g(a)g(b)|g(c)$. Therefore g does not behave as rer_{d_2} on S. Hence g behaves as id_{d_2} on S. Thus, $(\{g(d_1), g(d_2)\} \cup \bigcup_{j=i+1}^k g(X_j))|g(X_i)$ for all $i \in \{1, \ldots, k\}$. Since $g(d_1)g(d_2)g(d_2)g(d_4)$, we therefore must have that $(\{g(d_3)\} \cup \bigcup_{j=i+1}^k g(X_j))|g(X_i)$ for all $i \in \{1, \ldots, k\}$. Since $(\mathbb{L}; C)$ embeds all finite leaf structures, there are subsets Z_1, \ldots, Z_k of \mathbb{L} and $z \in \mathbb{L}$ such that $z|\bigcup_{j=1}^k Z_j$, $(\bigcup_{j=1}^{i-1} Z_j)|Z_i$, and $(Z_i; C)$ is isomorphic to $(g(X_i); C)$ for all $i \in \{1, \ldots, k\}$. The situation is illustrated in Figure 4. By the homogeneity of $(\mathbb{L}; Q)$ there is a $\beta \in \operatorname{Aut}(\mathbb{L}; Q)$ such that $\beta(g(d_3)) = z$, β preserves C on each $g(X_i)$, and $\beta(g(X_i)) = Z_i$ for all $i \in \{1, \ldots, k\}$. Then $\beta \circ g \circ \alpha$ is in M and behaves as cut_u on U. Again, Lemma 52 implies that Γ is preserved by lin.

We can now prove Theorem 1 and Corollary 3 that have already been stated in Section 2.

Proof of Theorem 1. Let Γ be a reduct of $(\mathbb{L}; C)$. We apply Proposition 69 and consider the following cases.

- All endomorphisms of Γ preserve C. Then $\operatorname{End}(\Gamma) \subseteq \operatorname{End}(\mathbb{L}; C)$; we claim that the opposite inclusion holds as well. Since $\operatorname{End}(\Gamma)$ is closed, it suffices to show that for every $e \in \operatorname{End}(\mathbb{L}; C)$ and every finite $S \subset \mathbb{L}$ there exists an $f \in \operatorname{End}(\Gamma)$ such that f(s) = e(s) for all $s \in S$. Since e preserves C, $e|_S$ is a partial isomorphism from (S; C) to (e(S); C) by Lemma 10. By homogeneity, $e|_S$ can be extended to an automorphism $f \in \operatorname{Aut}(\mathbb{L}; C)$. Since $\operatorname{Aut}(\mathbb{L}; C) \subseteq$ $\operatorname{End}(\Gamma)$, we have $f \in \operatorname{End}(\Gamma)$.
- Γ has a constant endomorphism. Then there is nothing to show since the second item of the statement applies.
- Γ is preserved by lin. Then Lemma 50 shows that Γ is homomorphically equivalent to a reduct of $(\mathbb{L}; =)$ and the third item of the statement applies.
- End(Γ) contains End($\mathbb{L}; Q$). This case implies that Γ is a reduct of ($\mathbb{L}; Q$). If Γ has a constant endomorphism or Γ is preserved by lin, then we are done as the above cases. Otherwise, by Proposition 71 all endomorphisms of Γ preserve Q. This means that End(Γ) \subseteq End(Q). Hence we have End(Γ) = End(Q), therefore the fourth item applies.

By Proposition 69, these four cases are exhaustive.

One may observe at this point that the proof of Theorem 1 does not rely on any results concerning Jordan permutation groups. We finally show that every reduct of Γ is existentially interdefinable with $(\mathbb{L}; C)$, with $(\mathbb{L}; Q)$, or with $(\mathbb{L}; =)$.

Proof of Corollary 3. Let Γ be a reduct of $(\mathbb{L}; C)$. Let Γ' be the expansion of Γ by the relations defined by negations of atomic formulas over Γ , including the equality relation (for example, when R is a ternary relation of Γ , the structure Γ' contains the binary relation defined by $\neg R(x, x, y)$). We apply Theorem 1 to Γ' . Since for every atomic formula ϕ over Γ' the signature of Γ' also contains a relation symbol for $\neg \phi$, all endomorphisms of Γ' must be embeddings, and therefore item 2 of Theorem 1 is impossible. If Γ' has the same endomorphisms as $(\mathbb{L}; C)$ or $(\mathbb{L}; Q)$, then by Proposition 2 the structure Γ' is existentially positively interdefinable with $(\mathbb{L}; C)$ or with $(\mathbb{L}; Q)$; hence, Γ is existentially interdefinable with one of those structures and we are done. Otherwise, Γ' is homomorphically equivalent with a reduct Δ of $(\mathbb{L}; =)$. Again, the homomorphism from Γ' to Δ must in fact be an embedding. Hence, Γ' is isomorphic to a substructure of Δ . Since Δ is preserved by all permutations, so is this substructure, and so is Γ' . It follows that Γ is preserved by all permutations, so Γ is a reduct of $(\mathbb{L}; =)$ by Proposition 19. In fact, Γ is even preserved by all injective maps from \mathbb{L} to \mathbb{L} and therefore by all self-embeddings of $(\mathbb{L}; =)$. Hence, Proposition 2 shows that Γ has an existential definition over $(\mathbb{L}; =)$. Conversely, $(\mathbb{L}; =)$ has an existential definition in every structure with domain \mathbb{L} , so Γ is existentially interdefinable with $(\mathbb{L};=).$

References

- S. A. Adeleke and D. Macpherson. Classification of infinite primitive Jordan permutation groups. Proceedings of the London Mathematical Society, 72(1):63–123, 1996.
- [2] S. A. Adeleke and P. M. Neumann. Primitive permutation groups with primitive Jordan sets. Proceedings of the London Mathematical Society, 53(2):209–229, 1994.
- [3] S. A. Adeleke and P. M. Neumann. *Relations related to betweenness: their structure and automorphisms*, volume 623 of *Memoirs of the AMS*. American Mathematical Society, 1998.

- [4] A. Aho, Y. Sagiv, T. Szymanski, and J. Ullman. Inferring a tree from lowest common ancestors with an application to the optimization of relational expressions. SIAM Journal on Computing, 10(3):405–421, 1981.
- [5] M. Bhattacharjee, D. Macpherson, R. G. Möller, and P. M. Neumann. Notes on Infinite Permutation Groups. Springer Lecture Notes in Mathematics, 1998.
- [6] M. Bodirsky. Complexity classification in infinite-domain constraint satisfaction. Mémoire d'habilitation à diriger des recherches, Université Diderot – Paris 7. Available at arXiv:1201.0856, 2012.
- [7] M. Bodirsky. Ramsey classes: Examples and constructions. To appear in the Proceedings of the 25th British Combinatorial Conference; arXiv:1502.05146, 2015.
- [8] M. Bodirsky, H. Chen, and M. Pinsker. The reducts of equality up to primitive positive interdefinability. *Journal of Symbolic Logic*, 75(4):1249–1292, 2010.
- [9] M. Bodirsky, M. Hils, and B. Martin. On the scope of the universal-algebraic approach to constraint satisfaction. In *Proceedings of the Annual Symposium on Logic in Computer Science (LICS)*, pages 90–99. IEEE Computer Society, July 2010.
- [10] M. Bodirsky, P. Jonsson, and T. V. Pham. The complexity of phylogeny constraint satisfaction. Preprint arXiv:1503.07310, 2015.
- [11] M. Bodirsky and J. Kára. The complexity of equality constraint languages. Theory of Computing Systems, 3(2):136–158, 2008. A conference version appeared in the proceedings of Computer Science Russia (CSR'06).
- [12] M. Bodirsky and J. Kára. The complexity of temporal constraint satisfaction problems. In C. Dwork, editor, *Proceedings of the Annual Symposium on Theory of Computing (STOC)*, pages 29–38. ACM, May 2008.
- [13] M. Bodirsky and J. Kára. The complexity of temporal constraint satisfaction problems. *Journal of the ACM*, 57(2):1–41, 2009. An extended abstract appeared in the Proceedings of the Symposium on Theory of Computing (STOC'08).
- [14] M. Bodirsky and J. K. Mueller. Rooted phylogeny problems. Logical Methods in Computer Science, 7(4), 2011. An extended abstract appeared in the proceedings of ICDT'10.
- [15] M. Bodirsky and D. Piguet. Finite trees are Ramsey with respect to topological embeddings. Technical report, arXiv:1002.1557, 2010.
- [16] M. Bodirsky and M. Pinsker. Reducts of Ramsey structures. AMS Contemporary Mathematics, vol. 558 (Model Theoretic Methods in Finite Combinatorics), pages 489–519, 2011.
- [17] M. Bodirsky and M. Pinsker. Minimal functions on the random graph. Israel Journal of Mathematics, 200(1):251–296, 2014.
- [18] M. Bodirsky and M. Pinsker. Schaefer's theorem for graphs. Journal of the ACM, 62(3):#19, 52 pages, 2015. A conference version appeared in the Proceedings of STOC 2011, pages 655–664.
- [19] M. Bodirsky, M. Pinsker, and A. Pongrácz. The 42 reducts of the random ordered graph. Proceedings of the LMS, 2015. to appear. Preprint available from arXiv:1309.2165.
- [20] M. Bodirsky, M. Pinsker, and T. Tsankov. Decidability of definability. Journal of Symbolic Logic, 78(4):1036–1054, 2013. A conference version appeared in the Proceedings of LICS 2011.

- [21] M. Bodirsky and M. Wrona. Equivalence constraint satisfaction problems. In *Proceedings of Computer Science Logic*, volume 16 of *LIPICS*, pages 122–136. Dagstuhl Publishing, September 2012.
- [22] D. Bryant. Building trees, hunting for trees, and comparing trees. PhD-thesis at the University of Canterbury, 1997.
- [23] D. Bryant and M. Steel. Extension operations on sets of leaf-labelled trees. Advances in Applied Mathematics, 16:425–453, 1995.
- [24] P. J. Cameron. Transitivity of permutation groups on unordered sets. Mathematische Zeitschrift, 148:127–139, 1976.
- [25] P. J. Cameron. Some treelike objects. Quart. J. Math. Oxford Ser. (2), 38(150):155–183, 1987.
- [26] P. J. Cameron. Oligomorphic permutation groups. Cambridge University Press, Cambridge, 1990.
- [27] G. Cantor. Über unendliche, lineare Punktmannigfaltigkeiten. Mathematische Annalen, 23:453–488, 1884.
- [28] W. Deuber. A generalization of Ramsey's theorem for regular trees. Journal of Combinatorial Theory, Series B, 18:18–23, 1975.
- [29] C. Frasnay. Quelques problèmes conbinatoires concernant les ordres totaux et les relations monomorphes. Annales de l'Institut Fourier (Grenoble), 15:415524, 1965.
- [30] D. Haskell and D. Macpherson. Cell decompositions of C-minimal structures. Annals of Pure and Applied Logic, 66:113–162, 1994.
- [31] M. Henzinger, V. King, and T. Warnow. Constructing a tree from homeomorphic subtrees, with applications to computational evolutionary biology. In *Proceedings of the 7th Symposium* on Discrete Algorithms (SODA'96), pages 333–340, 1996.
- [32] B. Herwig, H. D. Macpherson, G. Martin, A. Nurtazin, and J. K. Truss. On ℵ₀-categorical weakly o-minimal structures. Ann. Pure Appl. Logic, 101(1):65–93, 2000.
- [33] W. Hodges. A shorter model theory. Cambridge University Press, Cambridge, 1997.
- [34] M. Junker and M. Ziegler. The 116 reducts of $(\mathbb{Q}, \langle a \rangle)$. Journal of Symbolic Logic, 74(3):861–884, 2008.
- [35] A. Kechris, V. Pestov, and S. Todorcevic. Fraissé limits, Ramsey theory, and topological dynamics of automorphism groups. *Geometric and Functional Analysis*, 15(1):106–189, 2005.
- [36] D. Macpherson. A survey of Jordan groups. In 'Automorphisms of first-order structures', R. Kaye and D. Macpherson, Oxford University Press, pages 73–110, 1994.
- [37] D. Macpherson. A survey of homogeneous structures. Discrete Mathematics, 311(15):1599– 1634, 2011.
- [38] D. Macpherson and C. Steinhorn. On variants of o-minimality. Ann. Pure Appl. Logic, 79(2):165–209, 1996.
- [39] K. R. Milliken. A Ramsey theorem for trees. Journal of Combinatorial Theory, Series A, 26(3):215 – 237, 1979.
- [40] J. Nešetřil. Ramsey classes and homogeneous structures. Combinatorics, Probability & Computing, 14(1-2):171–189, 2005.

- [41] M. P. Ng, M. Steel, and N. C. Wormald. The difficulty of constructing a leaf-labelled tree including or avoiding given subtrees. *Discrete Applied Mathematics*, 98:227–235, 2000.
- [42] P. P. Pach, M. Pinsker, G. Pluhár, A. Pongrácz, and C. Szabó. Reducts of the random partial order. Advances in Mathematics, 267:94–120, 2014.
- [43] F. P. Ramsey. On a problem of formal logic. Proceedings of the LMS (2), 30(1):264–286, 1930.
- [44] M. Steel. The complexity of reconstructing trees from qualitative charaters and subtrees. Journal of Classification, 9:91–116, 1992.
- [45] S. Thomas. Reducts of the random graph. Journal of Symbolic Logic, 56(1):176–181, 1991.