## WEIHRAUCH GOES BROUWERIAN

### VASCO BRATTKA AND GUIDO GHERARDI

ABSTRACT. We prove that the Weihrauch lattice can be transformed into a Brouwer algebra by the consecutive application of two closure operators in the appropriate order: first completion and then parallelization. The closure operator of completion is a new closure operator that we introduce. It transforms any problem into a total problem on the completion of the respective types, where we allow any value outside of the original domain of the problem. This closure operator is of interest by itself, as it generates a total version of Weihrauch reducibility that is defined like the usual version of Weihrauch reducibility, but in terms of total realizers. From a logical perspective completion can be seen as a way to make problems independent of their premises. Alongside with the completion operator and total Weihrauch reducibility we need to study precomplete representations that are required to describe these concepts. In order to show that the parallelized total Weihrauch lattice forms a Brouwer algebra, we introduce a new multiplicative version of an implication. While the parallelized total Weihrauch lattice forms a Brouwer algebra with this implication, the total Weihrauch lattice fails to be a model of intuitionistic linear logic in two different ways. In order to pinpoint the algebraic reasons for this failure, we introduce the concept of a Weihrauch algebra that allows us to formulate the failure in precise and neat terms. Finally, we show that the Medvedev Brouwer algebra can be embedded into our Brouwer algebra, which also implies that the theory of our Brouwer algebra is Jankov logic.

**Keywords:** Weihrauch complexity, computable analysis, Brouwer algebra, intuitionistic and linear logic.

MSC classifications: 03B30, 03D30, 03D78, 03F52, 03F60, 06D20.

### Contents

1.	Introduction	2
2.	Closure Operators and Weihrauch Algebras	5
3.	Precomplete Representations	8
4.	Total Weihrauch Reducibility	11
5.	Completion	15
6.	Algebraic Operations	18
7.	Co-Residual Operations	25
8.	The Brouwer Algebra of Parallelizable Total Degrees	31
9.	Conclusion	34
Acknowledgments		34
References		35

Date: August 7, 2020.

#### 1. Introduction

Over the previous ten years Weihrauch complexity has been developed as a computability theoretic approach to classify the uniform computational content of theorems. A survey article that summarizes some of the current research directions in Weihrauch complexity can be found in [6]. The advantage of this approach is that it provides a direct computability theoretic way to classify problems, while heuristic observation shows that the approach can be seen as a uniform version of reverse mathematics in the sense of Friedman and Simpson [21].

Weihrauch complexity is based on Weihrauch reducibility  $\leq_{\rm W}$  that induces a lattice structure. Beyond the lattice operations the Weihrauch lattice is equipped with a number of additional algebraic operations. Early on it was noticed that the semantics of these operations has the flavor of linear logic. Table 1 provides a dictionary that shows how the usual symbols for operations on problems in the Weihrauch lattice are translated into operations of linear logic.

logical operation in linear logic	algebraic operation on problems
$\otimes$ multiplicative conjunction	$\times$ product
& additive conjunction	$\sqcup$ coproduct
$\oplus$ additive disjunction	$\sqcap$ infimum
? multiplicative disjunction	+ sum
! bang	^ parallelization

Table 1. Linear logic versus the algebra of problems

However, so far no satisfactory interpretation of the Weihrauch lattice as a model of (intuitionistic) linear logic has been found. This is partially due to the lack of an internal implication operation that corresponds to the linear implication  $\multimap$ . Such an implication would have to fulfill

$$(g \multimap f) \leq_{\mathbf{W}} h \iff f \leq_{\mathbf{W}} g \times h$$

and it can be proved that such an implication does not exist, given  $\leq_{\mathbf{W}}$  and  $\times$  [8, Proposition 37]. However, Weihrauch reducibility  $f \leq_{\mathbf{W}} g$  can be seen at least as an external implication operation  $f \Leftarrow g$ .

The Weihrauch lattice has also additional algebraic operations such as the compositional product  $\star$ , which can be seen as a non-commutative version of conjunction. Here  $f \star g$  captures what can be computed by first using the problem g and then the problem f, possibly with some intermediate computation. There is an implication operation  $g \to f$  in the Weihrauch lattice that is a right co-residual operation of  $\star$  [8], i.e., we have

$$(g \to f) \mathop{\leq_{\mathrm{W}}} h \iff f \mathop{\leq_{\mathrm{W}}} g \star h.$$

However, this setting does not provide a model for classical linear logic, since the operation  $\star$  is not commutative.<sup>2</sup>

While the connections to linear logic might not be as tight as one wishes, there is still hope that there is a close connection to intuitionistic logic. In linear logic intuitionistic implication is represented by  $!A \multimap B$ . Hence, it is to be expected that the parallelized Weihrauch reducibility  $f \leq_{\mathrm{W}} \widehat{g}$  gives us an external form of intuitionistic implication. This could theoretically be substantiated by showing that

<sup>&</sup>lt;sup>1</sup>A comprehensive up-to-date bibliography is maintained at the following web page: http://cca-net.de/publications/weibib.php

<sup>&</sup>lt;sup>2</sup>Girard also proposed a less known non-commutative version of linear logic, but also this logic does not seem to fit to our model [28].

the resulting structure is a Brouwer algebra, since Brouwer algebras are models for intermediate propositional logics in between classical and intuitionistic logic. However, also this hope did not materialize as Higuchi and Pauly proved that the parallelized Weihrauch lattice is not a Brouwer algebra [13].

In this article we prove that one does obtain a Brouwer algebra if one combines two closure operators in the Weihrauch lattice in the appropriate order: first completion  $f\mapsto \overline{f}$  and then parallelization  $f\mapsto \widehat{f}$ . While parallelization is a well understood operation [3] that corresponds somewhat to the usage of countable choice in constructive mathematics, completion is a new operation that we introduce in this article. Formally, the completion  $\overline{f}: \overline{X} \rightrightarrows \overline{Y}$  of a problem  $f: \subseteq X \rightrightarrows Y$  is defined by

$$\overline{f}(x) := \left\{ \begin{array}{ll} \underline{f}(x) & \text{if } x \in \text{dom}(f) \\ \overline{Y} & \text{otherwise} \end{array} \right.,$$

i.e., by a totalization of f on the completions  $\overline{X}, \overline{Y}$  of the corresponding types.<sup>3</sup>

Logically, completion can be seen as a way to make problems independent of their premises. In general, a logical statement of the form

$$(\forall x \in X)(x \in D \Longrightarrow (\exists y \in Y) \ P(x,y))$$

is translated into a problem  $f:\subseteq X \Rightarrow Y$  in the Weihrauch lattice by setting  $\operatorname{dom}(f) = D$  and  $\underline{f}(x) := \{y \in Y : P(x,y)\}$  for all  $x \in \operatorname{dom}(f)$ . Now the transition to the completion  $\overline{f}$  corresponds to the statement

$$(\forall x \in \overline{X})(\exists y \in \overline{Y})(x \in D \Longrightarrow P(x,y)),$$

where the existence is required independent of the premise  $x \in D$ . The completion of the data types is relevant here, as it guarantees the existence of total representations of the underlying types.

The completion operation  $f \mapsto \overline{f}$  is of interest by itself as it is a closure operator that yields a total version of Weihrauch reducibility  $\leq_{tW}$  by  $f \leq_{tW} g \iff f \leq_{W} \overline{g}$ . Total Weihrauch reducibility  $\leq_{tW}$  can also be defined directly almost as the usual reducibility  $\leq_{W}$ , but in terms of total realizers instead of partial realizers. In this case the completion of the types features again, since one needs to consider so-called precomplete representations for the underlying types.

Among other things we prove that total Weihrauch reducibility induces a lattice structure with operations induced by the original operations of the Weihrauch lattice. The lattice structure of the total Weihrauch lattice is somewhat different from the original Weihrauch lattice, but it does not change all too dramatically as many problems are actually complete, i.e., Weihrauch equivalent to their own completion. We list some examples of complete and incomplete problems:

- Complete problems: LPO, LLPO,  $\lim$ , J, WKL, SORT, IVT, PA, MLR,  $DNC_n$ .
- Incomplete problems:  $C_{\mathbb{N}}, C_{\mathbb{N}^{\mathbb{N}}}, WWKL$ .

The reader who does not know these problems will find relevant definitions of some of them later. The topic of completion of choice problems is subject of an entirely separate article [4].

When we move to the total Weihrauch lattice  $W_{tW}$  of total Weihrauch reducibility  $\leq_{tW}$ , then we can introduce a new implication  $f \rightarrow g$  that can almost be seen as a multiplicative co-residual of  $\times$ . However, also in this case we fail to obtain a model for intuitionistic linear logic.

In order to make the spectacular twofold failure of obtaining a model of intuitionistic linear logic more understandable, we introduce the concept of a Weihrauch

<sup>&</sup>lt;sup>3</sup>We were inspired to continue the study of completions by recent work of Dzhafarov who used them to show that strong Weihrauch reducibility induces a lattice structure [10].




FIGURE 1. Different types of algebras as models of logic

algebra in the following section 2. These are lattice-ordered monoids with some additional implication operation. The total Weihrauch lattice  $\mathcal{W}_{tW}$  is a commutative Weihrauch algebra with respect to  $\times$ ,  $\rightarrow$  and a deductive Weihrauch algebra with respect to  $\star$ ,  $\rightarrow$ . However, none of these Weihrauch algebras is commutative and deductive simultaneously, which is what is required in order to obtain, in our terms, a Troelstra algebra<sup>4</sup>, i.e., a model of some form of intuitionistic linear logic. See the diagram in Figure 1 for an illustration of the situation.

When we apply parallelization after completion, then we obtain the parallelized total Weihrauch lattice  $W_{ptW}$  which then leads to a Brouwer algebra, i.e., a Troelstra algebra where the monoid structure is merged with the lattice structure (in our terms  $\times$  and  $\sqcup$  are merged). In section 8 we prove that one can embed the Medvedev Brouwer algebra [24] into our Brouwer algebra. Like in the case of the Medvedev Brouwer algebra we obtain Jankov logic as the theory of our algebra.

In the following section 2 we provide some very basic lattice theoretic results regarding closure operators that are helpful for our study, and we define Weihrauch and Troelstra algebras alongside with Brouwer algebras. In section 3 we study precomplete representations and the data type of completion that is needed to introduce the closure operator of completion and total Weihrauch reducibility. In section 4 we introduce total Weihrauch reducibility and we prove some basic properties of it. In section 5 we introduce and study the closure operator of completion. Section 6 provides results that show how the algebraic operations of the Weihrauch lattice interact with completion. In particular, we prove that total Weihrauch reducibility actually yields a lattice structure. In section 7 we review the operations \* and  $\rightarrow$  and study their interaction with completion and we also introduce the new implication operation -. Finally, in section 8 we prove that the parallelized total Weihrauch lattice  $W_{\text{ptW}}$  is a Brouwer algebra with the implication derived from  $\rightarrow$ . We also discuss the embedding of the Medvedev lattice. We close this article with a brief survey on the classification of concrete problems in the parallelized total Weihrauch lattice.

<sup>&</sup>lt;sup>4</sup>This is the dual structure of what Troelstra called an *intuitionistic linear algebra* [26].

### 2. Closure Operators and Weihrauch Algebras

In this section we prepare some basic order theoretic concepts that we are going to use frequently. We recall that a  $preorder \leq on$  a set X is a binary relation on X that is reflexive and transitive. We also speak of a preordered space  $(X, \leq)$  in this context. An equivalence  $relation \equiv on$  a set X is a binary relation on X that is reflexive, symmetric and transitive. In the following we will have to deal with several closure operators.

**Definition 2.1** (Closure operator). Let  $(X, \leq)$  be a preordered space together with a map  $c: X \to X$ . Then c is called a *closure operator*, if

- (1)  $x \leq c(x)$ ,
- (2)  $cc(x) \le c(x)$  and
- (3)  $x \le y \Longrightarrow c(x) \le c(y)$

hold for all  $x, y \in X$ . We say that  $x \in X$  is closed if  $c(x) \leq x$ .

We call a map  $c: X \to X$  monotone, if  $x \leq y \Longrightarrow c(x) \leq c(y)$  holds and antitone, if  $x \leq y \Longrightarrow c(y) \leq c(x)$  holds. We use the same terminology for binary maps  $\square: X \times X \to X$  with respect to individual arguments. We use the usual concepts of a suprema (also called a least upper bound) and an infima (also called a greatest lower bound) for preordered sets in the usual way, and we note that on a preordered space they are only uniquely determined up to equivalence in the case of existence. If one has a preordered space  $(X, \leq)$  and one identifies all equivalent elements with each other, then one obtains a quotient structure  $(X/\equiv, \leq)$ , which is a partially ordered space, i.e., the resulting order is a preorder that is additionally anti-symmetric. A lattice  $(X, \leq, \wedge, \vee)$  is a partially ordered set together with a supremum operation  $\vee$  and an infimum operation  $\wedge$ . If  $\leq_c$  is a preorder on X and  $c: X \to X$  a map, then we say that c generates  $\leq_c$  on  $(X, \leq)$  if  $x \leq_c y \iff x \leq c(y)$  holds for all  $x, y \in X$ . The following result is straightforward to prove. It shows how closure operators act on lattices and preordered spaces.

**Proposition 2.2** (Closure operators). Let  $(X, \leq)$  be a preordered space with two closure operators  $c, c': X \to X$  and binary operations  $\square, \vee, \wedge: X \times X \to X$ . Then

- (1)  $x \leq_c y : \iff x \leq c(y) \iff c(x) \leq c(y)$  defines a preorder that satisfies  $x \leq y \implies x \leq_c y$  for all  $x, y \in X$ ,
- (2)  $x \equiv_c y : \iff (x \leq_c y \text{ and } y \leq_c x) \text{ defines an equivalence relation,}$
- (3)  $\square_c: X \times X \to X, (x,y) \mapsto c(x)\square c(y)$  shares corresponding monotonicity properties as  $\square$ , more precisely:
  - (a) if  $\square$  is monotone (antitone) in one argument with respect to  $\leq$ , then so is  $\square_c$  in the same argument with respect to  $\leq_c$ ,
  - (b) if  $\wedge$  is an infimum with respect to  $\leq$ , then so is  $\wedge_c$  with respect to  $\leq_c$ ,
  - (c) if  $\vee$  is a supremum with respect to  $\leq$ , then so is  $\vee_c$  with respect to  $\leq_c$ .
- (4) If  $(X, \leq, \wedge, \vee)$  is a lattice, then so is  $(X/\equiv_c, \leq_c, \wedge_c, \vee_c)$ .
- (5)  $c' \circ c: X \to X$  is monotone with respect  $to \leq and \leq_c$ .

*Proof.* (1) Reflexivity of  $\leq_c$  follows from  $x \leq c(x)$ , transitivity from monotonicity of c together with  $cc(x) \leq c(x)$ . It is also clear that  $x \leq c(y) \iff c(x) \leq c(y)$  holds. Finally,  $x \leq y \implies x \leq_c y$  holds as c is monotone.

- (2) Is obvious.
- (3) (a) Suppose  $\square$  is antitone in the first argument and  $x_1, x_2, y \in X$  with  $x_1 \leq_c x_2$ . Then  $c(x_1) \leq c(x_2)$  and hence  $c(x_2) \square c(y) \leq c(x_1) \square c(y)$ , since  $\square$  is antitone in the first argument. Hence  $x_2 \square_c y \leq x_1 \square_c y \leq c(x_1 \square_c y)$ , which means  $x_2 \square_c y \leq_c x_1 \square_c y$ , i.e.,  $\square_c$  is antitone in the first argument with respect to  $\leq_c$ . The other cases are treated analogously.
- (b), (c) If  $\wedge$  is an infimum with respect to  $\leq$  and  $x,y\in X$ , then  $x\wedge y\leq x$  and

 $x \wedge y \leq y$  and hence  $x \wedge y \leq_c x$  and  $x \wedge y \leq_c y$ . Hence  $x \wedge y$  is a lower bound of x and y with respect  $\leq_c$ . Let now  $z \in X$  be such that  $z \leq_c x$  and  $z \leq_c y$ . Then  $z \leq c(x)$  and  $z \leq c(y)$ , which implies  $z \leq c(x) \wedge c(y)$  and hence  $z \leq_c x \wedge_c y$ . This means that  $x \wedge_c y$  is above every lower bound of x and y with respect to  $\leq_c$  and hence it is an infimum with respect to  $\leq_c$ . The statement for suprema can be proved analogously.

- (4) This follows from (1)–(3).
- (5) If  $x \leq y$ , then  $c'c(x) \leq c'c(y)$  follows. If  $x \leq_c y$ , then  $cc'c(x) \leq cc'c(y)$  follows and hence  $c'c(x) \leq_c c'c(y)$ .

We also need to deal with situations where a closure operator respects certain underlying algebraic operations or other closure operators. Hence, we use the following terminology.

**Definition 2.3** (Preservation). Let  $(X, \leq)$  be a preordered space with closure operators  $c, c': X \to X$  and a binary operation  $\Box: X \times X \to X$ .

- (1) We say that c is preserved by  $\square$  if  $c(x\square y) \le c(x)\square c(y)$  for all  $x, y \in X$ .
- (2) We say that c is co-preserved by  $\square$  if  $c(x)\square c(y) \le c(x\square y)$  for all  $x, y \in X$ .
- (3) We say that c is preserved by c' if  $c \circ c'(x) \le c' \circ c(x)$  for all  $x \in X$ .

Whenever a closure operator is preserved by a certain operation, then we can draw certain conclusions. The proof of the following result is straightforward.

**Proposition 2.4** (Preservation). Let  $(X, \leq)$  be a preordered space with closure operators  $c, c': X \to X$  and a binary monotone operation  $\square: X \times X \to X$ .

(1) If c is preserved by  $\square$ , then for all  $x, y \in X$ 

$$c(x\Box y) < c(x)\Box c(y) \equiv c(c(x)\Box c(y)).$$

In particular,  $x \square y$  is closed if x and y are.

(2) If c is co-preserved by  $\square$ , then for all  $x, y \in X$ 

$$c(x)\Box c(y) \le c(x\Box y) \le c(c(x)\Box c(y)).$$

(3) If c is preserved by c', then for all  $x \in X$ 

$$cc'(x) < c'c(x) \equiv cc'c(x)$$
.

In particular, c'c is a closure operator with respect to  $\leq_c$  and  $\leq$ , and c'(x) is closed with respect to c if x is so.

*Proof.* The equivalences in (1) and (3) are consequences of the respective first relations and the fact that c is a closure operator. For the second relation in (2) we just use that c is a closure operator. It is clear that c'(x) is closed if x is closed. That c'c is monotone with respect to  $\leq_c$  follows from Proposition 2.2. Clearly, also  $x \leq_c c'c(x)$  holds. Finally,  $c'cc'c(x) \leq c'c'(x) \leq c'c(x)$  and hence  $c'cc'c(x) \leq_c c'c(x)$ .

If the binary operation is a supremum or an infimum operation, then it is always preserved in certain ways.

**Proposition 2.5** (Preservation of suprema and infima). Let  $(X, \leq)$  be a preordered space with a closure operator  $c: X \to X$ , and binary operations  $\vee, \wedge: X \times X \to X$ .

- (1) If  $\vee$  is a supremum operation, then it co-preserves c.
- (2) If  $\wedge$  is an infimum operation, then it preserves c.

In particular,  $x \vee y \equiv_c x \vee_c y$ , if  $\vee$  is a supremum.

*Proof.* Since  $x \vee y$  is a supremum, we obtain  $c(x) \leq c(x \vee y)$  and  $c(y) \leq c(x \vee y)$  due to monotonicity of c. Hence  $c(x) \vee c(y) \leq c(x \vee y)$ , which means that  $\vee$  co-preserves c. The statement for  $\wedge$  can be proved analogously. That  $\vee$  co-preserves c means  $x \vee_c y \leq_c x \vee y$ . We also have  $x \vee y \leq c(x) \vee c(y)$ , i.e.,  $x \vee y \leq_c x \vee_c y$ .

We note that this result implies that the we can replace  $\vee_c$  by  $\vee$  in Proposition 2.2.

In the following we will have to deal with lattices that have some additional algebraic operations and we propose the following concept that encapsulates a structure that we will see in different variations.

**Definition 2.6** (Weihrauch algebra). We call  $(X, \leq, \land, \lor, \cdot, \rightarrow, 1, \bot, \top)$  a Weihrauch algebra if the following hold:

- (1)  $(X, \leq, \wedge, \vee)$  is a bounded lattice with bottom  $\bot$  and top  $\top$ . (Lattice)
- (2)  $(X, \cdot, 1)$  is a monoid with neutral element 1. (Monoid)
- (3) :  $X \times X \to X$  is monotone in both components. (Monotonicity)
- (4)  $\rightarrow: X \times X \to X$  is monotone in the second component, antitone in the first component. (Monotonicity)
- (5)  $x \le y \cdot z \Longrightarrow (y \to x) \le z$  holds for all  $x, y, z \in X$ . (Implication)

A Weihrauch algebra is called *commutative*, if  $\cdot$  is commutative, and it is called *deductive*, if " $\iff$ " holds instead of " $\implies$ " in (5).

One could add additional distributivity requirements to this definition. Structures that satisfy (1), (2) and (3) have also been called *lattice-ordered monoids*. Using these building blocks, we can define structures that have been already considered for other purposes.

**Definition 2.7** (Algebras). Let  $\mathcal{X} = (X, \leq, \land, \lor, \cdot, \rightarrow, 1, \bot, \top)$  be a Weihrauch algebra. We call  $\mathcal{X}$  a *Troelstra algebra* if it is commutative and deductive. If, additionally,  $\cdot = \lor$  and  $1 = \bot$ , then  $\mathcal{X}$  is called a *Brouwer algebra*.

If we denote a Brouwer algebra as a tuple, then we omit the double occurrence of  $\cdot = \vee$  and  $1 = \bot$ , respectively. What we call a Troelstra algebra is exactly what Troelstra [25] called an *intuitionistic linear algebra*, except that the order is reversed. A bottom element in our sense is not required in Troelstra's axioms, but it always exists by [25, Lemma 8.3]. The relevance of Troelstra algebras is that they form sound and complete models of intuitionistic linear logic [25, Theorem 8.15]. In an analogous sense Brouwer algebras (that are just defined dually to Heyting algebras<sup>5</sup>) are known as models of intermediate logics, i.e., predicate logics between classical logic and intuitionistic logic [12].

A Brouwer algebra embedding is an injective map from one Brouwer algebra to another one that is monotone in both directions, preserves suprema, infima, implications and the bottom and top elements.

In the case of a deductive Weihrauch algebra the condition (5) can be seen as a law of (co-)residuation. We need to add the prefix "co-" as residuation is normally considered in the opposite order [12].

**Definition 2.8** (Co-residuation). Let  $(X, \leq)$  be a preordered set with a binary operation  $\cdot: X \times X \to X$ . Then we call  $\cdot$  right co-residuated, if there is a binary operation  $\to: X \times X \to X$  such that

$$x \le y \cdot z \iff (y \to x) \le z$$

<sup>&</sup>lt;sup>5</sup>The term Brouwer algebra is used in different versions in different references, we mean by a *Brouwer algebra* just the dual concept of a Heyting algebra, as usual in computability theory [24].

holds for all  $x, y, z \in X$ . Analogously, we call  $\cdot$  left co-residuated, if an analogous condition holds with  $z \cdot y$  in place of  $y \cdot z$ . Finally,  $\cdot$  is called co-residuated if and only if it is left and right co-residuated.

Hence, a deductive Weihrauch algebra is a right co-residuated lattice-ordered monoid and a Troelstra algebra is a co-residuated lattice-ordered monoid.

### 3. Precomplete Representations

We will need some pairing functions in the following. Firstly, we define a pairing function  $\pi: \mathbb{N}^{\mathbb{N}} \times \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}, (p,q) \mapsto \langle p,q \rangle$  by  $\langle p,q \rangle (2n) := p(n)$  and  $\langle p,q \rangle (2n+1) := q(n)$  for  $p,q \in \mathbb{N}^{\mathbb{N}}$  and  $n \in \mathbb{N}$ . We define a pairing function of type  $\langle , \rangle : (\mathbb{N}^{\mathbb{N}})^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  by  $\langle p_0, p_1, p_2, ... \rangle \langle n, k \rangle := p_n(k)$  for all  $p_i \in \mathbb{N}^{\mathbb{N}}$  and  $n, k \in \mathbb{N}$ , where  $\langle n, k \rangle$  is the standard Cantor pairing defined by  $\langle n, k \rangle := \frac{1}{2}(n+k+1)(n+k)+k$ . Finally, we note that by np we denote the concatenation of a number  $n \in \mathbb{N}$  with a sequence  $p \in \mathbb{N}^{\mathbb{N}}$ . By  $\pi_i : \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}, \langle p_0, p_1, p_2, ... \rangle \mapsto p_i$  we denote the projection on the i-th component of a tuple and we also use the binary tupling functions  $\pi_1 \langle p, q \rangle = p$  and  $\pi_2 \langle p, q \rangle = q$ . It will always be clear from the context whether we apply these functions in a countable or binary setting.

We recall that a represented space  $(X, \delta)$  is a set X together with a surjective (partial) map  $\delta : \subseteq \mathbb{N}^{\mathbb{N}} \to X$ , called the representation of X. For the purposes of our topic so-called precomplete representations are important. They were introduced by Kreitz and Weihrauch [15] following the concept of a precomplete numbering, that was originally introduced by Eršov [11].

**Definition 3.1** (Precompleteness). A representation  $\delta :\subseteq \mathbb{N}^{\mathbb{N}} \to X$  is called *precomplete*, if for any computable function  $F :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  there exists a total computable function  $G : \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  such that  $\delta F(p) = \delta G(p)$  for all  $p \in \text{dom}(F)$ .

In this situation we also say that the represented space  $(X, \delta)$  is precomplete. We point out that we demand that the equation in the definition holds for all  $p \in \text{dom}(F)$ , not only for  $p \in \text{dom}(\delta F)$ . The precomplete representations are exactly those that satisfy a certain version of the recursion theorem [15]. For us they are relevant since we are going to work with total functions. It is clear that not all represented spaces are precomplete. By  $\text{id}: \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  we denote the identity of Baire space. For other sets X we usually add an index X and write the identity as  $\text{id}_X: X \to X$ . By  $\widehat{n} := nnn... \in \mathbb{N}^{\mathbb{N}}$  we denote the constant sequence with value  $n \in \mathbb{N}$ .

**Example 3.2.** There are partial computable functions  $F :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  without total computable extension, such as the function defined by  $F(p) = \widehat{n} : \iff p$  starts with exactly n digits 0, where  $\text{dom}(F) = \{0^n p : n \in \mathbb{N}, p \in \mathbb{N}^{\mathbb{N}}, p(0) \neq 0\}$ . This shows that the represented space  $(\mathbb{N}^{\mathbb{N}}, \text{id})$  is not precomplete.

However, it is not too hard to see that in every equivalence class of representations there is a precomplete representation. We recall that for two representations  $\delta_1, \delta_2$  of the same set X we say that  $\delta_1$  is computably reducible to  $\delta_2$ , in symbols  $\delta_1 \leq \delta_2$ , if and only if there is a computable  $F : \subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  such that  $\delta_1 = \delta_2 F$ . We denote the corresponding equivalence by  $\equiv$ . For  $p \in \mathbb{N}^{\mathbb{N}}$  we denote by  $p-1 \in \mathbb{N}^{\mathbb{N}} \cup \mathbb{N}^*$  the sequence or word that is formed as concatenation of p(0)-1, p(1)-1, p(2)-1,... with the understanding that  $-1 = \varepsilon$  is the empty word.

**Definition 3.3** (Precompletion). Let  $(X, \delta_X)$  be a represented space. Then the precompletion  $\delta_X^{\wp}$  of  $\delta_X$  is defined by  $\delta_X^{\wp}(p) := \delta_X(p-1)$  for all  $p \in \mathbb{N}^{\mathbb{N}}$  such that  $p-1 \in \text{dom}(\delta_X)$ .

 $<sup>^6</sup>$ This result is due to Matthias Schröder (personal communication 2009), see the construction in [20, Lemmas 4.2.10, 4.2.11, Section 4.2.5].

We note that the identity id :  $\mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ , considered as a representation of  $\mathbb{N}^{\mathbb{N}}$ , has the precompletion id  $\mathcal{P} :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  with id  $\mathcal{P}(p) := p-1$  and in general  $\delta_X^{\wp} = \delta_X \circ \mathrm{id}^{\wp}$ . Now we can prove the following result.

**Proposition 3.4** (Precompleteness). Let  $(X, \delta_X)$  be a represented space. The precompletion  $\delta_X^{\wp}$  of  $\delta_X$  is precomplete and satisfies  $\delta_X^{\wp} \equiv \delta_X$ .

Proof. The computable function  $F: \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}, p \mapsto p+1$  satisfies  $\delta_X(p) = \delta_X^{\wp} F(p)$  and hence it witnesses  $\delta_X \leq \delta_X^{\wp}$ . The computable function  $G: \subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}, p \mapsto p-1$  satisfies  $\delta_X^{\wp}(p) = \delta_X G(p)$  and hence it witnesses  $\delta_X^{\wp} \leq \delta_X$ . Altogether  $\delta_X^{\wp} \equiv \delta_X$ . We need to prove that  $\delta_X^{\wp}$  is precomplete. Let  $F: \subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  be computable and let M be a Turing machine that computes F. We modify this machine such that it never halts and after every n steps for some suitable fixed number  $n \in \mathbb{N}$  the machine writes a 0 on the output tape, irrespective of the input. Otherwise the machine is left unchanged. Then the modified machine computes a total function  $G: \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  with F(p) - 1 = G(p) - 1 and hence  $\delta_X^{\wp} F(p) = \delta_X^{\wp} G(p)$  for all  $p \in \text{dom}(F)$ .

We will also need the fact that other classes of functions can be extended to total ones under precomplete representations. Hence we introduce the following concept.

**Definition 3.5** (Respect for precompleteness). We say that a set P of functions  $F :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  respects precompleteness, if for every precomplete representation  $\delta$  and any function  $F \in P$  there exists a total function  $G \in P$  such that  $\delta F(p) = \delta G(p)$  for all  $p \in \text{dom}(F)$ .

It is clear that the set of computable functions respects precompleteness by definition. However, also other classes of functions do. Some of them, simply because they can already be extended to total functions in the same class irrespectively of the representation. We provide a number of examples. We call a function non-uniformly computable if it maps all computable inputs in its domain to computable outputs. By  $J: \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ ,  $p \mapsto p'$  we denote the Turing jump operator and by  $U:\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  a universal computable function such that for every continuous function  $F:\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  there is a  $q \in \mathbb{N}^{\mathbb{N}}$  with  $F(p) = U\langle q, p \rangle$  for all  $p \in \text{dom}(F)$  [27, Theorem 2.3.8].

**Proposition 3.6** (Respect for precompleteness). The following classes of partial functions  $F :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  respect precompleteness: computable, continuous, limit computable, Borel measurable and non-uniformly computable.

Proof. The statement for computable functions is a consequence of the definition of precompleteness. Let  $\delta$  be a precomplete representation and let  $F:\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ be a continuous function. Then there is a  $q \in \mathbb{N}^{\mathbb{N}}$  such that F(p) = U(q, p) for all  $p \in \text{dom}(F)$ . By precompleteness of  $\delta$ , there is a total function  $V: \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ with  $\delta V(p) = \delta U(p)$  for all  $p \in \text{dom}(U)$ . Hence  $G(p) := V\langle q, p \rangle$  defines a total continuous function with  $\delta G(p) = \delta F(p)$  for all  $p \in \text{dom}(F)$ . This shows that the class of continuous functions respects precompleteness. For every limit computable function  $F:\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  there exists a computable function  $H:\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  such that  $F = H \circ J$  [1, Theorem 14]. By precompleteness of  $\delta$  there exists a total computable function  $I: \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  such that  $\delta I(p) = \delta H(p)$  for all  $p \in \text{dom}(H)$ . Hence  $G := I \circ J$  is a total function that is limit computable and satisfies  $\delta F(p) =$  $\delta G(p)$  for all  $p \in \text{dom}(F)$ . Hence the class of limit computable functions respects precompleteness. Every partial Borel measurable function  $F:\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  can be extended to a total Borel measurable function by a theorem of Kuratowski (see [14, Theorem 2.2]). The class of non-uniformly computable functions respects precompleteness since every non-uniformly computable function  $F:\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  can simply be extended to a total non-uniformly computable function  $G: \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  by defining  $G(p) = \widehat{0} = 000...$  for all  $p \in \mathbb{N}^{\mathbb{N}} \setminus \text{dom}(F)$ .

The proof for limit computable functions (which are exactly the effectively  $\Sigma_2^0$ computable functions) can easily be extended to any finite level of the Borel hierarchy. We prove in [4, Corollaries 8.4, 9.3] that functions that are computable with finitely many mind changes and low functions do not respect precompleteness.

We also need to study how certain algebraic constructions on represented spaces behave with respect to precompleteness. For any sets X and Y we denote by  $X \times Y$ and  $X^{\mathbb{N}}$  the usual products, by  $X \sqcup Y := (\{0\} \times X) \cup (\{1\} \times Y)$  the disjoint union of X and Y, by  $X^* := \bigcup_{i=0}^{\infty} (\{i\} \times X^i)$  the set of words over X, where  $X^i$  denotes the *i*-fold product of X with itself, and  $X^0 := \{0\}$ . By  $\overline{X} := X \cup \{\bot\}$  we denote the completion X, where we assume that  $\bot \notin X$ .

**Definition 3.7** (Constructions on representations). Let  $(X, \delta_X)$  and  $(Y, \delta_Y)$  be represented spaces. We define

- $\begin{array}{l} (1) \ \delta_{X\times Y}:\subseteq \mathbb{N}^{\mathbb{N}} \to X\times Y, \ \delta_{X\times Y}\langle p,q\rangle:=(\delta_X(p),\delta_Y(q)) \\ (2) \ \delta_{X\sqcup Y}:\subseteq \mathbb{N}^{\mathbb{N}} \to X\sqcup Y, \ \delta_{X\sqcup Y}(0p):=(0,\delta_X(p)) \ \text{and} \ \delta_{X\sqcup Y}(1p):=(1,\delta_Y(p)) \\ (3) \ \delta_{X^*}:\subseteq \mathbb{N}^{\mathbb{N}} \to X^*, \ \delta_{X^*}(n\langle p_1,p_2,...,p_n\rangle):=(n,(\delta_X(p_1),\delta_X(p_2),...,\delta_X(p_n))) \\ (4) \ \delta_{X^{\mathbb{N}}}:\subseteq \mathbb{N}^{\mathbb{N}} \to X^{\mathbb{N}}, \ \delta_{X^{\mathbb{N}}}\langle p_0,p_1,p_2,...\rangle:=(\delta_X(p_n))_{n\in\mathbb{N}} \\ (5) \ \delta_{\overline{X}}:\mathbb{N}^{\mathbb{N}} \to \overline{X}, \ \delta_{\overline{X}}(p):=\delta_X^{\mathcal{G}}(p) \ \text{if} \ p\in \mathrm{dom}(\delta_X^{\mathcal{G}}) \ \text{and} \ \delta_{\overline{X}}(p):=\bot \ \text{otherwise}. \end{array}$

We warn the reader that all these constructions on represented spaces preserve equivalence of representations, except the last one for the completion. In other words, the equivalence class of  $\delta_{\overline{Y}}$  does not only depend on the equivalence class of  $\delta_X$ , but on the concrete representative  $\delta_X$  itself. For our applications this does not cause any problems (see the remark after Corollary 5.3; the problem could also be circumvented by moving to multi-valued representations [20, Lemma 4.2.11]).

The next observation is that finite and countable products preserve precomplete-

**Proposition 3.8** (Products and precompleteness). Let  $(X, \delta_X)$  and  $(Y, \delta_Y)$  be precomplete represented spaces. Then so are  $(X \times Y, \delta_{X \times Y})$  and  $(X^{\mathbb{N}}, \delta_{X^{\mathbb{N}}})$ .

*Proof.* If  $F:\subseteq \mathbb{N}^{\mathbb{N}}\to \mathbb{N}^{\mathbb{N}}$  is computable, then so are the projections  $F_i=\pi_i\circ F$ for  $i \in \{1,2\}$  with  $\pi_1\langle p,q\rangle = p$  and  $\pi_2\langle p,q\rangle = q$ . Hence, by precompleteness there are total computable functions  $G_i: \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  with  $\delta_X F_1(p) = \delta_X G_1(p)$ for all  $p \in \text{dom}(F_1)$  and with an analogous statement for  $\delta_Y, F_2$  and  $G_2$ . Let  $G(p) = \langle G_1(p), G_2(p) \rangle$  for all  $p \in \mathbb{N}^{\mathbb{N}}$ . Then  $G : \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  is computable and total, and we obtain  $\delta_{X\times Y}F(p)=\delta_{X\times Y}G(p)$  for all  $p\in \text{dom}(F)$ . Hence  $\delta_{X\times Y}$  is precomplete. If  $F:\subseteq\mathbb{N}^{\mathbb{N}}\to\mathbb{N}^{\mathbb{N}}$  is computable, then so is the function  $H:\subseteq\mathbb{N}^{\mathbb{N}}\to\mathbb{N}^{\mathbb{N}}, \langle i,p\rangle\mapsto \pi_i\circ F(p)$ , where  $\pi_i:\mathbb{N}^{\mathbb{N}}\to\mathbb{N}^{\mathbb{N}}, \langle p_0,p_1,p_2,...\rangle\mapsto p_i$  denotes the i–th projection. Due to precompleteness of  $\delta_X$  there is a total computable function  $I: \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  with  $\delta_X H(p) = \delta_X I(p)$  for all  $p \in \text{dom}(H)$ . Then also the function  $G: \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}, p \mapsto \langle I(0,p), I(1,p), I(2,p), ... \rangle$  is computable and total and satisfies  $\delta_{X^{\mathbb{N}}}F(p) = \delta_{X^{\mathbb{N}}}G(p)$  for all  $p \in \text{dom}(F)$ . This shows that  $\delta_{X^{\mathbb{N}}}$  is precomplete.

The coproduct constructions for  $X \sqcup Y$  and  $X^*$  are less nicely behaved with respect to precompleteness. One problem is that also the natural number component that selects the argument has to be handled in a precomplete manner. One can modify the definition of  $\delta_{X \sqcup Y}$  and  $\delta_{X^*}$  to take this into account. However, even then it is not clear why the construction should preserve precompleteness. We just obtain that if  $\delta_X$  and  $\delta_Y$  are the precompletions according to Proposition 3.4, then  $\delta_{X \sqcup Y}$  and  $\delta_{X^*}$  are precomplete in the modified definition. We formulate this more formally. We use the total representation  $\delta_{\mathbb{N}}$  of  $\mathbb{N}$  given by  $\delta_{\mathbb{N}}(p) := p(0)$ .

**Proposition 3.9** (Coproducts and precompleteness). Let  $(X, \delta_X)$  and  $(X_i, \delta_{X_i})$  be represented spaces for  $i \in \{0, 1\}$ .

(1) We define a representation  $\delta$  of  $X_0 \sqcup X_1$  by

$$\delta\langle q, p \rangle := (\delta_{\mathbb{N}}^{\wp}(q), \delta_{X_i}^{\wp}(p))$$

for all  $q, p \in \mathbb{N}^{\mathbb{N}}$  such that  $\delta_{\mathbb{N}}^{\wp}(q) = i \in \{0, 1\}$  and  $p \in \text{dom}(\delta_{X_i}^{\wp})$ . Then  $\delta$  is precomplete and  $\delta \equiv \delta_{X_0 \sqcup X_1}$ .

(2) We define a representation  $\delta$  of  $X^*$  by

$$\delta\langle q, \langle p_1, ..., p_n \rangle \rangle := (\delta_{\mathbb{N}}^{\wp}(q), (\delta_{\mathbb{X}}^{\wp}(p_1), ..., \delta_{\mathbb{X}}^{\wp}(p_n)))$$

for all  $q, p_1, ..., p_n \in \mathbb{N}^{\mathbb{N}}$  such that  $\delta_{\mathbb{N}}^{\wp}(q) = n$  and  $p_i \in \text{dom}(\delta_X^{\wp})$  for i = 1, ..., n. Then  $\delta$  is precomplete and  $\delta \equiv \delta_{X^*}$ .

Proof. The proof is similar to the proof of Proposition 3.4. We only consider the case of  $X^*$  and leave the case  $X_0 \sqcup X_1$  to the reader. Given a  $\delta_{X^*}$ -name  $\langle n, \langle p_1, ..., p_n \rangle \rangle$  of  $x \in X^*$  we compute  $q := \widehat{n} = nnn...$  and then  $\langle q+1, \langle p_1+1, ..., p_n+1 \rangle \rangle$  is a  $\delta$ -name of the same point x. Since  $r \mapsto r+1$  is computable, we obtain  $\delta_{X^*} \leq \delta$ . Given a  $\delta$ -name  $\langle q, \langle p_1, ..., p_n \rangle \rangle$  of a point  $x \in X^*$ , we can search for the first non-zero value  $k \in \mathbb{N}$  in q, in which case we know that n = k-1, and then we can compute  $\langle n, \langle p_1-1, ..., p_n-1 \rangle \rangle$ , which is a  $\delta_{X^*}$ -name of the same point x. Since  $r \mapsto r-1$  is computable on sequences such that  $r-1 \in \mathbb{N}^{\mathbb{N}}$ , we obtain  $\delta \leq \delta_{X^*}$ . Any machine that computes a function  $F : \subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  can be modified as in the proof of Proposition 3.4 such that it computes a total function  $G : \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ , potentially with extra zeros on the output side and such that  $\delta F(p) = \delta G(p)$  for all  $p \in \text{dom}(F)$ .

We mention that the completion  $(\overline{X}, \delta_{\overline{X}})$  of a represented space is always precomplete. This follows like in the proof of Proposition 3.4. The only additional observation required in the proof is that if  $\delta_{\overline{X}}F(p)=\bot$ , then also  $\delta_{\overline{X}}G(p)=\bot$ . We recall that a *computable embedding*  $f:X\to Y$  is a computable function that is injective and whose partial inverse  $f^{-1}:\subseteq Y\to X$  is computable too.

**Corollary 3.10** (Completion).  $(\overline{X}, \delta_{\overline{X}})$  is a precomplete represented space for every represented space  $(X, \delta_X)$  and  $\iota : X \to \overline{X}, x \mapsto x$  is a computable embedding.

# 4. Total Weihrauch Reducibility

In this section we are going to introduce a total variant of Weihrauch reducibility that behaves very similarly to the usual reducibility from a practical perspective, but that has different algebraic properties.

By a problem  $f :\subseteq X \Rightarrow Y$  we mean a partial multi-valued map  $f :\subseteq X \Rightarrow Y$  on represented spaces  $(X, \delta_X)$  and  $(Y, \delta_Y)$ . We recall that *composition* of problems  $f :\subseteq X \Rightarrow Y$  and  $g :\subseteq Y \Rightarrow Z$  is defined by

$$g \circ f(x) := \{ z \in Z : (\exists y \in f(x)) \ z \in g(y) \}$$

for all  $x \in \text{dom}(g \circ f) := \{x \in \text{dom}(f) : f(x) \subseteq \text{dom}(g)\}$ . For two problems  $f :\subseteq X \rightrightarrows Y$  and  $g :\subseteq X \rightrightarrows Z$  with identical source space X we define the juxtaposition  $(f,g) :\subseteq X \rightrightarrows Y \times Z$  by  $(f,g)(x) := f(x) \times g(x)$  and  $\text{dom}(f,g) := \text{dom}(f) \cap \text{dom}(g)$ . If  $f,g :\subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$  are problems on Baire space, then we also call  $\langle f,g \rangle := \langle \ \rangle \circ (f,g)$  the juxtaposition of f and g and  $\langle f \times g \rangle$  defined by  $\langle f \times g \rangle \langle p,q \rangle := \langle f(p),g(q) \rangle$  for all  $p,q \in \mathbb{N}^{\mathbb{N}}$  the product of f and g.

We say that a function  $F : \subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  is a realizer of f, if  $\delta_Y F(p) \in f\delta_X(p)$  for all  $p \in \text{dom}(f\delta_X)$ . We denote this by  $F \vdash f$ . We say that f is computable if it has a computable realizer. Other notions, such as continuity, Borel measurability

and so forth that are well-defined for functions  $F:\subseteq \mathbb{N}^{\mathbb{N}}\to \mathbb{N}^{\mathbb{N}}$  are transferred in an analogous manner to problems  $f :\subseteq X \rightrightarrows Y$ .

We write  $F \vdash_{\mathbf{t}} f$ , if F is a total realizer of f. We now recall the definition of ordinary and strong Weihrauch reducibility on problems f, g, which is denoted by  $f \leq_{W} g$  and  $f \leq_{sW} g$ , respectively, and we introduce two new concepts of total Weihrauch reducibility and strong total Weihrauch reducibility, which are denoted by  $f \leq_{\text{tW}} g$  and  $f \leq_{\text{stW}} g$ , respectively.

**Definition 4.1** (Weihrauch reducibility). Let  $f :\subseteq X \rightrightarrows Y$  and  $g :\subseteq U \rightrightarrows V$  be problems. We define:

- (1)  $f \leq_{\mathbf{W}} g : \iff (\exists \text{ computable } H, K : \subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}})(\forall G \vdash g) \ H \langle \text{id}, GK \rangle \vdash f.$
- (1)  $f \subseteq_{\text{sW}} g : \iff (\exists \text{ computable } H, K : \subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}})(\forall G \vdash g) H \land K, GK \not\vdash f.$ (2)  $f \subseteq_{\text{sW}} g : \iff (\exists \text{ computable } H, K : \subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}})(\forall G \vdash_{\text{t}} g) H \land id, GK \land \vdash_{\text{t}} f.$ (3)  $f \subseteq_{\text{tW}} g : \iff (\exists \text{ computable } H, K : \subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}})(\forall G \vdash_{\text{t}} g) H \land id, GK \land \vdash_{\text{t}} f.$ (4)  $f \subseteq_{\text{stW}} g : \iff (\exists \text{ computable } H, K : \subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}})(\forall G \vdash_{\text{t}} g) H \land GK \land \vdash_{\text{t}} f.$

For (3) and (4) we assume that we replace each of the given representations of X, Y, U and V by a computably equivalent precomplete representation of the corresponding set.

We call the reducibilities  $\leq_{W}$  and  $\leq_{sW}$  partial<sup>7</sup> in order to distinguish them from their total counterparts  $\leq_{tW}$  and  $\leq_{stW}$ . We note that precompleteness is not required or relevant in the partial case, but it can be assumed without loss of generality since the concept of partial (strong) Weihrauch reducibility is invariant under computably equivalent representations [3, Lemma 2.11]. In the total cases (3) and (4), however, precompleteness is essential, since otherwise these definitions would not be invariant under computably equivalent representations. By Proposition 3.4 we can always choose precomplete representations that are computably equivalent to the given representations of the spaces X, Y, U and V. But we still need to show that the definition of  $\leq_{tW}$  and  $\leq_{stW}$  does not depend on this choice.

We will prove a slightly more general result that highlights the places where precompleteness is actually needed. For this purpose we introduce the following terminology: we say that  $f \leq_{tW} g$  holds with respect to  $(\delta_X, \delta_Y, \delta_U, \delta_V)$ , if Definition 4.1 (3) holds as it stands but exactly for the given representations of X, Y, Uand V, respectively, and these representations are not required to be precomplete. Hence the statement defined here is weaker than  $f \leq_{tW} g$  in the sense defined above. We use a corresponding terminology for  $\leq_{\text{stW}}$ . Now we obtain the following result.

**Lemma 4.2** (Invariance under representations). Let  $f :\subseteq X \rightrightarrows Y$ ,  $g :\subseteq U \rightrightarrows V$  be problems on represented spaces  $(X, \delta_X)$ ,  $(Y, \delta_Y)$ ,  $(U, \delta_U)$  and  $(V, \delta_V)$ . Let  $\delta_X'$ ,  $\delta_Y'$ ,  $\delta_U'$ and  $\delta'_V$  be further representations of the given sets, respectively, such that

- (1)  $\delta_X' \leq \delta_X$ ,  $\delta_Y \leq \delta_Y'$ ,  $\delta_U \leq \delta_U'$  and  $\delta_V' \leq \delta_V$ , (2)  $\delta_V$ ,  $\delta_Y'$  and  $\delta_U'$  are precomplete.

If  $f \leq_{tW} g$  holds with respect to  $(\delta_X, \delta_Y, \delta_U, \delta_V)$ , then it also holds with respect to  $(\delta'_X, \delta'_Y, \delta'_U, \delta'_U)$ . An analogous statement holds for  $f \leq_{\text{stW}} g$ .

Proof. We follow the construction as outlined in the proof of [3, Lemma 2.11]. Since  $\delta'_U$  and  $\delta'_V$  are precomplete according to (2), we can additionally assume that the computable functions  $S,T:\mathbb{N}^{\mathbb{N}}\to\mathbb{N}^{\mathbb{N}}$  in that proof are total. In that proof it is shown that whenever  $G' \vdash g$  holds with respect to  $(\delta'_U, \delta'_V)$ , then  $G := TG'S \vdash g$ holds with respect to  $(\delta_U, \delta_V)$ . Due to totality of T, S, the same holds true if we replace  $\vdash$  by  $\vdash_{t}$  in both occurrences. If we assume that  $H(\mathrm{id}, GK) \vdash_{t} f$  holds with respect to  $(\delta_X, \delta_Y)$ , then we obtain as in the proof mentioned above that  $H'(\mathrm{id}, G'K') \vdash f$  holds with respect to  $(\delta'_X, \delta'_Y)$ . Due to precompleteness of  $\delta'_U$  and

<sup>&</sup>lt;sup>7</sup>This is not related to the preorders being partial or total in an order theoretic sense; they are both partial in that sense.

 $\delta'_Y$  according to (2), we can always assume that H', K' are even total computable functions. Hence, we even obtain  $H'\langle \operatorname{id}, G'K' \rangle \vdash_{\operatorname{t}} f$ , which completes the proof. The proof for  $\leq_{\operatorname{stW}}$  is analogous.

If  $f \leq_{\text{tW}} g$  holds with respect to  $(\delta_X, \delta_Y, \delta_U, \delta_V)$  and at least  $\delta_V$  is precomplete among these representations, then according to Lemma 4.2 we can always replace the non-precomplete representations by equivalent precomplete ones and  $f \leq_{\text{tW}} g$  holds with respect to these precomplete representations and hence  $f \leq_{\text{tW}} g$  holds in terms of Definition 4.1.

For the moment Lemma 4.2 is useful as it implies that  $\leq_{tW}$  and  $\leq_{stW}$  are well-defined and invariant under computably equivalent representations.

**Corollary 4.3** (Invariance under equivalent representations). Let  $f :\subseteq X \rightrightarrows Y$  and  $g :\subseteq U \rightrightarrows V$  be problems. The relations  $f \subseteq_W g$ ,  $f \subseteq_{sW} g$ ,  $f \subseteq_{tW} g$  and  $f \subseteq_{stW} g$  remain unchanged if we replace the representations of X, Y, U and V by computably equivalent ones.

We note that the statement for  $\leq_W$  and  $\leq_{sW}$  was proved in [3, Lemma 2.11]. The following example shows that precompleteness in Definition 4.1 cannot be omitted if one wants to achieve invariance under equivalent representations.

**Example 4.4.** Every computable function  $F:\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  without total computable extension (see Example 3.2) has a total computable realizer with respect to (id, id $^{\wp}$ ), but not with respect to (id, id). Hence  $F \leq_{\mathrm{tW}} F$  does not hold with respect to (id, id, id, id $^{\wp}$ ). Clearly,  $F \leq_{\mathrm{tW}} F$  holds with respect to (id $^{\wp}$ , id $^{\wp}$ , id $^{\wp}$ , id $^{\wp}$ ) and hence  $F \leq_{\mathrm{tW}} F$  holds in terms of Definition 4.1.

The argument used in the proof of Lemma 4.2 concerning H' and K' also allows us to slightly rephrase Definition 4.1. Due to precompleteness we can demand total H, K (and replace  $\vdash_t$  by  $\vdash$  on the right-hand side.)

**Lemma 4.5** (Weihrauch reducibility). Let  $f :\subseteq X \Rightarrow Y$  and  $g :\subseteq U \Rightarrow V$  be problems. We choose precomplete representations that are computably equivalent to the given representations of X, Y, U and V. Then:

- (1)  $f \leq_{\mathbf{W}} g \iff (\exists computable H, K : \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}})(\forall G \vdash g) H \langle id, GK \rangle \vdash f.$
- $(2) \ f \leq_{\mathrm{sW}} g \iff (\exists \ computable \ H, K : \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}) (\forall G \vdash g) \ HGK \vdash f.$
- (3)  $f \leq_{\mathsf{tW}} g \iff (\exists \ computable \ H, K : \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}) (\forall G \vdash_{\mathsf{t}} g) \ H \langle \mathrm{id}, GK \rangle \vdash f.$
- (4)  $f \leq_{\text{stW}} g \iff (\exists computable H, K : \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}})(\forall G \vdash_{\text{t}} g) HGK \vdash f.$

The proof of the backward direction is immediate and the forward direction follows from precompleteness of the representations of U and Y, respectively.

In [3, Lemma 2.4] we have proved that  $\leq_{\rm W}$  and  $\leq_{\rm sW}$  are preorders, i.e., they are reflexive and transitive. The associated equivalences are denoted by  $\equiv_{\rm W}$  and  $\equiv_{\rm sW}$ , respectively. Using Lemma 4.5 we can now easily transfer these proofs to the case of the total reducibilities.

**Proposition 4.6** (Preorders). The relations  $\leq_{tW}$  and  $\leq_{stW}$  are preorders on the class of problems.

*Proof.* We follow the proof of [3, Lemma 2.4] and the notations used therein. Reflexivity is obvious as the corresponding functions H, K are total. For the transitivity proof, we assume that the reductions  $f \leq_{\text{tW}} g$  and  $g \leq_{\text{tW}} h$  are given by total H, K, H', K'. Then the corresponding functions H'' and K'' constructed in the proof of [3, Lemma 2.4] are also total and hence the claim follows from Lemma 4.5.

By  $\equiv_{tW}$  and  $\equiv_{stW}$  we denote the equivalence relations that are associated with  $\leq_{tW}$  and  $\leq_{stW}$ , respectively. If the different versions of Weihrauch reducibility are

expressed as in Lemma 4.5, then it is immediately clear that a partial reduction implies the corresponding total reduction. Using Lemma 4.5, Corollary 4.3 and Proposition 3.4 obtain the following corollary.

Corollary 4.7 (Partial and total Weihrauch reducibility). Let f and g be problems. Then  $f \leq_{W} g \Longrightarrow f \leq_{tW} g$  and  $f \leq_{sW} g \Longrightarrow f \leq_{stW} g$ .

This means that all positive results that hold for a partial version of Weihrauch reducibility can be transferred to the corresponding total variant. Together with the obvious other implications we obtain the diagram for the logical relations between different versions of Weihrauch reducibility that is displayed in Figure 2. The diagram is complete up to transitivity (see Example 4.8). The diagram also shows the generating closure operators of cylindrification and completion that we discuss later.




FIGURE 2. Implications between notions of reducibility

**Example 4.8.** Let  $f: \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  denote a constant function with computable value. Then  $\mathrm{id} \equiv_{\mathrm{W}} f$ , but  $\mathrm{id} \not\leq_{\mathrm{stW}} f$ . Let  $0:\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  denote the nowhere defined function. Then  $\mathrm{id} \equiv_{\mathrm{tW}} 0$ , but  $\mathrm{id} \not\leq_{\mathrm{W}} 0$ . Let  $\mathrm{id}|_{\{p\}} :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  be the identity restricted to a non-computable  $p \in \mathbb{N}^{\mathbb{N}}$ . Then  $\mathrm{id} \equiv_{\mathrm{stW}} \mathrm{id}|_{\{p\}} \times \mathrm{id}$ , but  $\mathrm{id} \not\leq_{\mathrm{W}} \mathrm{id}|_{\{p\}} \times \mathrm{id}$ .

We note that the reducibilities  $\leq_{\text{tW}}$  and  $\leq_{\text{stW}}$  share similar properties as  $\leq_{\text{W}}$  and  $\leq_{\text{sW}}$  when it comes to the preservation of computability or other properties. We say that a class  $\mathcal C$  of problems is *preserved downwards* by a reducibility  $\leq$  for problems if  $f \leq g$  and  $g \in \mathcal C$  imply  $f \in \mathcal C$ .

**Proposition 4.9** (Downwards preservation). Computability, continuity, limit computability, Borel measurability and non-uniform computability are preserved downwards by  $\leq_{tW}$ .

*Proof.* Let  $\mathcal{C}$  be the class of computable, continuous, limit computable, Borel measurable or non-uniformly computable problems. We choose precomplete representations and total computable H, K that witness  $f \leq_{\mathrm{tW}} g$  according to Lemma 4.5. If  $g \in \mathcal{C}$ , then it has a realizer  $G :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  that is in  $\mathcal{C}$ . Since the target space of g is represented with a precomplete representation, we can assume without loss of generality that G is total by Proposition 3.6. Hence  $H\langle \mathrm{id}, GK \rangle$  is a (even total) realizer of f that is also in the class  $\mathcal{C}$ . This proves that  $f \in \mathcal{C}$ .

Any class  $\mathcal{C}$  of functions  $F :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  constitutes a property of problems that is preserved downwards by total Weihrauch reducibility if the following conditions are satisfied:  $\mathcal{C}$  contains the identity, is closed under composition with computable functions, is closed under juxtaposition with the identity and  $\mathcal{C}$  respects precompleteness. In [4, Corollaries 6.2, 7.4, 8.3] we prove that finite mind change computability and Las Vegas computability is not preserved downwards by  $\leq_{\mathrm{tW}}$ , whereas non-deterministic computability is preserved.

It is known that the class of the nowhere defined problems (often denoted by  $\mathbf{0}$ ) forms the bottom element of the Weihrauch lattice [3, Lemma 2.7], while the Weihrauch equivalence class of id (often denoted by  $\mathbf{1}$ ) is the class of all computable problems with at least one computable point in the domain [3, Lemma 2.8]. Moreover, a problem f is computable if and only if  $f \leq_{\mathbf{W}} \mathrm{id}$ . The statement about the nowhere defined function  $0 :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  in Example 4.8, namely that  $\mathrm{id} \equiv_{\mathrm{tW}} 0$ , together with Proposition 4.9 show that the minimal equivalence class with respect to total Weihrauch reducibility is the class of all computable problems.

Corollary 4.10 (Minimal total degree). The equivalence class of all computable problems forms the minimal element with respect to total Weihrauch reducibility.

This already shows that the algebraic structure induced by total Weihrauch reducibility is significantly different from the structure induced by partial Weihrauch reducibility. In between **0** and **1** one obtains a complicated structure for partial Weihrauch reducibility, and among other results one can show that one can embed the entire Medvedev lattice (and hence the Turing semi-lattice) in an order-reversing way into the Weihrauch lattice between **0** and **1** [13, Lemma 5.6]. In contrast to this the two degrees **0** and **1** fall together with respect to total Weihrauch reducibility.

Strictly speaking, the class of problems is not a set, but we can always consider representatives of problems on Baire space to obtain a set as underlying structure. This is known for  $\leq_{\rm W}$  and  $\leq_{\rm sW}$  (see [6, Lemma 3.8]) and holds correspondingly for  $\leq_{\rm tW}$  and  $\leq_{\rm stW}$ .

**Corollary 4.11** (Realizer version). Let  $f :\subseteq X \rightrightarrows Y$  be a problem on represented spaces  $(X, \delta_X)$  and  $(Y, \delta_Y)$ . Then  $f^r := \delta_Y^{-1} \circ f \circ \delta_X :\subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$  satisfies  $f^r \equiv_{\operatorname{stW}} f$ .

*Proof.* By  $f^r \equiv_{sW} f$  holds according to [6, Lemma 3.8] (and is easy to see, since  $f^r$  and f share exactly the same realizers). Hence  $f^r \equiv_{stW} f$  follows by Corollary 4.7.

We note that we do not need to assume that  $\delta_X$  and  $\delta_Y$  are precomplete in this result. However, for  $f^r :\subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$  we need to use precomplete representations of  $\mathbb{N}^{\mathbb{N}}$  for the total versions of Weihrauch reducibility.

## 5. Completion

In this section we discuss the closure operation of completion  $f \mapsto \overline{f}$  that generates  $\leq_{tW}$  on  $\leq_{w}$  and  $\leq_{stW}$  on  $\leq_{sw}$ . For the definition of the completion  $\overline{f}$  we use the completion  $\overline{X}$  of a represented space according to Definition 3.7.

**Definition 5.1** (Completion). Let  $f :\subseteq X \Rightarrow Y$  be a problem. We define the *completion* of f by

$$\overline{f}: \overline{X} \rightrightarrows \overline{Y}, x \mapsto \left\{ \begin{array}{ll} f(x) & \text{if } x \in \text{dom}(f) \\ \overline{Y} & \text{otherwise} \end{array} \right.$$

We note that the completion  $\overline{f}$  is always *pointed*, i.e., it has a computable point in its domain. This is because  $\bot \in \overline{X}$  is always computable (as it has the constant zero sequence as a name).

Sometimes it is useful to think of  $\overline{f}$  in terms of its realizer version  $\overline{f}^{\,\mathrm{r}}:\mathbb{N}^{\mathbb{N}}\rightrightarrows\mathbb{N}^{\mathbb{N}}$ , which is given by

$$\overline{f}^{\,\mathrm{r}}(p) = \delta_{\overline{Y}}^{-1} \circ \overline{f} \circ \delta_{\overline{X}}(p) = \left\{ \begin{array}{ll} (\delta_{Y}^{\wp})^{-1} \circ f \circ \delta_{X}^{\wp}(p) & \text{if } p \in \mathrm{dom}(f \circ \delta_{X}^{\wp}) \\ \mathbb{N}^{\mathbb{N}} & \text{otherwise} \end{array} \right.$$

Since  $\overline{f}$  has exactly the same realizers as  $\overline{f}^{r}$ , one can deduce from this formula that the realizers of  $\overline{f}$  are exactly the total realizers of f with respect to  $\delta_{X}^{\wp}$  and  $\delta_{Y}^{\wp}$ , which immediately yields the following conclusion with the help of Lemma 4.5.

**Lemma 5.2** (Completion and total Weihrauch reducibility). For all problems f, g:  $f \leq_{\operatorname{W}} \overline{g} \iff \overline{f} \leq_{\operatorname{W}} \overline{g} \iff f \leq_{\operatorname{tW}} g \text{ and } f \leq_{\operatorname{sW}} \overline{g} \iff \overline{f} \leq_{\operatorname{sW}} \overline{g} \iff f \leq_{\operatorname{stW}} g.$ 

Thus, we could define total Weihrauch reducibility also using the completion operation and partial Weihrauch reducibility. Lemma 5.2 also shows that the total Weihrauch degrees can be order theoretically embedded into the pointed partial Weihrauch degrees. Together with Corollary 4.7 we obtain that completion is monotone.

Corollary 5.3 (Monotonicity of completion). Let f and g be problems. Then

- $\begin{array}{ll} (1) & f \leq_{\mathcal{W}} g \Longrightarrow \overline{f} \leq_{\mathcal{W}} \overline{g}, \\ (2) & f \leq_{s\mathcal{W}} g \Longrightarrow \overline{f} \leq_{s\mathcal{W}} \overline{g}. \end{array}$

We note that this result also implies that completion is a well-defined operation on (strong) Weihrauch degrees: if  $f_1, f_2$  are identical problems with possibly different but computably equivalent representations on the input and output side, respectively, then  $f_1 \equiv_{sW} f_2$  and hence  $\overline{f_1} \equiv_{sW} \overline{f_2}$  follows. This is so, even so the representations of the corresponding completions of the spaces on the input and output side are not necessarily computably equivalent (see the remark after Definition 3.7). Now we can see that completion is a closure operator.

**Proposition 5.4** (Completion as closure operator). Completion  $f \mapsto \overline{f}$  is a closure operator on  $\leq_{W}$  and  $\leq_{sW}$ .

*Proof.* By Lemma 5.2  $f \leq_{sW} \overline{f}$  is equivalent to  $f \leq_{stW} f$ , which holds since  $\leq_{stW}$ is reflexive by Proposition 4.6. By Lemma 5.2  $\overline{f} \leq_{\text{sW}} \overline{f}$  is equivalent to  $\overline{f} \leq_{\text{sW}} \overline{f}$ , which holds since  $\leq_{sW}$  is reflexive. Completion is monotone with respect to  $\leq_{w}$ and  $\leq_{sW}$  according to Corollary 5.3. Altogether completion is a closure operator with respect to  $\leq_{W}$  and  $\leq_{sW}$ .

We have used properties of  $\leq_{tW}$  and  $\leq_{stW}$  in order to obtain properties of completion. Vice versa Proposition 5.4 and Lemma 5.2 also imply Proposition 4.6 and Corollary 4.7. Hence, these concepts yield different perspectives on the same properties.

It is clear that every f is strongly totally equivalent to its completion by Lemma 5.2 and Proposition 5.4.

Corollary 5.5.  $f \equiv_{\text{stW}} \overline{f}$  for every problem f.

In the study of total Weihrauch reducibility the degrees that have identical cones with respect to partial and total Weihrauch reducibility play an important role. Hence, we introduce a name for such degrees.

**Definition 5.6** (Complete problems). A problem f is called *complete* if  $f \equiv_{\mathbf{W}} \overline{f}$ and strongly complete if  $f \equiv_{sW} \overline{f}$ .

Now we obtain the following straightforward characterization of completeness.

**Theorem 5.7** (Completeness). Let g be a problem. Then the following hold:

- (1)  $g \ complete \iff (\forall \ problems \ f)(f \leq_{W} g \iff f \leq_{tW} g).$
- (2)  $g \text{ strongly complete} \iff (\forall \text{ problems } f)(f \leq_{\text{sW}} g \iff f \leq_{\text{stW}} g).$

*Proof.* If g is (strongly) complete, then the respective given equivalence holds by Lemma 5.2. On the other hand, if  $f \leq_{W} g \iff f \leq_{tW} g$  holds for all f, then  $\overline{g} \equiv_{W} g$  follows from Corollary 5.5. The case of strong completeness can be handled analogously.

Examples of complete problems are abundant. We study a number of landmarks in the Weihrauch lattice, among them the Turing jump operator J and and the binary sorting problem SORT that was introduced and studied by Neumann and Pauly [18]. The problems  $WBWT_2$ ,  $ACC_X$ , PA and MLR were studied for instance in [7]. We identify  $X \in \mathbb{N}$  with the set  $X = \{0, 1, ..., X - 1\}$ . Many further completeness questions regarding choice are studied in [4].

**Proposition 5.8** (Complete problems). The following problems are all strongly complete:

```
(1) id: \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}, p \mapsto p,
```

(2) 
$$J: \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}, p \mapsto p',$$

(2) 
$$J: \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}, p \mapsto p',$$
  
(3)  $\lim_{n \to \infty} \subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}, \langle p_0, p_1, p_2, ... \rangle \mapsto \lim_{n \to \infty} p_n,$ 

$$(4) \ \mathsf{LPO} : \mathbb{N}^{\mathbb{N}} \to \{0,1\}, \mathsf{LPO}(p) = 0 : \iff (\exists n \in \mathbb{N}) \ p(n) = 0,$$

(5) SORT:  $2^{\mathbb{N}} \to 2^{\mathbb{N}}$  with

$$\mathsf{SORT}(p) := \left\{ \begin{array}{ll} 0^k \widehat{1} & \textit{if $p$ contains exactly $k \in \mathbb{N}$ zeros} \\ \widehat{0} & \textit{if $p$ contains infinitely many zeros} \end{array} \right.$$

- (6)  $\mathsf{WBWT}_2: 2^{\mathbb{N}} \rightrightarrows 2^{\mathbb{N}}, p \mapsto \{q \in 2^{\mathbb{N}} : \lim_{n \to \infty} q(n) \text{ is a cluster point of } p\}.$

*Proof.* (1) Follows since  $\overline{\mathrm{id}} \leq_{\mathrm{sW}} \mathrm{id}_{\overline{\mathrm{NN}}} \leq_{\mathrm{sW}} \mathrm{id}$ .

- (2) There is a total computable function  $H: \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  such that  $H \circ \mathsf{J}(p) =$ J(p-1)+1 for all p with  $p-1\in\mathbb{N}^{\mathbb{N}}$ . This can be proved using the smn-Theorem. Together with the identity K this function H witnesses the reduction  $\overline{J} \leq_{sW} J$ .
- (3) Follows by Corollary 5.3 since  $\lim \equiv_{sW} J$  holds (see [6, Theorem 6.7]).
- (4) Given a name  $p \in \mathbb{N}^{\mathbb{N}}$  of a point in  $\overline{\mathbb{N}^{\mathbb{N}}}$  with respect to  $\delta_{\overline{\mathbb{N}^{\mathbb{N}}}}$ , we can compute K(p) as follows:  $K(p)(n) = 0 : \iff p(n) = 1$  and K(p)(n) := 1 otherwise. If  $\delta_{\overline{\mathbb{N}^{\mathbb{N}}}}(p) = q \in \mathbb{N}^{\mathbb{N}}$ , then  $\mathsf{LPO} \circ K(p) = \mathsf{LPO}(q)$ . Hence, together with H(r) := r + 1the functions H, K witness  $\overline{\mathsf{LPO}} \leq_{\mathrm{sW}} \mathsf{LPO}$ .
- (5) As always we assume that  $2^{\mathbb{N}}$  is represented by  $\delta_{2^{\mathbb{N}}} :\subseteq \mathbb{N}^{\mathbb{N}} \to 2^{\mathbb{N}}, p \mapsto p$  with  $\operatorname{dom}(\delta_{2^{\mathbb{N}}}) = 2^{\mathbb{N}}$ . Given a name  $p \in \mathbb{N}^{\mathbb{N}}$  of some  $q \in \overline{2^{\mathbb{N}}}$ , i.e.,  $\delta_{2^{\mathbb{N}}}(p) = q$  we can compute K(p) as follows:  $K(p)(n) = 0 : \iff p(n) = 1$  and K(p)(n) := 1 otherwise. Then  $\mathsf{SORT} \circ K(p) = \mathsf{SORT}(q)$  if  $q \in 2^{\mathbb{N}}$ . Hence, H(r) := r + 1 and K witness  $\overline{\mathsf{SORT}}\mathop{\leq_{\mathrm{sW}}}\mathsf{SORT}.$
- (6) We represent  $2^{\mathbb{N}}$  as above. Given a name  $p \in \mathbb{N}^{\mathbb{N}}$  of some  $q \in \overline{2^{\mathbb{N}}}$  we can compute K(p) as follows, we let K(p)(n) := p(n) - 1 if  $p(n) \neq 0$  and we let K(p)(n) = ifor the number  $i \in \{0,1\}$  such that i+1 appears a maximal number of times within p(0),...,p(n) (and we choose i=0 if 1 and 2 appear equally often). This construction guarantees that we do not generate any additional cluster points, i.e.,  $\mathsf{WBWT}_2K(p) = \mathsf{WBWT}_2(q)$  for  $q \in 2^{\mathbb{N}}$ . Similarly as in the other cases above, this proves  $\overline{\mathsf{WBWT}_2} \leq_{\mathrm{sW}} \mathsf{WBWT}_2$ .
- (7) Given some name  $p \in \mathbb{N}^{\mathbb{N}}$  of a point  $q \in \overline{\mathbb{N}^{\mathbb{N}}}$  we compute K(p) as follows: we let K(p)(n) := k + 1 if k + 2 = p(n) is the first number larger than 1 among p(0),...,p(n) and  $k \in X$ . Otherwise, we let K(p)(n) := 0. This guarantees that  $ACC_XK(p) = ACC_X(q)$ , if  $q \in dom(ACC_X)$ . Similarly as in the other cases above, this proves  $\overline{\mathsf{ACC}_X} \leq_{\mathrm{sW}} \mathsf{ACC}_X$ .
- (8), (9) We use  $K: \mathbb{N}^{\mathbb{N}} \to 2^{\mathbb{N}}, p \mapsto 0^{p(0)+1}10^{p(1)+1}10^{p(2)+1}...$ , which is total computable. It is straightforward to see that every problem  $F:2^{\mathbb{N}}\rightrightarrows 2^{\mathbb{N}}$  that is antitone in the sense that  $p \leq_T q$  implies  $F(q) \subseteq F(p)$  is strongly complete. This is because

 $p-1 \leq_{\mathrm{T}} p \equiv_{\mathrm{T}} K(p)$  if  $p \in \mathbb{N}^{\mathbb{N}}$  is such that  $p-1 \in 2^{\mathbb{N}}$ , and hence  $FK(p) \subseteq F(p-1)$ . This proves  $\overline{F} \leq_{\text{sW}} F$ . This applies in particular to PA and MLR.

These results show that the cones below the given problems are identical in the total and partial Weihrauch lattices. It is known, for instance, that f is limit computable if and only if  $f \leq_{\mathbf{W}} \lim [6]$ . Hence, an analogous statement holds for

## 6. Algebraic Operations

In this section we want to discuss properties of certain algebraic operations and we want to prove that the total versions of Weihrauch reducibility yield lattice structures. We start recalling the usual algebraic operations on the Weihrauch lattice [6].

**Definition 6.1** (Algebraic operations). Let  $f:\subseteq X \rightrightarrows Y$  and  $g:\subseteq U \rightrightarrows V$  be multi-valued functions. We define the following operations:

- (1)  $f \times g :\subseteq X \times U \rightrightarrows Y \times V, (f \times g)(x, u) := f(x) \times g(u)$  and  $dom(f \times g) := dom(f) \times dom(g)$ (product)
- $(2) \ \ f \sqcup g \stackrel{\sim}{:}\subseteq X \sqcup U \rightrightarrows Y \sqcup V, \ (f \sqcup g)(0,x) := \{0\} \times f(x), \ (f \sqcup g)(1,u) := \{1\} \times g(u)$ and  $dom(f \sqcup g) := dom(f) \sqcup dom(g)$
- $(3) \ f \boxplus g : \subseteq \overrightarrow{X} \sqcup \overrightarrow{U} \rightrightarrows \overrightarrow{Y} \times \overrightarrow{V}, (f \boxplus g)(0, x) := f(x) \times \overrightarrow{V}, (f \boxplus g)(1, u) := \overrightarrow{Y} \times g(u)$ and  $dom(f \boxplus g) := dom(f) \sqcup dom(g)$ (box sum)
- (4)  $f \sqcap g :\subseteq X \times U \Rightarrow Y \sqcup V, (f \sqcap g)(x, u) := f(x) \sqcup g(u)$  and  $dom(f \sqcap g) := dom(f) \times dom(g)$
- (5)  $f + g :\subseteq X \times U \Rightarrow \overline{Y} \times \overline{V}, (f + g)(x, u) := (f(x) \times \overline{V}) \cup (\overline{Y} \times g(u))$  and  $dom(f+g) := dom(f) \times dom(g)$ (sum)
- (finite parallelization)
- $\begin{aligned}
  \operatorname{dom}(f+g) &:= \operatorname{dom}(f) \times \operatorname{dom}(g) \\
  (6) \quad f^* &:\subseteq X^* \rightrightarrows Y^*, f^*(i,x) := \{i\} \times f^i(x) \text{ and } \\
  \operatorname{dom}(f^*) &:= \operatorname{dom}(f)^* \\
  (7) \quad \widehat{f} &:\subseteq X^{\mathbb{N}} \rightrightarrows Y^{\mathbb{N}}, \widehat{f}(x_n)_{n \in \mathbb{N}} := \mathsf{X}_{i \in \mathbb{N}} f(x_i) \text{ and } \\
  \operatorname{dom}(\widehat{f}) &:= \operatorname{dom}(f)^{\mathbb{N}}
  \end{aligned}$ (parallelization)

For every operation  $\square \in \{\times, \sqcup, \boxplus, \sqcap, +\}$  we define its completion  $\overline{\square}$  by  $f\overline{\square}g :=$  $\overline{f} \square \overline{g}$ . It follows from Lemma 5.2 that these operations are monotone with respect to total Weihrauch reducibility, since the underlying operations □ are monotone with respect to partial Weihrauch reducibility by [6, Proposition 3.6].

Corollary 6.2 (Monotonicity).  $(f,g) \mapsto f\overline{\square}g$  for  $\square \in \{\times, \sqcup, \boxplus, \sqcap, +\}, f \mapsto \widehat{\overline{f}}$  and  $f \mapsto \overline{f}^*$  are monotone with respect to  $\leq_{\text{tW}}$  and  $\leq_{\text{stW}}$ .

*Proof.* By Lemma 5.2 completion generates  $\leq_{tW}$  on  $\leq_{W}$  (and  $\leq_{stW}$  on  $\leq_{sW}$ ). By [6, Proposition 3.6] the given operations  $\square$  are monotone with respect to  $\leq_{\rm W}$  and  $\leq_{\rm sW}$ , respectively. The claim now follows with Proposition 2.2.

Now we prove that the algebraic operations preserve completeness in the sense of Definition 2.3. It is clear by Proposition 2.5 that we also get co-preservation for suprema (see Proposition 6.5). Later we will show that this also holds for + (see Proposition 6.8).

**Proposition 6.3** (Completion and algebraic operations). Let f and q be problems. We obtain

- $(1) \ \overline{f \square g} \leq_{\mathrm{sW}} \overline{f} \square \overline{g} \equiv_{\mathrm{sW}} \overline{\overline{f} \square \overline{g}} \ \text{for} \ \square \in \{\times, \sqcup, \boxplus, \sqcap, +\},$
- (2)  $\overline{\widehat{f}} \leq_{sW} \widehat{\overline{f}} \equiv_{sW} \overline{\widehat{\overline{f}}},$ (3)  $\overline{f^*} \leq_{sW} \overline{f}^* \equiv_{sW} \overline{\overline{f}^*}.$

In particular, if f and g are (strongly) complete, then so are  $f \times g$ ,  $f \sqcup g$ ,  $f \boxtimes g$ ,  $f \cap g$ , f + g,  $\widehat{f}$  and  $f^*$ .

Proof. We consider problems  $f:\subseteq X\rightrightarrows Y$  and  $g:\subseteq U\rightrightarrows V$  and  $\square\in\{\times,\sqcup,\boxplus,\sqcap,+\}$ . Since  $X\sqcup U\subseteq \overline{X}\sqcup \overline{U}$  and  $X\times U\subseteq \overline{X}\times \overline{U}$ , it follows that  $\mathrm{dom}(f\square g)\subseteq \mathrm{dom}(\overline{f}\square \overline{g})$ , and restricted to  $x\in \mathrm{dom}(f\square g)$  we have  $\overline{f\square g}(x)=(f\square g)(x)\subseteq (\overline{f}\square \overline{g})(x)$ . The " $\subseteq$ " is even an equality in the cases  $\square\in\{\times,\sqcup,\sqcap\}$ . In the other cases it is not an equality simply because  $\overline{V}\subsetneq\overline{V}$  and  $\overline{Y}\subsetneq\overline{V}$ . We can also assume that the representations of  $\overline{X}\times\overline{U}$  and  $\overline{X}\sqcup\overline{U}$  are total (since the representations of  $\overline{X}$  and  $\overline{U}$  are so). Hence every realizer of  $\overline{f}\square \overline{g}$  is total. By Corollary 3.10  $\iota:Z\to\overline{Z},z\mapsto z$  is computable for every represented space Z, hence it follows that  $\overline{f}\square \overline{g}\leq_{\mathrm{sW}} \overline{f}\square \overline{g}$ , since a realizer for  $\overline{f}\square g$  can choose any value outside of  $\mathrm{dom}(f\square g)$ . This also holds in the cases where we only have " $\subseteq$ " above, since the representation of  $\overline{V}$  is total, every name of a point in  $\overline{V}$  is also a name of some point in  $\overline{V}$  and an analogous statement holds for Y. The proofs for the unary operations are analogous. We have  $X^{\mathbb{N}}\subseteq \overline{X}^{\mathbb{N}}$  and  $X^*\subseteq \overline{X}^*$  and hence  $\overline{f}\leq_{\mathrm{sW}}\widehat{f}$  and  $\overline{f^*}\leq_{\mathrm{sW}}\overline{f}^*$ . The remaining claims follow by Proposition 2.4 as completion is a closure operator by Proposition 5.4.  $\square$ 

The closure properties of complete problems are very useful. For instance, it is known that  $\lim_{s \to w} \widehat{\mathsf{LPO}}$  [6] and hence the statement on  $\lim$  in Proposition 5.8 could also be derived from the statement on  $\mathsf{LPO}$ . Likewise, we obtain a number of further complete problems in this way. We refrain from giving exact definitions of the listed problems, but we rather point the reader to [7] were all stated equivalences have been proved [7, Theorem 5.2, Corollary 5.3, Proposition 14.10]. For the purpose of this article, the equivalences can be read as definitions.

Corollary 6.4 (Complete problems). WKL  $\equiv_{sW} C_{2^{\mathbb{N}}} \equiv_{sW} \widehat{\mathsf{ACC}_2}$ ,  $\mathsf{DNC}_X \equiv_{sW} \widehat{\mathsf{ACC}_X}$  for  $X \in \mathbb{N}$  with  $X \geq 2$  or  $X = \mathbb{N}$  are strongly complete, and  $\mathsf{COH} \equiv_{\mathsf{W}} \widehat{\mathsf{WBWT}_2}$  is complete.

In [3, Proposition 3.11] we proved that  $\sqcap$  is the infimum operation with respect to  $\leq_{sW}$  and  $\leq_{W}$ . That  $\sqcup$  is the supremum operation with respect to  $\leq_{W}$  was first proved by Pauly [19, Theorem 4.5] (see also [6, Theorem 3.9]). Dzhafarov proved that  $\boxplus$  is a supremum operation for  $\leq_{sW}$  [10] and he also showed  $f \boxplus g \equiv_{W} f \sqcup g$ . Using Propositions 2.2 and 2.5 we can transfer these results to the total versions of Weihrauch reducibility.

**Proposition 6.5** (Infima and suprema). Let f, g be problems. Then

- (1)  $\overline{f} \cap \overline{g}$  is an infimum of f and g with respect to  $\leq_{tW}$  and  $\leq_{stW}$ .
- (2)  $f \sqcup g$  is a supremum of f and g with respect to  $\leq_{tW}$ .
- (3)  $f \boxplus g$  is a supremum of f and g with respect to  $\leq_{\text{stW}}$ .
- (4)  $\overline{f \sqcup g} \equiv_{\mathbf{W}} \overline{f} \sqcup \overline{g}$  and hence  $f \sqcup g \equiv_{\mathbf{tW}} \overline{f} \sqcup \overline{g} \equiv_{\mathbf{tW}} f \boxplus g$ .
- (5)  $\overline{f \boxplus g} \equiv_{\text{sW}} \overline{f} \boxplus \overline{g} \text{ and hence } f \boxplus g \equiv_{\text{stW}} \overline{f} \boxplus \overline{g}.$

In Lemma 6.9 we will see that the equivalences in (4) cannot be strengthened to strong equivalences.

By a (strong) total Weihrauch degree we mean an equivalence class with respect to  $\leq_{\rm tW}$  (or with respect to  $\leq_{\rm stW}$  in the strong case). We denote the corresponding classes by  $\mathcal{W}_{\rm tW}$  and  $\mathcal{W}_{\rm stW}$ . Strictly speaking, these are not sets, but every equivalence class has a representative on Baire space according to Corollary 4.11, and if desired, we can turn the classes  $\mathcal{W}_{\rm tW}$  and  $\mathcal{W}_{\rm stW}$  into sets of such representatives. The same applies to further classes of degrees that we consider in the following. We can extend the reducibilities  $\leq_{\rm tW}$  and  $\leq_{\rm stW}$  to the corresponding degrees and

any monotone algebraic operation too. By Proposition 6.5 ( $W_{tW}$ ,  $\leq_{tW}$ ,  $\square$ ,  $\sqcup$ ) yields a lattice structure.

It was first proved by Pauly [19, Theorem 4.22] that the Weihrauch lattice is distributive. In fact, he proved that it is a distributive join semi-lattice, which implies distributivity as a lattice. That is, we have  $f \sqcup (g \sqcap h) \equiv_{\mathrm{W}} (f \sqcup g) \sqcap (f \sqcup h)$  and  $f \sqcap (g \sqcup h) \equiv_{\mathrm{W}} (f \sqcap g) \sqcup (f \sqcap h)$  [8, Theorem 31]. Also the total Weihrauch degrees form a distributive lattice.

**Theorem 6.6** (Total Weihrauch lattice).  $(\mathcal{W}_{tW}, \leq_{tW}, \overline{\sqcap}, \sqcup)$  is a distributive lattice.

*Proof.* By Proposition 6.5 we obtain

$$f\overline{\sqcap}(g\sqcup h)=\overline{f}\sqcap\overline{(g\sqcup h)}\equiv_{\mathcal{W}}\overline{f}\sqcap(\overline{g}\sqcup\overline{h})\equiv_{\mathcal{W}}(\overline{f}\sqcap\overline{g})\sqcup(\overline{f}\sqcap\overline{h})=(f\overline{\sqcap}g)\sqcup(f\overline{\sqcap}h)$$

and hence  $f \overline{\sqcap}(g \sqcup h) \equiv_{\text{tW}} (f \overline{\sqcap} g) \sqcup (f \overline{\sqcap} h)$  by Corollary 4.7. With Proposition 6.5 and Corollary 5.5 we obtain similarly as above

$$\begin{array}{ll} f \sqcup (g \overline{\sqcap} h) & \equiv_{\mathrm{tW}} & \overline{f} \sqcup (g \overline{\sqcap} h) = \overline{f} \sqcup (\overline{g} \sqcap \overline{h}) \equiv_{\mathrm{W}} (\overline{f} \sqcup \overline{g}) \sqcap (\overline{f} \sqcup \overline{h}) \\ & \equiv_{\mathrm{W}} & \overline{(f \sqcup g)} \sqcap \overline{(f \sqcup h)} = (f \sqcup g) \overline{\sqcap} (f \sqcup h) \end{array}$$

and hence  $f \sqcup (g \overline{\sqcap} h) \equiv_{\mathrm{tW}} (f \sqcup g) \overline{\sqcap} (f \sqcup h)$ . Altogether, this shows that the total Weihrauch lattice is distributive.

Proposition 6.5 implies that  $W_{stW}$  is a lattice. Dzhafarov proved that the lattice  $W_{sW}$  is not distributive [10, Theorem 4.4]. We can transfer his proof to  $W_{stW}$ .

**Theorem 6.7** (Strong total Weihrauch lattice).  $(W_{stW}, \leq_{stW}, \overline{\sqcap}, \boxplus)$  is a lattice, which is not distributive.

*Proof.* Proposition 6.5 implies that  $W_{\text{stW}}$  is a lattice. Suppose that this lattice is distributive. Then, in particular again by Proposition 6.5

$$(\overline{f} \boxplus \overline{g}) \sqcap \overline{h} \equiv_{\mathrm{sW}} \overline{(f \boxplus g)} \sqcap \overline{h} = (f \boxplus g) \overline{\sqcap} h \leq_{\mathrm{stW}} (f \overline{\sqcap} h) \boxplus (g \overline{\sqcap} h) = (\overline{f} \sqcap \overline{h}) \boxplus (\overline{g} \sqcap \overline{h}),$$
  
i.e.,  $(\overline{f} \boxplus \overline{g}) \sqcap \overline{h} \leq_{\mathrm{stW}} (\overline{f} \sqcap \overline{h}) \boxplus (\overline{g} \sqcap \overline{h}),$  which by Lemma 5.2, Propositions 6.5 and 6.3 is equivalent to

$$(\overline{f} \boxplus \overline{g}) \sqcap \overline{h} \leq_{\mathrm{sW}} \overline{(\overline{f} \sqcap \overline{h}) \boxplus (\overline{g} \sqcap \overline{h})} \equiv_{\mathrm{sW}} \overline{(\overline{f} \sqcap \overline{h})} \boxplus \overline{(\overline{g} \sqcap \overline{h})} \equiv_{\mathrm{sW}} (\overline{f} \sqcap \overline{h}) \boxplus (\overline{g} \sqcap \overline{h}).$$

Hence, it suffices to provide a counterexample for  $(\overline{f} \boxplus \overline{g}) \cap \overline{h} \leq_{\mathrm{sW}} (\overline{f} \cap \overline{h}) \boxplus (\overline{g} \cap \overline{h})$ . We use the proof idea of [10, Theorem 4.4] and we consider the constant problems  $c_{p,q}:\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}, p \mapsto q \text{ with } \operatorname{dom}(c_{p,q})=\{p\} \text{ for } p,q\in \mathbb{N}^{\mathbb{N}}. \text{ Let } p_i,q_i\in \mathbb{N}^{\mathbb{N}}$ for  $i \in \{0,1,2\}$  be mutually Turing incomparable and such that none of these points can be computed from the supremum of the others (this is possible, see for instance [22, Exercise 2.2 in Chapter VII]). We choose  $f := c_{p_0,q_0}, g := c_{p_1,q_1}$  and  $h:=c_{p_2,q_2}.$  We recall that  $\overline{\mathbb{N}^{\mathbb{N}}}=\mathbb{N}^{\mathbb{N}}\cup\{\bot\}$  is represented with a precomplete representation  $\delta$ , defined by  $\delta(p)=\operatorname{id}^{\wp}(p)=p-1$  for  $p-1\in\mathbb{N}^{\mathbb{N}}$  and  $\delta(p)=\bot$ otherwise. Now assume that  $(\overline{f} \boxplus \overline{g}) \sqcap \overline{h} \leq_{\mathrm{sW}} (\overline{f} \sqcap \overline{h}) \boxplus (\overline{g} \sqcap \overline{h})$  via computable H, K. We claim that  $K\langle\langle i, p_i+1\rangle, p_2+1\rangle = \langle i, \langle p_i', p_2'\rangle\rangle$  for  $i \in \{0, 1\}$  with names  $p_k'$  of  $p_k$ for  $k \in \{0, 1, 2\}$ . Firstly, if  $K(\langle i, p_i + 1 \rangle, p_2 + 1) = \langle j, \langle r, s \rangle \rangle$  such that r is not a name of  $p_j$  or s is not a name of  $p_2$ , then a realizer of  $e := (\overline{f} \cap \overline{h}) \boxplus (\overline{g} \cap \overline{h})$  on  $\langle j, \langle r, s \rangle \rangle$ could return any value, for instance a computable one, and in this case H could neither compute  $q_i$  nor  $q_2$  from this result. Hence  $K\langle\langle i, p_i+1\rangle, p_2+1\rangle = \langle j, \langle p'_j, p'_2\rangle\rangle$ with  $j \in \{0,1\}$  and  $p'_k$  a name for  $p_k$  for  $k \in \{0,1,2\}$ . Secondly, if  $j \neq i$ , then a realizer of e upon input of  $\langle j, \langle p_j', p_2' \rangle \rangle$  could return a name of  $q_j$  together with some computable values, from which H can neither compute  $q_i$  nor  $q_2$ . This proves the claim above. Now on input  $\langle 0, \langle p'_0, p'_2 \rangle \rangle$  as above, a realizer of e can produce  $r := \langle \langle 0, q'_0 \rangle + 1, 0 \rangle$  with a name  $q'_0$  of  $q_0$ . Suppose  $H(r) = \langle 0, s \rangle$  with some  $s \in \mathbb{N}^{\mathbb{N}}$ . Since H is continuous, a certain prefix of r is sufficient to produce the output 0

in the first component. Now on input  $\langle 1, \langle p'_1, p'_2 \rangle \rangle$  a realizer of e can produce the output  $t := \langle \langle 0, c \rangle + 1, 0^n (\langle 1, q_2' \rangle + 1) \rangle$  with a computable c that shares a sufficiently long prefix with  $q'_0$  and a sufficiently large  $n \in \mathbb{N}$  and a name  $q'_2$  of  $q_2$ . Then  $H(t) = \langle 0, s' \rangle$  with some  $s' \in \mathbb{N}^{\mathbb{N}}$ . However, s' is computable from  $q_2$  and hence it can neither compute  $q_0$  nor  $q_1$ , which is a contradiction. Hence  $H(r) = \langle 1, s \rangle$  with some  $s \in \mathbb{N}^{\mathbb{N}}$ . Again, due to continuity of H, some prefix of the input is sufficient to produce the component 1 on the output side. On input  $\langle 1, \langle p'_1, p'_2 \rangle \rangle$  a realizer of e can now produce the output  $t := \langle \langle 0, q'_0 \rangle + 1, 0^n (\langle 0, q'_1 \rangle + 1) \rangle$  for sufficiently large  $n \in \mathbb{N}$  and  $H(t) = \langle 1, s' \rangle$  with  $s' \in \mathbb{N}^{\mathbb{N}}$ . However, since s' is computable from  $q_0$ and  $q_1$ , it cannot compute  $q_2$ , which is a contradiction. 

We are going to prove that + also co-preserves completion with respect to  $\leq_{tW}$ and  $\leq_{\text{stW}}$ .

**Proposition 6.8** (Sums).  $\overline{f+g} \equiv_{sW} \overline{f} + \overline{g}$  and hence  $f+g \equiv_{stW} \overline{f} + \overline{g}$  for all problems f, g.

*Proof.* We consider problems  $f:\subseteq X \Rightarrow Y$  and  $g:\subseteq U \Rightarrow V$ . We obtain the problems  $\overline{f} + \overline{g} : \overline{X} \times \overline{U} \rightrightarrows \overline{\overline{Y}} \times \overline{\overline{V}}$  with

$$(\overline{f} + \overline{g})(x, u) = \begin{cases} (f(x) \times \overline{\overline{V}}) \cup (\overline{\overline{Y}} \times g(u)) & \text{if } (x, u) \in \text{dom}(f) \times \text{dom}(g) \\ (f(x) \times \overline{\overline{V}}) \cup (\overline{\overline{Y}} \times \overline{V}) & \text{if } x \in \text{dom}(f) \text{ and } u \not \in \text{dom}(g) \\ (\overline{Y} \times \overline{\overline{V}}) \cup (\overline{\overline{Y}} \times g(u)) & \text{if } x \not \in \text{dom}(f) \text{ and } u \in \text{dom}(g) \\ (\overline{Y} \times \overline{\overline{V}}) \cup (\overline{\overline{Y}} \times \overline{V}) & \text{otherwise} \end{cases}$$

and  $\overline{f+q}: \overline{X\times U} \rightrightarrows \overline{\overline{Y}\times \overline{V}}$  with

$$(\overline{f+g})(z) = \begin{cases} \frac{(f(x) \times \overline{V}) \cup (\overline{Y} \times g(u))}{\overline{Y} \times \overline{V}} & \text{if } z = (x,u) \in \text{dom}(f) \times \text{dom}(g) \\ & \text{otherwise} \end{cases}$$

And we also consider  $h: \overline{X} \times \overline{U} \rightrightarrows \overline{Y} \times \overline{V}$  with

$$h(x,u) := \left\{ \begin{array}{ll} (f(x) \times \overline{V}) \cup (\overline{Y} \times g(u)) & \text{if } (x,u) \in \text{dom}(f) \times \text{dom}(g) \\ \overline{Y} \times \overline{V} & \text{otherwise} \end{array} \right.$$

Then we have  $h(x,u)\subseteq (\overline{f}+\overline{g})(x,u)$  for all  $(x,u)\in \overline{X}\times \overline{U}$  and hence together with Proposition  $6.3 \frac{f}{f} + g \leq_{\text{sW}} \frac{f}{f} + \overline{g} \leq_{\text{sW}} h$ . On the other hand, there is a computable function  $s: \overline{Y} \times \overline{V} \to \overline{Y} \times \overline{V}$  with s(y,v) = (y,v) for all  $(y,v) \in \overline{Y} \times \overline{V}$ . Namely, one can just consider  $S: \subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}, p \mapsto p-1$  and extend this to a total computable realizer of s under the representation of  $\overline{Y} \times \overline{V}$ , which is possible, since this space has a precomplete representation by Proposition 3.8. Analogously to s, there is also a computable function  $\iota: \overline{X} \times \overline{U} \to \overline{X \times U}$  with  $\iota(x,u) = (x,u)$  for  $(x,u) \in X \times U$ . Then  $h = s \circ \overline{(f+g)} \circ \iota$  and hence  $h \leq_{sW} \overline{f+g}$ .

The following example shows that  $\times$  and  $\sqcap$  do not co-preserve completion with respect to  $\leq_{\rm W}$  and that  $\sqcup$  does not co-preserve completion with respect to  $\leq_{\rm sW}$ .

**Lemma 6.9.** There are problems  $f, g :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  such that

- (1)  $\overline{f} \times \overline{g} \not\leq_{W} \overline{f \times g}$ , and hence  $\overline{f} \times \overline{g} \not\leq_{tW} f \times g$ , (2)  $\overline{f} \sqcap \overline{g} \not\leq_{W} \overline{f \sqcap g}$ , and hence  $\overline{f} \sqcap \overline{g} \not\leq_{tW} f \sqcap g$ , (3)  $\overline{f} \sqcup \overline{g} \not\leq_{sW} \overline{f \sqcup g}$ , and hence  $\overline{f} \sqcup \overline{g} \not\leq_{stW} f \sqcup g$ .

*Proof.* We consider the constant problems  $c_{p,q}:\subseteq \mathbb{N}^{\mathbb{N}}\to\mathbb{N}^{\mathbb{N}}, p\mapsto q$  with  $\mathrm{dom}(c_{p,q})=$  $\{p\}$  for  $p,q\in\mathbb{N}$ . Let  $p,q,r,s\in\mathbb{N}^{\mathbb{N}}$  be mutually Turing incomparable. We choose  $f:=c_{p,q}$  and  $g:=c_{r,s}$ . We recall that  $\overline{\mathbb{N}^{\mathbb{N}}}=\mathbb{N}^{\mathbb{N}}\cup\{\bot\}$  is represented with a precomplete representation  $\delta$ , defined by  $\delta(p) = \mathrm{id}^{\wp}(p) = p-1$  for  $p-1 \in \mathbb{N}^{\mathbb{N}}$  and  $\delta(p) = \bot$  otherwise. We only need to prove the former statements regarding  $<_{\rm W}$ , since the latter statements regarding  $\leq_{tW}$  follow in each case with Lemma 5.2.

- (1) holds since a name for the input pair  $(p,p) \in \text{dom}(\overline{c_{p,q}} \times \overline{c_{r,s}})$  can only be mapped computably to a name of an input outside of  $\text{dom}(c_{p,q} \times c_{r,s}) = \{(p,r)\}$  since r is not computable from p, and a realizer for  $\overline{c_{p,q} \times c_{r,s}}$  can map such a name to any name, for instance a computable name. From a computable name and a name for (p,p) one cannot compute q.
- (2) Let us assume that  $\overline{c_{p,q}} \sqcap \overline{c_{r,s}} \leq_{\mathrm{W}} \overline{c_{p,q}} \sqcap c_{r,s}$  is witnessed by computable H, K. We consider the name p+1 of p and the name  $\widehat{0}$  of  $\bot$ . Since  $(p,\bot) \in \mathrm{dom}(\overline{c_{p,q}} \sqcap \overline{c_{r,s}})$ ,  $K\langle p+1,\widehat{0}\rangle$  has to be defined, but it cannot be a name of a point in  $\mathrm{dom}(c_{p,q} \sqcap c_{r,s}) = \{p\} \times \{r\}$ . Let G be a realizer of  $\overline{c_{p,q}} \sqcap \overline{c_{r,s}}$  that maps every name of a point outside of  $\mathrm{dom}(c_{p,q} \sqcap c_{r,s})$  to  $\widehat{0}$ . Then  $H\langle \langle p+1,\widehat{0}\rangle, GK\langle p+1,\widehat{0}\rangle\rangle = H\langle \langle p+1,\widehat{0}\rangle, \widehat{0}\rangle = \langle 1,t\rangle$  for some  $t\in\mathbb{N}^{\mathbb{N}}$ , since it cannot be equal to  $\langle 0,u\rangle$  for some  $u\in\mathbb{N}^{\mathbb{N}}$  because q cannot be computed from p and  $H\langle \mathrm{id}, GK\rangle$  has to be a realizer of  $\overline{c_{p,q}} \sqcap \overline{c_{r,s}}$ . Due to continuity of H the output 1 in the first component is determined already by a prefix of the input, say by  $w\sqsubseteq p+1$  and  $0^n\sqsubseteq \widehat{0}$ . Hence, on the names  $w\widehat{0}$  and  $0^n(r+1)$  of  $\bot$  and r, respectively, the function H will also produce 1 in the first component. Moreover  $K\langle w\widehat{0}, 0^n(r+1)\rangle$  is also a name of a point outside of  $\mathrm{dom}(c_{p,q} \sqcap c_{r,s}) = \{p\} \times \{r\}$  and hence  $GK\langle w\widehat{0}, 0^n(r+1)\rangle = \widehat{0}$ . In this case we must have  $H\langle \langle w\widehat{0}, 0^n(r+1)\rangle$ ,  $GK\langle w\widehat{0}, 0^n(r+1)\rangle = H\langle \langle w\widehat{0}, 0^n(r+1)\rangle$ ,  $\widehat{0}\rangle = \langle 1,t\rangle$  with a name t of s, which is impossible, since s cannot be computed from r.
- (3) Let us assume that  $\overline{c_{p,q}} \sqcup \overline{c_{r,s}} \leq_{\mathrm{sW}} \overline{c_{p,q}} \sqcup \overline{c_{r,s}}$  is witnessed by computable H, K. Upon input of the name  $\langle i, \widehat{0} \rangle$  of  $(i, \bot) \in \mathrm{dom}(\overline{c_{p,q}} \sqcup \overline{c_{r,s}})$  with  $i \in \{0, 1\}$  the function K cannot produce a name of a point in  $\mathrm{dom}(c_{p,q} \sqcup c_{r,s}) = \{(0,p),(1,r)\}$ . There is a realizer G of  $\overline{f \sqcup g}$  that produces the name  $\widehat{0}$  of  $\bot$  on any input outside of the domain of  $\mathrm{dom}(c_{p,q} \sqcup c_{r,s})$  and hence  $HGK\langle i, \widehat{0} \rangle = \langle j, t \rangle$  for some fixed  $j \in \{0, 1\}$  and  $t \in \mathbb{N}^{\mathbb{N}}$  and both values  $i \in \{0, 1\}$ . The fixed j can only be correct for one of the values i, since we need i = j for the correctness of H, K, which is impossible.  $\square$

With the help of Corollary 5.5 it follows that  $\times$  and  $\sqcap$  are not monotone with respect to the total versions of Weihrauch reducibility.

**Corollary 6.10.**  $\times, \sqcap$  are neither monotone with respect to  $\leq_{\mathrm{tW}}$  nor with respect to  $\leq_{\mathrm{stW}}$ , and  $\sqcup$  is not monotone with respect to  $\leq_{\mathrm{stW}}$ .

Many further algebraic properties of the Weihrauch lattice have been studied in [8]. Some of these results can be transferred to the total case by Corollary 4.7. In some cases we can also transfer results for pointed problems, since the completion  $\overline{f}$  of any problem is always pointed. For instance, the completions of the algebraic operations are ordered in the following way, as the corresponding reductions hold more generally for pointed problems (by [6, Proposition 5.7], and that  $f^* \leq_{\mathrm{W}} \widehat{f}$  holds for pointed f, is easy to see).

**Corollary 6.11** (Order of operations). For all problems f and g we obtain:  $\overline{f} + \overline{g} \leq_{sW} \overline{f} \sqcap \overline{g} \leq_{sW} \overline{f} \boxplus \overline{g} \leq_{sW} \overline{f} \sqcup \overline{g} \leq_{sW} \overline{f} \times \overline{g}$ ,  $\overline{f} \boxplus \overline{g} \leq_{sW} \overline{f} \times \overline{g}$  and  $\overline{f}^* \leq_{w} \overline{f}$ .

Now we study the completions of parallelization  $f \mapsto \widehat{\overline{f}}$  and finite parallelization  $f \mapsto \overline{f}^*$ . In [3, Proposition 4.2] we proved that  $f \mapsto \widehat{f}$  is a closure operator for  $\leq_{\mathrm{W}}$  and  $\leq_{\mathrm{sW}}$  and Pauly proved in [19, Theorem 6.2] that  $f \mapsto f^*$  is a closure operator for (the topological version of)  $\leq_{\mathrm{W}}$ . We note that the latter one is not a closure operator for  $\leq_{\mathrm{sW}}$ . Nevertheless, the completions of both operators are closure operators for  $\leq_{\mathrm{tW}}$  and  $\leq_{\mathrm{stW}}$ . In order to prove this, we need the following additional lemma.

**Lemma 6.12** (Arno Pauly<sup>8</sup>).  $f^{**} \equiv_{sW} f^*$  for all pointed problems f.

*Proof.* It is easy to see that  $f \leq_{\text{sW}} f^*$  holds for all problems f, in particular, we obtain  $f^* \leq_{\text{sW}} f^{**}$ . For the inverse reduction we assume that f is pointed. Let  $p_0$ be a computable name of a point in dom(f). We use the computable functions K, H with

$$K\langle n, \langle \langle i_1, \langle p_{1,1}, ..., p_{1,i_1} \rangle \rangle, \langle i_2, \langle p_{2,1}, ..., p_{2,i_2} \rangle \rangle, ..., \langle i_n, \langle p_{n,1}, ..., p_{n,i_n} \rangle \rangle \rangle \rangle$$

$$:= \left\langle k, \langle p_{1,1}, ..., p_{1,i_1}, p_{2,1}, ..., p_{2,i_2}, ...., p_{n,1}, ..., p_{n,i_n}, \underbrace{p_0, ..., p_0}_{\text{en times}} \right\rangle$$

where  $k := \langle n, \langle i_1, ..., i_n \rangle \rangle \geq i_1 + ... + i_n$  and  $m := k - (i_1 + ... + i_n)$ , and for arbitrary  $k = \langle n, \langle i_1, ..., i_n \rangle \rangle \in \mathbb{N}$  and  $j := i_1 + ... + i_n \leq k$  we define

$$H\langle k, \langle q_1, ..., q_k \rangle \rangle$$

$$:= \langle n, \langle \langle i_1, \langle q_1, ..., q_{i_1} \rangle \rangle, \langle i_2, \langle q_{i_1+1}, ..., q_{i_1+i_2} \rangle \rangle, ..., \langle i_n, \langle q_{i_1+...+i_{n-1}+1}, ..., q_i \rangle \rangle \rangle \rangle.$$

Then H, K are computable and witness  $f^{**} \leq_{sW} f^*$ .

Now we are prepared to prove the following result.

**Proposition 6.13** (Parallelization).  $f \mapsto \widehat{\overline{f}}$  and  $f \mapsto \overline{f}^*$  are closure operators for  $\leq_{\mathrm{tW}}$  and  $\leq_{\mathrm{stW}}$  (and also for  $\leq_{\mathrm{W}}$  and  $\leq_{\mathrm{sW}}$ ).

*Proof.* Since parallelization  $f \mapsto \widehat{f}$  and completion  $f \mapsto \overline{f}$  are both closure operators for  $\leq_{sW}$  and  $\leq_{W}$  by [3, Proposition 4.2] and Proposition 5.4, and parallelization preserves completion by Proposition 6.3, the claim follows from Propositions 2.2 and 2.4. The claim for  $f \mapsto \overline{f}^*$  with respect to  $\leq_{tW}$  follows analogously. In order to prove the claim for  $\leq_{\text{stW}}$ , we note that  $f \mapsto f^*$  is a closure operator with respect to  $\leq_{\rm sW}$  restricted to pointed problems. This follows from Corollary 6.2, Lemma 6.12 and since  $f \leq_{\text{sW}} f^*$  obviously holds true. Hence, we also obtain that  $f \mapsto \overline{f}^*$ is a closure operator with respect to  $\leq_{\text{stW}}$ , since all problems of the form  $\overline{f}$  are pointed. 

With the following counterexamples we show that (finite) parallelization does not co-preserve completion. Some of the statements can be seen as a strengthening of the first statement in Lemma 6.9.

**Lemma 6.14.** There is a problem f with  $\overline{f} \times \overline{f} \leq_{\mathbf{W}} \overline{\widehat{f}}$  and  $\overline{f} \times \overline{f} \leq_{\mathbf{W}} \overline{f^*}$ . This implies

- (1)  $\overline{f} \times \overline{f} \not\leq_{\mathbf{W}} \overline{f \times f}$ , and hence  $\overline{f} \times \overline{f} \not\leq_{\mathbf{tW}} f \times f$ , (2)  $\overline{f}^* \not\leq_{\mathbf{W}} \overline{f^*}$ , and hence  $\overline{f}^* \not\leq_{\mathbf{tW}} f^*$ ,
- (3)  $\overline{f} \not\leq_{\mathbf{W}} \overline{\widehat{f}}$ , and hence  $\overline{\widehat{f}} \not\leq_{\mathbf{tW}} \widehat{f}$ .

*Proof.* We consider the function  $f:\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  with f(p)=q, f(r)=s, dom(f)=s $\{p,r\}$  and pairwise Turing incomparable  $p,q,r,s\in\mathbb{N}^{\mathbb{N}}$  such that none of these is computable from the supremum of the others (this is possible, see for instance [22, Exercise 2.2 in Chapter VII]). We recall that  $\overline{\mathbb{N}^{\mathbb{N}}} = \mathbb{N}^{\mathbb{N}} \cup \{\bot\}$  is represented with a precomplete representation  $\delta$ , defined by  $\delta(p) = \mathrm{id}^{\wp}(p) = p-1$  for  $p-1 \in \mathbb{N}^{\mathbb{N}}$  and  $\delta(p) = \bot$  otherwise. Let us assume that the reduction  $\overline{f} \times \overline{f} \leq_{\mathbf{W}} \widehat{f}$  holds, witnessed by computable H, K. The names p + 1, r + 1 of p, q are mapped by K to a name  $K\langle p+1,r+1\rangle$  of a point  $(q_n)_{n\in\mathbb{N}}$  in  $\mathrm{dom}(\widehat{f})$ , since a realizer of  $\overline{\widehat{f}}$  can choose a computable output outside of  $dom(\hat{f})$  and the result (q,s) cannot be computed from p, r alone. For the same reason  $q_n = r$  for at least one  $n \in \mathbb{N}$  and hence  $\pi_n K(p+1,r+1)$  is a name  $r_0$  of r. Due to continuity of K there are prefixes

<sup>&</sup>lt;sup>8</sup>By personal communication 2018

 $w \sqsubseteq p+1$  and  $v \sqsubseteq r+1$  that are sufficient for K to produce a prefix  $u \sqsubseteq r_0$  that is long enough so that it cannot be extended to a name of p. We can now replace r+1 by  $t=v\widehat{0}$ , which is a name of  $\bot \in \mathrm{dom}(\overline{f})$ . Now  $\pi_n K\langle p+1,t\rangle$  cannot be a name of r, since r cannot be computed from p and t and it cannot be a name of p either, since  $u \sqsubseteq \pi_n K\langle p+1,t\rangle$ . Hence  $K\langle p+1,t\rangle$  is a name for a point outside of  $\mathrm{dom}(\widehat{f})$  and a realizer of  $\overline{f}$  can choose a computable result c on this name. But  $H\langle\langle p+1,t\rangle,c\rangle$  cannot compute q, which is required by the assumption. This proves  $\overline{f}\times\overline{f}\not\leq_W \widehat{f}$ . The second statement can be proved analogously, one has to choose w,v such that also the natural number component of the name of an output in  $(\mathbb{N}^{\mathbb{N}})^*$  is fixed.

All other statements that involve  $\leq_{\mathrm{W}}$  are consequences since  $\overline{f} \times \overline{f} \leq_{\mathrm{W}} \overline{f}^* \leq_{\mathrm{W}} \overline{f}$ ,  $\overline{f} \times \overline{f} \leq_{\mathrm{W}} \overline{f}^*$  and  $\overline{f} \times \overline{f} \leq_{\mathrm{W}} \overline{f}^*$ . These reductions follow since obviously  $g \times g \leq_{\mathrm{W}} g^*$  and  $g \times g \leq_{\mathrm{W}} \widehat{g}$  for any problem g, completion is a closure operator by Proposition 5.4, and by Corollary 6.11, since  $\overline{f}$  is pointed. The statements that involve  $\leq_{\mathrm{tW}}$  follow from Lemma 5.2.

As an immediate consequence of these counterexamples we can conclude that parallelization and finite parallelization are not monotone operations for the total variants of Weihrauch reducibility. Since  $\overline{f} \leq_{\text{stW}} f$  holds by Corollary 5.5, we obtain the following conclusion using Lemma 6.14.

Corollary 6.15.  $f \mapsto \widehat{f}$  and  $f \mapsto f^*$  are neither monotone with respect to  $\leq_{\text{tW}}$  nor with respect to  $\leq_{\text{stW}}$ .

Another consequence of Lemma 6.14 is that completion does neither preserve idempotency nor parallelizability. We recall that a problem f is called *idempotent*, if  $f \equiv_{\mathbf{W}} f \times f$  and it is called *parallelizable*, if  $\widehat{f} \equiv_{\mathbf{W}} f$ . If we consider the problem f from Lemma 6.14, then we can take  $f^*$  and  $\widehat{f}$  as examples to obtain the following result.

Corollary 6.16 (Idempotency and parallelizability).

- (1) There is an idempotent problem f such that  $\overline{f}$  is not idempotent.
- (2) There is a parallelizable problem f such that  $\overline{f}$  is not parallelizable.

In the next step we want to clarify the relation between  $\leq_{\text{tW}}$  and  $\leq_{\text{stW}}$  and for this purpose we need to study cylinders. We recall that a problem f is called cylinder if id  $\times f \leq_{\text{sW}} f$  holds, and id  $\times f$  is called the cylindrification of f [3]. It follows from [4, Proposition 4.16] that "total cylinders" are exactly the usual cylinders.

Corollary 6.17 (Total cylinders).  $id \times f \leq_{sW} f \iff id \times f \leq_{stW} f \text{ holds for all problems } f$ .

It is known that g is a cylinder if and only if  $f \leq_{W} g \iff f \leq_{sW} g$  holds for all problems f [3, Proposition 3.5, Corollary 3.6]. We provide a similar result for the total variant of Weihrauch reducibility.

**Proposition 6.18** (Cylinder). A problem g is a cylinder if and only if for every problem f one has  $f \leq_{\text{tW}} g \iff f \leq_{\text{stW}} g$ .

*Proof.* Let us assume that  $f \leq_{\text{tW}} g \iff f \leq_{\text{stW}} g$  holds for every problem f. It is clear that  $\text{id} \times g \equiv_{\text{W}} g$  and hence  $\text{id} \times g \equiv_{\text{tW}} g$  by Corollary 4.7. By the assumption this implies  $\text{id} \times g \leq_{\text{stW}} g$  and hence  $\text{id} \times g \leq_{\text{sW}} g$  by Corollary 6.17. This shows that g is a cylinder.

For the other direction, let us now assume that g is a cylinder, i.e., id  $\times g \leq_{\text{sW}} g$  and hence id  $\times g \leq_{\text{stW}} g$  by Corollary 4.7. We only need to prove that  $f \leq_{\text{tW}} g$ 

implies  $f \leq_{\text{stW}} g$ . Let us assume that  $f \leq_{\text{tW}} g$  holds. Since  $f \leq_{\text{sW}} \text{id} \times f$ , we obtain  $f \leq_{\text{stW}} \text{id} \times f$  by Lemma 5.2. Now it suffices to show id  $\times f \leq_{\text{stW}} \text{id} \times g$ . But this can be done by using the construction of the proof of [3, Proposition 3.5]. By Lemma 4.5 it suffices to note that if H, K from the proof of [3, Proposition 3.5] are total, then also the H', K' constructed in the first half of that proof are total.  $\square$ 

Hence, the relations between strong and weak versions of the reducibility can be expressed in the same way in the partial and the total case, respectively.

We can also say something on the interaction between cylindrification and completion. While the completion of a cylinder f is only a cylinder in the trivial case that the original problem f is already strongly complete, the cylindrification of a complete problem is always complete.

**Proposition 6.19** (Completion and cylindrification). Let f be a problem. Then

- (1)  $\overline{f}$  is a cylinder  $\iff$  f is strongly complete and a cylinder,
- (2)  $id \times f$  is complete  $\iff$  f is complete.

The implication "\( "=" in (2) also holds for strongly complete instead of complete."

*Proof.* (1) If  $f \equiv_{sW} \overline{f}$  and f is a cylinder, then clearly  $id \times \overline{f} \leq_{sW} id \times f \leq_{sW} \overline{f}$ and hence  $\overline{f}$  is a cylinder. If, on the other hand,  $\overline{f}$  is a cylinder, then id  $\times \overline{f} \leq_{sW} \overline{f}$ . Hence id  $\times \overline{f} \leq_{\text{stW}} f$  and since id  $\times \overline{f}$  is diverse, we obtain by [4, Proposition 4.16] that id  $\times \overline{f} \leq_{\underline{sW}} f$ . This implies id  $\times f \leq_{\underline{sW}} id \times \overline{f} \leq_{\underline{sW}} f$ , which means that f is a cylinder and  $\overline{f} \leq_{\text{sW}} \text{id} \times \overline{f} \leq_{\text{sW}} f$ , which means that f is strongly complete. (2) If f is complete, then  $\overline{\mathrm{id} \times f} \leq_{\mathrm{W}} \overline{\mathrm{id}} \times \overline{f} \leq_{\mathrm{W}} \mathrm{id} \times f$  by Propositions 6.3 and 5.8,

which means that id  $\times f$  is complete. The proof in the strong case is analogous. If on the other hand, id  $\times f$  is complete, then  $\overline{f} \leq_{\mathbf{W}} \overline{\mathrm{id} \times f} \leq_{\mathbf{W}} \mathrm{id} \times f \leq_{\mathbf{W}} f$ , where the first reduction holds since  $f \leq_{\mathbf{W}} \mathrm{id} \times f$  and completion is a closure operator.

## 7. Co-Residual Operations

In this section we will discuss certain algebraic operations that are co-residual operations. In this context we have to deal with a top element in the Weihrauch lattice. The Weihrauch lattice has no natural top element, but we can just attach a top element  $\infty$  to it. The algebraic operations are then naturally extended to the top element, so that the lattice structure and the order among the operations is preserved. We are led to the following choice of values for all problems f including  $\infty$  (see also the discussion in [8]):

- (1)  $f \sqcap \infty = \infty \sqcap f = f$ ,
- $(2) \ f \sqcup \infty = \infty \sqcup f = \infty,$
- (3)  $f \times \infty = \infty \times f = \infty$ ,
- (4)  $f + \infty = \infty + f = f$ , (5)  $\overline{\infty} = \widehat{\infty} = \infty^* = \infty$ .

One arguable alternative could be to choose  $0 \times \infty = 0$ , given that  $0 \times f \equiv_W 0$  for all  $f \neq \infty$ . However this seems to be less natural for our purposes. It is consistent with our usage of the term to say that a problem f is pointed, if  $1 \leq_W f$  holds. According to this definition  $\infty$  is pointed too.

Using our universal function  $U:\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ , we can define a representation  $\Phi$ of certain continuous functions by  $\Phi_q(p) := U\langle q, p \rangle$  for all  $p, q \in \mathbb{N}^{\mathbb{N}}$ . Then any continuous  $F:\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  has an extension of the form  $\Phi_a:\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  and for a computable F we can choose a computable q (see [27]). From this representation we can derive a Gödel numbering  $\varphi$  of the computable  $F:\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ , i.e., for every computable F there is some  $n \in \mathbb{N}$  such that  $\varphi_n :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  extends F. We also assume that  $\varphi$  satisfies suitable utm- and smn-Theorems (see [27] for details). We use  $\Phi$  and  $\varphi$  to define the compositional product and two implications.

The compositional product  $f \star g$  was originally defined in [5] using the property (1) stated in Fact 7.2 below. It expresses a problem that can be obtained by first applying g and then f with some possible intermediate computation. A corresponding compositional implication operation  $g \to f$  was introduced and studied in [8]. It characterizes the minimal problem h such that  $f \leq_{\mathbf{W}} g \star h$  (see Fact 7.2). Here we phrase these operations in a type free version on Baire space (as in [6]). We also introduce a multiplicative implication  $g \to f$ , which is supposed to capture a problem simpler than every h such that  $f \leq_{\mathbf{W}} g \times h$  (see Proposition 7.9).

**Definition 7.1** (Compositional product and implications). Let f, g be problems. We define problems  $f \star g$ ,  $(g \to f)$ ,  $(g \twoheadrightarrow f) :\subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$  by

```
(1) \ (f \star g) \langle q, p \rangle := \langle \operatorname{id} \times f^{r} \rangle \circ \Phi_{q} \circ g^{r}(p),
```

- $(2) (g \to f)(p) := \{ \langle t, q \rangle : \emptyset \neq \Phi_t \circ g^{\mathbf{r}}(q) \subseteq f^{\mathbf{r}}(p) \},$
- $(3) (g \to f)(p) := \{ \langle n, k, q \rangle : \emptyset \neq \varphi_n \langle q, g^{\mathsf{r}} \circ \varphi_k(p) \rangle \subseteq f^{\mathsf{r}}(p) \},$

where we assume for (2) and (3) that  $dom(g) \neq \emptyset$  or  $dom(f) = \emptyset$ . In the case of special constants we define:

```
(1) f \star \infty := \infty \star f := \infty,
```

- (2)  $(g \to 0) := (g \to 0) := 0, (0 \to f) := (0 \to f) := \infty \text{ for } f \not\equiv_{\mathbf{W}} 0,$
- (3)  $(\infty \to f) := (\infty \twoheadrightarrow f) := 0, (g \to \infty) := (g \twoheadrightarrow \infty) := \infty \text{ for } g \neq \infty.$

We call  $f \star g$  the compositional product,  $(g \to f)$  the compositional implication and  $(g \to f)$  the multiplicative implication.

The definition of  $(g \to 0) := (g \to 0) := 0$  is consistent with what is defined in the first two items (2) and (3) above. The domains in the first items (1)–(3) are always meant to be maximal. For instance  $\operatorname{dom}(g \to f) = \operatorname{dom}(f^r)$  if g is somewhere defined. The fact that we use Gödel numbers  $n, k \in \mathbb{N}$  for  $(g \to f)$  actually has some reason: the crucial properties of this implication are computability theoretic ones (see Proposition 7.11) and do not relativize to a topological version in an obvious way. However, the fact that we use Gödel numbers makes the domain of  $(g \to f)$  relatively complicated. If g is somewhere defined, then

$$dom(g \twoheadrightarrow f) = \{ p \in dom(f^{r}) : (\exists q \in dom(g^{r})) \ q \leq_{T} p \}.$$

For pointed g (that have a computable point in the domain) the domain is more natural and we obtain  $dom(g \rightarrow f) = dom(f^r)$ . The following facts were proved in [8, Corollaries 18 and 25, Theorem 24, Proposition 31].

**Fact 7.2** (Compositional product and implication). For all problems f and g including  $\infty$ :

```
(1) f \star g \equiv_{\mathbf{W}} \max_{\leq_{\mathbf{W}}} \{ f_0 \circ g_0 : f_0 \leq_{\mathbf{W}} f, g_0 \leq_{\mathbf{W}} g \},
```

- (2)  $(g \to f) \equiv_{\mathbf{W}} \min_{\leq_{\mathbf{W}}} \{ h : f \leq_{\mathbf{W}} g \star h \},$
- (3)  $(g \to f) \leq_{\mathbf{W}} h \iff f \leq_{\mathbf{W}} g \star h$ ,
- (4)  $\star$  is monotone with respect to  $\leq_{W}$  in both components,
- (5)  $\rightarrow$  is monotone with respect to  $\leq_W$  in the second component and antitone in the first component.

We note that for (3) to be correct in the case of  $dom(g) = \emptyset$  and  $dom(f) \neq \emptyset$ , we actually use  $(g \to f) = \infty$  and  $f \star \infty = \infty \star f = \infty$ .

By W we denote the class of Weihrauch degrees including  $\infty$ . We extend all the algebraic operations to degrees in the usual way without introducing a new notation. It is known that the underlying structure is a lattice [8] and together with Fact 7.2 (3) we obtain the following conclusion.

**Corollary 7.3** (Weihrauch algebra).  $(W, \leq_W, \sqcap, \sqcup, \star, \to, 0, 1, \infty)$  is a deductive Weihrauch algebra that is not commutative.

For instance  $\lim \star WKL \equiv_W \lim <_W WKL \star \lim$  and hence  $\star$  is clearly not commutative.

We can interpret  $(f \to \infty) = (f \to \infty)$  as negation operation in the Weihrauch lattice and we formally define negation correspondingly.

**Definition 7.4** (Negation). For every problem f we define its negation  $\neg f$  by  $\neg f := \infty \text{ for } f \neq \infty \text{ and } \neg \infty := 0 \text{ (the nowhere defined problem } 0 : \subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}).$ 

It is then obvious that our negation behaves as in Jankov logic.

Corollary 7.5 (Jankov rule).  $\neg \neg f \sqcap \neg f \equiv_{\mathbf{W}} 0$  is computable.

We note that  $\neg \overline{f} \equiv_{W} \neg f \leq_{W} \overline{\neg f}$ , but equivalence does not hold as we obtain  $\neg \overline{\infty} = 0 <_{W} 1 \equiv_{W} \overline{\neg \infty}$ . Here we are in particular interested in how the compositional product and the implications interact with completion in general. We show that  $\star$  co-preserves completion with respect to  $\leq_{\mathrm{sW}}$  and  $\to$  preserves completion with respect to  $\leq_{W}$ .

**Proposition 7.6** (Completion and compositional products and implication). For all problems f, g including  $\infty$ :

$$(1) \ \overline{f \star g} \leq_{\mathrm{sW}} \overline{f} \star \overline{g} \equiv_{\mathrm{sW}} \overline{\overline{f} \star \overline{g}}.$$

(1) 
$$\overline{f \star g} \leq_{\text{sW}} \overline{f} \star \overline{g} \equiv_{\text{sW}} \overline{\overline{f} \star \overline{g}}$$
.  
(2)  $(\overline{g} \to \overline{f}) \leq_{\text{W}} (\overline{g} \to \overline{f}) \leq_{\text{W}} (g \to f)$ .

In particular  $f \star g$  is (strongly) complete, if f and g are so.

*Proof.* (1) It is routine to check the claim for the special cases where the problem  $\infty$ is involved. Otherwise, it suffices to consider  $f,g:\subseteq\mathbb{N}^{\mathbb{N}}\rightrightarrows\mathbb{N}^{\mathbb{N}}$  and for such problems we have  $f \star g = \langle \operatorname{id} \times f \rangle \circ U \circ \langle \operatorname{id} \times g \rangle$ . Hence,  $\overline{f \star g} = \overline{\langle \operatorname{id} \times f \rangle} \circ \overline{U} \circ \overline{\langle \operatorname{id} \times g \rangle}$ . Since  $\overline{U}$  is computable and id is complete, this implies by Proposition 6.3

$$\overline{f \star g} \leq_{\mathbf{W}} \overline{(\mathrm{id} \times f)} \star \overline{(\mathrm{id} \times g)} \leq_{\mathbf{W}} (\mathrm{id} \times \overline{f}) \star (\mathrm{id} \times \overline{g}) \leq_{\mathbf{W}} \overline{f} \star \overline{g}.$$

Since every compositional product is a cylinder by [8, Lemma 17], we even obtain the strong Weihrauch reduction. The equivalence follows as in Proposition 6.3.

(2) Since  $f \leq_{\mathbf{W}} g \star (g \to f)$  by Fact 7.2 and completion is a closure operator, we obtain with (1)

$$\overline{f} \leq_{\mathcal{W}} \overline{g \star (g \to f)} \leq_{\mathcal{W}} \overline{g} \star \overline{(g \to f)}.$$

Hence Fact 7.2 implies  $(\overline{g} \to \overline{f}) \leq_{\mathbf{W}} \overline{(g \to f)}$ , which in turn implies the statement, as completion is a closure operator.

We note that neither of the reductions in (2) are equivalences in general, as the following examples show:

$$(1) \ \ \underline{(\overline{\infty} \to \overline{\infty})} \equiv_W 0 <_W 1 \equiv_W \overline{(\overline{\infty} \to \overline{\infty})},$$

(2) 
$$\overline{(\overline{0} \to \overline{1})} \equiv_{\mathrm{W}} 1 <_{\mathrm{W}} \infty \equiv_{\mathrm{W}} \overline{(0 \to 1)}.$$

In particular  $(\overline{g} \to \overline{f})$  does not need to be complete, even though  $\overline{g}$  and  $\overline{f}$  are.

We now want to study the multiplicative implication  $(g \rightarrow f)$  somewhat further. We first study its monotonicity properties.

**Proposition 7.7** (Monotonicity of multiplicative implication). Let  $f_i, g_i$  be problems for  $i \in \{0,1\}$  including  $\infty$ . If  $f_0 \leq_W f_1$ ,  $g_0 \leq_W g_1$  and  $g_0$  is pointed, then  $(g_1 \twoheadrightarrow f_0) \leq_{\mathbf{W}} (g_0 \twoheadrightarrow f_1).$ 

*Proof.* It is routine to check that the claim holds in those cases where the implication takes the values 0 or  $\infty$ . This includes the cases where  $\infty$  is among  $f_i, g_i$ . We break the proof for the other cases into two manageable pieces, where we either fix  $f = f_0 = f_1$  or  $g = g_0 = g_1$ . It suffices to consider problems  $g_i, f_i, g, f :\subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ 

(1) Let  $g_0 \leq_W g_1$  hold via computable H, K. We prove  $(g_1 \twoheadrightarrow f) \leq_W (g_0 \twoheadrightarrow f)$ .

Let us assume that  $g_0$  is pointed. This implies that  $g_1$  is also pointed and we also obtain  $dom(g_0 \rightarrow f) = dom(g_1 \rightarrow f) = dom(f)$ . By the smn-Theorem there are computable functions  $r, h : \mathbb{N} \to \mathbb{N}$  such that

- $\varphi_{h\langle n,k\rangle}\langle\langle p,q\rangle,t\rangle = \varphi_n\langle q,H\langle \pi_1\varphi_k(p),t\rangle\rangle,$
- $\varphi_{r(k)}(p) = K \circ \pi_2 \varphi_k(p)$

for all  $n, k \in \mathbb{N}$  and  $p, q, t \in \mathbb{N}^{\mathbb{N}}$ . Let  $p \in \text{dom}(g_1 \twoheadrightarrow f) = \text{dom}(g_0 \twoheadrightarrow f) = \text{dom}(f)$ . Let  $\langle n, k, q \rangle \in (g_0 \rightarrow f)(p)$ . Then we obtain

```
\varphi_{h\langle n,k\rangle}\langle\langle p,q\rangle, g_1\circ\varphi_{r(k)}(p)\rangle = \varphi_n\langle q,H\langle\pi_1\varphi_k(p),g_1\circ\varphi_{r(k)}(p)\rangle\rangle
                                                                            = \varphi_n \langle q, H \langle \pi_1 \varphi_k(p), g_1 \circ K \circ \pi_2 \varphi_k(p) \rangle \rangle
                                                                             = \varphi_n \langle q, H \langle id, q_1 \circ K \rangle \circ \varphi_k(p) \rangle
                                                                             \subseteq \varphi_n\langle q, g_0 \circ \varphi_k(p) \rangle \subseteq f(p).
```

This means  $\langle h\langle n,k\rangle, r(k), \langle p,q\rangle\rangle \in (g_1 \twoheadrightarrow f)(p)$ . Since the function H' with  $H'\langle p,\langle n,k,q\rangle\rangle:=\langle h\langle n,k\rangle,r(k),\langle p,q\rangle\rangle$  is computable, we obtain the desired conclusion  $(g_1 \twoheadrightarrow f) \leq_{\mathbf{W}} (g_0 \twoheadrightarrow f)$ .

- (2) Let now  $f_0 \leq_W f_1$  hold via computable functions H, K. We prove that we obtain  $(g \to f_0) \leq_{W} (g \to f_1)$ . By the smn-Theorem there are computable functions  $r, h : \mathbb{N} \to \mathbb{N}$  such that
  - $\varphi_{h\langle n,k\rangle}\langle\langle p,q\rangle,t\rangle = H\langle p,\varphi_n\langle q,t\rangle\rangle,$   $\varphi_{r(k)}(p) = \varphi_k \circ K(p)$

for all  $n, k \in \mathbb{N}$  and  $p, q, t \in \mathbb{N}^{\mathbb{N}}$ . Since g is pointed, we have  $dom(g \to f_i) = dom(f_i)$ for  $i \in \{0,1\}$ . Let  $p \in \text{dom}(g \twoheadrightarrow f_0)$ . Then  $K(p) \in \text{dom}(g \twoheadrightarrow f_1)$ . Let  $\langle n, k, q \rangle \in \text{dom}(g \twoheadrightarrow f_1)$ .  $(g \to f_1)K(p)$ . This means that we have  $\emptyset \neq \varphi_n\langle q, g\varphi_kK(p)\rangle \subseteq f_1K(p)$ . Then we

$$\emptyset \neq \varphi_{h\langle n,k\rangle}\langle\langle p,q\rangle, g\varphi_{r(k)}(p)\rangle = H\langle p,\varphi_n\langle q,g\varphi_kK(p)\rangle\rangle \subseteq H\langle p,f_1K(p)\rangle \subseteq f_0(p),$$
  
i.e.,  $\langle h\langle n,k\rangle, r(k), \langle p,q\rangle\rangle \in (g \twoheadrightarrow f_0)(p)$ . This proves  $(g \twoheadrightarrow f_0) \leq_{\mathcal{W}} (g \twoheadrightarrow f_1)$ .

The pointedness assumption is not necessary when we deal with total Weihrauch reducibility. Hence, analogously to the proof of Proposition 2.2 we can obtain the following conclusion.

Corollary 7.8 (Monotonicity of multiplicative implication).  $\rightarrow$  is monotone in the second argument and antitone in the first argument with respect to  $\leq_{tW}$ .

Now we would like to have an analog of Fact 7.2 (3) for  $\rightarrow$ . Unfortunately, this is not possible, but we can say at least the following.

**Proposition 7.9** (Multiplicative implication). For all problems f, g including  $\infty$ :

- $(1) \ f \leq_{\mathbf{W}} g \times h \Longrightarrow (g \twoheadrightarrow f) \leq_{\mathbf{W}} h,$
- (2)  $(g \twoheadrightarrow f) \leq_{\mathbf{W}} h \Longrightarrow f \leq_{\mathbf{W}} g \star h$ , provided that g is pointed,
- (3)  $(g \to f) \leq_{\mathbf{W}} (g \twoheadrightarrow f)$ , provided that g is pointed.

*Proof.* It is routine to check that the claim holds in those cases where the implication takes the values 0 or  $\infty$ . This includes the cases where  $\infty$  is among f, q, h. Otherwise, it suffices to consider problems  $f, g, h :\subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ .

(1) Let  $f \leq_{\mathbf{W}} g \times h$  be witnessed by computable functions H and K. Then there are  $n, k \in \mathbb{N}$  with  $\varphi_n\langle\langle p, r \rangle, s \rangle = H\langle p, \langle s, r \rangle \rangle$  and  $\varphi_k = \pi_1 K$ . We need to prove  $(g \twoheadrightarrow f) \leq_{\mathbf{W}} h$ . We define  $K', H' :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  by  $K' := \pi_2 K$  and  $H'\langle p, r \rangle := \langle n, k, \langle p, r \rangle \rangle$  for all  $p, r \in \mathbb{N}^{\mathbb{N}}$  and  $n, k \in \mathbb{N}$ . Given an input  $p \in \text{dom}(g \twoheadrightarrow f)$ we claim that  $H'(p, hK'(p)) \subseteq (g \to f)(p)$ , i.e., H', K' witness  $(g \to f) \leq_W h$ : if  $\langle n, k, \langle p, r \rangle \rangle \in H'\langle p, hK'(p) \rangle$ , then  $r \in h\pi_2K(p)$  and hence

$$\varphi_n\langle\langle p,r\rangle,g\varphi_k(p)\rangle\subseteq H\langle p,\langle g\pi_1K(p),h\pi_2K(p)\rangle\rangle=H\langle p,\langle g\times h\rangle\circ K(p)\rangle\subseteq f(p).$$

This means  $\langle n, k, \langle p, r \rangle \rangle \in (g \rightarrow f)(p)$ , which proves the claim.

- (2) This follows from (3) together with Fact 7.2.
- (3) Given a  $p \in \text{dom}(g \to f)$  we can use  $(g \twoheadrightarrow f)$  in order to determine a  $\langle n, k, q \rangle \in (g \twoheadrightarrow f)(p)$ . Here we use that g is pointed and hence  $\text{dom}(f \twoheadrightarrow g) = \text{dom}(f \to g)$ . We can then compute a  $t \in \mathbb{N}^{\mathbb{N}}$  with  $\Phi_t(r) = \varphi_n \langle q, r \rangle$  for all  $\langle q, r \rangle \in \text{dom}(\varphi_n)$ . We claim that  $\langle t, \varphi_k(p) \rangle \in (g \to f)(p)$ :

$$\Phi_t \circ g \circ \varphi_k(p) = \varphi_n \langle q, g \circ \varphi_k(p) \rangle \subseteq f(p).$$

This proves the claim.

Again the pointedness assumptions can be removed when we deal with total Weihrauch reducibility and the corresponding completions of operations. In this way Proposition 7.9 shows that we have an instance of a commutative Weihrauch algebra. We formulate this result together with the deductive Weihrauch algebra whose existence follows from Fact 7.2 (3).

Corollary 7.10 (Weihrauch algebra of total Weihrauch degrees). The total Weihrauch degrees give rise to the following Weihrauch algebras:

- (1)  $(\mathcal{W}_{tW}, \leq_{tW}, \overline{\sqcap}, \sqcup, \overline{\times}, \xrightarrow{\twoheadrightarrow}, 1, 1, \infty)$  is a commutative Weihrauch algebra.
- (2)  $(\mathcal{W}_{tW}, \leq_{tW}, \overline{\sqcap}, \sqcup, \overline{\star}, \overline{\to}, 1, 1, \infty)$  is a deductive Weihrauch algebra.

It would be desirable to have an equivalence in Proposition 7.9 (1) instead of just an implication, which would mean that  $\rightarrow$  is a co-residual operation of  $\times$  in the same way as  $\rightarrow$  is a co-residual of  $\star$ . However, in [8, Proposition 37] it was proved that there is no such co-residual operation to  $\times$ . The following result shows that  $\rightarrow$  has such a co-residual property at least restricted to special problems.

**Proposition 7.11** (Multiplicative deduction).  $(g \to f) \leq_W h \Longrightarrow f \leq_W \widehat{\overline{g}} \times h$  for all problems f, g, h including  $\infty$ , such that g is pointed.

Proof. It is routine to check the claim for the special cases where the problem  $\infty$  is involved. Otherwise, it suffices to consider problems  $f,g,h:\subseteq\mathbb{N}^\mathbb{N} \rightrightarrows \mathbb{N}^\mathbb{N}$ . Let g be pointed and let  $(g \twoheadrightarrow f) \leq_{\mathbb{W}} h$  hold via computable H,K. Then given a point  $p \in \mathrm{dom}(f) = \mathrm{dom}(g \twoheadrightarrow f)$  any  $\langle n,k,q \rangle \in H\langle p,hK(p) \rangle$  satisfies  $\emptyset \neq \varphi_n\langle q,g \circ \varphi_k(p) \rangle \subseteq f(p)$ . Since  $\overline{\mathbb{N}^\mathbb{N}}$  has a precomplete representation  $\delta_{\overline{\mathbb{N}^\mathbb{N}}}$ , it follows that there is a total computable universal function  $u:\mathbb{N}^\mathbb{N} \to \mathbb{N}^\mathbb{N}$  with  $\delta_{\overline{\mathbb{N}^\mathbb{N}}} \circ u\langle k,p \rangle = \delta_{\overline{\mathbb{N}^\mathbb{N}}}(\varphi_k(p)+1) = \varphi_k(p)$  for all  $k \in \mathbb{N}$  and  $p \in \mathrm{dom}(\varphi_k)$ . We define a total computable function  $K'(p) := \langle \langle u\langle 0,p \rangle, u\langle 1,p \rangle, u\langle 2,p \rangle, \ldots \rangle, K(p) \rangle$  and a computable function  $H'\langle p, \langle \langle q_0,q_1,q_2,\ldots \rangle,r \rangle\rangle := \varphi_n\langle q,q_k-1 \rangle$  where  $\langle n,k,q \rangle = H\langle p,r \rangle$ . Whenever G is a realizer of  $\widehat{\overline{g}}$ , with respect to  $\delta_{\overline{\mathbb{N}^\mathbb{N}^\mathbb{N}}}$ , then we obtain

$$H'\langle p, \langle G \times h \rangle \circ K'(p) \rangle = H'\langle p, \langle G \langle u \langle 0, p \rangle, u \langle 1, p \rangle, u \langle 2, p \rangle, \ldots \rangle, hK(p) \rangle \rangle$$

$$\subseteq \bigcup \{ \varphi_n \langle q, g \circ \varphi_k(p) \rangle : \langle n, k, q \rangle \in H\langle p, hK(p) \rangle \}$$

$$\subseteq f(p),$$

i.e., 
$$f \leq_{\mathbf{W}} \widehat{\overline{g}} \times h$$
.

The basic idea of the proof is that using the parallelization we can evaluate  $\overline{g}$  on all possible inputs  $\varphi_k(p)$  with Gödel numbers  $k \in \mathbb{N}$  and only after we learn the result of h we know which of these values is actually needed. The completion guarantees that all these values actually exist.

A similar idea as in the proof of Proposition 7.11 has been independently used by Neumann and Pauly [18, Proposition 31] to prove the following result, which we rephrase in terms of our terminology.<sup>9</sup>

<sup>&</sup>lt;sup>9</sup>The notion of precompleteness used by Neumann and Pauly is not the usual one; what is required is rather a uniform version of completeness, which is satisfied by our completion  $\overline{g}$ .

**Proposition 7.12** (Neumann and Pauly 2018).  $g \star h \leq_{\mathbf{W}} \widehat{\overline{g}} \times h$  for all problems gand  $h :\subseteq X \Longrightarrow \mathbb{N}$ .

This result yields a similar transition from  $g \star h$  to  $\widehat{g} \times h$  as the one that happens from Proposition 7.9 to 7.11, except that we do not need problems h with natural number output for the latter transition. We obtain the following obvious corollary of Proposition 7.11.

Corollary 7.13 (Multiplicative deduction).  $(g \rightarrow f) \leq_{W} h \iff f \leq_{W} g \times h$  for all problems f, g, h including  $\infty$  and such that g is parallelizable and complete.

This is the key observation that is used in the next section in order to show that the parallelized total Weihrauch degrees form a Brouwer algebra. We note that by [8, Proposition 37] it is known that there is no way to define → such that the statement in Corollary 7.13 holds for all problems g. This remains so, even if we replace Weihrauch reductions  $\leq_W$  by total Weihrauch reductions  $\leq_{tW}$  and the product  $\times$  by its completion  $\overline{\times}$ , as a refined version of the argument from [8, Proposition 37] shows.

**Proposition 7.14.** The operation  $\overline{\times}$  is not co-residuated and  $\overline{\star}$  is not left coresiduated with respect to  $\leq_{tW}$ .

Proof. We have

- $(1) \ \mathsf{C}_{2^{\mathbb{N}}} \times \overline{\mathsf{C}_{\mathbb{N}}} \leq_{\mathrm{W}} \overline{\mathsf{C}_{\mathbb{N}}} \times (\mathsf{C}_{2^{\mathbb{N}}} \sqcup \overline{\mathsf{C}_{\mathbb{N}}}),$

While (1) and (2) are clear, it remains to justify (3) and (4). We obtain (3) since  $C_{2^{\mathbb{N}}} \star C_{2^{\mathbb{N}}} \equiv_{W} C_{2^{\mathbb{N}}}$  and by distributivity properties of  $\star$  [8, Proposition 39]

$$\begin{split} (\mathsf{C}_{2^{\mathbb{N}}} \sqcap \overline{\mathsf{C}_{\mathbb{N}}}) \star (\mathsf{C}_{2^{\mathbb{N}}} \sqcup \overline{\mathsf{C}_{\mathbb{N}}}) & \equiv_{W} \quad ((\mathsf{C}_{2^{\mathbb{N}}} \sqcap \overline{\mathsf{C}_{\mathbb{N}}}) \star \mathsf{C}_{2^{\mathbb{N}}}) \sqcup ((\mathsf{C}_{2^{\mathbb{N}}} \sqcap \overline{\mathsf{C}_{\mathbb{N}}}) \star \overline{\mathsf{C}_{\mathbb{N}}}) \\ & \leq_{W} \quad (\mathsf{C}_{2^{\mathbb{N}}} \sqcap (\overline{\mathsf{C}_{\mathbb{N}}} \star \mathsf{C}_{2^{\mathbb{N}}})) \sqcup ((\mathsf{C}_{2^{\mathbb{N}}} \star \overline{\mathsf{C}_{\mathbb{N}}}) \sqcap (\overline{\mathsf{C}_{\mathbb{N}}} \star \overline{\mathsf{C}_{\mathbb{N}}})) \\ & \leq_{W} \quad \mathsf{C}_{2^{\mathbb{N}}} \sqcup (\overline{\mathsf{C}_{\mathbb{N}}} \star \overline{\mathsf{C}_{\mathbb{N}}}). \end{split}$$

Now we need to justify why (4) holds. Since  $C_{2^{\mathbb{N}}}$  is a fractal by [2, Corollary 5.6], [5, Fact 3.2] and  $\overline{C_N}$  is a fractal as proved in [4, Lemma 8.7], it follows that  $C_{2^N} \times \overline{C_N}$ is a fractal and hence join irreducible by [5, Proposition 2.6] . This means that  $C_{2^{\mathbb{N}}} \times \overline{C_{\mathbb{N}}} \leq_{W} C_{2^{\mathbb{N}}} \sqcup (\overline{C_{\mathbb{N}}} \star \overline{C_{\mathbb{N}}}) \text{ would imply that } C_{2^{\mathbb{N}}} \times \overline{C_{\mathbb{N}}} \leq_{W} C_{2^{\mathbb{N}}} \text{ or } C_{2^{\mathbb{N}}} \times \overline{C_{\mathbb{N}}} \leq_{W} \overline{C_{\mathbb{N}}} \star \overline{C_{\mathbb{N}}}$ holds. The latter is impossible, as  $C_{2^N}$  has computable inputs without computable solutions, while  $\overline{C_{\mathbb{N}}} \star \overline{C_{\mathbb{N}}}$  has computable solutions for all inputs. The former is impossible as even  $C_{\mathbb{N}} \not\leq_{\mathrm{W}} C_{2^{\mathbb{N}}}$ .

By Propositions 6.3 and 7.6 all degrees that appear in (1)–(4) are complete, as  $C_{2^{\mathbb{N}}}$  is complete by Corollary 6.4. Hence, all the statements (1)-(4) hold true if we replace  $\leq_{\mathrm{W}}$  by  $\leq_{\mathrm{tW}}$ . Suppose now a binary operation  $\square$  would exist such that  $(g \Box f) \leq_{\text{tW}} h \iff f \leq_{\text{tW}} h \overline{\times} g$  holds for all problems f, g, h. We consider g := $(\mathsf{C}_{2^{\mathbb{N}}}\sqcup\overline{\mathsf{C}_{\mathbb{N}}})$  and  $h_1:=\overline{\mathsf{C}_{\mathbb{N}}},\,h_2:=\mathsf{C}_{2^{\mathbb{N}}}$  and  $f:=\mathsf{C}_{2^{\mathbb{N}}}\times\overline{\mathsf{C}_{\mathbb{N}}}.$  Then by (1)–(4)  $f\leq_{\mathrm{tW}}h_1\overline{\times}g$ and  $f \leq_{\text{tW}} h_2 \overline{\times} g$ , but  $f \not\leq_{\text{tW}} (h_1 \overline{\sqcap} h_2) \overline{\star} g$ , which also implies  $f \not\leq_{\text{tW}} (h_1 \overline{\sqcap} h_2) \overline{\times} g$  This simultaneously shows that  $\square$  does not exist and also a corresponding operation for  $\overline{\star}$  does not exist.

The Weihrauch algebra of total Weihrauch degrees fails in two different ways being a model of some intuitionistic linear logic. The multiplicative and compositional versions of the algebra both fail to be Troelstra algebras, the former is not deductive, the latter is not commutative.

Corollary 7.15. The Weihrauch algebras from Corollary 7.10 are not Troelstra algebras, i.e.,

- (1)  $(W_{tW}, \leq_{tW}, \overline{\sqcap}, \sqcup, \overline{\times}, \xrightarrow{\twoheadrightarrow}, 1, 1, \infty)$  is not deductive,
- (2)  $(\mathcal{W}_{tW}, \leq_{tW}, \overline{\sqcap}, \sqcup, \overline{\star}, \overline{\to}, 1, 1, \infty)$  is not commutative.
  - 8. The Brouwer Algebra of Parallelizable Total Degrees

In [3] we have already studied parallelized Weihrauch reducibility  $\leq_{pW}$ , which is the reducibility that is generated by the closure operator of parallelization on  $\leq_{W}$ . Likewise we want to study parallelized total Weihrauch reducibility  $\leq_{ptW}$ .

**Definition 8.1** (Parallelized Weihrauch reducibility). For problems f, g we write

- $(1) \ f \leq_{\mathrm{pW}} g : \iff f \leq_{\mathrm{W}} \widehat{g} \qquad \qquad \text{(parallelized Weihrauch reducibility)}$
- (2)  $f \leq_{\text{ptW}} g : \iff f \leq_{\text{W}} \widehat{\overline{g}}$  (parallelized total Weihrauch reducibility)

Analogously, we write  $\equiv_{pW}$  and  $\equiv_{ptW}$  for the corresponding equivalences.

It is clear that  $\leq_{\text{pW}}$  and  $\leq_{\text{ptW}}$  are actually preorders by Propositions 2.2, as completion and parallelized completion are closure operators (the latter by Proposition 6.13). We note that we also have  $f \leq_{\text{ptW}} g \iff f \leq_{\text{tW}} \widehat{\overline{g}}$  by Proposition 6.3. It is important to mention that the order in which we apply the closure operators matters. While  $\widehat{\overline{g}}$  is always complete and parallelizable,  $\widehat{\overline{g}}$  is always complete, but not necessarily parallelizable (see Lemma 6.14).

For each operation  $\Box \in \{\times, \sqcup, \boxplus, \sqcap, +, \star, \to, \twoheadrightarrow\}$  we define its parallelized completion  $\widehat{\Box}$  by  $f\widehat{\Box}g := \widehat{\overline{f}}\Box\widehat{\overline{g}}$ . Since parallelized completion is a closure operator for  $\leq_{\mathrm{W}}$  by Proposition 6.13, we straightforwardly obtain the following by Proposition 2.2.

Corollary 8.2 (Monotonicity).

- (1)  $(f,g) \mapsto f \widehat{\Box} g \text{ for } \Box \in \{\times, \sqcup, \boxplus, \sqcap, +, \star\} \text{ is monotone with respect to } \leq_{\text{ptW}}.$
- (2)  $(f,g) \mapsto f \widehat{\Box} g$  for  $\Box \in \{\rightarrow, \twoheadrightarrow\}$  is monotone with respect to  $\leq_{\text{ptW}}$  in the second argument and antitone in the first argument.

*Proof.* The corresponding monotonicity properties with respect to  $\leq_{\mathrm{W}}$  are known by [6, Proposition 3.6], except for  $\twoheadrightarrow$ :  $\overline{\twoheadrightarrow}$  is monotone with respect to  $\leq_{\mathrm{tW}}$  by Corollary 7.8. Hence the claims follow from Proposition 2.2.

An interesting property of parallelized (total) Weihrauch reducibility is that suprema and products are merged in a certain sense. We summarize some facts regarding preservation and co-preservation of parallelization that were proved in [3, Propositions 4.5, 4.8, 4.9] and [8, Propositions 41, 44].

Fact 8.3 (Parallelization and algebraic operations). For all problems f, g including  $\infty$ :

- $(1) \widehat{f \times g} \equiv_{\mathrm{sW}} \widehat{f} \times \widehat{g} \leq_{\mathrm{W}} \widehat{f} \sqcup \widehat{g},$
- (2)  $\widehat{f} \sqcup \widehat{g} \leq_{\mathbf{W}} \widehat{f} \sqcup \widehat{g} \leq_{\mathbf{W}} \widehat{\widehat{f}} \sqcup \widehat{g}$ ,
- $(3) \widehat{f \cap g} \leq_{\mathrm{sW}} \widehat{f} \cap \widehat{g} \equiv_{\mathrm{sW}} \widehat{f} \cap \widehat{g},$
- $(4) \widehat{f \star g} \leq_{\mathrm{sW}} \widehat{f} \star \widehat{g} \equiv_{\mathrm{sW}} \widehat{\widehat{f}} \star \widehat{\widehat{g}},$
- (5)  $\widehat{\overline{f}} \times \overline{g} \equiv_{\text{sW}} \widehat{\overline{f}} \times \widehat{\overline{g}} \equiv_{\text{W}} \widehat{\overline{f}} \sqcup \overline{g} \equiv_{\text{W}} \widehat{\overline{f}} \sqcup g$

Hence,  $\overline{\times}$  and  $\sqcup$  are equivalent operations under parallelized total Weihrauch reducibility. This follows from Fact 8.3 and Proposition 2.5.

Corollary 8.4 (Products and coproducts).  $\overline{f} \times \overline{g} \equiv_{\text{ptW}} \widehat{\overline{f}} \times \widehat{\overline{g}} \equiv_{\text{ptW}} \widehat{\overline{f}} \sqcup \widehat{\overline{g}} \equiv_{\text{ptW}} f \sqcup g$  for all problems f, g.

By  $W_{ptW}$  we denote the class of parallelized total Weihrauch degrees including  $\infty$ . We use the same notation  $\leq_{ptW}$  for the order on degrees and we consider the

operations to be extended to these degrees. In order to avoid too clumsy notation we use the abbreviation  $\Rightarrow$  for  $\stackrel{\frown}{\Rightarrow}$  in the following. We prove that the parallelized total Weihrauch degrees form a Brouwer algebra.

**Theorem 8.5** (Brouwer algebra).  $(W_{ptW}, \leq_{ptW}, \overline{\sqcap}, \sqcup, \Rightarrow, 1, \infty)$  is a Brouwer alge-

*Proof.*  $(W_{ptW}, \leq_{ptW}, \overline{\Box}, \sqcup)$  is a lattice by Proposition 2.2 as parallelized completion is a closure operator. We obtain by Corollary 7.13 and Fact 8.3

$$(g \Rrightarrow f) \leq_{\mathrm{ptW}} h \quad \Longleftrightarrow \quad (\widehat{\overline{g}} \twoheadrightarrow \widehat{\overline{f}}) \leq_{\mathrm{W}} \widehat{\overline{h}}$$

$$\Longleftrightarrow \quad \widehat{\overline{f}} \leq_{\mathrm{W}} \widehat{\overline{g}} \times \widehat{\overline{h}}$$

$$\Longleftrightarrow \quad \widehat{\overline{f}} \leq_{\mathrm{W}} \widehat{\overline{g} \sqcup \overline{h}}$$

$$\Longleftrightarrow \quad f \leq_{\mathrm{ptW}} g \sqcup h.$$

This proves the claim.

In [3] we have proved that the Medvedev lattice can be embedded into the parallelized Weihrauch lattice. This embedding can actually be extended to a Brouwer algebra embedding into the parallelized total Weihrauch lattice. We recall some basic definitions for the Medvedev lattice [24]. Let  $A, B \subseteq \mathbb{N}^{\mathbb{N}}$ . Then A is said to be Medvedev reducible to B, in symbols  $A \leq_{\mathbf{M}} B$ , if there is a computable function  $F:\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  such that  $B\subseteq \text{dom}(F)$  and  $F(B)\subseteq A$ . We recall the definition of the algebraic operations of the Medvedev lattice:

- (1)  $A \otimes B := 0A \cup 1B = \langle A \sqcup B \rangle$ ,
- $(2) \ A \oplus B := \langle A \times B \rangle,$
- $(3) B \to A := \{ \langle n, q \rangle \in \mathbb{N}^{\mathbb{N}} : (\forall p \in B) \varphi_n \langle q, p \rangle \in A \}.$

By  $\mathcal{M}$  we denote the set of Medvedev degrees. We identify degrees with their members and use the same notation for the algebraic operations on degrees. Medvedev [17] proved that  $(\mathcal{M}, \otimes, \oplus, \to, \mathbb{N}^{\mathbb{N}}, \emptyset)$  is a Brouwer algebra (see [24, Theorem 9.1]). In [3] we have considered the constant problems

$$c_A: \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}, p \mapsto A$$

for every non-empty  $A \subseteq \mathbb{N}^{\mathbb{N}}$  and  $c_{\emptyset} = \infty$ . The following facts were proved in [3, Theorem 5.1].

**Fact 8.6** (Medvedev embedding). For all  $A, B \subseteq \mathbb{N}^{\mathbb{N}}$ :

- $(1) A \leq_{\mathbf{M}} B \iff c_A \leq_{\mathbf{W}} c_B,$
- (2)  $c_{A\otimes B} \equiv_{sW} c_A \sqcap c_B$ , (3)  $c_{A\oplus B} \equiv_{sW} c_A \times c_B \equiv c_A \star c_B$ .

The equivalence  $c_A \times c_B \equiv_{\mathrm{W}} c_A \star c_B$ , was not proved in the references, but it is easy to see. For one,  $f \times g \leq_W f \star g$  holds in general and on the other hand,  $c_A \star c_B = \langle \operatorname{id} \times c_A \rangle \circ U \circ \langle \operatorname{id} \times c_B \rangle \leq_W c_A \times c_B$ , as the output of  $c_A$  does not depend on the input. Here we add the observation that also the implication is preserved. In fact, since the product and the compositional product for problems of the form  $c_A$  coincide, also the multiplicative and compositional implications coincide.

**Lemma 8.7** (Medvedev implication).  $c_{B\to A} \equiv_{\mathcal{W}} (c_B \to c_A) \equiv_{\mathcal{W}} (c_B \to c_A)$  for all  $A, B \subseteq \mathbb{N}^{\mathbb{N}}$ .

*Proof.* It is routine to check the special cases of problems that involve  $A, B \in$  $\{\emptyset, \mathbb{N}^{\mathbb{N}}\}$ . Since the Medvedev lattice is a Brouwer algebra by [24, Theorem 9.1], we have  $A \leq_{\mathrm{M}} B \oplus (B \to A)$ . With the help of Proposition 7.9 and Fact 8.6 we obtain

$$A \leq_{\mathrm{M}} B \oplus (B \to A) \implies c_A \leq_{\mathrm{W}} c_{B \oplus (B \to A)} \equiv_{\mathrm{W}} c_B \times c_{B \to A}$$
$$\implies (c_B \twoheadrightarrow c_A) \leq_{\mathrm{W}} c_{B \to A}.$$

We can also prove  $c_{B\to A} \leq_{\mathrm{W}} (c_B \twoheadrightarrow c_A)$ . Given a  $p \in \mathbb{N}^{\mathbb{N}}$  we obtain

$$\langle n, k, q \rangle \in (c_B \twoheadrightarrow c_A)(p) \iff \emptyset \neq \varphi_n \langle q, c_B \circ \varphi_k(p) \rangle \subseteq c_A(p)$$
  
 $\iff \langle n, q \rangle \in c_{B \to A}(p).$ 

Hence,  $c_{B\to A} \leq_{\mathrm{W}} (c_B \to c_A)$  follows. We have  $(c_B \to c_A) \leq_{\mathrm{W}} (c_B \to c_A)$  by Proposition 7.9. We also obtain  $(c_B \to c_A) \leq_{\mathrm{W}} (c_B \to c_A)$ . To this end, let h be a problem such that  $c_A \leq_{\mathrm{W}} c_B \star h$ . Like above we obtain  $c_B \star h \leq_{\mathrm{W}} c_B \times h$ , since the output of  $c_B$  does not depend on its input. That means  $c_A \leq_{\mathrm{W}} c_B \times h$  and hence  $(c_B \to c_A) \leq_{\mathrm{W}} h$  by Proposition 7.9. However, if g is a problem such that  $c_A \leq_{\mathrm{W}} c_B \star h$  implies  $g \leq_{\mathrm{W}} h$  for every h, then  $g \leq_{\mathrm{W}} (c_B \to c_A)$  follows. Hence,  $(c_B \to c_A) \leq_{\mathrm{W}} (c_B \to c_A)$ .

Hence the map  $A \mapsto c_A$  is a lattice embedding from the Medvedev lattice into the Weihrauch lattice that also preserves the corresponding implications (even though the Weihrauch lattice itself is not a Brouwer algebra). It is easy to see that every Weihrauch degree of the form  $c_A$  with  $A \subseteq \mathbb{N}^{\mathbb{N}}$  is parallelizable and complete, i.e.,  $\overline{c_A} \equiv_{\mathbb{W}} c_A$ . Hence the above embedding is also an embedding into the parallelized total Weihrauch degrees. We note that  $c_{\emptyset} = \infty$  and  $c_{\mathbb{N}^{\mathbb{N}}} \equiv_{\mathbb{W}} 1$ . Hence, we obtain a Brouwer algebra embedding, i.e., a lattice embedding that preserves the implication and the lower and upper bound.

**Theorem 8.8** (Embedding of the Medvedev lattice).  $c: \mathcal{M} \to \mathcal{W}_{ptW}, A \mapsto c_A$  is a Brouwer algebra embedding.

The fact that the parallelized total Weihrauch lattice is a Brouwer algebra implies that it is a model for some intermediate logic (i.e., some propositional logic intermediate between intuitionistic logic and classical logic). The existence of an embedding from the Medvedev lattice into the parallelized total Weihrauch lattice allows us to conclude that the logic of the parallelized total Weihrauch lattice is Jankov logic, i.e., the deductive closure of intuitionistic logic together with the weak principle of excluded middle  $\neg \neg A \lor \neg A$ . We follow Sorbi [23, 24] for a formal definition of the theory of a Brouwer algebra. Let Form denote the set of well formed propositional formulas. Then we call a map  $v : \text{Form} \to \mathcal{W}_{\text{ptW}}$  valuation if it satisfies the following for all  $A, B \in \text{Form}$ :

- (1)  $v(A \vee B) = v(A) \widehat{\Box} v(B)$ ,
- (2)  $v(A \wedge B) = v(A) \sqcup v(B)$ ,
- (3)  $v(A \to B) = (v(A) \Longrightarrow v(B)),$
- (4)  $v(\neg A) = (v(A) \Rightarrow \infty)$ .

We write  $\mathcal{W}_{\text{ptW}} \vDash A$  if v(A) = 1 for all valuations v. Then the set of formulas  $\text{Th}(\mathcal{W}_{\text{ptW}}) := \{A \in \text{Form} : \mathcal{W}_{\text{ptW}} \vDash A\}$  is called the *theory* of  $\mathcal{W}_{\text{ptW}}$ . It was proved by Medvedev [16] (see [24, Corollary 6.4]) that the theory of the Brouwer algebra  $\mathcal{M}$  is Jankov logic. We obtain the same result for our Brouwer algebra  $\mathcal{W}_{\text{ptW}}$ . For one, it contains Jankov logic by Corollary 7.5. On the other hand, it cannot validate any additional propositional formulas as the Medvedev Brouwer algebra is embeddable by Theorem 8.8.

Corollary 8.9 (Theory of the parallelized complete Weihrauch degrees). The theory of the Brouwer algebra  $W_{ptW}$  is Jankov logic.

We note that Higuchi and Pauly proved [13, Theorems 4.1, 4.2] that neither the Weihrauch lattice by itself nor the parallelized Weihrauch lattice (restricted to the pointed problems) is a Brouwer algebra. Hence, the closure operator of completion seems to be essential in order to obtain a Brouwer algebra.

In view of Corollary 7.10 one could obtain a way to transform the total Weihrauch lattice into a Troelstra algebra by restricting it to a linear fragment. We call

 $\mathcal{L} \subseteq \mathcal{W}_{\text{ptW}}$  linear if  $f \times g \equiv_{\text{tW}} f * g$  holds for all  $f, g \in \mathcal{L}$ . If there would be any linear sublattice of interest that also preserves the monoid structure, then that would be a potential candidate for a Troelstra algebra. We note that the constant multi-valued problems  $c_A$  used for the embedding of the Medvedev lattice form a linear subset of the total Weihrauch degrees by Fact 8.6, however, this is not a sublattice and leads directly to a Brouwer algebra, i.e., a trivial example of a Troelstra algebra.




Figure 3. Problems in the parallelized total Weihrauch lattice  $W_{ptW}$ 

## 9. Conclusion

We have proved that the Weihrauch lattice can be transformed into a Brouwer algebra by completion followed by parallelization. It would be desirable to understand the structure of this Brouwer algebra somewhat better. Is it isomorphic to the Medvedev Brouwer algebra? Presumably not, as the Medvedev algebra considers only problems that are independent of the input. However, we need more structural information on the lattices and algebras in order to prove such properties. The Medvedev lattice has, for instance, a second smallest degree, called 0', which consists of all non-computable  $p \in \mathbb{N}^{\mathbb{N}}$ . Is there such a second smallest degree in the parallelized total Weihrauch lattice? Or is the structure dense? We do not even know the answer to this question for the ordinary Weihrauch lattice or its total variant. What we can say, though, is that the parallelized total Weihrauch lattice is still inhabited by a variety of interesting problems. The diagram in Figure 3 shows a number of problems (that are taken without further explanation from [7] and [9]), and that inhabit  $W_{ptW}$ . Even though a lot of problems that are normally separated in the Weihrauch lattice are identified in  $\mathcal{W}_{ptW}$ , the structure is still rich and non-linear.

## ACKNOWLEDGMENTS

We would like to thank Paulo Oliva for discussions of models of intuitionistic linear logic at the Logic Colloquium 2018 in Udine that have helped us to identify the relevance of Troelstra and Weihrauch algebras.

#### References

- [1] Vasco Brattka. A Galois connection between Turing jumps and limits. Logical Methods in Computer Science, 14(3:13):1–37, August 2018.
- [2] Vasco Brattka, Matthew de Brecht, and Arno Pauly. Closed choice and a uniform low basis theorem. Annals of Pure and Applied Logic, 163:986–1008, 2012.
- [3] Vasco Brattka and Guido Gherardi. Weihrauch degrees, omniscience principles and weak computability. The Journal of Symbolic Logic, 76(1):143–176, 2011.
- [4] Vasco Brattka and Guido Gherardi. Completion of choice. arXiv 1910.13186, 2019.
- [5] Vasco Brattka, Guido Gherardi, and Alberto Marcone. The Bolzano-Weierstrass theorem is the jump of weak Kőnig's lemma. Annals of Pure and Applied Logic, 163:623–655, 2012.
- [6] Vasco Brattka, Guido Gherardi, and Arno Pauly. Weihrauch complexity in computable analvsis. Accepted for publication, 2018.
- [7] Vasco Brattka, Matthew Hendtlass, and Alexander P. Kreuzer. On the uniform computational content of computability theory. Theory of Computing Systems, 61(4):1376–1426, 2017.
- [8] Vasco Brattka and Arno Pauly. On the algebraic structure of Weihrauch degrees. Logical Methods in Computer Science, 14(4:4):1–36, 2018.
- [9] Vasco Brattka and Tahina Rakotoniaina. On the uniform computational content of Ramsey's theorem. *Journal of Symbolic Logic*, 82(4):1278–1316, 2017.
- [10] Damir D. Dzhafarov. Joins in the strong Weihrauch degrees. Mathematical Research Letters, 26(3):749-767, 2019.
- [11] Ju. L. Eršov. Theory of numberings. In Edward R. Griffor, editor, Handbook of Computability Theory, volume 140 of Studies in Logic and the Foundations of Mathematics, pages 473–503. Elsevier, Amsterdam, 1999.
- [12] Nikolaos Galatos, Peter Jipsen, Tomasz Kowalski, and Hiroakira Ono. Residuated lattices: an algebraic glimpse at substructural logics, volume 151 of Studies in Logic and the Foundations of Mathematics. Elsevier B. V., Amsterdam, 2007.
- [13] Kojiro Higuchi and Arno Pauly. The degree structure of Weihrauch reducibility. Log. Methods Comput. Sci., 9(2):2:02, 17, 2013.
- [14] Alexander S. Kechris. Classical Descriptive Set Theory, volume 156 of Graduate Texts in Mathematics. Springer, Berlin, 1995.
- [15] Christoph Kreitz and Klaus Weihrauch. Theory of representations. Theoretical Computer Science, 38:35–53, 1985.
- [16] Ju. T. Medvedev. Finitive problems. Doklady Akademii Nauk SSSR, 142:1015-1018, 1962.
- [17] Yu. T. Medvedev. Degrees of difficulty of the mass problem. Doklady Akademii Nauk SSSR, 104:501–504, 1955.
- [18] Eike Neumann and Arno Pauly. A topological view on algebraic computation models. *Journal of Complexity*, 44(Supplement C):1–22, 2018.
- [19] Arno Pauly. On the (semi)lattices induced by continuous reducibilities. Mathematical Logic Quarterly, 56(5):488–502, 2010.
- [20] Matthias Schröder. Admissible Representations for Continuous Computations. PhD thesis, Fachbereich Informatik, FernUniversität Hagen, 2002.
- [21] Stephen G. Simpson. Subsystems of Second Order Arithmetic. Perspectives in Logic, Association for Symbolic Logic. Cambridge University Press, Poughkeepsie, second edition, 2009.
- [22] Robert I. Soare. Recursively Enumerable Sets and Degrees. Perspectives in Mathematical Logic. Springer, Berlin, 1987.
- [23] Andrea Sorbi. Embedding Brouwer algebras in the Medvedev lattice. Notre Dame Journal of Formal Logic, 32(2):266–275, 1991.
- [24] Andrea Sorbi. The Medvedev lattice of degrees of difficulty. In Computability, enumerability, unsolvability, volume 224 of London Math. Soc. Lecture Note Ser., pages 289–312. Cambridge Univ. Press, Cambridge, 1996.
- [25] A. S. Troelstra. Lectures on linear logic, volume 29 of CSLI Lecture Notes. Stanford University, Center for the Study of Language and Information, Stanford, CA, 1992.
- [26] A.S. Troelstra. Comparing the theory of representations and constructive mathematics. In E. Börger, G. Jäger, H. Kleine Büning, and M.M. Richter, editors, Computer Science Logic, volume 626 of Lecture Notes in Computer Science, pages 382–395, Berlin, 1992. Springer. Proceedings of the 5th Workshop, CSL'91, Berne Switzerland, October 1991.
- [27] Klaus Weihrauch. Computable Analysis. Springer, Berlin, 2000.
- [28] David N. Yetter. Quantales and (noncommutative) linear logic. The Journal of Symbolic Logic, 55(1):41-64, 1990.

Faculty of Computer Science, Universität der Bundeswehr München, Germany and Department of Mathematics & Applied Mathematics, University of Cape Town, South Africa $^{10}$ 

 $E\text{-}mail\ address{:}\ {\tt Vasco.Brattka@cca-net.de}$ 

DIPARTIMENTO DI FILOSOFIA E COMUNICAZIONE, UNIVERSITÀ DI BOLOGNA, ITALY

 $E\text{-}mail\ address: \verb"Guido.Gherardi@unibo.it"$ 

 $<sup>^{10}\</sup>mathrm{Vasco}$ Brattka has received funding from the National Research Foundation of South Africa