DEFINABILITY OF HENSELIAN VALUATIONS BY CONDITIONS ON THE VALUE GROUP

LOTHAR SEBASTIAN KRAPP, SALMA KUHLMANN, AND MORITZ LINK

Abstract. Given a henselian valuation, we study its definability (with and without parameters) by examining conditions on the value group. We show that any henselian valuation whose value group is not closed in its divisible hull is definable in the language of rings, using one parameter. Thereby we strengthen known definability results. Moreover, we show that in this case, one parameter is optimal in the sense that one cannot obtain definability without parameters. To this end, we present a construction method for a *t*-henselian non-henselian ordered field elementarily equivalent to a henselian field with a specified value group.

§1. Introduction. In recent years, the study of definable henselian valuations has received a considerable amount of attention and several definability conditions have been established. Early results of this study were related to decidability questions (cf. e.g. [26, 2]), whereas a more recent motivation is due to the Shelah–Hasson Conjecture on the classification of NIP fields (cf. e.g. [27, 17, 18, 5, 10, 11, 12, 22]). We refer the reader to [9] for a more detailed survey on the definability of henselian valuations. While some of these results deal with the existence of non-trivial definable henselian valuations and others consider the quantifier complexity of defining formulas (cf. [9, Sections 2 & 3]), in this paper we are interested in the definability of a given henselian valuation with a certain value group.¹

The strongest currently known definability results which only pose conditions on the value group of a given henselian valuation are exhibited in [14] and [21]. In the following, we state the results under consideration for this paper and outline how we strengthen these.

FACT 1.1. [14, Corollary 2]. Let (K, v) be a henselian valued field such that the value group vK is discretely ordered. Then v is \mathcal{L}_r -definable with one parameter from K.

For densely ordered value groups, the following is established.

The first author was partially supported by Werner und Erika Messmer Stiftung.

During parts of this project, the second and third authors were supported by the German Excellence Strategy and the German Research Foundation.

¹There are also conditions for the definability of a given henselian valuation only depending on its residue field (see our Concluding Remark).

FACT 1.2. [14, Theorem 3]. Let (K, v) be a henselian valued field such that vK is densely ordered. Assume that vK contains a convex subgroup that is p-regular but not p-divisible, for some prime $p \in \mathbb{N}$. Then v is \mathcal{L}_r -definable with one parameter from K.

While in [14] definability is treated in the language of rings \mathcal{L}_r , the definability results in [21] are stated for ordered fields in the language of ordered rings \mathcal{L}_{or} .

FACT 1.3. [21, Theorem 5.3 (2)]. Let (K, <, v) be an ordered henselian valued field such that vK is not closed in its divisible hull. Then v is \mathcal{L}_{or} -definable with one parameter from K.

After introducing preliminary notions in Section 2, we show in Section 3 that the hypothesis on vK in Fact 1.2 is strictly stronger than that in Fact 1.3 (see Proposition 3.3 and Example 3.4). Thus, we obtain the following strengthening of both Fact 1.2 and Fact 1.3:²

THEOREM A. Let (K, v) be a henselian valued field such that vK is not closed in its divisible hull. Then v is \mathcal{L}_r -definable with one parameter from K.

In Section 4, we examine definability without parameters of the valuations. To this end, in Construction 4.5 – the technical heart of this paper – we refine [25, page 338] to obtain a t-henselian non-henselian field (see Definition 4.1) with prescribed value groups. More precisely, given an arbitrary family of non-trivial ordered abelian groups $\{A_{\gamma} \mid \gamma \in -\omega\}$, Construction 4.5 produces a t-henselian non-henselian ordered field (K, <) as well as, for each $n \in \mathbb{N}$, a convex valuation u_n on K whose value group is the Hahn sum (see page 3) over $\{A_{\gamma} \mid \gamma \leq -n\}$.

By applying Construction 4.5 to the specific case where all A_{γ} are identical, we obtain the following:³

THEOREM B. Let $A \neq \{0\}$ be an ordered abelian group. Then there exists an ordered henselian valued field (L, <, v) such that vL is the Hahn sum $\coprod_{-\omega} A$ and the only parameter-free \mathcal{L}_{or} -definable henselian valuation on (L, <) is the trivial valuation. In particular, v is not \mathcal{L}_{or} -definable without parameters.

Finally, by choosing appropriately the group in Theorem B, we can show that Fact 1.1 - 1.3 and Theorem A are optimal in the sense that, in general, one cannot obtain definability of the valuations *without* parameters (see Example 4.9 and Example 4.10).

We conclude this work in Section 5 by collecting open questions motivated by our results.

§2. Preliminaries. More details on the model and valuation theoretic notions we use can be found in [24, 6, 23].

We denote by \mathbb{N} the set of natural numbers without 0 and by ω the set of natural numbers with 0. All topological properties are considered with respect to the order topology. Whenever an algebraic structure carries a standard linear ordering, we refer to this specific ordering if not explicitly stated otherwise.

 $^{^2\}mathrm{Theorem}$ A will be restated as Theorem 3.1.

³Theorem B will be restated as Theorem 4.7.

Let K be a field and let v be a valuation on K. We denote the valuation ring of v (i.e. the subring $\{a \in K \mid v(a) \geq 0\}$ of K) by \mathcal{O}_v and its valuation ideal (i.e. its maximal ideal $\{a \in \mathcal{O}_v \mid v(a) > 0\}$) by \mathcal{M}_v . Moreover, we denote the value group of v by vK and the residue field $\mathcal{O}_v/\mathcal{M}_v$ by Kv. An element $a + \mathcal{M}_v \in Kv$ with $a \in \mathcal{O}_v$ is also denoted by av. Suppose that K admits an ordering <. Then we say that v is **convex** (with respect to <) if \mathcal{O}_v is a convex subset of K (with respect to the ordering <). By [19, Lemma 2.1], any henselian valuation on an ordered field is convex. Given a valued field K we denote by $K^{\rm h}$ its henselisation.

Let $\mathcal{L}_{og} = \{+, -, 0, <\}$ be the **language of ordered groups**, $\mathcal{L}_{r} = \{+, -, \cdot, 0, 1\}$ the **language of rings** and $\mathcal{L}_{or} = \mathcal{L}_{r} \cup \{<\}$ the **language of ordered rings**. If no confusion is likely to arise, we denote an \mathcal{L}_{og} -structure (G, +, -, 0, <) simply by G, an \mathcal{L}_{r} -structure $(R, +, -, \cdot, 0, 1)$ by R and an \mathcal{L}_{or} -structure $(R, +, -, \cdot, 0, 1, <)$ by (R, <). We say that a valuation v on a field K (an ordered field (K, <)) is \mathcal{L}_{r} -definable (\mathcal{L}_{or} -definable) if \mathcal{O}_{v} is an \mathcal{L}_{r} -definable (\mathcal{L}_{or} -definable) subset of K. Here, definability is meant with parameters. We say that v is $0-\mathcal{L}_{r}$ -definable $(0-\mathcal{L}_{or}$ -definable) if there is a defining formula without parameters from K.

If we expand $\mathcal{L}_{\mathbf{r}}$ by a unary predicate \mathcal{O}_v , we obtain the **language of valued** fields $\mathcal{L}_{\mathrm{vf}}$. Similarly, $\mathcal{L}_{\mathrm{ovf}} = \mathcal{L}_{\mathrm{or}} \cup \{\mathcal{O}_v\}$ denotes the **language of ordered** valued fields. Let (K, \mathcal{O}_v) be a valued field. An atomic formula of the form $v(t_1) \geq v(t_2)$, where t_1 and t_2 are $\mathcal{L}_{\mathbf{r}}$ -terms, is equivalent to $t_1 = t_2 = 0 \lor (t_2 \neq 0 \land \frac{t_1}{t_2} \in \mathcal{O}_v)$. Hence, by abuse of notation, we can also write (K, v) for the $\mathcal{L}_{\mathrm{vf}}$ -structure of a valued field. Likewise the $\mathcal{L}_{\mathrm{ovf}}$ -structure of an ordered valued field is denoted by (K, <, v).

Let $(\Gamma, <)$ be a linearly ordered set, and for each $\gamma \in \Gamma$, let $A_{\gamma} \neq \{0\}$ be an abelian group (expressed additively). The corresponding **Hahn product** over $\{A_{\gamma} \mid \gamma \in \Gamma\}$ is given by

$$\underset{\gamma \in \Gamma}{\mathbf{H}} A_{\gamma} = \left\{ s \in \prod_{\gamma \in \Gamma} A_{\gamma} \mid \text{supp}(s) \text{ is well-ordered} \right\},$$

where $\prod_{\gamma \in \Gamma} A_{\gamma}$ denotes the group product over $(A_{\gamma})_{\gamma \in \Gamma}$ and $\operatorname{supp}(s)$ denotes the **support** of *s* given by the set $\{\gamma \in \Gamma \mid s(\gamma) \neq 0\}$. The Hahn product is an abelian group under pointwise addition. It carries a valuation with value set Γ given by $v_{\min} : s \mapsto \min \operatorname{supp}(s)$ for $s \neq 0$. Any $s \in \mathbf{H}_{\gamma \in \Gamma} A_{\gamma}$ can be expressed as a sum $s = \sum_{\gamma \in \Gamma} s_{\gamma} \mathbb{1}_{\gamma}$, where $s_{\gamma} = s(\gamma)$ and $\mathbb{1}_{\gamma}$ is the characteristic function mapping γ to some fixed $\mathbb{1}_{\gamma} \in A_{\gamma}$ (which in our application is always clear from the context) and everything else to 0. The valued subgroup

$$\coprod_{\gamma \in \Gamma} A_{\gamma} = \{ s \in \mathbf{H}_{\gamma \in \Gamma} A_{\gamma} \mid \operatorname{supp}(s) \text{ is finite} \}$$

is called the **Hahn sum** over $(\Gamma, (A_{\gamma})_{\gamma \in \Gamma})$. If Γ is a convex subset of \mathbb{Z} with $\min \Gamma = a$ and $\max \Gamma = b$ (where we set $a = -\infty$ if Γ is not bounded from below, and $b = \infty$ if Γ is not bounded from above), then we express $\coprod_{\gamma \in \Gamma} A_{\gamma}$ as $\coprod_{\gamma = a}^{b} A_{\gamma}$ if $a \neq -\infty$, and as $\coprod_{\gamma > -\infty}^{b} A_{\gamma}$ if $a = -\infty$. For an ordinal α we have that $\coprod_{\gamma \in -\alpha} A_{\gamma} = \mathbf{H}_{\gamma \in -\alpha} A_{\gamma}$. If there is an ordered abelian group A such that for each $\gamma \in \Gamma$ we have $A_{\gamma} = A$, we simply write $\mathbf{H}_{\Gamma}A$ ($\coprod_{\Gamma} A$) for the Hahn

product (Hahn sum) above. If A_{γ} is an ordered abelian group for each $\gamma \in \Gamma$, then also $\mathbf{H}_{\gamma \in \Gamma} A_{\gamma}$ is an ordered abelian group with the ordering given as follows: for any non-zero $s = \sum_{\gamma \in \Gamma} s_{\gamma} \mathbb{1}_{\gamma} \in \mathbf{H}_{\gamma \in \Gamma} A_{\gamma}$ we have s > 0 if and only if $s_{\gamma_0} > 0$, where $\gamma_0 = v_{\min}(s)$.

Let k be a field and let G be an ordered abelian group. The **Hahn field** (also called **power series field**) k((G)) has as underlying additive group the Hahn product $\mathbf{H}_G k$. Multiplication on k((G)) is defined by $(rs)_h = \sum_{g \in G} r_g s_{h-g}$ for any $h \in G$. To distinguish between elements of Hahn products and Hahn fields, we write an element $s \in k((G))$ as sum $s = \sum_{g \in G} s_g t^g$, where t^g is the characteristic function with $t^g(g) = 1$ and $t^g(h) = 0$ for any $h \neq g$. The valuation v_{\min} on k((G)) is henselian. Again, if k is ordered, then also k((G)) carries a natural ordering as described above making it an ordered field. We denote by k[G] the **group ring** with coefficient field k and value group G. It consists of all elements of k((G)) with finite support. Its quotient field, denoted by k(G), is generated in k((G)) by the set of all monic monomials, i.e. $k(G) = k(t^g | g \in G)$.

Let G be an ordered abelian group. We say that G is **discretely ordered** if there exists some $1 \in G$ with 0 < 1 and there is no element in G strictly between 0 and 1. If G is not discretely ordered, then it is **densely ordered**. Let $n \in \mathbb{N}$. Then G is called *n*-regular if every infinite convex subset of G contains at least one *n*-divisible element. If G is *n*-regular for every $n \in \mathbb{N}$, then we say that it is regular. We denote the **divisible hull** of G by G^{div} .

§3. Definable valuations. We start by stating and proving the main result of this section. Subsequently, we will discuss how this generalises the known definability results from Fact 1.2 and Fact 1.3.

THEOREM 3.1. Let (K, v) be a henselian valued field. Suppose that vK is not closed in vK^{div} . Then v is \mathcal{L}_r -definable with one parameter.

Theorem 3.1 can essentially be proven by the methods in [14, page 15] and [13, page 19].⁴ However, due to the more general context there are certain adjustments to be made, which is why we present the proof in detail.

PROOF. We set G = vK. Let $\varepsilon \in K$ and let $n \in \mathbb{N}$ such that $v(\varepsilon)$ is not *n*-divisible in G but $\frac{v(\varepsilon)}{n}$ is a limit point of G in G^{div} . Since $\frac{v(\varepsilon)}{n}$ is a limit point of G, also $-\frac{v(\varepsilon)}{n}$ is a limit point of G. Thus, by replacing ε by ε^{-1} if necessary, we may assume that $\frac{v(\varepsilon)}{n}$ is a left-sided limit point of G, i.e. for any positive $h \in G^{\text{div}}$ the half-open interval $\left(\frac{v(\varepsilon)}{n} - h, \frac{v(\varepsilon)}{n}\right]$ in G^{div} contains some element from G.

Set $\Phi_{\varepsilon} = \{x \in K \mid v(\varepsilon x^n) > 0\}$. We show that the formula $\varphi(x, \varepsilon)$ given by

$$\exists y \ (y^n - y^{n-1} = \varepsilon x^n)$$

defines Φ_{ε} . Let $x \in \Phi_{\varepsilon}$. Then $\varepsilon x^n \in \mathcal{M}_v$ and by Hensel's Lemma there exists $y \in K$ such that $y^n - y^{n-1} = \varepsilon x^n$, i.e. $K \models \varphi(x, \varepsilon)$. Conversely, let $x \in K$ such that $K \models \varphi(x, \varepsilon)$, i.e. there exists $y \in K$ such that $y^{n-1}(y-1) = \varepsilon x^n$. By

⁴We thank Blaise Boissonneau for pointing out this method to the first author.

applying v, we obtain

$$(n-1)v(y) + v(y-1) = v(\varepsilon) + nv(x)$$

and distinguish three cases.

Case 1: v(y) > 0. Then v(y-1) = 0 and thus $0 < (n-1)v(y) = v(\varepsilon) + nv(x) = v(\varepsilon x^n)$, as required.

Case 2: v(y) = 0. Then $v(y-1) = v(\varepsilon) + nv(x)$. If v(y-1) > 0, then we are done immediately. Otherwise, v(y-1) = 0 and we obtain $\frac{v(\varepsilon)}{n} = -v(x)$, a contradiction to the fact that $v(\varepsilon)$ is not n-divisible in G.

Case 3: v(y) < 0. Then we obtain $v(\varepsilon) = n(v(y) - v(x))$, again a contradiction, as $v(\varepsilon)$ is not *n*-divisible.

Now set

$$\Psi_{\varepsilon} = \{ \varepsilon x^n \mid x \in \Phi_{\varepsilon} \} = \{ \varepsilon x^n \mid x \in K, v(\varepsilon x^n) > 0 \}.$$

Again, Ψ_{ε} is \mathcal{L}_{r} -definable only using the parameter ε . Finally, set

$$\Omega_{\varepsilon} = \left\{ x^n - x^{n-1} \mid x \in K, K \models \exists y \; \exists (z \in \Psi_{\varepsilon}) \; z(y^n - y^{n-1}) = x^n - x^{n-1} \right\}.$$

We show that $\Omega_{\varepsilon} = \mathcal{M}_{v}$, as then \mathcal{O}_{v} is defined by $\forall (u \in \Omega_{\varepsilon}) xu \in \Omega_{\varepsilon}$.

First let $a \in \Omega_{\varepsilon}$ and fix $x \in K$ such that $x^n - x^{n-1} = a$. Then by definition of Ω_{ε} there are $y \in K$ and $z \in \Psi_{\varepsilon}$ such that $z(y^n - y^{n-1}) = a$. Let $r \in K$ with $z = \varepsilon r^n \in \mathcal{M}_v$. Assume, for a contradiction, that $v(a) \leq 0$. If v(a) < 0, then

$$0 > v(x^{n-1}(x-1)) = (n-1)v(x) + v(x-1) = nv(x)$$

and 0 > v(z) + (n-1)v(y) + v(y-1) > (n-1)v(y) + v(y-1) = nv(y).

Since

$$\varepsilon = zr^{-n} = (x^n - x^{n-1})[(y^n - y^{n-1})r^n]^{-1},$$

we obtain

$$v(\varepsilon) = v(x^n - x^{n-1}) - v(y^n - y^{n-1}) - nv(r) = n(v(x) - v(y) - v(r)),$$

contradicting the choice of ε . If v(a) = 0, then 0 = (n-1)v(x) + v(x-1) and thus v(x) = v(x-1) = 0. Moreover, 0 = v(z) + (n-1)v(y) + v(y-1) and thus 0 > -v(z) = (n-1)v(y) + v(y-1) = nv(y). This implies

$$v(\varepsilon) = v(x^n - x^{n-1}) - v(y^n - y^{n-1}) - nv(r) = n(v(y) - v(r)),$$

which again contradicts the fact that $v(\varepsilon)$ is not *n*-divisible. This shows that v(a) > 0, i.e. $a \in \mathcal{M}_v$.

For the converse, let $a \in \mathcal{M}_v$. By Hensel's Lemma there exists $x \in K$ such that $x^n - x^{n-1} = a$. Since $\frac{v(\varepsilon)}{n}$ is a left-sided limit point of G, we obtain some $b \in K^{\times}$ with

$$\frac{v(\varepsilon)}{n} - \frac{v(a)}{n} < v(b) < \frac{v(\varepsilon)}{n}.$$

Hence, $-v(\varepsilon) < -nv(b) < v(a) - v(\varepsilon)$. We set $z = \varepsilon b^{-n}$ and obtain that 0 < v(z) < v(a), whence $z \in \Psi_{\varepsilon}$ and $v(\frac{a}{z}) > 0$. Again by applying Hensel's Lemma, we find some $y \in K$ such that $y^n - y^{n-1} = \frac{a}{z}$. This yields $a \in \Omega_{\varepsilon}$ and therefore $\Omega_{\varepsilon} = \mathcal{M}_v$, as required.

REMARK 3.2. Note that the class of non-trivial ordered abelian groups that are closed in their divisible hull is elementary in the language of ordered groups. Indeed, it is axiomatised by the axiom scheme

$$\forall a \ (\forall \ (b > 0) \ \exists g \ |a - ng| < b \to \exists h \ a = nh),$$

 \dashv

 \dashv

one for each $n \in \mathbb{N}$.

Our next aim is to show that Theorem 3.1 implies Fact 1.2. Note that in both of these statements the value group is densely ordered; in Fact 1.2 by assumption and in Theorem 3.1 since any discretely ordered abelian group is already closed in its divisible hull.

PROPOSITION 3.3. Let G be a densely ordered abelian group. Suppose that there exists some $n \in \mathbb{N}$ such that G contains a convex subgroup C that is n-regular but not n-divisible. Then G is not closed in G^{div} .

PROOF. Note that since G contains a convex subgroup that is not n-divisible, we have that G is a non-divisible ordered abelian group. Assume, for a contradiction, that G is closed in G^{div} . Since C is not n-divisible, there exists $a \in C$ such that $\frac{a}{n} \in G^{\text{div}} \setminus G$. Moreover, since G is closed in G^{div} , there exists a positive $\delta \in G^{\text{div}}$ such that

$$\left(\frac{a}{n}-\delta,\frac{a}{n}+\delta\right)\cap G=\emptyset.$$

As 0 is a limit point of G in G^{div} (cf. [21, Lemma 3.6]), there exists $b \in G$ with $0 < b < \delta$ and hence

(1)
$$\left(\frac{a}{n} - \frac{b}{n}, \frac{a}{n} + \frac{b}{n}\right) \cap G = \emptyset.$$

By convexity of C, we may choose $b \in C^{>0}$. Further, since G is densely ordered and thus C is densely ordered, the interval (a-b, a+b) is an infinite convex subset of C. By *n*-regularity of C, there is some $z \in C$ such that $a - b \leq nz \leq a + b$. This is equivalent to

$$\frac{a-b}{n} \le z \le \frac{a+b}{n},$$

a contradiction to (1). Hence, G is not closed in G^{div} .

Proposition 3.3 shows that the condition on the value group in Fact 1.2 is covered by the condition in Theorem 3.1. The converse of Proposition 3.3 does not hold, as verified by the following example.

EXAMPLE 3.4. We consider the ordered abelian group

$$G = \left(\prod_{\omega} \mathbb{Z}\right) + a\mathbb{Z} \subseteq \underset{\omega}{\mathbf{H}} \mathbb{Z}, \text{ where } a = \sum_{n \in \omega} 2\mathbb{1}_n.$$

Note that G consists of all elements $s = \sum_{n \in \omega} s_n \mathbb{1}_n \in \mathbf{H}_{\omega} \mathbb{Z}$ for which the following holds: there exist $m \in \mathbb{Z}$ and $n_0 \in \omega$ such that for any $n \geq n_0$ we have $s_n = 2m$. We now show that G is not closed in G^{div} but for any $n \in \mathbb{N}$ and any convex subgroup C of G, if C is n-regular, then it is already n-divisible.

By [21, Lemma 3.6], we have that G is densely ordered, as its value set $v_{\min}G = \omega$ does not have a maximum. We first note that the divisible hull of G is given by

$$G^{\operatorname{div}} = \left(\coprod_{\omega} \mathbb{Q}\right) + a\mathbb{Q}.$$

Let $s = \sum_{n \in \omega} \mathbb{1}_n \in G^{\text{div}}$. By the description of the elements of G above, one immediately sees that $s \notin G$. We now show that s is a limit point of G in G^{div} , whence G is not closed in G^{div} . Let $b \in G^{\text{div}}$ be positive and set $\ell = v_{\min}(b)$. Further, let $g = \sum_{n=0}^{\ell} \mathbb{1}_n \in G$ and note that g < s. As $b_{\ell} > 0$, we obtain that $0 < s - g = 0 \cdot \mathbb{1}_{\ell} + \sum_{n \geq \ell+1} \mathbb{1}_n < b$. We have thus found an element $g \in G$ with s - b < g < s, showing that s is a limit point of G in G^{div} .

Let $n \in \mathbb{N}$ with $n \geq 2$. We show that G contains no non-trivial *n*-regular convex subgroup. Thus, in particular, the only *n*-regular convex subgroup of G is the trivial group, which is *n*-divisible.

Let C be a non-trivial convex subgroup of G. Then there exists $\ell \in \omega$ such that C is of the form

$$C = \{g \in G \mid v_{\min}(g) \ge \ell\} = \left(\prod_{\gamma=\ell}^{\infty} \mathbb{Z}\right) + \left\{\sum_{i \ge \ell} k \mathbb{1}_i \mid k \in 2\mathbb{Z}\right\}$$

(cf. [23, page 50 f.]). Now let $c, d \in C$ with $c = q \mathbb{1}_{\ell}$ and $d = q \mathbb{1}_{\ell} + \mathbb{1}_{\ell+1}$, where $q \in \mathbb{N}$ is some prime with q > n. Then c < d and (c, d) is an infinite convex subset of the densely ordered group C. Since for any $z \in (c, d)$ we have that $z_{\ell} = q$ and $n \nmid q$ in \mathbb{Z} , there is no *n*-divisible element in (c, d). Hence, C is not *n*-regular.

Example 3.4 shows that the condition in Theorem 3.1 is indeed more general than the condition in Fact 1.2. More precisely, let k be any field and let G be the ordered abelian group constructed in Example 3.4; then Theorem 3.1 implies that the henselian valuation v_{\min} on k((G)) is \mathcal{L}_r -definable (with one parameter), whereas Fact 1.2 cannot be applied directly to obtain the \mathcal{L}_r -definability of v_{\min} .

In conclusion, Theorem 3.1 is a strict generalisation of both Fact 1.2 and Fact 1.3 for densely ordered value groups: Theorem 3.1 clearly generalises Fact 1.3, as it has the same condition on the value group but does not rely on the existence of an ordering on the field. Moreover, Proposition 3.3 shows that the value group condition of Theorem 3.1 implies that of Fact 1.2 but Example 3.4 illustrates that the converse does not hold.

§4. Main construction and parameter-free definability. The main aim of this section is to show that the conclusions of Fact 1.1 and Theorem 3.1 can, in general, not be strengthened to $0-\mathcal{L}_{\rm r}$ -definability. More precisely, we construct a class of ordered henselian valued fields that do not admit any non-trivial 0- $\mathcal{L}_{\rm or}$ -definable henselian valuation (see Theorem 4.7). This class contains ordered henselian valued fields (K, <, v) with discretely ordered value group vK (see Example 4.9) as well as such with a value group vK not closed in $vK^{\rm div}$ (see Example 4.10).

The method for constructing an *ordered* henselian valued field (K, <, v) that does not admit a non-trivial 0- \mathcal{L}_{or} -definable henselian valuation relies on the construction of a *t*-henselian non-henselian *ordered* field.⁵ We start by clarifying the notion of *t*-henselianity, which was first introduced in [25, Section 7].⁶

DEFINITION 4.1. We call a field **henselian** if it admits a non-trivial henselian valuation. A field K is *t*-henselian if there exists a henselian field L such that $K \equiv L$ (in the language \mathcal{L}_r).

We now establish a sufficient condition for t-henselianity based on n_{\leq} -henselian valuations: Let $n \in \mathbb{N}$. Following the terminology of [8, Definition 6.1], we say that a valued field (K, v) is n_{\leq} -henselian if Hensel's Lemma holds in (K, v)for all polynomials of degree at most n. Note that (K, v) is henselian if and only if it is n_{\leq} -henselian for any $n \in \mathbb{N}$. Arguing similarly as in the proof of [6, Theorem 4.1.3], one can show that (K, v) is n_{\leq} -henselian if and only if every polynomial $X^{\ell} + X^{\ell-1} + a_{\ell-2}X^{\ell-2} + \ldots + a_0 \in \mathcal{O}_v[X]$ with $\ell \leq n$ and $a_{\ell-2}, \ldots, a_0 \in \mathcal{M}_v$ has a zero in K. Moreover, by the arguments in the proof of [19, Lemma 2.1], for $n \geq 2$ any n_{\leq} -henselian valuation on a real field K is convex with respect to any ordering on K.

LEMMA 4.2. Let K be a field. Suppose that for any $n \in \mathbb{N}$, there exists a non-trivial n_{\leq} -henselian valuation on K. Then K is t-henselian.

PROOF. Let h_n be the \mathcal{L}_{vf} -sentence stating that v is a non-trivial valuation for which Hensel's Lemma holds for polynomials of degree at most n. Then any finite subset of the \mathcal{L}_{vf} -theory $\operatorname{Th}(K) \cup \{h_n \mid n \in \mathbb{N}\}$ is satisfied by (K, v_m) for some $m \in \mathbb{N}$, where v_m is m_{\leq} -henselian. Hence, by the Compactness Theorem, there exists a henselian field L with $L \models \operatorname{Th}(K)$, i.e. $L \equiv K$.

Our aim is to refine the construction of a *t*-henselian non-henselian field in [25, page 338] by gaining influence on the resulting value group. More precisely, starting with a family of non-trivial ordered abelian groups $\{A_{\gamma} \mid \gamma \in -\omega\}$, we construct a non-henselian ordered field (K, <) such that for any $n \in \mathbb{N}$, there is an n_{\leq} -henselian valuation v_n on K with $v_n K = \coprod_{\gamma > -\infty}^{-n} A_{\gamma}$.

In the following, for an ordered abelian group G we consider G^{div} also as a \mathbb{Q} -vector space with the usual scalar multiplication. For an algebraic field extension $K \subseteq L$ and an element $\alpha \in L$, we denote by $\deg_K(\alpha)$ the degree of the minimal polynomial of α over K.

Lemma 4.3 and Lemma 4.4 below will be used in Construction 4.5. Note that the statement of Lemma 4.4 can be adjusted for an arbitrary field k of characteristic 0 (instead of an ordered field).

LEMMA 4.3. Let k be a field of characteristic 0 and let G be an ordered abelian group. Moreover, let $a \in k[G]$ such that for some $n \ge 2$ we have $\operatorname{supp}(a) = \{g_1, \ldots, g_n, 0\}$ and g_1, \ldots, g_n are linearly independent in the Q-vector space G^{div} . Then a is prime in k[G].

 $^{{}^{5}}$ Our construction is inspired by the arguments in [25, page 338], which have also been refined in [8, Section 6], [16, Section 6] and [1, Section 4].

 $^{^{6}}$ Several characterisations for a field to be t-henselian are given in [8, page 354].

PROOF. First note that the group of units of k[G] is given by the multiplicative group of monomials of k((G)), i.e. $k[G]^{\times} = \{ct^g \mid c \in k^{\times}, g \in G\}$. By [3, Proposition 4.1], every irreducible element of k[G] is prime. It thus suffices to show that a is irreducible. Let $r, s \in k[G]$ with a = rs. We obtain from [3, Proposition 4.8] that a is prime in $k[G^{\text{div}}]$ and hence irreducible in $k[G^{\text{div}}]$ (as $k[G^{\text{div}}]$ is an integral domain). We may thus assume that r is a unit in $k[G^{\text{div}}]$. Hence, $r \in k[G] \cap \{ct^g \mid c \in k^{\times}, g \in G^{\text{div}}\} = k[G]^{\times}$. This shows that r is a unit in k[G], as required.

For two ordered abelian groups A and B, we denote by A II B the abelian group $A \times B$ ordered lexicographically.

LEMMA 4.4. Let (k, <) be an ordered field and let $A_1, A_2 \neq \{0\}$ be ordered abelian groups. Moreover, set $K = k(A_1 \amalg A_2)$ and let $(K^h, <)$ be a henselisation of K in $(k((A_1 \amalg A_2)), <, v_{\min})$. Then for any $n \in \mathbb{N}$, there exists an intermediate field extension $K \subseteq L \subsetneq K^h$ such that (L, v_{\min}) is n_{\leq} -henselian but not henselian.

PROOF. Let $n \in \mathbb{N}$ and let $p \in \mathbb{N}$ be prime with p > n. Since p_{\leq} -henselianity implies n_{\leq} -henselianity, it suffices to construct an ordered valued field $(L, <, v_{\min})$ with $(K, <, v_{\min}) \subseteq (L, <, v_{\min}) \subsetneq (K^{\mathrm{h}}, <, v_{\min})$ such that for any polynomial of the form

$$f(T) = T^m + a_{m-1}T^{m-1} + \ldots + a_0 \in L[T],$$

where $m \leq p$, and any $\alpha \in K^{\rm h}$ with $f(\alpha) = 0$, we have that $\alpha \in L$. Let M be the set of all natural numbers m such that for any prime number $q \in \mathbb{N}$ dividing m we have $q \leq p$. We set $L = K(\alpha \in K^{\rm h} | \deg_K(\alpha) \in M)$, i.e. L is generated by all elements in $K^{\rm h}$ whose degree over K only has prime factors less than or equal to p. Note that since M is closed under multiplication and divisors, $\deg_K(x) \in M$ for any $x \in L$. Let $m \leq p$ and let $f(T) = T^m + a_{m-1}T^{m-1} + \ldots + a_0 \in L[T]$. By construction of L, we have $\deg_K(a_i) \in M$ for any $i \in \{0, \ldots, m-1\}$. Moreover, since $[K(a_0, \ldots, a_{m-1}) : K]$ divides the product $[K(a_0) : K] \ldots [K(a_{m-1}) : K] = \deg_K(a_0) \ldots \deg_K(a_{m-1})$, we obtain $[K(a_0, \ldots, a_{m-1}) : K] \in M$. Suppose that $\alpha \in K^{\rm h}$ is a zero of f. Since also $[K(a_0, \ldots, a_{m-1}, \alpha) : K(a_0, \ldots, a_{m-1})] \leq m \leq p$, we obtain

$$[K(a_0, \dots, a_{m-1}, \alpha) : K] = [K(a_0, \dots, a_{m-1})] \cdot [K(a_0, \dots, a_{m-1})] : K] \in M.$$

Finally, since $\deg_K(\alpha) = [K(\alpha) : K]$ divides $[K(a_0, \ldots, a_{m-1}, \alpha) : K]$, we obtain $\deg_K(\alpha) \in M$ and thus $\alpha \in L$.

It remains to show that (L, v_{\min}) is not henselian, i.e. that there exists $y \in K^{\mathrm{h}}$ such that $y \notin L$. Let $q \in \mathbb{N}$ be a prime number greater than p. We set

$$a = t^{(b_1,0)} + t^{(0,b_2)} + 1 \in k[A_1 \amalg A_2],$$

where $b_1 \in A_1$ and $b_2 \in A_2$ are fixed positive elements, and observe that a is prime in $k[A_1 \amalg A_2]$ by Lemma 4.3. Further, let

$$g(T) = T^{q} + aT^{q-1} + at^{(b_{1},0)}T^{q-2} + \dots + at^{(b_{1},0)}T + at^{(b_{1},0)} \in K[T].$$

Note that $v_{\min}(a) = 0$ and $v_{\min}(at^{(b_1,0)}) = (b_1,0) > 0$. Hence, by applying the residue map, we obtain the polynomial $gv_{\min}(T) = T^q + T^{q-1} \in k[T]$. Since $gv_{\min}(T)$ has a simple zero in k, we obtain that g has a zero $y \in K^{\mathrm{h}}$. Moreover, by Eisenstein's Criterion we obtain that g is irreducible over K and thus q = [K(y):K]. Since q does not divide any element in M, we obtain by construction of L that $y \notin L$. Hence, $L \subsetneq K^{\mathrm{h}}$ as required.

We now proceed with our main construction.

CONSTRUCTION 4.5 (Main Construction). Let $\{A_{\gamma} \mid \gamma \in \omega\}$ be a family of non-trivial ordered abelian groups and set $G := \coprod_{\gamma \in -\omega} A_{-\gamma}$. We construct an ordered field (K, <) such that for any $n \in \mathbb{N}$ there exists a convex n_{\leq} -henselian valuation u_n on K with value group $u_n K = \coprod_{\gamma > -\infty}^{-n} A_{-\gamma}$ and K does not admit a non-trivial henselian valuation. In particular, K is t-henselian (by Lemma 4.2) but not henselian.

By induction on $m \in \omega$ we construct a chain of ordered fields $(K_m, <)$ each endowed with a valuation v_{\min}^m . From this the ordered field (K, <) will be obtained by an inverse limit construction, as well as a chain of convex valuations v_m on K for $m \in \omega$. Ultimately, for each $n \in \omega$ we set u_{2n} to be v_n and u_{2n+1} to be a certain coarsening of u_{2n} .

To initiate the induction, we set $(K_0, <) = (\mathbb{R}, <)$. Now let $m \in \mathbb{N}$ and suppose that the ordered field $(K_{m-1}, <)$ has already been defined. We denote the valuation given by $s \mapsto \min \operatorname{supp}(s)$ on $K_{m-1}((A_{2m-1} \amalg A_{2(m-1)}))$ by v_{\min}^m . By Lemma 4.4, there is an ordered valued field K_m with

$$K_{m-1} (A_{2m-1} \amalg A_{2(m-1)}) \subseteq K_m$$

$$\subseteq K_{m-1} (A_{2m-1} \amalg A_{2(m-1)})^h \subseteq K_{m-1} ((A_{2m-1} \amalg A_{2(m-1)})),$$

where the ordering is inherited from $K_{m-1}((A_{2m-1} \amalg A_{2(m-1)}))$ and the valuation is v_{\min}^m , such that any zero in $K_{m-1}(A_{2m-1} \amalg A_{2(m-1)})^h$ of a polynomial over K_m of degree at most 2m+1 already lies in K_m . Since the extensions above are immediate, we have $K_m v_{\min}^m = K_{m-1}$. Let \mathcal{O}'_m and \mathcal{M}'_m be the valuation ring and valuation ideal of v_{\min}^m in K_m and let $\pi_m \colon \mathcal{O}'_m \to K_{m-1}$ denote the corresponding residue map $a \mapsto a v_{\min}^m$.

For $i, j \in \omega$ with $j \leq i$ we define the map $\psi_{i,j} \colon K_i \cup \{\infty\} \to K_j \cup \{\infty\}$ as follows: We set

$$\psi_{i,i} = \mathrm{id}_{K_i \cup \{\infty\}} \text{ and } \psi_{i,i-1}(x) = \begin{cases} \pi_i(x) & \text{if } x \in \mathcal{O}'_i, \\ \infty & \text{if } x \notin \mathcal{O}'_i. \end{cases}$$

For $0 \leq j < i-1$ we set $\psi_{i,j} = \psi_{j+1,j} \circ \cdots \circ \psi_{i,i-1}$. Now we define I to be the inverse limit of $(K_m \cup \{\infty\})_m$, i.e. I is the following set of sequences:

$$I = \left\{ (x_m) \in \prod_{m \in \omega} (K_m \cup \{\infty\}) \ \middle| \ \psi_{i,j}(x_i) = x_j \text{ for all } i, j \in \omega \text{ with } j \le i \right\}.$$

Note that if for a given sequence $(x_m) \in I$, we have $x_{\ell} = 0$ (respectively $x_{\ell} = \infty$) for some $\ell \in \omega$, then $x_k = 0$ (respectively $x_k = \infty$) for any $k \leq \ell$. On the other hand, if $x_{\ell} \notin \{0, \infty\}$ for some $\ell \in \omega$, then

(2)
$$x_k \in \mathcal{O}'_k \setminus \mathcal{M}'_k$$

i.e. $v_{\min}^k(x_k) = 0$, for all $k > \ell$. Therefore, if $x_\ell \notin \{0, \infty\}$ and $x_{\ell-1} \in \{0, \infty\}$ for some $\ell \in \mathbb{N}$, then $v_{\min}^\ell(x_\ell) \neq 0$.

CLAIM 1. The set $K := I \setminus \{(\infty)\}$ can be made an ordered field.

PROOF OF CLAIM 1. Addition, multiplication and an ordering on K can be defined by a standard procedure⁷ making K an ordered field: Let $(x_m), (y_m) \in K$ and let $\ell \in \omega$ be least such that $x_\ell \neq \infty$ and $y_\ell \neq \infty$. For any $i \geq \ell$ set $z_i = x_i + y_i$. For any $i < \ell$ set $z_i = \psi_{\ell,i}(x_\ell + y_\ell)$. Then $(x_m) + (y_m)$ is given by $(z_m) \in K$. Likewise, multiplication can be defined via the multiplication operations on the fields K_m . Finally, a non-zero element (x_m) of K is positive if and only if x_i is non-negative for any $i \in \omega$ with $x_i \neq \infty$.

For any $n \in \omega$ we define the projection map $\psi_n \colon K \to K_n \cup \{\infty\}, (x_m) \mapsto x_n$. This gives rise to the following commutative diagram and equation for $j \leq i$:

$$K \xrightarrow{\psi_i} K_i \cup \{\infty\}$$

$$\downarrow_{\psi_j} \qquad \qquad \qquad \downarrow_{\psi_{i,j}}$$

$$K_j \cup \{\infty\}$$

(3) $\psi_{i,j} \circ \psi_i = \psi_j$

One can easily verify that ψ_n is a place, and therefore $\mathcal{O}_n = \psi_n^{-1}(K_n)$ defines a valuation ring on K with maximal ideal $\mathcal{M}_n = \psi_n^{-1}(\{0\})$ and residue field K_n . By (3) we compute for any $a \in \mathcal{M}_{n+1}$:

$$\psi_n(a) = \psi_{n+1,n}(\psi_{n+1}(a)) = \psi_{n+1,n}(0) = 0.$$

Therefore we have $a \in \mathcal{M}_n$, and thus $\mathcal{O}_n \subseteq \mathcal{O}_{n+1}$.

We denote the valuation on K with valuation ring \mathcal{O}_n by v_n and fix $n \in \mathbb{N}$.

CLAIM 2. v_n is a $(2n+1) \leq$ -henselian valuation on K.

We observe that, if Claim 2 is established, then we also have that v_n is $(2n)_{\leq}$ -henselian and convex (as $2n + 1 \geq 2$).

PROOF OF CLAIM 2. Let

$$f(T) = T^{2n+1} + T^{2n} + a^{(2n-1)}T^{2n-1} + \dots + a^{(0)} \in K[T],$$

where $a^{(2n-1)}, \ldots, a^{(0)} \in \mathcal{M}_n$. We need to find a zero $(y_m) \in K$ of f. For any $i \in \omega$, set $a_i^{(2n-1)} = \psi_i(a^{(2n-1)}), \ldots, a_i^{(0)} = \psi_i(a^{(0)}) \in K_i$ and

$$f_i(T) = T^{2n+1} + T^{2n} + a_i^{(2n-1)}T^{2n-1} + \dots + a_i^{(0)} \in K_i[T].$$

First let $i \leq n$. Then $a_i^{(2n-2)} = \cdots = a_i^{(0)} = 0$ and therefore $f_i(T) = T^{2n+1} + T^{2n}$. Set $y_i = -1$, which is a simple zero of f_i . Now let i > n. We construct y_i iteratively. To this end, assume that $y_{i-1} \in K_{i-1}$ is a simple zero of f_{i-1} .

⁷See [6, page 114 ff.] for analogous construction in the group case.

with $\psi_{i-1,j}(y_{i-1}) = y_j$ for $0 \le j \le i-1$. Applying π_i to the coefficients of $f_i(T) \in K_i[T]$ we obtain the residue polynomial

$$T^{2n+1} + T^{2n} + a_{i-1}^{(2n-1)}T^{2n-1} + \dots + a_{i-1}^{(0)} = f_{i-1}(T) \in K_{i-1}[T].$$

Thus, f_i has a simple zero y_i in the henselisation of K_i with $\pi_i(y_i) = y_{i-1}$. By construction of K_i (as i > n), we obtain $y_i \in K_i$, as required. Finally, one can verify by direct computation that the element $(y_m) \in K$ we obtained from this construction is a zero of f.

The argument above shows that v_n satisfies the condition for $(2n + 1) \leq$ -henselianity only for polynomials of degree equal to 2n + 1. However, since the coarsening of a $(2n+1)\leq$ -henselian valuation is also $(2n+1)\leq$ -henselian (see the proof of [20, Lemma A.4.24]), we immediately obtain $(2n + 1)\leq$ -henselianity of v_n .

It remains to show that v_n is not henselian.

CLAIM 3. v_n is not henselian.

PROOF OF CLAIM 3. Assume for a contradiction that v_n is henselian. Let $m \in \mathbb{N}$ and let

$$g(X) = X^m + X^{m-1} + a_{m-2}X^{m-2} + \dots + a_0 \in K_{n+1}[X]$$

for some $a_{m-2}, \ldots, a_0 \in \mathcal{M}'_{n+1}$. Let $a^{(m-2)}, \ldots, a^{(0)} \in K$ such that $\psi_{n+1}(a^{(i)}) = a_i$ for $0 \leq i \leq m-2$. Then $a^{(m-2)}, \ldots, a^{(0)} \in \psi_n^{-1}(\{0\}) = \mathcal{M}_n$. As v_n is henselian, there exists $x \in \mathcal{O}_n$ such that $x^m + x^{m-1} + a^{(m-2)}x^{m-2} + \cdots + a^{(0)} = 0$. A direct computation shows that $\psi_{n+1}(x)$ is a zero of the polynomial g(X). Thus, we have that $(K_{n+1}, v_{\min}^{n+1})$ is henselian, which is a contradiction to the construction of K_{n+1} .

So far we have seen that for any $n \in \mathbb{N}$ there exists a non-henselian but $(2n + 1) \leq$ -henselian valuation v_n on K. It remains to show that K does not admit a non-trivial henselian valuation.

CLAIM 4. K does not admit a non-trivial henselian valuation.

PROOF OF CLAIM 4. Let u be any non-trivial convex valuation on K and pick $a = (a_m) \in K$ with u(a) < 0, i.e. $a \notin \mathcal{O}_u$. Let $\ell \in \mathbb{N}$ be least such that $a_\ell \neq \infty$. Then $\psi_\ell(a) = a_\ell \in K_\ell$, whence $a \in \mathcal{O}_\ell$. Since v_ℓ is convex, we obtain that v_ℓ is a proper coarsening of u. Therefore, since v_ℓ is not henselian, also u is not henselian. Finally, as any henselian valuation on an ordered field is convex (cf. [19, Lemma 2.1]), this suffices to show that K does not admit a non-trivial henselian valuation. \diamond

Finally, we show inductively that $v_n K = \coprod_{\gamma > -\infty}^{-2n} A_{-\gamma}$. Since v_n is a coarsening of v_0 for all $n \in \mathbb{N}$, we have that $v_n K$ is the quotient of $v_0 K$ by $v_0(\mathcal{O}_n^{\times})$. Therefore our next task is to compute $v_0 K$ and $v_0(\mathcal{O}_n^{\times})$. In order to do so, let us first recall the construction of the fields K_m .

We have the inclusions as ordered fields

$$\mathbb{R}(A_1 \amalg A_0) \subseteq K_1 \subseteq \mathbb{R}((A_1 \amalg A_0)) \subseteq \mathbb{R}((G)),$$

12

where the last inclusion is obtained by embedding $A_1 \amalg A_0$ into G as its convex subgroup $\coprod_{\gamma=-1}^{0} A_{-\gamma}$. In the next step of our iterative field construction we obtained the ordered field inclusions

 $\mathbb{R}(A_1 \amalg A_0)(A_3 \amalg A_2) \subseteq K_2 \subseteq \mathbb{R}((A_1 \amalg A_0))((A_3 \amalg A_2)) \subseteq \mathbb{R}((G)).$

The last inclusion is given via the isomorphism⁸

 $\mathbb{R}((A_1 \amalg A_0))((A_3 \amalg A_2)) \cong \mathbb{R}(((A_3 \amalg A_2) \amalg (A_1 \amalg A_0)))$

and the embedding of $(A_3 \amalg A_2) \amalg (A_1 \amalg A_0)$ into G as its convex subgroup $\coprod_{\gamma=-3}^{0} A_{-\gamma}$. Continuing this process, we have that

$$\mathbb{R}\left(\prod_{\gamma=-(2m-1)}^{0} A_{-\gamma}\right) \subseteq K_m \subseteq \mathbb{R}\left(\left(\prod_{\gamma=-(2m-1)}^{0} A_{-\gamma}\right)\right) \subseteq \mathbb{R}((G)).$$

We denote by v_{\min} the standard valuation $s \mapsto \min \operatorname{supp}(s)$ on $\mathbb{R}((G))$. By the inclusion described above, v_{\min} restricted to K_m defines a valuation with value group $v_{\min}K_m = \coprod_{\gamma=-(2m-1)}^0 A_{-\gamma}$. For any $m \in \omega$ we have that $v_{\min}K_m$ can be regarded as a convex subgroup of G. Note that the valuation v_{\min} restricted to K_m is a refinement of v_{\min}^m : Both v_{\min} and v_{\min}^m are convex, $v_{\min}K_m = \coprod_{\gamma=-(2m-1)}^0 A_{-\gamma}$ and $v_{\min}^m K_m = (A_{2m-1} \amalg A_{2(m-1)})$. Thus, v_{\min}^m is the coarsening of v_{\min} induced by the convex subgroup $\coprod_{\gamma=-(2m-3)}^0 A_{-\gamma}$ of $v_{\min}K_m$.

We now want to define a map v on K such that v is a valuation on K and $\mathcal{O}_v = \mathcal{O}_0$. We set $v(0) = \infty$.

CLAIM 5. For any element $(x_m) \in K^{\times}$ and for any $x_i, x_j \notin \{0, \infty\}$ we have that $v_{\min}(x_i) = v_{\min}(x_j)$.

PROOF OF CLAIM 5. Let $\ell \in \omega$ be maximal with $x_{\ell-1} \in \{0, \infty\}$, if such ℓ exists, and $\ell = 0$ otherwise. It suffices to show that $v_{\min}(x_k) = v_{\min}(x_\ell)$ for any $k \geq \ell$. We proceed by induction on k. For $k = \ell$ there is nothing to prove. Now assume that $v_{\min}(x_\ell) = v_{\min}(x_k)$ for some fixed $k \geq \ell$. We first express $x_k \in K_k$ as

$$x_k = \sum_{g \in G} c_g t^g \in \mathbb{R}((G)).$$

Now set $g_0 = v_{\min}(x_k)$. Then $v_{\min}(x_k) = v_{\min}(x_\ell) \in \coprod_{\gamma=-(2\ell-1)}^0 A_{-\gamma} \subseteq G$ if $\ell \ge 1$, and $g_0 = v_{\min}(x_0) = 0$ if $\ell = 0$. Further we set

$$g_0 = \sum_{\gamma = -(2\ell - 1)}^0 h_{\gamma} \mathbb{1}_{\gamma},$$

where $h_{\gamma} \in A_{-\gamma}$ for each γ . By (2) we have that $v_{\min}^{k+1}(x_{k+1}) = 0$ and by construction $x_{k+1}v_{\min}^{k+1} = x_k$. Therefore we can set

$$x_{k+1} = c_{g_0} t^{g_0} + \sum_{g > g_0} c_g t^g + \varepsilon,$$

⁸Generally, for any ordered abelian groups C and D and any ordered field (k, <) we have $(k((C))((D)), <) \cong (k((D \amalg C)), <)$ via the isomorphism of ordered fields induced by $at^{c}t^{d} \mapsto at^{(d,c)}$ for any $a \in k$ and $(c,d) \in C \times D$.

for some $\varepsilon \in \mathcal{M}'_{k+1}$. Considering ε as an element of $\mathbb{R}((G))$ we can write

ε

$$\varepsilon = \sum_{g \in \coprod_{\gamma=-(2k+1)}^{0} A_{-\gamma}} \varepsilon_g t^g.$$

Let $\delta = v_{\min}(\varepsilon)$ and write

$$\delta = \delta_{-(2k+1)} \mathbb{1}_{-(2k+1)} + \delta_{-2k} \mathbb{1}_{-2k} + \sum_{\gamma = -(2k-1)}^{-2\ell} \delta_{\gamma} \mathbb{1}_{\gamma} + \sum_{\gamma = -(2\ell-1)}^{0} \delta_{\gamma} \mathbb{1}_{\gamma}.$$

Now since $\varepsilon \in \mathcal{M}'_{k+1}$, we have that $\delta_{-(2k+1)}\mathbb{1}_{-(2k+1)} + \delta_{-2k}\mathbb{1}_{-2k} > 0$. Hence,

$$\delta - g_0 = \underbrace{\delta_{-(2k+1)} \mathbb{1}_{-(2k+1)} + \delta_{-2k} \mathbb{1}_{-2k}}_{>0} + \sum_{\gamma = -(2k-1)}^{-2\ell} \delta_{\gamma} \mathbb{1}_{\gamma} + \sum_{\gamma = -(2\ell-1)}^{0} (\delta_{\gamma} - h_{\gamma}) \mathbb{1}_{\gamma}$$

and therefore $\delta > g_0$. Thus, $v_{\min}(x_{k+1}) = v_{\min}(x_k + \varepsilon) = v_{\min}(x_k) = v_{\min}(x_\ell)$, as required.

Let $(x_m) \in K^{\times}$. We define

$$v((x_m)) := v_{\min}(x_i) \in G$$

for any $x_i \notin \{0, \infty\}$.

CLAIM 6. The map v defines a valuation on K with vK = G.

PROOF OF CLAIM 6. Note that by Claim 5, v is well-defined. Let $(x_m), (y_m) \in K^{\times}$ and let i be large enough such that $x_i, y_i \notin \{0, \infty\}$. We obtain

$$v((x_m)(y_m)) = v_{\min}(x_i y_i) = v_{\min}(x_i) + v_{\min}(y_i) = v((x_m)) + v((y_m)).$$

Similarly, we prove that $v((x_m) + (y_m)) \ge \min\{v((x_m)), v((y_m))\}.$

Moreover, vK = G: For any $g \in G$ we can choose the least ℓ such that there is an element $a_{\ell} \in K_{\ell}$ with $v_{\min}(a_{\ell}) = g$. Now for any $(x_m) \in K$ with $x_{\ell} = a_{\ell}$ a direct calculation shows $v((x_m)) = v_{\min}(a_{\ell}) = g$.

We now show that $\mathcal{O}_v = \mathcal{O}_0$, this will yield that $v_0 K$ is (isomorphic to) G, as required.

CLAIM 7. $\mathcal{O}_v = \mathcal{O}_0$.

PROOF OF CLAIM 7. Let $(x_m) \in K$. First suppose that $(x_m) \notin \mathcal{O}_0 = \psi_0^{-1}(K_0)$. Then $x_0 = \infty$. Let *i* be maximal such that $x_i = \infty$. Then $v((x_m)) = v_{\min}(x_{i+1}) < 0$. Hence, $(x_m) \notin \mathcal{O}_v$. For the converse, suppose that $(x_m) \in \mathcal{O}_0 \setminus \{0\}$. Then $x_0 \neq \infty$. If $x_0 = 0$, then let $\ell \in \omega$ be maximal with $x_{\ell-1} = 0$. We obtain $v((x_m)) = v_{\min}(x_\ell) > 0$ and thus $(x_m) \in \mathcal{O}_v$. If $x_0 \neq 0$, then $x_i \neq 0$ for every $i \in \omega$. Hence, $v((x_m)) = 0$, also showing $(x_m) \in \mathcal{O}_v$.

CLAIM 8. For any $n \in \omega$ we have $v_0(\mathcal{O}_n^{\times}) = \coprod_{\gamma=-2n+1}^0 A_{-\gamma}$ and therefore

(4)
$$v_n K = \left(\prod_{\gamma > -\infty}^{0} A_{-\gamma}\right) / \left(\prod_{\gamma = -2n+1}^{0} A_{-\gamma}\right) = \prod_{\gamma > -\infty}^{-2n} A_{-\gamma}.$$

PROOF OF CLAIM 8. Let $(x_m) \in \mathcal{O}_1^{\times}$. Then $x_1 \in K_1 \setminus \{0\}$. Hence, by the inclusion of ordered fields above, we have $x_1 \in \mathbb{R}\left(\left(\coprod_{\gamma=-1}^0 A_{-\gamma}\right)\right)$ and thus $v_0((x_m)) = v_{\min}(x_1) \in \coprod_{\gamma=-1}^0 A_{-\gamma}$. By iteration of this argument, we obtain (4) for any $n \in \omega$.

To complete our construction, set $u_{2n} = v_n$ for any $n \in \omega$. Claim 2 and Claim 8 yield that u_{2n} is a $(2n+1)_{\leq}$ -henselian (and thus $(2n)_{\leq}$ -henselian) valuation with $u_{2n}K = \coprod_{\gamma > -\infty}^{-2n} A_{-\gamma}$. Now let u_{2n+1} be the coarsening of u_{2n} induced by the convex subgroup A_{2n} of $u_{2n}K$. Since u_{2n+1} is the coarsening of a $(2n+1)_{\leq}$ -henselian valuation, it is itself also $(2n+1)_{\leq}$ -henselian. By definition of u_{2n+1} , its value group is given by $u_{2n+1}K = \coprod_{\gamma > -\infty}^{-2n-1} A_{-\gamma}$, as required.

REMARK 4.6. In Construction 4.5 we could have started with any ordered field K_0 (instead of \mathbb{R}) to obtain the required *t*-henselian non-henselian ordered field. In fact, one can also start with any field K_0 of characteristic 0 (so Lemma 4.3 is applicable) to obtain a *t*-henselian non-henselian field (without an ordering) with the specified value groups.

THEOREM 4.7. Let $A \neq \{0\}$ be an ordered abelian group. Then there exists an ordered henselian valued field (L, <, v) such that $vL = \coprod_{-\omega} A$ and (L, <)admits no non-trivial 0- \mathcal{L}_{or} -definable henselian valuation. In particular, v is not 0- \mathcal{L}_{or} -definable.

PROOF. For any \mathcal{L}_{og} -sentence σ , let σ^* be the \mathcal{L}_{vf} -sentence such that for any valued field (F, w) we have $wF \models \sigma$ if and only if $(F, w) \models \sigma^*$.⁹ Let $G = \coprod_{-\omega} A$ and let $\operatorname{Th}(G)$ be the complete \mathcal{L}_{og} -theory of G. We set Σ to be the \mathcal{L}_{vf} -theory $\{\sigma^* \mid \sigma \in \operatorname{Th}(G)\}$.

Let (K, <) and u_n (for any $n \in \mathbb{N}$) be as in Construction 4.5 for the case $A_{\gamma} = A$ for any $\gamma \in \omega$. For any $n \in \mathbb{N}$ we let h_n be the \mathcal{L}_{vf} -sentence stating that v is a non-trivial n_{\leq} -henselian valuation. Finally, let $\mathrm{Th}(K, <)$ be the complete \mathcal{L}_{or} -theory of (K, <).

For any $n \in \mathbb{N}$, the ordered valued field $(K, <, u_n)$ satisfies $\{h_i \mid i \leq n\}$. Since $u_n K = G$, we also have $(K, u_n) \models \Sigma$. Hence, any finite subset of the \mathcal{L}_{ovf} -theory $\Sigma \cup \{h_n \mid n \in \mathbb{N}\} \cup \text{Th}(K, <)$ is satisfiable. By the Compactness Theorem, we obtain an ordered valued field (F, <, w) such that $(F, <) \models \text{Th}(K, <)$, $wF \models \text{Th}(G)$ and w is non-trivial, n_{\leq} -henselian for any $n \in \mathbb{N}$ and thus henselian. We set L = Fw((G)) and $v = v_{\min}$ on L. Since Fw = Lv and $wF \equiv G = vL$, the Ax–Kochen–Ershov principle for ordered fields (cf. [7, Corollary 4.2]) implies $(L, <, v) \equiv (F, <, w)$. As $(F, <) \equiv (K, <)$, we also obtain $(L, <) \equiv (K, <)$.

By elementary equivalence, any non-trivial $0-\mathcal{L}_{\text{or}}$ -definable henselian valuation in (L, <) corresponds to a non-trivial $0-\mathcal{L}_{\text{or}}$ -definable henselian valuation in (K, <). However, since (K, <) does not admit any non-trivial henselian valuation, this shows the required conclusion.

⁹Syntactically, the sentence σ^* is obtained from σ as follows: Any instance of quantification Qx (where $Q \in \{\exists, \forall\}$ and x is a variable) is replaced by $Q(x \neq 0)$, any occurrence of the constant symbol 0 is replaced by v(1), any occurrence of a variable x is replaced by v(x) and finally any expression of the form v(t) + v(s) is replaced by v(ts).

REMARK 4.8. 1. Let $A \neq \{0\}$ be an ordered abelian group. In order to obtain an ordered henselian valued field (L, <, v) such that $vL = \coprod_{-\omega} A$ and v is not $0 \cdot \mathcal{L}_{or}$ -definable, one can also proceed as follows:¹⁰ Let $L = K((\coprod_{\mathbb{Z}} A))$ for some ordered field (K, <). Now let v be the coarsening of v_{\min} induced by the convex subgroup $\coprod_{\mathbb{N}} A$ of $v_{\min} L$. Then $Lv = K((\coprod_{\mathbb{N}} A))$. Now let w be the strict coarsening of v induced by the convex subgroup $\coprod_{\{0\}} A$ of vL. Then $wL = \coprod_{-\mathbb{N}} A \cong vL$ and $Lw = K((\coprod_{\omega} A)) \cong Lv$. We fix $a \in \mathcal{O}_w \setminus \mathcal{O}_v$. Assume, for a contradiction, that there is a parameter-free \mathcal{L}_{or} -formula $\varphi(x)$ defining \mathcal{O}_v . Then we have that

$$(L, <) \models \neg \varphi(a)$$

By the Ax-Kochen-Ershov principle for ordered fields we have that $(L, <, v) \equiv (L, <, w)$, whence $\varphi(x)$ also defines \mathcal{O}_w . Since $a \in \mathcal{O}_w$, we obtain

$$(L, <) \models \varphi(a),$$

a contradiction.

- 2. Let (L, <, v) be as in Theorem 4.7. Since any non-trivial coarsening w of v is also henselian, also w is not $0-\mathcal{L}_{or}$ -definable. Moreover, for any convex subgroup $B \subseteq A$, the corresponding coarsening w of v with value group $(\coprod_{-\mathbb{N}} A) \amalg (A/B)$ is not $0-\mathcal{L}_{or}$ -definable.
- 3. Theorem 4.7 stands in contrast to [14, Theorem 4], which implies that any henselian valuation v such that vK is non-divisible and archimedean is already $0-\mathcal{L}_r$ -definable. In this regard, note that the rank $-\omega$ of $\coprod_{-\omega} A$ has no minimum, whence the valuation v in Theorem 4.7 has no coarsening with archimedean value group.

We can now use Theorem 4.7 to show that the conclusion in Fact 1.1 cannot be strengthened to parameter-free definability.

EXAMPLE 4.9. The ordered abelian group $\coprod_{-\omega} \mathbb{Z}$ is discretely ordered (with $\mathbb{1}_0$ as least positive element). By Theorem 4.7, there exists an ordered henselian valued field (L, <, v) with $vL = \coprod_{-\omega} \mathbb{Z}$ such that v is not $0-\mathcal{L}_{or}$ -definable. \dashv

Similarly, we show that Fact 1.2, Fact 1.3 and Theorem 3.1 cannot be strengthened to parameter-free definability.

EXAMPLE 4.10. Consider the densely ordered abelian group

$$A = \left\{ \frac{a}{2^n} \mid a, n \in \mathbb{Z} \right\} \subseteq \mathbb{Q}.$$

Then A is regular but not 3-divisible, and is (isomorphic to) a convex subgroup of $G := \coprod_{-\omega} A$. Proposition 3.3 yields that G is not closed in G^{div} . By Theorem 4.7, there exists an ordered field (L, <) and a henselian valuation v on L such that vL = G and v is not 0- \mathcal{L}_{or} -definable.

REMARK 4.11. By [13, Lemma 2.3.7], the following holds:¹¹ Let (K, v) be a henselian valued field and let $p \in \mathbb{N}$ be prime. Suppose that vK is densely ordered and p-regular but not p-divisible. Then v is $0-\mathcal{L}_{r}$ -definable.

 $^{^{10}}$ We thank the referee for this argument and for the permission to include it in our paper.

¹¹A similar argument can also be found in [16, Proof of Proposition 3.7].

In contrast to this result, Example 4.10 shows that the mere existence of a convex p-regular subgroup that is not p-divisible does, in general, not ensure definability without parameters.

§5. Further work. We conclude this work by collecting open questions which may serve as a possible starting point for future work on topics related to this paper.

For a class C of non-trivial ordered abelian groups we consider the following:

 $(\dagger)_0$ If (K, v) is a henselian valued field such that $vK \in \mathcal{C}$, then v is $0-\mathcal{L}_{r}$ -definable.

(†) If (K, v) is a henselian valued field such that $vK \in C$, then v is \mathcal{L}_r -definable.

The corresponding properties $(\not{t})_0$ and (\not{t}) stand for the negations of the above; that is, a class \mathcal{C} of non-trivial ordered abelian groups satisfies $(\not{t})_0$ (respectively (\not{t})) if there exists a henselian valued field (K, v) such that $vK \in \mathcal{C}$ but v is not $0-\mathcal{L}_r$ -definable (respectively \mathcal{L}_r -definable).

To summarise the results from the literature as well as this work, we distinguish between discretely and densely ordered abelian groups.

Discrete case. We consider the following two elementary classes of ordered abelian groups.

 $\mathcal{C}_0^{\text{discr}}$: all regular discretely ordered abelian groups;

 $\mathcal{C}_1^{\text{discr}}$: all discretely ordered abelian groups.

The class C_0^{discr} consists of all ordered abelian groups that are elementarily equivalent to \mathbb{Z} as \mathcal{L}_{og} -structures. These are also called \mathbb{Z} -groups (cf. e.g. [21, Fact 3.4]). By [14, Theorem 4], C_0^{discr} satisfies (†)₀.

The ordered abelian group $\mathbb{Z} \amalg \mathbb{Z}$ serves as an example of a non-regular discretely ordered abelian group, showing that C_0^{discr} is strictly contained in C_1^{discr} . By Fact 1.1 the class C_1^{discr} satisfies (†), and due to Example 4.9 this class does not satisfy (†)₀. We thus obtain the following picture:

$$\underbrace{\mathcal{C}_{0}^{\text{discr}}}_{(\dagger)_{0}} \quad \subsetneq \underbrace{\mathcal{C}_{1}^{\text{discr}}}_{(\dagger) \text{ and } (\dagger)_{0}}$$

Dense case. We consider the following four elementary classes of ordered abelian groups C_i^{dense} consisting of all densely ordered abelian groups G satisfying the condition c_i , where c_i is given as follows:

- c_0 : for some prime p, the group G is p-regular but not p-divisible;
- c_1 : for some prime p, some convex subgroup of G is p-regular but not p-divisible;
- c_2 : G is not closed in G^{div} ;
- c_3 : for some prime p, the group G contains no non-trivial p-divisible convex subgroup.

While C_0^{dense} satisfies $(\dagger)_0$, the class C_1^{dense} satisfies (\dagger) but not $(\dagger)_0$ (see Example 4.10 and Remark 4.11). By Proposition 3.3 and Example 3.4, the class C_1^{dense} is strictly contained in C_2^{dense} . Moreover, Theorem 3.1 yields that C_2^{dense} satisfies (\dagger) .

Further, C_2^{dense} is strictly contained in C_3^{dense} (cf. [21, Proposition 3.18 and Example 3.20]). The remarks in [4, page 1147] show that for any densely ordered abelian group *G* not satisfying condition c_3 above there exists an (almost real closed) valued field (K, v) with vK = G such that v is not \mathcal{L}_r -definable. Hence, any class of densely ordered abelian groups \mathcal{D} that strictly contains C_3^{dense} does not satisfy (†).

As a result of the discussion above, we obtain the following picture:

$$\underbrace{ \mathcal{C}_{0}^{\text{dense}}}_{(\dagger)_{0}} \subsetneq \underbrace{ \mathcal{C}_{1}^{\text{dense}} \subsetneq \mathcal{C}_{2}^{\text{dense}}}_{(\dagger) \text{ and } (f)_{0}} \varsubsetneq \underbrace{ \mathcal{C}_{3}^{\text{dense}}}_{(f)}$$

Open questions. While from the observations above it is clear that the class \mathcal{G}_0 of ordered abelian groups for which $(\dagger)_0$ holds is a *proper* subclass of the class \mathcal{G} of ordered abelian groups satisfying (\dagger) , none of the described dividing lines are yet proven to be sharp. Therefore, we pose the following questions:

QUESTION 5.1. (i) What is the largest class of ordered abelian groups satisfying (\dagger) , respectively $(\dagger)_0$?

(ii) Is there a characterisation of \mathcal{G}_0 inside \mathcal{G} ?

Concluding Remark. As noted in the introduction, in this paper we only consider conditions on the value group. There are also conditions for the definability of a given henselian valuation only depending on its residue field (cf. e.g. [15, Proposition 3.1 and Corollaries 3.3 & 3.8] and [21, Theorem 5.3 (3)]). In particular, the analogous of (\dagger) and $(\dagger)_0$ can be formulated for residue fields. Such conditions will be the subject of a further publication.

References

[1] S. ANSCOMBE and F. JAHNKE, 'Henselianity in the language of rings', Ann. Pure Appl. Logic 169 (2018) 872–895, doi:10.1016/j.apal.2018.04.008.

[2] J. Ax, 'On the undecidability of power series fields', Proc. Amer. Math. Soc. 16 (1965), 846, doi:10.2307/2033940.

[3] D. BILJAKOVIC, M. KOCHETOV and S. KUHLMANN, 'Primes and irreducibles in truncation integer parts of real closed fields', *Logic in Tehran*, Lect. Notes Log. 26 (eds A. Enayat, I. Kalantari and M. Moniri; Assoc. Symbol. Logic, La Jolla, CA, 2006) 42–65.

[4] F. DELON and R. FARRÉ, 'Some model theory for almost real closed fields', J. Symb. Log. 61 (1996) 1121–1152, doi:10.2307/2275808.

[5] K. DUPONT, A. HASSON and S. KUHLMANN, 'Definable valuations induced by multiplicative subgroups and NIP fields', Arch. Math. Logic 58 (2019) 819–839, doi:10.1007/s00153-019-00661-2.

[6] A. J. ENGLER and A. PRESTEL, Valued Fields, Springer Monogr. Math. (Springer, Berlin, 2005).

[7] R. FARRÉ, 'A transfer theorem for Henselian valued and ordered fields', J. Symb. Log. 58 (1993) 915–930, doi:10.2307/2275104.

[8] A. FEHM and F. JAHNKE, 'On the quantifier complexity of definable canonical Henselian valuations', *MLQ Math. Log. Q.* 61 (2015) 347–361, doi:10.1002/malq.201400108.

[9] A. FEHM and F. JAHNKE, 'Recent progress on definability of Henselian valuations', Ordered Algebraic Structures and Related Topics, Contemp. Math. 697 (eds F. Broglia, F. Delon, M. Dickmann, D. Gondard-Cozette and V. A. Powers; Amer. Math. Soc., Providence, RI, 2017), 135–143, doi:10.1090/conm/697/14049. [10] Y. HALEVI and A. HASSON, 'Eliminating field quantifiers in strongly dependent henselian fields', *Proc. Amer. Math. Soc.* 147, 2213–2230 (2019), doi:10.1090/proc/14203.

[11] Y. HALEVI and A. HASSON, 'Strongly Dependent Ordered Abelian Groups and Henselian Fields', Israel J. Math. 232, 719–758 (2019), doi:10.1007/s11856-019-1885-3.

[12] Y. HALEVI, A. HASSON and F. JAHNKE, 'A conjectural classification of strongly dependent fields', *Bull. Symb. Log.* 25, 182–195 (2019), doi:10.1017/bsl.2019.13.

[13] J. HONG, 'Immediate expansions by valuations of fields', Doctoral Thesis, McMaster University, 2013.

[14] J. HONG, 'Definable non-divisible Henselian valuations', Bull. Lond. Math. Soc. 46 (2014) 14–18, doi:10.1112/blms/bdt074.

[15] F. JAHNKE and J. KOENIGSMANN, 'Definable Henselian valuations', J. Symb. Log. 80 (2015) 85–99, doi:10.1017/jsl.2014.64.

[16] F. JAHNKE and J. KOENIGSMANN, 'Defining coarsenings of valuations', Proc. Edinb. Math. Soc. (2) 60 (2017) 665–687, doi:10.1017/S0013091516000341.

[17] F. JAHNKE, P. SIMON and E. WALSBERG, 'Dp-minimal valued fields', J. Symb. Log. 82 (2017) 151–165, doi:10.1017/jsl.2016.15.

[18] W. JOHNSON, 'The canonical topology on dp-minimal fields', J. Math. Log. 18 (2018) 1850007, doi:10.1142/S0219061318500071

[19] M. KNEBUSCH and M. J. WRIGHT, 'Bewertungen mit reeller Henselisierung', J. Reine Angew. Math. 286/287 (1976) 314–321, doi:10.1515/crll.1976.286-287.314.

[20] L. S. KRAPP, 'Algebraic and Model Theoretic Properties of O-minimal Exponential Fields', Doctoral Thesis, Universität Konstanz, 2019, urn:nbn:de:bsz:352-2-166bghaubh8tf9.

[21] L. S. KRAPP, S. KUHLMANN and G. LEHÉRICY, 'Ordered Fields Dense in Their Real Closure and Definable Convex Valuations', *Forum Math.* 33 (2021) 953–972, doi:10.1515/forum-2020-0030.

[22] L. S. KRAPP, S. KUHLMANN and G. LEHÉRICY, 'Strongly NIP almost real closed fields', MLQ Math. Log. Q. 67 (2021) 321–328, doi:10.1002/malq.202000060.

[23] S. KUHLMANN, Ordered Exponential Fields, Fields Inst. Monogr. 12 (Amer. Math. Soc., Providence, RI, 2000), doi:10.1090/fim/012.

[24] D. MARKER, Model Theory: An Introduction, Grad. Texts in Math. 217 (Springer, New York, 2002), doi:10.1007/b98860.

[25] A. PRESTEL and M. ZIEGLER, 'Model-theoretic methods in the theory of topological fields', J. Reine Angew. Math. 299/300 (1978) 318–341, doi:10.1515/crll.1978.299-300.318.

[26] J. ROBINSON, 'Definability and decision problems in arithmetic', J. Symb. Log. 14 (1949) 98–114, doi:10.2307/2266510.

[27] S. SHELAH, 'Strongly dependent theories', Israel J. Math. 204 (2014) 1–83, doi:10.1007/s11856-014-1111-2.

FACHBEREICH MATHEMATIK UND STATISTIK UNIVERSITÄT KONSTANZ

78457 KONSTANZ, GERMANY *E-mail*: sebastian.krapp@uni-konstanz.de *URL*: http://www.math.uni-konstanz.de/~krapp/

FACHBEREICH MATHEMATIK UND STATISTIK UNIVERSITÄT KONSTANZ

78457 KONSTANZ, GERMANY

E-mail: salma.kuhlmann@uni-konstanz.de *URL*: https://www.mathematik.uni-konstanz.de/kuhlmann/

FACHBEREICH MATHEMATIK UND STATISTIK

UNIVERSITÄT KONSTANZ

78457 KONSTANZ, GERMANY

E-mail:moritz.link@uni-konstanz.de

URL: https://www.mathematik.uni-konstanz.de/volkwein/members/research-group/

doctoral-students/moritzlink/