Classification of one dimensional dynamical systems by countable structures

Henk Bruin

Benjamin Vejnar*

Faculty of Mathematics
University of Vienna
Oskar Morgensternplatz 1
1090 Vienna, Austria
henk.bruin@univie.ac.at

Faculty of Mathematics and Physics Charles University Prague, Czechia

vejnar@karlin.mff.cuni.cz

September 5, 2022

Abstract

We study the complexity of the classification problem of conjugacy on dynamical systems on some compact metrizable spaces. Especially we prove that the conjugacy equivalence relation of interval dynamical systems is Borel bireducible to isomorphism equivalence relation of countable graphs. This solves a special case of the Hjorth's conjecture which states that every orbit equivalence relation induced by a continuous action of the group of all homeomorphisms of the closed unit interval is classifiable by countable structures. We also prove that conjugacy equivalence relation of Hilbert cube homeomorphisms is Borel bireducible to the universal orbit equivalence relation.

Mathematics Subject Classification (2010): Primary: 54H20 Secondary: 03E15,

Keywords. Conjugacy, classification, Borel reduction, countable structures, universal orbit equivalence relation

1 Introduction

Measuring the complexity of relations on structures is a very general task. In this paper we use the notion of *Borel reducibility* (see Definition 1) and the results of invariant descriptive set theory to compare the complexities of classification problems. For more

^{*}This work has been supported by Charles University Research Centre program No.UNCE/SCI/022.

details on invariant descriptive set theory we refer to the book by Gao [Gao09]. For a short and nice introduction to the theory of Borel reductions we refer to a paper by Foreman [For18].

Several equivalence relations became milestones in this theory. Let us mention four of those, which describe an increasing chain of complexities:

- the equality on an uncountable Polish space,
- the equality of countable sets of real numbers,
- the S_{∞} -universal orbit equivalence relation (S_{∞} is the group of permutations on \mathbb{N}),
- the universal orbit equivalence relation.

Let us give several examples to make the reader more familiar with the above relations. A classical example is a result of Gromov (see e.g. [Gao09, Theorem 14.2.1]) who proved that the isometry equivalence relation of compact metric spaces is a smooth equivalence relation, which means that it is Borel reducible to the equality of real numbers (or equivalently of an uncountable Polish space). The isomorphism relation of countable graphs or the isomorphism relation of countable linear orders are Borel bireducible to the S_{∞} -universal orbit equivalence. The homeomorphism equivalence relation of compact metrizable spaces and the isometry relation of separable complete metric spaces were proved by Zielinski in [Zie16] and by Melleray in [Mel07], respectively, to be Borel bireducible to the universal orbit equivalence relation (see the survey paper by Motto Ros [MR17]).

In order to capture all the structures in one space we need some sort of coding. This can be done by considering some universal space (e.g. the Hilbert cube or the Urysohn space) and all its subspaces with some natural Polish topology or Borel structure (e.g the hyperspace topology or the Effros Borel structure). Sometimes there are other natural ways to encode a given structure. For example the class of separable complete metric spaces can be coded by the set of all metrics on N where two metrics are defined to be equivalent if the completions of the respective spaces are isometric. Fortunately in this case, by [Gao09, Theorem 14.1.3] it does not matter which of the two coding we choose. It is generally believed that this independence on a natural coding is common to other structures and thus the statements are usually formulated for all structures without mentioning the current coding. Nevertheless, for the formal treatment some coding is always necessary.

The aim of this paper is to determine the complexity of some classification problems of dynamical systems up to conjugacy. Dynamical systems of a fixed compact metrizable space X can be naturally coded as a space of continuous mappings of X into itself, with the uniform topology. This one as well as the subspace of all self-homeomorphisms is well known to be a Polish space.

Let us mention several results which are dealing with the complexity of conjugacy equivalence relation. It was proved by Hjorth that conjugacy equivalence relation of homeomorphisms of [0,1] is classifiable by countable structures [Hjo00, Section 4.2] (in fact Borel

bireducible to the universal S_{∞} -orbit equivalence relation) but conjugacy of homeomorphisms of $[0,1]^2$ is not [Hjo00, Section 4.3]. By a result of Camerlo and Gao, conjugacy equivalence relation of both selfmaps and homeomorphisms of the Cantor set are Borel bireducible to the S_{∞} -universal orbit equivalence relation [CG01, Theorem 5]. Kaya proved that conjugacy of pointed minimal Cantor dynamical systems is Borel bireducible to the equality of countable subsets of reals [Kay17b]. Conjugacy of odometers is smooth due to Buescu and Stewart [BS95]. The complexity of conjugacy of Toepliz subshifts was treated several times – by Thomas, Sabok and Tsankov, and by Kaya [Tho13, ST17, Kay17a]. Conjugacy of two-sided subshifts is Borel bireducible to the universal countable Borel equivalence relation due to Clemens [Cle09]. There is an extensive exposition of results on the complexity of conjugacy equivalence relation on subshifts of 2^G for a countable group G in the book by Gao, Jackson and Seward [GJS16, Chapter 9]. Recently, during the 8th Visegrad Conference on Dynamical Systems in 2019 it was announced by Dominik Kwietniak that conjugacy of shifts with specification is Borel bireducible to the universal countable Borel equivalence relation.

In this paper, we deal with some of the missing parts. By mainly elementary and standard tools (excluding the complexity level of countable structures), we prove:

Theorem A (see Theorem 17). The conjugacy equivalence relation of interval maps is Borel bireducible to the S_{∞} -universal orbit equivalence relation.

Also, we prove:

Theorem B (see Theorem 27). The conjugacy equivalence relation of homeomorphisms as well as conjugacy of selfmaps of the Hilbert cube is Borel bireducible to the universal orbit equivalence relation.

To this end we use some tools of infinite dimensional topology and a result of Zielinski on the complexity of homeomorphism equivalence relation of metrizable compacta [Zie16] combining with some ideas of P. Krupski and the second author [KV20]. Finally we make a small overview on the complexity of conjugacy equivalence relation of dynamical systems on the Cantor set, on the interval, on the circle and on the Hilbert cube.

2 Definitions and notations

Let us define some standard notions from descriptive set theory (see e.g. [Kec95]). A Polish space is a separable completely metrizable topological space. Recall that a standard Borel space is a measurable space (X, S) such that there is a Polish topology τ on X for which the family of Borel subsets of (X, τ) is equal to S. In order to compare the complexities of equivalence relations we use the notion of Borel reducibility.

Definition 1. Suppose that X and Y are sets and let E, F be equivalence relations on X and Y respectively. We say that E is *reducible* to F, and we denote this by $E \leq F$, if there exists a mapping $f: X \to Y$ such that

$$xEx' \iff f(x)Ff(x'),$$

for every $x, x' \in X$. The mapping f is called a reduction of E into F. If the sets X and Y are endowed with Polish topologies (or standard Borel structures), we say that E is Borel reducible to F, and we write $E \leq_B F$, if there is a reduction $f: X \to Y$ of E into F which is Borel measurable. We say that E is Borel bireducible to F, and we write $E \sim_B F$, if E is Borel reducible to F and F is Borel reducible to E.

In a similar fashion we define being *continuously reducible* if in addition X and Y are Polish spaces and f is continuous.

In the whole paper we set \mathbb{N} for all positive integers, I = [0, 1] and we denote the closure operator by Cl. For a separable metric space X we denote by $\mathcal{K}(X)$ the hyperspace of all compacta in X with the Hausdorff distance d_H and the corresponding Vietoris topology. If X is a Polish space $\mathcal{K}(X)$ is known to be Polish. For compact metric spaces X, Y we consider the space C(X,Y) of all continuous mappings of X into Y with the supremum metric. In this way we get a Polish space. Especially the collection of all continuous selfmaps of X is denoted shortly by C(X). We also denote by Inj(X,Y) the collection of all embeddings of X into Y and by $\mathcal{H}(X)$ the collection of all homeomorphisms of X. These are again known to be Polish spaces.

The equality equivalence relation of real numbers is denoted as $E_{=}$. We denote by $E_{=+}$ the equivalence relation on $\mathbb{R}^{\mathbb{N}}$ defined by $(a_n)E_{=+}(b_n)$ if and only if $\{a_n : n \in \mathbb{N}\} = \{b_n : n \in \mathbb{N}\}$. The last equivalence relation is called the *equality of countable sets*.

We say that an equivalence relation E defined on a standard Borel space X is classifiable by countable structures if there is a countable relational language \mathcal{L} such that E is Borel reducible to the isomorphism relation of \mathcal{L} -structures whose underlying set is \mathbb{N} . An equivalence relation E on a standard Borel space X is said to be an orbit equivalence relation if there is a Borel action of a Polish group G on X such that xEx' if and only if there is some $g \in G$ for which gx = x'.

Let \mathcal{C} be a class of equivalence relations on standard Borel spaces. An element $E \in \mathcal{C}$ is called universal for \mathcal{C} if $F \leq_B E$ for every $F \in \mathcal{C}$. It is known that for every Polish group G there is an equivalence relation (denoted by E_G) on a standard Borel space that is universal for all orbit equivalence relations given by continuous G-actions. We are particularly interested in the universal S_{∞} -equivalence relation $E_{S_{\infty}}$, where S_{∞} is the group of all permutations of \mathbb{N} . It is known that an equivalence relation is classifiable by countable structures if and only if it is Borel reducible to $E_{S_{\infty}}$. Moreover $E_{S_{\infty}}$ is known to be Borel bireducible to isomorphism equivalence relation of countable graphs. Also there exists a universal orbit equivalence relation which is denoted by $E_{G_{\infty}}$. We should also

note that all the mentioned equivalence relations are analytic sets, i.e. images of standard Borel spaces with respect to a Borel measurable map. We have a chain of complexities

$$E_{=} \leq_{B} E_{=+} \leq_{B} E_{S_{\infty}} \leq_{B} E_{G_{\infty}}$$

and it is known that none of these Borel reductions can be reversed.

3 Interval dynamical systems

In this section we prove that conjugacy of interval dynamical systems is classifiable by countable structures. The strategy of our proof is as follows. In the first part we describe a natural reduction of interval dynamical systems to some kind of countable structures. We assign to every $f \in C(I)$ a countable invariant set $C_f \subseteq I$ of some dynamically exceptional points for f. Since the set C_f does not need to be dense in I we do not have enough information to capture the dynamics of f by restricting to C_f . On the other hand the dynamics on the maximal open intervals of $I \setminus C_f$ is quite simple. Hence it will be enough to define an invariant countable dense subset D_f in $I \setminus Cl(C_f)$ arbitrarily. Consequently, we get that for f conjugate to g there exists a conjugacy of f to g which sends the set $C_f \cup D_f$ onto $C_g \cup D_g$. Finally it is enough to assign to every $f \in C(I)$ a countable structure $\Psi(f)$ whose underlying set is $C_f \cup D_f$ and which is equipped with one binary relation $\leq \upharpoonright_{C_f \cup D_f}$ and one mapping $f \upharpoonright_{C_f \cup D_f}$ (which can be as usual considered as a binary relation). We will prove then that if two such structures $\Psi(f)$ and $\Psi(g)$ are isomorphic then f and g are conjugate.

In the second part we prove that this reduction can be modified using some sort of coding so that the assigned countable structures share the same support and so that the new reduction is Borel. To this end we use Lusin-Novikov selection theorem [Kec95, Theorem 18.10] several times.

For $g \in C(I)$ we denote by Fix(g) the set of fixed points of g, i.e. those points for which g(x) = x. We omit the proof of the following "folklore" lemma. The key idea of the proof is the back and forth argument.

Lemma 2. Let $f, g \in C(I)$ be increasing homeomorphisms such that $Fix(f) = Fix(g) = \{0,1\}$ and let $A, B \subseteq (0,1)$ be countable dense sets that are invariant in both directions for f and g respectively. Then there is a conjugacy h of f and g satisfying h(A) = B.

Definition 3. For $f \in C(I)$ let us say that a point $z \in I$ is a *left sharp local maximum* of f if there is some $\delta > 0$ such that f(x) < f(z) for $x \in (z - \delta, z)$ and $f(x) \le f(z)$ for $x \in (z, z + \delta)$. In a similar fashion we define *left sharp local minimum*, right sharp local minimum and right sharp local maximum.

Notation 4. Let M_f be the union of $\{0,1\}$ and the set of all left and right sharp local maxima and minima. It is easily shown that the set M_f is countable. For a closed set

 $F \subseteq I$ denote by Acc(F) the set of all accessible points of F in \mathbb{R} , i.e. those points $x \in F$ for which there exists an open interval $(a,b) \subseteq \mathbb{R} \setminus F$ for which x=a or x=b.

For every $f \in C(I)$ let us denote by C_f the smallest set such that

- a) $M_f \subseteq C_f$,
- b) if $f^{-1}(y)$ contains an interval then $y \in C_f$,
- c) if $n \in \mathbb{N}$ then $Acc(Fix(f^n)) \subseteq C_f$,
- d) $f(C_f) \subseteq C_f$,
- e) if $y \in C_f$ then $Acc(f^{-1}(y)) \subseteq C_f$.

Lemma 5. The set C_f is countable for every $f \in C(I)$.

Proof. Let S_1 be the union of M_f , all the values of f at locally constant points and all the sets $Acc(Fix(f^n))$ for $n \in \mathbb{N}$. Clearly S_1 is countable. Let $S_{i+1} = S_i \cup f(S_i) \cup \{Acc(f^{-1}(y)): y \in S_i\}$. Clearly $C_f = \bigcup \{S_i: i \in \mathbb{N}\}$ and thus it is countable. \square

Note that C_f depends only on the topological properties of I and the dynamics of f. That is if f and g are conjugate by some homeomorphism h, then $h(C_f) = C_g$. This is clear because h maps M_f onto M_g , locally constant intervals of f to locally constant intervals of g and periodic points of g.

Let us denote by \mathcal{J}_f be the collection of all maximal open subintervals of $I \setminus C_f$.

Lemma 6. Let $J \in \mathcal{J}_f$. Then either $f \upharpoonright_J$ is constant or $f \upharpoonright_J$ is one to one and in this case $f(J) \in \mathcal{J}_f$. Also $f^{-1}(J)$ is the finite union (possibly the empty union) of elements of \mathcal{J}_f .

Proof. Let us prove first that $f \upharpoonright_J$ is either constant or one-to-one. Suppose that the contrary holds. Then there are points $x, y, z \in J$ such that $f(x) = f(y) \neq f(z)$ and $x \neq y$. Let us suppose that x < y < z and f(x) < f(z) (the other possibilities are just easy modifications). Let $u = \min f \upharpoonright_{[x,z]}$ and let $v = \max(f^{-1}(u) \cap [x,z])$. It follows that $v \in (x,z)$ is a right sharp local minimum. By Notation 4 a) it follows that $v \in C_f$ which is a contradiction since J is disjoint from C_f .

Suppose now that $f \upharpoonright_J$ is one-to-one and let us prove that $f(J) \in \mathcal{J}_f$. Observe first that f(J) is disjoint from C_f , otherwise there would be a point $y \in f(J) \cap C_f$ and since $f^{-1}(y)$ is a closed set not containing the whole set J there will be a point in $\operatorname{Acc} f^{-1}(y) \cap J$ which is a contradiction with Notation 4 e). We need to prove that f(J) is a maximal interval disjoint from C_f . Suppose that J = (a, b). Then there are $a_n, b_n \in C_f$ such that $a_n \to a, b_n \to b$. By continuity of f it follows that $f(a_n) \to f(a)$ and $f(b_n) \to f(b)$. Also $f(a_n), f(b_n) \in C_f$ by Notation 4 d). Thus the maximality follows.

Observe first that $f^{-1}(J)$ is a countable union of disjoint collection of open intervals and if we prove that each of the intervals is mapped by f onto J it will follow by continuity that such a collection is in fact finite. Denote (a, b) = J and let (c, d) be a maximal interval in $f^{-1}(J)$. Clearly $(c,d) \cap C_f = \emptyset$ by Notation 4 d), so it is enough to prove that it is maximal with this property. Note that $f(c), f(d) \in \{a, b\}$ otherwise we get a contradiction with (c,d) being maximal interval in $f^{-1}(J)$. Also it can not happen that f(c)=f(d)otherwise there will be a point of left local maximum or minimum in (c,d) which would produce a point in $M_f \cap (c,d)$, which in turn would give a point in $C_f \cap J$, by Notation 4 a), d). Hence f((c,d)) = J. Moreover, by the first part of this proof we get that $f \upharpoonright_{(c,d)}$ is one-to-one and thus it is either increasing or decreasing. Without loss of generality suppose the first case. Let us distinguish several possibilities. If $f \geq f(c)$ on a left neighborhood of c then c is a point of right sharp local minimum and thus $c \in C_f$. Otherwise choose a sequence $a_n \in C_f$ such that $a_n \to a$. We define points $c_n = \max([0, c] \cap f^{-1}(a_n))$. These are eventually well defined, $c_n \to c$ and $c_n \in \mathrm{Acc} f^{-1}(a_n)$. Hence by Notation 4 e) $c_n \in C_f$. We can proceed in a similar way with the point d and thus the interval (c,d) is maximal subinterval of $I \setminus C_f$.

Example 7. For the tent map $f(x) = \min\{2x, 2(1-x)\}$, the set C_f contains all the dyadic numbers in I, thus C_f is a dense subset of I and hence $\mathcal{J}_f = \emptyset$. For the map $g = \frac{1}{4}f$ we have

$$C_g = \{2^{-n}, 1 - 2^{-n} : n \in \mathbb{N}\} \cup \{0, 1\},$$

 $\mathcal{J}_g = \{(2^{-n-1}, 2^{-n}), (1 - 2^{-n}, 1 - 2^{-n-1}) : n \in \mathbb{N}\}.$

Notation 8. Let G_f be a directed graph on \mathcal{J}_f where (J,K) forms an oriented edge if and only if f(J) = K. Note that for every $K \in \mathcal{J}_f$ there are only finitely many $J \in \mathcal{J}_f$ for which f(J) = K. Hence every vertex of the graph G_f admits only finitely many arrows to enter. Let $E_f = \mathbb{Q} \cap I \setminus \mathrm{Cl}(C_f)$ and let

$$D_f = \bigcup_{n \in \mathbb{Z}} f^n(E_f).$$

Note that the union is taken over all integers. In spite of that it follows by Lemma 6 that D_f is countable. Let us define

$$\Psi(f) = (C_f \cup D_f, \leq \upharpoonright_{C_f \cup D_f}, f \upharpoonright_{C_f \cup D_f}).$$

Theorem 9. The mapping Ψ is a reduction of orientation preserving conjugacy of interval dynamical systems to the isomorphism relation of countable structures.

Proof. Suppose first that f is conjugate to g via some increasing homeomorphism h, that is $f = h^{-1}gh$. We want to find an isomorphism $\varphi \colon \Psi(f) \to \Psi(g)$. Since h does not need

to map D_f to D_g , we need to do some more work. In fact we find a conjugacy \bar{h} of f and g such that $\bar{h}(C_f \cup D_f) = C_g \cup D_g$. Then it will be enough to define a mapping $\varphi \colon C_f \cup D_f \to C_g \cup D_g$ as the restriction of \bar{h} . We will define \bar{h} by parts. First of all we define \bar{h} on the set $\mathrm{Cl}(C_f)$ in the same way as h.

Clearly h induces an isomorphism of the graphs (\mathcal{J}_f, G_f) and (\mathcal{J}_g, G_g) . We will consider the components of the symmetrized graphs G_f and G_g . Note that $J, K \in \mathcal{J}_f$ are in the same component of G_f if there are $m, n \geq 0$ such that $f^m(J) = f^n(K)$.

Let us distinguish two cases for the components of G_f . If a component of G_f contains an oriented cycle, choose an element J in there (note that the cycle is unique). Hence there is $n \in \mathbb{N}$ such that $f^n(J) = J$. By using Notation 4 c) it follows that either all the points of J are fixed points for f^n or there are no fixed points of f^n in J and the same has to be true for g^n on h(J). In the first case we just let $\bar{h} \upharpoonright J$ to be any homeomorphism of $\mathrm{Cl}(J)$ and $\mathrm{Cl}(h(J))$, in the second case we obtain by Lemma 2 that there is a conjugacy $\bar{h} \upharpoonright_{\mathrm{Cl}(J)}$ of $f^n \upharpoonright_{\mathrm{Cl}(J)}$ and $g^n \upharpoonright_{\mathrm{Cl}(h(J))}$ sending $D_f \cap J$ onto $D_g \cap h(J)$. In components which do not contain an oriented cycle we choose J arbitrarily, and let $\bar{h} \upharpoonright_J$ be an arbitrary increasing homeomorphism $J \to h(J)$ which maps $J \cap D_f$ onto $h(J) \cap D_g$.

For any K that is in the same component as J find $m, n \geq 0$ such that $f^m(J) = f^n(K) \in \mathcal{J}_f$ and define \bar{h} on K using the definition of \bar{h} on J as

$$(g^{-n}\upharpoonright_{h(K)})g^m\bar{h}(f^{-m}\upharpoonright_J)f^n.$$

On the other hand suppose that φ is an isomorphism of the countable structure $\Psi(f)$ to $\Psi(g)$. Hence $\varphi \colon C_f \cup D_f \to C_g \cup D_g$ is a bijection preserving the order. Thus it can be extended to an increasing homeomorphism $\tilde{\varphi} \colon I \to I$. We claim that $\tilde{\varphi}$ conjugates f and g. Consider any point $x \in C_f \cup D_f$ and compute

$$g(\tilde{\varphi}(x)) = g(\varphi(x)) = \varphi(f(x)) = \tilde{\varphi}(f(x)),$$

where the middle equality follows form φ being an isomorphism of $\Psi(f)$ and $\Psi(g)$. Since the set $C_f \cup D_f$ is dense it follows by continuity that $g(\tilde{\varphi}(x)) = \tilde{\varphi}(f(x))$ for every $x \in I$. Hence f and g are conjugate.

3.1 Borel coding

We need to verify that the mapping Ψ that was proved in Theorem 9 to be a reduction can be coded in a Borel way. We use standard notation for the Borel hierarchy, especially Σ_1^0 is used for the collection of all open sets, Σ_2^0 is used for the collection of countable unions of closed sets etc. For a set $B \subseteq X \times Y$ and $x \in X$ let us denote by B_x the set $\{y \in Y : (x,y) \in B\}$ and call it *vertical section* of B.

The following seems to be folklore in descriptive set theory, but for the sake of completeness we include a proof.

Proposition 10. Let X, Y be Polish spaces and $B \subseteq X \times Y$ be a Borel set with countable vertical sections. Then the set $\bigcup_{x \in X} \{x\} \times Cl(B_x)$ is Borel as well.

Proof. Let \mathcal{B} be a countable base for the topology of Y. By the Lusin-Novikov selection theorem, we can assume that $B = \bigcup f_n$ for some Borel maps $f_n \colon X \to Y$. It follows that

$$(X \times Y) \setminus \left(\bigcup_{x \in X} \{x\} \times \operatorname{Cl}(B_x)\right) = \bigcup_{U \in \mathcal{B}} \bigcap_{n \in \mathbb{N}} ((X \setminus f_n^{-1}(U)) \times U).$$

Hence the set under discussion is Borel.

Let us denote $\Gamma = \{(K, a) \in \mathcal{K}(I) \times I : a \in Acc(K)\}.$

Lemma 11. The set Γ is a Σ_2^0 -set.

Proof. The sets

$$L_n = \{ (K, a) \in \mathcal{K}(I) \times I \colon a \in K, K \cap (a - 2^{-n}, a) = \emptyset \},$$

$$R_n = \{ (K, a) \in \mathcal{K}(I) \times I \colon a \in K, K \cap (a, a + 2^{-n}) = \emptyset \}$$

are closed for every $n \in \mathbb{N}$. Hence the set $\bigcup (L_n \cup R_n)$ is a Σ_2^0 -set.

Notation 12. For a set $B \subseteq C(I) \times I$ let us define

$$\begin{split} B^{\to} &= \{ (f, f(x)) \colon (f, x) \in B \}, \\ B^{\leftarrow} &= \{ (f, x) \colon (f, f(x)) \in B \}, \\ B^{\Leftarrow} &= \{ (f, x) \colon x \in Acc(f^{-1}(y)), (f, y) \in B \}. \end{split}$$

Lemma 13. Let $B \subseteq C(I) \times I$ be a Borel set with countable vertical sections. Then the sets B^{\rightarrow} , B^{\leftarrow} and B^{\leftarrow} are Borel as well.

Proof. The evaluation mapping $e: C(I) \times I \to I$, e(f,x) = f(x) is continuous. Hence the mapping $\Phi: (f,x) \mapsto (f,e(f,x))$ is continuous as well. Especially, the restriction of Φ to B is Borel and also countable-to-1. Since by [Kec95, 18.14] countable-to-1 image of a Borel set is Borel we conclude that $\Phi(B) = B^{\to}$ is Borel.

By the Lusin-Novikov selection theorem we can write $B = \bigcup F_n$ for some Borel maps F_n . It follows then that

$$B^{\leftarrow} = \bigcup_{n \in \mathbb{N}} \{ (f, x) \colon e(f, x) = F_n(f) \}$$

and thus it is a Borel set.

The mapping $p: C(I) \times I \to \mathcal{K}(I)$, $p(f,y) = f^{-1}(y)$ is upper semicontinuous and hence it is Borel by [Kec95, 25.14]. The set Γ is Borel by Lemma 11 and it has nonempty and countable vertical sections. Hence $\Gamma = \bigcup b_n$ for some Borel mappings $b_n: \mathcal{K}(I) \to I$, by the Lusin-Novikov selection theorem. The mapping $\Psi_n: (f,y) \mapsto (f,b_n(f^{-1}(y))) = (f,b_n(p(f,y)))$ is a Borel mapping and its restriction to B is countable-to-1. Hence by [Kec95, 18.14] the set $\bigcup \Psi_n(B) = B^{\Leftarrow}$ is Borel.

Lemma 14. Let X, Y, Z be standard Borel spaces, $f: X \to Y$ a Borel mapping and $R \subseteq Y \times Z$ a Borel binary relation. Then the set $R \circ f = \{(x, z) : (f(x), z) \in R\}$ is Borel.

Proof. Define $F: X \times Z \to Y \times Z$ by F(x,z) = (f(x),z). Clealy F is a Borel mapping and $R = F^{-1}(R)$ which is consequently a Borel set.

Lemma 15. The set

$$A = \{(f, x) \in C(I) \times I \colon x \in C_f \cup D_f\}$$

is a Borel subset of $C(I) \times I$.

Proof. Let us prove first that the set $B_a := \{(f, x) : x \in M_f\}$ is Borel. As the set $\{(f, x) : x \text{ is a left sharp local maximum}\}$ can be written in the form

$$\bigcup_{\varepsilon>0} \bigcap_{\eta>0} \bigcup_{\delta>0} \{(f,x) \colon \forall z \in [x-\varepsilon,x-\eta] \colon f(z) \le f(x) - \delta \& \forall z \in [x,x+\varepsilon] \colon f(z) \le f(x) \}$$

it follows that it is a Σ_3^0 set. By symmetry it follows that B_a is the union of four Σ_3^0 -sets and thus it is Borel. The set $B_b := \{(f,y) \colon f^{-1}(y) \text{ contains an interval}\}$ is a Σ_2^0 -set. Let $B_c := \{(f,x) \in C(I) \times I \colon x \in \operatorname{Acc}(\operatorname{Fix}(f^n)), n \in \mathbb{N}\}$. The mapping $F_n \colon C(I) \to \mathcal{K}(I), F_n(f) = \operatorname{Fix}(f^n)$ is upper semicontinuous (since if f_i converge uniformly to f and x_i converge to x with $f_i^n(x_i) = x_i$ then $f^n(x) = \lim_i f_i^n(\lim_j x_j) = \lim_j \lim_i f_i^n(x_j) = \lim_j f^n(x_j) = f^n(x)$ by the Moore-Osgood theorem) and thus it is Borel. Since Γ is a Borel set by Lemma 11 we conclude that the composition $\Gamma \circ F_n$ is Borel because the composition of a Borel binary relation and a Borel mapping (in that order) is a Borel relation by Lemma 14. Hence $B_c = \bigcup_{n \in \mathbb{N}} \Gamma \circ F_n$ is Borel. Hence the set $B = B_a \cup B_b \cup B_c$ is Borel. Define recursively $B_1 = B, B_{n+1} = B_n \cup B_n^{\to} \cup B_n^{\leftarrow}$ for $n \in \mathbb{N}$. All these sets are Borel by Lemma 13. It follows that $A_1 = \bigcup B_n = \{(f,x) \colon x \in C_f\}$ is Borel.

Since A_1 has countable vertical sections and it is Borel we conclude using Proposition 10 that $A_2 = \bigcup_{f \in C(I)} (\{f\} \times \operatorname{Cl}(A_{1,f}))$ is Borel as well. Consequently $A_3 = \{(f,x) \colon x \in E_f\} = (C(I) \times \mathbb{Q}) \setminus A_2$ is Borel. By Lemma 13 we conclude that all the sets $A_{n+1} = A_n \cup A_n^{\rightarrow} \cup A_n^{\leftarrow}$, $n \geq 3$ are Borel. Finally $A = A_1 \cup \bigcup_{n \geq 3} A_n$ is a Borel set.

Theorem 16. The orientation preserving conjugacy of interval dynamical systems is Borel bireducible to the S_{∞} -universal orbit equivalence relation.

Proof. By the result of [Hjo00, Section 4.2] orientation preserving conjugacy of increasing interval homeomorphisms is Borel bireducible to the S_{∞} -universal orbit equivalence relation. Hence especially the S_{∞} -universal orbit equivalence relation is Borel reducible to increasing conjugacy of orientation preserving interval dynamical systems.

Let us argue for the converse. The set A from Lemma 15 is Borel and it has nonempty and countable vertical sections. Hence by the Lusin-Novikov selection theorem we can find Borel mappings $F_n \colon C(I) \to I$ such that $\bigcup F_n = A$. Since all the vertical sections are infinite we can additionally suppose that for every pair $(f, x) \in A$ there is exactly one $n \in \mathbb{N}$ satisfying $F_n(f) = x$. Let

$$\Phi(f) = (\mathbb{N}, R, m),$$

where R is a binary relation and m is a unary function such that aRb iff $F_a(f) \leq F_b(f)$ and m(a) = b iff $f(F_a(f)) = F_b(f)$ for $a, b \in \mathbb{N}$. There is a natural isomorphism $\Phi(f) \to \Psi(f)$, $a \mapsto F_a(f)$. Hence clearly Φ is a reduction. It is routine to verify that Φ is Borel by the fact that the mappings F_n are Borel.

Let us note that the same conclusion as in the previous theorem can be proved without assuming orientation preserving conjugacy but with just conjugacy. The reason is that in the proofs of Theorem 16 and Theorem 9 we can simply consider a ternary betweenness relation T instead of the binary relation of linear order \leq , i.e. $(x, y, z) \in T$ if and only if y is an element of the smallest interval containing x and z. This ternary relation is clearly forgetting the order of I. Also by [Hjo00, Exercise 4.14] $E_{S_{\infty}}$ is Borel reducible to conjugacy of interval homeomorphisms. Thus we get the following result.

Theorem 17. The conjugacy of interval dynamical systems is Borel bireducible to the S_{∞} -universal orbit equivalence relation.

We note that Theorem 17 is a special case of Hjorth's conjecture [Hjo00, Conjecture 10.6] stating that every equivalence relation induced by a continuous action of the group $\mathcal{H}(I)$ of all interval homeomorphisms on a Polish space is classifiable by countable structures. In this case the homeomorphism group acts on the space of continuous selfmaps by conjugacy. Similarly one can prove that the orbit equivalence relations induced by natural left or right composition actions of the homeomorphism group on the space of continuous selfmaps is Borel reducible to the S_{∞} -universal equivalence relation. Also it is known that the orbit equivalence induced by the homeomorphism group action $\mathcal{H}(I)$ on the hyperspace $\mathcal{K}(I)$ is Borel bireducible to the S_{∞} -universal orbit equivalence relation (see [Hjo00, Exercise 4.13] or [CG19] for a proof). All these are special cases of Hjorth's conjecture.

4 Hilbert cube dynamical systems

Since the homeomorphism equivalence relation of metrizable compacta is known to be Borel bireducible to the universal orbit equivalence relation, it is not surprising that conjugacy of dynamical systems on the Hilbert cube is of the same complexity, which is the main result of this section. In a dynamical system (X, f), a point x is called a *locally attracting fixed point* if f(x) = x and there is a neighborhood U of x such that for every $z \in U$ the trajectory $(f^n(z))_{n \in \mathbb{N}}$ converges to x. The notion of a Z-set in the Hilbert cube Q plays an important role and it describes a kind of relative homotopical smallness.

Definition 18. A closed subset of a (separable metric) space X is called a Z-set in X if for every open cover \mathcal{U} of X and every continuous function f of the Hilbert cube Q into X there is a continuous function $g \colon Q \to X$ such that f and g are \mathcal{U} -close (i.e. for every $x \in X$ there is $U \in \mathcal{U}$ such that $f(x), g(x) \in U$) and $g(Q) \cap A = \emptyset$.

An introduction to this notion can be found in [vM01, Chapter 5]. Mostly, we will need the following properties on Z-sets in the Hilbert cube. First, every homeomorphism of Z-sets can be extended to a homeomorphism of the Hilbert cube [vM01, Theorem 5.3.7]. Second, the Hilbert cube $Q \times I$ contains a topological copy of itself $Q \times \{0\}$ as a Z-set [vM01, Lemma 5.1.3] and similarly the base in the cone of the Hilbert cube is a Z-set. Third, every closed subset of a Z-set in Q is a Z-set in Q [vM01, Lemma 5.1.2]. If follows from the first and second property that there is topologically just one way, how to embed the Hilbert cube into itself as a Z-set (namely $Q \times \{0\}$ included in $Q \times I$). For the purpose of this paper, an absolute retract is just a space homeomorphic to a retract of the Hilbert cube (which is equivalent to being a retract of every separable metric space, in which it is embedded). A space X is said to have the disjoint cell property if for every $\varepsilon > 0$, $n \in \mathbb{N}$ and continuous mappings $f, g: I^n \to X$ there are continuous mappings $f', g': I^n \to X$ with disjoint images such that f and f' as well as g and g' are ε -close. The last two notions give a topological characterization of the Hilbert cube.

Theorem 19 (Toruńczyk, see e.g. [vM01, Theorem 4.2.25]). A space X is homeomorphic to the Hilbert cube if and only if it is an absolute retract with the disjoint cell property.

The following proposition is a special case of [Zie16, Proposition 1] and it can be easily proved using the back and forth argument. Another reference for the proof is [Lor81, Proposition 9]. Our formulation is using a slightly different language.

Proposition 20. Let $K \subseteq A, L \subseteq B$ be four nonempty compact metrizable spaces such that $A \setminus K$ and $B \setminus L$ are dense sets of isolated points in A and B respectively. Then every homeomorphism of K onto L can be extended to a homeomorphism of A onto B.

The following will be useful in the proof of Theorem 27.

Proposition 21 ([GvM93, Theorem 2.6] or [vM01, Corollary 4.2.24]). If X is a non-degenerate Peano continuum then there exists a homotopy $H: \mathcal{K}(X) \times I \to \mathcal{K}(X)$ for which

- H(A,0) = A for every $A \in 2^X$,
- H(A,t) is finite for every t > 0 and $A \in 2^X$.

Recall that if $Y \subseteq X$ and $\varepsilon > 0$ we say that X is ε -deformable into Y if there exists a continuous mapping $\varphi \colon X \times I \to X$ such that $\varphi(x,0) = x$, $\varphi(x,1) \in Y$ and the diameter of $\varphi(\{x\} \times I)$ is at most ε for every $x \in X$. The following proposition was proved in [Kra76, 1.1 and 1.3].

Proposition 22. Let X be a compact space such that for every $\varepsilon > 0$ there exists an absolute retract $Y \subseteq X$ for which X is ε -deformable into Y. Then X is an absolute retract.

By a result of [And67] the union of two Hilbert cubes, whose intersection is a Z-set in each of the cubes and which is homeomorphic to the Hilbert cube, is the Hilbert cube again. By the result of [Han78], even a weaker condition is enough to get the same conclusion:

Proposition 23. Let X be a space which is the union of two Hilbert cubes Q_1 and Q_2 . Suppose that $Q_1 \cap Q_2$ is a Hilbert cube which is a Z-set in Q_1 . Then X is a Hilbert cube.

It should be noted here that a space which is the union of two Hilbert cubes intersecting in a Hilbert cube may not be a Hilbert cube [She77].

Lemma 24. Let X be a compact metric space which is the union of Hilbert cubes Q, Q_1, Q_2, \ldots such that $Q_i \cap Q_j = \emptyset$ and $Q \cap Q_i$ is a Z-set in Q_i for every $i, j \in \mathbb{N}$, $i \neq j$. Suppose moreover that the diameter of Q_i tends to zero. Then X is homeomorphic to the Hilbert cube as well.

Proof. Let us denote $X_i = Q \cup Q_1 \cup \cdots \cup Q_i$ and observe that it is homeomorphic to the Hilbert cube for every $i \in \mathbb{N}$ by an inductive usage of Proposition 23. To make the same conclusion for X we use the Toruńczyk's theorem.

There is topological just one way, how to embed the Hilbert cube into itself as a Z-set. Hence every pair $(Q_i, Q_i \cap Q)$ is equivalent to $(Q \times I, Q \times \{0\})$ because $Q_i \cap Q$ is a Z-set in Q_i and it is homeomorphic to the Hilbert cube. Thus simply there is a homotopy $h_i \colon Q_i \times I \to Q_i$ such that $h_i(x,t) = x$ for t = 0 or $x \in Q_i \cap Q$ and $h_i(x,1) \in Q \cap Q_i$. Let us denote

$$s_i(x,t) = \begin{cases} x, & x \in X_i, \\ h_j(x,t), & x \in Q_j, & j > i. \end{cases}$$

Since the diameter of Q_i tends to zero, the diameters of $s_i(\{x\} \times I)$ are sufficiently small for large i. Hence for every $\varepsilon > 0$ it follows that X is ε -deformable into X_i for some $i \in \mathbb{N}$.

Moreover, for every $i \in \mathbb{N}$, X_i is an absolute retract. Hence X is an absolute retract by Proposition 22.

Let us argue that X has the disjoint cell property (see [vM01, p. 294]). Denote $r_i(x) = s_i(x, 1)$. Then $r_i \colon X \to X_i$ is a retraction. Suppose that $f, g \colon I^n \to X$ are continuous mappings and $\varepsilon > 0$. Then for sufficiently large $i \in \mathbb{N}$ diameters of Q_j are smaller than ε for $j \geq i$. Hence r_i is ε -close to identity on X. Since X_i is homeomorphic to the Hilbert cube it has the disjoint cell property and thus there are continuous mappings $f', g' \colon I^n \to X_i$ with disjoint images such that $r_i \circ f$ and f' as well as $r_i \circ g$ and g' are ε -close. It follows that by the triangle inequality that f and f' as well as g and g' are g-close. Thus g has the disjoint cell property. As mentioned at the beginning of the proof, by Toruńczyk's theorem it follows that g is homeomorphic to the Hilbert cube. \square

An equivalence relation E on a Borel subset Y of a Polish space X is said to be *countably separated* if there is a sequence $(Z_n)_{n=1}^{\infty}$ of E-invariant Borel subsets of Y such that for all $x, y \in Y$, the points x and y are E-equivalent if and only if the sets $\{n \in \mathbb{N} : x \in Z_n\}$ and $\{n \in \mathbb{N} : y \in Z_n\}$ are equal. A *transversal* for an equivalence relation $E \subseteq X \times X$ is a set $T \subseteq X$ whose intersection with every E-equivalence class is a one point set.

The following proposition by Burgess is a special kind of a selection theorem and it will serve as a useful tool to complete the Borel coding argument, which is by no means straightforward.

Proposition 25 ([Bur79]). Let G be a Polish group, X a Polish space and let α be a continuous action of G on X. Denote by E the orbit equivalence relation induced by α and let Y be an E-invariant Borel subset of X. Let E_Y be the restriction of E to Y and assume that E_Y is countably separated. Then there is a Borel transversal for E_Y .

In the next proposition, we denote by $\mathcal{K}_X(Q)$ the collection of subspaces of Q which are homeomorphic to X. It is known for a long time that this is always a Borel set [RN65].

Proposition 26. There is a Borel mapping $\gamma \colon \mathcal{K}_Q(Q) \to Inj(Q,Q)$ such that the image of $\gamma(R)$ equals to R.

Proof. Let G be the homeomorphism group of Q and let us consider the action α of G on $\mathrm{Inj}(Q,Q)$ given by $g\cdot h=h\circ g$. It follows that the corresponding orbit equivalence relation E induced by α satisfies that fEg if and only if the images of f and g are equal. Moreover E is countably separated as if we consider a countable base \mathcal{B} of Q and $Z_B=\{f\in\mathrm{Inj}(Q,Q)\colon Im(f)\cap B\neq\emptyset\}$ for $B\in\mathcal{B}$ then fEg if and only if $\{B\in\mathcal{B}\colon f\in Z_B\}=\{B\in\mathcal{B}\colon g\in Z_B\}$ and also the sets Z_B are clearly invariant with respect to E. By a straightforward application of Proposition 25 we get that there is a Borel transversal T of E. As the mapping $\chi\colon f\mapsto Im(f)$, $\mathrm{Inj}(Q,Q)\to\mathcal{K}(Q)$ is Borel (even continuous), the graph of χ is a Borel subset of $\mathrm{Inj}(Q,Q)\times\mathcal{K}(Q)$. As moreover T is a Borel subset of the domain of χ and χ is one-to-one on T it follows that the mapping $\gamma=(\chi|T)^{-1}$ has a Borel




Figure 1: The compactum Q_K

graph and thus it is a Borel mapping. Clearly for every $R \in \mathcal{K}_Q(Q)$ we get that $\gamma(R)$ is an embedding of Q into Q whose image equals R.

Some ideas for the proof of the following comes from the paper [KV20].

Theorem 27. The conjugacy of Hilbert cube homeomorphisms (or selfmaps) is Borel bireducible to $E_{G_{\infty}}$.

Proof. For one direction it is enough to prove that the homeomorphism equivalence relation of metrizable compacta is Borel reducible to conjugacy of Hilbert cube homeomorphisms because the first relation is Borel bireducible to $E_{G_{\infty}}$ by the main result of [Zie16]. To this end let

$$Q = \{x \in \ell_2 : 0 \le x_n \le 1/n\},\$$

$$Q' = Q \times I \times \{0\},\$$

$$Q'' = Q \times I \times [-1, 1],\$$

$$Q'^- = Q \times I \times [-1, 0]$$

and let $\|\cdot\|$ be the usual norm on ℓ_2 .

Let us fix a homotopy $H: \mathcal{K}(Q) \times I \to \mathcal{K}(Q)$ given by Proposition 21 for the case X = Q. Let us fix $K \in \mathcal{K}(Q)$. We want to find a homeomorphism f_K of a Hilbert

cube $Q_K \subseteq Q''$ such that the topological information about K is somehow encoded in the dynamics of f_K . Let $D_n^K = H(K, 2^{-n})$, $n \in \mathbb{N}$. Let ε_n be the minimum of 1/n and the smallest distance of different points in D_n^K . For every $d \in D_n^K$ fix a set

$$B_n^d = \{(x, 2^{-n}, 0) \in Q'' : ||d - x|| \le \varepsilon_n/3\}.$$

It follows that B_n^d is always homeomorphic to the Hilbert cube since it is affinely homeomorphic to an infinite dimensional compact convex subset of a Hilbert space [Kel31]. Let C_n^d be the cone in Q'' with base B_n^d and with the vertex $(d, 2^{-n}, 2^{-n}), d \in D_n^K, n \in \mathbb{N}$, i.e., the union of all segments with end points $(d, 2^{-n}, 2^{-n})$ and $p, p \in B_n^d$. The cone over the Hilbert cube is homeomorphic to the Hilbert cube [vM01, Theorem 1.7.5], which applies to C_n^d . Let $Q_K^m = Q'^- \cup \bigcup \{C_n^d \colon n \in \mathbb{N}, n \leq m, d \in D_n^K\}$ and $Q_K = \bigcup \{Q_K^m \colon m \in \mathbb{N}\}$ (see Figure 1).

Since $Q'^- \cap C_n^d = B_n^d$ is homeomorphic to the Hilbert cube, which is a Z-set in C_n^d , we inductively obtain by Lemma 24 that Q_K is a Hilbert cube.

Let $h(x) = \sqrt{x}, x \in I$ or any fixed homeomorphism of I with two fixed points 0, 1; and 1 being a locally attracting fixed point. We define

$$f_K(x) = \begin{cases} x, & x \in Q'^-, \\ ((1-h(t))a + h(t)d, 2^{-n}, 2^{-n}h(t)), & x = ((1-t)a + td, 2^{-n}, 2^{-n}t) \in C_n^d, \\ d \in D_n^K, t \in I, n \in \mathbb{N}. \end{cases}$$

All the points in Q'^- are fixed points for f_K and these are clearly not attracting. All the points in $\bigcup D_n^K$ are fixed points of f_K and these are attracting. There are no other fixed points of f_K . It follows that K is homeomorphic (or even equal) to the set of fixed points that are limits of attracting points but not attracting by itself (and thus defined only by dynamical notions). Hence if f_K and f_L are conjugate then K and L are homeomorphic, $K, L \in \mathcal{K}(Q)$.

On the other hand if K, L are homeomorphic compacta in Q then the sets $K \cup \bigcup_{n \in \mathbb{N}} D_n^K \times \{2^{-n}\}$ and $L \cup \bigcup_{n \in \mathbb{N}} D_n^L \times \{2^{-n}\}$ are homeomorphic by Lemma 20. This homeomorphism can be simply extended to a homeomorphism

$$\varphi\colon (K\times\{(0,0)\})\cup \left[\ \left| \{B_n^d(K)\colon d\in D_n^K, n\in\mathbb{N}\} \to (L\times\{(0,0)\})\cup \left[\ \right| \{B_n^d(L)\colon d\in D_n^L, n\in\mathbb{N}\}. \right] \right]$$

Both the sets in the domain and range of φ are Z-sets in Q'^- since these are closed subsets of the Z-set $Q' \times \{0\}$ [vM01, Lemma 5.1.2, Lemma 5.1.3]. Hence φ can be extended to a homeomorphism $\varphi' \colon Q'^- \to Q'^-$ [vM01, Theorem 5.3.7]. It remains to extend φ' linearly on the cones to obtain a homeomorphism φ'' . It follows that φ'' conjugates f_K and f_L . Note that we can identify f_K with its graph and thus it can be considered as a closed subspace of $Q'' \times Q''$. To verify that the mapping $\chi \colon \mathcal{K}(Q) \to \mathcal{K}(Q'' \times Q'')$, $K \mapsto f_K$ is Borel is a routine which is usually omitted in this type of proofs.

However, we are still not done, since f_K is defined on the topological copy of the Hilbert cube Q_K which differs when changing K. Let us consider the Borel mapping γ given by Proposition 26. We redefine the mapping f_K by conjugating it via $\gamma(Q_K)$ in the following way. The mapping $K \mapsto \gamma(Q_K)^{-1} \circ f_K \circ (\gamma(Q_K))$, $\mathcal{K}(Q) \to \mathcal{H}(Q)$ is the desired Borel reduction.

To conclude the proof it is enough to Borel reduce conjugacy of Hilbert cube maps to $E_{G_{\infty}}$. Consider structures of the form (Q,R) where R is a closed binary relation on Q. Two such structures (Q,R) and (Q,S) are said to be isomorphic if there is a homeomorphisms $\psi \colon Q \to Q$ for which $(\psi \times \psi)(R) = S$. By a fairly more general result [RZ18] it follows that such isomorphism equivalence relation is Borel reducible to $E_{G_{\infty}}$. There is a Borel (even continuous) reduction which takes a continuous map $f \colon Q \to Q$ and assigns $(Q, \operatorname{graph}(f))$ to it. Combining the two reductions we get the desired one.

5 Concluding remarks and questions

Let us summarize some of the results on the complexity of conjugacy equivalence relation in Table 1 in which we consider conjugacy equivalence relation of maps, homeomorphisms, and pointed transitive homeomorphisms of the interval, circle, Cantor set and Hilbert cube, respectively. Let us recall that a pointed dynamical system is a triple (X, f, x), where (X, f) is a dynamical system and $x \in X$. We say, that a pointed dynamical system (X, f, x) is transitive if the forward orbit of x in (X, f) is dense. Two pointed dynamical systems (X, f, x) and (Y, g, y) are called conjugate if there is a conjugacy of (X, f) and (Y, g) mapping x to y. We proceed by a series of simple notes as comments on Table 1.

	Homeomorphisms/maps	Pointed transitive homeomorphisms
Interval	$E_{S_{\infty}}$ [Hjo00], Theorem 17	Ø Note 29
Circle	$E_{S_{\infty}}$ Note 30	$E_{=}$ Note 31
Cantor set	$E_{S_{\infty}}$ [CG01]	$E_{=+}$ [Kay17b], Note 32
Hilbert cube	$E_{G_{\infty}}$ Theorem 27	? Question 33

Table 1: The complexity of conjugacy equivalence relation.

Note 28. Conjugacy of pointed transitive maps of the interval is smooth; indeed it is enough to assign to every pointed transitive dynamical system (I, f, x) the $\mathbb{N} \times \mathbb{N}$ matrix of true and false: $(f^m(x) < f^n(x))_{m,n \in \mathbb{N}}$ which determines f uniquely up to increasing conjugacy.

Note 29. There are no transitive homeomorphisms on the interval.

Note 30. The complexity result by Hjorth [Hjo00, Section 4.2] that conjugacy of interval homeomorphisms is Borel bireducible to $E_{S_{\infty}}$, remains true for circle homeomorphisms

simply by a modification of the original proof. A modification of the proof of Theorem 17 will give a similar result for circle maps. The same method as for the interval case can be used just by considering left or right local maxima and minima defined in an obvious way and then iterating this set forward and backward in a similar manner as in Notation 4. Thus conjugacy of circle selfmaps is Borel bireducible to the S_{∞} -universal orbit equivalence relation.

Note 31. Transitive homeomorphisms of the circle are well known to be conjugate to irrational rotations. Hence the rotation number is a complete invariant and hence conjugacy of (pointed) transitive homeomorphisms of the circle is Borel bireducible to the equality on irrationals (or on an uncountable Polish space).

Note 32. By a result of Kaya [Kay17b], conjugacy of pointed minimal homeomorphisms of the Cantor set is Borel bireducible to the equality of countable sets $E_{=+}$. Note that his proof works in the same vein for pointed transitive homeomorphisms of the Cantor set. Let us recall the main part of his construction in this case. Let X be the Cantor set and \mathcal{B} the collection of all clopen sets in X. To a pointed transitive system (X, f, x) we assign the collection

$$Ret(f, x) = \{Ret_B(f, x) : B \in \mathcal{B}\},\$$

where $Ret_B(f,x) = \{n \in \mathbb{Z} : f^n(x) \in B\}$. It can be verified that the mapping Φ defined as

$$\Phi(f,x) = (\mathsf{Ret}_\mathsf{B}(f,x) \colon B \in \mathcal{B}) \in \mathcal{P}(\mathbb{Z})^{\mathcal{B}}$$

is a reduction of pointed transitive Cantor maps to the equality of countable sets in $\mathcal{P}(\mathbb{Z})^{\mathcal{B}}$, i.e., (f,x) is conjugate to (g,y) if and only if $\mathsf{Ret}(f,x) = \mathsf{Ret}(g,y)$.

The following question is the missing part to complete Table 1.

Question 33. What is the complexity of conjugacy of transitive pointed Hilbert cube homeomorphisms (or maps)?

It was explained to us by Burak Kaya, that conjugacy equivalence relation of pointed transitive Hilbert cube homeomorphisms is a Borel relation [Kay17c]. The main reason is that every conjugacy of such systems preserves the distinguished point and thus it is automatically prescribed on a dense subset. Hence there is at most one conjugacy between such systems. Let us note that neither $E_{S_{\infty}}$ nor $E_{G_{\infty}}$ is Borel and thus these equivalence relations can not answer Question 33.

Since triangular maps i.e., maps $f\colon I^2\to I^2$ of the form f(x,y)=(g(x),h(x,y)) for continuous maps $g\colon I\to I$ and $h\colon I^2\to I$, lie in between one-dimensional and two dimensional and there is a gap in the complexity of the last two mentioned equivalence relations, the following question is natural.

Question 34. What is the complexity of conjugacy of triangular maps? Is it Borel bireducible to $E_{S_{\infty}}$ or to $E_{G_{\infty}}$?

Positive answer to the next question would provide a strengthening of Theorem 17.

Question 35. Is conjugacy of closed binary relations on the closed interval Borel reducible to the S_{∞} -universal orbit equivalence relation?

The answer to the preceding question is affirmative if Hjorth's conjecture is true.

References

- [And67] Richard D. Anderson. Topological properties of the Hilbert cube and the infinite product of open intervals. *Trans. Amer. Math. Soc.*, 126:200–216, 1967.
- [BS95] Jorge Buescu and Ian Stewart. Liapunov stability and adding machines. *Ergodic Theory Dynam. Systems*, 15(2):271–290, 1995.
- [Bur79] John P. Burgess. A selection theorem for group actions. *Pacific J. Math.*, 80(2):333–336, 1979.
- [CG01] Riccardo Camerlo and Su Gao. The completeness of the isomorphism relation for countable Boolean algebras. *Trans. Amer. Math. Soc.*, 353(2):491–518, 2001.
- [CG19] Cheng Chang and Su Gao. The complexity of the classification problems of finite-dimensional continua. *Topology Appl.*, 267:106876, 18, 2019.
- [Cle09] John D. Clemens. Isomorphism of subshifts is a universal countable Borel equivalence relation. *Israel J. Math.*, 170:113–123, 2009.
- [For18] Matthew Foreman. What is a Borel reduction? Notices Amer. Math. Soc., 65(10):1263–1268, 2018.
- [Gao09] Su Gao. Invariant descriptive set theory, volume 293 of Pure and Applied Mathematics (Boca Raton). CRC Press, Boca Raton, FL, 2009.
- [GJS16] Su Gao, Steve Jackson, and Brandon Seward. Group colorings and Bernoulli subflows. *Mem. Amer. Math. Soc.*, 241(1141):vi+241, 2016.
- [GvM93] Helma Gladdines and Jan van Mill. Hyperspaces of Peano continua of Euclidean spaces. Fund. Math., 142(2):173–188, 1993.
- [Han78] Michael Handel. On certain sums of Hilbert cubes. General Topology and Appl., 9(1):19–28, 1978.

- [Hjo00] Greg Hjorth. Classification and orbit equivalence relations, volume 75 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2000.
- [Kay17a] Burak Kaya. The complexity of the topological conjugacy problem for Toeplitz subshifts. *Israel J. Math.*, 220(2):873–897, 2017.
- [Kay17b] Burak Kaya. The complexity of topological conjugacy of pointed Cantor minimal systems. Arch. Math. Logic, 56(3-4):215–235, 2017.
- [Kay17c] Burak Kaya. On the complexity of topological conjugacy of compact metrizable G-ambits. arXiv e-prints, page arXiv:1706.09821, June 2017.
- [Kec95] Alexander S. Kechris. Classical descriptive set theory, volume 156 of Graduate Texts in Mathematics. Springer-Verlag, New York, 1995.
- [Kel31] Ott-Heinrich Keller. Die Homoiomorphie der kompakten konvexen Mengen im Hilbertschen Raum. *Math. Ann.*, 105(1):748–758, 1931.
- [Kra76] Józef Krasinkiewicz. On a method of constructing ANR-sets. An application of inverse limits. Fund. Math., 92(2):95–112, 1976.
- [KV20] Pavel Krupski and Benjamin Vejnar. The complexity of the homeomorphism relations on some classes of compacta. J. Symb. Log., pages 1–19, 2020.
- [Lor81] Edgar R. Lorch. On some properties of the metric subalgebras of l^{∞} . Integral Equations Operator Theory, 4(3):422–434, 1981.
- [Mel07] Julien Melleray. Computing the complexity of the relation of isometry between separable Banach spaces. *MLQ Math. Log. Q.*, 53(2):128–131, 2007.
- [MR17] Luca Motto Ros. Can we classify complete metric spaces up to isometry? *Boll. Unione Mat. Ital.*, 10(3):369–410, 2017.
- [RN65] C. Ryll-Nardzewski. On a Freedman's problem. Fund. Math., 57:273–274, 1965.
- [RZ18] Christian Rosendal and Joseph Zielinski. Compact metrizable structures and classification problems. J. Symb. Log., 83(1):165–186, 2018.
- [She77] R. B. Sher. The union of two Hilbert cubes meeting in a Hilbert cube need not be a Hilbert cube. *Proc. Amer. Math. Soc.*, 63(1):150–152, 1977.
- [ST17] Marcin Sabok and Todor Tsankov. On the complexity of topological conjugacy of Toeplitz subshifts. *Israel J. Math.*, 220(2):583–603, 2017.

- [Tho13] Simon Thomas. Topological full groups of minimal subshifts and just-infinite groups. In *Proceedings of the 12th Asian Logic Conference*, pages 298–313. World Sci. Publ., Hackensack, NJ, 2013.
- [vM01] Jan van Mill. The infinite-dimensional topology of function spaces, volume 64 of North-Holland Mathematical Library. North-Holland Publishing Co., Amsterdam, 2001.
- [Zie16] Joseph Zielinski. The complexity of the homeomorphism relation between compact metric spaces. Adv. Math., 291:635–645, 2016.