On a Speculated Relation Between Chvátal-Sankoff Constants of Several Sequences

M. Kiwi^{*} J. Soto[†]

Abstract

It is well known that, when normalized by n, the expected length of a longest common subsequence of d sequences of length n over an alphabet of size σ converges to a constant $\gamma_{\sigma,d}$. We disprove a speculation by Steele regarding a possible relation between $\gamma_{2,d}$ and $\gamma_{2,2}$. In order to do that we also obtain some new lower bounds for $\gamma_{\sigma,d}$, when both σ and d are small integers.

1 Introduction

String matching is one of the most intensively analyzed problems in computer science. Among string matching problems the longest common subsequence problem (LCS) stands out. This problem consists of finding the longest subsequence common to all strings in a set of sequences (often just two). The LCS problem is the basis of Unix's diff command, has applications in bioinformatics, and also arises naturally in remarkably distinct domains such as cryptographic snooping, the mathematical analysis of bird songs, and comparative genomics. In addition, the LCS problem offers a concrete basis for the illustration and benchmarking of mathematical methods and tools such as subadditive methods and martingale inequalities; see for example Steele's monograph [Ste86].

Although the LCS problem has been studied under many different contexts there are several issues concerning it that are still unresolved. The most prominent of the outstanding questions relating to the LCS problem concerns the length $L_{n,\sigma,d}$ of a LCS of d sequences of n characters chosen uniformly and independently over some alphabet of size σ . Subadditivity arguments yield that for fixed d and n going to infinity, the expected value of $L_{n,\sigma,d}$ normalized by n converges to a constant $\gamma_{\sigma,d}$. For $d, \sigma \geq 2$, the precise value of $\gamma_{\sigma,d}$ is unknown. The constant $\gamma_{2,2}$ is referred to as the Chvátal-Sankoff constant. The calculation of its exact value is an over 3 decades old open problem. The determination of its value has received a fair amount of attention, starting with the work of Chvátal and Sankoff [CS75], encompassing among others [Dek79, Ale94, DP95, Dan98, BYNGS99, Lue03], and is explicitly stated in several well known texts such as the ones by Waterman [Wat95, § 11.1.3], Steele [Ste96, p. 3], Pevzner [Pev00, p. 107], and Szpankowski [Szp00, p. 109]. To the best of our knowledge the current sharpest bounds on $\gamma_{2,2}$ are due to Lueker [Lue03] who established that $0.788071 \leq \gamma_{2,2} \leq 0.826280$.

The starting point for this investigation is the following comment by Steele [Ste86]:

"It would be of interest to relate c_3 to c^2 , and one is tempted to speculate that $c_3 = c^2$ (and more generally that $c_k = c^{k-1}$). Computational evidence does not yet rule this out."

Here, Steele uses c to denote the limiting value of the longest common subsequence of two random sequences of length n normalized by n as n goes to infinity, and in general, he uses c_k to denote the analogous constant for k sequences. However, it is unclear if in this comment he uses c and c_k to denote the constants $\gamma_{2,2}$ and $\gamma_{k,2}$ (i.e. specifically for the case of alphabet size 2) or if he is generically denoting the constants for arbitrary alphabet size. Dančík [Dan98] cites

^{*}Departamento de Ingeniería Matemática. Centro de Modelamiento Matemático (UMI 2807, CNRS), U. Chile. Web: www.dim.uchile.cl/~mkiwi. Gratefully acknowledges the support of CONICYT via FONDAP in Applied Mathematics and Anillo en Redes ACT08.

[†]Department of Mathematics, MIT. jsoto@math.mit.edu. Gratefully acknowledges the support of CONICYT via Anillo en Redes ACT08.

the previous statement as a conjecture by Steele using the second interpretation, i.e., as the claim that for all $d \ge 3$ and $\sigma \ge 2$,

$$\gamma_{\sigma,d} = \gamma_{\sigma,2}^{d-1} \,. \tag{1}$$

Dančík [Dan98, Theorem 2.1, Corollary 2.1] shows that for $d \ge 2$

$$1 \leq \liminf_{\sigma \to \infty} \sigma^{1-1/d} \gamma_{\sigma,d} \leq \limsup_{\sigma \to \infty} \sigma^{1-1/d} \gamma_{\sigma,d} \leq e \,.$$

Hence, if (1) was true, then for $\epsilon > 0$ and σ sufficiently large,

$$1 - \epsilon \le \sigma^{1 - 1/d} \gamma_{\sigma, d} = \sigma^{1 - 1/d} \gamma_{\sigma, 2}^{d - 1} \le \sigma^{1 - 1/d} \left(\frac{e(1 + \epsilon)}{\sqrt{\sigma}}\right)^{d - 1}.$$

Dančík's results disprove (1) by observing that for d > 2 one may choose σ large enough so as to make the rightmost term of the last displayed equation arbitrarily close to 0.

If we use the first interpretation of Steele's speculation quoted above, i.e., considering only the case of binary alphabets as we believe it was intended, then (1) is not invalidated by Dančík's work.

In [Ste86], Steele does not justify his speculation. The following non-rigorous argument gives some indication that one should expect that $\gamma_{2,3}$ is strictly bigger than $\gamma_{2,2}^2$. Indeed, let A_1 , A_2 and A_3 be three independently and uniformly chosen binary sequences of length n. For $i \neq j$ and very large values of n one knows that a longest common subsequence $\ell_{i,j}$ of sequences A_i and A_j would be of length approximately $\gamma_{2,2}n$. One would expect (although we can not prove it) that $\ell_{i,j}$ would behave like a uniformly chosen binary string of length $\gamma_{2,2}n$. Sequences $\ell_{1,2}$ and $\ell_{2,3}$ are clearly correlated. However, one might guess that the correlation is weak (again, we can certainly neither formalize nor prove such a statement). The previously stated discussion suggests that a longest common subsequence $\ell_{1,2,3}$ of $\ell_{1,2}$ and $\ell_{2,3}$ should be of length approximately $\gamma_{2,2}^2 n$. Since $\ell_{1,2,3}$ is clearly a longest common subsequence of A_1 , A_2 and A_3 , one is led to conclude that

$$\gamma_{2,3} \ge \gamma_{2,2}^2$$
 (2)

However, there are two good reasons why one suspects that this last inequality should be strict:

- Since ℓ_{2,3} has only a fraction of A₃'s length, one expects that a longest common subsequence of ℓ_{1,2} and A₃ is significantly larger than a longest common subsequence of ℓ_{1,2} and ℓ_{2,3}.
- The longest common subsequence of A_1 , A_2 and A_3 might arise by taking a longest common subsequence on sub-optimal common subsequences $\ell'_{1,2}$ and $\ell'_{2,3}$ of A_1 and A_2 , and A_2 and A_3 , respectively.

This work's main contribution is to show that the inequality in (2) is indeed strict.

In Section 2 we give a simple argument that proves that when σ is fixed and d is large the identity $\gamma_{\sigma,d} = \gamma_{\sigma,2}^{d-1}$ does not hold. The underlying argument is essentially an application of the probabilistic method. However, it might still be possible that the relation would hold for some specific values of σ and d. Of particular interest is the case of binary sequences, i.e. $\sigma = 2$. In Section 3 we show that even this weaker identity does not hold, i.e. that $\gamma_{2,3} \neq \gamma_{2,2}^2$. To achieve this goal, we rely on Lucker's [Lue03] U = 0.826280 upper bound on $\gamma_{2,3}$ and determine a lower bound on $\gamma_{2,3}$ which is strictly larger than $U^2 \geq \gamma_{2,2}^2$. The lower bound on $\gamma_{2,3}$ is obtained by an approach similar to the one used by Lucker [Lue03] to lower bound $\gamma_{2,2}$, although in our case we have to consider a non-binary alphabet. Aside from the extra notation needed to handle the cases $\sigma, d > 2$, our treatment is a straightforward generalization of the approach used by Lucker. (In fact, in order to keep the exposition as clear as possible we do not even use the optimization tweaks implemented by Lucker in order to take advantage of the symmetries inherent to the problem and objects that arise in its analysis.) We conclude with some final comments in Section 4.

2 Disproving $\gamma_{\sigma,d} = \gamma_{\sigma,2}^{d-1}$ for large d

We start this section by introducing some notation. Given strings A_1, \ldots, A_d of length n, we denote by $L(A_1, \ldots, A_d)$ the length of the longest common subsequence of all A_i 's. Let $U_{n,\sigma}$ be the distribution of sequences of length n whose

characters are chosen uniformly and independently from $\Sigma = \{1, \ldots, \sigma\}$. We denote by $L_{n,\sigma,d}$ the random variable $L(A_1, \ldots, A_d)$ when all the A_i are chosen according to $\mathcal{U}_{n,\sigma}$. Finally, we let $\gamma_{\sigma,d}$ denote the limit of $\mathbf{E}L_{n,\sigma,d}/n$ when $n \to \infty$ (the existence of this limit follows from standard subadditivity arguments [CS75]).

In what follows, we give a lower bound for $\gamma_{\sigma,d}$ that is independent of d. This bound is based on the following simple fact: If X is chosen according to $\mathcal{U}_{n,\sigma}$ and n is large, then the number of occurrences of a fixed character in Σ is roughly n/σ . Intuitively, this means that for a set of d random strings of (very large) length n, with very high probability a sequence formed by roughly $\lfloor n/\sigma \rfloor$ equal characters will be a common subsequence of all the d random strings.

Lemma 1 For all d and σ , we have $\gamma_{\sigma,d} \geq 1/\sigma$.

Proof: Let A_1, \ldots, A_d be d independent random strings chosen according to $\mathcal{U}_{n,\sigma}$. Let X_i denote the number of times the character $c \in \Sigma$ appears in A_i , and $X = \min\{X_1, \ldots, X_d\}$. The string c^X formed by X copies of the character c is a common subsequence of all X_i 's. It follows that $L(A_1, \ldots, A_d) \ge X$.

Each X_i is a binomial variable with parameter $p = 1/\sigma$. By a standard Chernoff bound [JŁR00, Remark 2.5] we have that for any $0 < \varepsilon < 1$,

$$\Pr[X_i \le (1 - \varepsilon)np] \le \exp(-2n(p\varepsilon)^2).$$

Applying Markov's inequality, and recalling that the X_i 's are independent, it follows that:

$$\mathbf{E}X \ge (1-\varepsilon)np\Pr[X \ge (1-\varepsilon)np] \ge (1-\varepsilon)np[1-\exp(-2n(p\varepsilon)^2)]^d.$$

Letting n be sufficiently large so that $[1 - \exp(-2n(p\varepsilon)^2)]^d \ge (1 - 2\varepsilon)/(1 - \varepsilon)$, we obtain $\mathbf{E}X \ge np(1 - 2\varepsilon)$. Therefore:

$$\frac{\mathbf{E}L_{n,\sigma,d}}{n} = \frac{\mathbf{E}L(A_1,\ldots,A_d)}{n} \ge \frac{\mathbf{E}X}{n} \ge (1-2\varepsilon)p = \frac{1-2\varepsilon}{\sigma}.$$

It follows that $\gamma_{\sigma,d} \ge (1-2\varepsilon)/\sigma$. Since this is true for any $\varepsilon > 0$, we conclude that $\gamma_{\sigma,d} \ge 1/\sigma$.

It is now easy to disprove that $\gamma_{\sigma,d} = \gamma_{\sigma,2}^{d-1}$ for large d. Indeed, since $\gamma_{\sigma,2} < 1$ [CS75], then $\lim_{d\to\infty} \gamma_{\sigma,2}^{d-1} = 0$. On the other hand, the previous lemma asserts that $\gamma_{\sigma,d} \ge 1/\sigma$ for all d, hence for d large enough, $\gamma_{\sigma,2}^{d-1} < \gamma_{\sigma,d}$.

In particular, for the case $\sigma = 2$, Lueker [Lue03] proved that $\gamma_{2,2} \leq U$ for U = 0.826280. Thus, for all $d \geq 5$, we have the strict inequality

$$\gamma_{2,2}^{d-1} \le (0.826280)^{d-1} < 1/2 \le \gamma_{2,d}$$

3 Disproving $\gamma_{2,3} = \gamma_{2,2}^2$

3.1 Diagonal common subsequence

As already mentioned, the best known provable lower bound for $\gamma_{2,2}$ found so far is due to Lueker [Lue03]. The starting point of Lueker's lower bound technique is a result by Alexander [Ale94] who related the expected length of the LCS of two random strings of the same length n, to the expected length of the LCS of two random strings whose lengths sum up to 2n. Below, we establish an analog of Alexander's result but for the case of d randomly chosen sequences.

Let C[j..k] denote the substring C[j]C[j+1]...C[k] formed by all the characters between the j-th and k-th positions of C. Given strings $A_1, ..., A_d$ of length at least n, we say that B is an n-diagonal common subsequence of $A_1, ..., A_d$ if B is a common subsequence of a set of prefixes of $A_1, ..., A_d$ whose lengths sum to n, i.e., if for some indices $i_1, ..., i_d$ such that $i_1 + \cdots + i_d = n$, the string B is a common subsequence of $A_1[1..i_1], ..., A_d[1..i_d]$.

Let $D_n(A_1, \ldots, A_d)$ denote the length of a longest *n*-diagonal common subsequence of the strings A_1, \ldots, A_d . We denote by $D_{n,\sigma,d}$ the random variable $D_n(A_1, \ldots, A_d)$ where the strings A_1, \ldots, A_d are chosen according to $\mathcal{U}_{n,\sigma}$.

The main objective of this section is to prove the following extension of a result of Alexander [Ale94, Proposition 2.4] for the d = 2 case:

Theorem 1 For all $n \ge d$,

$$d \cdot \mathbf{E} D_{n,\sigma,d} - d^{3/2} \sqrt{2n \ln n} \le \mathbf{E} L_{n,\sigma,d} \le \mathbf{E} D_{nd,\sigma,d}$$

In particular, for all σ there exists $\delta_{\sigma,d}$ such that:

$$\delta_{\sigma,d} = \lim_{n \to \infty} \frac{\mathbf{E} D_{n,\sigma,d}}{n} = \frac{\gamma_{\sigma,d}}{d}.$$

For the sake of clarity of exposition, before proving Theorem 1 we establish some intermediate results.

Lemma 2 For all n and d, $\mathbf{E}L_{n,\sigma,d} \leq \mathbf{E}D_{nd,\sigma,d}$.

Proof: Let A_1, \ldots, A_d be random strings independently chosen according to $\mathcal{U}_{nd,\sigma}$. Since a longest common subsequence of $A_1[1..n], \ldots, A_d[1..n]$ is also an *nd*-diagonal common subsequence of A_1, \ldots, A_d ,

$$L(A_1[1..n], \ldots, A_d[1..n]) \le D_{nd}(A_1, \ldots, A_d).$$

Taking expectation on both sides of the previous inequality yields the desired conclusion.

Lemma 3 For all $n \ge d$,

$$d \cdot \mathbf{E} D_{n,\sigma,d} - d^{3/2} \sqrt{2n \ln n} \le \mathbf{E} L_{n,\sigma,d}$$

Proof: Let A_1, \ldots, A_d be a list of words of length n. Note that if we change one character of any word in the list, then the values $L(A_1, \ldots, A_d)$ and $D_n(A_1, \ldots, A_d)$ will change by at most one unit. It follows that the random variables $L_{n,\sigma,d}$ and $D_{n,\sigma,d}$ (seen as functions from $(\Sigma^n)^d$ to **R**) are both 1-Lipschitz. Applying Azuma's inequality (as treated in for example [JŁR00, § 2.4]) we get:

$$\Pr\left[D_{n,\sigma,d} \le \mathbf{E}D_{n,\sigma,d} - \sqrt{n/2}\right] \le \exp\left(-\frac{2(n/2)}{nd}\right) = e^{-1/d} < \frac{d}{d+1},$$

where the last inequality holds since $e^{-x} < 1/(x+1)$ for all x > 0.

Let $\lambda = \mathbf{E}D_{n,\sigma,d} - \sqrt{n/2}$. Since $D_{n,\sigma,d} > \lambda$ implies that there are positive indices i_1, \ldots, i_d such that $i_1 + \cdots + i_d = n$ and $L(A_1[1..i_1], \ldots, A_d[1..i_d]) \ge \lambda$,

$$\Pr[D_{n,\sigma,d} > \lambda] \le \sum_{\substack{0 < i_1, \dots, i_d < n, \\ i_1 + \dots + i_d = n}} \Pr[L(A_1[1..i_1], \dots, A_d[1..i_d]) > \lambda].$$

Let I be the number of summands in the right hand side. Note that $I = \binom{n-1}{d-1}$ since it counts the the number of ways of partitioning n into d positive summands. It follows that there exist positive j_1, \ldots, j_d summing to n such that:

$$\Pr[L(A_1[1..j_1], \dots, A_d[1..j_d]) > \lambda] > \frac{1}{I} \left(1 - \frac{d}{d+1}\right) = \frac{1}{I(d+1)}.$$

Note that the distribution of the random variable $L(A_1[1..j_1], \ldots, A_d[1..j_d])$ is the same as the distribution of $L(A_1[1..j_{\tau(1)}], \ldots, A_d[1..j_{\tau(d)}])$ for any permutation $\tau : [d] \to [d]$. It is also easy to see that the distribution of $L(A_1[a_1..b_1], \ldots, A_d[a_d..b_d])$ and $L(A_1[a'_1..b'_1], \ldots, A_d[a'_d..b'_d])$ are the same when $b_m - a_m = b'_m - a'_m$ for all $1 \le m \le d$.

Now, let τ be the cyclic permutation $(12 \dots d)$ and for $0 \le m \le d-1$ let \mathcal{E}_m denote the event

$$L\left(A_{1}\left[\sum_{l=0}^{m-1} j_{\tau^{l}(1)} + 1 \dots \sum_{l=0}^{m} j_{\tau^{l}(1)}\right], \dots, A_{d}\left[\sum_{l=0}^{m-1} j_{\tau^{l}(d)} + 1 \dots \sum_{l=0}^{m} j_{\tau^{l}(d)}\right]\right) > \lambda.$$

In particular, \mathcal{E}_0 is the event $\{L(A_1[1..j_1], \ldots, A_d[1..j_d]) > \lambda\}$ whose probability was bounded above. Note that the events $\mathcal{E}_0, \ldots, \mathcal{E}_{d-1}$ are equiprobable. Since each of the \mathcal{E}_m 's depends on a different set of characters, they are

independent. Moreover, if $\mathcal{E}_0, \ldots, \mathcal{E}_{d-1}$ simultaneously occur, then by concatenating the common subsequences of each block of characters we get that $L(A_1, \ldots, A_d) > d\lambda$. Hence,

$$\left(\frac{1}{I(d+1)}\right)^d < \prod_{m=0}^{d-1} \Pr[\mathcal{E}_m] = \Pr[\mathcal{E}_0, \mathcal{E}_1, \dots, \mathcal{E}_{d-1}] \le \Pr[L_{n,\sigma,d} > d\lambda].$$
(3)

Applying Azuma's inequality again, we have:

$$\Pr\left[L_{n,\sigma,d} \ge \mathbf{E}L_{n,\sigma,d} + \sqrt{\frac{nd^2\ln(I(d+1))}{2}}\right] \le \left(\frac{1}{I(d+1)}\right)^d.$$
(4)

Combining (3) and (4) and recalling that $\lambda = \mathbf{E}D_{n,\sigma,d} - \sqrt{n/2}$ we obtain:

$$\Pr\left[L_{n,\sigma,d} \ge \mathbf{E}L_{n,\sigma,d} + \sqrt{\frac{nd^2\ln(I(d+1))}{2}}\right] < \Pr\left[L_{n,\sigma,d} > d\mathbf{E}D_{n,\sigma,d} - d\sqrt{\frac{n}{2}}\right]$$

Hence:

$$\mathbf{E}L_{n,\sigma,d} + \sqrt{\frac{nd^2\ln(I(d+1))}{2}} \ge d\mathbf{E}D_{n,\sigma,d} - d\sqrt{\frac{n}{2}}$$

Since $2 \leq d \leq n$, $(d+1)I = (d+1) \binom{n-1}{d-1} \leq n^d$, and so:

$$d\mathbf{E}D_{n,\sigma,d} \le \mathbf{E}L_{n,\sigma,d} + d\sqrt{\frac{n}{2}} + \sqrt{\frac{nd^2\ln(I(d+1))}{2}} \le \mathbf{E}L_{n,\sigma,d} + d^{3/2}\sqrt{2n\ln(n)}$$

Proof of Theorem 1: Lemmas 2 and 3 already give the bounds on $\mathbf{E}L_{n,\sigma,d}$.

To complete the proof we need to show that $\lim_{n\to\infty} \mathbf{E}D_{n,\sigma,d}/n$ exists and that its value is $\gamma_{\sigma,d}/d$. By Lemmas 2 and 3 we have:

$$\mathbf{E}L_{n,\sigma,d} \le \mathbf{E}D_{nd,\sigma,d} \le \frac{1}{d}\mathbf{E}L_{nd,\sigma,d} + d^{1/2}\sqrt{2nd\ln(nd)}.$$

Dividing by n, it follows that $\lim_{n\to\infty} \mathbf{E}D_{nd,\sigma,d}/n = \gamma_{\sigma,d}$. Furthermore, $\mathbf{E}D_{n,\sigma,d}$ is non decreasing in n, so:

$$\frac{\lfloor n/d \rfloor}{n/d} \cdot \frac{\mathbf{E}D_{d\lfloor n/d \rfloor, \sigma, d}}{\lfloor n/d \rfloor} \le \frac{\mathbf{E}D_{n, \sigma, d}}{n/d} \le \frac{\lceil n/d \rceil}{n/d} \cdot \frac{\mathbf{E}D_{d\lceil n/d \rceil, \sigma, d}}{\lceil n/d \rceil}.$$

Since both the left hand side and right hand side terms above converge to $\gamma_{\sigma,d}$ when $n \to \infty$, the middle term also converges to that value, and so $\lim_{n\to\infty} \mathbf{E}D_{n,\sigma,d}/n = \gamma_{\sigma,d}/d$ as claimed.

3.2 Longest common subsequence of two words over a binary alphabet

In this section we describe Lueker's [Lue03] approach for finding a lower bound on $\gamma_{d,\sigma}$ when $d = \sigma = 2$. Later on, we will generalize Lueker's technique to the cases of arbitrary d and σ .

Let X_1 and X_2 be two random sequences chosen from $U_{n,2}$, i.e. strings of length n such that all their characters are chosen uniformly and independently from the binary alphabet $\{0, 1\}$. Lucker defines, for any two strings A and Bover the binary alphabet, the quantity

$$W_n(A,B) = \mathbf{E}\left[\max_{i+j=n} L(AX_1[1..i], BX_2[1..j])\right]$$

Informally, $W_n(A, B)$ represents the expected length of a LCS of two strings with prefixes A and B respectively and suffixes formed by uniformly and independently choosing n characters in $\{0, 1\}$. It is easy to see that $W_n(A, B)$

behaves as $D_{n,2,2}$ as $n \to \infty$. Hence, applying Alexander's d = 2 version of Theorem 1, Lucker observes that for all $A, B \in \{0, 1\}^*$,

$$\gamma_{2,2} = \lim_{n \to \infty} \frac{W_{2n}(A,B)}{n}$$

A natural idea is to approximate $\gamma_{2,2}$ by $W_{2n}(A, B)/n$. Fix the length $l \in \mathbb{N}$ of the strings A and B and denote by w_n the 2^{2l} dimensional vector whose coordinates correspond to the values $W_n(A, B)$ when A and B vary over all binary sequences of length l. For example, when l = 2 the vector w_n has the following form:

$$w_n = \begin{pmatrix} w_n[00,00] \\ w_n[00,01] \\ \vdots \\ w_n[11,10] \\ w_n[11,11] \end{pmatrix} = \begin{pmatrix} W_n(00,00) \\ W_n(00,01) \\ \vdots \\ W_n(11,10) \\ W_n(11,11) \end{pmatrix}$$

Lueker established a lower bound for each component of w_n as a function of the components of w_{n-1} and w_{n-2} . To reproduce that lower bound, we need to introduce some more notation. If $A = A[1]A[2] \dots A[l]$ is a sequence of length $l \ge 2$, let h(A) denote the *head* of A, i.e. its first character, and let T(A) denote its *tail*, i.e. the substring obtained from A by removing its head. In other words, h(A) = A[1] and T(A) = A[2...l]. It is easy to see that the following relations among w_n, w_{n-1} and w_{n-2} hold:

• If
$$h(A) = h(B)$$
, then

$$w_n[A, B] \ge 1 + \frac{1}{4} \sum_{(c,c') \in \{0,1\}^2} w_{n-2}[T(A)c, T(B)c'].$$

• If $h(A) \neq h(B)$, then

$$w_n[A, B] \ge \frac{1}{2} \max\left\{ \sum_{c \in \{0,1\}} w_{n-1}[T(A)c, B], \sum_{c \in \{0,1\}} w_{n-1}[A, T(B)c] \right\}.$$

Using the previous inequalities one can define a function $F : \mathbf{R}^{2^{2l}} \times \mathbf{R}^{2^{2l}} \to \mathbf{R}^{2^{2l}}$ such that for all $n \ge 2$, we have $w_n \ge F(w_{n-1}, w_{n-2})$. Furthermore, the function F can be decomposed in two simpler functions F_{\pm} and F_{\neq} such that if Π_{\pm} and Π_{\neq} are the projections of the vectors onto the coordinates corresponding to the pairs of words with the same and different heads respectively, then:

$$\Pi_{=}(w_n) \ge F_{=}(w_{n-2}), \quad \text{and} \quad \Pi_{\neq}(w_n) \ge F_{\neq}(w_{n-1}).$$

It might be useful to see some examples of these transformations. For instance, to obtain a lower bound of $w_n[001, 011]$, one considers:

$$w_n[001, 011] \ge F_{=}(w_{n-2})[001, 011]$$

= 1 + $\frac{1}{4}(w_{n-2}[010, 110] + w_{n-2}[010, 111] + w_{n-2}[011, 110] + w_{n-2}[011, 111]).$

And to bound $w_n[001, 111]$,

$$w_n[001, 111] \ge F_{\ne}(w_{n-1})[001, 111]$$

= $\frac{1}{2} \max \{ w_{n-1}[010, 111] + w_{n-1}[011, 111], w_{n-1}[001, 110] + w_{n-1}[001, 111] \}$

3.3 Longest common subsequence of d words over general alphabets

In this section we extend Lueker's lower bound arguments as described in the previous section to the general case of d strings whose characters are uniformly and independently chosen over an alphabet of size σ .

Let X_1, \ldots, X_d be a collection of d independent random strings chosen according to $\mathcal{U}_{n,\sigma}$ and let A_1, \ldots, A_d be a collection of d finite sequences over the same alphabet. We now consider:

$$W_n(A_1,...,A_d) = \mathbf{E}\left[\max_{i_1+...+i_d=n} L(A_1X_1[1..i_1],...,A_dX_d[1..i_d])\right]$$

This quantity represents the expected length of a LCS of d words with prefixes A_1, \ldots, A_d respectively and d suffixes whose lengths sum up to n and whose characters are uniformly and independently chosen in $\Sigma = \{1, \ldots, \sigma\}$. Since $W_n(A_1, \ldots, A_d)$ and $D_{n,\sigma,d}$ behave similarly as $n \to \infty$, Theorem 1 implies that for all A_1, \ldots, A_d ,

$$\gamma_{\sigma,d} = \lim_{n \to \infty} \frac{W_{nd}(A_1, \dots, A_d)}{n}.$$
(5)

Just as in the d = 2 case, fix $l \in \mathbb{N}$ and denote by w_n the σ^{ld} dimensional vector whose coordinates are all the values of $W_{nd}(A_1, \ldots, A_d)$ when A_1, \ldots, A_d vary over all sequences in Σ^l . We again seek a lower bound for w_n as a function of vectors w_m , with m < n.

It is easy to see that if all the strings A_1, \ldots, A_d start with the same character, then:

$$w_n[A_1, \dots, A_d] \ge 1 + \frac{1}{|\Sigma^d|} \sum_{\vec{c} \in \Sigma^d} w_{n-d}[T(A_1)c(1), T(A_2)c(2), \dots, T(A_d)c(d)].$$

Informally, the previous inequality asserts that if all the words start with the same character then the expected length of the LCS of all of them, allowing n random extra characters, is at least 1 (the first character) plus the average of the expected length of the LCS of the words obtained by eliminating the first character and "borrowing" d of the n random characters.

If not all the words start with the same character, we can still find a lower bound, but to write it down we need to introduce some additional notation. For any two sets X and Y we follow the standard convention of denoting by Y^X the set of all mappings from X to Y. Also, for a d-tuple of strings $A = (A_1, \ldots, A_d)$ and $z \in \Sigma$ we denote by $N_z(A)$ the set of indices $j \in \{1, \ldots, d\}$ such that A_j 's head is not equal to z, i.e. to the set of string indices not starting with z. For a mapping $c : N_z(A) \to \Sigma$ we define $\tau_z(A, c)$ as the the d-tuple of strings obtained from A by replacing each string A_i that does not start with z by the sequence obtained by eliminating its first character and adding the character c(i) at its tail. Formally, $\tau_z(A, c) = (A'_1, \ldots, A'_d)$ where

$$A'_{i} = \begin{cases} A_{i}, & \text{if } h(A_{i}) = z, \\ T(A_{i})c(i), & \text{if } h(A_{i}) \neq z. \end{cases}$$

A crucial fact is that for a d-tuple of strings A, if its coordinates do not all start with the same character, then

$$w_n[A] \ge \max_{z \in \Sigma} \frac{1}{|\Sigma^{N_z(A)}|} \sum_{c \in \Sigma^{N_z(A)}} w_{n-|N_z(A)|}[\tau_z(A, c)].$$

Informally, each term over which the maximum is taken corresponds to the expected length of the LCS of the strings one would obtain by disregarding all first characters of sequences not starting with z, and concatenating to the tail of these strings an element randomly chosen over the alphabet Σ .

For the sake of illustration, consider the following example of the derived inequalities when $\sigma = 2$ and d = 4:

$$w_n[001, 011, 101, 001] \ge \max\left\{\frac{1}{2} \sum_{c \in \{0,1\}^{\{3\}}} w_{n-1}[001, 011, 01c(3), 001], \\ \frac{1}{2^3} \sum_{c \in \{0,1\}^{\{1,2,4\}}} w_{n-3}[01c(1), 11c(2), 101, 01c(4)]\right\}$$

In the previous example only the third string over which w_n is evaluated does not start with 0. Hence, the first term over which the maximum is taken is the average of the values of w_{n-1} evaluated at the two possible 4-tuples of strings obtained from A by removing the initial 1 from the third string and adding a 0 or 1 final character. On the other hand, w_n is evaluated at three strings that do not start with a 1. Hence, the second term over which the maximum is taken is the average of the values of w_{n-3} over all the 4-tuples of strings obtained from A by removing all the initial 0's and adding a 0 or 1 final character to those same strings.

Expressing all the derived inequalities in vector form we have that there is a function $F : (\mathbf{R}^{\sigma^{ld}})^d \to \mathbf{R}^{\sigma^{ld}}$ such that

$$w_n \ge F(w_{n-1}, w_{n-2}, \dots, w_{n-d}).$$
 (6)

For the ensuing discussion it will be convenient to rewrite F in an alternative way. For each $z \in \Sigma$ we define the linear transformation $F_z : (\mathbf{R}^{\sigma^{ld}})^d \to \mathbf{R}^{\sigma^{ld}}$ such that

$$F_{z}(v_{1},\ldots,v_{d})[A] = \begin{cases} \frac{1}{|\Sigma^{N_{z}(A)}|} \sum_{c \in \Sigma^{N_{z}(A)}} v_{|N_{z}(A)|}[\tau_{z}(A,c)], & \text{if } |N_{z}(A)| \neq 0, \\ 0, & \text{if } |N_{z}(A)| = 0. \end{cases}$$
(7)

Then, if we let $b \in \mathbf{R}^{\sigma^{ld}}$ be the vector with value 1 in the coordinates associated to *d*-tuples of strings of length *l* starting all with the same character and 0 in the rest of the coordinates, *F* can be expressed as:

$$F(v_1, \dots, v_d) = b + \max_{z \in \Sigma} F_z(v_1, \dots, v_d).$$

$$\tag{8}$$

3.4 Finding a lower bound for $\gamma_{\sigma,d}$

In the preceding section we established that for any *d*-tuple of strings $A = (A_1, \ldots, A_d)$, each of length *l*, we have $\gamma_{\sigma,d} = \lim_{n \to \infty} w_{nd}[A]/n$. To lower bound this latter quantity one is tempted to try the following approach: (1) For a fixed word length *l*, compute explicitly w_0, \ldots, w_{d-1} , and, (2) Define a new sequence of vectors $(v_n)_{n \in \mathbb{N}}$ as $v_i = w_i$ for $0 \le i \le d-1$, and then iteratively define $v_n = F(v_{n-1}, v_{n-2}, \ldots, v_{n-d})$, for all $n \ge d$. Since *F* is monotone and by (6), we have that $v_n \le w_n$ for every $n \in \mathbb{N}$. It is natural to fix an arbitrary *d*-tuple of strings $A = (A_1, \ldots, A_d)$ and estimate a lower bound for $\gamma_{\sigma,d}$ by $\lim_{n\to\infty} v_{nd}[A]/n$ for large enough *n*.

Unfortunately, for the approach discussed in the previous paragraph to work one would need to determine for which values of n the quantity $v_{nd}[A]/n$ is effectively a lower bound for $\gamma_{\sigma,d}$. Indeed, $v_{nd}[A]/n$ does not even need to be increasing and $w_{nd}[A]/n$ equals $\gamma_{\sigma,d}$ only in the limit when $n \to \infty$. We will pursue a different approach that relies on the next lemma which is a generalization of an observation by Lueker [Lue03] for the $d = \sigma = 2$ case.

Lemma 4 Let $\mathcal{F} : (\mathbf{R}^{\sigma^{ld}})^d \to \mathbf{R}^{\sigma^{ld}}$ be a transformation that satisfies the following properties:

- 1. Monotonicity: If the inequality $(v_1, v_2, ..., v_d) \le (w_1, w_2, ..., w_d)$ holds component-wise, then the inequality $\mathcal{F}(v_1, v_2, ..., v_d) \le \mathcal{F}(w_1, w_2, ..., w_d)$ also holds component-wise.
- 2. Translation invariance: Let 1 be the vector of ones in $\mathbf{R}^{\sigma^{ld}}$ and $\vec{1} = (1, ..., 1)$ be the vector of ones in $(\mathbf{R}^{\sigma^{ld}})^d$. Then, for any $r \in \mathbf{R}$ and for all $(v_1, v_2, ..., v_d) \in (\mathbf{R}^{\sigma^{ld}})^d$,

$$\mathcal{F}((v_1, v_2, \dots, v_d) + r\mathbf{\vec{1}}) = \mathcal{F}(v_1, \dots, v_d) + r\mathbf{1}.$$

3. *Feasibility:* There exists a feasible triplet for \mathcal{F} , *i.e.* $a(u, r, \varepsilon)$ with $u \in \mathbf{R}^{\sigma^{ld}}$, $r \in \mathbf{R}$, and $0 \le \varepsilon \le r$ such that:

$$\mathcal{F}(u+(d-1)r\mathbf{1},\ldots,u+2r\mathbf{1},u+r\mathbf{1},u) \ge u+(dr-\varepsilon)\mathbf{1}$$

Then, for any sequence $(v_n)_{n \in \mathbb{N}}$ of vectors in $\mathbb{R}^{\sigma^{ld}}$ such that $v_n \geq \mathcal{F}(v_{n-1}, \ldots, v_{n-d})$ for all $n \geq d$, there exists a vector u_0 in $\mathbb{R}^{\sigma^{ld}}$ such that for all $n \geq 0$,

$$v_n \ge u_0 + n(r - \varepsilon)\mathbf{1}.\tag{9}$$

Proof: Let \mathcal{F} be a transformation satisfying the hypothesis of the lemma and (u, r, ε) a feasible triplet for \mathcal{F} . Let $(v_n)_{n \in \mathbb{N}}$ be a sequence of vectors as in the lemma's statement and let $\alpha \in \mathbb{R}$ be large enough so that for all $j \leq d-1$,

$$v_j + \alpha \mathbf{1} \ge u + j(r - \varepsilon) \mathbf{1}.$$

For example, set α to be the largest component of the vector $\max_{0 \le j \le d-1} (u + j(r-\varepsilon)\mathbf{1} - v_j)$.

Note that $u_0 = u - \alpha \mathbf{1}$ satisfies (9) for all $n \le d - 1$. We will prove by induction that this holds for all $n \in \mathbf{N}$. Suppose that (9) holds up to n - 1. Using the inductive hypothesis we have:

$$\begin{aligned} &(v_{n-1},\ldots,v_{n-d}) \\ &\geq (u_0+(n-1)(r-\varepsilon)\mathbf{1},\ldots,u_0+(n-j)(r-\varepsilon)\mathbf{1},\ldots,u_0+(n-d)(r-\varepsilon)\mathbf{1}) \\ &= (u+(d-1)r\mathbf{1},\ldots,u+(d-j)r\mathbf{1}+(j-1)\varepsilon\mathbf{1},\ldots,u+(d-1)\varepsilon\mathbf{1}) + \\ &((n-d)(r-\varepsilon)-(d-1)\varepsilon-\alpha)\vec{\mathbf{1}} \\ &\geq (u+(d-1)r\mathbf{1},\ldots,u+(d-j)r\mathbf{1},\ldots,u) + ((n-d)(r-\varepsilon)-(d-1)\varepsilon-\alpha)\vec{\mathbf{1}}. \end{aligned}$$

Evaluating \mathcal{F} at the terms on both sides of the previous inequality we get, by monotonicity and translation invariance, that

$$v_n \ge \mathcal{F}(v_{n-1}, \dots, v_{n-d})$$

$$\ge \mathcal{F}(u + (d-1)r\mathbf{1}, \dots, u + (d-j)r\mathbf{1}, \dots, u) + ((n-d)(r-\varepsilon) - (d-1)\varepsilon - \alpha)\mathbf{1}.$$

Since (u, r, ε) is a feasible triplet, it follows that:

$$v_n \ge u + (dr - \varepsilon)\mathbf{1} + ((n-d)(r-\varepsilon) - (d-1)\varepsilon - \alpha)\mathbf{1}$$

= $u - \alpha \mathbf{1} + n(r-\varepsilon)\mathbf{1} = u_0 + n(r-\varepsilon)\mathbf{1}.$

This completes the proof.

From F's definition it easily follows that F is monotone and invariant under translations. If we find a feasible triplet (u, r, ε) for F then, by Lemma 4, we can conclude that the sequence of vectors $(w_n)_{n \in \mathbb{N}}$ satisfy $w_n \ge u_0 + n(r - \varepsilon)\mathbf{1}$ for all n. It follows from (5) that:

$$\gamma_{\sigma,d} \ge d(r-\varepsilon).$$

The key point we are trying to make is that in order to establish a good lower bound for $\gamma_{\sigma,d}$ one only needs to exhibit a good feasible triplet, namely one such that $(r - \varepsilon)$ is as large as possible.

Empirically, one observes that for any set of initial vectors v_0, \ldots, v_{d-1} , if one makes $v_{n+d} = F(v_{n+d-1}, \ldots, v_n)$ for all $n \in \mathbb{N}$, then the sequence $(v_n)_{n \in \mathbb{N}}$ is such that v_n/n seems to converge to a vector with all its components taking the same value. In fact, one observes that for large values of n the vectors v_n and v_{n+1} differ essentially by a constant (independent of n) times the all ones vector. Roughly, there exists a real value r such that $v_{n+1} - v_n$ is approximately $r\mathbf{1}$ for all large enough n. Since, by definition $v_{n+d} = F(v_{n+d-1}, \ldots, v_{n+1}, v_n)$, this implies that

$$F(v_n + (d-1)r\mathbf{1}, v_n + (d-2)r\mathbf{1}, \dots, v_n + r\mathbf{1}, v_n) \sim v_n + dr\mathbf{1}$$

It follows that one possible approach to find a feasible triplet is to consider an n large enough so that the difference between v_n and v_{n-1} is essentially a constant times the all ones vector. Then, set $u = v_n$, and define r as the maximum value such that $v_n - v_{n-1} \ge r\mathbf{1}$ and ε as the minimum possible value such that the triplet (u, r, ε) is feasible for F. The following result validates the approach just described.

Lemma 5 Let $\mathcal{F}: (\mathbf{R}^{\sigma^{ld}})^d \to \mathbf{R}^{\sigma^{ld}}$ be a monotone and translation invariant transformation. Let $v_0, \ldots, v_{d-1} \in \mathbf{R}^{\sigma^{ld}}$ and $v_{n+d} = \mathcal{F}(v_{n+d-1}, \ldots, v_{n+1}, v_n)$ for all $n \in \mathbf{N}$. If for some $r \in \mathbf{R}$, $n_0 \ge 1$ and $\varepsilon > 0$ we have $||v_{n+1} - v_n - r\mathbf{1}||_{\infty} \le \varepsilon/2d$ for all $n \in \{n_0, \ldots, n_0+d-1\}$, then $(v_{n_0}, r, \varepsilon)$ is a feasible triplet for \mathcal{F} .

Proof: First, observe that the monotonicity and translation invariance property of \mathcal{F} implies that

$$||\mathcal{F}(x_0,\ldots,x_{d-1}) - \mathcal{F}(y_0,\ldots,y_{d-1})||_{\infty} \le \max_{i=0,\ldots,d-1} ||x_i - y_i||_{\infty}$$

Let $u = v_{n_0}$ and note that $||v_{n_0+i} - (u + ir\mathbf{1})||_{\infty} \le i\varepsilon/2d < \varepsilon/2$ for $0 \le i \le d$. Hence, by definition of v_{n_0+d} ,

$$||v_{n_0+d} - \mathcal{F}(u + (d-1)r\mathbf{1}, u + (d-2)r\mathbf{1}, \dots, u + r\mathbf{1}, u)||_{\infty} \le \varepsilon/2.$$

Since $||v_{n_0+d} - (u + dr\mathbf{1})||_{\infty} \le \varepsilon/2$ it follows that

$$||(u+dr\mathbf{1}) - \mathcal{F}(u+(d-1)r\mathbf{1}, u+(d-2)r\mathbf{1}, \dots, u+r\mathbf{1}, u)||_{\infty} \le \varepsilon$$

In other words, (u, r, ε) is a feasible triplet for \mathcal{F} .

It is easy to check that F satisfies the hypothesis of Lemma 5. This justifies, together with the empirical observation that $v_{n+1} - v_n$ is approximately $r\mathbf{1}$ for large values of n, the general approach described in this section for finding a feasible triplet for F, and thus a lower bound for $\gamma_{\sigma,d}$. It is important to stress here that there is no need to prove the convergence of v_n/n to $r\mathbf{1}$ in order to establish the lower bound $\gamma_{\sigma,d} \ge d(r - \varepsilon)$. We only need to find a feasible triplet (u, r, ε) for F. The characteristics of F, empirical observations and Lemma 5, efficiently lead to such feasible triplets.

3.5 Implementation and results. New bounds

In this section we describe the procedure we implemented in order to find a feasible triplet (u, r, ε) for F and, as a corollary, a lower bound for $\gamma_{\sigma,d}$. The procedure is called FEASIBLETRIPLET, it is parameterized in terms of the number of sequences d and the alphabet Σ , and its pseudocode is given in Algorithm 1. In order to implement F we

Algorithm 1 Procedure for computing a feasible triple for F	
1: procedure FEASIBLETRIPLET _{d,Σ} (l, n)	$\triangleright l \in \mathbf{N}$ parameter, $n \in \mathbf{N}$ iteration steps
2: for $i = 0,, d - 1$ do	
3: $v_i \leftarrow 0$	\triangleright Where 0 denotes the vector of zeros in $\mathbf{R}^{\sigma^{ld}}$
4: end for	
5: $(u, r, \varepsilon) \leftarrow (v_0, 0, 0)$	
6: for $i = d,, n$ do	
7: $v_i \leftarrow F(v_{i-1}, v_{i-2}, \dots, v_{i-d})$	
8: $R \leftarrow \max_{A \in (\Sigma^l)^d} (v_i - v_{i-1})[A]$	
9: $W \leftarrow v_i + dR1 - F(v_i + (d-1)R1, \dots, v_i + R1, v_i)$	
10: $E \leftarrow \max\{0, \max_{A \in (\Sigma^l)^d} W[A]\}$	
11: if $R - E \ge r - \varepsilon$ then	
12: $(u, r, \varepsilon) \leftarrow (v_i, R, E)$	
13: end if	
14: end for	
15: return (u, r, ε)	
16: end procedure	

rely on the characterization given by (7) and (8). Since the F_z 's are linear transformations, they can be represented as matrices. This allows for fast evaluation of the F_z 's, but requires a prohibitively large amount of main memory for all but small values of σ , l and d. In order to optimize memory usage, we use the fact that by distinguishing (7) according to the cardinality of $N_z(A)$ where $A \in (\Sigma^l)^d$, F_z can be written as:

$$F_{z}(v_{1},\ldots,v_{d}) = \frac{1}{\sigma^{1}}F_{z,1}(v_{1}) + \ldots + \frac{1}{\sigma^{d}}F_{z,d}(v_{d}),$$

where

$$F_{z,i}(v_i)[A] = \begin{cases} \sum_{c \in \Sigma^{N_z(A)}} v_i[\tau_z(A,c)], & \text{if } |N_z(A)| = i\\ 0, & \text{otherwise.} \end{cases}$$

Note in particular that every $F_{z,i}$ can be represented as a 0-1 sparse matrix.

In our experiments we ran Algorithm 1 for different values of l and alphabet sizes σ . As one would expect, the derived lower bounds improve as l grows. However, the memory resources required to perform the computation also increases. Indeed, throughout the second loop of Algorithm 1 we need to store d vectors of dimension σ^{ld} . Also, a simple analysis of the definition of the sparse matrix $F_{z,i}$ shows that it has $\binom{d}{i}\sigma^{(l-1)d}(\sigma-1)^i\sigma^i$ non-zero entries. It follows that a sparse matrix representation of F_z has roughly $\sigma^{ld}(\sigma-1)^d$ non-zero entries. Hence, the necessary computations are feasible only for small values of σ , l and d, unless additional features of the matrices involved are taken advantage of in order to optimize memory usage.

Table 1 summarizes the lower bounds we obtain for $\gamma_{\sigma,2}$ and contrasts them with previously derived ones. To the best of our knowledge, for the d = 2 case and alphabet sizes 3, 4, 5, and 6, this work provides the currently best known lower bounds for $\gamma_{\sigma,2}$. It might be worth mentioning that, as can be seen in that table, the bound of [Dan94, Dek79] is better than the bound of the more recent work of [BYNGS99] for alphabet size 6, and that for bigger alphabet sizes, the bound of [Dan94, Dek79] is still better than ours.

The best known lower bound for $\gamma_{2,2}$ is still the one established by Lueker [Lue03]. Table 2 lists the distinct choices of σ and d for which we could execute Algorithm 1 and indicates the value of the parameter l giving rise to the reported lower bound.

	$\gamma_{\sigma,2}$						
σ	This work	Baeza et. al. lower bound [BYNGS99]	Dančík-Deken's lower bound [Dan94, Dek79]				
3	0.671697	0.63376	0.61538				
4	0.599248	0.55282	0.54545				
5	0.539129	0.50952	0.50615				
6	0.479452	0.46695	0.47169				
7	0.444577	-	0.44502				
8	0.356545	-	0.42237				
9	0.327935	-	0.40321				
10	0.303490	-	0.38656				

Table 1: Best known lower bounds for $\gamma_{\sigma,2}$ (in boldface).

3.6 Disproving Steele's $\gamma_{2,2} = \gamma_{2,3}^2$ speculation

We showed in Section 2 that $\gamma_{2,d} > \gamma_{2,2}^{d-1}$ for all $d \ge 5$. We now establish that this is also the case when d = 3 and d = 4. Recall that Lucker [Lue03] proved that $\gamma_{2,2} \le U$ for U = 0.826280. From Table 2 we see that for d = 3 and d = 4, the indicated lower bound for $\gamma_{2,d}$ is strictly greater than U^{d-1} , and therefore, is also strictly greater than $\gamma_{2,2}^{d-1}$. This implies that $\gamma_{2,d} > \gamma_{2,2}^{d-1}$ for d = 4 and d = 3 as claimed. Together with the results of Section 2 this establishes that $\gamma_{2,d} > \gamma_{2,2}^{d-1}$ for all $d \ge 3$.

4 Final comments

As already mentioned at the start of this paper, Steele [Ste86] pointed out that it would be of interest to find relations between the values of the $\gamma_{\sigma,d}$'s, especially between $\gamma_{2,2}$ and $\gamma_{2,3}$. We think it would be very interesting if such a relation would exist. In fact, it might shed some light upon the longstanding open problem of determining the exact value of the Chvátal-Sankoff constant.

	Alphabet size σ =] [Alphabet size $\sigma = 5$		
d	L such that $\gamma_{2,d} \geq L$	Parameter <i>l</i>		d	L such that $\gamma_{5,d} \geq L$	Parameter <i>l</i>
2	0.781281	10		2	0.539129	4
3	0.704473	7		3	0.356717	2
4	0.661274	5		4	0.289398	1
5	0.636022 4			5	0.273884	1
6	0.617761 3		•			
7	0.602493 2				Alphabet size $\sigma = 6$	
8	0.594016	2		d	L such that $\gamma_{6,d} \geq L$	Parameter l
9	0.587900	2		2	0.479452	3
10	0.570155	1		3	0.309424	2
11	0.570155	1		4	0.245283	1
12	0.563566	1				
13	0.563566	1			Alphabet size σ =	= 7
14	0.558494	1		d	L such that $\gamma_{7,d} \geq L$	Parameter l
-			_	2	0.444577	3
	Alphabet size σ =	= 3		3	0.234567	1
d	L such that $\gamma_{3,d} \geq L$	Parameter <i>l</i>		4	0.212786	1
2	0.671697					
3	0.556649	4		Alphabet size $\sigma = 8$		
4	0.498525	3		d	L such that $\gamma_{8,d} \geq L$	Parameter <i>l</i>
5	0.461402	2		2	0.356545	2
6	0.421436	1		3	0.207547	1
7	0.413611	1				
8	0.405539 1				Alphabet size σ =	= 9
		,		d	L such that $\gamma_{9,d} \ge L$	Parameter <i>l</i>
	Alphabet size σ =	= 4		2	0.327935	2
d	L such that $\gamma_{4,d} \geq L$	Parameter <i>l</i>		3	0.186104	1
2	0.599248	5				10
3	0.457311	3		Alphabet size $\sigma = 10$		
4	0.389008	2		d	L such that $\gamma_{10,d} \ge L$	Parameter <i>l</i>
5	0.335517	1		2	0.303490	2
6	0.324014	1		3	0.168674	1

Table 2: Lower bounds for $\gamma_{\sigma,d}$

Lacking a relation among the $\gamma_{\sigma,d}$'s it would still be interesting to relate these terms to some other constants that arise in connection with other combinatorial problems. A step in this direction was taken by Kiwi, Loebl and Matoušek [KLM05] who showed that $\sqrt{\sigma}\gamma_{\sigma,2} \rightarrow c_2$ when $\sigma \rightarrow \infty$, where c_2 is a constant that turns up in the study of the Longest Increasing Sequence (LIS) problem (also known as Ulam's problem). Specifically, c_2 is the limit to which the expected length of a LIS of a randomly chosen permutation of $\{1, \ldots, n\}$ converges when normalized by \sqrt{n} . Logan and Shepp [LS77] and Vershik and Kerov [VK77] showed that $c_2 = 2$. Consider now the following experiment: Choose *n* points in a unit *d*-dimensional cube $[0, 1]^d$ and let $H_d(n)$ be the random variable corresponding to the length of a longest chain (for the standard partial order in \mathbb{R}^d) of the *n* chosen points. Bollobás and Winkler [BW88] proved that there are constants c'_2, c'_3, \ldots such that $c'_d < e$, $\lim_{d\to\infty} c'_d = e$ and $\lim_{n\to\infty} H_d(n)/n^{1/d} = c'_d$. By labeling a set *S* of points in $[0, 1]^2$ in increasing order of their *x* coordinate and reading the labels in the order of their *y* coordinates one can associate a permutation π to the set *S*. It is easy to see that a chain of points in *S* is in one to one correspondence to an increasing sequence of π . Hence, it follows that $c'_2 = c_2$. Soto [Sot06] extended the results of [KLM05] and showed that $\sigma^{1-1/d}\gamma_{\sigma,d} \to c'_d$ when $\sigma \to \infty$. We think that any similar type of result, or even a reasonable conjecture, that would hold for fixed σ and *d* would also be quite interesting.

References

- [Ale94] K. S. Alexander. The rate of convergence of the mean of the longest common subsequence. *Annals of Applied Probability*, 4:1074–1083, 1994.
- [BW88] B. Bollobás and P. Winkler. The longest chain among random points in euclidean space. Proceedings of the American Mathematical Society, 103(2):347–353, June 1988.
- [BYNGS99] R. Baeza-Yates, G. Navarro, R. Gavaldá, and R. Schehing. Bounding the expected length of the longest common subsequences and forests. *Theory of Computing Systems*, 32(4):435–452, 1999.
- [CS75] V. Chvátal and D. Sankoff. Longest common subsequences of two random sequences. J. of Applied Probability, 12:306–315, 1975.
- [Dan94] V. Dančík. Expected Length of Longest Common Subsequences. PhD thesis, Department of Computer Science, University of Warwick, September 1994.
- [Dan98] V. Dančík. Common subsequences and supersequences and their expected length. Combinatorics, Probability and Computing, 7(4):365–373, 1998.
- [Dek79] J. P. Deken. Some limit results for longest common subsequences. *Discrete Mathematics*, 26:17–31, 1979.
- [DP95] V. Dančík and M. Paterson. Upper bounds for the expected length of a longest common subsequence of two binary sequences. *Random Structures & Algorithms*, 6(4):449–458, 1995.
- [JŁR00] S. Janson, T. Łuczak, and A. Rucinski. *Random Graphs*. Wiley, 2000.
- [KLM05] M. Kiwi, M. Loebl, and J. Matoušek. Expected length of the longest common subsequence for large alphabets. Advances in Mathematics, 197(2):480–498, 2005.
- [LS77] B. Logan and L. Shepp. A variational problem of random Young tableaux. *Advances in Mathematics*, 26:206–222, 1977.
- [Lue03] G. Lueker. Improved bounds on the average length of longest common subsequences. In *Proceedings* of the 14th Annual ACM/SIAM Symposium on Discrete Algorithms, pages 130–131, 2003.
- [Pev00] P. Pevzner. Computational Molecular Biology: An Algorithmic Approach. MIT Press, 2000.
- [Sot06] J. Soto. Variantes aleatorias de la subsecuencia común más grande. Departamento de Ingeniería Matemática, U. Chile, Aug 2006. (In Spanish).
- [Ste86] J. M. Steele. An Efron–Stein inequality for nonsymmetric statistics. *The Annals of Statistics*, 14(2):753–758, 1986.
- [Ste96] J. M. Steele. *Probability Theory and Combinatorial Optimization*. CBMS-NSF Regional Conference Series in Applied Mathematics. SIAM, 1996.
- [Szp00] W. Szpankowski. *Average Case Analysis of Algorithms and Sequences*. Series in Discrete Mathematics and Optimization. Wiley Inter-Science, 2000.
- [VK77] A. Vershik and S. Kerov. Asymptotics of the Plancherel measure of the symmetric group and the limiting form of Young tableaux. *Doklady Akademii Nauk SSSR*, 233:1024–1028, 1977.
- [Wat95] M. Waterman. Introduction to Computational Biology: Average Case Analysis of Algorithms and Sequences. Series in Discrete Mathematics and Optimization. Chapman & Hall/CRC, first CRC press reprint edition, 1995.