The Copernican Multiverse of Sets

Paul K. Gorbow and Graham E. Leigh

University of Gothenburg

Department of Philosophy, Linguistics, and Theory of Science
Box 200, 405 30 GÖTEBORG, Sweden

Abstract

We develop an untyped framework for the multiverse of set theory. ZF is extended with semantically motivated axioms utilizing the new symbols $\mathsf{Uni}(\mathcal{U})$ and $\mathsf{Mod}(\mathcal{U},\sigma)$, expressing that \mathcal{U} is a universe and that σ is true in the universe \mathcal{U} , respectively. Here σ ranges over the augmented language, leading to liar-style phenomena that are analysed. The framework is both compatible with a broad range of multiverse conceptions and suggests its own philosophically and semantically motivated multiverse principles. In particular, the framework is closely linked with a deductive rule of Necessitation expressing that the multiverse theory can only prove statements that it also proves to hold in all universes. We argue that this may be philosophically thought of as a Copernican principle that the background theory does not hold a privileged position over the theories of its internal universes.

Our main mathematical result is a lemma encapsulating a technique for locally interpreting a wide variety of extensions of our basic framework in more familiar theories. We apply this to show, for a range of such semantically motivated extensions, that their consistency strength is at most slightly above that of the base theory ZF, and thus not seriously limiting to the diversity of the set-theoretic multiverse. We end with case studies applying the framework to two multiverse conceptions of set theory: arithmetic absoluteness and Joel D. Hamkins' multiverse theory.

1 Introduction

ZF set theory serves as a foundation for mathematics, but has also turned out to be interesting in itself as a field of mathematical study. Much of the interest lies in that it raises questions that are not only undecidable, but also lacking clear-cut intuitive answers and demanding deep mathematical developments. The continuum hypothesis is a primary historical example: It seems implausible to reach a consensus on affirming or denying it, and it motivated two techniques central to set theory: the inner model and forcing constructions. It is natural to view these techniques as enabling constructions of set-theoretic universes from other set-theoretic universes, thus taking a multiverse view of the subject matter of set theory, rather than adopting the universe view that there is a single absolute universe of sets. In the words of Hamkins, an advocate of the multiverse view (2012, p. 418):

A large part of set theory over the past half-century has been about constructing as many different models of set theory as possible /.../ Would you like to live in a universe where CH holds, but \diamondsuit fails? Or where $2^{\aleph_n} = \aleph_{n+2}$ for every natural number n? Would you like to have rigid Suslin trees? Would you like every Aronszajn tree

to be special? Do you want a weakly compact cardinal κ for which $\diamond_{\kappa}(REG)$ fails? Set theorists build models to order.

Hamkins follows this perspective on set-theoretic practice with his argument for adopting the multiverse view:

This abundance of set-theoretic possibilities poses a serious difficulty for the universe view, for if one holds that there is a single absolute background concept of set, then one must explain or explain away as imaginary all of the alternative universes that set theorists seem to have constructed. This seems a difficult task, for we have a robust experience in those worlds, and they appear fully set theoretic to us. The multiverse view, in contrast, explains this experience by embracing them as real, filling out the vision hinted at in our mathematical experience, that there is an abundance of set-theoretic worlds into which our mathematical tools have allowed us to glimpse.

Methodologically, it makes sense to represent the universes of sets as models of a theory of sets, as that makes them accessible to the well-developed techniques of model theory. This raises two questions:

- 1. Which set theory?
- 2. Which models of that set theory?

In answer to the first question the authors have decided on limiting scope to ZF. This is the most utilized set theory, established in the mathematical community as a robust foundation of mathematics. All of the results of this paper go through for extensions of ZF (axiom of choice, large cardinals, ...). Although one development of this paper uses full ZF,¹ the authors conjecture that for most of the results much weaker fragments suffice.

On the second question this paper takes a liberal approach. In particular, scope is not limited to well-founded models. The framework is intended to be applicable to a wide range of multiverse conceptions, for example the conceptions that every universe is of the form V_{α} , that every universe is well-founded, that every universe is countable and recursively saturated (and therefore ill-founded), or that every model is a universe.

Why consider ill-founded universes? There is a sense in which a model \mathcal{M} of set theory can be situated as an element in two different models \mathcal{N}_0 and \mathcal{N}_1 of set theory, such that \mathcal{N}_0 satisfies that \mathcal{M} is well-founded while \mathcal{N}_1 satisfies that \mathcal{M} is ill-founded.² Thus, we may think of the property of well-foundedness as depending on the set-theoretic background. In Hamkins's multiverse conception this is a key feature (Hamkins, 2012, p. 438-9):

The concept of well-foundedness [...] depends on the set-theoretic background, for different models of set theory can disagree on whether a structure is well-founded. [...] Indeed, every set-theoretic argument can take place in a model, which from the inside appears to be totally fine, but actually, the model is seen to be ill-founded from the external perspective of another, better model. Under the universe view, this problem terminates in the absolute set-theoretic background universe, which provides an accompanying absolute standard of well-foundedness. But the multiverse view allows for many different set-theoretic backgrounds, with varying concepts of the well-founded, and there seems to be no reason to support an absolute notion of well-foundedness.

¹The proof of Theorem 2.4 requires full ZF. This is used in the proof of Theorem 5.7, a consistency result. ²Using Definition 2.1, the precise sense is that $\mathcal{M} = \mathcal{A}_{\mathcal{N}_0} = \mathcal{B}_{\mathcal{N}_1}$, for some $\mathcal{A} \in \mathcal{N}_0$ and $\mathcal{B} \in \mathcal{N}_1$, such that \mathcal{N}_0 satisfies that \mathcal{A} is well-founded while \mathcal{N}_1 satisfies that \mathcal{B} is ill-founded.

To approach the set-theoretic multiverse mathematically, we need a foundational theory to situate the universes in. Just as the foundational background theory of ZF is useful for studying groups and topological spaces, it is useful for studying set-theoretic universes. So we find ourselves in a situation of studying models of ZF from the background theory ZF. The multiverse theorist may extend the background theory of ZF to a multiverse theory (in an expanded language), with axioms specifying properties of the multiverse. In such a background multiverse theory, it is natural to consider the universes as themselves being models of multiverse theories, having their own internal universes, and so on. This raises an important question:

Main Question. What is the relationship between the external universe of the background multiverse theory, and the universes internal to the background theory? Similarly, what is the relationship between each universe and the universes within that universe?

We shall investigate several responses to the Main Question. Most fundamentally, the authors propose that the background multiverse theory obeys the following principle:

Copernican Principle. The background theory of the multiverse should not have a privileged position compared to the multiverse theories of the internal universes; specifically, if the background multiverse theory proves a statement, it should also prove that holds in all universes.

We have an analogy with the heliocentric model of the solar system: Earth corresponds to the universe of the background multiverse theory, as the basis for our point of view. The geocentric model gives earth a privileged central position as an absolute reference point, while the heliocentric model puts earth on a par with all of the planets. So the heliocentric model differs from the geocentric model in that it obeys the principle that for any appropriately fundamental assumption we make about our point of view, we are committed to making the same assumption about every other plausible point of view. Similarly, in the context of set theory, the authors propose the Copernican Principle as the constraint that for every assumption introduced by a multiverse theorist, s/he is committed to that it holds from the vantage point of an arbitrary universe of sets. The name is borrowed from a modern principle in physics, which Peacock states as "that humans are not privileged observers". Peacock applies the principle arguing "if the universe appears isotropic about our position, it would also appear isotropic to observers in other galaxies" (1998, p. 66). So for the physicist, the principle is a conceivably falsifiable statement about the uniformity of the physical universe; while for the theorist of the multiverse of sets, it is an a priori postulate. Below we explicate a formal deductive rule, NEC, expressing this principle.

To approach the Main Question, we require a framework that makes sense of the notion of truth-in-a-universe. If the universes are mere models of ZF, then the usual satisfaction-relation expressed in the language of set theory suffices. But as soon as we consider each universe to contain a multiverse in its own right, it is more natural to consider the universes as structures in the language of the multiverse theory.

The main contribution of this paper is an untyped framework for handling the notion of truth-in-a-ZF-universe. It is intended to be applicable to just about any theory of the multiverse of sets. A primitive predicate $Uni(\mathcal{U})$ is introduced to express that \mathcal{U} is a universe, and the primitive relation $Mod(\mathcal{U}, \sigma)$ is introduced to express that the $\mathcal{L}_{Uni,Mod}$ -statement σ is true in the universe \mathcal{U} , where $\mathcal{L}_{Uni,Mod}$ is the language of set theory, \mathcal{L} , augmented with the symbols Uni and Mod. The multiverse theories considered in this paper are expressed in $\mathcal{L}_{Uni,Mod}$.

1.1 Multiverse principles

Now that the language $\mathcal{L}_{Uni,Mod}$ and the intuitive intended meaning of its symbols has been briefly explained, the next task is to give natural and useful axioms for Uni and Mod. Axioms for

Uni specify what universes comprise the multiverse, what closure properties it satisfies, etc.; for example the multiverse axioms of Hamkins (2012). In this paper we are focused on semantically motivated axioms, meant to be applicable to a wide range of multiverse conceptions. The application of this framework to Hamkins's multiverse is discussed in §6.2. Since Mod is an untyped semantic relation, it is not surprising that it is exposed to liar-style phenomena. Our Theorem 4.6 shows, e.g., that the schema $(\forall \mathcal{U} (\mathsf{Uni}(\mathcal{U}) \to \mathsf{Mod}(\mathcal{U}, \ulcorner \sigma \urcorner)) \to \sigma$, over $\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}$ -statements σ , expressing that whatever holds in every universe also holds in the background universe, is inconsistent with a natural and mild theory in $\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}$. However, this contradiction is not derivable when this schema is restricted to \mathcal{L} .

The basic theory introduced is called CM^- (Compositional satisfaction for the Multiverse).³ CM^- is formed by adding compositional semantically motivated $\mathcal{L}_{Uni,Mod}$ -axioms to the background theory ZF, for each logical connective and quantifier, and by extending the Separation and Replacement schemas of ZF to $\mathcal{L}_{Uni,Mod}$. For example, the compositional axiom for \wedge is

```
if \theta \in \mathcal{L}_{\mathsf{Uni},\mathsf{Mod}} is the conjunction of \phi and \psi, then \mathsf{Mod}(\mathcal{U},\theta) \iff \mathsf{Mod}(\mathcal{U},\phi) \land \mathsf{Mod}(\mathcal{U},\psi).
```

These are also called the Tarskian laws of satisfaction. The analogue in the present framework of the well-known Tarskian schema $\operatorname{Tr}(\lceil \sigma \rceil) \leftrightarrow \sigma$ (for $\sigma \in \mathcal{L}$) would say roughly that σ is true in every universe if, and only if, it holds in the background universe. So this would say that the multiverse is not very diverse, and certainly not closed under forcing, for example. But analogues of $\operatorname{Tr}(\lceil \sigma \rceil) \to \sigma$ (for $\sigma \in \mathcal{L}$) and the rule of Necessitation $\vdash \sigma \Rightarrow \vdash \operatorname{Tr}(\lceil \sigma \rceil)$ (for $\sigma \in \mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}$) are highly relevant.

In CM $^-$ we can prove the soundness principle that the set of statements true in any particular universe is deductively closed. CM is CM $^-$ plus an axiom called Multiverse_{ZF} saying that every universe satisfies ZF, which is just intended to set the scope of the present treatment. (For most of the results, the authors believe that natural generalizations to weak fragments of ZF are possible.) Theorem 5.6 shows that an extension of CM interpreting the Gödel-Löb modal logic is conservative over ZF. So by the soundness principle, Multiverse_{ZF}, and Gödel's second incompleteness theorem, CM does not prove the statement $\exists \mathcal{U} \, \mathsf{Uni}(\mathcal{U})$, saying that there exists a universe.

A flexible revision-semantic technique for expanding models of the background theory ZF to models of extensions of CM⁻ is developed. This technique builds on ideas from Gupta (1982), Herzberger (1982a) and Herzberger (1982b), for circumventing truth-theoretic paradoxes. In short, one starts by setting parameters specifying the particular multiverse conception desired. Among other things, this pins down the interpretation of Uni. Then the interpretation of Mod is determined by a revision-semantic process. Intuitively, a basic definition of truth-in-a-universe is supplied among the parameters, and this definition is revised step-by-step to more adequate definitions. Theorems 5.4 and 5.5 show that some natural settings of the parameters lead to that further semantically motivated axioms and deductive rules are validated in the constructed model, more on this further below.

We introduce several axioms and deductive rules in response to the Main Question. The most fundamental such principle for this framework is the deductive rule of Necessitation, NEC, which is a formal expression of the Copernican Principle:

If
$$\sigma$$
 is provable, then $\forall \mathcal{U} (\mathsf{Uni}(\mathcal{U}) \to \mathsf{Mod}(\mathcal{U}, \lceil \sigma \rceil))$ is provable,

where $\lceil \sigma \rceil$ is the Gödel code of σ . Under mild assumptions on the parameters, NEC is validated in the revision-semantic model construction. Theorem 4.8 shows that CM + NEC is conservative over ZF.

³The precise specification of CM⁻ is given in System 4.1.

Dually, the deductive rule of Co-Necessitation, CONEC, states:

If $\forall \mathcal{U} (\mathsf{Uni}(\mathcal{U}) \to \mathsf{Mod}(\mathcal{U}, \lceil \sigma \rceil))$ is provable, then σ is provable.

In the context of NEC as formalizing the Copernican Principle, CONEC may be thought of as expressing that the theory is maximal within the bounds of the Copernican Principle. On the other hand, as a stand-alone principle, CONEC can be used to boost the expressive power: For example, we will consider CM extended by CONEC and the statement that no universe satisfies a Σ_1^0 -statement that does not already hold in the standard model of arithmetic in the background theory; in other words, a Turing machine that does not halt in the background theory, halts in no universe. In CM extended with this axiom we can use basic model-theoretic considerations to prove that every universe satisfies the Reflection schema iterated ω_1^{CK} times over ZF.⁴ Now, by adding CONEC we can prove ω_1^{CK} -iterated Reflection schema over ZF outright in the background theory. So in general, CONEC enables outright proofs of statements that are provably satisfied across a model-theoretically delimited multiverse, thus in some sense "extracting the deductively accessible content" of higher-order non-recursive properties.

We write MS (Multiverse theory of Satisfaction) for the theory CM + NEC + CONEC. This theory is analogous to the Firedman—Sheard theory of truth (FS) from Friedman & Sheard (1987). A revision-semantic technique for constructing models of CM + NEC and/or CONEC (building on a technique from the aforementioned two papers) is embodied in the Main Lemma (in §5) and its Corollary 5.2.

We now proceed to discuss three axioms motivated by the Main Question that have a reflective character in that they assert that the background universe is in some sense reflected in the multiverse. We will establish bounds on the consistency strength of these in terms of iterated reflection principles. The reader is referred to Systems 2.8 and 5.3 for the definition of these principles.

A very basic multiverse axiom is Non-Triviality, $\exists \mathcal{U} \, \mathsf{Uni}(\mathcal{U})$, saying that there is a universe. In the presence of NEC, this also yields that every universe contains a universe, and so on. We show in Theorem 5.4 (and in Theorem 5.6) that $\mathsf{CM} + \mathsf{Non-Triviality} + \mathsf{NEC}$ is locally interpreted in (and conservative over) the theory of iterated consistency over ZF .

A stronger axiom motivated by the Main Question, called Self-Perception, expresses that the background universe is isomorphic (over the set theoretic language) to one of the internal universes. This embodies the idea that the universe of the background theory should also be available in its internal multiverse, and has a distinct reflective character. It turns out to be convenient to take the universes to be countable recursively saturated models when modelling CM + Self-Perception + NEC, a phenomenon that corresponds to the multiverse model of Gitman and Hamkins in (2010). This suggests that their multiverse theory would harmonize well with CM + Self-Perception + NEC, a hypothesis we explore briefly in subsection 6.2. In Theorem 5.7, we use the revision-semantic technique to interpret CM + Self-Perception + NEC in the theory of ω -iterated Global Reflection over ZF. The latter is a natural untyped theory of truth, that mildly strengthens ZF. All universes are countable recursively saturated in this interpretation.

Self-Perception is closely related to the notion of condensible models studied by Enayat in (2020). The definition of condensability is somewhat technical, involving an infinitary language: A model \mathcal{M} of ZF is condensible, if there is some ordinal $\alpha \in \mathcal{M}$ such that $\mathcal{M} \cong \mathcal{M}(\alpha) \prec_{\mathbb{L}_{\mathcal{M}}} \mathcal{M}$, where $\mathcal{M}(\alpha)$ denotes the substructure of \mathcal{M} of ranks below α and $\mathbb{L}_{\mathcal{M}}$ denotes the intersection of $\mathcal{L}_{\omega_1,\omega}$ with the well-founded part of \mathcal{M} . In particular, Enayat positively answers a question that sprung from the present paper: Is there an ω -standard model of ZF with unboundedly many ordinals α such that $\mathcal{M} \cong \mathcal{M}(\alpha) \prec \mathcal{M}$? Note that recursively saturated models are

⁴See System 5.3 for the definition of this theory.

 ω -non-standard. Enayat's result means that the door also appears to be open for models of CM + Self-Perception + NEC with ω -standard universes.

We also introduce the axiom schema of Multiverse Reflection, stating for each sentence σ in the language \mathcal{L} of set theory: $(\forall \mathcal{U} (\mathsf{Uni}(\mathcal{U}) \to \mathsf{Mod}(\mathcal{U}, \lceil \sigma \rceil))) \to \sigma$. Over CM, this principle is implied by Self-Perception and implies Non-Triviality. Using the revision-semantic technique, we show in Theorem 5.5 (and Theorem 5.6) that MS + Multiverse Reflection is locally interpreted in (and conservative over) the theory of ω -iterated proof-theoretic reflection schema over ZF.

The body of the paper ends with case studies, where we look at two independent multiverse conceptions through the lens of the framework we have developed. The first of these is a conception of the multiverse as being arithmetically absolute, in the sense that arithmetic truth does not vary across the multiverse. The second is a conception due to Hamkins which is fundamentally based on the principles that the multiverse is closed under the forcing and inner model techniques, and that every universe is countable and ω -non-standard from the perspective of some other universe. We close with a concluding section reflecting on the contributions of the paper and the relevance of this framework for future research on the set-theoretic multiverse.

2 Preliminaries

2.1 A term-calculus for representation of syntax

We shall work with various recursively enumerable set theories, in languages obtained by adding finitely many new non-logical symbols to the usual language of set theory on the signature $\{\in\}$. We define a *set theory* to be any recursively enumerable system proving Mac Lane set theory (excluding Choice)⁵ or proving Kripke–Platek set theory with Infinity⁶, in a language with finitely many non-logical symbols including a term-calculus for arithmetic and Gödel coding explained below. Our specific choice of set theories underlying this definition is somewhat arbitrary (even weaker theories may well suffice). Both of these theories are sufficient for constructing the structure of the natural numbers and implementing basic model theory; these are the important features for this paper.

Since we will be reasoning about syntactic objects, it is convenient to employ a Gödel coding of syntax. Let K be a language with finitely many non-logical symbols. In any set theory T (in language L) under consideration, we can define the arithmetic functions needed to formulate a natural $G\"{o}del$ coding in T of terms and formulas of K. Through the $G\"{o}del$ coding, the "grammatical structure" of K is coherently represented in T. The complicated details of this procedure are described in any rigorous account of $G\"{o}del$'s incompleteness theorems. The gist is that for each syntactic object (symbol, term or formula) s of K, there is a definable number $\lceil s \rceil$ in L (the $G\"{o}del$ code of s), which represents s in T, and there are operations definable in T corresponding to syntactic operations on such objects. The authors trust that the reader is familiar with this.

It is customary in set theory to informally introduce defined constant, relation and function symbols to the language, in order to make the presentation more readable. For example, one may use a function symbol +, as if it belonged to the language and there was an axiom expressing that + is addition on the finite von Neumann ordinals. In this paper we assume that a finite number of such symbols needed for arithmetic and Gödel coding are already present in the language of every set theory, and that the appropriate axioms regulating them are available in every set

⁵Its axioms are Extensionality, Null set, Pair, Union, Power set, Separation for Δ_0 -formulas, and Infinity.

⁶Its axioms are Extensionality, Null set, Pair, Union, Separation for Δ_0 -formulas, Collection for Δ_0 -formulas, Foundation for Π_1 -formulas, and Infinity.

theory. Here follows a semi-formal account of some of the main principles of this expanded language L for a set theory T, also serving to specify the notation:

- 1. We have a constant $\underline{0}$ and function symbols $S, +, \times$ for the successor, addition and multiplication operations in arithmetic. For each $n \in \mathbb{N}$, n is shorthand for $S^n(0)$.
- 2. Each variable, constant, relation or function symbol s of K is represented in T by a numeral $\lceil s \rceil$ in L.
- 3. Recursively, each term $f(t_1, \ldots, t_n)$ of K is represented by the term $\lceil f \rceil (\lceil t_1 \rceil, \ldots, \lceil t_n \rceil)$ of L. Formally, the term $\lceil f \rceil (\lceil t_1 \rceil, \ldots, \lceil t_n \rceil)$ is the result of applying a function symbol of L to the numerals $\lceil f \rceil, \lceil t_1 \rceil, \ldots, \lceil t_n \rceil$. Moreover, $\lceil f(t_1, \ldots, t_n) \rceil$ denotes a numeral that T proves to equal $\lceil f \rceil (\lceil t_1 \rceil, \ldots, \lceil t_n \rceil)$.
- 4. Each atomic formula $R(t_1, \ldots, t_n)$ of K is represented by the numeral $\lceil R(t_0, \ldots, t_n) \rceil$ of L. Analogous remarks apply as in the case of terms described above.
- 5. The syntactic operations, standardly used to build up complex formulas from atomic formulas, are all available. For example:
 - (a) L has a function symbol \neg , such that for each ϕ in K, $\neg \ulcorner \phi \urcorner$ represents $\neg \phi$. Moreover, T proves that $\ulcorner \neg \phi \urcorner = \neg \ulcorner \phi \urcorner$.
 - (b) L has a function symbol $\dot{\wedge}$, such that for each ϕ and each ψ in K, $\lceil \phi \rceil \dot{\wedge} \lceil \psi \rceil$ represents $\phi \wedge \psi$. Moreover, T proves that $\lceil \phi \wedge \psi \rceil = \lceil \phi \rceil \dot{\wedge} \lceil \psi \rceil$.
 - (c) L has a function symbol \forall , such that for each variable v and each formula ϕ in K, $\forall \ulcorner v \urcorner \ulcorner \phi \urcorner$ represents $\forall v \phi$. Moreover, T proves that $\ulcorner \forall v \phi \urcorner = \forall \ulcorner v \urcorner \ulcorner \phi \urcorner$.
 - (d) For any ϕ in K, $\phi[t/x]$ denotes the formula obtained from ϕ by replacing each free occurrence of the variable x by the term t (if t has variables, then their bound occurrences in ϕ are renamed as necessary). L has a function symbol (written -[-/-]) which represents this primitive recursive substitution operation. Moreover, T proves that $\lceil \phi(x) \lceil y/x \rceil \rceil = \lceil \phi(x) \rceil \lceil y \rceil / \lceil x \rceil$. Somewhat less formally, if ϕ has been introduced as $\phi(x)$, we may write $\phi(t)$ for the formula $\phi[t/x]$.

In the context of a set theory T in a set-theoretic language L, Σ_n^0 , Π_n^0 and Δ_n^0 denote the usual arithmetic hierarchy as defined for L-formulas (all quantifiers are bounded to \mathbb{N}), up to equivalence in T. It is well-known that for any recursive system S, there is a Σ_1^0 -formula \Pr_S , representing S-provability in T. We write Con_S for the sentence $\neg \Pr_S(\ulcorner \bot \urcorner)$, expressing that S is consistent. In both cases, the dot under S is sometimes omitted, when it can be inferred from the context.

As an example, consider this consequence of Gödel's second incompleteness theorem:

$$\mathsf{ZF} \nvdash \mathsf{Con}_{\mathsf{ZF}}$$

From the perspective of the meta-theory, "ZF" refers to a set of sentences (the object theory of ZF) whereas "ZF" refers to a formula representing the recursive set of Gödel codes of that set in the object theory ZF.

Suppose now that a set theory T' in language L' is represented in a set theory T in language L. Then L' is Gödel coded in T, as explained above. But T', in turn, also Gödel codes languages; say that T' Gödel codes the language L''. Note that the whole Gödel coding of L'' in T' is then carried along by the representation of T' in T. For example, if ϕ is a formula in L'', then there is a term $\lceil \phi \rceil$ in L' which represents ϕ in T'. If $\psi(x)$ is a formula of L', we can then form the

formula $\psi[\lceil \phi \rceil/x]$ of L'. This formula, in turn, is represented in T by an L-term $\lceil \psi[\lceil \phi \rceil/x] \rceil$. So if $\theta(y)$ is an L-formula, we can form the L-formula $\theta[\lceil \psi[\lceil \phi \rceil/x] \rceil/y]$. Thus, Gödel codes may be nested, as a set theory represents a set theory, which in turn represents a language.

2.2 Miscellaneous logical preliminaries

 \mathcal{L} is the language with the symbol " \in " along with a finite number of arithmetic and syntactic symbols as explained in Subsection 2.1. We assume that ZF is formulated as an \mathcal{L} -theory, with the natural axioms for defining the arithmetic and syntactic symbols of \mathcal{L} . \mathcal{L}^+ denotes any extension of \mathcal{L} with a finite number of new symbols.

If L is a language and S_1, \dots, S_n are symbols, then L_{S_1,\dots,S_n} denotes the language obtained by augmenting L with S_1,\dots,S_n . The Separation schema applying to all formulas of a language L is denoted Sep(L), and the Replacement schema applying to all formulas of a language L is denoted Rep(L).

Any set theory suffices as meta-theory. Suppose that in the meta-theory we consider a definable set $A = \{x \mid \phi(x)\}$, such as a theory. We may then refer to the corresponding set within an object-theory, for example as follows: Using the symbol A somewhat ambiguously, we write a statement of the form $\mathsf{ZF} \vdash \cdots \supset \mathsf{M} \models A \cdots$ for the more formally precise statement of the form $\mathsf{ZF} \vdash \cdots \supset \mathsf{X} \ (\forall x (x \in X \leftrightarrow \phi(x)) \land \forall x \in \mathsf{X} \ (\mathcal{M} \models x)) \cdots$. The dot under A is occasionally omitted, when clear from the context. To illustrate, we might express a special case of Gödel's completeness theorem within $A \supset \mathsf{TF} \ (\mathsf{Con}_{\mathsf{ZF}} \leftrightarrow \exists \mathcal{M} \ (\mathcal{M} \models \mathsf{ZF}))$. We say that a theory $A \supset \mathsf{TF} \ (\mathsf{Con}_{\mathsf{ZF}} \to \mathsf{TF})$ bounds the consistency strength of (or has at least as high consistency strength as) a theory $A \supset \mathsf{TF} \ (\mathsf{Con}_{\mathsf{ZF}} \to \mathsf{TF})$ if the consistency of $A \supset \mathsf{TF}$ implies the consistency of $A \supset \mathsf{TF} \ (\mathsf{Con}_{\mathsf{ZF}} \to \mathsf{TF})$.

An interpretation \mathcal{I} from a language L_0 to a language L_1 is a function $\mathcal{I}: L_0 \to L_1$ which is generated (by structural recursion) from the non-logical symbols of L_0 . Moreover, we say that \mathcal{I} interprets or validates the L_0 -system T_0 in the L_1 -system T_1 if for any L_0 -formula ϕ , $T_1 \vdash \mathcal{I}(\phi)$ whenever $T_0 \vdash \phi$.

As default, we work with first-order languages and classical logic, but we will consider additional deductive rules (NEC and CONEC). $\phi \equiv \psi$ is the statement that ϕ and ψ are identical formulas. If S and T are systems in languages both including L, then $S \equiv_L T$ is the statement that S and T have the same L-theorems. If S is a system involving deductive rules, and A is an axiom, then S + A denotes the natural extension of S in which these deductive rules may be applied to proofs also involving A. For example, in $\mathsf{MS} + \exists x \, \mathsf{Uni}(x)$, we may use NEC to derive $\forall \mathcal{U} \in \mathsf{Uni} \, \mathsf{Mod}(\mathcal{U}, \ulcorner \exists x \, \mathsf{Uni}(x) \urcorner)$.

It is sometimes notationally convenient to introduce classes of the form $A = \{x \mid \phi(x)\}$ (the class of all sets x such that $\phi(x)$), where ϕ is an \mathcal{L}^+ -formula. Then $x \in A$ may be regarded as an alternative notation for $\phi(x)$. Thus, we have no need to specify a formal class theory. For example $V = \{x \mid \top\}$ is the class of all sets.

Formally, Var is the set of variables $\{x_1, x_2, \dots\}$, indexed by the positive natural numbers. But we use other symbols informally (such as $x, y, p, f, \mathcal{U}, \dots$) for variables as well. VA is the class of variable-assignments, $\{f \mid f : \text{Var} \to V\}$. If a is a set (or a structure), then VA^a is the set $\{f \mid f : \text{Var} \to a\}$ of all variable-assignments to elements of (the underlying set of) a. If f is a variable-assignment and v is a variable, then $\text{VA}_{f,v}$ is the set of all variable-assignments g, such that for all $u \in \text{Var}, u \neq v \to g(u) = f(u)$. Suppose that we are working in a set theory T in a language L containing terms $t_1, \dots t_n$. Note that for $n < \omega$, T proves from $v_1 \in \text{Var}, \dots, v_n \in \text{Var}$ that there is a primitive recursive variable-assignment f satisfying $f(v_1) = t_1, \dots, f(v_n) = t_n$, and $\forall m \in \mathbb{N} \ (m > n \to f(v_m) = \underline{0})$. Such a variable assignment f is denoted $\langle v_1, \dots v_n \rangle \mapsto \langle t_1, \dots t_n \rangle$ (or just $v_1 \mapsto t_1$ in the case n = 1).

We assume that model theory is set up so that any structure \mathcal{M} uniquely determines its

language, which we denote by $L(\mathcal{M})$, and we take the symbol " \mathcal{M} " to refer ambiguously to both the structure and its domain. Let \mathcal{M} be an L-structure, ϕ a formula in L and $f \in \mathrm{VA}^{\mathcal{M}}$. We use the arrow-notation \vec{a} for finite tuples $\langle a_1, \cdots, a_n \rangle$, and the shorthand $\vec{a} \in \mathcal{M}$ for that each component a_i of \vec{a} is an element of \mathcal{M} . We write $\mathcal{M} \models (\phi, f)$ for the statement " ϕ is true in \mathcal{M} under the variable-assignment f", as defined in the usual Tarskian semantics of first-order logic. If $\vec{a} \in \mathcal{M}$ and $\psi(\vec{x}) \in L$, then we write $\mathcal{M} \models \psi(\vec{a})$ for $\mathcal{M} \models (\psi, \vec{x} \mapsto \vec{a})$. We write $\mathcal{M} \models \phi$ for $\forall f \in \mathrm{VA}^{\mathcal{M}} \mathcal{M} \models (\phi, f)$. If K is a sublanguage of L, then $\mathcal{M} \upharpoonright_K$ denotes the reduct of \mathcal{M} to K. We write $\mathcal{M} \equiv_K \mathcal{N}$ for the statement that \mathcal{M} satisfy the same K-sentences as \mathcal{N} . We write $\mathcal{M} \cong_K \mathcal{N}$ for the statement that $\mathcal{M} \upharpoonright_K$ is isomorphic to $\mathcal{N} \upharpoonright_K$. When the subscripts are dropped, they are assumed to be $L(\mathcal{M})$.

We use abbreviations for certain variations of the quantifiers:

- 1. $\exists x \in y \ \phi \text{ stands for } \exists x \ (x \in y \land \phi)$.
- 2. $\forall x \in y \ \phi \text{ stands for } \forall x \ (x \in y \to \phi).$
- 3. $\exists x \phi$ stands for $\exists x (\phi(x) \land \forall y (\phi(y) \to x = y))$.

If P is a predicate symbol, we may write $x \in P$ for P(x). Similarly, we write $\exists x \in P \phi$ for $\exists x (P(x) \land \phi)$, and so on.

We will introduce primitive relation symbols "Sat", "Uni" and "Mod". Informally, $\mathsf{Sat}(\phi, f)$ expresses that ϕ is satisfied under the variable assignment f; $\mathsf{Uni}(\mathcal{U})$ expresses that \mathcal{U} is a universe; and $\mathsf{Mod}(\mathcal{U}, \phi, f)$ expresses that ϕ is satisfied in the universe \mathcal{U} under the variable assignment f. Recall that $\mathcal{L}_{\mathsf{Sat}}$ denotes the language \mathcal{L} augmented with the symbol Sat , while $\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}$ denotes \mathcal{L} augmented with Mod and Uni .

Again, we introduce some abbreviations:

- 1. $\mathsf{Sat}(\phi)$ and $\mathsf{Tr}(\phi)$ stand for $\forall f \in \mathsf{VA} \, \mathsf{Sat}(\phi, f)$.
- 2. $\mathsf{Mod}(\mathcal{U}, \phi)$ stands for $\forall f \in \mathsf{VA}^{\mathcal{U}} \, \mathsf{Mod}(\mathcal{U}, \phi, f)$.
- 3. $\mathsf{Tr}^{\square}(\phi)$ stands for $\forall \mathcal{U} \in \mathsf{Uni} \, \forall f \in \mathsf{VA}^{\mathcal{U}} \, \mathsf{Mod}(\mathcal{U}, \phi, f)$.
- 4. $\operatorname{Tr}^{\diamondsuit}(\phi)$ stands for $\neg \operatorname{Tr}^{\square}(\dot{\neg}\phi)$.

If X is a formula, term or definable object in the language L of the structure \mathcal{M} , then $X^{\mathcal{M}}$ denotes its interpretation in \mathcal{M} ; e.g. $\phi^{\mathcal{M}} =_{\mathrm{df}} \{ \vec{a} \in \mathcal{M} \mid \mathcal{M} \models \phi(\vec{a}) \}.$

Informally speaking, if \mathcal{M} is a model of set theory, $\mathcal{N} \in \mathcal{M}$, and $\mathcal{M} \models$ " \mathcal{N} is a model of set theory", so that \mathcal{N} is an internal model of \mathcal{M} , then we may need to extract \mathcal{N} into an external model that can be studied on a par with \mathcal{M} . This is achieved by the following formal definition.

Definition 2.1. If \mathcal{M} is a model of set theory and $a \in \mathcal{M}$, then

$$a_{\mathcal{M}} =_{\mathrm{df}} \{ x \in \mathcal{M} \mid \mathcal{M} \models x \in a \}.$$

This notation is generalized in the cases that a is considered as a relation or as a structure: If \mathcal{M} is a model of set theory, $R \in \mathcal{M}$ and \mathcal{M} satisfies that R is an \underline{n} -ary relation (for a natural number n under consideration), then

$$R_{\mathcal{M}} =_{\mathrm{df}} \big\{ \langle x_1, \cdots, x_n \rangle \mid \exists p \in \mathcal{M} \left[\mathcal{M} \models p \in R \land \right] \\ \bigwedge_{1 \le i \le n} \text{``} x_i \text{ is the } \underline{i} \text{:th component of } p'' \big] \big\}.$$

If \mathcal{M} is a model of set theory, $\mathcal{N}, N, R_1, \dots, R_n$ are elements of \mathcal{M} , and \mathcal{M} satisfies that \mathcal{N} is a structure with domain N and relations R_1, \dots, R_n (of arities r_1, \dots, r_n , respectively), then

$$\mathcal{N}_{\mathcal{M}} =_{\mathrm{df}} \langle N_{\mathcal{M}}, (R_1)_{\mathcal{M}}, \cdots, (R_n)_{\mathcal{M}} \rangle.$$

In either of the above cases $a_{\mathcal{M}}$ is called the \mathcal{M} -externalization of a.

For example, $\omega^{\mathcal{M}}$ denotes the *element* of \mathcal{M} such that $\mathcal{M} \models \phi(\omega)$, where ϕ defines ω . On the other hand, $(\omega^{\mathcal{M}})_{\mathcal{M}}$ denotes the *subset* $\{a \in \mathcal{M} \mid \mathcal{M} \models a \in \omega\}$ of \mathcal{M} , consisting of all a such that $\mathcal{M} \models a \in \omega$.

2.3 Recursive saturation

A type $p(\vec{x})$, over a theory T in a language L, is a set of L-formulas such that $T \cup p$ is consistent when the variables \vec{x} are considered as fresh constant symbols. If $\mathcal{M} \models T$, then $p(\vec{x})$ is realized in \mathcal{M} if there is $\vec{a} \in \mathcal{M}$, such that for all $\phi(\vec{x}) \in p$, we have $\mathcal{M} \models \phi(\vec{a})$. A type $p(\vec{x}, \vec{b})$, over \mathcal{M} with parameters $\vec{b} \in \mathcal{M}$, is a type over $\text{Th}(\mathcal{M}, \vec{b})$ (the theory of \mathcal{M} with parameters \vec{b}). Such a type $p(\vec{x}, \vec{b})$ is recursive if it is a recursive set (under some fixed Gödel coding of the formulas as natural numbers). A structure \mathcal{M} is recursively saturated if it realizes every recursive type over \mathcal{M} . A crsm is a countable recursively saturated model.

Theorem 2.2 (Completeness of the crsm-semantics). Let \mathcal{M} be a countable model in a recursive language. There is a countable recursively saturated elementary extension of \mathcal{M} . In particular, every consistent theory in a recursive language is modelled by a crsm.

Proof. See the proof of Theorem 2.4.1 in Chang & Keisler (1990).

Let \mathcal{M} be a model of a set theory T. The interpretations of the numerals in \mathcal{M} are called the *standard natural numbers of* \mathcal{M} . For each $n < \omega$, let us make the identification $n = \underline{n}^{\mathcal{M}}$. We say that \mathcal{M} is ω -non-standard if there is $c \in (\omega^{\mathcal{M}})_{\mathcal{M}} \setminus \omega$. Such a c is said to be a non-standard number of \mathcal{M} .

Proposition 2.3. If \mathcal{M} is a recursively saturated model of a set theory, then \mathcal{M} is ω -non-standard.

Proof. By recursive saturation, \mathcal{M} realizes the type $\{x \in \mathbb{N}\} \cup \{\underline{n} < x \mid n \in \mathbb{N}\}.$

Suppose that \mathcal{M} is ω -non-standard. We say that a subset A of ω is coded in \mathcal{M} , if there is (a code) $a \in (\omega^{\mathcal{M}})_{\mathcal{M}}$, such that $A = \{n \in \mathbb{N} \mid \mathcal{M} \models \underline{n} < a\}$. We define the standard system of \mathcal{M} as

$$SSy(\mathcal{M}) =_{df} \{ A \subseteq \omega \mid \text{``A is coded in } \mathcal{M}'' \}.$$

The following result is due to Wilmers (1975), employing Friedman's back-and-forth technique (1973):

Theorem 2.4 (Canonicity of countable recursively saturated models). Let \mathcal{M} and \mathcal{N} be crsms modelling ZF. If $\mathcal{M} \equiv_{\mathcal{L}} \mathcal{N}$ and $SSy(\mathcal{M}) = SSy(\mathcal{N})$, then $\mathcal{M} \cong_{\mathcal{L}} \mathcal{N}$.

Proof. See the proof of Theorem 7.14 in Gorbow (2019).

Remark. As far as the authors can see, the proof of the above Theorem requires both the full Separation and Replacement schemas of ZF, as well as its Foundation axiom.

We define the class

$$\mathbf{crsm} =_{\mathrm{df}} \{ \mathcal{M} \mid \mathcal{M} \upharpoonright_{\mathcal{L}} \models \mathsf{ZF} \wedge \text{``} \mathcal{M} \text{ is a crsm''} \}.$$

2.4 Systems of satisfaction over ZF

Before embarking on developing a framework for a notion of truth-in-a-universe relevant to the set-theoretic multiverse, we shall go through some related systems of truth over ZF. An intuitive philosophical perspective is that systems of truth capture various absolute notions of truth, as motivated by the universe view of set theory, while our framework for truth-in-a-universe captures various relative notions of truth, as motivated by the multiverse view of set theory. From a mathematical perspective, it is interesting to relate these two approaches. Moreover, since we will generalize techniques that have been developed for studying systems of truth, these provide a relevant context for viewing our results.

Right at the start of the endeavour to axiomatize truth, one faces the choice between introducing (to the base language of set theory) a unary truth-predicate $\operatorname{Tr}(\sigma)$, applying to sentences σ , or a binary satisfaction-relation $\operatorname{Sat}(\phi,f)$, applying to formulas ϕ and variable-assignments $f:\operatorname{Var}\to V$. In the former option, σ needs to range over a class-sized language where there is a constant-symbol c_x corresponding to each $x\in V$. The authors have chosen the latter option. As a general heuristic, one is justified to expect any theory of satisfaction to be interpretable in a corresponding theory of truth; the idea being to interpret $\operatorname{Sat}(\phi(x),f)$ by $\operatorname{Tr}(\phi[c_{f(x)}/x])$. Fujimoto has made a comprehensive study of theories of truth over set theory, following the former option (2012).

System 2.5 (CT). Let S be a set theory in \mathcal{L}^+ . The system $\mathsf{CT}(S) \upharpoonright$ (for Compositional Truth) consists of these axioms in the language $\mathcal{L}^+_{\mathsf{Sat}}$:

```
\begin{array}{ll} \mathsf{Base} & S \\ \mathsf{CT}_{=} & \forall y_0, y_1 \left( \mathsf{Sat}(\ulcorner x_0 = x_1 \urcorner, \langle \ulcorner x_0 \urcorner, \ulcorner x_1 \urcorner \rangle \mapsto \langle y_0, y_1 \rangle) \leftrightarrow y_0 = y_1 \right) \\ \mathsf{CT}_{\in} & \forall y_0, y_1 \left( \mathsf{Sat}(\ulcorner x_0 \in x_1 \urcorner, \langle \ulcorner x_0 \urcorner, \ulcorner x_1 \urcorner \rangle \mapsto \langle y_0, y_1 \rangle) \leftrightarrow y_0 \in y_1 \right) \\ \mathsf{CT}_{\neg} & \forall \phi \in \mathcal{L}_{\mathsf{Sat}}^+ \forall f \in \mathsf{VA} \left( \mathsf{Sat}(\lnot \phi, f) \leftrightarrow \lnot \mathsf{Sat}(\phi, f) \right) \\ \mathsf{CT}_{\land} & \forall \phi, \psi \in \mathcal{L}_{\mathsf{Sat}}^+ \forall f \in \mathsf{VA} \left( \mathsf{Sat}(\phi \land \psi, f) \leftrightarrow \mathsf{Sat}(\phi, f) \land \mathsf{Sat}(\psi, f) \right) \\ \mathsf{CT}_{\forall} & \forall \phi \in \mathcal{L}_{\mathsf{Sat}}^+ \forall f \in \mathsf{VA} \left( \mathsf{Sat}(\forall u \phi, f) \leftrightarrow \forall g \in \mathsf{VA}_{f,u} \, \mathsf{Sat}(\phi, g) \right) \end{array}
```

We write $\mathsf{CT} \upharpoonright \mathsf{for} \; \mathsf{CT}(\mathsf{ZF}) \upharpoonright$. The axioms of the form CT_{-} are called *compositional axioms*. By basic logic, $\mathsf{CT} \upharpoonright \mathsf{also}$ proves the axioms CT_{\lor} , CT^{\to} and CT_{\exists} (analogously defined). We use phrases such as "Sat is \lor -compositional" to express that we have CT_{\lor} , for example.

```
\mathsf{CT} \text{ is } \mathsf{CT} \upharpoonright + \mathsf{Sep}(\mathcal{L}_{\mathsf{Sat}}) + \mathsf{Rep}(\mathcal{L}_{\mathsf{Sat}}).
```

A routine induction argument in the meta-theory shows this Proposition:

Proposition 2.6. For all
$$\mathcal{L}$$
-formulas $\phi(x)$, $\mathsf{CT} \upharpoonright \mathsf{Fat}(\lceil \phi(x) \rceil, \lceil x \rceil \mapsto y) \leftrightarrow \phi(y)$.

The theory of satisfaction $\mathsf{CT} \upharpoonright + \mathsf{Sep}(\mathcal{L}_{\mathsf{Sat}})$ corresponds to the theory of truth $\mathsf{TC} \upharpoonright + \mathsf{Sep}^+$ in (Fujimoto, 2012, §4). It is straightforward to interpret the former in the latter. Using this, it follows from Theorem 20 in (Fujimoto, 2012, §4.1) that $\mathsf{CT} \upharpoonright + \mathsf{Sep}(\mathcal{L}_{\mathsf{Sat}})$ is conservative over ZF . In contrast, in CT we have access to the Reflection theorem for $\mathcal{L}_{\mathsf{Sat}}$ -formulas, enabling us to prove that there is a V_α modelling ZF . See (Fujimoto, 2012, §4.1) for more details and refinements.

 $\mathbf{Proposition~2.7.~CT} \upharpoonright + \mathsf{Sep}(\mathcal{L}_{\mathsf{Sat}}) \vdash \text{``Transfinite Induction over $\mathcal{L}_{\mathsf{Sat}}$''}$

Proof. Let $\phi(x) \in \mathcal{L}_{\mathsf{Sat}}$. Assuming $\neg \phi(\alpha)$, for some ordinal α , we shall refute the corresponding induction hypothesis. By $\mathsf{Sep}(\mathcal{L}_{\mathsf{Sat}})$, the set $S = \{\xi \leq \alpha \mid \neg \phi(\xi)\}$ exists, and by assumption it is non-empty. Let β be the least ordinal in S. Then $\neg \phi(\beta)$ and $\forall \xi < \beta \phi(\xi)$, refuting the induction hypothesis.

System 2.8 (GR^{ω}). Let S be a set theory in \mathcal{L}^+ . Here we present the systems of iterated Global Reflection over S, denoted $\mathsf{GR}^{\alpha}(S)$, for ordinals $\alpha \leq \omega$. For any set theory T in the language $\mathcal{L}_{\mathsf{Sat}}$, the axiom of Global Reflection over T is

$$\mathsf{GR}_T \quad \forall \phi \in \mathcal{L}^+_{\mathsf{Sat}} \left(\mathrm{Pr}_T(\phi) \to \mathsf{Tr}(\phi) \right).$$

(The dot under T is sometimes omitted, when it can be inferred from the context.) Recursively, for each $\alpha \leq \omega$, we define the system $\mathsf{GR}^{\alpha}(S)$:

$$\begin{split} \mathsf{GR}^0(S) =_{\mathrm{df}} \mathsf{CT}(S)\!\!\upharpoonright + \mathsf{Sep}(\mathcal{L}_{\mathsf{Sat}}^+) \\ \mathsf{GR}^{\alpha+1}(S) =_{\mathrm{df}} \mathsf{GR}^{\alpha}(S) + \mathsf{GR}_{\mathsf{GR}^{\alpha}} \\ \mathsf{GR}^{\omega}(S) =_{\mathrm{df}} \bigcup_{n<\omega} \mathsf{GR}^n(S) \end{split}$$

We write GR^{α} for $\mathsf{GR}^{\alpha}(\mathsf{ZF})$.

Remark. Observe that GR^α is defined with $\mathsf{CT} \upharpoonright + \mathsf{Sep}(\mathcal{L}_\mathsf{Sat})$ as base case, thus without Replacement for formulas with the Satisfaction predicate. The reason for this is that it is intended to express iterated Global Reflection over ZF , which $\mathsf{CT} \upharpoonright + \mathsf{Sep}(\mathcal{L}_\mathsf{Sat})$ conservatively extends. If it were defined with CT as base case, then it would not be morally "over ZF ", since CT proves strong reflection principles of its own.

System 2.9 (FS). The systems FS\ and FS (for Friedman–Sheard) are obtained by adding these rules of proof to CT\ and CT, respectively:

NEC For each
$$\phi \in \operatorname{Sent}(\mathcal{L}_{\mathsf{Sat}})$$
: If $\mathsf{FS} \vdash \phi$, then $\mathsf{FS} \vdash \mathsf{Tr}(\lceil \phi \rceil)$.

CONEC For each $\phi \in \operatorname{Sent}(\mathcal{L}_{\mathsf{Sat}})$: If $\mathsf{FS} \vdash \mathsf{Tr}(\lceil \phi \rceil)$, then $\mathsf{FS} \vdash \phi$.

Recall that $Tr(\theta)$ is defined as $\forall f \in VA Sat(\theta, f)$.

Given a set-theoretic system S in language L, the following rule will be considered:

Reflection rule For each
$$\phi$$
 in L : If $S \vdash \Pr_S(\lceil \phi \rceil)$, then $S \vdash \phi$.

Definition 2.10. A set-theoretic system S in language L is ω -inconsistent if there is an L-formula $\delta(x)$ such that:

$$S \vdash \forall x \ (\Im(x) \to x \in \mathbb{N})$$

$$S \vdash \exists x \ \Im(x)$$
 For each $n \in \mathbb{N}, \ S \vdash \neg \Im(\underline{n})$

We say that $\mathfrak{F}(x)$ witnesses the ω -inconsistency of S^{7}

Proposition 2.11. If ZF is ω -consistent, then ZF is closed under the Reflection rule.

Proof. Suppose $\mathsf{ZF} \not\vdash \phi$. Then $\mathsf{ZF} \vdash \neg \mathsf{Pr}_{\mathsf{ZF}}(\underline{n}, \lceil \phi \rceil)$, for every standard $n \in \mathbb{N}$ (where $\mathsf{Pr}_{\mathsf{ZF}}(\underline{n}, \lceil \phi \rceil)$ is the formula expressing that \underline{n} is the Gödel code of a proof of ϕ). Now, by ω -consistency, $\mathsf{ZF} \vdash \neg \mathsf{Pr}_{\mathsf{ZF}}(\lceil \phi \rceil)$.

Let c be a fresh constant symbol. Note that the schema $\{\underline{m} < c < \omega \mid m \in \mathbb{N}\}$, expressing that there is a non-standard natural number, yields ω -inconsistency when added to a set-theoretic system (proof: take x = c as δ).

⁷The symbol " δ " (Uranus) used here is meant to help the reader remember that its extension consists of non-standard numbers: The arrow may be taken to point outwards from the standard model \mathbb{N} .

Proposition 2.12. GR^{ω} and FS are ω -inconsistent. $\mathsf{GR}^{\omega} + \mathsf{GR}_{\mathsf{GR}^{\omega}}$ and $\mathsf{FS} + \mathsf{GR}_{\mathsf{FS}}$ are inconsistent.

Proof. This is a corollary of McGee's paradox, see McGee (1985), and can be proved analogously as Theorem 13.9 and Corollary 14.39 in Halbach (2014), respectively. The arguments in Halbach (2014) are written for theories of truth over PA, not for theories of satisfaction over ZF. But they go through with these natural modifications:

- 1. Replacing instances of the truth predicate "T" by our defined predicate "Tr", except for instances quantifying over terms, of the form $\forall \vec{t} T(\phi[\vec{t}/\!\!\!/ \vec{x} \ \!\!\!])$, which are replaced by $\forall \vec{y} \in \omega \operatorname{\mathsf{Sat}}(\phi, \lceil \vec{x} \rceil \mapsto \vec{y})$, where \vec{y} is fresh.
- 2. Replacing quantifiers of the form " $\forall x$ " by " $\forall x \in \omega$ ".

These arguments rely on that GR^ω admits NEC, which we proceed to show: Let $\sigma \in \mathsf{Sent}(\mathcal{L}_{\mathsf{Sat}})$ and suppose that $\mathsf{GR}^\omega \vdash \sigma$. Then there is $k < \omega$, such that $\mathsf{GR}^k \vdash \sigma$. since this proof can be represented in GR^ω , we have $\mathsf{GR}^\omega \vdash \mathsf{Pr}_{\mathsf{GR}^k}(\lceil \sigma \rceil)$, so that by Global Reflection, $\mathsf{GR}^\omega \vdash \mathsf{Tr}(\lceil \sigma \rceil)$. Since $\mathsf{GR}^\omega \vdash \mathsf{CT}$, it follows that $\mathsf{GR}^\omega \vdash \sigma$, as desired.

So if we were to naturally extend the definition of GR^{α} to all ordinals α , then we would get that GR^{α} is inconsistent for all $\alpha > \omega$.

Later on we will introduce a multiverse axiom, called Self-Perception, to the effect that the universe of the background theory is isomorphic (over \mathcal{L}) to one of its internal universes; this axiom is motivated by the idea that the universe of the background theory should be available in its multiverse. The following lemmas establish technical results needed to validate that axiom.

Lemma 2.13. Let \mathcal{U} be a crsm of GR_0 and let $\mathcal{V} \in \mathcal{U}$, such that \mathcal{U} satisfies

$$\mathcal{V} \models \{ \sigma \in \operatorname{Sent}(\mathcal{L}_{\mathsf{Sat}}) \mid \mathsf{Tr}(\sigma) \}.$$

Then $\mathcal{U} \cong_{\mathcal{L}} \mathcal{V}_{\mathcal{U}}$.

Recall that $\mathcal{V}_{\mathcal{U}}$ is the \mathcal{U} -externalization of \mathcal{V} , see Definition 2.1.

Proof. We shall establish $\mathcal{U} \cong_{\mathcal{L}} \mathcal{V}_{\mathcal{U}}$ by invoking Theorem 2.4. Thus we need to show that $\mathcal{V}_{\mathcal{U}}$ is a crsm, that $SSy(\mathcal{U}) = SSy(\mathcal{V}_{\mathcal{U}})$ and that $\mathcal{U} \equiv_{\mathcal{L}} \mathcal{V}_{\mathcal{U}}$.

Note that \mathcal{U} is ω -non-standard, by $\mathcal{U} \in \mathbf{crsm}$ and Proposition 2.3. That $\mathcal{V}_{\mathcal{U}}$ is a crsm now follows from Lemma 2.2 in Gitman & Hamkins (2010).⁸

 $(\omega^{\mathcal{U}})_{\mathcal{U}}$ is mapped initially into $(\omega^{(\mathcal{V}_{\mathcal{U}})})_{(\mathcal{V}_{\mathcal{U}})}$ by an embedding j (for each $x \in (\omega^{\mathcal{U}})_{\mathcal{U}}$, j(x) is defined as the unique $y \in \mathcal{U}$ such that $\mathcal{U} \models y = \underline{x}^{\mathcal{V}}$). Therefore, we obtain $SSy(\mathcal{U}) = SSy(\mathcal{V}_{\mathcal{U}})$ as follows: Let $A \in SSy(\mathcal{U})$, coded by $a \in (\omega^{\mathcal{U}})_{\mathcal{U}}$. Since j is an embedding, j(a) is a code for A in $(\omega^{(\mathcal{V}_{\mathcal{U}})})_{(\mathcal{V}_{\mathcal{U}})}$. Conversely, let $B \in SSy(\mathcal{V}_{\mathcal{U}})$, coded by $b \in (\omega^{(\mathcal{V}_{\mathcal{U}})})_{(\mathcal{V}_{\mathcal{U}})}$. Since j is initial and \mathcal{U} is ω -non-standard, there is a non-standard $c \in (\omega^{\mathcal{U}})_{\mathcal{U}}$, such that $j(c) \leq^{\mathcal{V}_{\mathcal{U}}} b$. So since j is an embedding, c is a code for b in $(\omega^{\mathcal{U}})_{\mathcal{U}}$.

To see $\mathcal{U} \equiv_{\mathcal{L}} \mathcal{V}_{\mathcal{U}}$, let ϕ be a sentence of \mathcal{L} . By absoluteness of \models for standard formulas,

$$\mathcal{V}_{\mathcal{U}} \models \phi \iff \mathcal{U} \models "\mathcal{V} \models (\lceil \phi \rceil)".$$

Since Sat satisfies the Tarski-biconditionals for \mathcal{L} , we have

$$\mathcal{U} \models \phi \iff \mathcal{U} \models \mathsf{Tr}(\lceil \phi \rceil).$$

Moreover, by CT¬ and the condition of the Lemma,

$$\mathcal{U} \models \mathsf{Tr}(\lceil \phi \rceil) \iff \mathcal{U} \models "\mathcal{V} \models (\lceil \phi \rceil)".$$

By combining these we obtain $\mathcal{U} \equiv_{\mathcal{L}} \mathcal{V}_{\mathcal{U}}$, as desired.

⁸That Lemma is stated for ZFC, but it is easily seen that its proof does not make use of Choice.

Remark. $\mathcal{U} \models \mathsf{ZF}$ is needed for the proof of this lemma, as it relies on Theorem 2.4. Thus, the authors do not expect it to generalize to $\mathsf{GR}_0(S)$, unless $S \vdash \mathsf{ZF}$.

Lemma 2.14. Let $k < \alpha \le \omega$, and let $\mathcal{U} \models \mathsf{GR}^{\alpha}$. Then there is $\mathcal{V} \in \mathcal{U}$, such that \mathcal{U} satisfies

$$\mathcal{V} \in \mathbf{crsm} \wedge \mathcal{V} \models \{ \sigma \in \mathrm{Sent}(\mathcal{L}_{\mathsf{Sat}}) \mid \mathsf{Tr}(\sigma) \}.$$

In particular, \mathcal{U} satisfies that $\mathcal{V} \models \mathsf{GR}^k$. Moreover, if $\mathcal{U} \in \mathbf{crsm}$, then $\mathcal{U} \cong_{\mathcal{L}} \mathcal{V}_{\mathcal{U}}$.

Proof. We work in \mathcal{U} . From $\mathsf{ZF} + \mathsf{Sep}(\mathcal{L}_{\mathsf{Sat}})$ we get that the set $\mathbf{Tr} = \{\phi \in \mathsf{Sent}(\mathcal{L}_{\mathsf{Sat}}) \mid \mathsf{Tr}(\phi)\}$ exists. For the first statement, by completeness of the crsm-semantics (Theorem 2.2), it suffices to establish $\mathsf{Con}(\mathbf{Tr})$; and for this it suffices to establish $\mathsf{Con}(\sigma)$, where σ is an arbitrary finite conjunction of sentences in \mathbf{Tr} . By \land -compositionality of Sat , we have $\mathsf{Sat}(\sigma)$. By $\mathsf{GR}_{\mathsf{ZF}+\mathsf{Sep}(\mathcal{L}_{\mathsf{Sat}})}$, we have $\mathsf{Pr}_{\mathsf{ZF}+\mathsf{Sep}(\mathcal{L}_{\mathsf{Sat}})}(\neg\sigma) \to \mathsf{Sat}(\neg\sigma)$, and by \neg -compositionality of Sat , we have $\mathsf{Sat}(\neg\sigma) \leftrightarrow \neg\mathsf{Sat}(\sigma)$. So since $\mathsf{Sat}(\sigma)$, we obtain $\mathsf{Pr}_{\mathsf{ZF}+\mathsf{Sep}(\mathcal{L}_{\mathsf{Sat}})}(\neg\sigma) \to \bot$, whence $\mathsf{Con}(\sigma)$. By Theorem 2.2, we can let $\mathcal V$ be a model of Tr in crsm .

It follows from $\mathsf{GR}_{\mathsf{GR}^k}$ that $\mathsf{GR}^k \subseteq \mathbf{Tr}$, yielding the second statement.

The last statement now follows from Lemma 2.13.

Axiom 2.15. Let ι be a function symbol and self be a constant symbol. Iso(x) denotes an \mathcal{L}_{ι}^+ -formula expressing that x is an \mathcal{L}^+ -structure and that ι is an \in -isomorphism from the universe V onto x. We shall study this axiom in $\mathcal{L}_{\iota,\text{self}}^+$:

(This formulation is chosen over $\exists x \operatorname{Iso}(x)$, as it is convenient to have a reference to a witness.)

By the \in -isomorphism property, and the absoluteness of \models for standard formulas, we have:

Proposition 2.16. For each $\phi(\vec{x}) \in \mathcal{L}$, $\mathsf{ZF} + \mathsf{Iso}(\mathsf{self}) \vdash (\mathsf{self} \models \lceil \phi(\iota(\vec{x})) \rceil) \leftrightarrow \phi(\vec{x})$.

System 2.17. Let S be a set theory in \mathcal{L}^+ . By recursion, for each $\alpha \leq \omega$, we define the system $\mathsf{SP}^{\alpha}(S)$ (standing for Self-Perception):

$$\begin{split} \mathsf{SP}^0(S) =_{\mathrm{df}} \mathsf{GR}^0(S) &= \mathsf{CT}(S) \!\!\upharpoonright + \mathsf{Sep}(\mathcal{L}^+_{\mathsf{Sat}}) \\ \mathsf{SP}^{\alpha+1}(S) =_{\mathrm{df}} \mathsf{GR}^{\alpha+1}(S) + \mathsf{self} \in \mathbf{crsm} + \mathsf{Iso}(\mathsf{self}) + \mathsf{self} \models \mathbf{Tr} \cup \mathsf{S}\underline{\mathsf{P}}^{\underline{\alpha}}(S) \\ \mathsf{SP}^{\omega}(S) =_{\mathrm{df}} \bigcup_{n < \omega} \mathsf{SP}^n(S), \end{split}$$

where $\mathbf{Tr} =_{\mathrm{df}} \{ \sigma \in \mathrm{Sent}(\mathcal{L}_{\mathsf{Sat}}) \mid \mathsf{Tr}(\sigma) \}$. We write SP^{α} for $\mathsf{SP}^{\alpha}(\mathsf{ZF})$.

Remark. A clarificatory note on the role of the languages in SP^{α} . Let $\alpha \geq 1$. The language of SP^{α} is $\mathcal{L}_{\mathsf{Sat},\iota,\mathsf{self}}$. SP^{α} proves Separation for $\mathcal{L}_{\mathsf{Sat},\mathsf{self}}$, and it proves Replacement for $\mathcal{L}_{\mathsf{self}}$ (since self can be treated as a parameter in these schemas).

Lemma 2.18. Let $\alpha \leq \omega$. Every $\operatorname{crsm} \mathcal{U}$ of $\operatorname{\mathsf{GR}}^{\alpha}$ expands to a model \mathcal{U}^* of $\operatorname{\mathsf{SP}}^{\alpha}$. Moreover, if \mathcal{U} is a definable model, then we can also obtain that \mathcal{U}^* is definable.

Proof. We start by showing the case $\alpha < \omega$ by induction. The base case $\alpha = 0$ is trivial. The induction hypothesis is that if $\mathcal{U} \models \mathsf{GR}^{\alpha}$, then \mathcal{U} expands to a model of SP^{α} ; and if \mathcal{U} is definable, then the expansion is definable. Let $\mathcal{U} \models \mathsf{GR}^{\alpha+1}$. Applying Lemma 2.14, we find a crsm \mathcal{V} in \mathcal{U} , such that $\mathcal{U} \cong_{\mathcal{L}} \mathcal{V}_{\mathcal{U}}$ (as witnessed by an \mathcal{L} -isomorphism $i: \mathcal{U} \to \mathcal{V}_{\mathcal{U}}$) and \mathcal{U} satisfies that $\mathcal{V} \models \mathbf{Tr} \cup \mathsf{GR}^{\underline{\alpha}}$. So by the induction hypothesis applied in \mathcal{U} , \mathcal{U} satisfies that \mathcal{V} expands

to a model W of $SP^{\underline{\alpha}}$. Let U^* be the model obtained from U by interpreting self by W and interpreting ι by i. It is immediate from the construction that $U^* \models SP^{\alpha+1}$.

Now to the case that $\alpha = \omega$: Assume that $\mathcal{U} \models \mathsf{GR}^{\omega}$. Working in \mathcal{U} , let $\mathbf{Tr} = \{\phi \in \mathsf{Sent}(\mathcal{L}_{\mathsf{Sat}}) \mid \mathsf{Tr}(\phi)\}$. By the above,

$$\{x \in \mathbf{crsm} \land x \models \mathbf{Tr} \cup \mathsf{SP}^{\underline{n}} \mid n < \omega\}$$

is a recursive type over \mathcal{U} . So since \mathcal{U} is recursively saturated, it is realized by some \mathcal{W} in \mathcal{U} . It now follows from Lemma 2.13 that $\mathcal{U} \cong_{\mathcal{L}} \mathcal{W}_{\mathcal{U}}$. So just as in the former case, \mathcal{U} can be expanded to a model \mathcal{U}^* of SP^ω .

Assume now that \mathcal{U} is definable. \mathcal{W} can then be defined as the least element of a definable enumeration of \mathcal{U} that satisfies the appropriate conditions. An isomorphism witnessing $\mathcal{U} \cong_{\mathcal{L}} \mathcal{W}_{\mathcal{U}}$ can also be defined: As seen from a close look at the proof of Theorem 2.4, the isomorphism is constructed by recursion on enumerations of \mathcal{U} and $\mathcal{W}_{\mathcal{U}}$, both of which can be chosen definable since \mathcal{U} and $\mathcal{W}_{\mathcal{U}}$ are definable and countable.

Lemma 2.19. GR^{ω} interprets SP^{ω} .

Proof. Since GR^{ω} is ω -inconsistent, there is a formula $\mathfrak{F}(x)$ such that $\mathsf{GR}^{\omega} \vdash \exists x < \omega \, \mathfrak{F}(x)$, but for each $n \in \mathbb{N}$, $\mathsf{GR}^{\omega} \vdash \neg \mathfrak{F}(n)$.

We start by working in GR^{ω} . By $\mathsf{ZF} + \mathsf{Sep}(\mathcal{L}_{\mathsf{Sat}})$, the theory $\mathsf{Tr} = \{\sigma \in \mathsf{Sent}(\mathcal{L}_{\mathsf{Sat}}) \mid \mathsf{Tr}(\sigma)\}$ of truth is a set; and by the argument starting the proof of Lemma 2.14, it is a consistent theory. So there is a definable $\mathcal{S} \in \mathsf{crsm}$ that models truth. Since \mathcal{S} is a model of truth, the Global Reflection axioms allow us to prove $\mathcal{S} \models \mathsf{GR}^{\underline{n}}$, for each standard natural number n. Let $d < \omega$ be the minimal number such that $\mathfrak{S}(d)$, and let c be the maximal number such that $c \leq d$ and $\mathcal{S} \models \mathsf{GR}^c$. Note that for each standard natural number n, we can prove $\underline{n} < c$. By Lemma 2.18, \mathcal{S} expands to a definable model \mathcal{S}' of SP^c .

Working in the meta-theory, it follows that GR^{ω} interprets SP^{ω} by an interpretation \mathcal{J} mapping each sentence σ in the language of SP^{ω} to the $\mathcal{L}_{\mathsf{Sat}}$ -sentence $\mathcal{S}' \models \ulcorner \sigma \urcorner$.

3 Revision-semantic truth-in-a-universe

We shall now go through the key revision-semantic technique introduced in this paper, which may be used to construct a variety of untyped truth-in-a-universe relations for the multiverse of set theory. Revision-semantics was independently invented in Gupta (1982), Herzberger (1982a) and Herzberger (1982b). In Friedman & Sheard (1987), the axiomatic theory of truth FS was presented and shown to be validated by a model constructed through such a revision process. The revision process starts with an arbitrary extension S_0 of truth, and recursively defines S_{n+1} as the theory of the structure (\mathbb{N}, S_n) . In particular, the theory of \mathbb{N} is a subset of S_1 , and the liar sentence is in S_{n+1} iff it is not in S_n .

The construction in this paper is somewhat different in that it is intensional. We start with a more-or-less arbitrary formula *defining* truth-in-a-universe, and revise the definition in a revision-semantic fashion. The construction can be modified by adjusting parameters. For example, we shall see that certain conditions on the parameters result in that the eventual definition of truth-in-a-universe validates the multiverse theory MS, introduced in System 4.5. This theory is analogous to FS, but is actually weaker. The parameters need to satisfy some basic conditions as specified in this definition:

⁹This follows from the proof of Theorem 2.4.1 in Chang & Keisler (1990). The key observation to see that the model is definable is that it is essentially a Henkin-construction by recursion on an enumeration of a recursive language and on an enumeration of the set of all recursive subsets of that language, both of which can be chosen definable.

Definition 3.1. Let $T_n(\phi)$, $Uni_n(\mathcal{U})$ and $Mod_0(\mathcal{U}, \phi, f)$ be formulas of the meta-language (\mathcal{L}) , in the free variables $\{n, \phi\}$, $\{n, \mathcal{U}\}$ and $\{\mathcal{U}, \phi, f\}$, respectively.¹⁰ For each $n \in \mathbb{N}$:

$$\mathbf{T}_{n} =_{\mathrm{df}} \{ \phi \mid \mathbf{T}_{n}(\phi) \}$$

$$\mathbf{T}_{\omega} =_{\mathrm{df}} \bigcup_{n < \omega} \mathbf{T}_{n}$$

$$\mathbf{Uni}_{n} =_{\mathrm{df}} \{ \mathcal{U} \mid \mathbf{Uni}_{n}(\mathcal{U}) \}$$

$$\mathbf{Uni}_{\omega} =_{\mathrm{df}} \bigcap_{n < \omega} \mathbf{Uni}_{n}$$

 $(\mathbf{T}_n)_{n\in\mathbb{N}}$ is intended to be a sequence of first-order set theories, and $(\mathbf{Uni}_n)_{n\in\mathbb{N}}$ is intended to be a sequence of classes of models, as formally specified below. Let \mathcal{L}^T , \mathcal{L}^{Rev} be recursive languages. We say that T, Uni, Mod₀ are revision parameters (in \mathcal{L}^T , \mathcal{L}^{Rev}) if the following conditions (closed under $\forall n \in \mathbb{N}$, where appropriate) are provable in the meta-theory and in a given set theory (as object theory):

- 1. $\mathcal{L} \subseteq \mathcal{L}^{Rev} \subseteq \mathcal{L}^{T}$
- 2. "The symbols Uni, Mod do not appear in \mathcal{L}^{T} ."
- 3. " \mathbf{T}_0 is a set theory in \mathcal{L}^{T} ."
- 4. $\mathbf{T}_{n+1} \vdash \mathbf{T}_n$
- 5. $Uni_0(\mathcal{U}) \to "\mathcal{U} \text{ is an } \mathcal{L}^T\text{-structure."}$
- 6. $\mathbf{Uni}_{n+1} \subseteq \mathbf{Uni}_n$
- 7. $\operatorname{Mod}_0(\mathcal{U}, \phi, f) \to \operatorname{Uni}_0(\mathcal{U}) \land \phi \in \mathcal{L}^{\operatorname{Rev}}_{\operatorname{Uni} \operatorname{\mathsf{Mod}}} \land f \in \operatorname{VA}^{\mathcal{U}}$

In the construction below we shall see how, given revision parameters, an untyped revision-semantic truth-in-a-universe predicate can be defined as an \mathcal{L} -formula $\mathrm{Mod}_n(\mathcal{U}, \phi, f)$, with this intended reading of the variables: n is the stage in the revision process, \mathcal{U} is a universe, ϕ is a formula in (the representation of) $\mathcal{L}^{\mathrm{Rev}}_{\mathsf{Uni},\mathsf{Mod}}$ and f is an assignment of variables to elements of \mathcal{U} . So \mathcal{L}^{T} is the language of the theories \mathbf{T}_n , $\mathcal{L}^{\mathrm{Rev}}$ is any sublanguage of \mathcal{L}^{T} , and $\mathcal{L}^{\mathrm{Rev}}_{\mathsf{Uni},\mathsf{Mod}}$ is the language undergoing revision. The \mathcal{L} -formula $\mathsf{Mod}^{\circlearrowright}$ is also introduced as a variant of Mod_n . Actually, only the Uni_n and Mod_0 parameters influence the construction. The T_n parameter comes into play later on, in the Main Lemma (in §5), where we show (under certain conditions on the revision parameters) that the Mod_n formula satisfies desirable semantically motivated axioms when constructed in the theory T_ω .

The construction may intuitively be thought of as a recursive procedure, where Mod_0 is a more-or-less arbitrary truth-in-a-universe relation and each Mod_{n+1} revises Mod_n into a more adequate relation. It turns out to be efficient to perform the construction using Gödel's fixed-point lemma. It is in fact possible to choose Mod_0 such that it gets revised to itself (see the definition of $\text{Mod}^{\circlearrowright}$ below). This phenomenon contrasts with the revision-semantics ordinarily used to construct a model of the Friedman–Sheard theory of truth, where the revision-operation has no fixed-point (see Lemma 14.9(iii) in Halbach (2014)). A key difference between the present revision-process and that one is that the former is intensional and the latter is extensional. In the present framework we start with an arbitrary formula defining truth-in-a-universe and revise it to

 $^{^{10}}$ Even though the n is notationally in subscript-position, it is a free variable of the formulas T and Uni. This pattern will also be used for the formula Mod introduced below.

more adequate definitions, while in the other framework one starts with an arbitrary *extension* of truth and revise it more adequate extensions. The move from extensional to intensional revision-semantics is highly relevant for the present framework.

Construction 3.2 (Construction of Revision-semantics for the Multiverse). Let $T_n(\phi)$, $Uni_n(\mathcal{U})$ and $Mod_0(\mathcal{U}, \phi, f)$ be revision parameters.

By Gödel's fixed-point lemma, there is an \mathcal{L} -formula $\operatorname{Mod}_n(\mathcal{U}, \phi, f)$, in the free variables n, \mathcal{U}, ϕ, f , such that provably:

$$\operatorname{Mod}_{n}(\mathcal{U}, \phi, f) \leftrightarrow \begin{pmatrix} n \in \mathbb{N} \wedge \operatorname{Uni}_{n}(\mathcal{U}) \wedge \phi \in \mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}^{\mathsf{Rev}} \wedge f \in \operatorname{VA}^{\mathcal{U}} \\ \wedge (n = 0 \to \operatorname{Mod}_{0}(\mathcal{U}, \phi, f)) \\ \wedge (n > 0 \to \langle \mathcal{U} |_{\mathcal{L}^{\mathsf{Rev}}}, \lceil \operatorname{Uni}_{\underline{n-1}} \rceil^{\mathcal{U}}, \lceil \operatorname{Mod}_{\underline{n-1}} \rceil^{\mathcal{U}} \rangle \models (\phi, f)) \end{pmatrix}$$
 (†)

Recall that if $\phi(\vec{x})$ is a formula in the language of a structure \mathcal{M} , then $\phi^{\mathcal{M}} = \{\vec{a} \in \mathcal{M} \mid \mathcal{M} \models \phi(\vec{a})\}$. Above, this notation is used for formulas in a represented language, hence the Gödel quotes, $\ulcorner \urcorner$. Working in ZF , $\langle \mathcal{U} \upharpoonright_{\mathcal{L}^{\mathsf{Rev}}}, \ulcorner \mathsf{Uni}_{\underline{n-1}} \urcorner^{\mathcal{U}}, \ulcorner \mathsf{Mod}_{\underline{n-1}} \urcorner^{\mathcal{U}} \rangle$ is the expansion of $\mathcal{U} \upharpoonright_{\mathcal{L}^{\mathsf{Rev}}}$ to $\mathcal{L}^{\mathsf{Rev}}_{\mathsf{Uni},\mathsf{Mod}}$, interpreting $\ulcorner \mathsf{Mod} \urcorner$ by $\{\vec{a} \in \mathcal{U} \mid \mathcal{U} \models \ulcorner \mathsf{Mod}_{\underline{n-1}} \urcorner(\vec{a})\}$, and interpreting $\ulcorner \mathsf{Uni} \urcorner$ by $\{u \in \mathcal{U} \mid \mathcal{U} \models \ulcorner \mathsf{Uni}_{n-1} \urcorner(u)\}$.

Formally, we now have two references for the expression " Mod_0 ", the formula Mod_0 and the formula Mod_n , with the variable assignment $n \mapsto 0$. However, it is clear that these are equivalent.

The above construction works for a very wide range of choices for Mod_0 . But by the fixed-point lemma, we can choose Mod_0 to be "equivalent to its own revision", so that Mod_n turns out to be constant with respect to n. Indeed, there is an \mathcal{L} -formula $\text{Mod}^{\circlearrowleft}$, such that provably:

$$\operatorname{Mod}^{\circlearrowright}(\mathcal{U}, \phi, f) \leftrightarrow \begin{pmatrix} \operatorname{Uni}_{0}(\mathcal{U}) \land \phi \in \mathcal{L}^{\operatorname{Rev}}_{\operatorname{Uni}, \operatorname{\mathsf{Mod}}} \land f \in \operatorname{VA}^{\mathcal{U}} \\ \land \langle \mathcal{U} \upharpoonright_{\mathcal{L}^{\operatorname{Rev}}}, \ulcorner \operatorname{Uni}_{\underline{0}} \urcorner^{\mathcal{U}}, \ulcorner \operatorname{Mod}^{\circlearrowright \urcorner^{\mathcal{U}}} \rangle \models (\phi, f) \end{pmatrix}$$

End of Construction.

For the reader familiar with the extensional revision-procedure used to construct a model of FS (this construction is reasonably well-known for FS formulated over PA, see (Halbach, 2014, ch. 14.1) for a detailed treatment), note how the recursive call in the fixed-point formula of our construction operates on an intension (the formula Mod), which is interpreted in internal models. In contrast, the recursive call of the revision procedure for constructing a model of FS (say as a theory of truth over arithmetic) operates on an extension (the set of true sentences from the previous step), which is obtained from the external model. The authors take this to explain why it is possible to define a truth-in-a-universe relation Mod[©] which is fixed by the revision-procedure. In the extensional revision-semantics, this is not possible simply because the liar sentence must switch truth-value in the external model at every step of the revision.

Necessity was the mother of the intensional revision-semantics of this paper; the authors do not see any way to construct models of "the Copernican multiverse of sets" (as formalized by various theories in this paper, e.g. CM + Non-Triviality + NEC) by the extensional approach. Conversely, the authors do not see that the intensional approach can replace the extensional approach, as the former relies on that the intension acted upon in the recursive call is interpreted in an internal model. For arithmetic this may be a serious obstacle, as arithmetic does not have that kind of internal models.

Some conditions and rules for revision parameters, relevant for showing that the Mod_n formula constructed as above satisfies a desirable semantically motivated theory (see the Main Lemma in §5), are shown in Figure 3.1. If one of the rules holds, we say that the revision parameters admit

$\begin{array}{c} \text{Revision rules} \\ \text{NEC}^* & \forall n \in \mathbb{N} \ \forall \phi \in \mathcal{L}^{\text{Rev}} \left((\mathbf{T}_n \vdash \phi) \rightarrow (\mathbf{T}_{n+1} \vdash \forall \mathcal{U} \in \mathbf{Uni}_{n+1} \ (\mathcal{U} \models \ulcorner \phi \urcorner)) \right) \\ \text{CONEC}^* & \forall n \in \mathbb{N} \ \forall \phi \in \mathcal{L}^{\text{Rev}} \left((\mathbf{T}_{n+1} \vdash \forall \mathcal{U} \in \mathbf{Uni}_{n+1} \ (\mathcal{U} \models \ulcorner \phi \urcorner)) \rightarrow \mathbf{T}_{n+2} \vdash \phi \right) \\ \text{Reflection rule}^* & \forall n \in \mathbb{N} \ \forall \phi \in \mathcal{L}^{\text{Rev}} \left((\mathbf{T}_{n+1} \vdash \Pr_{\mathbf{T}_n} (\ulcorner \phi \urcorner)) \rightarrow \mathbf{T}_{n+2} \vdash \phi \right) \\ \\ \textbf{Revision conditions} \\ \text{Soundness}^* & \forall n \in \mathbb{N} \ \forall \mathcal{U} \in \mathbf{Uni}_{n+1} \ (\mathcal{U} \models \mathbf{T}_n) \\ \text{Completeness}^* & \forall n \in \mathbb{N} \ \forall \phi \in \mathcal{L}^{\text{Rev}} \left((\forall \mathcal{U} \in \mathbf{Uni}_{n+1} \ (\mathcal{U} \models \phi)) \rightarrow \mathbf{T}_n \vdash \phi \right) \\ \end{array}$

Figure 3.1: Rules and conditions for the revision parameters

it. Essentially, if the revision parameters admit NEC* or CONEC*, then the multiverse theory interpreted admits NEC or CONEC, respectively. In practice it is often easier to work with the other rule and conditions in Figure 3.1, using this Lemma:

Lemma 3.3. Let T, Uni, Mod_0 be revision parameters.

- (a) If Soundness* is provable, then the parameters admit NEC*.
- (b) If Completeness* is provable and the parameters admit the Reflection rule*, then the parameters admit CONEC*.
- *Proof.* (a) Let $n \in \mathbb{N}$ and $\phi \in \mathcal{L}^{Rev}$. Assume that Soundness* is provable (in the meta-theory ZF) and that $\mathbf{T}_n \vdash \phi$. By the latter, encoding the proof in ZF, we have $\mathsf{ZF} \vdash \mathsf{Pr}_{\mathbf{T}_n}(\lceil \phi \rceil)$. Combining these, we get $\mathsf{ZF} \vdash \forall \mathcal{U} \in \mathbf{Uni}_{n+1} \ (\mathcal{U} \models \lceil \phi \rceil)$. Since $\mathbf{T}_{n+1} \vdash \mathsf{ZF}$, we are done.
- (b) Let $n \in \mathbb{N}$ and $\phi \in \mathcal{L}^{Rev}$. Assume that Completeness* is provable and the parameters admit the Reflection rule*. Since $\mathbf{T}_{n+1} \vdash \mathsf{ZF}$, we have $\mathbf{T}_{n+1} \vdash \mathsf{Completeness}^*$. Now suppose that $\mathbf{T}_{n+1} \vdash \forall \mathcal{U} \in \mathbf{Uni}_{n+1} \ (\mathcal{U} \models \lceil \phi \rceil)$. Then $\mathbf{T}_{n+1} \vdash \Pr_{\mathbf{T}_n}(\lceil \phi \rceil)$. So by the Reflection rule*, $\mathbf{T}_{n+2} \vdash \phi$, as desired.

4 Theories of untyped satisfaction for the multiverse

Section 3 showed how a revision-semantic relation of truth-in-a-universe can be constructed in set theory. We turn now to the task of finding appropriate axioms for truth-in-a-universe that are validated by such revision-constructions.

System 4.1 (CM). CM⁻, standing for *Compositional satisfaction for the Multiverse*, is axiomatized as follows:

```
 \begin{array}{ll} \mathsf{Base} & \mathsf{ZF} + \mathsf{Sep}(\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}) + \mathsf{Rep}(\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}) \\ \mathsf{CM}_{=} & \forall \mathcal{U} \in \mathsf{Uni} \, \forall f \in \mathsf{VA}^{\mathcal{U}} \left( \mathsf{Mod}(\mathcal{U}, \ulcorner x = y \urcorner, f) \leftrightarrow f(x) = f(y)) \right) \\ \mathsf{CM}_{\neg} & \forall \mathcal{U} \in \mathsf{Uni} \, \forall \phi \in \mathcal{L}_{\mathsf{Uni},\mathsf{Mod}} \, \forall f \in \mathsf{VA}^{\mathcal{U}} \left( \mathsf{Mod}(\mathcal{U}, \lnot \phi, f) \leftrightarrow \lnot \mathsf{Mod}(\mathcal{U}, \phi, f) \right) \\ \mathsf{CM}_{\land} & \forall \mathcal{U} \in \mathsf{Uni} \, \forall \phi, \psi \in \mathcal{L}_{\mathsf{Uni},\mathsf{Mod}} \, \forall f \in \mathsf{VA}^{\mathcal{U}} \left( \mathsf{Mod}(\mathcal{U}, \phi \land \psi, f) \leftrightarrow (\mathsf{Mod}(\mathcal{U}, \phi, f) \land \mathsf{Mod}(\mathcal{U}, \psi, f)) \right) \\ \mathsf{CM}_{\forall} & \forall \mathcal{U} \in \mathsf{Uni} \, \forall \phi \in \mathcal{L}_{\mathsf{Uni},\mathsf{Mod}} \, \forall f \in \mathsf{VA}^{\mathcal{U}} \left( \mathsf{Mod}(\mathcal{U}, \psi, \phi, f) \leftrightarrow \forall g \in \mathsf{VA}^{\mathcal{U}}_{f,u} \, \mathsf{Mod}(\mathcal{U}, \phi, g) \right) \\ \end{array}
```

Define $ZF_{Uni,Mod} =_{df} ZF + Sep(\mathcal{L}_{Uni,Mod}) + Rep(\mathcal{L}_{Uni,Mod})$. We write CM for CM⁻ plus the axiom:

 $\mathsf{Multiverse_{\mathsf{ZF}}} \quad \forall \mathcal{U} \in \mathsf{Uni} \, \forall \sigma \in \mathsf{ZF}_{\mathsf{Uni},\mathsf{Mod}} \, \mathsf{Mod}(\mathcal{U},\sigma)$

If \mathcal{L}' expands \mathcal{L} , then we write $\mathsf{CM}^-(\mathcal{L}')$ and $\mathsf{CM}(\mathcal{L}')$ for the corresponding systems obtained by replacing all occurrences of $\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}$ in the axioms of the form CM_- above by the language $\mathcal{L}'_{\mathsf{Uni},\mathsf{Mod}}$. (So the Separation and Replacement schemas remain unchanged, ranging only over $\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}$.)

Remark. The natural analogue axioms CM_{\lor} , CM_{\to} , CM_{\exists} are easily derived in CM^{-} .

Remark. In CM⁻, each $\mathcal{U} \in \mathsf{Uni}$ may be viewed as an $\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}$ -structure, by performing this assignment:

$$\begin{split} & \in^{\mathcal{U}} =_{\mathrm{df}} \big\{ \langle a,b \rangle \mid \mathsf{Mod}(\mathcal{U}, \ulcorner x \in y \urcorner, \langle x,y \rangle \mapsto \langle a,b \rangle) \big\} \\ & \mathsf{Uni}^{\mathcal{U}} =_{\mathrm{df}} \big\{ a \mid \mathsf{Mod}(\mathcal{U}, \ulcorner \mathsf{Uni}(x) \urcorner, x \mapsto a) \big\} \\ & \mathsf{Mod}^{\mathcal{U}} =_{\mathrm{df}} \big\{ \langle a,b,c \rangle \mid \mathsf{Mod}(\mathcal{U}, \ulcorner \mathsf{Mod}(x,y,z) \urcorner, \langle x,y,z \rangle \mapsto \langle a,b,c \rangle) \big\} \end{split}$$

Accordingly, we will occasionally use the notation $\mathcal{U} \models \phi$ for satisfaction in that $\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}^-}$ structure. Using the compositional axioms of CM^- , it is easily shown that $\mathsf{Mod}(\mathcal{U},\phi,f) \iff \mathcal{U} \models (\phi,f)$.

In applications, it is natural to add further axioms to CM⁻, ensuring e.g. that we can prove:

Non-Triviality
$$\exists \mathcal{U} \mathsf{Uni}(\mathcal{U})$$

Note that over CM^- , Non-Triviality is equivalent to $Tr^{\square}(\lceil \bot \rceil) \to \bot$. Recall that the formulas $Tr^{\square}(\sigma)$ and $Tr^{\diamondsuit}(\sigma)$ are defined as $\forall \mathcal{U} \in \mathsf{Uni}\,(\mathsf{Mod}(\mathcal{U},\sigma))$ and $\neg \mathsf{Tr}^{\square}(\neg \sigma)$, respectively. We may naturally consider the interpretation of the modal operators \square, \diamondsuit , generated by interpreting $\square \sigma$ by $Tr^{\square}(\lceil \sigma \rceil)$. Therefore, it is useful to exhibit some compositional conditions easily provable for Tr^{\square} in CM^- and $CM^- + \mathsf{Non-trivility}$:

Proposition 4.2. CM⁻ proves:

```
 \begin{array}{ll} \mathsf{CM}^{\square}_{\to} & \forall \phi, \psi \in \mathrm{Sent}(\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}) \left( \mathsf{Tr}^{\square}(\phi {\to} \psi) \to (\mathsf{Tr}^{\square}(\phi) \to \mathsf{Tr}^{\square}(\psi)) \right) \\ \mathsf{CM}^{\square}_{\to} & \forall \phi, \psi \in \mathrm{Sent}(\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}) \left( \mathsf{Tr}^{\square}(\phi {\leftrightarrow} \psi) \to (\mathsf{Tr}^{\square}(\phi) \leftrightarrow \mathsf{Tr}^{\square}(\psi)) \right) \\ \mathsf{CM}^{\square}_{\wedge} & \forall \phi, \psi \in \mathrm{Sent}(\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}) \left( \mathsf{Tr}^{\square}(\phi {\wedge} \psi) \leftrightarrow (\mathsf{Tr}^{\square}(\phi) \wedge \mathsf{Tr}^{\square}(\psi)) \right) \\ \diamondsuit_{\mathsf{CM}} & \forall \phi \in \mathrm{Sent}(\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}) \left( \mathsf{Tr}^{\diamondsuit}(\phi) \leftrightarrow \exists \mathcal{U} \in \mathsf{Uni} \, \mathsf{Mod}(\mathcal{U},\phi) \right) \end{array}
```

Proposition 4.3. $CM^- + Non-Triviality proves:$

$$\begin{array}{ll} \mathsf{CM}_{\bot}^{\square} & \mathsf{Tr}^{\square}(\ulcorner \bot \urcorner) \leftrightarrow \bot \\ \mathsf{CM}_{\lnot}^{\square} & \forall \phi \in \mathrm{Sent}(\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}) \left(\mathsf{Tr}^{\square}(\lnot \phi) \to \lnot \mathsf{Tr}^{\square}(\phi)\right) \\ \mathsf{D}_{\mathsf{CM}} & \forall \phi \in \mathrm{Sent}(\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}) \left(\mathsf{Tr}^{\square}(\phi) \to \mathsf{Tr}^{\diamondsuit}(\phi)\right) \end{array}$$

Lemma 4.4 (Soundness Lemma). CM⁻ proves that for all $\mathcal{U} \in \mathsf{Uni}$, $\{\phi \in \mathcal{L}_{\mathsf{Uni},\mathsf{Mod}} \mid \mathsf{Mod}(\mathcal{U},\phi)\}$ is deductively closed.

Proof. Using the compositional axioms CM_{\neg} , CM_{\land} and CM_{\forall} , this is proved just like the soundness theorem for the usual semantics of first-order logic.

The theory MS (standing for Multiverse theory of Satisfaction) is analogous to the Friedman–Sheard theory of truth FS:

System 4.5 (MS). Consider these rules of proof:

```
NEC For each \phi \in \operatorname{Sent}(\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}): If \mathsf{MS} \vdash \phi, then \mathsf{MS} \vdash \mathsf{Tr}^{\square}(\lceil \phi \rceil). CONEC For each \phi \in \operatorname{Sent}(\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}): If \mathsf{MS} \vdash \mathsf{Tr}^{\square}(\lceil \phi \rceil), then \mathsf{MS} \vdash \phi.
```

```
Scope and Completeness axioms
                                                                     Given a set theory S in language \mathcal{L}^+:
                                                                                   \begin{array}{l} \forall \phi \in \mathcal{L}^+_{\mathsf{Uni},\mathsf{Mod}} \left( \phi \in S \to \forall \mathcal{U} \in \mathsf{Uni} \, \mathsf{Mod}(\mathcal{U},\phi) \right) \\ \forall \phi \in \mathcal{L}^+_{\mathsf{Uni},\mathsf{Mod}} \left( \forall \mathcal{U} \in \mathsf{Uni} \, \mathsf{Mod}(\mathcal{U},\phi) \to \mathrm{Pr}_S(\phi) \right) \end{array} 
                     Multiverse<sub>S</sub>
                      Completeness<sub>S</sub>
                                                                                                     Reflective axioms
                                          Non-Triviality
                                                                                                                             \exists \mathcal{U} \mathsf{Uni}(\mathcal{U})
                                          Multiverse Reflection \forall \sigma \in \operatorname{Sent}(\mathcal{L}) : \operatorname{Tr}^{\square}(\lceil \sigma \rceil) \to \sigma
                                          Self-Perception
                                                                                                                             Iso(self) + Uni(self)
                                                                                                            Modal axioms
                                                                                                                           \begin{split} & \operatorname{Tr}^\square(\ulcorner\sigma\to\tau\urcorner)\to \left(\operatorname{Tr}^\square(\ulcorner\sigma\urcorner)\to\operatorname{Tr}^\square(\ulcorner\tau\urcorner)\right) \\ & \operatorname{Tr}^\square(\ulcorner\sigma\urcorner)\to\operatorname{Tr}^\lozenge(\ulcorner\sigma\urcorner) \\ & \operatorname{Tr}^\square(\ulcorner\sigma\urcorner)\to\sigma \\ & \operatorname{Tr}^\square(\ulcorner\sigma\urcorner)\to\operatorname{Tr}^\square(\ulcorner\operatorname{Tr}^\square(\ulcorner\sigma\urcorner)\urcorner) \\ & \left(\operatorname{Tr}^\square(\ulcorner\operatorname{Tr}^\square(\ulcorner\sigma\urcorner)\to\operatorname{Tr}^\square(\ulcorner\sigma\urcorner)\right) \end{split}
K_{CM}
                              \forall \sigma, \tau \in \text{Sent}(\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}):
                              \forall \sigma \in \operatorname{Sent}(\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}):
D_{CM}
                              \forall \sigma \in \operatorname{Sent}(\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}):
T_{CM}
                              \forall \sigma \in \operatorname{Sent}(\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}):
4_{CM}
                              \forall \sigma \in \mathrm{Sent}(\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}):
Löb<sub>CM</sub>
                                                                                             Gödel-Löb multiverse
                                  \mathsf{Comp}_\mathsf{CM} \quad \mathsf{CM}^- + \mathsf{Multiverse}_\mathsf{Comp}_\mathsf{CM} + \mathsf{Completeness}_\mathsf{Comp}_\mathsf{CM}
```

Figure 4.1: Semantically motivated multiverse axioms

The system MS^- is $CM^- + NEC + CONEC$ and the system MS is CM + NEC + CONEC.

If \mathcal{L}' expands \mathcal{L} , then we write $MS^-(\mathcal{L}')$ and $MS(\mathcal{L}')$ for the corresponding systems obtained by replacing all occurrences of $\mathcal{L}_{Uni,Mod}$, in the axioms of the form CM_- and in the rules NEC, CONEC, by the language $\mathcal{L}'_{Uni,Mod}$.

Recall that if S is a system involving deductive rules, and A is an axiom, then S + A denotes the natural extension of S in which these deductive rules may be applied to proofs also involving A. For example, in $MS + \exists x \, Uni(x)$ we may use NEC to derive $\forall \mathcal{U} \in Uni \, Mod(\mathcal{U}, \, \exists x \, Uni(x) \,)$.

Figure 4.1 displays reflective axioms, modal axioms and the system $\mathsf{GL}_{\mathsf{CM}^-}$ interpreting Gödel-Löb logic.

The reflective axioms may be viewed as statements, of increasing strength, that the universe of the background theory is reflected in the multiverse: Non-Triviality just says that there is a universe in the multiverse; Multiverse Reflection is equivalent to that any \mathcal{L} -sentence holding in the background universe also holds in some universe; and Self-Perception goes as far as saying that the background universe is isomorphic to a universe in the multiverse.

The next theorem applies standard arguments from axiomatic theories of truth to exhibit semantically motivated axioms that turn out to be paradoxical.

Theorem 4.6. (a) The following axiom schema is inconsistent over $CM^- + NEC$:

$$\mathsf{T}_{\mathsf{CM}} \quad \forall \sigma \in \mathsf{Sent}(\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}), \; \mathsf{Tr}^{\square}(\lceil \sigma \rceil) \to \sigma$$

(b) The following axiom schema is inconsistent over CM⁻ + NEC + Non-Triviality:

$$4_{\mathsf{CM}} \quad \forall \sigma \in \mathrm{Sent}(\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}), \ \mathsf{Tr}^{\square}(\ulcorner \sigma \urcorner) \to \mathsf{Tr}^{\square}(\ulcorner \mathsf{Tr}^{\square}(\ulcorner \sigma \urcorner) \urcorner)$$

Remark. Note that T_{CM} is the untyped version of Multiverse Reflection.

Proof. By Gödel diagonalization, there is an $\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}$ -sentence λ , such that

$$\mathsf{CM} \vdash \lambda \leftrightarrow \neg \mathsf{Tr}^{\square}(\lceil \lambda \rceil).$$

By T_{CM} , $Tr^{\square}(\lceil \lambda \rceil) \to \lambda$, so we get $\neg Tr^{\square}(\lceil \lambda \rceil)$, and therefore λ . Now $Tr^{\square}(\lceil \lambda \rceil)$ follows by NEC, a contradiction.

By 4_{CM} , $Tr^{\square}(\lceil \lambda \rceil) \to Tr^{\square}(\lceil Tr^{\square}(\lceil \lambda \rceil) \rceil)$, so by $CM_{\leftrightarrow}^{\square}$, $Tr^{\square}(\lceil \lambda \rceil) \to Tr^{\square}(\lceil \neg \lambda \rceil)$, and by CM_{\neg}^{\square} (using Non-Triviality), $Tr^{\square}(\lceil \lambda \rceil) \to \neg Tr^{\square}(\lceil \lambda \rceil)$. So we get $\neg Tr^{\square}(\lceil \lambda \rceil)$ and therefore λ . Now $Tr^{\square}(\lceil \lambda \rceil)$ follows by NEC, a contradiction.

The following Proposition relates the natural systems obtained by adding reflective axioms to $\mathsf{CM}^- + \mathsf{NEC}$.

Proposition 4.7. Over $CM^- + NEC$:

- (a) Multiverse Reflection ⊢ Non-Triviality
- (b) Self-Perception ⊢ Multiverse Reflection

Proof. (a) From $\mathsf{Tr}^{\square}(\lceil \bot \rceil) \to \bot$ we get $\neg \forall \mathcal{U} \in \mathsf{Uni} \, \mathsf{Mod}(\mathcal{U}, \lceil \bot \rceil)$, whence Non-Triviality.

(b) Let $\sigma \in \mathcal{L}$, and assume $\mathsf{Tr}^{\square}(\lceil \sigma \rceil)$. Then $\mathsf{Mod}(\mathsf{self}, \lceil \sigma \rceil)$. So by Proposition 2.16, we have σ .

 $\mathsf{Comp}_\mathsf{CM}$ in Figure 4.1 includes two axioms that refer to the theory $\mathsf{Comp}_\mathsf{CM}$. These axioms can be constructed by Gödel's fixed-point lemma. The following theorem shows that $(\Box \mapsto \mathsf{Tr}^\Box)$ generates an interpretation of Gödel-Löb provability logic in $\mathsf{Comp}_\mathsf{CM}$, which in turn is interpretable in ZF .

Theorem 4.8. (a) There is an interpretation \mathcal{B} of the system K of modal propositional logic¹¹ in CM⁻ + NEC, satisfying any given assignment of the propositional variables, and

$$\mathcal{B}(\Box \sigma) = \mathsf{Tr}^{\Box}(\mathcal{B}(\ulcorner \sigma \urcorner)), \text{ for each modal propositional formula } \sigma.$$

(b) \mathcal{B} above interprets the system GL of modal predicate $logic^{12}$ in $Comp_{CM}$. In particular,

$$Comp_{CM} \vdash K_{CM} + 4_{CM} + L\ddot{o}b_{CM} + NEC.$$

- (c) ZF interprets Comp_{CM}. If ZF is closed under the Reflection rule (or if ZF is ω -consistent), then ZF interprets Comp_{CM} + CONEC.
- *Proof.* (a) The interpretation \mathcal{B} can be constructed by primitive recursion, using the technique described in Halbach (2014)[Ch. 5.3]. By NEC, \mathcal{B} validates the modal necessitation rule, and by $\mathsf{CM}^{\square}_{\to}$ (from which K_{CM} is easily derived), it validates K .

¹¹The system K is regulated by the rule of modal necessitation, $\vdash \sigma \Rightarrow \vdash \Box \sigma$, and the axiom schema K, $\Box(\sigma \to \theta) \to (\Box \sigma \to \Box \theta)$.

¹²The system GL extends K with the axiom schema 4, $\Box \sigma \rightarrow \Box \Box \sigma$, and with Löb's schema, $\Box (\Box \sigma \rightarrow \sigma) \rightarrow \Box \sigma$.

(b) By the Soundness Lemma,

$$\mathsf{Comp}_{\mathsf{CM}} \vdash \forall \sigma \in \mathsf{Sent}(\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}) \, (\mathsf{Pr}_{\mathsf{GL}_{\mathsf{CM}}}(\sigma) \leftrightarrow \mathsf{Tr}^{\square}(\sigma)). \tag{*}$$

We apply the Hilbert-Bernays-Löb provability conditions. For NEC, note that for each $\sigma \in \mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}$, $\mathsf{Comp}_\mathsf{CM} \vdash \sigma \Rightarrow \mathsf{Comp}_\mathsf{CM} \vdash \mathsf{Pr}_{\mathsf{Comp}_\mathsf{CM}}(\ulcorner \sigma \urcorner)$, and apply (*). For 4_CM , note that for each $\sigma \in \mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}$, $\mathsf{Comp}_\mathsf{CM} \vdash \mathsf{Pr}_\mathsf{Comp}_\mathsf{CM}(\ulcorner \sigma \urcorner) \to \mathsf{Pr}_\mathsf{Comp}_\mathsf{CM}(\ulcorner \mathsf{Pr}_\mathsf{Comp}(\ulcorner \sigma \urcorner))$, and apply (*) both externally and internally. By Löb's Theorem (see Lemma 13.7 in Halbach (2014)) and the preceding item, we are done.

(c) Let \mathcal{C} be the interpretation generated by:

$$\mathsf{Uni}(\mathcal{U}) \mapsto \mathcal{U} \models \mathsf{Comp}_\mathsf{CM}$$

$$\mathsf{Mod}(\mathsf{Uni}, \phi, f) \mapsto \mathcal{U} \models (\phi, f)$$

By the Tarskian conditions of satisfaction, $\mathsf{ZF} \vdash \mathcal{C}(\mathsf{CM}^-)$. By construction of \mathcal{C} , $\mathsf{ZF} \vdash \mathcal{C}(\mathsf{Multiverse}_{\mathsf{Comp}_{\mathsf{CM}}})$. By construction of \mathcal{C} and the Completeness theorem, $\mathsf{ZF} \vdash \mathcal{C}(\mathsf{Completeness}_{\mathsf{Comp}_{\mathsf{CM}}})$. From (*), it is easily seen that the Reflection rule yields CONEC. By Proposition 2.11, ω -consistency suffices.

5 Interpreting the Copernican multiverse of sets

Now we proceed to lay forth a technique for validating the theories in §4 by means of the revision-semantic construction in §3. The Main Lemma establishes that a variety of Copernican multiverse theories can by interpreted in a suitable hierarchy of theories.

First we need a lemma establishing a normal form for derivation in MS. The analogous result for the case of the Friedman–Sheard theory of truth (over arithmetic) was established by Broberg (2021). The authors are grateful to Broberg for allowing the inclusion of his proof re-worked for the system MS. We write $MS_{NC}^- + S$ for the system whose theorems are the conclusions of Hilbert-style proofs in $MS^- + S$ such that all applications of NEC are before all applications of CONEC.

Lemma 5.1. Let S be a theory. If $\mathsf{MS}^- + S \vdash \psi$, then there exists χ such that $\mathsf{CM}^- + \mathsf{NEC} + S \vdash \chi$ and $\mathsf{CM}^- + \mathsf{CONEC} + S + \chi \vdash \psi$.

Proof. For simplicity, we assume that all formulas in all theories and proofs considered are sentences; there is an adequate Hilbert-style proof system meeting this assumption. It suffices to show that $\mathsf{MS}^-_{\mathrm{NC}} + S \vdash \psi$.

Let ρ be a Hilbert-style proof, with $\psi_0, \dots, \psi_{l-1}$ as rows, of ψ_{l-1} in $\mathsf{MS}^- + S$. By induction, we may assume that $\mathsf{MS}^-_{\mathrm{NC}} + S \vdash \psi_r$, for each r < l-1. There are four cases to consider as to how the last row of ρ is obtained:

(Axiom) ψ_{l-1} is an axiom.

(First-order) ψ_{l-1} is derived by a rule of inference of first-order logic.

(CONEC) ψ_{l-1} is derived from ψ_r by CONEC, for some r < l - 1.

(NEC) ψ_{l-1} is derived from ψ_r by NEC, for some r < l - 1.

We proceed to establish $MS_{NC}^- + S \vdash \psi_{l-1}$ for each case.

- (Axiom) In this case ψ_{l-1} is also an axiom of $MS_{NC}^- + S$ (a proof of length 1).
- (First-order) In this case ψ_{l-1} is also derived by the same rule in $MS_{NC}^- + S$, utilizing the induction hypothesis.
- (CONEC) In this case ψ_{l-1} is also derived by CONEC in $MS_{NC}^- + S$, utilizing the induction hypothesis and that this application of CONEC is right at the end, after all applications of NEC.
- (NEC) This is the case requiring work. We have that ψ_{l-1} is the sentence $\mathsf{Tr}^{\square}(\lceil \psi_r \rceil)$ and that $\mathsf{MS}^{-}_{\mathrm{NC}} + S \vdash \psi_r$. Let $\theta_0, \dots, \theta_{k-1} = \psi_r$ be the rows of a Hilbert-style proof π witnessing this. We proceed to show $\mathsf{MS}^{-}_{\mathrm{NC}} + S \vdash \mathsf{Tr}^{\square}(\lceil \theta_q \rceil)$, for each $q \leq k-1$. By induction, we may assume that this holds for each q < k-1. Again there are four cases to consider as to how the last row of π is obtained:

(Axiom') θ_{k-1} is an axiom.

(First-order') θ_{k-1} is derived by a rule of inference of first-order logic.

(CONEC') θ_{k-1} is derived from θ_q by CONEC, for some q < k-1.

(NEC') θ_{k-1} is derived from θ_q by NEC, for some q < k-1.

We proceed to establish $\mathsf{MS}_{\mathrm{NC}}^- + S \vdash \mathsf{Tr}^{\square}(\lceil \theta_{k-1} \rceil)$ for each case.

- (Axiom') In this case we apply NEC to θ_{k-1} to obtain a proof of $\mathsf{Tr}^{\square}(\lceil \theta_{k-1} \rceil)$ (of length 2).
- (First-order') We have $q_0 < \cdots < q_n < k-1$, such that $\{\theta_{q_0}, \cdots, \theta_{q_n}\} \vdash \theta_{k-1}$. By the induction hypothesis, $\mathsf{MS}_{\mathrm{NC}}^- + S \vdash \mathsf{Tr}^\square(\ulcorner \theta_{q_i} \urcorner)$, for each $0 \le i \le n$. Clearly, these proofs can be merged (respecting the requirement on the order of the applications of NEC and CONEC) into one Hilbert-style proof in $\mathsf{MS}_{\mathrm{NC}}^- + S$ in which $\mathsf{Tr}^\square(\ulcorner \theta_{q_0} \urcorner), \cdots, \mathsf{Tr}^\square(\ulcorner \theta_{q_n} \urcorner)$ are derived.

Now note that by the Soundness Lemma,

$$\mathsf{CM}^- + \{\mathsf{Tr}^\square(\lceil \theta_{q_0} \rceil), \cdots, \mathsf{Tr}^\square(\lceil \theta_{q_n} \rceil)\} \vdash \mathsf{Tr}^\square(\lceil \theta_{k-1} \rceil).$$

We add a proof of that (which has no applications of NEC or CONEC) to the end of the previous proof, obtaining $\mathsf{MS}^-_{\mathsf{NC}} + S \vdash \mathsf{Tr}^\square(\lceil \theta_{k-1} \rceil)$, as desired.

(CONEC') In this case $\mathsf{Tr}^{\square}(\lceil \theta_{k-1} \rceil)$ equals θ_q , which we already have a proof of.

(NEC') In this case the last step of π is obtained by NEC, so there cannot be any application of CONEC in π . Hence, we can make an extra application of NEC at the end of π to obtain a proof of $\operatorname{Tr}^{\square}(\lceil \theta_{k-1} \rceil)$ in $\operatorname{MS}^{-}_{\operatorname{NC}} + S$.

Before embarking on proving the Main Lemma of the paper, we introduce notation for the kind of interpretations involved. The Main Lemma encapsulates the revision-semantic construction of a model of "the Copernican multiverse of sets".

Let T, Uni, Mod_0 be revision parameters. Let S be a set theory in a language L and let t be an L-term such that $S \vdash t \in \mathbb{N}$. Then $\mathcal{I}_{L,t}^{\mathrm{Uni}, Mod_0}$ denotes the interpretation of the language $L_{\mathrm{Uni}, Mod}$ into L generated by interpreting Mod as Mod_t (obtained from Construction 3.2) and Uni as Uni_t . (Although the full notation is $\mathcal{I}_{L,t}^{\mathrm{Uni}, Mod_0}$, with L, Uni , Mod fixed, for example, we denote this interpretation by \mathcal{I}_t .) Note that this interpretation fixes each formula in L.

More generally, if $S \vdash \exists^! x \ (\phi(x) \land x \in \mathbb{N})$, then \mathcal{I}_{ϕ} denotes the interpretation generated by interpreting Mod by $\forall x \in \mathbb{N} \ (\phi(x) \to \operatorname{Mod}_x)$ and Uni by $\forall x \in \mathbb{N} \ (\phi(x) \to \operatorname{Uni}_x)$. We may expand

the language with a constant symbol c_{ϕ} and extend \mathcal{S} with the axiom $\forall x \, (\phi(x) \leftrightarrow x = c_{\phi})$, to produce an interpretation $\mathcal{I}_{c_{\phi}}$ equivalent to \mathcal{I}_{ϕ} .

Let T_0 and T_1 be theories in the languages L_0 and L_1 , respectively. Let \mathfrak{F} be a family (set) of interpretations from L_0 to L_1 . We say that \mathfrak{F} is a local interpretation of T_0 in T_1 if for any finite set T'_0 of consequences of T_0 , there is an $\mathcal{I} \in \mathfrak{F}$ which interprets T'_0 in T_1 . Alternatively, we say that T_1 locally interprets T_0 by \mathfrak{F} , respectively.

Given revision parameters T, Uni, Mod₀, recall that \mathcal{L}^{T} is the language of the theories \mathbf{T}_{n} and that \mathcal{L}^{Rev} is any sublanguage of \mathcal{L}^{T} .

Main Lemma. Let T, Uni and Mod₀ be revision parameters such that $\mathbf{T}_{\omega} \vdash \mathsf{ZF}$, let $\mathfrak{F} = \{\mathcal{I}_{\underline{k}} \mid k \in \mathbb{N}\}$ and let S be an $\mathcal{L}^{\mathrm{Rev}}_{\mathsf{Uni},\mathsf{Mod}}$ -theory such that for any finite $\Gamma \subseteq S$,

$$\exists A \in \mathbb{N} \, \forall k \in \mathbb{N} \, [A \leq k \to \mathbf{T}_k \vdash \mathcal{I}_k^{\mathrm{Uni}, \mathrm{Mod}_0}(\Gamma)].$$

- (a) If T, Uni, Mod₀ admit NEC*, then T_{ω} locally interprets CM⁻ + NEC + S by \mathfrak{F} .
- (b) If T, Uni, Mod₀ admit CONEC*, then T_{ω} locally interprets CM⁻ + CONEC + S by \mathfrak{F} .
- (c) If T, Uni, Mod₀ admit NEC* and CONEC*, then T_{ω} locally interprets $MS^- + S$ by \mathfrak{F} .

Proof. We prove the latter, most complicated assertion; the other two assertions follow by restricting the proof to the appropriate cases. Assume that T, Uni, Mod_0 are revision parameters admitting NEC^* and $CONEC^*$.

Assume that $\mathsf{MS}^- + S \vdash \psi$. By Lemma 5.1, we have that $\mathsf{MS}^-_{\mathrm{NC}} + S \vdash \psi$. Let ρ be a linear Hilbert-style proof witnessing this. Let Γ be the axioms of $\mathsf{CM}^- + S$ occurring in ρ . We shall start by showing that

$$\exists A' \in \mathbb{N} \ \forall k \in \mathbb{N} \ [A' \le k \to \mathbf{T}_k \vdash \mathcal{I}_{\underline{k}}^{\mathrm{Uni}, \mathrm{Mod_0}}(\Gamma)]. \tag{*}$$

We have by the assumption of the lemma that there is $A<\omega$, such that for any $k<\omega$ with $A\leq k$, we have $\mathbf{T}_k\vdash\mathcal{I}_{\underline{k}}^{\mathsf{Uni},\mathsf{Mod_0}}(\phi)$, for every axiom ϕ of S in Γ . Moreover, note that for any $k<\omega$, and any axiom ϕ of $\mathsf{ZF}+\mathsf{Sep}(\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}})+\mathsf{Rep}(\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}})$, $\mathcal{I}_{\underline{k}}^{\mathsf{Uni},\mathsf{Mod_0}}(\phi)$ is an axiom of ZF . So since $\mathbf{T}_\omega\vdash\mathsf{ZF}$, there is $A'\geq 1$ with $A\leq A'<\omega$, such that for any $k<\omega$ with $A'\leq k$, we have $\mathbf{T}_k\vdash\mathcal{I}_{\underline{k}}^{\mathsf{Uni},\mathsf{Mod_0}}(\phi)$, for every axiom ϕ of $\mathsf{ZF}+\mathsf{Sep}(\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}})+\mathsf{Rep}(\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}})$ in Γ .

Suppose that ϕ is a compositional axiom (of the form CM_). Then $\forall A' \leq k < \omega \left[\mathbf{T}_k \vdash \mathcal{I}_{\underline{k}}^{\mathrm{Uni},\mathrm{Mod_0}}(\phi) \right]$ follows from that for all \mathcal{L}^{T} -structures \mathcal{U} and all $\mathcal{L}^{\mathrm{Rev}}_{\mathrm{Uni},\mathrm{Mod}}$ -formulas ϕ , it is provable that for all $A' \leq k < \omega$,

$$\operatorname{Mod}_{k}(\mathcal{U}, \phi) \iff \langle \mathcal{U} |_{\mathcal{L}^{\operatorname{Rev}}}, \lceil \operatorname{Uni}_{k-1} \rceil^{\mathcal{U}}, \lceil \operatorname{Mod}_{k-1} \rceil^{\mathcal{U}} \rangle \models \phi$$

(using $1 \leq A'$), and from that the corresponding compositional conditions hold for \models .

Hence, A' satisfies (*) as desired. We introduce shifted parameters T', Uni', Mod'_0 defined by $T'_k \equiv_{df} T_{k+\underline{A'}}$, $Uni'_k \equiv_{df} Uni_{k+\underline{A'}}$ and $Mod'_k \equiv_{df} Mod_{k+\underline{A'}}$. Note that Mod' also satisfies (†) in Construction 3.2. It is easily seen that T', Uni', Mod'_0 are revision parameters admitting NEC^* and $CONEC^*$, and that $T'_{\omega} = T_{\omega}$. Note that for all ϕ , it is provable that for all $k < \omega$, $\mathcal{I}_{\underline{k}}^{Uni'}, Mod'_0$ (ϕ) $\leftrightarrow \mathcal{I}_{\underline{k}+\underline{A'}}^{Unii}, Mod_0$ (ϕ).

We index the sequence of steps in the proof by numbers $0, 1, \dots, l-1$, where l is the length of ρ . For each q < l, let ψ_q be the derived formula (or axiom) at step q of ρ (so $\psi = \psi_{l-1}$), let N_q be the number of applications of NEC in the derivation of ψ_q , and let C_q be the number of applications of CONEC in the derivation of ψ_q . It suffices to show that there are natural numbers m, n, such that $\mathbf{T}'_m \vdash \mathcal{I}^{\mathrm{Uni'}, \mathrm{Mod'_0}}_{\underline{n}}(\psi_q)$, for each q < l. We do so by induction on the steps of ρ . Let r < l. Here is our induction hypothesis:

(IH) $\mathbf{T}'_{2C_q+k} \vdash \mathcal{I}_{\underline{k}}^{\mathrm{Uni'},\mathrm{Mod}'_0}(\psi_q)$, for any step q < r, and for any $k \geq N_q + 1$.

We need to show that $\mathbf{T}'_{2C_r+k} \vdash \mathcal{I}_{\underline{k}}^{\mathrm{Uni'},\mathrm{Mod'_0}}(\psi_r)$, whenever $N_r+1 \leq k$. There are four cases, as to which rule of inference (if any) is applied to obtain ψ_r from $\{\psi_q \mid q < r\}$:

(Axiom) ψ_r is an axiom in Γ .

(First-order) ψ_r is derived by a rule of inference of first-order logic.

(NEC) ψ_r is derived from $\psi_{r'}$ by NEC, for some r' < r.

(CONEC) ψ_r is derived from $\psi_{r'}$ by CONEC, for some r' < r.

Let $k \geq N_r + 1$. We proceed to show $\mathbf{T}'_{2C_r+k} \vdash \mathcal{I}^{\mathrm{Uni'},\mathrm{Mod'}_0}_{\underline{k}}(\psi_r)$ in each of the above cases. The fact that $\mathbf{T}'_a \vdash \mathbf{T}'_b$, for any $0 \leq b \leq a \in \mathbb{N}$, will be used repeatedly without mention.

(Axiom) Suppose that ψ_r is an axiom in Γ . We have $\mathbf{T}_k' \vdash \mathcal{I}_{\underline{k}}^{\mathrm{Uni}',\mathrm{Mod}_0'}(\phi_r)$, since $\mathbf{T}_k' \vdash \mathbf{T}_{k+A'}$ and $\mathbf{T}_{k+A'} \vdash \mathcal{I}_{k+A'}^{\mathrm{Uni},\mathrm{Mod}_0}(\phi_r)$.

(First-order) By (IH), $\mathbf{T}'_{2C_r+k} \vdash \mathcal{I}_{\underline{k}}^{\mathrm{Uni'},\mathrm{Mod'_0}}(\phi_q)$, for each q < r. Since $\mathcal{I}_{\underline{k}}^{\mathrm{Uni'},\mathrm{Mod'_0}}$ is an interpretation, it respects the inference rules of first-order logic. Therefore, $\mathbf{T}'_{2C_r+k} \vdash \mathcal{I}_{\underline{k}}^{\mathrm{Uni'},\mathrm{Mod'_0}}(\psi_r)$.

(NEC) In this case ψ_r is $\forall \mathcal{U} \in \mathsf{Uni}(\mathsf{Mod}(\mathcal{U}, \lceil \psi_{r'} \rceil))$. Note that $C_r = 0$ and $N_{r'} < N_r$.

$$\begin{split} \mathbf{T}_{k-1}' &\vdash \mathcal{I}_{\underline{k-1}}^{\mathrm{Uni'},\mathrm{Mod'_0}}(\psi_{r'}) & \text{(IH)} \\ \mathbf{T}_{k}' &\vdash \forall \mathcal{U} \in \mathbf{Uni'_k}(\mathcal{U} \models \ulcorner \mathcal{I}_{\underline{k-1}}^{\mathrm{Uni'},\mathrm{Mod'_0}}(\psi_{r'}) \urcorner) & \text{NEC*} \\ \mathbf{T}_{k}' &\vdash \forall \mathcal{U} \in \mathbf{Uni'_k}(\mathrm{Mod'_k}(\mathcal{U}, \ulcorner \psi_{r'} \urcorner)) & (\dagger) \text{ in Construction } 3.2 \\ \mathbf{T}_{k}' &\vdash \mathcal{I}_{\underline{k}}^{\mathrm{Uni'},\mathrm{Mod'_0}} \big(\forall \mathcal{U} \in \mathsf{Uni}\left(\mathsf{Mod}(\mathcal{U}, \ulcorner \psi_{r'} \urcorner)\right) \big) & \text{Definition of } \mathcal{I}_{\underline{k}}^{\mathrm{Uni'},\mathrm{Mod'_0}} \end{split}$$

(CONEC) In this case $\psi_{r'}$ is $\forall \mathcal{U} \in \mathsf{Uni}(\mathsf{Mod}(\mathcal{U}, \lceil \psi_r \rceil))$.

$$\begin{split} \mathbf{T}_{2C_{r'}+k+1}' &\vdash \mathcal{I}_{\underline{k+1}}^{\mathrm{Uni'},\mathrm{Mod'_0}} \big(\forall \mathcal{U} \in \mathrm{Uni} \left(\mathrm{Mod}(\mathcal{U}, \ulcorner \psi_r \urcorner) \right) \big) & \text{(IH)} \\ \mathbf{T}_{2C_{r'}+k+1}' &\vdash \forall \mathcal{U} \in \mathbf{Uni'_{k+1}} \left(\mathrm{Mod'_{k+1}}(\mathcal{U}, \ulcorner \psi_r \urcorner) \right) & \text{Definition of } \mathcal{I}_{\underline{k+1}}^{\mathrm{Uni'},\mathrm{Mod'_0}} \big(\mathbf{V}_r) \urcorner \big) \\ \mathbf{T}_{2C_{r'}+k+1}' &\vdash \forall \mathcal{U} \in \mathbf{Uni'_{k+1}} \left(\mathcal{U} \models \ulcorner \mathcal{I}_{\underline{k}}^{\mathrm{Uni'},\mathrm{Mod'_0}} (\psi_r) \urcorner \right) & \text{(†) in Construction 3.2} \\ \mathbf{T}_{2C_{r'}+k+1}' &\vdash \forall \mathcal{U} \in \mathbf{Uni'_{2C_{r'}+k+1}} \left(\mathcal{U} \models \ulcorner \mathcal{I}_{\underline{k}}^{\mathrm{Uni'},\mathrm{Mod'_0}} (\psi_r) \urcorner \right) & \mathbf{Uni'_{2C_{r'}+k+1}} \subseteq \mathbf{Uni'_{k+1}} \\ \mathbf{T}_{2C_{r'}+k+2}' &\vdash \mathcal{I}_{\underline{k}}^{\mathrm{Uni'},\mathrm{Mod'_0}} (\psi_r) & \mathrm{CONEC}^* \\ \mathbf{T}_{2C_{r}+k}' &\vdash \mathcal{I}_{\underline{k}}^{\mathrm{Uni'},\mathrm{Mod'_0}} (\psi_r) & C_{r'} < C_r \end{split}$$

This completes the proof of the lemma. We close by recording the more detailed statement that we have actually proved: If N, C are the number of applications of NEC and CONEC, respectively, in a proof of ψ in $\mathsf{MS}^-_{NC} + S$, then

$$\mathbf{T}'_{2C+N+1+A'} \vdash \mathcal{I}^{\mathrm{Uni'},\mathrm{Mod}'_0}_{\underline{N+1+A'}}(\psi).$$

Corollary 5.2. Let T, Uni and Mod_0 be revision parameters such that for some $B \in \mathbb{N}$, $T_B \vdash \mathsf{ZF}$. Let $\mathfrak{F} = \{\mathcal{I}_{\underline{k}} \mid k \in \mathbb{N}\}$ and let S be a theory (possibly with Uni, Mod in its language) such that for any finite $\Gamma \subseteq S$,

$$\exists A \in \mathbb{N} \, \forall k \in \mathbb{N} \, [A \leq k \to \mathbf{T}_k \vdash \mathcal{I}_{\underline{k}}^{\mathrm{Uni}, \mathrm{Mod}_0}(\Gamma)].$$

- (a) If T, Uni, Mod₀ admit NEC*, then T_{ω} locally interprets CM + NEC + S by \mathfrak{F} .
- (b) If T, Uni, Mod₀ admit CONEC*, then T_{ω} locally interprets CM + CONEC + S by \mathfrak{F} .
- (c) If T, Uni, Mod₀ admit NEC* and CONEC*, then T_{ω} locally interprets MS + S by \mathfrak{F} .

Proof. Let $B \in \mathbb{N}$, such that $\mathbf{T}_B \vdash \mathsf{ZF}$. Since T, Uni and Mod₀ are revision parameters, we have

$$\forall k \in \mathbb{N} [B+1 \leq k \to \mathbf{T}_k \vdash \forall \mathcal{U} \in \mathbf{Uni}_k (\mathcal{U} \models \mathsf{ZF})].$$

So by the definition of $\mathcal{I}^{\mathrm{Uni},\mathrm{Mod}_0}$ and (†) in Construction 3.2,

$$\forall k \in \mathbb{N} \ [B+1 \leq k \to \mathbf{T}_k \vdash \mathcal{I}_k^{\mathrm{Uni}, \mathrm{Mod}_0}(\mathsf{Multiverse_{ZF}})].$$

Applying the Main Lemma with $S + \mathsf{Multiverse_{ZF}}$ for S, and the maximum of A and B + 1 for A, we obtain the desired result.

The following systems, along with GR^α from Definition 2.8, are useful for measuring the consistency strength of various extensions of CM .

System 5.3. Let S be a set-theoretic system. For any set theory T in language L, R_T is the so called *(proof-theoretic) Reflection schema*:

$$\mathsf{R}_T \quad \{ \mathsf{Pr}_T(\lceil \phi \rceil) \to \phi \mid \phi \in L \}.$$

(The dot under T is sometimes omitted, when it is clear from the context.)

We recursively define, for recursive ordinals¹³ α , the theories $\mathsf{Con}^{\alpha}(S)$ and $\mathsf{R}^{\alpha}(S)$, of α -iterated Consistency over S and α -iterated Reflection schema over S, respectively:

$$\begin{split} &\operatorname{\mathsf{Con}}^0(S) =_{\operatorname{df}} S \\ &\operatorname{\mathsf{Con}}^{\alpha+1}(S) =_{\operatorname{df}} \operatorname{\mathsf{Con}}^{\alpha}(S) + \operatorname{\mathsf{Con}}_{\operatorname{\mathsf{Con}}^{\alpha}(S)} \\ &\operatorname{\mathsf{Con}}^{\alpha}(S) =_{\operatorname{df}} \bigcup_{\xi < \alpha} \operatorname{\mathsf{Con}}^{\xi}(S), \text{ for } \alpha \text{ a limit ordinal;} \\ &\operatorname{\mathsf{R}}^0(S) =_{\operatorname{df}} S \\ &\operatorname{\mathsf{R}}^{\alpha+1}(S) =_{\operatorname{df}} \operatorname{\mathsf{R}}^{\alpha}(S) + \operatorname{\mathsf{R}}_{\operatorname{\mathsf{R}}^{\alpha}(S)} \\ &\operatorname{\mathsf{R}}^{\alpha}(S) =_{\operatorname{df}} \bigcup_{\xi < \alpha} \operatorname{\mathsf{R}}^{\xi}(S), \text{ for } \alpha \text{ a limit ordinal.} \end{split}$$

We use the notations Con^{α} and R^{α} for $Con^{\alpha}(ZF)$ and $R^{\alpha}(ZF)$, respectively.

Recall System 2.8, where GR^{α} is defined, using the axiom GR_T of Global Reflection over a set theory T extending $\mathsf{CT} \upharpoonright$ (in some language L_{Sat} with a satisfaction predicate):

$$\forall \phi \in L_{\mathsf{Sat}} \left(\Pr_T(\phi) \to \mathsf{Tr}(\phi) \right)$$

Comparing R_T with GR_T , note that R_T is a schema, with a separate axiom for each formula of the form ϕ in the meta-language, while GR_T quantifies internally over all formulas in the object-language; the latter is made possible by the satisfaction/truth-predicate.

¹³An ordinal α is recursive if there is a Σ_1^0 -formula defining a well-ordering of a subset of $\mathbb N$ of order-type α . These are precisely the ordinals below ω_1^{CK} .

Remark. Let us pause to measure the consistency strengths of GR^ω , R^ω and Con^ω : The consistency strength of GR^ω is bounded by that of $\mathsf{MK} + \mathsf{GC}$ (Morse-Kelley class theory with Global Choice), which, in turn, is far less than that of $\mathsf{ZFC} +$ "there exists an inaccessible cardinal". The consistency strength of R^ω is bounded by that of GR^1 . The consistency strength of Con^ω is bounded by that of R^1 . Moreover, it is easily observed that for each $n \in \mathbb{N}$: GR^{n+1} , R^{n+1} and Con^{n+1} proves the consistency of GR^n , R^n and Con^n , respectively.

Theorem 5.4. Con $^{\omega}$ locally interprets CM + NEC + Non-Triviality.

Proof. We can choose revision parameters T, Uni, Mod₀, such that for each $n \in \mathbb{N}$:

$$\mathbf{T}_n = \mathsf{Con}^n$$
 $\mathbf{Uni}_{n+1} = \{\mathcal{U} \mid \mathcal{U} \models \mathbf{T}_n\}$

Clearly, these are revision parameters provably satisfying Soundness*, and thereby admitting NEC*. Moreover, we have for each $k \in \mathbb{N}$:

$$\begin{aligned} \mathbf{T}_{k+1} &\vdash \mathbf{Uni}_{\underline{k+1}} \neq \varnothing & \text{Definition of } \mathbf{T}_{k+1}, \mathbf{Uni}_{k+1} \\ \mathbf{T}_{k+1} &\vdash \mathcal{I}_{\underline{k+1}}^{\mathrm{Uni}, \mathrm{Mod}_0}(\mathsf{Non-Triviality}) & \text{Definition of } \mathcal{I}_{\underline{k+1}}^{\mathrm{Uni}, \mathrm{Mod}_0} \end{aligned}$$

So the result follows from Corollary 5.2, by setting $S = \{\text{Non-Triviality}\}.$

Remark. Under the mild meta-theoretic assumption that each Con^n is closed under the Reflection rule, it follows from Lemma 3.3 that the revision parameters in the above proof admit CONEC^* , yielding that Con^ω locally interprets $\mathsf{CM} + \mathsf{NEC} + \mathsf{Non-Triviality}$. This meta-theoretic assumption follows from the assumption that Con^ω is ω -consistent, which in turn follows from the existence of an ω -standard model of ZF .

Remark. Using the fine-grained result obtained at the end of the proof of the Main Lemma, we can show by an overspill-argument that $\mathsf{Con}^\omega + \{\underline{n} < c \mid n \in \mathbb{N}\}$ (for a fresh constant c) interprets $\mathsf{CM} + \mathsf{NEC} + \mathsf{Non-Triviality}$ (not just locally). This raises:

Question. Is CM + NEC + Non-Triviality ω -inconsistent?

Theorem 5.5. R^{ω} locally interprets MS + Multiverse Reflection.

Proof. We can choose revision parameters T, Uni, Mod₀, such that for each $n \in \mathbb{N}$:

$$\mathbf{T}_n = \mathsf{R}^n$$
 $\mathbf{Uni}_{n+1} = \{\mathcal{U} \mid \mathcal{U} \models \mathbf{T}_n\}$

¹⁴The following argument indicates that the consistency strength of GR^ω is far less than that of $\mathsf{MK}+\mathsf{GC}$. We rely on Fujimoto (2012): Fujimoto shows in his Theorem 70 that the consistency strength of his theory of truth, FS, is equal that of NBG_ω , which is a subtheory of $\mathsf{NBG}_{< E_0}$ introduced in Jäger & Krähenbühl (2010). By Theorem 15 in (ibid.), and by Fujimoto's Proposition 4, $\mathsf{NBG}_{< E_0}$ is a subsystem of $\mathsf{MK}+\mathsf{GC}$. (All of the relevant definitions are found in Fujimoto's §3.1.) Moreover, the proof of Fujimoto's Proposition 21 provides the base step, and NEC provides the induction step, to show that his FS proves the version of GR^ω for truth (even with the Replacement schema extended to the language with the truth predicate). Since that version of GR^ω interprets our GR^ω , the consistency strength of our GR^ω is bounded by the consistency strength of NBG_ω (and since our GR^ω does not have the Replacement schema extended to the language with the satisfaction relation, this bound is probably not tight), which in turn is bounded by the consistency strength of $\mathsf{MK}+\mathsf{GC}$.

¹⁵If κ is an inaccessible cardinal, then $V_{\kappa+1}$ provides a natural model of MK + GC.

¹⁶This is shown by a routine argument, using Global Reflection to prove each iteration of the Reflection schema.

¹⁷This is shown by a routine argument, using the Reflection schema to prove each iteration of Consistency.

It is easily seen that these are revision parameters provably satisfying Soundness*, and thereby admitting NEC*. Similarly, it is easily seen that they admit the Reflection rule* and satisfy Completeness*, so that they admit CONEC*. Moreover, we have for each $k \in \mathbb{N}$ and each $\phi \in \mathcal{L}$:

$$\mathbf{T}_{k+1} \vdash \left(\forall \mathcal{U} \in \mathbf{Uni}_{\underline{k+1}} \, \mathrm{Mod}_{\underline{k+1}} (\mathcal{U}, \lceil \phi \rceil) \right) \to \phi \qquad \text{The completeness theorem and}$$

$$\text{the definition of } \mathbf{T}_{k+1}, \mathbf{Uni}_{k+1}$$

$$\mathbf{T}_{k+1} \vdash \mathcal{I}_{\underline{k+1}}^{\mathrm{Uni}, \mathrm{Mod}_0} (\mathsf{Tr}^{\square}(\lceil \phi \rceil) \to \phi) \qquad \text{Definition of } \mathcal{I}_{\underline{k+1}}^{\mathrm{Uni}, \mathrm{Mod}_0}$$

So the result follows from Corollary 5.2, by setting S = Multiverse Reflection.

Remark. The technique for obtaining full (not just local) interpretability, mentioned in the second remark following Theorem 5.4, does not work for the above theorem, because Multiverse Reflection is not finitely axiomatizable (as far as the authors can see). The overspill-argument must be carried out on a single formula, not a schema.

Theorem 5.6. The following conservativity results hold:

- (a) $\mathsf{Comp}_{\mathsf{CM}} \equiv_{\mathcal{L}} \mathsf{ZF}$
- (b) $CM + NEC + Non-Triviality \equiv_{\mathcal{L}} Con^{\omega}$
- (c) MS + Multiverse Reflection $\equiv_{\mathcal{L}}$ CM + NEC + Multiverse Reflection $\equiv_{\mathcal{L}}$ R $^{\omega}$

Proof. (a) This is immediate from Theorem 4.8(c) and that the interpretation used in its proof restricts to the identity on \mathcal{L} .

(b) The right-to-left direction follows from Theorem 5.4, observing that the interpretation is the identity on \mathcal{L} .

For the left-to-right direction, suppose as induction hypothesis that we have proved Con^n in $\mathsf{CM} + \mathsf{NEC} + \mathsf{Non-Triviality}$. By NEC , we have $\mathsf{Tr}^{\square}(\mathsf{Con}^{\underline{n}})$. So by $\mathsf{Non-Triviality}$ and Lemma 4.4, we can prove $\mathsf{Con}^n + \mathsf{Con}(\mathsf{Con}^{\underline{n}})$, which is Con^{n+1} , as desired.

(c) That R^{ω} proves every \mathcal{L} -theorem of MS + Multiverse Reflection follows from Theorem 5.5, observing that the interpretation is the identity on \mathcal{L} .

That MS + Multiverse Reflection $\vdash CM + NEC + Multiverse$ Reflection is trivial.

For $\mathsf{CM} + \mathsf{NEC} + \mathsf{Multiverse}$ Reflection $\vdash \mathsf{R}^\omega$, we shall show that for each $n \in \mathbb{N}$, $\mathsf{CM} + \mathsf{NEC} + \mathsf{Multiverse}$ Reflection $\vdash \mathsf{Tr}^\square(\Breve{\mathbb{R}}^n)$. Then the result follows from Multiverse Reflection. We proceed by induction. By Multiverseze, we have the base case: $\mathsf{MS} + \mathsf{Multiverse}$ Reflection $\vdash \mathsf{Tr}^\square(\Breve{\mathbb{R}}^0)$. So suppose as induction hypothesis that

$$MS + Multiverse Reflection \vdash Tr^{\square}(\underline{R}^{\underline{k}}).$$

Let $\sigma \in \text{Sent}(\mathcal{L})$. By NEC we have

$$MS + Multiverse Reflection \vdash Tr^{\square}(\lceil Tr^{\square}(R^{\underline{k}})\rceil)$$
 (1)

$$MS + Multiverse Reflection \vdash Tr^{\square}(CM)$$
 (2)

$$MS + Multiverse Reflection \vdash Tr^{\square}(\lceil Tr^{\square}(\lceil \sigma \rceil) \to \sigma^{\rceil}), \tag{3}$$

since (by Multiverse_{ZF}) NEC only needs to be applied to finitely many axioms of CM. We work in MS + Multiverse Reflection. Let $\mathcal{U} \in \mathsf{Uni}$. Assume $\mathsf{Mod}(\mathcal{U}, \lceil \Pr_{R^{\underline{k}}}(\lceil \sigma \rceil) \rceil)$. We shall show $\mathsf{Mod}(\mathcal{U}, \lceil \sigma \rceil)$. By (2), we can apply the Soundness Lemma in \mathcal{U} to obtain $\mathsf{Mod}(\mathcal{U}, \lceil \mathsf{Tr}^{\square}(\lceil \sigma \rceil) \rceil)$ from (1) and $\mathsf{Mod}(\mathcal{U}, \lceil \Pr_{R^{\underline{k}}}(\lceil \sigma \rceil) \rceil)$. Now by CM_{\to} and (3), we obtain $\mathsf{Mod}(\mathcal{U}, \lceil \sigma \rceil)$, as desired.

Theorem 5.7. GR^{ω} interprets $\mathsf{CM}(\mathcal{L}_{\iota,\mathsf{self}}) + \mathsf{NEC} + \mathsf{Self-Perception}$.

Proof. Set $\mathcal{L}^{T} = \mathcal{L}_{\mathsf{Sat},\iota,\mathsf{self}}$ and set \mathcal{L}^{Rev} to $\mathcal{L}_{\iota,\mathsf{self}}$. We can choose revision parameters T, Uni, Mod₀, such that for each $n \in \mathbb{N}$:

$$\mathbf{T}_n = \mathsf{SP}^n$$
 $\mathbf{Uni}_{n+1} = \{\mathcal{U} \mid \mathcal{U} \models \mathbf{T}_n \land \mathcal{U} \in \mathbf{crsm}\}$

Clearly these are revision parameters satisfying Soundness*, thus admitting NEC*. Moreover, note that for each $k < \omega$, $\mathbf{T}_k \vdash \mathrm{Uni}_k(\mathsf{self})$. So for each $k < \omega$,

$$\mathbf{T}_k dash \mathcal{I}_{\underline{k}}^{\mathrm{Uni},\mathrm{Mod}_0}(\mathsf{Self ext{-}Perception}).$$

Thus, it follows from Corollary 5.2 that $\mathfrak{F} = \{\mathcal{I}_{\underline{k}} \mid k < \omega\}$ is a local interpretation of $\mathsf{CM}(\mathcal{L}_{\iota,\mathsf{self}}) + \mathsf{NEC} + \mathsf{Self}\text{-}\mathsf{Perception} \text{ in } \mathsf{SP}^{\omega}.$

Recall from Lemma 2.19 that there is an interpretation $\mathcal J$ of SP^ω in GR^ω . For each $n<\omega$, let $\mathcal K_{\underline n}=\mathcal J\circ\mathcal I_{\underline n}$. Then $\mathfrak G=\{\mathcal K_{\underline k}\mid k<\omega\}$ is a local interpretation of $\mathsf{CM}(\mathcal L_{\iota,\mathsf{self}})+\mathsf{NEC}+\mathsf{Self}$ -Perception in GR^ω . There is a technical hurdle later on in this proof caused by the fact that the image of each $\mathcal K_{\underline n}$ is not included in $\mathcal L$, but includes the instance of Tr needed to define the set $\mathsf{Tr}=\{\sigma\in\mathsf{Sent}(\mathcal L_{\mathsf{Sat}})\mid \mathsf{Tr}(\sigma)\}$ for the construction of $\mathcal J$ in the proof of Lemma 2.19. To overcome this, we construct the functions $\mathcal K'_{\underline n}(\phi)$, for each $n<\omega$, replacing each occurrence of the form " $\mathsf{Tr}(t)$ " in the values of $\mathcal K_{\underline n}$ by " $t\in y$ ", where y is assumed to be fresh. Then we have for each ϕ that

$$\mathsf{GR}^{\omega} \vdash \mathcal{K}(\phi) \leftrightarrow (\mathcal{K}'(\phi))[\mathbf{Tr}/y].$$

Let f be a function enumerating the theorems of $\mathsf{CM}(\mathcal{L}_{\iota,\mathsf{self}}) + \mathsf{NEC} + \mathsf{Self}\text{-Perception}$, such that the length of the theorems is non-decreasing. Then, for each $i < \omega$ there is σ , such that $\mathsf{GR}^{\omega} \vdash f(\underline{i}) = \ulcorner \sigma \urcorner$. Let $\theta(x,z)$ be the formula expressing

$$x<\omega\wedge\exists j<\omega\,\forall i\leq x\,\mathsf{Sat}\big(\dot{\mathcal{K}}_{\underline{j}}'(f(i)),\ulcorner y\urcorner\mapsto z\big).$$

Since the values of the $\mathcal{K}'_{\underline{n}}$ are in \mathcal{L} , we have by Proposition 2.6 that for each $j < \omega$ and each $\phi \in \mathcal{L}^{\mathrm{Rev}}_{\mathsf{Uni},\mathsf{Mod}}$:

$$\mathsf{GR}^\omega \vdash \mathsf{Sat}(\Breve{\mathcal{K}}_{\underline{j}}'(\ulcorner \phi \urcorner), \ulcorner y \urcorner \mapsto z) \leftrightarrow \Breve{\mathcal{K}}_{\underline{j}}'(\phi)[z/y].$$

So since \mathfrak{G} is a local interpretation, we have for each $n < \omega$ that

$$\mathsf{GR}^{\omega} \vdash \theta(n, \mathbf{Tr}).$$

Since GR^{ω} is ω -inconsistent, there is a formula $\mathfrak{F}(x)$ such that $\mathsf{GR}^{\omega} \vdash \exists x < \omega \, \mathfrak{F}(x)$, but for each $n \in \mathbb{N}$, $\mathsf{GR}^{\omega} \vdash \neg \mathfrak{F}(\underline{n})$. Working in GR^{ω} , employing Proposition 2.7, we obtain a maximal number $d < \omega$ such that $\theta(d, \mathbf{Tr}) \land \neg \mathfrak{F}(d)$. Now there is a minimal number $e < \omega$ such that

$$\forall i \leq d \operatorname{Sat} \big(\underline{\mathcal{K}}_{\underline{e}}'(f(i)), \lceil y \rceil \mapsto \mathbf{Tr} \big).$$

So GR^{ω} defines e by a formula ψ . It follows from Proposition 2.6 that \mathcal{K}_{ψ} is an interpretation of $\mathsf{CM}(\mathcal{L}_{\iota,\mathsf{self}}) + \mathsf{NEC} + \mathsf{Self}\text{-Perception}$ in GR^{ω} .

Remark. Under the meta-theoretic assumption that the revision parameters used in the above proof admit the Reflection-rule*, GR^ω interprets $\mathsf{MS}(\mathcal{L}_{\iota,\mathsf{self}}) + \mathsf{Self}\text{-Perception}$. Another potential approach to validating CONEC would be to employ revision parameters based on a hierarchy of theories converging to $\mathsf{FS}^{\upharpoonright} + \mathsf{Sep}(\mathcal{L}_{\mathsf{Sat}})$, rather than to GR^ω (but the authors are not certain that this approach would work).

Axioms of Arithmetic Absoluteness

```
\begin{array}{ll} \Sigma_1^{\mathsf{PA}}\text{-}\mathsf{Absoluteness} & \forall \sigma \in \mathrm{Sent}(\Sigma_1^{\mathsf{PA}}): \, \forall \mathcal{U} \in \mathsf{Uni} \left(\mathsf{Mod}(\mathcal{U}, \ulcorner \sigma^{\mathbb{N} \urcorner}) \leftrightarrow \sigma^{\mathbb{N}}\right) \\ \mathsf{Arithmetic} \,\, \mathsf{Absoluteness} & \forall \sigma \in \mathrm{Sent}(\mathcal{L}^{\mathsf{PA}}): \, \forall \mathcal{U} \in \mathsf{Uni} \left(\mathsf{Mod}(\mathcal{U}, \ulcorner \sigma^{\mathbb{N} \urcorner}) \leftrightarrow \sigma^{\mathbb{N}}\right) \\ \mathsf{Arithmetic} \,\, \mathsf{Compositionality} & \forall \mathcal{U} \in \mathsf{Uni} \, \forall \phi \in \mathcal{L}_{\mathsf{Uni},\mathsf{Mod}} \left(\mathsf{Mod}(\mathcal{U}, \ulcorner \forall x \in \mathbb{N} \urcorner \phi) \leftrightarrow \forall n \in \mathbb{N} \, \mathsf{Mod}(\mathcal{U}, \phi, \ulcorner x \urcorner \mapsto \underline{n}^{\mathcal{U}})\right) \end{array}
```

Figure 6.1: Axioms of Arithmetic Absoluteness

Question. Does FS\[\times + Sep(\mathcal{L}_{Sat})\] interpret MS(\mathcal{L}_{\ell,self}) + Self-Perception?

Remark. There is another proof of the above theorem that utilizes the fine-grained result at the end of the proof of the Main Lemma. That technique only works when the theory S of the Main Lemma is finitely axiomatized (as here where $S = \{Self-Perception\}$). But the technique used in the above proof works also for non-finitely axiomatized S.

Remark. Note the contrast that CM includes $ZF + Sep(\mathcal{L}_{Uni,Mod}) + Rep(\mathcal{L}_{Uni,Mod})$ while GR^{ω} only includes $ZF + Sep(\mathcal{L}_{Sat})$. The essential reason why GR^{ω} interprets $Rep(\mathcal{L}_{Uni,Mod})$ is that the interpretations $\mathcal{I}_{\underline{n}}$ in the above proof map Mod and Uni to the \mathcal{L} -formulas $Mod_{\underline{n}}$ and $Uni_{\underline{n}}$, respectively.

In light of Theorems 5.6 and 5.7, the authors ask:

Question. Is it the case that $GR^{\omega} \equiv_{\mathcal{L}} CM(\mathcal{L}_{\iota,self}) + NEC + Self-Perception? If not, what is the precise consistency strength of <math>CM(\mathcal{L}_{\iota,self}) + NEC + Self-Perception?$

The fact that the extended schema $Sep(\mathcal{L}_{Sat})$ in GR^{ω} was only used to obtain the set of true sentences, suggests a negative answer; GR^{ω} may have strictly higher consistency strength than $CM(\mathcal{L}_{\iota,self}) + NEC + Self-Perception$.

6 Case studies

This section examines how the framework introduced above can be applied to two rather different conceptions of the set-theoretic multiverse.

6.1 Multiverse conceptions of arithmetical absoluteness

One may feel confident in adopting a universe view on arithmetic, appealing to the general acceptance of an intended model consisting of the finite ordinals, while having a multiverse view of set theory, where agreement on an intended model is lacking. This subsection therefore explores how the techniques of this paper may be applied to a conception of the multiverse where arithmetic is more or less fixed throughout the universes.

Let $\mathcal{L}^{\mathsf{PA}}$ be the language of arithmetic, and let Σ^{PA}_n be the usual complexity hierarchy of arithmetic formulas over PA. Given $\phi \in \mathcal{L}^{\mathsf{PA}}$, $\phi^{\mathbb{N}}$ denotes the corresponding \mathcal{L} -formula obtained by restricting the quantifiers to \mathbb{N} . Figure 6.1 exhibits axioms, of increasing strength, expressing the absoluteness of arithmetic. The strongest of these is Arithmetic Absoluteness, which expresses that the bounded quantifier " $\forall x \in \mathbb{N}$ " commutes with the Mod-relation, for the untyped language $\mathcal{L}_{\mathsf{Uni},\mathsf{Mod}}$.

Proposition 6.1. $CM^- + Arithmetic Compositionality \vdash Arithmetic Absoluteness$

Proof. Let $\sigma \in \operatorname{Sent}(\mathcal{L}^{\mathsf{PA}})$, such that $\sigma^{\mathbb{N}}$ holds. Let $\mathcal{U} \in \mathsf{Uni}$. We show by induction on the syntactic structure that $\mathsf{Mod}(\mathcal{U}, \ulcorner \sigma^{\mathbb{N} \urcorner}) \leftrightarrow \sigma^{\mathbb{N}}$. For atomic sentences it follows from that arithmetic equations are decidable. For the propositional connectives, the induction step follows from the axioms of the form CM_{-} ; let us look at $\sigma^{\mathbb{N}} \equiv \neg \phi$ for example:

$$\mathsf{Mod}(\mathcal{U}, \lceil \neg \phi \rceil) \iff \neg \mathsf{Mod}(\mathcal{U}, \lceil \phi \rceil) \iff \neg \phi$$

The first equivalence holds by CM_{\neg} and the second by the induction hypothesis. For the quantifier case, suppose that $\sigma^{\mathbb{N}} \equiv \forall x \in \mathbb{N} \ \phi(x)$. We calculate:

$$\mathsf{Mod}(\mathcal{U}, \lceil \forall x \in \mathbb{N} \, \phi(x) \rceil) \iff \forall n \in \mathbb{N} \, \mathsf{Mod}(\mathcal{U}, \lceil \phi(n) \rceil) \iff \forall n \in \mathbb{N} \, \phi(n) \iff \sigma^{\mathbb{N}}$$

The first equivalence holds by Arithmetic Compositionality, the second by the induction hypothesis, and the third by the fact that for all $n \in \mathbb{N}$, $\mathbb{N} \models n = \underline{n}$.

The following proposition shows the reflective power of Arithmetic Absoluteness, and exhibits a scenario where CONEC is useful.

Proposition 6.2. CM + Σ_1^{PA} -Absoluteness + CONEC $\vdash R^{\omega_1^{CK}}$

Proof. By CONEC it suffices to prove $\operatorname{Tr}^{\square}(\dot{\mathbb{R}}^{\omega_1^{\operatorname{CK}}})$. Naturally, we do so by transfinite induction. The base case follows from Multiversez, an axiom of CM. For the successor case, assume that $\operatorname{Tr}^{\square}(\dot{\mathbb{R}}^{\alpha})$ for some $\alpha < \omega_1^{\operatorname{CK}}$. Suppose $\mathcal{U} \in \operatorname{Uni}$, let $\sigma \in \mathcal{L}$ and assume that $\operatorname{\mathsf{Mod}}(\mathcal{U}, \lceil \operatorname{Pr}_{\dot{\mathbb{R}}^{\alpha}}(\sigma) \rceil)$. Since α is recursive, $\operatorname{Pr}_{\dot{\mathbb{R}}^{\alpha}}$ is Σ_1^0 , so by $\Sigma_1^{\operatorname{PA}}$ -Absoluteness, we have $\operatorname{Pr}_{\dot{\mathbb{R}}^{\alpha}}(\sigma)$. It now follows from $\operatorname{Tr}^{\square}(\dot{\mathbb{R}}^{\alpha})$ and the Soundness Lemma that $\operatorname{\mathsf{Mod}}(\mathcal{U}, \lceil \sigma \rceil)$. So by $\operatorname{\mathsf{CM}}_{\to}$, we have $\operatorname{\mathsf{Mod}}(\mathcal{U}, \dot{\mathbb{R}}^{\alpha+1})$, as desired. The limit case is immediate from the definition of $\operatorname{\mathsf{R}}^{\alpha}$.

Remark. Note that the above proof argues model-theoretically on an arbitrary universe. But even though $CM + \Sigma_1^{PA}$ -Absoluteness + CONEC proves a fair amount of reflection, it is not clear to the authors whether it proves that there is a universe (Non-Triviality).

This section raises questions about the consistency (strength) of combinations of Copernican multiverse theories and axioms of arithmetic absoluteness, in particular:

Question. Is $CM + \Sigma_1^{PA}$ -Absoluteness + CONEC consistent relative to $R^{\omega_1^{CK}}$?

Question. Is $CM^{-}(\mathcal{L}_{\iota.self}) + NEC + Self-Perception + Arithmetic Compositionality consistent?$

The following Proposition may be viewed as a partial answer to the second question, but the authors do not consider it to suggest an ultimately negative answer:

Proposition 6.3. The system $CM^- + NEC + Arithmetic Compositionality + Non-Triviality is <math>\omega$ -inconsistent.

Proof. This is a corollary of McGee's paradox, see McGee (1985).

6.2 The Hamkins multiverse

Hamkins (2012) introduced a conception of the set-theoretic multiverse, which informally is based on four over-arching principles:

- 1. The multiverse is a non-empty collection of models of ZFC.
- 2. The multiverse is closed under the usual techniques for constructing models of set theory from other models of set theory, such as forcing extensions and inner models.

- 3. Every universe is countable and ω -non-standard from the perspective of another universe.
- 4. The multiverse is closed under iterating large cardinal embeddings backward.

Definition 1.1 in (Gitman & Hamkins, 2010, pp. 475–6) gives succinct formulations of axioms encapsulating this conception. The third (and possibly the fourth) principle is more controversial than the first two. For example, the Well-foundedness Mirage axiom states that for every universe \mathcal{U} there is a universe \mathcal{V} which thinks \mathcal{U} is ω -non-standard.

Philsophical argmuents in support for the axioms are provided in (Hamkins, 2012, §9); and a model of them is constructed by Gitman & Hamkins (2010), essentially taking the multiverse to consist of the countable recursively saturated models of ZFC. We call this the *Gitman-Hamkins* model of the multiverse.

Gitman & Hamkins (2010) consider a weak and strong form of Well-foundedness Mirage. In the terminology of this paper, these are formally stated as follows:

```
\begin{array}{ll} \mathsf{WM}_{\mathsf{weak}} & \forall \mathcal{U} \in \mathsf{Uni} \ \exists \mathcal{V} \in \mathsf{Uni} \ \exists u \in \mathcal{V} \ \big( u_{\mathcal{V}} = \mathcal{U} \land \\ & \land \mathcal{V} \models \text{``} u \ \mathrm{is} \ \omega\text{-non-standard''} \big) \\ \\ \mathsf{WM}_{\mathsf{strong}} & \forall \mathcal{U} \in \mathsf{Uni} \ \exists \mathcal{V} \in \mathsf{Uni} \ \exists u \in \mathcal{V} \ \big( u_{\mathcal{V}} = \mathcal{U} \land \\ & \land \mathcal{V} \models \text{``} u \ \mathrm{is} \ \mathrm{an} \ \omega\text{-non-standard model of ZFC''} \big) \end{array}
```

The distinction between the axioms is discussed in (Gitman & Hamkins, 2010, pp. 479-480), where a reflection assumption is introduced to ensure that the stronger axiom gets validated in the model. Their multiverse conception is flat in the sense that it does not consider the universes as themselves being models of the multiverse axioms. However, the move from WM_{weak} to WM_{strong} may naturally be viewed as a step in that direction. Accordingly, we may reformulate Well-founded Mirage in $\mathcal{L}_{Uni,Mod}$ as:

```
WM \forall \mathcal{U} \in \mathsf{Uni} \,\exists \mathcal{V} \in \mathsf{Uni} \,\exists u \in \mathcal{V} \, (u_{\mathcal{V}} = \mathcal{U} \land \land \mathsf{Mod}(\mathcal{V}, "u \text{ is an } \omega\text{-non-standard model in } \mathbf{Uni}"))
```

Informally speaking this says not only "for every universe \mathcal{U} there is a universe \mathcal{V} that thinks that \mathcal{U} is ω -non-standard", but also " \mathcal{V} thinks that \mathcal{U} is a universe". Note that over CM + Choice + NEC,¹⁸ we get WM_{strong} from WM, and also iterated forms of WM, starting with $\forall \mathcal{W} \in$ Uni Mod(\mathcal{W} , WM). The authors take this to be a natural way for Well-founded Mirage to manifest in the multiverse of sets. The analogous modification can also be made to the Countability axiom in Definition 1.1 of Gitman & Hamkins (2010).

We write HM (the Hamkins Multiverse) for the $\mathcal{L}_{Uni,Mod}$ -theory obtained by extending CM with the axioms of Definition 1.1 in Gitman & Hamkins (2010) reformulated so that \models is replaced by Mod and the Well-founded Mirage and Countability axioms are modified as above. In general, it is natural to add the closure condition NEC to this system, since that gives us a Copernican conception which ensures that the backgound universe does not have a privileged point of outlook over the multiverse. In particular, this yields a natural strengthening of Well-founded Mirage (and Countability), as explained in the previous paragraph. Moreover, the techniques used to validate Self-Perception in this paper are closely related to the validation of Well-founded Mirage by Gitman & Hamkins (2010). Therefore, the authors champion $CM(\mathcal{L}_{\iota,self}) + HM + Self-Perception$ as a theory of the multiverse.

Note that the choice of \mathbf{Uni}_n in the proof of Theorem 5.7 is the collection of countable recursively saturated models, just as in the Gitman-Hamkins model of the multiverse. A plausible

¹⁸Choice is added here because Hamkins's conception of the multiverse is formulated for ZFC.

approach to proving the consistency of the above theory is therefore to use the construction from the proof of Theorem 5.7, setting up the revision parameters T, Uni, Mod₀ so that for each $n \in \mathbb{N}$:

$$\mathbf{T}_n = \mathsf{SP}^n(\mathsf{ZFC})$$
 $\mathbf{Uni}_{n+1} = \{\mathcal{U} \mid \mathcal{U} \models \mathbf{T}_n \land \mathcal{U} \mid_{\mathcal{L}} \in \mathbf{crsm}\}$

Conjecture. $\mathsf{GR}^{\omega}(\mathsf{ZFC})$ interprets $\mathsf{CM}(\mathcal{L}_{\iota.\mathsf{self}}) + \mathsf{NEC} + \mathsf{HM} + \mathsf{Self}\text{-Perception}$.

Addressing this conjecture falls outside the scope of this paper. What needs to be done is essentially to verify that the proof of the Main Theorem by Gitman & Hamkins (2010) generalizes from models of ZFC to models of $SP^n(ZFC)$.

7 Conclusion

We have developed a framework of satisfaction for the multiverse of set theory, with two sides: A revision-semantic construction of an increasingly adequate definition of truth-in-a-universe, and a family of axiomatic theories validated by the revision construction. We have shown how the construction can be adjusted, by tuning the revision parameters, in order to validate various multiverse axioms.

The basic theory of satisfaction for the multiverse is CM, which extends ZF with axioms expressing that truth-in-a-universe is compositional with respect to the logical connectives and quantifiers. Adding the deductive rule NEC yields a system with the closure condition that whatever is provable in the multiverse theory also provably holds in each universe. So such a system respects the Copernican Principle that the background universe should not have a privileged point of outlook over the multiverse. Adding also the dual principle of CONEC, yields the system MS. MS is in a sense analogous to the Friedman–Sheard theory of truth (FS), but in Theorem 4.8 we saw that, unlike FS, it is conservative over the base theory, under the meta-theoretic assumption that ZF is closed under the Reflection rule (which follows if ZF is ω -consistent, and in particular if there is an ω -standard model of ZF).

The choice of ZF as base theory for our framework is important for the axiom of Self-Perception. The proof of theorem 5.7 yields a model of $CM(\mathcal{L}_{Sat}) + NEC + Self-Perception$ satisfying that each universe is a countable recursively saturated model of ZF (under Mod). As far as the authors can see, both the full Separation and Replacement schemas of ZF, as well as its Foundation axiom, are needed both for the background theory and for the internal theory of each universe, due to the application of Theorem 2.4 in the proofs of Lemma 2.14 and Lemma 2.18. In contrast, for the weaker extensions of CM introduced in this paper, the authors do not see any need for full ZF. For example, the authors believe that most of the results of this paper hold (with minor modifications) also when taking Mac Lane set theory¹⁹, or Kripke–Platek set theory with Infinity²⁰, as base theory and as theory for the internal universes. Due to the close connection between Mac Lane set theory and topoi, this suggests that the framework can be adapted to give an analogous multiverse framework for topos theory.

We explored adding axioms of a reflective character, asserting that the universe of the background multiverse theory is reflected in the multiverse. Whereas Non-Triviality merely states the existence of a universe in the multiverse, Multiverse Reflection can be viewed as expressing that for every formula in the base language that holds in the background universe, it holds in some universe. Self-Perception goes as far as expressing that the background universe is isomorphic

¹⁹This set theory is ZF minus Foundation and Replacement, and with Separation only for Δ_0 -formulas.

 $^{^{20}}$ This set theory is ZF minus Powerset, and with Separation and Collection (instead of Replacement) only for Δ_0 -formulas.

to a universe in the multiverse. These axioms were interpreted in systems of various types of iterated reflection over ZF, all of which are very mild in terms of consistency strength; indeed their consistency strengths are all bounded by Morse-Kelley class theory with Global Choice (and the remark following System 5.3 indicates that they are far weaker), which in turn is far weaker than ZFC + "there exists an inaccessible cardinal".

Having this framework available, the multiverse theorist can proceed to make use of its untyped relation of truth-in-a-universe. Apart from the light it sheds on the axioms above, a concrete value added, compared to the usual \models -relation, is that it makes it possible to express multiverse principles that reach arbitrarily deep into the structure of universes, universes within universes, universes within universes, etc. We give the final word to Tomas Tranströmer, through his poem "Romanska bågar" (translation by Robert Bly):

Inne i dig öppnar sig valv bakom valv oändligt. Du blir aldrig färdig, och det är som det skall.

Inside you one vault after another opens endlessly. You'll never be complete, and that's as it should be.

Acknowledgements

This research was supported by the Swedish Research Council (VR) [2017-05111], and by the Knut and Alice Wallenberg Foundation (KAW) [2015.0179]. The authors are grateful for the comments and suggestions of the anonymous referee, to Anton Broberg for permitting us to incorporate his argument for Lemma 5.1, and to Ali Enayat for answering our question on the relationship between Self-Perception and recursive saturation (see §1.1).

References

- A. Broberg. Private communication (2021).
- C. C. Chang and H. J. Keisler. Model Theory. Elsevier Science Publishers (1990).
- A. Enayat. Condensable models of set theory. arXiv:1910.04029 [math.LO]
- H. Friedman. Countable models of set theories. In A.R.D. Mathias and H. Rogers, eds., Cambridge Summer School in Mathematical Logic. Springer-Verlag (1973).
- H. Friedman and M. Sheard. An axiomatic approach to self-referential truth. Annals of Pure and Applied Logic, 33 (1987), pp. 1–21.
- K. Fujimoto. Classes and truths in set theory. Annals of Pure and Applied Logic, 163 (2012), pp. 1484–1523.
- V. Gitman and J. D. Hamkins. A natural model of the multiverse axioms. Notre Dame Journal of Formal Logic, no. 4 (2010), pp. 475–484.
- P. K. Gorbow. Rank-initial embeddings of non-standard models of set theory. Arch. Math. Logic, 59 (2020), pp. 517–563.
- A. Gupta. Truth and paradox, Journal of Philosophical Logic, 11 (1982), pp. 1-60.

- G. Jäger, J. Krähenbühl. Σ_1^1 choice in a theory of sets and classes. In Ways of Proof Theory, R. Schindler (ed.), Ontos Verlag, Frankfurt (2010), pp. 283–314.
- V. Halbach. Axiomatic theories of truth, Revised edition. Cambridge University Press (2014).
- V. Halbach, H. Leitgeb and P. Welch. Possible Worlds Semantics for Predicates. In Intensionality, R. Kahle (ed.), Lecture Notes in Logic 22, Association of Symbolic Logic, A K Peters, Wellesley (2005), pp. 20-41.
- J. D. Hamkins. *The set-theoretic multiverse*. The Review of Symbolic Logic, vol. 5, no. 3 (September 2012), pp. 416–449.
- H. G. Herzberger. *Notes on naive semantics*. Journal of Philosophical Logic, 11 (1982), pp. 61–102.
- H. G. Herzberger. Naive semantics and the liar paradox. Journal of Philosophy, 79 (1982), pp. 479–497.
- V. McGee. How truthlike can a predicate be? A negative result. Journal of Philosophical Logic, 14 (1985), pp. 399–410.
- J. Peacock. Cosmological Physics. Cambridge University Press (1998).
- J. -P. Ressayre. *Modèles non Standard et Sous-Systèmes Remarquables de ZF*. In Modèles non Standard en arithmétique et théorie des ensambles. Publications mathématiques de l'Université Paris VII, no. 22, U.E.R. de Mathématiques, Paris (1987), pp. 47-147.
- G. M. Wilmers. Ph.D. Thesis, Oxford (1975).