## WEAKLY 2-RANDOMS AND 1-GENERICS IN SCOTT SETS

LINDA BROWN WESTRICK

ABSTRACT. Let S be a Scott set, or even an  $\omega$ -model of WWKL. Then for each  $A \in S$ , either there is  $X \in S$  that is weakly 2-random relative to A, or there is  $X \in S$  that is 1-generic relative to A. It follows that if  $A_1, \ldots, A_n \in S$  are non-computable, there is  $X \in S$  such that each  $A_i$  is Turing incomparable with X, answering a question of Kučera and Slaman. More generally, any  $\forall \exists$  sentence in the language of partial orders that holds in  $\mathcal{D}$  also holds in  $\mathcal{D}^S$ , where  $\mathcal{D}^S$  is the partial order of Turing degrees of elements of S.

Kučera and Slaman [3] showed that for every Scott set S and every non-computable  $A \in S$ , there is  $X \in S$  such that X and A are Turing incomparable. Their non-uniform proof hinged on whether A was a K-trivial set. Conidis [1] generalized this result to require only that S is an  $\omega$ -model of WWKL (meaning that for every  $A \in S$ , every  $\Pi_1^0(A)$  class of positive measure has an element in S). Kučera and Slaman left open whether, given non-computable  $A, B \in S$ , one could find  $X \in S$  incomparable with both. We answer that question in the affirmative, generalizing both of the above results (Corollary 1).

This is related to the question, asked by Kučera and Slaman, of which  $\forall \exists$  sentences (in the language of partial orders) hold in  $\mathcal{D}^{\mathcal{S}}$ , the partial order induced by  $\leq_T$  on the degrees of  $\mathcal{S}$ . One direction of the proof of the decidability of the  $\forall \exists$  theory of the Turing degrees  $\mathcal{D}$  rests on a technical result (see [4, Theorem II.4.11]) which we generalize from  $2^{\omega}$  to an arbitrary Scott set  $\mathcal{S}$  in Theorem 3. Whenever Theorem 3 holds for some  $\mathcal{S}$ , then every  $\forall \exists$  sentence that holds in  $\mathcal{D}$  also holds in  $\mathcal{D}^{\mathcal{S}}$  (Corollary 2). Li and Slaman [5] recently showed that whenever  $\mathcal{S} \subseteq \Delta_2^0$  is a Scott set and  $A \in \mathcal{S}$ , then there is  $G \in \mathcal{S}$  such that G is 1-generic relative to A. Theorem 3 and Corollary 2 for the case where  $\mathcal{S} \subseteq \Delta_2^0$  follow immediately.

Theorem 2 is the new result which allows the usual proof of Theorem 3 for  $S = 2^{\omega}$  to lift to arbitrary  $\omega$ -models of WWKL. Corollary 1 is also a special case of Theorem 3.

The author would like to thank Uri Andrews, Joe Miller, Noah Schweber and Mariya Soskova, with whom some partial results were obtained by a different method during the author's visit to UW-Madison. This note would not exist without the benefit of those discussions.

To prove Theorem 2, we relativize the following theorem characterizing the weakly 2-random (W2R) sets in the Martin-Löf random (MLR) sets. The relativizations of these notions to a set A are denoted by  $W2R^A$  and  $MLR^A$  respectively.

**Theorem 1** (Hirschfeldt and Miller, see [6, Thm 5.3.16]; Downey, Nies, Weber and Yu [2]). An MLR set is W2R if and only if it does not compute any non-computable c.e. set.

**Theorem 2.** For any Scott set S and any  $A \in S$ , either there is  $X \in S$  that is weakly 2-random relative to A, or there is  $X \in S$  that is 1-generic relative to A.

*Proof.* Let  $\mathcal{P}$  be a  $\Pi_1^0(A)$  class consisting entirely of  $MLR^A$  reals. Let  $X \in S \cap \mathcal{P}$ , and suppose that X is not  $W2R^A$ . By the relativization of Theorem 1 to A, there is a set  $D \leq_T X \oplus A$  such that D is c.e. in A and  $D \not\leq_T A$ . Relativizing the proof that every non-computable c.e. set computes a 1-generic, there is a  $G \leq_T D \oplus A$  such that G is 1-generic relative to A.

Because  $\mathcal{P}$  has positive measure, Theorem 2 and all the other results here hold even if  $\mathcal{S}$  is only an  $\omega$ -model of WWKL. If every element of  $\mathcal{S}$  is  $\Delta_2^0$ , then no element is W2R, and if every element of  $\mathcal{S}$  is hyperimmune-free, then no element is 1-generic, so the result is sharp. **Corollary 1.** For any Scott set S and any non-computable  $A_1, \ldots, A_n \in S$ , there is  $X \in S$  such that X is Turing incomparable with each  $A_i$ .

*Proof.* Let  $A = \bigoplus_{i \leq n} A_i$ . If some  $X \in S$  is 1-generic relative to A, then X is 1-generic relative to each  $A_i$ , which suffices. So suppose some  $X \in S$  is  $W2R^A$ . Then for each  $i, X \not\geq_T A_i$ , because  $\{Y : Y \geq_T A_i\}$  is a  $\Sigma_3^0(A_i)$  set of measure 0.

The same method gives a more general extension of embeddings result, entirely analogous to the extension of embeddings result that holds in  $\mathcal{D}$ . The only addition is the observation that a weakly 2-random can play the same role as a 1-generic in the usual proof of this theorem in  $\mathcal{D}$ .

If  $\mathcal{H} = (H, \leq_H), \mathcal{M} = (M, \leq_M)$  are partial orders, then  $\mathcal{H} \prec \mathcal{M}$  means that  $H \subseteq M$  and  $\leq_H$ and  $\leq_M$  agree on elements of H. We say that  $\mathcal{M}$  is an *end-extension* of  $\mathcal{H}$  if for every  $h \in H$  and  $m \in M \setminus H$ , we have  $m \not\leq_M h$ . If  $\mathcal{H}$  is an upper semi-lattice with join  $\vee_H$ , we say that  $\mathcal{M}$  respects the joins of  $\mathcal{H}$  if for every  $h, k \in H$  and  $m \in M \setminus H$ , if  $m \geq_M h$  and  $m \geq_M k$ , then  $m \geq_M h \vee_H k$ .

**Theorem 3.** Let  $\mathcal{H} \prec \mathcal{M}$  be finite partial orders, where  $\mathcal{H}$  is an upper semi-lattice. Let  $\mathcal{S}$  be a Scott set, and let  $f : \mathcal{H} \to \mathcal{S}$  be an upper semi-lattice embedding of  $\mathcal{H}$  into  $(\mathcal{S}, \leq_T, \oplus)$ . Suppose  $\mathcal{M}$  is an end-extension of  $\mathcal{H}$  which respects the joins of  $\mathcal{H}$ . Then f can be extended to  $\hat{f} : \mathcal{M} \to \mathcal{S}$  which is a partial order embedding of  $\mathcal{M}$  into  $(\mathcal{S}, \leq_T)$ .

*Proof.* For  $h \in H$ , let  $A_h = f(h)$ . Let  $G \in S$  be either 1-generic or weakly 2-random relative to  $\bigoplus_{h \in H} A_h$ . Interpret G as the join of finitely many columns, one for each  $m \in M \setminus H$ , so that the reals  $G_m$  are defined by  $G = \bigoplus_{m \in M \setminus H} G_m$ . For each  $m \in M \setminus H$ , define

$$A_m = \bigoplus_{\substack{h \in H \\ h \leq Mm}} A_h \oplus \bigoplus_{\substack{\ell \in M \setminus H \\ \ell \leq Mm}} G_\ell.$$

We claim that  $\hat{f}(m) := A_m$  satisfies the conclusion of the theorem. If G is 1-generic, we have just repeated the usual construction. If G is weakly 2-random, there is only one verification step that is different: we claim that if  $h \in H$  and  $m \in M \setminus H$  with  $h \not\leq_M m$ , then  $A_h \not\leq A_m$ . Let

$$\mathcal{B} = \{ Z : \bigoplus_{\substack{k \in H \\ k \le M m}} A_k \oplus Z \ge_T A_h \}$$

If  $\mathcal{B}$  had positive measure, a majority vote argument would provide a computation to show  $\bigoplus_{\substack{k \in H \\ k \leq Mm}} A_k \geq_T A_h$ . This is not possible because  $m \not\geq_M h$  and  $\mathcal{M}$  respects the joins of  $\mathcal{H}$ . Therefore  $\mathcal{B}$  is null, and it is  $\Sigma_3^0(\bigoplus_{k \in H} A_k)$ , so  $G \notin \mathcal{B}$ , so  $A_m \notin \mathcal{B}$ .

All the other steps in the verification are the same after the observation that because G is  $MLR^{\bigoplus_{k\in H}A_k}$ , we have  $\bigoplus_{h\in H}A_h\oplus\bigoplus_{\ell\neq m}G_\ell \not\geq_T G_m$  for all  $m\in M\setminus H$ .

We can now give a partial answer to the question asked in [3] of whether the  $\forall \exists$  theory (in the language of partial orders) of the degrees of a Scott set coincides with the  $\forall \exists$  theory of  $\mathcal{D}$ .

**Corollary 2.** For any Scott set S, any  $\forall \exists$  sentence that holds in  $\mathcal{D}$  also holds in  $\mathcal{D}^S$ .

Proof. An extension sentence is a sentence of the form  $\forall \bar{x} \exists \bar{y}(\theta(\bar{x}) \to \bigvee_{j < n} \eta_j(\bar{x}, \bar{y}))$ , where  $\theta(\bar{x})$ and each  $\eta_j(\bar{x}, \bar{y})$  are conjunctions of atomic formulas and negated atomic formulas describing the complete  $\leq$ -diagram on their inputs, where the diagram described by  $\theta$  is an upper semi-lattice, and where each  $\eta_j$  describes a diagram extending the diagram given by  $\theta$ . It is known (see [4, Theorem VII.4.4]) that for any  $\forall \exists$  sentence  $\psi$ , there is a finite conjunction of extension sentences  $\phi$ , such that in any upper semi-lattice,  $\psi$  holds if and only if all the conjuncts hold. Furthermore, in  $\mathcal{D}$ , an extension sentence  $\phi$  holds if and only if there is a j such that the diagram  $\mathcal{M}_j$  described by  $\eta_j$  is an end extension of the diagram  $\mathcal{H}$  described by  $\theta$ , and  $\mathcal{M}$  respects the joins of  $\mathcal{H}$  (see [4, Theorems II.4.11, VII.4.1]). Applying Theorem 3, if an extension sentence  $\phi$  holds in  $\mathcal{D}$ , then  $\phi$  also holds in  $\mathcal{D}^{\mathcal{S}}$ . Therefore, if  $\psi$  holds in  $\mathcal{D}$ , then all the conjuncts  $\phi$  hold in both structures, so  $\psi$  holds in  $\mathcal{D}^{\mathcal{S}}$  as well.  $\Box$ 

It remains open whether all Scott sets satisfy exactly these  $\forall \exists$  sentences.

## References

- Chris J. Conidis. A measure-theoretic proof of Turing incomparability. Ann. Pure Appl. Logic, 162(1):83–88, 2010.
- [2] Rod Downey, Andre Nies, Rebecca Weber, and Liang Yu. Lowness and Π<sup>0</sup><sub>2</sub> nullsets. J. Symbolic Logic, 71(3):1044– 1052, 2006.
- [3] Antonín Kučera and Theodore A. Slaman. Turing incomparability in Scott sets. Proc. Amer. Math. Soc., 135(11):3723–3731, 2007.
- [4] Manuel Lerman. Degrees of unsolvability. Perspectives in Mathematical Logic. Springer-Verlag, Berlin, 1983. Local and global theory.
- [5] Wei Li and Theodore A. Slaman. Private communication.
- [6] André Nies. Computability and randomness, volume 51 of Oxford Logic Guides. Oxford University Press, Oxford, 2009.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CONNECTICUT, STORRS, CONNECTICUT U.S.A. *E-mail address*: westrick@uconn.edu