
ON THE COMMUTATIVITY OF PULL-BACK AND PUSH-FORWARD FUNCTORS ON MOTIVIC CONSTRUCTIBLE FUNCTIONS

by

Jorge Cely & Michel Raibaut

Abstract. — In this article, we study the commutativity between the pull-back and the push-forward functors on constructible functions in Cluckers–Loeser motivic integration.

Introduction

Let k be a characteristic zero field. By analogy with integration over local fields, Kontsevich introduced in [16] an integration theory on finite dimensional vector spaces over $k((t))$, called *motivic integration*, with values in a Grothendieck ring of varieties $K_0(\text{Var}_k)$. Later, Cluckers – Loeser in [8] (announced in [4], [5]) generalized that construction allowing in particular integrals with parameters in the context of henselian valued fields of equal characteristic zero, and in [14], Hrushovski – Kazhdan treated the case of algebraically closed valued fields of equal characteristic zero.

For any *definable sets* X , given by a first order formula in the Denef-Pas language, Cluckers – Loeser construct in [8] an algebra $C(X)$ of *constructible motivic functions* defined on X , and in [7] and [9] they enlarge it in an algebra $C(X)^{\text{exp}}$ of *exponential constructible functions*. Moreover, for any definable function $f : X \rightarrow Y$, they define a pull-back functor $f^* : C(Y)^{\text{exp}} \rightarrow C(X)^{\text{exp}}$, an abelian subgroup $I_Y C(f)^{\text{exp}}$ of $C(X)^{\text{exp}}$ of f -integrable constructible functions and a push-forward functor $f_! : I_Y C(f)^{\text{exp}} \rightarrow C(Y)^{\text{exp}}$ which corresponds to an integration along fibers of f . Roughly speaking, for any definable set X , there exists integers m , n and r such that for any k -field extension K , the set of K -rational points $X(K)$ is contained in $K((t))^m \times K^n \times \mathbb{Z}^r$. Such definable set admits a cell decomposition and similarly to the construction of the integration against Euler characteristic in the real semi-algebraic setting, the construction of the functor $f_!$ is given by an induction process on the valued field dimension. We recall in the first section of this article main ideas, definitions and results of these constructions.

In [19] the second author introduced a notion of *definable distributions* in Cluckers–Loeser motivic setting. He introduced also a notion of *motivic wave front set*, which

allows him, as in the real setting [13] or in the p-adic setting [12] and [3], to study the pull-back of a distribution by a function which requires the natural following commutativity relation between pull-back and push-forward functors

Theorem. — Let X , W and W' be definable sets over k , let γ be a definable morphism from W to W' . We denote by π_W the projection from $W \times X$ to W and by $\pi_{W'}$ the projection from $W' \times X$ to W' . Let $[\varphi]$ be a constructible exponential function in $C(W' \times X)^{exp}$.

1. If $[\varphi]$ is $\pi_{W'}$ -integrable then $[(\gamma \times Id_X)^*\varphi]$ is π_W -integrable. Furthermore, if γ is onto then this implication is an equivalence.
2. If $[\varphi]$ satisfies the condition (1) then

$$\pi_{W!}[(\gamma \times Id_X)^*\varphi] = \gamma^*(\pi_{W'!}[\varphi]).$$

In this article, we start in section 1 by recalling main definitions and ideas of Cluckers–Loeser motivic integration. Then, in section 2, we prove above theorem in a slightly more general context (lemma 1.18 and theorem 1.19) following all different steps of the construction of the theory as the induction process on the valued field dimension using cell decompositions and the computation at the residue and value group levels.

1. Motivic integration and constructible motivic functions

For the reader's convenience we shall start by recalling briefly some definitions, notations and constructions from [8] and [9] that will be used in this article. For an introduction to this circle of ideas we refer to the surveys [6], [2] and [11] and the notes [4], [5] and [7].

1.1. Denef-Pas, Presburger language. — We fix a field k of characteristic zero and we denote by Field_k the category of fields containing k . For any field K in this category we consider the field of Laurent series $K((t))$ endowed with its natural *valuation*

$$\text{ord} : K((t)) \setminus \{0\} \longrightarrow \mathbb{Z}$$

extended by $\text{ord } 0 = +\infty$, and with the *angular component* mapping

$$\overline{\text{ac}} : K((t)) \rightarrow K$$

defined by $\overline{\text{ac}}(x) = xt^{-\text{ord } x} \pmod t$ if $x \neq 0$ and $\overline{\text{ac}}(0) = 0$.

We shall use the three sorted language introduced by Denef and Pas in [17]

$$\mathcal{L}_{DP,P} = (\mathbf{L}_{\text{Val}}, \mathbf{L}_{\text{Res}}, \mathbf{L}_{\text{Ord}}, \text{ord}, \overline{\text{ac}})$$

with sorts corresponding respectively to *valued field*, *residue field* and *value group* variables. The languages \mathbf{L}_{Val} and \mathbf{L}_{Res} are the ring language $\mathbf{L}_{\text{Rings}} = (+, -, \cdot, 0, 1)$ and the language \mathbf{L}_{Ord} is the Presburger language

$$\mathbf{L}_{\text{PR}} = \{+, -, 0, 1, \leq\} \cup \{\equiv_n \mid n \in \mathbb{N}, n > 1\},$$

with \equiv_n symbols interpreted as equivalence relation modulo n . Symbols ord and $\overline{\text{ac}}$ will be interpreted respectively as valuation and angular component, so that for any K in Field_k the triple $(K((t)), K, \mathbb{Z})$ is a structure for $\mathcal{L}_{DP,P}$. We shall also add constant symbols in the Val-sort and in the Res-sort for elements of $k((t))$, resp. of k .

We will work with the $\mathcal{L}_{DP,P}$ -theory $H_{\overline{\text{ac}},0}$ of structures whose valued field is Henselian, with characteristic zero residue field, and with value group \mathbb{Z} . Denef and Pas proved in [17] the following theorem on elimination of valued field quantifiers.

Theorem 1.1 (Denef-Pas [17], Presburger [18]). — *Every formula $\phi(x, \xi, \alpha)$ without parameters in the $\mathcal{L}_{DP,P}$ -language, with x variables in the Val-sort, ξ variables in the Res-sort and α variables in the Ord-sort is $H_{\overline{\text{ac}},0}$ -equivalent to a finite disjunction of formulas of the form*

$$\psi(\overline{\text{ac}} f_1(x), \dots, \overline{\text{ac}} f_k(x), \xi) \wedge \eta(\text{ord } f_1(x), \dots, \text{ord } f_k(x), \alpha),$$

with ψ a \mathbf{L}_{Res} -formula, η a \mathbf{L}_{Ord} -formula without quantifiers and f_1, \dots, f_k polynomials in $\mathbb{Z}[x]$. The theory $H_{\overline{\text{ac}},0}$ admits elimination of quantifiers in the valued field sort.

1.2. Definable subassignments. — From now on we will work with the Denef-Pas language enriched with constant symbols in the Val-sort and in the Res-sort for elements of $k((t))$, resp. of k , and we will denote this language also by $\mathcal{L}_{DP,P}$.

1.2.1. Definable subassignments and definable morphisms. — Let φ be a formula respectively in m , n and r free variables in the various sorts. For every field K in Field_k , we denote by $h_\varphi(K)$ the subset of

$$h[m, n, r](K) := K((t))^m \times K^n \times \mathbb{Z}^r$$

consisting of points satisfying φ . The assignment $K \mapsto h_\varphi(K)$ is called a *definable subassignment* or *definable set*. For instance we will denote by $\{*\}$ the definable subassignment $h[0, 0, 0]$ defined by $K \mapsto \text{Spec } K$. A *definable morphism* F between two definable subassignments h_φ and h_ψ is a collection of applications parametrized by K in Field_k

$$F(K) : h_\varphi(K) \rightarrow h_\psi(K)$$

such that the graph map $K \mapsto \text{Graph } F(K)$ is a definable subassignment. Definable subassignments and definable morphisms are precisely objects and morphisms of the category of definable subassignments over k denoted by Def_k . More generally, for any definable subassignment S in Def_k , we will consider the category Def_S of definable subassignments over S whose objects are definable morphisms θ_Z in Def_k from a definable Z to S and morphisms are definable maps $g : Y \rightarrow Z$ such that $\theta_Y = \theta_Z \circ g$. Sometimes, instead of using θ_Z , we will simply say that Z is a definable set in Def_S .

1.2.2. Finiteness of some definable functions. — We deduce from theorem 1.1 on quantifier elimination the following corollary

Corollary 1.2. — *For non negative integers m, n and r , every definable map from $h[0, n, 0]$ to $h[m, 0, r]$ or from $h[0, 0, r]$ to $h[m, n, 0]$ or from $h[0, n, r]$ to $h[m, 0, 0]$ takes finitely many values.*

1.2.3. Points and fibers. — A point x of a definable set X is by definition a couple $x = (x_0, K)$ where K is an extension of k and x_0 is a point of $X(K)$. The field K will be denoted by $k(x)$ and called *residue field* of x .

Let f be a definable morphism from a subassignment of $h[m, n, r]$ denoted by X to a subassignment of $h[m', n', r']$ denoted by Y . Let $\varphi(x, y)$ be the formula which describes the graph of f , where x runs over $h[m, n, r]$ and y runs over $h[m', n', r']$. For every point $y = (y_0, k(y))$ of Y , the *fiber* X_y is the object of $\text{Def}_{k(y)}$ defined by the formula $\varphi(x, y_0)$ which has coefficients in $k(y)$ and $k(y)((t))$. Taking fibers at y gives rise to a functor $i_y^* : \text{Def}_Y \rightarrow \text{Def}_{k(y)}$.

1.2.4. Dimension. — For any positive integer m , an algebraic subvariety Z of $\mathbb{A}_{k((t))}^m$ induces a definable subassignment h_Z of $h[m, 0, 0]$ with $h_Z(K)$ equal to $Z(K((t)))$ for any extension K of k . The *Zariski closure* of a subassignment S of $h[m, 0, 0]$ is by definition the subassignment of the intersection W of all algebraic subvarieties Z of $\mathbb{A}_{k((t))}^m$ such that h_Z contained S . The *dimension* $\text{Kdim } S$ of S is naturally defined as $\dim W$. More generally, the dimension of a subassignment S of $h[m, n, r]$ is defined as the dimension $\text{Kdim } p(S)$ where p is the projection from $h[m, n, r]$ to $h[m, 0, 0]$. It is proved in [8], using results of Pas [17] and van den Dries [20], that isomorphic definable subassignments in Def_k have the same dimension.

1.3. Grothendieck rings and exponentials. —

1.3.1. The category $\text{RDef}_k^{\text{exp}}$. — For any definable subassignment Z in Def_k , the subcategory RDef_Z of Def_Z whose objects are definable morphisms π_Y , with Y a subassignment of a product $Z \times h[0, n, 0]$, n a non negative integer and π_Y the canonical projection on Z , has been introduced in [8].

Example 1.3. — If Z is the point $h[0, 0, 0]$, then the subcategory RDef_Z is the category of definable sets in the ring language with coefficients from k .

More generally, in [9] motivic additive characters were considered in this context through the category $\text{RDef}_Z^{\text{exp}}$ whose objects are triples (π_Y, ξ, g) with π_Y a definable set in RDef_Z , ξ a definable morphism from Y to $h[0, 1, 0]$ and g a definable morphism from Y to $h[1, 0, 0]$. A morphism from $(\pi_{Y'}, \xi', g')$ to (π_Y, ξ, g) in $\text{RDef}_Z^{\text{exp}}$ is a morphism h from Y' to Y satisfying the equalities

$$\pi_{Y'} = \pi_Y \circ h, \quad \xi' = \xi \circ h, \quad g' = g \circ h.$$

Remark 1.4. — The functor $\pi_Y \mapsto (\pi_Y, 0, 0)$ allows to identify RDef_Z as a full subcategory of $\text{RDef}_Z^{\text{exp}}$.

1.3.2. The Grothendieck ring $K_0(\text{RDef}_Z^{\text{exp}})$. — As an abelian group it is the free abelian group over symbols $[\pi_Y, \xi, g]$ modulo the following relations :

Isomorphism. For any isomorphic (π_Y, ξ, g) and $(\pi_{Y'}, \xi', g')$,

$$(R1) \quad [\pi_Y, \xi, g] = [\pi_{Y'}, \xi', g']$$

Additivity. For π_Y and $\pi_{Y'}$ definable subassignments of some π_X in RDef_Z and for ξ and g defined on $Y \cup Y'$

$$(R2) \quad [\pi_{Y \cup Y'}, \xi, g] + [\pi_{Y \cap Y'}, \xi|_{Y \cap Y'}, g|_{Y \cap Y'}] = [\pi_Y, \xi|_Y, g|_Y] + [\pi_{Y'}, \xi|_{Y'}, g|_{Y'}].$$

Compatibility with reduction. For any π_Y in Def_Z , for any definable morphism f from Y to $h[1, 0, 0]$ with $\text{ord } f(y) \geq 0$ for any $y \in Y$ we have

$$(R3) \quad [\pi_Y, \xi, g + f] = [\pi_Y, \xi + \bar{f}, g]$$

with \bar{f} the reduction of f modulo (t) .

Sum over the line. Let p be the canonical projection from $Y[0, 1, 0]$ to $h[0, 1, 0]$. If the morphisms $\pi_{Y[0, 1, 0]}$, g and ξ all factorize through the canonical projection from $Y[0, 1, 0]$ to Y then

$$(R4) \quad [Y[0, 1, 0] \rightarrow Z, \xi + p, g] = 0.$$

This Grothendieck group is endowed with a ring structure by setting

$$(R5) \quad [\pi_Y, \xi, g] \cdot [\pi_{Y'}, \xi', g'] = [\pi_{Y \otimes_Z Y'}, \xi \circ p_Y + \xi' \circ p_{Y'}, g \circ p_Y + g' \circ p_{Y'}]$$

where $Y \otimes_Z Y'$ is the fiber product of Y and Y' above Z , p_Y is the projection to Y and $p_{Y'}$ is the projection to Y' . The element $[\text{Id}_Z, 0, 0]$ is the multiplicative unit of $K_0(\text{RDef}_Z^{\text{exp}})$. The Grothendieck ring $K_0(\text{RDef}_Z)$ is defined as above and the functor defined in Remark 1.4 induces an injection $K_0(\text{RDef}_Z) \rightarrow K_0(\text{RDef}_Z^{\text{exp}})$.

Remark 1.5. — The element $[\pi_Y, \xi, g]$ of the Grothendieck ring $K_0(\text{RDef}_Z^{\text{exp}})$ will be denoted by $e^\xi E(g)[\pi_Y]$. We will abbreviate $e^0 E(g)[\pi_Y]$ by $E(g)[\pi_Y]$, $e^0 E(0)[\pi_Y]$ by $[\pi_Y]$ and $e^0 E(g)[\text{Id}_Z]$ by $E(g)$.

Remark 1.6 (Interpretation of E). — The element $E(g)$ in $K_0(\text{RDef}_Z^{\text{exp}})$ can be viewed as the exponential (at the valued field level of the definable morphism g from Z to $h[1, 0, 0]$, said otherwise, it is a motivic additive character on the valued field evaluated in g . More precisely, by relations (R3) and (R5), E can be interpreted as a universal additive character which is trivial on the maximal ideal of the valuation ring. This is compatible with specialization to p -adic fields as explained in Section 9 of [9].

Remark 1.7 (Interpretation of e). — The element $e(\xi)$ in $K_0(\text{RDef}_Z^{\text{exp}})$ can be considered as the exponential (at the residue field level) of the definable morphism ξ from Z to $h[0, 1, 0]$. By relation (R4), e can be interpreted as a universal additive character on the residue field. For instance in the case where Z is the point, the relation $[h[0, 1, 0] \rightarrow \{*\}, p, 0] = 0$ should be interpreted as an abstraction of the classical nullity of the sum of a non trivial character over elements of a finite field. Relation (R3) expresses compatibility under reduction modulo the uniformizing parameter between the exponentials over the valued field and over the residue field.

1.3.3. Pull-back and push-forward. — For $f : Z \rightarrow Z'$ in Def_k there is a natural pull-back morphism $f^* : K_0(\text{RDef}_{Z'}^{\text{exp}}) \rightarrow K_0(\text{RDef}_Z^{\text{exp}})$, induced by the fiber product. Furthermore, if f is a morphism in $\text{RDef}_{Z'}$, then composition with f induces a morphism $f_! : K_0(\text{RDef}_Z^{\text{exp}}) \rightarrow K_0(\text{RDef}_{Z'}^{\text{exp}})$.

1.4. Constructible exponential functions. —

1.4.1. Constructible motivic functions. — In the p -adic context (see [10], [15], [9] and [1]), for instance over the field \mathbb{Q}_p itself, one fixes an additive character $\Psi : K \rightarrow \mathbb{C}^\times$ trivial on $p\mathbb{Z}_p$ and non trivial on the set $\text{ord } x = 0$ and one denotes by \mathbb{A}_p the ring $\mathbb{Z}[1/p, 1/(1-p^{-i})]$. For any X contained in some \mathbb{Q}_p^m and definable for the Macintyre language, it is natural to define the \mathbb{A}_p -algebra of constructible functions on X denoted by $\mathcal{C}(X)$ and generated by function of the form $|f| \text{ord}(h)$ where f and h are definable functions from X to \mathbb{Q}_p and h does not vanish. In [9], also a variant with additive characters is introduced, called constructible exponential functions on X and denoted by $\mathcal{C}(X)^{\text{exp}}$. The algebra $\mathcal{C}(X)^{\text{exp}}$ is generated by $\mathcal{C}(X)$ and functions of the form $\psi(g)$ with $g : X \rightarrow \mathbb{Q}_p$ with ψ a nontrivial additive character on \mathbb{Q}_p .

Analogously, in [8] Cluckers and Loeser consider the ring

$$\mathbb{A} = \mathbb{Z} \left[\mathbb{L}, \mathbb{L}^{-1}, \left(\frac{1}{1 - \mathbb{L}^{-i}} \right)_{i>0} \right],$$

where \mathbb{L} is a symbol, and they define the ring $\mathcal{C}(Z)$ of *constructible motivic functions* on a definable set Z by

$$\mathcal{C}(Z) := K_0(\text{RDef}_Z) \otimes_{\mathcal{P}^0(Z)} \mathcal{P}(Z),$$

where $\mathcal{P}(Z)$, called ring of *Presburger constructible functions*, is the subring of the ring of functions from the set of points of Z to \mathbb{A} , generated by constant functions, definable functions from Z to \mathbb{Z} and functions of the form \mathbb{L}^β with β a definable function from Z to \mathbb{Z} . Here, $\mathcal{P}^0(Z)$ is the subring of $\mathcal{P}(Z)$ generated by the constant function \mathbb{L} and the characteristic functions 1_Y of definable subsets Y of the base Z . The tensor product is given by the morphism from $\mathcal{P}^0(Z)$ to $K_0(\text{RDef}_Z)$ sending 1_Y to the class $[Y \rightarrow Z]$ of the canonical injection from Y to Z and sending \mathbb{L} to the class $[Z[0, 1, 0] \rightarrow Z]$ of the canonical projection to Z .

The following proposition [8, Proposition 5.3.1] allows to dissociate the Res-sort with the value group sort.

Proposition 1.8. — *Let S be a definable subassignment of $h[0, n, 0]$.*

– *Let W be a definable subassignment of $h[0, n, 0]$. The canonical morphism*

$$\mathcal{P}(S) \otimes_{\mathcal{P}^0(S)} K_0(\text{RDef}_{S \times W}) \rightarrow \mathcal{C}(S \times W)$$

is an isomorphism.

– *Let W be a definable subassignment of $h[0, 0, r]$. The canonical morphism*

$$K_0(\text{RDef}_S) \otimes_{\mathcal{P}^0(S)} \mathcal{P}(S \times W) \rightarrow \mathcal{C}(S \times W)$$

is an isomorphism.

1.4.2. Constructible exponential functions. — For any definable set Z in Def_k , the ring $\mathcal{C}(Z)^{\text{exp}}$ of *constructible exponential functions* is defined in [9] by

$$\mathcal{C}(Z)^{\text{exp}} := \mathcal{C}(Z) \otimes_{K_0(\text{RDef}_Z)} K_0(\text{RDef}_Z^{\text{exp}}),$$

where we use the morphism $a \mapsto a \otimes 1_Z$ from $K_0(\text{RDef}_Z)$ to $\mathcal{C}(Z)$. For any integer d , we denote by $\mathcal{C}^{\leq d}(Z)^{\text{exp}}$ the ideal of *constructible functions of K -dimension $\leq d$* ,

namely the ideal generated by the characteristic functions $1_{Z'}$ of subassignments Z' of Z of dimension at most d . A constructible function has *K-dimension* d , if it belongs to $\mathcal{C}^{\leq d}(Z)^{\text{exp}} \setminus \mathcal{C}^{\leq d-1}(Z)^{\text{exp}}$. This family of ideals is a filtration of the ring $\mathcal{C}(Z)^{\text{exp}}$ and the graded ring associated

$$C(Z)^{\text{exp}} = \bigoplus_{d \in \mathbb{N}} \mathcal{C}^{\leq d}(Z)^{\text{exp}} / \mathcal{C}^{\leq d-1}(Z)^{\text{exp}}$$

is called ring of *constructible exponential Functions*.

Remark 1.9. — Constructible Functions can be compared to the equivalence classes of Lebesgue measurable functions (equality up to a zero measure set). In this article we will just write function for Function; the difference still being visible in the notation $C(Z)^{\text{exp}}$ versus $\mathcal{C}(Z)^{\text{exp}}$.

1.5. Cell decomposition. — Let C be a definable subassignment in Def_k . Let $\alpha : C \rightarrow \mathbb{Z}$, $\xi : C \rightarrow h[0, 1, 0] \setminus \{0\}$ and $c : C \rightarrow h[1, 0, 0]$ be definable morphisms.

- The cell $Z_{C,c,\alpha,\xi}$ with basis C , center c , order α and angular component ξ , is

$$Z_{C,c,\alpha,\xi} = \left\{ (y, z) \in C[1, 0, 0] \left| \begin{array}{l} \text{ord}(z - c(y)) = \alpha(y) \\ \overline{\text{ac}}(z - c(y)) = \xi(y) \end{array} \right. \right\}$$

Note that this definable set is a family of balls $B(c(y) + \xi(y)t^{\alpha(y)}, \alpha(y) + 1)$ parametrized by the base C . The axiom (Axiom 8), below gives the push-forward morphism corresponding to the projection of this cell on its base C , that is, integration in the fibers of this projection map.

- The cell $Z_{C,c}$ with basis C and center c is

$$Z_{C,c} = \{(y, z) \in C[1, 0, 0] \mid z = c(y)\}.$$

The change of variables formula (theorem 1.14) gives in particular, the push-forward morphism corresponding to the projection of that cell on its base.

More generally, a definable subassignment Z of $S[1, 0, 0]$ for some S in Def_k is called a *1-cell* or a *0-cell* if there exists a definable isomorphism

$$\lambda : Z \rightarrow Z_{C,c,\alpha,\xi} \subset S[1, s, r] \text{ or } \lambda : Z \rightarrow Z_{C,c} \subset S[1, s, 0],$$

called *presentation* of the cell Z , where the base C is contained in $S[0, s, r]$ and such that the morphism $\pi \circ \lambda$ is the identity on Z with π the projection to $S[1, 0, 0]$.

Let us state a variant of Denef-Pas Cell Decomposition theorem [17], theorem 7.2.1 of [9], that will be used in the proof of the projection lemma 1.18 in subsection 2.3.

Theorem 1.10 (Cell decomposition). — *Let X be a definable subassignment of $S[1, 0, 0]$ with S in Def_k .*

1. *The definable set X is a finite disjoint union of cells.*
2. *For every φ in $\mathcal{C}(X)$ there exists a finite partition of X into cells Z_i with presentation (λ_i, Z_{C_i}) such that $\varphi|_{Z_i} = \lambda_i^* p_i^*(\psi_i)$, with ψ_i in $\mathcal{C}(C_i)$ and $p_i : Z_{C_i} \rightarrow C_i$ the projection. Similar statements hold for φ in $\mathcal{P}(X)$ and in $K_0(\text{RDef}_X)$.*

1.6. Pull-back of constructible exponential functions. — A definable map $f : Z \rightarrow Z'$ in Def_k induces a pull-back morphism (cf. [8, §5.4] and in [9, §3.4])

$$f^* : \mathcal{C}(Z')^{\text{exp}} \rightarrow \mathcal{C}(Z)^{\text{exp}}.$$

Indeed, the fiber product along f induces a pull-back morphism from $K_0(\text{RDef}_{Z'})^{\text{exp}}$ to $K_0(\text{RDef}_Z)^{\text{exp}}$ and the composition by f induces also a pull-back morphism from $\mathcal{P}(Z')$ to $\mathcal{P}(Z)$. These pull-backs are compatible with their tensor product.

Remark 1.11. — A constructible exponential function $E(g)e(\xi) \otimes \alpha \mathbb{L}^\beta$ can be thought of as

$$z \in Z \mapsto E(g(z))e(\xi(z)) \otimes \alpha(z) \mathbb{L}^{\beta(z)} \in \mathcal{C}(\{z\})^{\text{exp}}.$$

More generally, the constructible exponential function $[\pi_Y]E(g)e(\xi) \otimes \alpha \mathbb{L}^\beta$ can be thought of as

$$z \in Z \mapsto [Y_z]E(g|_{Y_z})e(\xi|_{Y_z}) \otimes \alpha|_{Y_z} \mathbb{L}^{\beta|_{Y_z}} \in \mathcal{C}(\{z\})^{\text{exp}}.$$

By corollary 1.2, the restrictions $\alpha|_{Y_z}$ and $\beta|_{Y_z}$ take finitely many values, but $[Y_z]$ should be thought of as a kind of motive standing for a possibly infinite sum over elements in Y_z , which is a definable subset of some power of the residue field. With E and e , the expression $[\pi_Y]E(g)e(\xi)$ is a kind of exponential motive, standing for possibly infinite exponential sums. In the p -adic case, the finiteness of the residue field allows one to see $[Y_z]$ as a finite sum again.

1.7. Push-forward of constructible exponential functions. — For S in Def_k , Cluckers and Loeser construct in [8] and [9] a functor I_S^{exp} from the category Def_S to the category $\underline{\text{Ab}}$ of abelian groups:

$$I_S^{\text{exp}} : \begin{cases} \text{Def}_S \rightarrow \underline{\text{Ab}} \\ (\theta_Z : Z \rightarrow S) \mapsto (I_S C(\theta_Z)^{\text{exp}} \subset C(Z)^{\text{exp}}) \\ (\theta_Z \xrightarrow{f} \theta_Y) \mapsto (I_S C(\theta_Z)^{\text{exp}} \xrightarrow{f!} I_S C(\theta_Y)^{\text{exp}}) \end{cases}$$

satisfying natural axioms implying its uniqueness, see theorems 10.1.1 and 13.2.1 in [8] and theorem 4.1.1 in [9]. The elements of $I_S C(\theta_Z)^{\text{exp}}$ are called θ_Z -integrable motivic constructible exponential functions on Z or simply θ_Z -integrable functions.

Example 1.12. — The ring $I_S C(\text{Id}_S)^{\text{exp}}$ is all of $C(S)^{\text{exp}}$, namely, every function in $C(S)^{\text{exp}}$ is already integrable up to S itself, with the identity map $S \rightarrow S$ as structural morphism.

The functor I_S^{exp} and the integrable functions are constructed simultaneously. The functor I_S is first defined in [8] in the setting without exponential and extended in [9] in the exponential setting to I_S^{exp} . In particular, for any $\theta_Z : Z \rightarrow S$ in Def_S , $I_S C(\theta_Z)^{\text{exp}}$ is a graded subgroup of $C(Z)^{\text{exp}}$ defined as

$$I_S C(\theta_Z)^{\text{exp}} := I_S C(\theta_Z) \otimes_{K_0(\text{RDef}_Z)} K_0(\text{RDef}_Z)^{\text{exp}}.$$

Remark 1.13. — Sometimes we will simply say S -integrable instead of θ_Z -integrable and write $I_S C(Z)^{\text{exp}}$ when the structural morphism θ_Z is implicitly clear.

The natural morphism of graded groups from $I_S C(\theta_Z)$ to $I_S C(\theta_Z)^{\text{exp}}$ is injective. We will use the following axioms (see theorem 10.1.1 in [8] and §13.2 in [9]):

Axiom 1 (Fubini). — Let S be in Def_k . Let $f : \theta_X \rightarrow \theta_Y$ be a definable morphism in Def_S . A constructible function φ on X is θ_X -integrable if and only if φ is f -integrable and $f_! \varphi$ is θ_Y -integrable namely:

$$\varphi \in I_S C(\theta_X)^{\text{exp}} \Leftrightarrow \varphi \in I_Y C(f)^{\text{exp}} \text{ and } f_! \varphi \in I_S C(\theta_Y)^{\text{exp}}.$$

Axiom 2 (Additivity). — Let Z be a definable subassignment in Def_S . Assume Z is the disjoint union of two definable subassignments Z_1 and Z_2 . Then, for every morphism $f : Z \rightarrow Y$ in Def_S , the isomorphism $C(Z) \simeq C(Z_1) \oplus C(Z_2)$ induces an isomorphism $I_S C(Z) \simeq I_S C(Z_1) \oplus I_S C(Z_2)$ under which we have $f_! = f_{1!} \oplus f_{2!}$.

Axiom 3 (Projection formula). — Let S be in Def_k . For every morphism f from θ_Z to θ_Y in Def_S , and every α in $\mathcal{C}(Y)^{\text{exp}}$ and β in $I_S C(\theta_Z)^{\text{exp}}$, if $f^*(\alpha)\beta$ belongs to $I_S C(\theta_Z)^{\text{exp}}$, then $f_!(f^*(\alpha)\beta) = \alpha f_!(\beta)$.

Axiom 4 (Push-forward for inclusions). — Let S be in Def_k . Let $\theta_T : T \rightarrow S$ be a definable set in Def_S . Let Z and Z' two definable subassignments of T with $Z \subset Z'$. Let $i : Z \rightarrow Z'$ be the corresponding inclusion and θ_Z and $\theta_{Z'}$ the corresponding restriction of θ_T to Z and Z' . We have $\theta_Z = \theta_{Z'} \circ i$. Composition with i induces a morphism $i_! : K_0(\text{RDef}_Z) \rightarrow K_0(\text{RDef}_{Z'})$. The extension by zero induces a morphism $i_! : \mathcal{P}(Z) \rightarrow \mathcal{P}(Z')$. By compatibility with the tensor product, we get a morphism

$$(1.1) \quad i_! : \mathcal{C}(Z) \rightarrow \mathcal{C}(Z').$$

For every constructible function φ in $\mathcal{C}(Z)$, the class $[\varphi]$ lies in $I_S C(\theta_Z)$ if and only if $[i_!(\varphi)]$ belongs to $I_S C(\theta_{Z'})$. If this is the case then $i_!([\varphi]) = [i_!(\varphi)]$.

The morphism (1.1) is also compatible with the morphism $i_!$ from $K_0(\text{RDef}_Z^{\text{exp}})$ to $K_0(\text{RDef}_{Z'}^{\text{exp}})$ also defined by extension by zero. We obtain in such a way a morphism $i_! : \mathcal{C}(Z)^{\text{exp}} \rightarrow \mathcal{C}(Z')^{\text{exp}}$. As i sends subassignments of Z to subassignments of Z' of the same dimension, there are group morphisms $i_! : \mathcal{C}^{\leq d}(Z)^{\text{exp}} \rightarrow \mathcal{C}^{\leq d}(Z')^{\text{exp}}$ and a graded group morphisms $i_! : \mathcal{C}(Z)^{\text{exp}} \rightarrow \mathcal{C}(Z')^{\text{exp}}$ which restricts to a morphism $i_! : I_S C(\theta_Z)^{\text{exp}} \rightarrow I_S C(\theta_{Z'})^{\text{exp}}$.

Axiom 5 (Projection along k -variables). — Let S be a definable subassignment in Def_k . Let $\theta_Y : Y \rightarrow S$ be in Def_S . Let $n \geq 0$ be an integer. We denote by Z the definable set $Y[0, n, 0]$, by $\pi : Z \rightarrow Y$ the canonical projection, and by θ_Z the structural map $\pi \circ \theta_Y$. By proposition 1.8, there is a canonical isomorphism

$$\mathcal{C}(Z) \simeq K_0(\text{RDef}_Z) \otimes_{\mathcal{P}(0)(Y)} \mathcal{P}(Y)$$

which allows to define a ring morphism $\pi_! : \mathcal{C}(Z) \rightarrow \mathcal{C}(Y)$ by sending $\sum_i a_i \otimes \varphi_i$ to $\sum_i \pi_!(a_i) \otimes \varphi_i$ with a_i in $K_0(\text{RDef}_Z)$, φ_i in $\mathcal{P}(Y)$ and $\pi_!(a_i)$ defined in subsection 1.3.3. For any constructible function φ in $\mathcal{C}(Z)$, the class $[\varphi]$ is π -integrable and

$$\pi_!([\varphi]) = [\pi_!(\varphi)]$$

where $\pi_!$ is defined above.

The map $\pi_! : K_0(\text{RDef}_Z^{\text{exp}}) \rightarrow K_0(\text{RDef}_Y^{\text{exp}})$ induces a ring morphism

$$\pi_! : \mathcal{C}(Z)^{\text{exp}} \rightarrow \mathcal{C}(Y)^{\text{exp}}.$$

Furthermore, as π sends subassignments of Z to subassignments of Y of the same dimension, there are group morphisms $\pi_! : \mathcal{C}^{\leq d}(Z)^{\text{exp}} \rightarrow \mathcal{C}^{\leq d}(Y)^{\text{exp}}$ for all d , and a graded group morphism $\pi_! : \mathcal{C}(Z)^{\text{exp}} \rightarrow \mathcal{C}(Y)^{\text{exp}}$ which restricts to a morphism

$$\pi_! : I_S \mathcal{C}(\theta_Z)^{\text{exp}} \rightarrow I_S \mathcal{C}(\theta_Y)^{\text{exp}}.$$

Axiom 6 (Projection along \mathbb{Z} -variables). — Let S be a definable subassignment in Def_k . Let $\theta_Y : Y \rightarrow S$ be in Def_S . Let $r \geq 0$ be an integer. We denote by Z the definable set $Y[0, 0, r]$, by $\pi : Z \rightarrow Y$ the canonical projection, and by θ_Z the structural map $\pi \circ \theta_Y$.

Let φ be a constructible function in $\mathcal{C}(Z)$. The class $[\varphi]$ is π -integrable if and only if φ can be written as $\varphi = \varphi_Y \otimes \varphi_{\mathcal{P}}$, where φ_Y is a constructible function in $\mathcal{C}(Y)$ and $\varphi_{\mathcal{P}}$ is a Presburger function in $I_Y \mathcal{P}(Z)$, namely $\varphi_{\mathcal{P}}$ is a Y -integrable Presburger function on Z : for any $q > 1$, for any y in Y the family $\sum_{x \in \mathbb{Z}^r} \nu_q(\varphi_{\mathcal{P}}(y, x))$ is summable with $\nu_q : \mathbb{A} \rightarrow \mathbb{R}$ is the unique morphism of rings mapping \mathbb{L} to q . In that case we have

$$\pi_![\varphi] = [\varphi_Y \otimes \pi_!(\varphi_{\mathcal{P}})],$$

where $\pi_!(\varphi_{\mathcal{P}})$ is the unique constructible function in $\mathcal{C}(Y)$ such that for any y in Y , for any $q > 1$

$$\nu_q((\pi_! \varphi_{\mathcal{P}})(y)) = \sum_{x \in \mathbb{Z}^r} \nu_q(\varphi_{\mathcal{P}}(y, x)).$$

This defines a morphism $\pi_! : I_S \mathcal{C}(\theta_Z) \rightarrow I_S \mathcal{C}(\theta_Y)$, which induces by tensor product a graded group morphism

$$\pi_! : I_S \mathcal{C}(\theta_Z)^{\text{exp}} \rightarrow I_S \mathcal{C}(\theta_Y)^{\text{exp}}$$

using the fact that the canonical morphism

$$K_0(\text{RDef}_Y^{\text{exp}}) \otimes_{\mathcal{P}^0(Y)} \mathcal{P}^0(Z) \rightarrow K_0(\text{RDef}_Z^{\text{exp}}),$$

is an isomorphism.

Axiom 7 (Volume of graph; 0-cell). — Let θ_Y be in Def_S , and

$$Z = \{(y, z) \in Y[1, 0, 0] \mid z = c(y)\}$$

where $c : Y \rightarrow h[1, 0, 0]$ is a definable morphism. Denote by $f : Z \rightarrow Y$ the morphism induced by the projection from $Y \times h[1, 0, 0]$ to Y , and θ_Z its composition with θ_Y . Then, the constructible function $[1_Z]$ is θ_Z -integrable if and only if $\mathbb{L}^{(\text{ord } j \text{ acf}) \circ f^{-1}}$ is θ_Y -integrable. In that case, in the ring $I_S \mathcal{C}(Y)^{\text{exp}}$ we have the equality

$$f_!([1_Z]) = \mathbb{L}^{(\text{ord } j \text{ acf}) \circ f^{-1}}.$$

Axiom 8 (Volume of balls; 1-cell). — Let θ_Y be in Def_S , and

$$Z = \{(y, z) \in Y[1, 0, 0] \mid \text{ord}(z - c(y)) = \alpha(y), \overline{\text{ac}}(z - c(y)) = \xi(y)\}$$

where $\alpha : Y \rightarrow \mathbb{Z}$, $\xi : Y \rightarrow h[0, 1, 0] \setminus \{0\}$ and $c : Y \rightarrow h[1, 0, 0]$ are definable morphisms. Denote by $f : Z \rightarrow Y$ the morphism induced by the projection from

$Y \times h[1, 0, 0]$ to Y , and θ_Z its composition with θ_Y . Then, the constructible function $[1_Z]$ is θ_Z -integrable if and only if $\mathbb{L}^{-\alpha-1}[1_Y]$ is θ_Y -integrable. In that case, in the ring $I_S C(Y)^{\text{exp}}$ we have the equality $f_!([1_Z]) = \mathbb{L}^{-\alpha-1}[1_Y]$.

By Axiom 8, the volume of a ball $\{z \in h[1, 0, 0] \mid \text{ord}(z - c) = \alpha, \overline{\text{ac}}(z - c) = \xi\}$ with α in \mathbb{Z} , c in $k((t))$ and ξ in k , $\xi \neq 0$ is $\mathbb{L}^{\alpha-1}$. This is natural by analogy with the p -adic case.

Axiom 9 (Relative balls of large volume). — Let θ_Y be in Def_S and

$$Z = \{(y, z) \in Y[1, 0, 0] \mid \text{ord } z = \alpha(y), \overline{\text{ac}} z = \xi(y)\}$$

where $\alpha : Y \rightarrow \mathbb{Z}$, $\xi : Y \rightarrow h[0, 1, 0] \setminus \{0\}$ are definable morphisms. Let $f : Z \rightarrow Y$ be the morphism induced by the projection from $Y[1, 0, 0]$ to Y . Suppose that the constructible function $[1_Z]$ is $(\theta_Y \circ f)$ -integrable and moreover $\alpha(y) < 0$ holds for every y in Y , then $f_!(E(z)[1_Z]) = 0$.

The previous axiom is also natural by analogy with the p -adic case, where an additive character evaluated in the identity function and integrated over a large ball is naturally zero.

Theorem 1.14 (Change of variables formula, theorem 5.2.1 of [9])

Let $f : X \rightarrow Y$ be a definable isomorphism between definable subassignments of dimension d . Let φ be in $C^{\leq d}(Y)^{\text{exp}}$ with a nonzero class in $C^d(Y)^{\text{exp}}$. Then $[f^*(\varphi)]$ belongs to $I_Y C^d(f)^{\text{exp}}$ and

$$f_!([f^*(\varphi)]) = \mathbb{L}^{\text{ord}(\text{Jac}f) \circ f^{-1}}[\varphi].$$

We give some ideas of the construction of this push-forward and refer to [8] and [9] and to the surveys [2], [6] and [11] for further details. For instance, we fix a base S , we consider a definable morphism $f : Y \rightarrow S$ where Y is a subassignment of some $h[m, n, r]$ and we denote by Γ_f the graph of f . By functionality the morphism $f_!$ is the composition $p_! \circ i_!$ where $i : Y \rightarrow \Gamma_f$ and $p : \Gamma_f \rightarrow S$ are the canonical injection and projection. Thus, it is enough to know how to construct the push-forward morphisms for injections and projections. The case of definable injection is done using extension by zero of constructible functions, and an adjustment with a Jacobian to match the induced measures. Using the axiom of the volume of balls and the change of variables formula, we observe that the construction of the push-forward morphism for a projection is done by induction on the valued field dimension. For instance, Γ_f can be seen as a definable subassignment of $S'[1, 0, 0]$ where S' is the definable set $S[m-1, n, r]$ and the push-forward $p_!$ will be the composition $p_!^{(m-1)} \circ \pi_!$ where $\pi : \Gamma_f \rightarrow S'$ and $p^{(m-1)} : S' \rightarrow S$ are canonical projections. The construction does not depend on the order of such projections and the main tool is the cell decomposition theorem stated above. Once the valuative dimension is zero we have to define a push-forward of a projection from some $S[0, n', r']$ to S . This is done using the independence between the residue field and the value group, coming from theorem 1.1, see for instance Proposition 1.8. The push-forward along residue variables is simply the push-forward induced by composition at the level of Grothendieck ring cf.

Axiom 5 (and [8, §5.6]). The integration along \mathbb{Z} -variables corresponds to summing over the integers, cf. Axiom 6 (and [8, §4.5]).

Example 1.15. — (Integration of constructible functions, [8, §11.1]).

In this example, we illustrate the computation of the integration along a projection $f : S \times Y \rightarrow S$ with $Y = h[1, n, r]$. We will use this computation in subsection 2.3. Let φ be a constructible function in $\mathcal{C}(S[1, n, r])$. By the cell decomposition theorem (theorem 1.10), there is a cell decomposition of $S[1, n, r]$ adapted to φ denoted by $(Z_i)_{i \in I}$. For any i in I , the cell Z_i has a presentation (λ_i, Z_{C_i}) , and there is a constructible function ψ_i in $\mathcal{C}(C_i)$ such that

$$(1.2) \quad \varphi|_{Z_i} = \lambda_i^* p_i^* (\psi_i) 1_{Z_i}$$

where Z_{C_i} is a subassignment of some $h[1, n + n_i, r + r_i]$, C_i is a subassignment of $h[0, n + n_i, r + r_i]$, $p_i : Z_{C_i} \rightarrow C_i$ and $q_i : C_i \rightarrow S$ are the projections. By a refinement of the cell decomposition we can assume that for any i , the restriction $\varphi|_{Z_i}$ is either zero or has the same K -dimension as Z_i . By the additivity axiom (Axiom 2) φ will be f -integrable if and only for any i in I the restriction $\varphi|_{Z_i}$ is f -integrable and in that case

$$f! \varphi = \sum_{i \in I} f! (\varphi|_{Z_i}).$$

For any i in I , we consider the commutative diagram

$$\begin{array}{ccc} Z_i & \xrightarrow[\simeq]{\lambda_i} & Z_{C_i} \\ f \downarrow & & \downarrow p_i \\ S & \xleftarrow{q_i} & C_i \end{array}$$

Using equation (1.2), the projection axiom (Axiom 3) and the Fubini axiom (Axiom 1), the following statement are equivalent

- the restriction $\varphi|_{Z_i}$ is f -integrable,
- the constructible function $p_i^* (\psi_i) [1_{Z_{C_i}}]$ is $f \circ \lambda_i^{-1}$ -integrable namely $q_i \circ p_i$ -integrable,
- the constructible function $\psi_i p_{i!} [1_{Z_{C_i}}]$ is q_i -integrable with $p_{i!} [1_{Z_{C_i}}]$ is an integration of a 0-cell given by (Axiom 7) or a 1-cell given by (Axiom 8).

The q_i -integrability condition and the q_i -integration can be treated by proposition 1.8 and (Axioms 5 and 6). If all these constructible functions are integrable then

$$f! \varphi = \sum_{i \in I} q_{i!} (\psi_i p_{i!} [1_{Z_{C_i}}]).$$

Example 1.16. — (Integration of exponential constructible functions in [9])

We consider two cases.

- Integration along the projection $\pi_S^{S[1,0,0]} : S[1, 0, 0] \rightarrow S$.

Let S be a definable set in Def_k . Let φ be a constructible function in $\mathcal{C}(S[1, 0, 0])^{\text{exp}}$. We can write

$$\varphi = [f : Y \rightarrow S[1, 0, 0]] e^\xi E(g) \otimes \tilde{\varphi}$$

with $Y \subset S[1, n_Y, 0]$. In the construction [9, §5.1], using a cell decomposition adapted to $\tilde{\varphi}$ the authors consider a definable isomorphism

$$\lambda : Y \rightarrow Y' \subset Y[0, n, r]$$

and the following diagram

$$\begin{array}{ccccc} Y & \xrightarrow{f} & S[1, 0, 0] & \xrightarrow{\pi_S} & S \\ \lambda \downarrow \simeq & & & \nearrow p' & \\ Y' & \xrightarrow{\pi'} & S[0, n + n_Y, r] & & \end{array}$$

where π' and p' are projections. They define $\xi' = \xi \circ \lambda^{-1}$, $g' = g \circ \lambda^{-1}$ and $\varphi' = \lambda^{-1*} f^*(\tilde{\varphi})$. By construction there exists also a unique definable function $\tilde{\xi}' : S[0, n_Y + n, r] \rightarrow h[0, 1, 0]$ such that $\xi' = \tilde{\xi}' \circ \pi'$.

We consider an element of Y' as a couple (x, y) with x in $S[0, n_Y, 0]$ and y in $h[1, 0, 0]$. Following from its construction, the authors decompose the definable set Y' as union $A \cup B$, such that $g'(x, \cdot) : y \rightarrow g'(x, y)$ is constant on each fiber B_x and non constant and injective on each fiber A_x where for each x ,

$$A_x = \{y \in h[1, 0, 0] \mid (x, y) \in A\} \text{ and } B_x = \{y \in h[1, 0, 0] \mid (x, y) \in B\}.$$

As g' is constant along fibers of B_x , we denote by $\tilde{g}' : \pi'(B) \rightarrow h[1, 0, 0]$ the unique definable function such that $g'|_B = \tilde{g}' \circ \pi'|_B$. They refined the above partition, decomposing A as the union $A_1 \cup A_2$ with

$$A_1 := \{(x, y) \in A \mid g'(x, \cdot) \text{ maps } A_x \text{ onto a ball of volume } \mathbb{L}^{-j} \text{ with } j \leq 0\}$$

and

$$A_2 := \{(x, y) \in A \mid g'(x, \cdot) \text{ maps } A_x \text{ onto a ball of volume } \mathbb{L}^{-j} \text{ with } j > 0\}$$

Finally by their construction they consider two definable morphisms

$$r' : S[0, n_Y + n, r] \rightarrow h[1, 0, 0] \text{ and } \eta' : S[0, n_Y + n, r] \rightarrow h[0, 1, 0]$$

such that for any (x, y) in A_2

$$g'(x, y) - r'(x) = \eta(x) \pmod{(t)}.$$

Then, using all these notations they define

$$(1.3) \quad \pi_{S!}^{S[0,1,0]}([\varphi]) := p'_! \left(e^{\tilde{\xi}'} E(\tilde{g}') \pi'_!([1_B \varphi']) + e^{\tilde{\xi}'+\eta} E(r) \pi'_!([1_{A_2} \varphi']) \right)$$

- Integration along $f : Z \rightarrow Y$. Let S be a definable set in Def_k . Let $f : Z \rightarrow Y$ be a morphism in Def_S . Let φ in $I_S C(Z)^{\text{exp}}$ be of the form

$$\varphi = E(g) e^\eta [p : X \rightarrow Z] \varphi_0$$

with $p : X \rightarrow Z$ in RDef_Z , $g : X \rightarrow h[1, 0, 0]$ and $\eta : \rightarrow h[0, 1, 0]$ definable morphisms and φ_0 in $I_S C(Z)$. We denote by $\delta_{f,g,\eta} : X \rightarrow Y[1, 1, 0]$ the morphism sending x to $((f \circ p)(x), g(x), \eta(x))$. We denote by

$$\pi_{Y[0,1,0]}^{Y[1,1,0]} : Y[1, 1, 0] \rightarrow Y[0, 1, 0] \text{ and } \pi_Y^{Y[0,1,0]} : Y[0, 1, 0] \rightarrow Y$$

the projections, and by x and ξ the canonical coordinates on the fibers of $\pi_{Y[0,1,0]}^{Y[1,1,0]}$ and $\pi_{Y^{[0,1,0]}}$. Then, using integration along a residue variable (see Axiom 5) and integration along one valued field variable explained above, Cluckers and Loeser define

$$f_!(\varphi) := \pi_{Y^{[0,1,0]}}^{Y[0,1,0]} \left(\pi_{Y[0,1,0]}^{Y[1,1,0]} (E(x) e^\xi \delta_{f,g,\eta!}(p^* \varphi_0)) \right).$$

1.8. Commutativity of pull-back and push-forward functors. —

Notations 1.17. — Let A, B, C, D some sets. Let $f : A \rightarrow C$ and $g : B \rightarrow D$ be some applications. We denote by $f \times g$ the application from $A \times B$ to $C \times D$ which maps (a, b) to $(f(a), g(b))$. Let $\varphi : A \times B \rightarrow C$ and $\psi : A \times B \rightarrow D$ be some applications. We denote by (φ, ψ) the application from $A \times B$ to $C \times D$ which maps (a, b) to $(\varphi(a, b), \psi(a, b))$.

In [19], the second author introduced a notion of *definable distributions* in Cluckers-Loeser motivic setting. He introduced also a notion of *motivic wave front set* of a definable distribution, which allows him, as in the real setting [13] or in the p-adic setting [12] and [3], to study the pull-back of a distribution by a function which requires the natural following commutativity relation between pull-back and push-forward functors that we prove in section 2.

Lemma 1.18. — *Let S be a definable set in Def_k and $\gamma : W \rightarrow W'$ be a definable morphism in Def_S . Let X be a definable set in Def_S . We denote by π_W the projection from $W \times_S X$ to W and by $\pi_{W'}$ the projection from $W' \times_S X$ to W' . Let φ be a constructible exponential function in $\mathcal{C}(W' \times_S X)^{\text{exp}}$.*

1. *If $[\varphi]$ is $\pi_{W'}$ -integrable then $[(\gamma \times \text{Id}_X)^* \varphi]$ is π_W -integrable. Furthermore, if γ is onto then this implication is an equivalence.*
2. *If $[\varphi]$ satisfies the condition (1) then*

$$(1.4) \quad \pi_{W!} [(\gamma \times \text{Id}_X)^* \varphi] = \gamma^* (\pi_{W'!} [\varphi]).$$

This lemma can be generalized in the following way

Theorem 1.19. — *Let S be a definable set in Def_k and $\gamma : W \rightarrow W'$ be a definable morphism in Def_S . Let $f : X \rightarrow Y$ be a definable morphism in Def_S . We denote by π_W the projection from $W \times_S X$ to W and by $\pi_{W'}$ the projection from $W' \times_S X$ to W' . Let φ be a constructible function in $\mathcal{C}(W' \times_S X)^{\text{exp}}$.*

1. *If $[\varphi]$ is $(\pi_{W'} \times f)$ -integrable then $[(\gamma \times \text{Id}_X)^* \varphi]$ is $(\pi_W \times f)$ -integrable. Furthermore, if γ is onto then this implication is an equivalence.*
2. *If $[\varphi]$ satisfies the condition (1) then*

$$(1.5) \quad (\pi_W \times f)_! [(\gamma \times \text{Id}_X)^* \varphi] = (\gamma \times \text{Id}_Y)^* ((\pi_{W'} \times f)_! [\varphi]).$$

2. Proofs

In subsection 2.1, we state and prove three lemmas allowing us to prove lemma 1.18 and theorem 1.19 in an inductive way following step by step the motivic integration construction in [8] and [9]. In subsection 2.2 we prove theorem 1.19 as a corollary of 1.18. In subsection 2.3 we give a proof of the projection lemma in the case of constructible functions without exponential, then in subsection 2.4 we give the proof of the general case with exponential.

2.1. Some preliminary lemmas. —

2.1.1. Extension lemma. —

Lemma 2.1. — *Let S be a definable set and $\gamma : W \rightarrow W'$ be a definable morphism in Def_S . Let X and Y be two definable sets in Def_S such that $W \times_S X$ and $W' \times_S X$ are respectively definable subassignments of $W \times_S Y$ and $W' \times_S Y$. Let i_W and $i_{W'}$ be the corresponding canonical injections. For any constructible exponential function φ in $\mathcal{C}(W' \times_S X)^{\text{exp}}$, $[\varphi]$ is $i_{W'}$ -integrable and $[(\gamma \times \text{Id}_X)^* \varphi]$ is i_W -integrable. Furthermore, φ satisfies the equality*

$$(2.1) \quad (\gamma \times \text{Id}_Y)^*(i_{W'}[\varphi]) = i_W[(\gamma \times \text{Id}_X)^* \varphi].$$

Proof. — Let φ be an exponential constructible function in $\mathcal{C}(W' \times_S X)^{\text{exp}}$. By Axiom 4 (see in particular [8, §5.5] and [9, §3.5]), the classes $[\varphi]$ and $[(\gamma \times \text{Id}_X)^* \varphi]$ are respectively $i_{W'}$ -integrable and i_W -integrable with the equalities

$$i_{W'}[\varphi] = [i_{W'}(\varphi)] \text{ and } i_W[(\gamma \times \text{Id}_X)^* \varphi] = [i_W((\gamma \times \text{Id}_X)^* \varphi)].$$

The equality (2.1) is then implied by the equality

$$(2.2) \quad (\gamma \times \text{Id}_Y)^*(i_{W'}(\varphi)) = i_W((\gamma \times \text{Id}_X)^* \varphi).$$

which comes from the definition of the pull-back morphism in [8, §5.1, §5.4] and [9, §3.2, §3.4] (see in particular subsection 1.3.3) and the push-forward morphism for an injection [8, §5.5] and [9, §3.5] (see also Axiom 4). Indeed, we write the exponential constructible function φ as

$$\varphi = [p : Z \rightarrow W' \times_S X] E(g) e(\xi) \otimes \varphi_P$$

where Z in $\text{RDef}_{W' \times_S X}$, $g : Z \rightarrow h[1, 0, 0]$ and $\xi : Z \rightarrow h[0, 1, 0]$ are two definable maps and φ_P is a Presburger function in $\mathcal{P}(W' \times_S X)$. Then, the main point is that the fiber product $Z \times_{(W' \times_S X)} (W \times_S X)$ of p and $\gamma \times \text{Id}_X$ is isomorphic to the fiber product of $i_{W'} \circ p$ and $\gamma \times \text{Id}_Y$. Indeed, this is a consequence from a direct computation or from the fact that $(W \times_S X, i_W, \gamma \times \text{Id}_X)$ is isomorphic to the fiber product of $\gamma \times \text{Id}_Y$ and $i_{W'}$ and the result follows from the classical pull-back lemma

in the following diagram

$$\begin{array}{ccc}
 Z \times_{(W' \times_S Y)} (W \times_S Y) \simeq Z \times_{(W' \times_S X)} (W \times_S X) & \xrightarrow{\quad} & Z \\
 \downarrow & \searrow & \downarrow p \\
 W \times_S X & \xrightarrow{\gamma \times Id_X} & W' \times_S X \\
 \downarrow i_W & \searrow & \downarrow i_{W'} \\
 W \times_S Y & \xrightarrow{\gamma \times Id_Y} & W' \times_S Y
 \end{array}$$

□

2.1.2. *Splitting lemma.* —

Lemma 2.2. — *Let S be a definable set and $\gamma : W \rightarrow W'$ be a definable morphism in Def_S . Let X, Y and Z be definable sets in Def_S . We consider the following commutative diagram composed with definable morphisms in Def_S*

$$\begin{array}{ccc}
 W \times_S X & \xrightarrow{\gamma \times Id_X} & W' \times_S X \\
 f \downarrow & & \downarrow f' \\
 W \times_S Y & \xrightarrow{\gamma \times Id_Y} & W' \times_S Y \\
 g \downarrow & & \downarrow g' \\
 W \times_S Z & \xrightarrow{\gamma \times Id_Z} & W' \times_S Z
 \end{array}$$

Assume

- for any constructible exponential function φ in $\mathcal{C}(W' \times_S X)^{exp}$, if $[\varphi]$ is f' -integrable then $(\gamma \times Id_X)^*[\varphi]$ is f -integrable (with equivalence if γ is onto) and in that case

$$(2.3) \quad (\gamma \times Id_X)^*(f'_![\varphi]) = f_![(\gamma \times Id_X)^* \varphi].$$

- for any constructible exponential function ψ in $\mathcal{C}(W' \times_S Y)^{exp}$, if $[\psi]$ is g' -integrable then $(\gamma \times Id_Y)^*[\psi]$ is g -integrable (with equivalence if γ is onto) and in that case

$$(2.4) \quad (\gamma \times Id_Z)^*(g'_![\psi]) = g_![(\gamma \times Id_Y)^* \psi].$$

then, for any constructible exponential function φ in $\mathcal{C}(W' \times_S X)^{exp}$, if $[\varphi]$ is $(g' \circ f')$ -integrable then $(\gamma \times Id_X)^*[\varphi]$ is $(g \circ f)$ -integrable (with equivalence if γ is onto) and in that case

$$(2.5) \quad (\gamma \times Id_Z)^*((g' \circ f')_![\varphi]) = (g \circ f)_![(\gamma \times Id_X)^* \varphi].$$

Proof. — The lemma follows from Fubini axiom (see Axiom 1) and the assumptions.

Indeed, let φ be a constructible exponential function in $\mathcal{C}(W' \times_S X)^{exp}$. By Fubini axiom, $[\varphi]$ is $(g' \circ f')$ -integrable if and only if $[\varphi]$ is f' -integrable and $f'_![\varphi]$ is g' -integrable. Then, using assumptions we observe that if $[\varphi]$ is $(g' \circ f')$ -integrable then $(\gamma \times Id_X)^*[\varphi]$ is f -integrable and $(\gamma \times Id_Y)^*(f'_![\varphi])$ is g -integrable (with equivalence

if γ is onto). By Fubini axiom, $(\gamma \times Id_X)^*[\varphi]$ is f -integrable and $f_!(\gamma \times Id_X)^*[\varphi]$ is g -integrable if and only if $(\gamma \times Id_X)^*[\varphi]$ is $(g \circ f)$ -integrable. Then, equation (2.4) implies the result. Furthermore, we obtain the equality (2.5) by a direct computation using (2.3), (2.4) and Fubini axiom

$$(\gamma \times Id_Z)^*((g' \circ f')_![\varphi]) = (\gamma \times Id_Z)^*(g'_!(f'_![\varphi]))$$

with

$$(\gamma \times Id_Z)^*(g'_!(f'_![\varphi])) = g_!((\gamma \times Id_Y)^*(f'_![\varphi])) = g_!(f_![(\gamma \times Id_X)^*\varphi]),$$

then

$$(\gamma \times Id_Z)^*((g' \circ f')_![\varphi]) = (g \circ f)_![(\gamma \times Id_X)^*\varphi].$$

□

2.1.3. *Reduction lemma.* —

Remark 2.3. — Let S be a definable set and W be a definable set in Def_S . Let m, n and r be some non negative integers. The fiber product $W \times_S S[m, n, r]$ is isomorphic to $W[m, n, r]$ and we identify them in the following.

Lemma 2.4. — Let S be a definable set and $\gamma : W \rightarrow W'$ be a definable morphism in Def_S . Let m, n and r be some non negative integers. Assume lemma 1.18 true in the $\mathcal{C}(W'[m, n, r])^{exp}$ case then it is true in the $\mathcal{C}(W' \times_S X)^{exp}$ case for any definable subassignment X of $S[m, n, r]$.

Proof. — Considering the assumption and the diagram

$$\begin{array}{ccc} W \times_S X & \xrightarrow{\gamma \times Id_X} & W' \times_S X \\ \downarrow i & & \downarrow i' \\ W[m, n, r] & \xrightarrow{\gamma \times Id_{S[m, n, r]}} & W'[m, n, r] \\ \downarrow \pi_W & & \downarrow \pi_{W'} \\ W & \xrightarrow{\gamma} & W' \end{array}$$

the lemma follows from the extension lemma 2.1 and the splitting lemma 2.2. □

2.2. Proof of theorem 1.19. — In this subsection, we assume lemma 1.18 true and we prove theorem 1.19 as a consequence of the extension lemma 2.1 and the splitting lemma 2.2.

Proof. — Let S be a definable set in Def_k and $\gamma : W \rightarrow W'$ be a definable morphism in Def_S . Let $f : X \rightarrow Y$ be a definable morphism in Def_S . We denote by Γ_f the graph of f , by $i_f : X \rightarrow \Gamma_f$ the canonical injection. In the following we will identify canonically $W \times_S \Gamma_f$ and $W' \times_S \Gamma_f$ with $(W \times_S Y) \times_Y X$ and $(W' \times_S Y) \times_Y X$. We consider the following commutative diagram

$$\begin{array}{ccccc}
& W \times_S X & \xrightarrow{\gamma \times Id_X} & W' \times_S X & \\
Id_W \times i_f \swarrow & \downarrow & & \downarrow & \searrow Id_{W'} \times i_f \\
W \times_S \Gamma_f & \xrightarrow{\gamma \times Id_{\Gamma_f}} & & W' \times_S \Gamma_f & \\
\cong & Id_W \times f & & Id_{W'} \times f & \cong \\
(W \times_S Y) \times_Y X & \xrightarrow{\gamma \times Id_Y \times Id_X} & & (W' \times_S Y) \times_Y X & \\
\pi_{W \times_S Y} \searrow & \downarrow & & \downarrow & \swarrow \pi_{W' \times_S Y} \\
& W \times_S Y & \xrightarrow{\gamma \times Id_Y} & W' \times_S Y &
\end{array}$$

The theorem follows from this diagram and the splitting lemma 2.2 (or very similar arguments) whose assumptions are satisfied by application of

- the extension lemma 2.1 for the diagram

$$\begin{array}{ccc}
W \times_S X & \xrightarrow{\gamma \times Id_X} & W' \times_S X \\
Id_W \times i_f \downarrow & & \downarrow Id_{W'} \times i_f \\
W \times_S \Gamma_f & \xrightarrow{\gamma \times Id_{\Gamma_f}} & W' \times_S \Gamma_f
\end{array}$$

- the projection lemma 1.18 (relatively to Y) for the diagram

$$\begin{array}{ccc}
(W \times_S Y) \times_Y X & \xrightarrow{\gamma \times Id_Y \times Id_X} & (W' \times_S Y) \times_Y X \\
\pi_{W \times_S Y} \downarrow & & \downarrow \pi_{W' \times_S Y} \\
W \times_S Y & \xrightarrow{\gamma \times Id_Y} & W' \times_S Y
\end{array}$$

□

2.3. Proof of the projection lemma for constructible functions without exponentials. — We prove in this subsection the projection lemma 1.18 for constructible functions in $C(W' \times_S X)$ -case. The exponential case will be proved in subsection 2.4.

2.3.1. Case $X = S[0, 0, r]$. — In this subsection, we prove lemma 1.18 in the case $X = S[0, 0, r]$. We use remark 2.3 and notations of lemma 1.18. Let φ be a constructible function in $\mathcal{C}(W' \times_S X)$. By proposition 1.8 we write $\varphi = \varphi_{W'} \otimes \varphi_P$, where $\varphi_{W'}$ is a constructible function in $\mathcal{C}(W')$ and φ_P is a Presburger function in $\mathcal{P}(W' \times_S X)$. By Axiom 6, $[\varphi]$ is $\pi_{W'}$ -integrable if and only if φ_P is W' -integrable. The pull-back $(\gamma \times Id_X)^* \varphi$ is equal to $(\varphi_{W'} \circ \gamma) \otimes (\varphi_P \circ (\gamma \times Id_X))$, then by Axiom 6, we deduce that if φ is $\pi_{W'}$ -integrable then $(\gamma \times Id_X)^* \varphi$ is π_W -integrable and furthermore

if γ is onto then this is an equivalence. Under the integrability assumption, $\pi_{W'}![\varphi]$ is equal to the class of $\varphi_{W'} \otimes \pi_{W'}!(\varphi_{\mathcal{P}})$, where $\pi_{W'}!(\varphi_{\mathcal{P}})$ is the unique Presburger function in $\mathcal{P}(W')$ such that for any w' in W' , for any $q > 1$

$$\nu_q((\pi_{W'}!(\varphi_{\mathcal{P}}))(w')) = \sum_{x \in \mathbb{Z}^r} \nu_q(\varphi_{\mathcal{P}}(w', x)).$$

In particular,

$$\gamma^*(\pi_{W'}![\varphi]) = [(\varphi_{W'} \circ \gamma) \otimes (\pi_{W'}!(\varphi_{\mathcal{P}}) \circ \gamma)].$$

As well, $\pi_{W'}![(\gamma \times Id_X)^*\varphi]$ is equal to the class $[(\varphi_{W'} \circ \gamma) \otimes \pi_{W'}!(\varphi_{\mathcal{P}} \circ (\gamma \times Id_X))]$ where $\pi_{W'}!(\varphi_{\mathcal{P}} \circ (\gamma \times Id_X))$ is the unique Presburger function in $\mathcal{P}(W)$ such that for any w in W , for any $q > 1$

$$\nu_q(\pi_{W'}!(\varphi_{\mathcal{P}} \circ (\gamma \times Id_X))(w)) = \sum_{x \in \mathbb{Z}^r} \nu_q((\varphi_{\mathcal{P}} \circ (\gamma \times Id_X))(w, x)).$$

But, for any $q > 1$ and w in W

$$\nu_q(\pi_{W'}!(\varphi_{\mathcal{P}} \circ (\gamma \times Id_X))(w)) = \sum_{x \in \mathbb{Z}^r} \nu_q(\varphi_{\mathcal{P}}(\gamma(w), x)) = \nu_q(\pi_{W'}!\varphi_{\mathcal{P}}(\gamma(w)))$$

then by uniqueness $\pi_{W'}!(\varphi_{\mathcal{P}} \circ (\gamma \times Id_X))$ is equal to $\pi_{W'}!(\varphi_{\mathcal{P}}) \circ \gamma$ and we deduce the equality 2 of lemma 1.18.

2.3.2. Case $X = S[0, n, 0]$. — In this subsection, we prove lemma 1.18 in the case where $X = S[0, n, 0]$. We use remark 2.3 and notations of lemma 1.18. Let φ be a constructible function in $\mathcal{C}(W' \times X)$. By proposition 1.8, φ can be written as $\varphi = [p : Y \rightarrow W' \times X] \otimes \varphi_{\mathcal{P}}$ with $[p : Y \rightarrow W' \times X]$ in $K_0(\text{RDef}_{W' \times X})$ and $\varphi_{\mathcal{P}}$ in $\mathcal{P}(W')$. By Axiom 5, the class $[\varphi]$ is $\pi_{W'}$ -integrable with

$$\pi_{W'}![\varphi] = [\pi_{W'} \circ p : Y \rightarrow W'] \otimes \varphi_{\mathcal{P}}$$

and $\gamma^*(\pi_{W'}![\varphi]) = [Y \times_{W'} W \rightarrow W] \otimes (\varphi_{\mathcal{P}} \circ \gamma)$. As well, $(\gamma \times Id_X)^*\varphi$ is equal to $[Y \times_{W' \times X} (W \times X) \rightarrow W \times X] \otimes (\varphi_{\mathcal{P}} \circ \gamma)$ and is π_W -integrable. As $(W \times X, \pi_W, \gamma \times Id_X)$ is isomorphic to the fiber product $W \times_{W'} (W' \times X)$ of γ and $\pi_{W'}$, we deduce similarly to the case $X = S[0, 0, r]$, by the classical pull-back theorem, that $Y \times_{W' \times X} (W \times X)$ and $Y \times_{W'} (W \times X)$ are isomorphic, which induces the equality 2 of lemma 1.18.

2.3.3. Case $X = S[0, n, r]$. — The lemma 1.18 in the case $X = S[0, n, r]$ follows immediately from the splitting lemma 2.2 applied to the case $X = S[0, 0, r]$ in 2.3.1 and the case $X = S[0, n, 0]$ in 2.3.2.

2.3.4. Case where X is a 0-cell and $\varphi = 1_{W' \times_S X}$. — Let X be a 0-cell of base Y in Def_S with center $c : Y \rightarrow h[1, 0, 0]$

$$X = \{(y, z) \in Y[1, 0, 0] \mid z = c(y)\}.$$

We denote by π the projection from X to Y which is an isomorphism. The product $W \times_S X$ is also a 0-cell of base $W \times_S Y$ and center $c_W = c \circ \pi_Y$ where π_Y is the projection from $W \times_S Y$ to Y . As well the product $W' \times_S X$ is a 0-cell of base

$W' \times_S Y$, center $c_{W'} = c \circ \pi'_Y$ where π'_Y is the projection from $W' \times_S Y$ to Y . We prove lemma 1.18 for the constructible function $\varphi = 1_{W' \times_S X}$ and the diagram

$$\begin{array}{ccc} W \times_S X & \xrightarrow{\gamma \times Id_X} & W' \times_S X \\ p_W = Id_W \times \pi \downarrow & & \downarrow p_{W'} = Id_{W'} \times \pi \\ W \times_S Y & \xrightarrow{\gamma \times Id_Y} & W' \times_S Y \end{array}.$$

By definition of the pull-back of a Presburger function we have

$$(\gamma \times Id_X)^* \varphi = \varphi \circ (\gamma \times Id_X) = 1_{W \times_S X}.$$

Then, by the change variable formula [8, Proposition 13.2.1] (see theorem 1.14), the class $[\varphi]$ is $p_{W'}$ -integrable and the class $[(\gamma \times Id_X)^* \varphi]$ is also p_W -integrable with

$$p_{W'}![\varphi] = \mathbb{L}^{\text{ord Jac } p_{W'} \circ p_{W'}^{-1}} \text{ and } p_W![(\gamma \times Id_X)^* \varphi] = \mathbb{L}^{\text{ord Jac } p_W \circ p_W^{-1}}.$$

Remark that by definition of p_W and $p_{W'}$, the order $\text{ord Jac } p_{W'} \circ p_{W'}^{-1}$ is equal to the order $\text{ord Jac } \pi \circ (\pi^{-1} \circ \pi'_Y)$. Similarly, the order $\text{ord Jac } p_W \circ p_W^{-1}$ is equal to the order $\text{ord Jac } \pi \circ (\pi^{-1} \circ \pi_Y)$. Then, the equality 2 of lemma 1.18 follows from the equality $\pi_Y = \pi'_Y \circ (\gamma \times Id_Y)$.

2.3.5. Case where X is 1-cell and $\varphi = 1_{W' \times_S X}$. — Let X be a 1-cell of base Y in Def_S with center $c : Y \rightarrow h[1, 0, 0]$ and data $\alpha : Y \rightarrow \mathbb{Z}$ and $\xi : Y \rightarrow h[0, 1, 0]$

$$X = \{(y, z) \in Y[1, 0, 0] \mid \text{ord}(z - c(y)) = \alpha(y), \overline{\alpha c}(z - c(y)) = \xi(y)\}.$$

We denote by π the projection from X to Y . The product $W \times_S X$ is still a 1-cell with base $W \times_S Y$, center $c_W = c \circ \pi_Y$ and data $\alpha_W = \alpha \circ \pi_Y$ and $\xi_W = \xi \circ \pi_Y$, with π_Y the projection from $W \times_S Y$ to Y . As well, the product $W' \times_S X$ is still a 1-cell with base $W' \times_S Y$, center $c_{W'} = c \circ \pi'_Y$ and data $\alpha_{W'} = \alpha \circ \pi'_Y$ and $\xi_{W'} = \xi \circ \pi'_Y$ with π'_Y the projection from $W' \times_S Y$ to Y . We prove lemma 1.18 for the constructible function $\varphi = 1_{W' \times_S X}$ and the diagram

$$\begin{array}{ccc} W \times_S X & \xrightarrow{\gamma \times Id_X} & W' \times_S X \\ p_W = Id_W \times \pi \downarrow & & \downarrow p_{W'} = Id_{W'} \times \pi \\ W \times_S Y & \xrightarrow{\gamma \times Id_Y} & W' \times_S Y \end{array}.$$

By definition of the pull-back of a Presburger function we have

$$(\gamma \times Id_X)^* \varphi = \varphi \circ (\gamma \times Id_X) = 1_{W \times_S X}.$$

Then, by Axiom 8, (see axiom (A7) of [8, theorem 10.1.1, Proposition 13.2.1]), the class $[\varphi]$ is $p_{W'}$ -integrable and the class $[(\gamma \times Id_X)^* \varphi]$ is also p_W -integrable with

$$p_{W'}![\varphi] = \mathbb{L}^{-\alpha_{W'} - 1}[1_{W' \times_S Y}] \text{ and } p_W![(\gamma \times Id_X)^* \varphi] = \mathbb{L}^{-\alpha_W - 1}[1_{W \times_S Y}].$$

Then, the equality 2 of lemma 1.18 follows from the equality $\pi_Y = \pi'_Y \circ (\gamma \times Id_Y)$.

2.3.6. *Proof of the projection lemma 1.18.* — Using the reduction lemma 2.4, it is enough to consider the case where X is equal to the definable set $S[m, n, r]$ for m, n and r some non-negative integers. We use remark 2.3 and notations of lemma 1.18. The projection lemma 1.18 is proved by induction on m . The base case $m = 0$ is ever considered in paragraph 2.3.3 and using the splitting lemma 2.2, it is enough to prove the projection lemma 1.18 for the diagram

$$(2.6) \quad \begin{array}{ccc} W \times_S X & \xrightarrow{\gamma \times Id_X} & W' \times_S X \\ \pi_{\overline{W}} \downarrow & & \downarrow \pi_{\overline{W'}} \\ \overline{W} & \xrightarrow{\gamma \times Id_{S[m-1, n, r]}} & \overline{W'} \end{array}$$

with $m \geq 1$, $\overline{W'} = W'[m-1, n, r]$, $\overline{W} = W[m-1, n, r]$ and where $\pi_{\overline{W}}$ and $\pi_{\overline{W'}}$ are the canonical projections.

We prove now the case of diagram (2.6) using the cell decomposition theorem 1.10 and the specific cases of 0-cell and 1-cell in paragraphs 2.3.4 and 2.3.5. As $m \geq 1$, we consider $W' \times_S X$ as the product $\overline{W'}[1, 0, 0]$ and $W \times_S X$ as the product $\overline{W}[1, 0, 0]$. Let φ be a constructible function in $\mathcal{C}(W' \times_S X)$. By theorem 1.10, we consider a cell decomposition $((W' \times_S X)_i)_{i \in I}$ of $W' \times_S X$ adapted to φ , with for any i in I , a presentation $(Z_{C'_i}, \lambda'_i)$ of the cell $(W' \times_S X)_i$

$$(2.7) \quad \begin{array}{ccc} (W' \times_S X)_i & \xrightarrow{\lambda'_i} & Z_{C'_i} \subset \overline{W'}[1, n_i, r_i] \\ \pi_{\overline{W'}} \downarrow & & \downarrow p'_i \\ \overline{W'} & \xleftarrow{\pi'_i} & C'_i \subset \overline{W'}[0, n_i, r_i] \end{array}$$

where the diagram is commutative, p'_i and π'_i are the canonical projections and

- if $(W' \times_S X)_i$ is a 0-cell, then the integer r_i is equal to 0 and the isomorphism λ'_i is equal to $Id_{(W' \times_S X)_i} \times \eta_i$ where $\eta_i : (W' \times_S X)_i \rightarrow h[0, n_i, 0]$ is a definable morphism. The jacobian order of the isomorphism λ'_i is 0.
- if $(W' \times_S X)_i$ is a 1-cell, then the isomorphism λ'_i is equal to $Id_{(W' \times_S X)_i} \times \eta_i \times l_i$ where $\eta_i : (W' \times_S X)_i \rightarrow h[0, n_i, 0]$ and $l_i : (W' \times_S X)_i \rightarrow h[0, 0, r_i]$ are two definable morphisms. The jacobian order of the isomorphism λ'_i is 0.

Taking a refinement of the cell decomposition, we may assume that for any i in I , $\varphi|_{(W' \times_S X)_i}$ is either zero or has the same K -dimension as $(W' \times_S X)_i$ (see 1.4.2 and also proof [8, §11.1]). Furthermore, for any i in I , there is a constructible function ψ'_i in $\mathcal{C}(C'_i)$ such that

$$(2.8) \quad \varphi|_{(W' \times_S X)_i} = (\lambda'_i)^* (p'_i)^* \psi'_i.$$

Taking the pull-back by the definable morphism $\gamma \times Id_X$, we deduce a cell decomposition $((W \times_S X)_i)_{i \in I}$ of $W \times_S X$ adapted to $(\gamma \times Id_X)^* \varphi$. More precisely, for any i in I , we define the cell $(W \times_S X)_i$ as the definable set $(\gamma \times Id_X)^{-1}(W' \times_S X)_i$. This

cell has a presentation (Z_{C_i}, λ_i) where

$$C_i = \{(w, x, \eta, l) \in W[m-1, n+n_i, r+r_i] \mid (\gamma(\omega), x, \eta, l) \in C'_i\},$$

and

- if $Z_{C'_i}$ is a 1-cell with center c'_i and data α'_i and ξ'_i , then Z_{C_i} is the 1-cell in $\overline{W}[1, n_i, r_i]$ with center $c_i = c'_i \circ (\gamma \times Id_{S[m-1, n+n_i, r_i]})$ and data

$$\alpha_i = \alpha'_i \circ (\gamma \times Id_{S[m-1, n+n_i, r_i]}) \text{ and } \xi_i = \xi'_i \circ (\gamma \times Id_{S[m-1, n+n_i, r_i]})$$

and the presentation morphism $\lambda_i : (W \times_S X)_i \rightarrow Z_{C_i}$ is

$$\lambda_i = Id_{(W \times_S X)_i} \times (\eta_i \circ (\gamma \times Id_X)) \times (l_i \circ (\gamma \times Id_X)).$$

- if $Z_{C'_i}$ is a 0-cell with center c'_i , then Z_{C_i} is the 0-cell in $\overline{W}[1, n_i, 0]$ with center

$$c_i = c'_i \circ (\gamma \times Id_{S[m-1, n+n_i, r_i]})$$

and the presentation morphism $\lambda_i : (W \times_S X)_i \rightarrow Z_{C_i}$ is

$$\lambda_i = Id_{(W \times_S X)_i} \times (\eta_i \circ (\gamma \times Id_X)).$$

For any i in I , we consider the constructible function

$$\psi_i := (\gamma \times Id_{S[m-1, n+n_i, r+r_i]})^*(\psi'_i)$$

and we have the equalities

$$(\gamma \times Id_X)^* 1_{(W' \times_S X)_i} = 1_{(W \times_S X)_i}$$

$$(2.9) \quad (\gamma \times Id_X)^* \varphi \cdot 1_{(W \times_S X)_i} = (\gamma \times Id_X)^* (\lambda_i'^* p_i'^* \psi'_i) = \lambda_i^* p_i^* \psi_i$$

thanks to the equality

$$p_i' \circ \lambda_i' \circ (\gamma \times Id_X) = (\gamma \times Id_{S[m-1, n+n_i, r+r_i]}) \circ p_i \circ \lambda_i$$

with p_i the canonical projection from $\overline{W}[1, n_i, r_i]$ to $\overline{W}[0, n_i, r_i]$.

By the additivity axiom (Axiom 2), as the cells $(W' \times_S X)_i$ and $(W \times_S X)_i$ are disjoint:

- the class $[\varphi]$ is $\pi_{\overline{W}'}^*$ -integrable if and only if for any i in I , $\varphi 1_{(W' \times_S X)_i}$ is $\pi_{\overline{W}'}^*$ -integrable, if and only if for any i in I , the class $[\psi'_i] p_i'^* [1_{Z_{C'_i}}]$ is $\pi_i'^*$ -integrable (applying to equation 2.8 and diagram 2.7, Fubini axiom (Axiom 1), change variable formula (theorem 1.14) and projection axiom (Axiom 3)), and in that case

$$(2.10) \quad \pi_{\overline{W}'}^* [\varphi \cdot 1_{(W' \times_S X)_i}] = \pi_i'^* ([\psi'_i] p_i'^* [1_{Z_{C'_i}}]).$$

- the class $[(\gamma \times Id_X)^* \varphi]$ is $\pi_{\overline{W}}^*$ -integrable if and only if for all i in I , the class of $(\gamma \times Id_X)^* \varphi \cdot 1_{(W \times_S X)_i}$ is $\pi_{\overline{W}}^*$ -integrable, if and only if for any i in I , $[\psi_i] p_i^* [1_{Z_{C_i}}]$ is π_i^* -integrable (by Fubini axiom (Axiom 1), change variable formula and projection axiom (Axiom 3) applied to equation 2.9), and in that case

$$(2.11) \quad \pi_{\overline{W}}^* [(\gamma \times Id_X)^* \varphi \cdot 1_{(W \times_S X)_i}] = \pi_i^* ([\psi_i] p_i^* [1_{Z_{C_i}}]).$$

But, by construction, $1_{Z_{C_i}} = (\gamma \times Id_{S[m, n+n_i, r+r_i]})^* 1_{Z_{C'_i}}$ and by the 0-cell and 1-cell case in sections 2.3.4 and 2.3.5 we have the identity

$$(\gamma \times Id_{S[m-1, n+n_i, r+r_i]})^* p'_{i!}[1_{Z_{C_i}}] = p_{i!}[1_{Z_{C_i}}]$$

which implies the equality

$$(2.12) \quad [\psi_i] p_{i!}[1_{Z_{C_i}}] = (\gamma \times Id_{S[m-1, n+n_i, r+r_i]})^* ([\psi'_i] p'_{i!}[1_{Z_{C'_i}}]).$$

Then, for any i in I , using subsection 2.3.3 for the diagram

$$\begin{array}{ccc} C_i \subset \overline{W}[0, n_i, r_i] & \xrightarrow{\gamma \times Id_{S[m-1, n+n_i, r+r_i]}} & C'_i \subset \overline{W'}[0, n_i, r_i] \\ \pi_i \downarrow & & \downarrow \pi'_i \\ \overline{W} & \xrightarrow{\gamma \times Id_{S[m-1, n, r]}} & \overline{W'} \end{array}$$

with π_i and π'_i the canonical projections, we deduce that if the class $[\psi'_i] p'_{i!}[1_{Z_{C'_i}}]$ is $\pi_{i'}$ -integrable then $(\gamma \times Id_{S[m-1, n+n_i, r+r_i]})^* ([\psi'_i] p'_{i!}[1_{Z_{C'_i}}])$ equal to $[\psi_i] p_{i!}[1_{Z_{C_i}}]$ is also π_i -integrable and if γ is onto, this is an equivalence. In that case we have the equality

$$(2.13) \quad \begin{aligned} & \pi_{i!}((\gamma \times Id_{S[m-1, n+n_i, r+r_i]})^* ([\psi'_i] p'_{i!}[1_{Z_{C'_i}}])) \\ &= \\ & (\gamma \times Id_{S[m-1, n, r]})^* (\pi_{i'!}([\psi'_i] p'_{i!}[1_{Z_{C'_i}}])). \end{aligned}$$

Then, we can conclude that for any i in I , if the class $[\varphi.1_{(W' \times_S X)_i}]$ is $\pi_{\overline{W'}}$ -integrable then the class $[(\gamma \times Id_X)^* \varphi.1_{(W \times_S X)_i}]$ is $\pi_{\overline{W}}$ -integrable and this is an equivalence in the case of γ onto. In that case by equations (2.10), (2.11), (2.12) and (2.13), we get for any i in I

$$\pi_{\overline{W}}[(\gamma \times Id_X)^* (\varphi.1_{(W' \times_S X)_i})] = (\gamma \times Id_{S[m-1, n, r]})^* \pi_{W'!}([\varphi.1_{(W' \times_S X)_i}])$$

By additivity Axiom 2 and summation we conclude that if $[\varphi]$ is $\pi_{\overline{W'}}$ -integrable then the class $[(\gamma \times Id_X)^* \varphi]$ is $\pi_{\overline{W}}$ -integrable and this is an equivalence in the case of γ onto, and in that case

$$\pi_{\overline{W}}[(\gamma \times Id_X)^* \varphi] = (\gamma \times Id_{S[m-1, n, r]})^* \pi_{W'!}([\varphi]).$$

Remark 2.5. — This achieves the proof of lemma 1.18 in the $C(W \times_S X)$ -case, which implies that theorem 1.19 is also true in this setting. We will use both of them in the proof of lemma 1.18 in the $C(W \times_S X)^{exp}$ -context.

2.4. $C(W' \times_S X)^{exp}$ -case. — .

Let X be a definable in Def_S , by the reduction lemma 2.4 we can assume that $X = S[m, n, r]$. Let $\gamma : W \rightarrow W'$ be a morphism in Def_S . Let $\pi_{W'} : W' \times_S X \rightarrow W'$ and $\pi_W : W \times_S X \rightarrow X$ be the canonical projections. We consider the diagram

$$\begin{array}{ccc} W \times_S X & \xrightarrow{\gamma \times Id_X} & W' \times_S X \\ \pi_W \downarrow & & \downarrow \pi_{W'} \\ W & \xrightarrow{\gamma} & W' \end{array}$$

Let φ be a constructible function in $\mathcal{C}(W' \times_S X)^{exp}$, then by definition φ can be written as

$$(2.14) \quad \varphi = E(g')e(\eta')[p' : Y' \rightarrow W' \times_S X] \otimes \varphi_0$$

where $g' : W' \times_S X \rightarrow h[1, 0, 0]$, $\eta' : W' \times_S X \rightarrow h[0, 1, 0]$, $p' : Y' \rightarrow W' \times_S X$ are definable functions with Y' a definable subset of some $(W' \times_S X)[0, n, 0]$ and φ_0 is a constructible function in $\mathcal{C}(W' \times_S X)$. In particular the pull-back by $\gamma \times Id_X$ of φ is

$$(2.15) \quad (\gamma \times Id_X)^* \varphi = E(g)e(\eta)[p : Y \rightarrow W \times_S X] \otimes (\gamma \times Id_X)^* \varphi_0$$

where $p : Y \rightarrow W \times X$, g and η are the pull-back of p' , g' and η' by $\gamma \times Id_X$.

Furthermore, by definition (see subsection 1.7), $[\varphi]$ is $\pi_{W'}$ -integrable if and only if $[\varphi_0]$ is $\pi_{W'}$ -integrable and in that case, it follows from theorem 4.1.1 in [9] and the uniqueness part of its proof §6.3 that

$$(2.16) \quad \pi_{W'}! [\varphi] = \left(\pi_{W'}^{W'[0,1,0]} \right)_! \left(\pi_{W'}^{W'[1,1,0]} \right)_! (E(x)e(\xi) \otimes \delta'_!(p'^*[\varphi_0])).$$

where we use the diagram

$$\begin{array}{ccc} Y' & \xrightarrow{\delta'} & W'[1, 1, 0] \\ \downarrow p' & & \downarrow \pi_{W'}^{W'[1,1,0]} \\ & & W'[0, 1, 0] \\ & & \downarrow \pi_{W'}^{W'[0,1,0]} \\ W \times_S X & \xrightarrow{\pi_{W'}} & W' \end{array}$$

where x and ξ are coordinate on the vector bundle $W'[1, 1, 0]$ and δ' is the definable function from Y' to $W'[1, 1, 0]$ equal to $(\pi_{W'} \circ p', g' \circ p', \eta' \circ p')$. In particular, we deduce from the case without exponential in section 2.3, that condition 1 of lemma 1.18 is satisfied. We have the following commutative diagram

$$\begin{array}{ccccc}
W[1, 1, 0] & \xrightarrow{\gamma \times Id_{S[1, 1, 0]}} & W'[1, 1, 0] & & \\
\downarrow \pi_W^{W[1, 1, 0]} & \swarrow \delta & \nearrow \delta' & & \downarrow \pi_{W'}^{W'[1, 1, 0]} \\
& Y & \xrightarrow{\gamma \times Id_{X[0, n, 0]}} & Y' & \\
& \downarrow p & & \downarrow p' & \\
W \times X & \xrightarrow{\gamma \times Id_X} & W' \times X & & \\
\downarrow \pi_W & \searrow & \swarrow & & \downarrow \pi_{W'} \\
W[0, 1, 0] & \xrightarrow{\gamma \times Id_{S[0, 1, 0]}} & W'[0, 1, 0] & & \\
\downarrow \pi_W^{W[0, 1, 0]} & & \downarrow \pi_{W'}^{W'[0, 1, 0]} & & \\
W & \xrightarrow{\gamma} & W' & &
\end{array}$$

Using the splitting lemma (lemma 2.2) to prove the equality 2 of lemma 1.18

$$\gamma^*(\pi_{W'}![\varphi]) = \pi_{W'}!((\gamma \times Id_X)^* \varphi)$$

it is enough to prove the result for the special cases

$$(2.17) \quad \begin{array}{ccc}
W[0, 1, 0] & \xrightarrow{\gamma \times Id_{S[0, 1, 0]}} & W'[0, 1, 0] \\
\downarrow \pi_W^{W[0, 1, 0]} & & \downarrow \pi_{W'}^{W'[0, 1, 0]} \\
W & \xrightarrow{\gamma} & W'
\end{array}$$

$$(2.18) \quad \begin{array}{ccc}
W[1, 1, 0] & \xrightarrow{\gamma \times Id_{S[1, 1, 0]}} & W'[1, 1, 0] \\
\downarrow \pi_W^{W[1, 1, 0]} & & \downarrow \pi_{W'}^{W'[1, 1, 0]} \\
W[0, 1, 0] & \xrightarrow{\gamma} & W'[0, 1, 0]
\end{array}$$

Indeed, by equation (2.16) we have

$$\gamma^*(\pi_{W'}![\varphi]) = \gamma^* \left(\left(\pi_{W'}^{W'[0, 1, 0]} \right)_! \left(\pi_{W'}^{W'[1, 1, 0]} \right)_! [E(x)e(\xi) \otimes \delta'_!(p'^*[\varphi_0])] \right).$$

Then, it follows from lemma 1.18 in the case of diagram (2.17) that

$$\gamma^*(\pi_{W'}![\varphi]) = \left(\pi_W^{W[0, 1, 0]} \right)_! \left[(\gamma \times Id_{S[0, 1, 0]})^* \left(\left(\pi_{W'}^{W'[1, 1, 0]} \right)_! [E(x)e(\xi) \otimes \delta'_!(p'^*[\varphi_0])] \right) \right],$$

then, it follows from lemma 1.18 in the case of diagram (2.18) that

$$\gamma^*(\pi_{W'!}[\varphi]) = \left(\pi_{W!}^{W[0,1,0]} \right)_! \left[\left(\pi_{W[0,1,0]}^{W[1,1,0]} \right)_! (\gamma \times Id_{S[1,1,0]})^* ([E(x)e(\xi) \otimes \delta'_!(p'^*[\varphi_0])]) \right],$$

and finally, as x and ξ are coordinates in the bundle $W'[1,1,0]$ and are independent from W' we have

$$\gamma^*(\pi_{W'!}[\varphi]) = \left(\pi_{W!}^{W[0,1,0]} \right)_! \left(\pi_{W[0,1,0]}^{W[1,1,0]} \right)_! [E(x)e(\xi) \otimes (\gamma \times Id_{S[1,1,0]})^* [\delta'_!(p'^*[\varphi_0])]].$$

Applying theorem 1.19 in the case without exponential (see remark 2.5) we obtain the equality

$$(\gamma \times Id_{S[1,1,0]})^* (\delta'_!(p'^*[\varphi_0])) = \delta'_! ((\gamma \times Id_{X[0,n,0]})^* (p'^*[\varphi_0]))$$

and by commutativity of the diagram we have

$$(\gamma \times Id_{S[1,1,0]})^* (\delta'_!(p'^*[\varphi_0])) = \delta'_! (p^*(\gamma \times Id_X)^*[\varphi_0])$$

and we can finally conclude

$$\gamma^*(\pi_{W'!}\varphi) = \left(\pi_{W!}^{W[0,1,0]} \right)_! \left(\pi_{W[0,1,0]}^{W[1,1,0]} \right)_! (E(x)e(\xi) \otimes \delta'_! [p^*(\gamma \times Id_X)^*\varphi_0]).$$

namely by equation (2.14)

$$\gamma^*(\pi_{W'!}[\varphi]) = \pi_{W!}([\gamma \times Id_X]^*\varphi).$$

We conclude now by the proof of the projection lemma 1.18 in the cases of diagrams (2.17) and (2.18).

- As in the paragraph 2.3.2 the case of diagram (2.17) follows by the definition of the push-forward in the residue variables in [9, §3.6].
- Taking pull-back of a cell decomposition as in the proof of the case without exponential (see section 2.3), the case of diagram 2.18 follows directly from the construction in [9, §5.1, lemma 5.1.1] (sketched in example 1.16) where the set of parameters is $W'[0,1,0]$ for the integration along $\pi_{W'[0,1,0]}^{W'[1,1,0]}$ and $W[0,1,0]$ for the integration along $\pi_{W[0,1,0]}^{W[1,1,0]}$. Without giving the details, the constructions (sketched in example 1.16) of $\pi_{W'[1,1,0]}^{W'[0,1,0]}([\varphi])$ and $\pi_{W[1,1,0]}^{W[0,1,0]}((\gamma \times Id_{S[0,1,0]})^*[\varphi])$ for a $\pi_{W'[1,1,0]}^{W'[0,1,0]}$ -integrable-constructible function φ in $\mathcal{C}(W'[1,1,0])^{\text{exp}}$ are step by step compatible with the base change from $W'[0,1,0]$ to $W[0,1,0]$, and the projection formula

$$(\gamma \times Id_{W'[0,1,0]})^* \left(\pi_{W'[1,1,0]}^{W'[0,1,0]}([\varphi]) \right) = \pi_{W[1,1,0]}^{W[0,1,0]}((\gamma \times Id_{S[0,1,0]})^*[\varphi])$$

follows from equation 1.3 using above ideas and the case without exponential.

Acknowledgments. — We are very grateful to Raf Cluckers, Immanuel Halupczok and François Loeser for inspiring discussions. The first author was supported by the European Research Council under the European Community’s Seventh Framework Programme (FP7/2007-2013) with ERC Grant Agreement no. 615722 MOTMEL-SUM, by the Labex CEMPI (ANR-11-LABX-0007-01). The first author would also like to thank VIASM in Hanoi for the hospitality and the great environment for work, during a three months visit in the summer 2018. The second author is partially supported by ANR-15-CE40-0008 (Defigeo) and by the ERC Advanced Grant NMNAG.

References

- [1] Raf Cluckers, Julia Gordon, and Immanuel Halupczok. Integrability of oscillatory functions on local fields: transfer principles. *Duke Math. J.*, 163(8):1549–1600, 2014.
- [2] Raf Cluckers, Thomas Hales, and François Loeser. Transfer principle for the fundamental lemma. In *On the stabilization of the trace formula*, volume 1 of *Stab. Trace Formula Shimura Var. Arith. Appl.*, pages 309–347. Int. Press, Somerville, MA, 2011.
- [3] Raf Cluckers, Immanuel Halupczok, François Loeser, and Michel Raibaut. Distributions and wave front sets in the uniform non-arhimedean setting. *Transaction of the London Mathematical Society*, 5(1):97–131, 2018.
- [4] Raf Cluckers and François Loeser. Fonctions constructibles et intégration motivique. I. *C. R. Math. Acad. Sci. Paris*, 339(6):411–416, 2004.
- [5] Raf Cluckers and François Loeser. Fonctions constructibles et intégration motivique. II. *C. R. Math. Acad. Sci. Paris*, 339(7):487–492, 2004.
- [6] Raf Cluckers and François Loeser. Ax-Kochen-Eršov theorems for p -adic integrals and motivic integration. In *Geometric methods in algebra and number theory*, volume 235 of *Progr. Math.*, pages 109–137. Birkhäuser Boston, Boston, MA, 2005.
- [7] Raf Cluckers and François Loeser. Fonctions constructibles exponentielles, transformation de Fourier motivique et principe de transfert. *C. R. Math. Acad. Sci. Paris*, 341(12):741–746, 2005.
- [8] Raf Cluckers and François Loeser. Constructible motivic functions and motivic integration. *Invent. Math.*, 173(1):23–121, 2008.
- [9] Raf Cluckers and François Loeser. Constructible exponential functions, motivic Fourier transform and transfer principle. *Ann. of Math. (2)*, 171(2):1011–1065, 2010.
- [10] J. Denef. The rationality of the Poincaré series associated to the p -adic points on a variety. *Invent. Math.*, 77(1):1–23, 1984.
- [11] Julia Gordon and Yoav Yaffe. An overview of arithmetic motivic integration. In *Ottawa lectures on admissible representations of reductive p -adic groups*, volume 26 of *Fields Inst. Monogr.*, pages 113–149. Amer. Math. Soc., Providence, RI, 2009.
- [12] D. B. Heifetz. p -adic oscillatory integrals and wave front sets. *Pacific J. Math.*, 116(2):285–305, 1985.
- [13] Lars Hörmander. The analysis of linear partial differential operators. I: Distribution theory and Fourier analysis. Grundlehren der Mathematischen Wissenschaften, 256. Berlin Heidelberg-New York - Tokyo: Springer-Verlag. IX, 391 p. DM 98.00; \$ 39.20 (1983)., 1983.
- [14] Ehud Hrushovski and David Kazhdan. Integration in valued fields. In *Algebraic geometry and number theory*, volume 253 of *Progr. Math.*, pages 261–405. Birkhäuser Boston, Boston, MA, 2006.

- [15] Jun-ichi Igusa. *An introduction to the theory of local zeta functions*, volume 14 of *AMS/IP Studies in Advanced Mathematics*. American Mathematical Society, Providence, RI; International Press, Cambridge, MA, 2000.
- [16] Kontsevich. Lecture at Orsay. Décembre 7, 1995.
- [17] Johan Pas. Uniform p -adic cell decomposition and local zeta functions. *J. Reine Angew. Math.*, 399:137–172, 1989.
- [18] M. Presburger. Über die Vollständigkeit eines gewissen Systems der Arithmetik.... . In *Comptes-rendus du I Congrès des Mathématiciens des Pays Slaves*, pp. 92–101. Warsaw (1929) .
- [19] Michel Raibaut. Motivic wave front sets. *arXiv:1810.10567*.
- [20] Lou van den Dries. Dimension of definable sets, algebraic boundedness and Henselian fields. *Ann. Pure Appl. Logic*, 45(2):189–209, 1989. Stability in model theory, II (Trento, 1987).

November 19, 2018

JORGE CELY, Bogotá, Colombia • *E-mail* : celyje@gmail.com

MICHEL RAIBAUT, Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, LAMA
Le Bourget-du-lac, 73376, France. • *E-mail* : Michel.Raibaut@univ-smb.fr
Url : raibautm.perso.math.cnrs.fr/site/MichelRaibaut.html