# BASES FOR FUNCTIONS BEYOND THE FIRST BAIRE CLASS

RAPHAËL CARROY AND BENJAMIN D. MILLER

ABSTRACT. We provide a finite basis for the class of Borel functions that are not in the first Baire class, as well as the class of Borel functions that are not  $\sigma$ -continuous with closed witnesses.

#### INTRODUCTION

A topological space is *analytic* if it is a continuous image of a closed subset of  $\mathbb{N}^{\mathbb{N}}$ . A subset of a topological space is *Borel* if it is in the  $\sigma$ -algebra generated by open sets,  $F_{\sigma}$  if it is a union of countably-many closed sets, and  $G_{\delta}$  if it is an intersection of countably-many open sets.

Suppose that X and Y are topological spaces. Given a family  $\Gamma$  of subsets of X, a function  $\phi: X \to Y$  is  $\Gamma$ -measurable if  $\phi^{-1}(V) \in \Gamma$  for every open set  $V \subseteq Y$ . A function is Borel if it is Borel-measurable, Baire class one if it is  $F_{\sigma}$ -measurable, and  $\sigma$ -continuous with closed witnesses if its domain is the union of countably-many closed sets on which it is continuous. A result of Jayne-Rogers (see [JR82, Theorem 1]) ensures that a function from an analytic metric space to a separable metric space has this property if and only if it is  $G_{\delta}$ -measurable.

A quasi-order on a set Z is a reflexive transitive binary relation  $\leq$  on Z. A set  $B \subseteq Z$  is a basis under  $\leq$  for Z if  $\forall z \in Z \exists b \in B \ b \leq z$ .

A closed continuous embedding of  $\phi: X \to Y$  into  $\phi': X' \to Y'$ consists of a pair of closed continuous embeddings  $\pi_X: X \to X'$  and  $\pi_Y: \overline{\phi(X)} \to \overline{\phi'(X')}$  such that  $\phi' \circ \pi_X = \pi_Y \circ \phi$ . Note that the existence of such a pair depends not only on the graphs of the functions  $\phi$  and  $\phi'$ , but on Y as well, since different choices of  $Y \supseteq \phi(X)$  can lead to different values of  $\overline{\phi(X)}$ . Here we establish the following results.

**Theorem 1.** There is a twenty-four-element basis under closed continuous embeddability for the class of non-Baire-class-one Borel functions between analytic metric spaces.

<sup>2010</sup> Mathematics Subject Classification. Primary 03E15, 26A21, 28A05, 54H05. Key words and phrases. Baire class, basis, embedding, sigma-continuous.

The authors were supported in part by FWF Grants P28153 and P29999.

#### R. CARROY AND B.D. MILLER

**Theorem 2.** There is a twenty-seven-element basis under closed continuous embeddability for the class of  $non-\sigma$ -continuous-with-closed-witnesses Borel functions between analytic metric spaces.

In §1, we discuss the compactification  $\mathbb{N}_*^{\leq \mathbb{N}}$  of  $\mathbb{N}^{\leq \mathbb{N}}$  underlying our arguments, as well as the corresponding compactification  $\mathbb{N}_*^{\mathbb{N}}$  of  $\mathbb{N}^{\mathbb{N}}$ . In §2, we discuss the endomorphisms of  $\mathbb{N}^{<\mathbb{N}}$  underlying our arguments. In §3, we provide a three-element basis for the class of Baire measurable functions from  $\mathbb{N}^{\mathbb{N}}$  to separable metric spaces. In §4, we provide a three-element basis for the class of non- $\sigma$ -continuous-withclosed-witnesses Baire-class-one functions from analytic metric spaces to separable metric spaces. In §5, we provide an eight-element basis for the class of all functions from  $\mathbb{N}_*^{\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}}$  to analytic metric spaces. And in §6, we establish Theorems 1 and 2.

## 1. A compactification of $\mathbb{N}^{\leq \mathbb{N}}$

We use  $s \sim t$  to denote the *concatenation* of sequences s and t, and we say that s is an *initial segment* of t, or  $s \sqsubseteq t$ , if there exists s' for which  $t = s \sim s'$ . Endow the set  $\mathbb{N}_*^{\leq \mathbb{N}} = \mathbb{N}^{\leq \mathbb{N}} \cup \{t \sim (\infty) \mid t \in \mathbb{N}^{<\mathbb{N}}\}$ with the smallest topology with respect to which the sets of the form  $\{t\}$  and  $\mathcal{N}_t = \{c \in \mathbb{N}_*^{\leq \mathbb{N}} \mid t \sqsubseteq c\}$ , where  $t \in \mathbb{N}^{<\mathbb{N}}$ , are clopen.

**Proposition 1.1.** The family  $\mathcal{B}$  of sets of the form  $\{t\}$  and  $\mathcal{N}_t \setminus (\{t\} \cup \bigcup_{i \leq i} \mathcal{N}_{t \cap (j)})$ , where  $i \in \mathbb{N}$  and  $t \in \mathbb{N}^{<\mathbb{N}}$ , is a clopen basis for  $\mathbb{N}_*^{\leq \mathbb{N}}$ .

Proof. Let  $\tau$  be the topology generated by  $\mathcal{B}$ . As every set in  $\mathcal{B}$  is clearly clopen, it is sufficient to show that the sets  $\{t\}$  and  $\mathcal{N}_t$  are  $\tau$ clopen for all  $t \in \mathbb{N}^{<\mathbb{N}}$ . As these sets are clearly  $\tau$ -open, we need only show that they are  $\tau$ -closed. As  $\mathcal{N}_{t^{\frown}(i)}$  is  $\tau$ -closed in  $\mathcal{N}_t$  for all  $i \in \mathbb{N}$ and  $t \in \mathbb{N}^{<\mathbb{N}}$ , a straightforward induction shows that  $\mathcal{N}_t$  is  $\tau$ -closed for all  $t \in \mathbb{N}^{<\mathbb{N}}$ . As  $\{t\}$  is  $\tau$ -closed in  $\mathcal{N}_t$  for all  $t \in \mathbb{N}^{<\mathbb{N}}$ , it follows that  $\{t\}$ is  $\tau$ -closed for all  $t \in \mathbb{N}^{<\mathbb{N}}$ .

**Proposition 1.2.** The space  $\mathbb{N}_*^{\leq \mathbb{N}}$  is compact.

*Proof.* Suppose, towards a contradiction, that there is an open cover  $\mathcal{U}$  of  $\mathbb{N}_*^{\leq \mathbb{N}}$  with no finite subcover.

**Lemma 1.3.** Suppose that  $t \in \mathbb{N}^{<\mathbb{N}}$  and no finite set  $\mathcal{V} \subseteq \mathcal{U}$  covers  $\mathcal{N}_t$ . Then there exists  $j \in \mathbb{N}$  such that no finite set  $\mathcal{V} \subseteq \mathcal{U}$  covers  $\mathcal{N}_{t \cap (j)}$ .

Proof. Fix  $U \in \mathcal{U}$  containing  $t \frown (\infty)$ . Proposition 1.1 then yields  $i \in \mathbb{N}$  with  $\mathcal{N}_t \setminus (\{t\} \cup \bigcup_{j < i} \mathcal{N}_{t \frown (j)}) \subseteq U$ , in which case no finite set  $\mathcal{V} \subseteq \mathcal{U}$  covers  $\bigcup_{j < i} \mathcal{N}_{t \frown (j)}$ , and it follows that there exists j < i for which no finite set  $\mathcal{V} \subseteq \mathcal{U}$  covers  $\mathcal{N}_{t \frown (j)}$ .

By recursively applying Lemma 1.3, we obtain  $b \in \mathbb{N}^{\mathbb{N}}$  such that for no  $i \in \mathbb{N}$  is there a finite set  $\mathcal{V} \subseteq \mathcal{U}$  covering  $\mathcal{N}_{b|i}$ . But Proposition 1.1 implies that every open neighborhood of b contains some  $\mathcal{N}_{b|i}$ .

Given a countable set I and a topological space X, we say that a sequence  $(x_i)_{i\in I} \in X^I$  converges to a point  $x \in X$ , or  $x_i \to x$ , if for every open neighborhood U of x there are only finitely many  $i \in I$  with  $x_i \notin U$ . When I and X are equipped with partial orders  $\leq_I$  and  $\leq_X$ , we say that  $(x_i)_{i\in I}$  is decreasing if  $i \leq_I j \implies x_j \leq_X x_i$  for all  $i, j \in I$ .

# **Proposition 1.4.** The space $\mathbb{N}_*^{\leq \mathbb{N}}$ has a compatible ultrametric.

*Proof.* Fix a decreasing sequence  $(\epsilon_t)_{t\in\mathbb{N}^{<\mathbb{N}}}$  of positive real numbers converging to zero. Set d(a, a) = 0 for all  $a \in \mathbb{N}_*^{\leq\mathbb{N}}$ , as well as  $d(a, b) = \max\{\epsilon_t \mid t \in \{a \restriction \min(|a|, i(a, b)), b \restriction \min(|b|, i(a, b))\} \cap \mathbb{N}^{<\mathbb{N}}\}$ for all distinct  $a, b \in \mathbb{N}_*^{\leq\mathbb{N}}$ , where  $i(a, b) = \min\{i \in \mathbb{N} \mid a \restriction i \neq b \restriction i\}$ .

To see that d is an ultrametric, suppose that  $a, b, c \in \mathbb{N}_*^{\leq \mathbb{N}}$  are pairwise distinct. Observe that if  $i(a, c) < \max\{i(a, b), i(b, c)\}$ , then  $d(a, c) \in \{d(b, c), d(a, b)\}$ , so  $d(a, c) \leq \max\{d(a, b), d(b, c)\}$ . And if  $i(a, c) = \max\{i(a, b), i(b, c)\}$ , then setting i = i(a, b) = i(a, c) = i(b, c), it follows that

$$d(a,c) = \max\{\epsilon_t \mid t \in \{a \upharpoonright i, c \upharpoonright i\} \cap \mathbb{N}^{<\mathbb{N}}\}$$
  
$$\leq \max\{\epsilon_t \mid t \in \{a \upharpoonright i, b \upharpoonright i, c \upharpoonright i\} \cap \mathbb{N}^{<\mathbb{N}}\}$$
  
$$= \max\{d(a,b), d(b,c)\}.$$

And if  $i(a,c) > \max\{i(a,b), i(b,c)\}$ , then setting  $\epsilon = d(a,b) = d(b,c)$ and  $t = a \upharpoonright i(a,b) = c \upharpoonright i(b,c)$ , it follows that  $d(a,c) \le \epsilon_t \le \epsilon$ , and therefore  $d(a,c) \le \max\{d(a,b), d(b,c)\}$ .

As  $\{t\} = \mathcal{B}(t, \epsilon_t)$  and  $\mathcal{N}_t \setminus \{t\} = \mathcal{B}(\mathcal{N}_t \setminus \{t\}, \epsilon_t)$  for all  $t \in \mathbb{N}^{<\mathbb{N}}$ , and  $\mathcal{N}_t \setminus (\{t\} \cup \bigcup_{j \leq i} \mathcal{N}_{t \cap (j)}) = \mathcal{B}(\mathcal{N}_t \setminus (\{t\} \cup \bigcup_{j \leq i} \mathcal{N}_{t \cap (j)}), \min(\{\epsilon_{t \cap (j)} \mid j \leq i\}))$ for all  $i \in \mathbb{N}$  and  $t \in \mathbb{N}^{<\mathbb{N}}$ , Proposition 1.1 ensures that every open subset of  $\mathbb{N}_*^{\leq \mathbb{N}}$  is *d*-open.

Given  $b \in \mathbb{N}^{\mathbb{N}}$  and  $\epsilon > 0$ , fix  $i \in \mathbb{N}$  with  $\epsilon_{b \mid i} < \epsilon$ , set  $t = b \mid i$ , and note that  $\mathcal{N}_t \subseteq \mathcal{B}(b,\epsilon)$ . Given  $t \in \mathbb{N}^{<\mathbb{N}}$  and  $\epsilon > 0$ , fix  $i \in \mathbb{N}$  with  $\epsilon_{t \cap (j)} < \epsilon$  for all  $j \ge i$ , and observe that  $\mathcal{N}_t \setminus (\{t\} \cup \bigcup_{j < i} \mathcal{N}_{t \cap (j)}) \subseteq \mathcal{B}(t \cap (\infty), \epsilon)$ . Thus every *d*-open subset of  $\mathbb{N}_*^{\leq \mathbb{N}}$  is open.

It follows that  $\mathbb{N}_*^{\leq \mathbb{N}}$  is Polish. As the space  $\mathbb{N}_*^{\mathbb{N}} = \mathbb{N}_*^{\leq \mathbb{N}} \setminus \mathbb{N}^{<\mathbb{N}}$  is a perfect subset of  $\mathbb{N}_*^{\leq \mathbb{N}}$ , a result of Brouwer's ensures that it is homeomorphic to  $2^{\mathbb{N}}$  (see, for example, [Kec95, Theorem 7.4]).

#### R. CARROY AND B.D. MILLER

#### 2. Meet embeddings

The meet of sequences  $s, t \in \mathbb{N}^{<\mathbb{N}}$  is the sequence  $r = s \wedge t$  of maximal length for which  $r \sqsubseteq s$  and  $r \sqsubseteq t$ . A  $\wedge$ -embedding is an injection  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  such that  $\pi(s \wedge t) = \pi(s) \wedge \pi(t)$  for all  $s, t \in \mathbb{N}^{<\mathbb{N}}$ .

**Proposition 2.1.** Suppose that  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$ . Then  $\pi$  is a  $\wedge$ -embedding if and only if the following conditions hold:

(1) 
$$\forall i \in \mathbb{N} \forall t \in \mathbb{N}^{<\mathbb{N}} \pi(t) \sqsubset \pi(t \frown (i)).$$
  
(2)  $\forall i, j \in \mathbb{N} \forall t \in \mathbb{N}^{<\mathbb{N}}$   
 $(i \neq j \implies \pi(t \frown (i))(|\pi(t)|) \neq \pi(t \frown (j))(|\pi(t)|)).$ 

Proof. Suppose first that  $\pi$  is a  $\wedge$ -embedding. To see that condition (1) holds, observe that if  $i \in \mathbb{N}$  and  $t \in \mathbb{N}^{<\mathbb{N}}$ , then  $\pi(t) = \pi(t) \wedge \pi(t \frown (i))$ , so  $\pi(t) \sqsubseteq \pi(t \frown (i))$ , thus  $\pi(t) \sqsubset \pi(t \frown (i))$ . And to see that condition (2) holds, observe that if  $i, j \in \mathbb{N}$  are distinct and  $t \in \mathbb{N}^{<\mathbb{N}}$ , then  $\pi(t) = \pi(t \frown (i)) \wedge \pi(t \frown (j))$ , so  $\pi(t \frown (i))(|\pi(t)|) \neq \pi(t \frown (j))(|\pi(t)|)$ .

Suppose now that  $\pi$  satisfies conditions (1) and (2). To see that  $\pi$  is a  $\wedge$ -embedding, suppose that  $s, t \in \mathbb{N}^{<\mathbb{N}}$  are distinct, and define  $r = s \wedge t$ . By reversing the roles of s and t if necessary, we can assume that |s| > |r|, so  $\pi(r) \sqsubset \pi(s)$ , thus either r = t or (|t| > |r| and  $\pi(s)(|\pi(r)|) \neq \pi(t)(|\pi(r)|)$ . In both cases, it follows that  $\pi(s) \neq \pi(t)$  and  $\pi(r) = \pi(s) \wedge \pi(t)$ .

**Remark 2.2.** In particular, it follows that if  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  has the property that  $\pi(t) \frown (i) \sqsubseteq \pi(t \frown (i))$  for all  $i \in \mathbb{N}$  and  $t \in \mathbb{N}^{<\mathbb{N}}$ , then  $\pi$  is a  $\wedge$ -embedding.

There is a simple but useful means of amalgamating appropriately indexed families of  $\wedge$ -embeddings.

**Proposition 2.3.** Suppose that  $(\pi_t)_{t\in\mathbb{N}^{<\mathbb{N}}}$  is a sequence of  $\wedge$ -embeddings with the property that  $\pi_t(\mathbb{N}^{<\mathbb{N}}) \subseteq \mathcal{N}_t$  for all  $t\in\mathbb{N}^{<\mathbb{N}}$ . Then the function  $\pi\colon\mathbb{N}^{<\mathbb{N}}\to\mathbb{N}^{<\mathbb{N}}$  given by  $\pi(t)=(\prod_{n\leq |t|}\pi_{t\restriction n})(t)$  is also a  $\wedge$ -embedding.

Proof. Note that if  $i \in \mathbb{N}$  and  $t \in \mathbb{N}^{<\mathbb{N}}$ , then  $t \frown (i) \sqsubseteq \pi_{t \frown (i)}(t \frown (i))$ , so Proposition 2.1 ensures that  $(\prod_{n \le |t|} \pi_{t \upharpoonright n})(t \frown (i)) \sqsubseteq \pi(t \frown (i))$ , thus  $\pi(t) \sqsubset (\prod_{n \le |t|} \pi_{t \upharpoonright n})(t \frown (i)) \sqsubseteq \pi(t \frown (i))$ . It also implies that if  $i \ne j$ , then  $(\prod_{n \le |t|} \pi_{t \upharpoonright n})(t \frown (i))(|\pi(t)|) \ne (\prod_{n \le |t|} \pi_{t \upharpoonright n})(t \frown (j))(|\pi(t)|)$ , so  $\pi(t \frown (i))(|\pi(t)|) \ne \pi(t \frown (j))(|\pi(t)|)$ . One last application of Proposition 2.1 therefore ensures that  $\pi$  is a  $\wedge$ -embedding.

We next consider the connection between  $\wedge$ -embeddings and closed continuous embeddings.

**Proposition 2.4.** Every  $\wedge$ -embedding  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  has a unique extension to a (necessarily injective) continuous map  $\overline{\pi} \colon \mathbb{N}_*^{\leq \mathbb{N}} \to \mathbb{N}_*^{\leq \mathbb{N}}$ , given by  $\overline{\pi}(b) = \bigcup_{i \in \mathbb{N}} \pi(b \upharpoonright i)$  and  $\overline{\pi}(t \frown (\infty)) = \pi(t) \frown (\infty)$  for all  $b \in \mathbb{N}^{\mathbb{N}}$  and  $t \in \mathbb{N}^{<\mathbb{N}}$ .

*Proof.* Suppose that  $\overline{\pi} \colon \mathbb{N}_*^{\leq \mathbb{N}} \to \mathbb{N}_*^{\leq \mathbb{N}}$  is a continuous extension of  $\pi$ . If  $b \in \mathbb{N}^{\mathbb{N}}$ , then  $b \upharpoonright i \to b$ , and since  $(\pi(b \upharpoonright i))_{i \in \mathbb{N}}$  is strictly increasing by Proposition 2.1, it follows that  $\overline{\pi}(b) = \bigcup_{i \in \mathbb{N}} \pi(b \upharpoonright i)$ . If  $t \in \mathbb{N}^{<\mathbb{N}}$ , then  $t \frown (i) \to t \frown (\infty)$ , and since  $\pi(t) = \pi(t \frown (i)) \land \pi(t \frown (j))$  for all distinct  $i, j \in \mathbb{N}$ , it follows that  $\overline{\pi}(t \frown (\infty)) = \pi(t) \frown (\infty)$ .

To see that these constraints actually define a continuous function, note that if  $t \in \mathbb{N}^{<\mathbb{N}}$ , then either  $\overline{\pi}^{-1}(\mathcal{N}_t) = \emptyset$  or there exists  $s \in \mathbb{N}^{<\mathbb{N}}$ of minimal length with  $t \sqsubseteq \pi(s)$ , in which case  $\overline{\pi}^{-1}(\mathcal{N}_t) = \mathcal{N}_s$ .

To see that  $\overline{\pi}$  is injective, it is enough to check that its restriction to  $\mathbb{N}^{\mathbb{N}}$  is injective. Towards this end, suppose that  $a, b \in \mathbb{N}^{\mathbb{N}}$  are distinct, fix  $i \in \mathbb{N}$  least for which  $a(i) \neq b(i)$ , set  $t = a \upharpoonright i = b \upharpoonright i$ , and observe that  $\pi(t \frown (a(i)))(|\pi(t)|) \neq \pi(t \frown (b(i)))(|\pi(t)|)$  by Proposition 2.1, thus  $\overline{\pi}(a)$  and  $\overline{\pi}(b)$  are distinct.

**Remark 2.5.** It follows that the extension associated with the composition of two  $\wedge$ -embeddings is the composition of their extensions.

Given a function  $\phi: X \to Y$  and sets  $X' \subseteq X$  and  $Y' \supseteq \phi(X')$ , let  $\phi \upharpoonright X' \to Y'$  denote the function  $\psi: X' \to Y'$  given by  $\phi(x) = \psi(x)$  for all  $x \in X'$ . Compactness ensures that if  $\pi$  is a  $\wedge$ -embedding, then  $\overline{\pi}$  and  $\overline{\pi} \upharpoonright \mathbb{N}^{\mathbb{N}}_*$  are closed continuous embeddings. The following observations show that so too are  $\overline{\pi} \upharpoonright \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  and  $\overline{\pi} \upharpoonright \mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}_*$ .

**Proposition 2.6.** Suppose that  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  is a  $\wedge$ -embedding. Then  $\overline{\pi} \upharpoonright \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$  is closed.

*Proof.* It is sufficient to show that every sequence  $(b_n)_{n\in\mathbb{N}}$  of elements of  $\mathbb{N}^{\mathbb{N}}$  for which  $(\overline{\pi}(b_n))_{n\in\mathbb{N}}$  converges to an element of  $\mathbb{N}^{\mathbb{N}}$  is itself convergent to an element of  $\mathbb{N}^{\mathbb{N}}$ . As  $(\overline{\pi}(b_n) \upharpoonright i)_{n\in\mathbb{N}}$  is eventually constant for all  $i \in \mathbb{N}$ , a simple induction shows that  $(b_n \upharpoonright i)_{n\in\mathbb{N}}$  is also eventually constant for all  $i \in \mathbb{N}$ , so  $(b_n)_{n\in\mathbb{N}}$  converges to an element of  $\mathbb{N}^{\mathbb{N}}$ .

**Proposition 2.7.** Suppose that  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  is a  $\wedge$ -embedding. Then  $\overline{\pi} \upharpoonright \mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}}$  is closed.

*Proof.* It is sufficient to show that every sequence  $(s_n)_{n\in\mathbb{N}}$  of elements of  $\mathbb{N}^{<\mathbb{N}}$  such that  $(\pi(s_n))_{n\in\mathbb{N}}$  converges to  $t \frown (\infty)$  for some  $t \in \mathbb{N}^{<\mathbb{N}}$ has a subsequence converging to an element of  $\mathbb{N}^{\mathbb{N}}_{\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}}$ . By passing to a subsequence, we can assume that  $\pi(s_m) \land \pi(s_n) = t$  for all distinct  $m, n \in \mathbb{N}$ . Let s be the  $\sqsubseteq$ -minimal element of  $\mathbb{N}^{<\mathbb{N}}$  for which  $t \sqsubseteq \pi(s)$ . Then  $s_m \land s_n = s$  for all distinct  $m, n \in \mathbb{N}$ , thus  $s_n \to s \frown (\infty)$ . A set  $T \subseteq \mathbb{N}^{<\mathbb{N}}$  is  $\sqsubseteq$ -dense if  $\forall s \in \mathbb{N}^{<\mathbb{N}} \exists t \in T \ s \sqsubseteq t$ . More generally, a set  $T \subseteq \mathbb{N}^{<\mathbb{N}}$  is  $\sqsubseteq$ -dense below  $r \in \mathbb{N}^{<\mathbb{N}}$  if  $\forall s \in \mathbb{N}^{<\mathbb{N}} \exists t \in T \ r \frown s \sqsubseteq t$ .

**Proposition 2.8.** Suppose that  $T \subseteq \mathbb{N}^{<\mathbb{N}}$ . Then there is a  $\wedge$ -embedding  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  such that  $\pi(\mathbb{N}^{<\mathbb{N}}) \subseteq T$  or  $\pi(\mathbb{N}^{<\mathbb{N}}) \subseteq \sim T$ .

*Proof.* Fix  $S \in \{T, \sim T\}$  which is  $\sqsubseteq$ -dense below some  $s \in \mathbb{N}^{<\mathbb{N}}$ , and recursively construct a function  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathcal{N}_s \cap S$  with the property that  $\pi(t) \frown (i) \sqsubseteq \pi(t \frown (i))$  for all  $i \in \mathbb{N}$  and  $t \in \mathbb{N}^{<\mathbb{N}}$ .

**Proposition 2.9.** Suppose that  $C \subseteq \mathbb{N}^{\mathbb{N}}$  is a non-meager set with the Baire property. Then there is a  $\wedge$ -embedding  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  with the property that  $\overline{\pi}(\mathbb{N}^{\mathbb{N}}) \subseteq C$ .

Proof. Fix  $s \in \mathbb{N}^{<\mathbb{N}}$  for which C is comeager in  $\mathcal{N}_s \cap \mathbb{N}^{\mathbb{N}}$ , as well as dense open sets  $U_n \subseteq \mathcal{N}_s \cap \mathbb{N}^{\mathbb{N}}$  with the property that  $\bigcap_{n \in \mathbb{N}} U_n \subseteq C$ . Set  $T_n = \{t \in \mathbb{N}^{<\mathbb{N}} \mid \mathcal{N}_t \cap \mathbb{N}^{\mathbb{N}} \subseteq U_n\}$  for all  $n \in \mathbb{N}$ , and recursively construct a function  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathcal{N}_s \cap \mathbb{N}^{<\mathbb{N}}$  such that  $\pi(\mathbb{N}^n) \subseteq T_n$  for all  $n \in \mathbb{N}$  and  $\pi(t) \cap (i) \sqsubseteq \pi(t \cap (i))$  for all  $i \in \mathbb{N}$  and  $t \in \mathbb{N}^{<\mathbb{N}}$ .

## 3. Baire measurable functions on $\mathbb{N}^{\mathbb{N}}$

Here we provide a basis for the class of Baire measurable functions from  $\mathbb{N}^{\mathbb{N}}$  to separable metric spaces.

**Proposition 3.1.** Suppose that X is a second countable topological space and  $\phi \colon \mathbb{N}^{\mathbb{N}} \to X$  is Baire measurable. Then there is a  $\wedge$ -embedding  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  for which  $\phi \circ \overline{\pi}$  is continuous.

*Proof.* Fix a comeager set  $C \subseteq \mathbb{N}^{\mathbb{N}}$  on which  $\phi$  is continuous, and appeal to Proposition 2.9 to obtain a  $\wedge$ -embedding  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  with the property that  $\overline{\pi}(\mathbb{N}^{\mathbb{N}}) \subseteq C$ .

**Proposition 3.2.** Suppose that X is a metric space and  $\phi \colon \mathbb{N}^{\mathbb{N}} \to X$  is continuous. Then there is a  $\wedge$ -embedding  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  with the property that diam  $\phi(\mathcal{N}_{\pi(t)}) \to 0$ .

*Proof.* Fix a sequence  $(\epsilon_t)_{t\in\mathbb{N}^{<\mathbb{N}}}$  of positive real numbers converging to zero, note that the continuity of  $\phi$  ensures that for all  $t\in\mathbb{N}^{<\mathbb{N}}$  the set  $T_t = \{s\in\mathbb{N}^{<\mathbb{N}} \mid \operatorname{diam} \phi(\mathcal{N}_s) < \epsilon_t\}$  is  $\sqsubseteq$ -dense, and recursively construct a function  $\pi\colon\mathbb{N}^{<\mathbb{N}}\to\mathbb{N}^{<\mathbb{N}}$  such that  $\pi(t)\in T_t$  for all  $t\in\mathbb{N}^{<\mathbb{N}}$  and  $\pi(t) \cap (i) \sqsubseteq \pi(t \cap (i))$  for all  $i\in\mathbb{N}$  and  $t\in\mathbb{N}^{<\mathbb{N}}$ .

Given a countable set I and a topological space X, we say that a sequence  $(X_i)_{i \in I}$  of subsets of X converges to a point  $x \in X$ , or  $X_i \to x$ , if for every open neighborhood U of x, all but finitely many  $i \in I$  have the property that  $X_i \subseteq U$ . We say that  $(X_i)_{i \in I}$  is discrete if for all

6

 $x \in X$  there is an open neighborhood U of x such that all but finitely many  $i \in I$  have the property that  $U \cap X_i = \emptyset$ .

**Proposition 3.3.** Suppose that X is a metric space and  $\phi \colon \mathbb{N}^{\mathbb{N}} \to X$ has the property that diam  $\phi(\mathcal{N}_{t_{n}(i)}) \to 0$  for all  $t \in \mathbb{N}^{<\mathbb{N}}$ . Then there is  $a \wedge \text{-embedding } \pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  such that  $(\phi(\mathcal{N}_{\pi(t_{n}(i))}))_{i\in\mathbb{N}}$  is convergent or discrete for all  $t \in \mathbb{N}^{<\mathbb{N}}$ .

Proof. For each  $t \in \mathbb{N}^{<\mathbb{N}}$ , the fact that diam  $\phi(\mathcal{N}_{t \frown (i)}) \to 0$  ensures that there is an injection  $\iota_t \colon \mathbb{N} \to \mathbb{N}$  for which  $(\phi(\mathcal{N}_{t \frown (\iota_t(i))}))_{i \in \mathbb{N}}$  is convergent or discrete. Define  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  by choosing  $\pi(\emptyset) \in \mathbb{N}^{<\mathbb{N}}$  arbitrarily and setting  $\pi(t \frown (i)) = \pi(t) \frown (\iota_{\pi(t)}(i))$  for all  $i \in \mathbb{N}$  and  $t \in \mathbb{N}^{<\mathbb{N}}$ .  $\boxtimes$ 

We say that a function  $\phi: X \to Y$  is *nowhere constant* if there is no non-empty open set  $U \subseteq X$  on which  $\phi$  is constant.

**Proposition 3.4.** Suppose that X is a metric space and  $\phi \colon \mathbb{N}^{\mathbb{N}} \to X$  is continuous and nowhere constant. Then there is a  $\wedge$ -embedding  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  such that

$$\forall i \in \mathbb{N} \forall t \in \mathbb{N}^{<\mathbb{N}} \ \overline{\phi(\mathcal{N}_{\pi(t \frown (i))})} \cap \overline{\bigcup_{j \in \mathbb{N} \setminus \{i\}} \phi(\mathcal{N}_{\pi(t \frown (j))})} = \emptyset.$$

*Proof.* Clearly each  $\phi(\mathcal{N}_t)$  is infinite.

**Lemma 3.5.** For all  $t \in \mathbb{N}^{<\mathbb{N}}$ , there is a function  $\iota_t \colon \mathbb{N} \to \mathbb{N}^{<\mathbb{N}} \setminus \{\emptyset\}$ such that  $(\iota_t(i)(0))_{i\in\mathbb{N}}$  is injective and the closures of  $\phi(\mathcal{N}_{t \frown \iota_t(i)})$  and  $\bigcup_{i\in\mathbb{N}\setminus\{i\}}\phi(\mathcal{N}_{t \frown \iota_t(j)})$  are disjoint for all  $i\in\mathbb{N}$ .

Proof. As each  $\phi(\mathcal{N}_{t^{(i)}})$  is infinite, there are extensions  $b_i \in \mathbb{N}^{\mathbb{N}}$  of  $t^{(i)}$  such that  $\phi(b_i) \notin \{\phi(b_j) \mid j < i\}$  for all  $i \in \mathbb{N}$ . Fix a subsequence  $(a_i)_{i \in \mathbb{N}}$  of  $(b_i)_{i \in \mathbb{N}}$  for which  $\{\phi(a_i) \mid i \in \mathbb{N}\}$  is discrete. For each  $i \in \mathbb{N}$ , fix  $\epsilon_i > 0$  such that  $\phi(a_j) \notin \mathcal{B}(\phi(a_i), \epsilon_i)$  for all  $j \in \mathbb{N} \setminus \{i\}$ , as well as  $\iota_t(i) \in \mathbb{N}^{\mathbb{N}} \setminus \{\emptyset\}$  with  $t \sim \iota_t(i) \sqsubseteq a_i$  and  $\phi(\mathcal{N}_{t \sim \iota_t(i)}) \subseteq \mathcal{B}(\phi(a_i), \epsilon_i/3)$ .

Suppose, towards a contradiction, that there exists  $i \in \mathbb{N}$  for which some  $x \in X$  is in the closures of  $\phi(\mathcal{N}_{t \sim \iota_t(i)})$  and  $\bigcup_{j \in \mathbb{N} \setminus \{i\}} \phi(\mathcal{N}_{t \sim \iota_t(j)})$ . Then there exist  $j \in \mathbb{N} \setminus \{i\}$  and  $y \in \phi(\mathcal{N}_{t \sim \iota_t(j)})$  with the property that  $d(x, y) \leq \epsilon_i/3$ , in which case

$$d(\phi(a_i), \phi(a_j)) \le d(\phi(a_i), x) + d(x, y) + d(y, \phi(a_j))$$
  
$$< \epsilon_i/3 + \epsilon_i/3 + \epsilon_j/3$$
  
$$\le \max\{\epsilon_i, \epsilon_j\},$$

so  $\phi(a_i) \in \mathcal{B}(\phi(a_j), \epsilon_j)$  or  $\phi(a_j) \in \mathcal{B}(\phi(a_i), \epsilon_i)$ , a contradiction.

Define  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  by choosing  $\pi(\emptyset) \in \mathbb{N}^{<\mathbb{N}}$  arbitrarily and setting  $\pi(t \frown (i)) = \pi(t) \frown \iota_{\pi(t)}(i)$  for all  $i \in \mathbb{N}$  and  $t \in \mathbb{N}^{<\mathbb{N}}$ .

We now obtain our main result stabilizing the topological behavior of Baire measurable functions from  $\mathbb{N}^{\mathbb{N}}$  to separable metric spaces.

**Theorem 3.6.** Suppose that X is a separable metric space and  $\phi \colon \mathbb{N}^{\mathbb{N}} \to X$  is Baire measurable. Then there is a  $\wedge$ -embedding  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  such that  $\phi \circ \overline{\pi}$  is constant or extends to a closed continuous embedding on  $\mathbb{N}^{\mathbb{N}}$  or  $\mathbb{N}^{\mathbb{N}}_{*}$ .

*Proof.* By Remark 2.5, we are free to replace  $\phi$  by its composition with the extension of any  $\wedge$ -embedding. For example, by Proposition 3.1, we can assume that  $\phi$  is continuous.

If there exists  $s \in \mathbb{N}^{<\mathbb{N}}$  for which  $\phi \upharpoonright \mathcal{N}_s$  is constant, then define  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  by  $\pi(t) = s \frown t$  for all  $t \in \mathbb{N}^{<\mathbb{N}}$ , so  $\phi \circ \overline{\pi}$  is constant. Otherwise, Propositions 2.8, 3.2, 3.3, and 3.4 yield a  $\wedge$ -embedding  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  such that diam  $\phi(\mathcal{N}_{\pi(t)}) \to 0$ ,  $(\phi(\mathcal{N}_{\pi(t\cap(i))}))_{i\in\mathbb{N}}$  is convergent for all  $t \in \mathbb{N}^{<\mathbb{N}}$  or discrete for all  $t \in \mathbb{N}^{<\mathbb{N}}$ , and

$$\forall i \in \mathbb{N} \forall t \in \mathbb{N}^{<\mathbb{N}} \ \overline{\phi(\mathcal{N}_{\pi(t \frown (i))})} \cap \overline{\bigcup_{j \in \mathbb{N} \setminus \{i\}} \phi(\mathcal{N}_{\pi(t \frown (j))})} = \emptyset$$

As  $\overline{\pi}(\mathcal{N}_t) \subseteq \mathcal{N}_{\pi(t)}$  for all  $t \in \mathbb{N}^{<\mathbb{N}}$ , it follows that

$$\forall i \in \mathbb{N} \forall t \in \mathbb{N}^{<\mathbb{N}} \ \overline{(\phi \circ \overline{\pi})(\mathcal{N}_{t \frown (i)})} \cap \overline{\bigcup_{j \in \mathbb{N} \setminus \{i\}} (\phi \circ \overline{\pi})(\mathcal{N}_{t \frown (j)})} = \emptyset.$$

So by replacing  $\phi$  with  $\phi \circ \overline{\pi}$ , we can assume that diam  $\phi(\mathcal{N}_t) \to 0$ ,  $(\phi(\mathcal{N}_{t \cap (i)}))_{i \in \mathbb{N}}$  is convergent for all  $t \in \mathbb{N}^{<\mathbb{N}}$  or discrete for all  $t \in \mathbb{N}^{<\mathbb{N}}$ , and

(†) 
$$\forall i \in \mathbb{N} \forall t \in \mathbb{N}^{<\mathbb{N}} \ \overline{\phi(\mathcal{N}_{t \frown (i)})} \cap \overline{\bigcup_{j \in \mathbb{N} \setminus \{i\}} \phi(\mathcal{N}_{t \frown (j)})} = \emptyset.$$

To see that  $\phi$  is injective, note that if  $a, b \in \mathbb{N}^{\mathbb{N}}$  are distinct, then there is a least  $i \in \mathbb{N}$  for which  $a(i) \neq b(i)$ . Setting  $t = a \upharpoonright i = b \upharpoonright i$ , it follows from (†) that  $\phi(\mathcal{N}_{t \cap (a(i))})$  and  $\phi(\mathcal{N}_{t \cap (b(i))})$  are disjoint, thus  $\phi(a)$ and  $\phi(b)$  are distinct.

We next check that if  $(\phi(\mathcal{N}_{t_{\frown}(i)}))_{i\in\mathbb{N}}$  is discrete for all  $t\in\mathbb{N}^{<\mathbb{N}}$ , then  $\phi$  is a closed continuous embedding. It is sufficient to show that every sequence  $(b_n)_{n\in\mathbb{N}}$  of elements of  $\mathbb{N}^{\mathbb{N}}$  for which  $(\phi(b_n))_{n\in\mathbb{N}}$  converges to some  $x\in X$  is itself convergent. But a straightforward recursive argument yields  $b\in\mathbb{N}^{\mathbb{N}}$  such that x is in the closure of  $\phi(\mathcal{N}_{b\restriction i})$  for all  $i\in\mathbb{N}$ , so  $(\dagger)$  ensures that x is not in the closure of  $\bigcup_{j\in\mathbb{N}\setminus\{b(i)\}}\phi(\mathcal{N}_{b\restriction i\frown(j)})$  for all  $i\in\mathbb{N}$ , thus  $(b_n\restriction i)_{n\in\mathbb{N}}$  is eventually constant with value  $b\restriction i$  for all  $i\in\mathbb{N}$ , hence  $b_n\to b$ .

It remains to check that if  $(\phi(\mathcal{N}_{t_{\frown}(i)}))_{i\in\mathbb{N}}$  is convergent for all  $t\in\mathbb{N}^{<\mathbb{N}}$ , then the extension of  $\phi$  to  $\mathbb{N}^{\mathbb{N}}_{*}$  given by  $\overline{\phi}(t\frown(\infty)) = \lim_{i\to\infty} \phi(\mathcal{N}_{t\frown(i)})$ for all  $t\in\mathbb{N}^{<\mathbb{N}}$  is a closed continuous embedding. To see that  $\overline{\phi}$  is injective, note that if  $c, d\in\mathbb{N}^{\mathbb{N}}_{*}$  are distinct, then there is a least  $i\in\mathbb{N}$  with  $c(i) \neq d(i)$ . By reversing the roles of c and d if necessary, we can assume that  $c(i) \neq \infty$ . Set  $t = c \upharpoonright i = d \upharpoonright i$ , and appeal to  $(\dagger)$  to see that  $\overline{\phi}(c)$  is in the closure of  $\phi(\mathcal{N}_{t \cap (c(i))})$  but  $\overline{\phi}(d)$  is not, so  $\overline{\phi}(c) \neq \overline{\phi}(d)$ . To see that  $\overline{\phi}$  is continuous, suppose that  $c \in \mathbb{N}^{\mathbb{N}}_{*}$  and Uis an open neighborhood of  $\overline{\phi}(c)$ , and fix an open neighborhood V of  $\overline{\phi}(c)$  whose closure is contained in U. If  $c \in \mathbb{N}^{\mathbb{N}}$ , then there exists  $i \in \mathbb{N}$ for which  $\phi(\mathcal{N}_{c \restriction i}) \subseteq V$ , thus  $\mathcal{N}_{c \restriction i}$  is an open neighborhood of c whose image under  $\overline{\phi}$  is contained in U. Otherwise, there exists  $t \in \mathbb{N}^{<\mathbb{N}}$  for which  $c = t \cap (\infty)$ , as well as  $i \in \mathbb{N}$  for which  $\phi(\mathcal{N}_t \setminus \bigcup_{j < i} \mathcal{N}_{t \cap (j)}) \subseteq V$ . Then  $\mathcal{N}_t \setminus \bigcup_{j < i} \mathcal{N}_{t \cap (j)}$  is an open neighborhood of c whose image under  $\overline{\phi}$  is contained in U.

For each topological space X, let  $c_X$  denote the unique function from X to the trivial topological space  $\{\infty\}$ . Given topological spaces  $X \subseteq Y$ , define  $\iota_{X,Y} \colon X \to Y$  by  $\iota_{X,Y}(x) = x$  for all  $x \in X$ .

**Proposition 3.7.** Suppose that X is a separable metric space,  $\phi \colon \mathbb{N}^{\mathbb{N}} \to X$  is Baire measurable,  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  is a  $\wedge$ -embedding, and  $\phi \circ \overline{\pi}$  is constant or extends to a closed continuous embedding on  $\mathbb{N}^{\mathbb{N}}$  or  $\mathbb{N}^{\mathbb{N}}_*$ . Then there exist  $\phi_0 \in \{c_{\mathbb{N}^{\mathbb{N}}}\} \cup \{\iota_{\mathbb{N}^{\mathbb{N}},Z} \mid Z \in \{\mathbb{N}^{\mathbb{N}}, \mathbb{N}^{\mathbb{N}}_*\}\}$  and  $\psi \colon \overline{\phi_0(\mathbb{N}^{\mathbb{N}})} \to \overline{\phi(\mathbb{N}^{\mathbb{N}})}$  with the property that  $(\overline{\pi} \upharpoonright \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}, \psi)$  is a closed continuous embedding of  $\phi_0$  into  $\phi$ .

*Proof.* If  $\phi \circ \overline{\pi}$  is constant, then set  $\phi_0 = c_{\mathbb{N}^{\mathbb{N}}}$  and let  $\psi$  be the unique function from  $c_{\mathbb{N}^{\mathbb{N}}}(\mathbb{N}^{\mathbb{N}})$  to  $(\phi \circ \overline{\pi})(\mathbb{N}^{\mathbb{N}})$ . If  $\phi \circ \overline{\pi}$  extends to a closed continuous embedding  $\psi$  on  $Z \in \{\mathbb{N}^{\mathbb{N}}, \mathbb{N}^{\mathbb{N}}_*\}$ , then set  $\phi_0 = \iota_{\mathbb{N}^{\mathbb{N}}, Z}$ .

# 4. Baire-class-one functions that are not $\sigma$ -continuous with closed witnesses

Here we strengthen [Sol98, Theorem 3.1] by providing a basis for the class of non- $\sigma$ -continuous-with-closed-witnesses Baire-class-one functions from analytic metric spaces to separable metric spaces.

**Proposition 4.1.** Suppose that X is a metric space and  $\phi \colon \mathbb{N}^{\mathbb{N}}_* \to X$ has the property that  $\phi \upharpoonright \mathbb{N}^{\mathbb{N}}$  is continuous. Then there is a  $\wedge$ -embedding  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  such that either  $(\phi \circ \overline{\pi})(\mathbb{N}^{\mathbb{N}}) \cap (\phi \circ \overline{\pi})(\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}}) = \emptyset$  or  $\phi \circ \overline{\pi}$  is continuous at every point of  $\mathbb{N}^{\mathbb{N}}$ .

*Proof.* We can assume that there is no  $s \in \mathbb{N}^{<\mathbb{N}}$  with the property that  $\inf\{d(\phi(s \frown b), \phi(s \frown t \frown (\infty))) \mid b \in \mathbb{N}^{\mathbb{N}} \text{ and } t \in \mathbb{N}^{<\mathbb{N}}\} > 0$ , since otherwise the  $\wedge$ -embedding  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  given by  $\pi(t) = s \frown t$  for all  $t \in \mathbb{N}^{<\mathbb{N}}$  has the property that  $\overline{(\phi \circ \overline{\pi})(\mathbb{N}^{\mathbb{N}})} \cap \overline{(\phi \circ \overline{\pi})(\mathbb{N}^{\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}})} = \emptyset$ .

**Lemma 4.2.** Suppose that  $\epsilon > 0$  and  $s \in \mathbb{N}^{<\mathbb{N}}$ . Then there exists  $t \in \mathbb{N}^{<\mathbb{N}}$  with  $d(\phi(s \frown t \frown b), \phi(s \frown t \frown (\infty))) < \epsilon$  for all  $b \in \mathbb{N}^{\mathbb{N}}$ .

Proof. Fix  $\delta < \epsilon$  and  $u \in \mathbb{N}^{<\mathbb{N}}$  with diam  $\phi(\mathcal{N}_{s \frown u} \cap \mathbb{N}^{\mathbb{N}}) < \delta$ , and  $b \in \mathbb{N}^{\mathbb{N}}$ and  $v \in \mathbb{N}^{<\mathbb{N}}$  with  $d(\phi(s \frown u \frown b), \phi(s \frown u \frown v \frown (\infty))) < \epsilon - \delta$ , and set  $t = u \frown v$ .

Fix a sequence  $(\epsilon_n)_{n\in\mathbb{N}}$  of positive real numbers converging to zero, and recursively construct a function  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  with the property that  $d(\phi(\pi(t) \frown b), \phi(\pi(t) \frown (\infty))) < \epsilon_{|t|}$  for all  $b \in \mathbb{N}^{\mathbb{N}}$  and  $t \in \mathbb{N}^{<\mathbb{N}}$ , and  $\pi(t) \frown (i) \sqsubseteq \pi(t \frown (i))$  for all  $i \in \mathbb{N}$  and  $t \in \mathbb{N}^{<\mathbb{N}}$ .

We say that a metric space is  $\epsilon$ -discrete if all distinct points have distance at least  $\epsilon$  from one another.

**Proposition 4.3.** Suppose that X is a metric space,  $\phi \colon \mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}} \to X$ ,  $\epsilon > 0$ , and  $t \in \mathbb{N}^{<\mathbb{N}}$ . Then there is a  $\wedge$ -embedding  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathcal{N}_t \cap \mathbb{N}^{<\mathbb{N}}$  with the property that  $\phi \circ \overline{\pi}$  is an injection into an  $\epsilon$ -discrete set or  $(\phi \circ \overline{\pi})(\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}})$  is contained in the  $\epsilon$ -ball around a point of  $\phi(\mathcal{N}_t)$ .

Proof. If for no finite set  $F \subseteq \phi(\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}})$  and extension u of t is it the case that  $\phi(\mathcal{N}_u) \subseteq \mathcal{B}(F, \epsilon)$ , then fix an enumeration  $(t_n)_{n \in \mathbb{N}}$  of  $\mathbb{N}^{<\mathbb{N}}$ with the property that  $t_m \sqsubseteq t_n \implies m \le n$  for all  $m, n \in \mathbb{N}$ , and recursively construct  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathcal{N}_t \cap \mathbb{N}^{<\mathbb{N}}$  such that  $\phi(\pi(t_n) \frown (\infty)) \notin \mathcal{B}(\{\phi(\pi(t_m) \frown (\infty)) \mid m < n\}, \epsilon)$  and  $\pi(t'_n) \frown (n) \sqsubseteq \pi(t_n)$  for all n > 0, where  $t'_n$  is the maximal proper initial segment of  $t_n$ .

Otherwise, there exists  $x \in \phi(\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}})$  with the property that the set  $S = \{s \in \mathbb{N}^{<\mathbb{N}} \mid \phi(s \frown (\infty)) \in \mathcal{B}(x,\epsilon)\}$  is  $\sqsubseteq$ -dense below some extension u of t, in which case we can recursively construct a function  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathcal{N}_u \cap S$  with the property that  $\pi(v) \frown (i) \sqsubseteq \pi(v \frown (i))$  for all  $i \in \mathbb{N}$  and  $v \in \mathbb{N}^{<\mathbb{N}}$ .

**Proposition 4.4.** Suppose that X is a metric space and  $\phi \colon \mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}} \to X$ . Then there is a  $\wedge$ -embedding  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  such that  $\phi \circ \overline{\pi}$  is an injection into an  $\epsilon$ -discrete set for some  $\epsilon > 0$  or diam  $(\phi \circ \overline{\pi})(\mathcal{N}_t) \to 0$ .

Proof. Suppose that for no  $\epsilon > 0$  is there a  $\wedge$ -embedding  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  such that  $\phi \circ \overline{\pi}$  is an injection into an  $\epsilon$ -discrete set, fix a sequence  $(\epsilon_t)_{t\in\mathbb{N}^{<\mathbb{N}}}$  of positive real numbers converging to zero, and recursively apply Proposition 4.3 to the functions  $\phi_t = \phi \circ \prod_{n < |t|} \overline{\pi_{t\mid n}}$  to obtain  $\wedge$ -embeddings  $\pi_t \colon \mathbb{N}^{<\mathbb{N}} \to \mathcal{N}_t \cap \mathbb{N}^{<\mathbb{N}}$  such that  $(\phi \circ \prod_{n \le |t|} \overline{\pi_{t\mid n}})(\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}})$  is contained in an  $\epsilon_t$ -ball for all  $t \in \mathbb{N}^{<\mathbb{N}}$ . Let  $\pi$  be the  $\wedge$ -embedding obtained from applying Proposition 2.3 to  $(\pi_t)_{t\in\mathbb{N}^{<\mathbb{N}}}$ , and observe that diam  $(\phi \circ \overline{\pi})(\mathcal{N}_t) \to 0$ .

10

Define  $p: \mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  by setting  $p(t \frown (\infty)) = t$  for all  $t \in \mathbb{N}^{<\mathbb{N}}$ . Let  $\mathbb{N}^{<\mathbb{N}}_* = \mathbb{N}^{<\mathbb{N}} \cup \{\infty\}$  denote the *one-point compactification* of  $\mathbb{N}^{<\mathbb{N}}$ .

**Theorem 4.5.** Suppose that X is an analytic metric space, Y is a separable metric space, and  $\phi: X \to Y$  is a Baire-class-one function that is not  $\sigma$ -continuous with closed witnesses. Then there exists  $\phi_0 \in \{c_{\mathbb{N}^{\mathbb{N}}}\} \cup \{\iota_{\mathbb{N}^{\mathbb{N}},Z} \mid Z \in \{\mathbb{N}^{\mathbb{N}}, \mathbb{N}^{\mathbb{N}}_*\}\}$  for which there is a closed continuous embedding of  $\phi_0 \cup p$  into  $\phi$ .

Proof. By the Jayne-Rogers theorem (see, for example, [JR82, Theorem 1]), we can assume that  $\phi$  is not  $G_{\delta}$ -measurable. Hurewicz's dichotomy theorem for  $F_{\sigma}$  sets then yields a closed continuous embedding  $\psi \colon \mathbb{N}^{\mathbb{N}}_* \to X$  with  $(\phi \circ \psi)(\mathbb{N}^{\mathbb{N}}) \cap (\phi \circ \psi)(\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}}) = \emptyset$  (see, for example, [CMS, Theorem 4.2]). As  $(\psi, \operatorname{id}_{\overline{(\phi \circ \psi)}(\mathbb{N}^{\mathbb{N}}_*)})$  is a closed continuous embedding of  $\phi \circ \psi$  into  $\phi$ , by replacing the latter with the former, we can assume that  $X = \mathbb{N}^{\mathbb{N}}_*$  and  $\overline{\phi}(\mathbb{N}^{\mathbb{N}}) \cap \phi(\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}}) = \emptyset$ . By Proposition 3.1, there is a  $\wedge$ -embedding  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  for which

By Proposition 3.1, there is a  $\wedge$ -embedding  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  for which  $(\phi \circ \overline{\pi}) \upharpoonright \mathbb{N}^{\mathbb{N}}$  is continuous. By composing  $\pi$  with the  $\wedge$ -embedding given by Proposition 4.1, we can assume that  $(\overline{\phi \circ \pi})(\mathbb{N}^{\mathbb{N}}) \cap (\overline{\phi \circ \pi})(\mathbb{N}^{\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}}) = \emptyset$  or  $\phi \circ \overline{\pi}$  is continuous at every point of  $\mathbb{N}^{\mathbb{N}}$ . As  $\phi$  is Baire class one, the former possibility would imply that the pre-images of  $(\overline{\phi \circ \pi})(\mathbb{N}^{\mathbb{N}})$ and  $(\overline{\phi \circ \pi})(\mathbb{N}^{\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}})$  under  $\phi \circ \overline{\pi}$  are disjoint dense  $G_{\delta}$  subsets of  $\mathbb{N}^{\mathbb{N}}_{*}$ , so the latter holds. By Proposition 4.4, we can assume that either there exists  $\epsilon > 0$  for which  $(\phi \circ \overline{\pi}) \upharpoonright \mathbb{N}^{\mathbb{N}}_{*} \setminus \mathbb{N}^{\mathbb{N}}$  is an injection into an  $\epsilon$ -discrete set, or diam  $(\phi \circ \overline{\pi})(\mathcal{N}_{t} \cap (\mathbb{N}^{\mathbb{N}}_{*} \setminus \mathbb{N}^{\mathbb{N}})) \to 0$ . As the former possibility contradicts the facts that  $(\phi \circ \overline{\pi})(\mathbb{N}^{\mathbb{N}}) \cap (\phi \circ \overline{\pi})(\mathbb{N}^{\mathbb{N}}_{*} \setminus \mathbb{N}^{\mathbb{N}}) = \emptyset$ and  $(\phi \circ \overline{\pi})(\mathbb{N}^{\mathbb{N}}) \subseteq \overline{(\phi \circ \overline{\pi})(\mathbb{N}^{\mathbb{N}}_{*} \setminus \mathbb{N}^{\mathbb{N}})}$ , it follows that the latter holds. By applying Proposition 4.3 with any  $\epsilon > 0$  and  $t \in \mathbb{N}^{<\mathbb{N}}$ , but replacing the given metric on X by one with respect to which all pairs of distinct points have distance at least  $\epsilon$  from one another, we can assume that  $(\phi \circ \overline{\pi}) \upharpoonright \mathbb{N}^{\mathbb{N}}_{*} \setminus \mathbb{N}^{\mathbb{N}}$  is either constant or injective.

**Lemma 4.6.** Suppose that  $(s_n)_{n \in \mathbb{N}}$  is an injective sequence of elements of  $\mathbb{N}^{<\mathbb{N}}$  and  $(b_n)_{n \in \mathbb{N}}$  is a sequence of elements of  $\mathbb{N}^{\mathbb{N}}$  such that  $s_n \sqsubseteq b_n$ for all  $n \in \mathbb{N}$ . Then  $d_X((\phi \circ \overline{\pi})(b_n), (\phi \circ \overline{\pi})(s_n \frown (\infty))) \to 0$ .

Proof. Simply note that  $(\phi \circ \overline{\pi})(b_n) \in \overline{(\phi \circ \overline{\pi})(\mathcal{N}_{s_n} \cap (\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}}))}$  for all  $n \in \mathbb{N}$  and diam  $(\phi \circ \overline{\pi})(\mathcal{N}_{s_n} \cap (\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}})) \to 0.$ 

Along with the facts that  $(\phi \circ \overline{\pi})(\mathbb{N}^{\mathbb{N}}) \cap (\phi \circ \overline{\pi})(\mathbb{N}^{\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}}) = \emptyset$  and  $(\phi \circ \overline{\pi})(\mathbb{N}^{\mathbb{N}}) \subseteq \overline{(\phi \circ \overline{\pi})(\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}})}$ , Lemma 4.6 ensures that  $(\phi \circ \overline{\pi}) \upharpoonright \mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}}$  is not constant, and is therefore injective. Along with the fact that  $(\overline{\phi \circ \overline{\pi})(\mathbb{N}^{\mathbb{N}})} \cap (\phi \circ \overline{\pi})(\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}}) = \emptyset$ , Lemma 4.6 ensures that  $(\phi \circ \overline{\pi})(\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}})$  is discrete.

By Theorem 3.6, we can assume that  $(\phi \circ \overline{\pi}) \upharpoonright \mathbb{N}^{\mathbb{N}}$  is constant or extends to a closed continuous embedding on  $\mathbb{N}^{\mathbb{N}}$  or  $\mathbb{N}^{\mathbb{N}}_*$ .

We will now complete the proof by showing that there exist  $\phi_0 \in \{c_{\mathbb{N}^{\mathbb{N}}}\} \cup \{\iota_{\mathbb{N}^{\mathbb{N}},Z} \mid Z \in \{\mathbb{N}^{\mathbb{N}}, \mathbb{N}_*^{\mathbb{N}}\}\}$  and  $\psi \colon \phi_0(\mathbb{N}_*^{\mathbb{N}}) \cup \mathbb{N}^{<\mathbb{N}} \to \overline{\phi(X)}$  for which  $(\overline{\pi} \upharpoonright \mathbb{N}_*^{\mathbb{N}} \to \mathbb{N}_*^{\mathbb{N}}, \psi)$  is a closed continuous embedding of  $\phi_0 \cup p$  into  $\phi$ .

If  $(\phi \circ \overline{\pi}) \upharpoonright \mathbb{N}^{\mathbb{N}}$  is constant with value  $y \in Y$ , then set  $\phi_0 = c_{\mathbb{N}^{\mathbb{N}}}$ , and note that the extension  $\psi$  of  $\phi \circ \overline{\pi} \circ p^{-1}$  to  $\mathbb{N}^{<\mathbb{N}}_*$  given by  $\psi(\infty) = y$ is injective. As Lemma 4.6 ensures that  $(\phi \circ \overline{\pi})(s_n \frown (\infty)) \to y$  for every injective sequence  $(s_n)_{n \in \mathbb{N}}$  of elements of  $\mathbb{N}^{<\mathbb{N}}$ , it follows that  $\psi$ is continuous, so the compactness of  $\mathbb{N}^{<\mathbb{N}}_*$  ensures that  $\psi$  is a closed continuous embedding.

If  $(\phi \circ \overline{\pi}) \upharpoonright \mathbb{N}^{\mathbb{N}}$  is a closed continuous embedding, then set  $\phi_0 = \iota_{\mathbb{N}^{\mathbb{N}},\mathbb{N}^{\mathbb{N}}}$ , and note that the extension  $\psi$  of  $\phi \circ \overline{\pi} \circ p^{-1}$  to  $\mathbb{N}^{\leq \mathbb{N}}$  given by  $\psi \upharpoonright \mathbb{N}^{\mathbb{N}} = (\phi \circ \overline{\pi}) \upharpoonright \mathbb{N}^{\mathbb{N}}$  is a continuous injection. To see that it is closed, it is enough to show that every injective sequence  $(a_n)_{n \in \mathbb{N}}$  of elements of  $\mathbb{N}^{\leq \mathbb{N}}$  for which  $(\psi(a_n))_{n \in \mathbb{N}}$  converges to some point  $y \in Y$  has a subsequence converging to a point of  $\mathbb{N}^{\mathbb{N}}$ . As  $\mathbb{N}_*^{\leq \mathbb{N}}$  is compact, by passing to a subsequence, we can assume that  $(a_n)_{n \in \mathbb{N}}$  converges to a point of  $\mathbb{N}_*^{\leq \mathbb{N}}$ . As every point of  $\mathbb{N}^{<\mathbb{N}}$  is isolated, it therefore converges to a point of  $\mathbb{N}_*^{\mathbb{N}}$ . And if there exists  $t \in \mathbb{N}^{<\mathbb{N}}$  for which  $a_n \to t \frown (\infty)$ , then there are extensions  $b_n \in \mathbb{N}^{\mathbb{N}}$  of  $a_n$  for all  $n \in \mathbb{N}$ , in which case  $b_n \to t \frown (\infty)$  and  $\psi(b_n) \to y$  by Lemma 4.6. Fix  $n \in \mathbb{N}$  sufficiently large that  $(\phi \circ \overline{\pi})(b_m) \neq y$  for all  $m \geq n$ , and observe that  $\{b_m \mid m \geq n\}$  is a closed subset of  $\mathbb{N}^{\mathbb{N}}$  whose image under  $\phi \circ \overline{\pi}$  is not closed, contradicting the fact that  $(\phi \circ \overline{\pi}) \upharpoonright \mathbb{N}^{\mathbb{N}}$  is closed.

If  $(\phi \circ \overline{\pi}) \upharpoonright \mathbb{N}^{\mathbb{N}}$  extends to a closed continuous embedding  $\psi'$  on  $\mathbb{N}_*^{\mathbb{N}}$ , then set  $\phi_0 = \iota_{\mathbb{N}^{\mathbb{N}}, \mathbb{N}_*^{\mathbb{N}}}$ , and note that the extension  $\psi$  of  $\phi \circ \overline{\pi} \circ p^{-1}$ to  $\mathbb{N}_*^{\leq \mathbb{N}}$  given by  $\psi \upharpoonright \mathbb{N}_*^{\mathbb{N}} = \psi' \upharpoonright \mathbb{N}_*^{\mathbb{N}}$  is injective. To see that it is continuous, suppose that  $(t_n)_{n \in \mathbb{N}}$  is an injective sequence of elements of  $\mathbb{N}^{<\mathbb{N}}$  converging to  $t \frown (\infty)$  for some  $t \in \mathbb{N}^{<\mathbb{N}}$ , fix  $b_n \in \mathcal{N}_{t_n} \cap \mathbb{N}^{\mathbb{N}}$  for all  $n \in \mathbb{N}$ , and observe that the continuity of  $\psi'$  ensures that  $\psi(b_n) \rightarrow$  $\psi(t \frown (\infty))$ , thus Lemma 4.6 implies that  $\psi(t_n) \rightarrow \psi(t \frown (\infty))$ . As  $\mathbb{N}_*^{\leq \mathbb{N}}$  is compact, it follows that  $\psi$  is a closed continuous embedding.

# 5. Functions on $\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}}$

Here we provide a basis for the class of all functions from  $\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}}$  to analytic metric spaces.

**Proposition 5.1.** Suppose that X is a topological space,  $\phi \colon \mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}} \to X$  is injective, and  $x \in X$ . Then there is a  $\wedge$ -embedding  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  such that  $x \notin (\phi \circ \overline{\pi})(\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}})$ .

12

*Proof.* Fix  $s \in \mathbb{N}^{<\mathbb{N}}$  such that  $x \notin \phi(\mathcal{N}_s)$ , and define  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  by  $\pi(t) = s \frown t$  for all  $t \in \mathbb{N}^{<\mathbb{N}}$ .

**Proposition 5.2.** Suppose that X is a metric space and  $\phi \colon \mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}} \to X$ . Then there is a  $\wedge$ -embedding  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  with the property that  $((\phi \circ \overline{\pi})(t \frown (i, \infty)))_{i \in \mathbb{N}}$  is convergent or  $\{(\phi \circ \overline{\pi})(t \frown (i, \infty)) \mid i \in \mathbb{N}\}$  is closed and discrete for all  $t \in \mathbb{N}^{<\mathbb{N}}$ .

Proof. For each  $t \in \mathbb{N}^{<\mathbb{N}}$ , there is an injection  $\iota_t \colon \mathbb{N} \to \mathbb{N}$  for which  $(\phi(t \frown (\iota_t(i), \infty)))_{i \in \mathbb{N}}$  is convergent or  $\{\phi(t \frown (\iota_t(i), \infty)) \mid i \in \mathbb{N}\}$  is closed and discrete. Define  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  by choosing  $\pi(\emptyset) \in \mathbb{N}^{<\mathbb{N}}$  arbitrarily and setting  $\pi(t \frown (i)) = \pi(t) \frown (\iota_{\pi(t)}(i))$  for all  $i \in \mathbb{N}$  and  $t \in \mathbb{N}^{<\mathbb{N}}$ , and note that  $(\phi \circ \overline{\pi})(t \frown (i, \infty)) = \phi(\pi(t \frown (i)) \frown (\infty)) = \phi(\pi(t) \frown (\iota_{\pi(t)}(i), \infty))$  for all  $i \in \mathbb{N}$  and  $t \in \mathbb{N}^{\mathbb{N}}$ .

**Proposition 5.3.** Suppose that X is a metric space,  $\phi \colon \mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}} \to X$ ,  $F \subseteq X$  is finite, and  $t \in \mathbb{N}^{<\mathbb{N}}$ . Then there is a  $\wedge$ -embedding  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathcal{N}_t \cap \mathbb{N}^{<\mathbb{N}}$  such that either  $((\phi \circ \overline{\pi})(u \frown (\infty)))_{u \in \mathbb{N}^{<\mathbb{N}}}$  converges to an element of F or the closure of  $(\phi \circ \overline{\pi})(\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}})$  is disjoint from F.

Proof. If the set  $S_{\epsilon} = \{s \in \mathbb{N}^{<\mathbb{N}} \mid \phi(s \frown (\infty)) \in \mathcal{B}(F, \epsilon)\}$  is  $\sqsubseteq$ -dense below t for all  $\epsilon > 0$ , then there exist an extension u of t and  $x \in$ F such that the set  $S_{\epsilon,x} = \{s \in \mathbb{N}^{<\mathbb{N}} \mid \phi(s \frown (\infty)) \in \mathcal{B}(x, \epsilon)\}$  is  $\sqsubseteq$ -dense below u for all  $\epsilon > 0$ . Fix a sequence  $(\epsilon_v)_{v \in \mathbb{N}^{<\mathbb{N}}}$  of positive real numbers converging to zero, and recursively construct a function  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathcal{N}_u \cap \mathbb{N}^{<\mathbb{N}}$  such that  $\pi(v) \in S_{\epsilon_v,x}$  for all  $v \in \mathbb{N}^{<\mathbb{N}}$  and  $\pi(v) \frown (i) \sqsubseteq \pi(v \frown (i))$  for all  $i \in \mathbb{N}$  and  $v \in \mathbb{N}^{<\mathbb{N}}$ , and observe that  $(\phi \circ \overline{\pi})(v \frown (\infty)) \to x$ .

Otherwise, fix  $\epsilon > 0$  and an extension u of t with the property that  $\mathcal{N}_u \cap S_{\epsilon} = \emptyset$ , define  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathcal{N}_u \cap \mathbb{N}^{<\mathbb{N}}$  by  $\pi(v) = u \frown v$ , and note that the closure of  $(\phi \circ \overline{\pi})(\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}})$  is disjoint from F.

For the rest of this section, it will be convenient to fix an enumeration  $(t_n)_{n\in\mathbb{N}}$  of  $\mathbb{N}^{<\mathbb{N}}$  such that  $t_m \sqsubseteq t_n \implies m \le n$  for all  $m, n \in \mathbb{N}$ .

**Proposition 5.4.** Suppose that X is a metric space and  $\phi: \mathbb{N}_*^{\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}} \to X$ . Then there is a  $\wedge$ -embedding  $\pi: \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  with the property that  $((\phi \circ \overline{\pi})(t \frown (\infty)))_{t \in \mathbb{N}^{<\mathbb{N}}}$  converges or for no natural numbers m < n is  $(\phi \circ \overline{\pi})(t_m \frown (\infty))$  or a limit point of  $\{(\phi \circ \overline{\pi})(t_m \frown (i, \infty)) \mid i \in \mathbb{N}\}$  in the closure of  $(\phi \circ \overline{\pi})(\mathcal{N}_{t_n})$ .

Proof. Suppose that for no  $\wedge$ -embedding  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  is the sequence  $((\phi \circ \overline{\pi})(t \frown (\infty)))_{t \in \mathbb{N}^{<\mathbb{N}}}$  convergent. By Proposition 5.2, we can assume that  $(\phi(t \frown (i, \infty)))_{i \in \mathbb{N}}$  is convergent or  $\{\phi(t \frown (i, \infty)) \mid i \in \mathbb{N}\}$  is closed and discrete for all  $t \in \mathbb{N}^{<\mathbb{N}}$ . By recursively applying Lemma

5.3 to the functions  $\phi_t = \phi \circ \prod_{k < |t|} \overline{\pi_{t \restriction k}}$ , we obtain  $\wedge$ -embeddings  $\pi_t \colon \mathbb{N}^{<\mathbb{N}} \to \mathcal{N}_t \cap \mathbb{N}^{<\mathbb{N}}$  such that for no natural numbers m < n is  $(\phi \circ \prod_{k \le |t_m|} \overline{\pi_{t_m \restriction k}})(t_m \frown (\infty))$  or a limit point of  $\{(\phi \circ \prod_{k \le |t_m|} \overline{\pi_{t_m \restriction k}})(t_m \frown (i, \infty)) \mid i \in \mathbb{N}\}$  in the closure of  $(\phi \circ \prod_{k \le |t_n|} \overline{\pi_{t_n \restriction k}})(\mathcal{N}_{t_n})$ . Let  $\pi$  be the  $\wedge$ -embedding obtained from applying Proposition 2.3 to  $(\pi_t)_{t \in \mathbb{N}^{<\mathbb{N}}}$ , and observe that for no natural numbers m < n is it the case that  $(\phi \circ \overline{\pi})(t_m \frown (\infty))$  or a limit point of  $\{(\phi \circ \overline{\pi})(t_m \frown (i, \infty)) \mid i \in \mathbb{N}\}$  in the closure of  $(\phi \circ \overline{\pi})(\mathcal{N}_{t_n})$ .

**Theorem 5.5.** Suppose that X is an analytic metric space and  $\phi \colon \mathbb{N}_*^{\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}} \to X$ . Then there is a  $\wedge$ -embedding  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  such that  $\phi \circ \overline{\pi}$  is constant,  $\phi \circ \overline{\pi}$  extends to a closed continuous embedding on  $\mathbb{N}_*^{\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}}$  or  $\mathbb{N}_*^{\mathbb{N}}$ , or  $\phi \circ \overline{\pi} \circ p^{-1}$  extends to a closed continuous embedding on  $\mathbb{N}^{<\mathbb{N}}$ ,  $\mathbb{N}_*^{<\mathbb{N}}$ ,  $\mathbb{N}_*^{\leq\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}}$ , or  $\mathbb{N}_*^{\leq\mathbb{N}}$ .

*Proof.* As before, we will repeatedly precompose  $\phi$  with appropriate  $\wedge$ -embeddings, albeit this time so as to stabilize the behavior of the function  $\psi = \phi \circ p^{-1}$ , as opposed to that of the function  $\phi$  itself. By applying Proposition 4.3 with any  $\epsilon > 0$  and  $t \in \mathbb{N}^{<\mathbb{N}}$ , but replacing the given metric on X by one with respect to which all pairs of distinct points have distance at least  $\epsilon$  from one another, we can assume that  $\psi$  is either constant or injective. As  $\phi$  is constant in the former case, we can assume that we are in the latter.

By Proposition 4.4, we can ensure that  $\psi(\mathbb{N}^{<\mathbb{N}})$  is closed and discrete or diam  $\psi(\mathcal{N}_t) \to 0$ . As  $\psi$  is a closed continuous embedding in the former case, we can assume that we are in the latter.

Let  $\overline{\psi}$  be the extension of  $\psi$  to a partial function on  $\mathbb{N}_*^{\leq \mathbb{N}}$  given by  $\overline{\psi}(b) = \lim_{i \to \infty} \psi(b \upharpoonright i)$  and  $\overline{\psi}(t \frown (\infty)) = \lim_{i \to \infty} \psi(t \frown (i))$  for all  $b \in \mathbb{N}^{\mathbb{N}}$  and  $t \in \mathbb{N}^{<\mathbb{N}}$ . By Proposition 5.2, we can assume that  $\{\psi(t \frown (i)) \mid i \in \mathbb{N}\}$  has a limit point  $\implies t \frown (\infty) \in \operatorname{dom}(\overline{\psi})$  for all  $t \in \mathbb{N}^{<\mathbb{N}}$ .

As each point of  $\mathbb{N}^{<\mathbb{N}}$  is isolated, diam  $\psi(\mathcal{N}_{b\restriction i}) \to 0$  for all  $b \in \mathbb{N}^{\mathbb{N}}$ , and diam  $\psi(\mathcal{N}_{t \cap (i)}) \to 0$  for all  $t \in \mathbb{N}^{<\mathbb{N}}$ , it follows that  $\overline{\psi}$  is continuous. To see that  $\overline{\psi}$  is closed, it is sufficient show that every injective sequence  $(c_n)_{n\in\mathbb{N}}$  of points in the domain of  $\overline{\psi}$  for which  $(\overline{\psi}(c_n))_{n\in\mathbb{N}}$  is convergent has a subsequence converging to a point in the domain of  $\overline{\psi}$ . By passing to a subsequence, we can assume that the sequence converges to a point of  $\mathbb{N}_*^{\leq\mathbb{N}}$ . As each point of  $\mathbb{N}^{<\mathbb{N}}$  is isolated, the sequence converges to a point of  $\mathbb{N}_*^{\mathbb{N}}$ , so the facts that diam  $\psi(\mathcal{N}_{b\restriction i}) \to 0$  for all  $b \in \mathbb{N}^{\mathbb{N}}$ , diam  $\psi(\mathcal{N}_{t \cap (i)}) \to 0$  for all  $t \in \mathbb{N}^{<\mathbb{N}}$ , and  $\{\psi(t \cap (i)) \mid i \in \mathbb{N}\}$  has a limit point  $\implies t \cap (\infty) \in \operatorname{dom}(\overline{\psi})$  for all  $t \in \mathbb{N}^{<\mathbb{N}}$  ensure that it converges to a point of the domain of  $\overline{\psi}$ . By Proposition 2.8, we can assume that one of the following holds:

- (1)  $\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}} \subseteq \operatorname{dom}(\overline{\psi}) \text{ and } \forall t \in \mathbb{N}^{<\mathbb{N}} \ \overline{\psi}(t) = \overline{\psi}(t \frown (\infty)).$ (2)  $\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}} \subseteq \operatorname{dom}(\overline{\psi}) \text{ and } \forall t \in \mathbb{N}^{<\mathbb{N}} \ \overline{\psi}(t) \neq \overline{\psi}(t \frown (\infty)).$
- (3)  $(\mathbb{N}^{\mathbb{N}}_{*} \setminus \mathbb{N}^{\mathbb{N}}) \cap \operatorname{dom}(\overline{\psi}) = \emptyset.$

As the domain of  $\overline{\psi}$  is analytic, so too is its intersection with  $\mathbb{N}^{\mathbb{N}}$ . It follows that the latter intersection has the Baire property, so Proposition 2.9 allows us to assume that one of the following holds:

- (a) The domain of  $\overline{\psi}$  is disjoint from  $\mathbb{N}^{\mathbb{N}}$ .
- (b) The domain of  $\overline{\psi}$  contains  $\mathbb{N}^{\mathbb{N}}$ .

In the special case that condition (b) holds, Theorem 3.6 allows us to assume that  $\overline{\psi} \upharpoonright \mathbb{N}^{\mathbb{N}}$  is either constant or injective.

Proposition 5.4 allows us to assume that  $(\psi(t))_{t\in\mathbb{N}^{\leq \mathbb{N}}}$  converges to some  $x \in X$  or for no natural numbers m < n is  $\psi(t_m)$  or  $\overline{\psi}(t_m \frown (\infty))$ in the closure of  $\psi(\mathcal{N}_{t_n})$ . In the former case, Proposition 5.1 allows us to assume that  $\psi(\mathbb{N}^{<\mathbb{N}})$  is discrete, so the extension of  $\psi$  to  $\mathbb{N}^{<\mathbb{N}}_*$ sending  $\infty$  to x is a closed continuous embedding, thus we can assume that we are in the latter.

**Lemma 5.6.** Suppose that  $c, d \in \text{dom}(\overline{\psi})$  are distinct but  $\overline{\psi}(c) = \overline{\psi}(d)$ . Then there exists  $t \in \mathbb{N}^{<\mathbb{N}}$  such that  $\{c, d\} = \{t, t \land (\infty)\}.$ 

*Proof.* To see that  $\overline{\psi} \upharpoonright \mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}}$  is injective, observe that if m < n, both  $t_m \sim (\infty)$  and  $t_n \sim (\infty)$  are in the domain of  $\overline{\psi}$ , and moreover  $\overline{\psi}(t_m \frown (\infty)) = \overline{\psi}(t_n \frown (\infty))$ , then  $\overline{\psi}(t_m \frown (\infty))$  is in the closure of  $\psi(\mathcal{N}_{t_n})$ , a contradiction.

To see that  $\overline{\psi} \upharpoonright \mathbb{N}^{\mathbb{N}}$  is injective when  $\mathbb{N}^{\mathbb{N}}$  is contained in the domain of  $\overline{\psi}$ , note that otherwise it is constant, and let x be this constant value. Then for each  $t \in \mathbb{N}^{<\mathbb{N}}$ , there is a sequence  $(u_i)_{i\in\mathbb{N}}$  of elements of  $\mathbb{N}^{<\mathbb{N}}$ such that  $\psi(t \cap (i) \cap (u_i)) \to x$ , so the fact that diam  $\psi(\mathcal{N}_{t \cap (i)}) \to 0$ ensures that  $\overline{\psi}(t \frown (\infty)) = x$ , contradicting the fact that  $\overline{\psi} \upharpoonright \mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}}$ is injective.

To see that  $\overline{\psi}(\mathbb{N}^{\mathbb{N}}) \cap \psi(\mathbb{N}^{<\mathbb{N}}) = \emptyset$ , note that if  $b \in \operatorname{dom}(\overline{\psi}) \cap \mathbb{N}^{\mathbb{N}}$ ,  $t \in \mathbb{N}^{<\mathbb{N}}$ , and  $\overline{\psi}(b) = \psi(t)$ , then there exist m < n with  $t_m = t$  and  $t_n \sqsubset b$ , so  $\psi(t_m)$  is in the closure of  $\psi(\mathcal{N}_{t_n})$ , a contradiction.

To see that  $\overline{\psi}(\mathbb{N}^{\mathbb{N}}) \cap \overline{\psi}(\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}}) = \emptyset$ , note that if  $b \in \operatorname{dom}(\overline{\psi}) \cap \mathbb{N}^{\mathbb{N}}$ ,  $t \in \mathbb{N}^{<\mathbb{N}}, t \frown (\infty) \in \operatorname{dom}(\overline{\psi}), \text{ and } \overline{\psi}(b) = \overline{\psi}(t \frown (\infty)), \text{ then there exist}$ m < n with  $t_m = t$  and  $t_n \sqsubset b$ , in which case  $\overline{\psi}(t_m \frown \infty)$  is in the closure of  $\psi(\mathcal{N}_{t_n})$ , a contradiction.

Observe finally that if  $s, t \in \mathbb{N}^{<\mathbb{N}}$  are distinct,  $t \frown (\infty) \in \operatorname{dom}(\overline{\psi})$ , and  $\psi(s) = \overline{\psi}(t \frown (\infty))$ , then there exist  $m \neq n$  such that  $t_m = s$  and  $t_n = t$ . Then  $\psi(t_m)$  is in the closure of  $\psi(\mathcal{N}_{t_n})$  and  $\overline{\psi}(t_n \frown (\infty))$  is in  $\psi(\mathcal{N}_{t_m})$ , a contradiction.

If (1a) or (1b) holds, then  $\overline{\psi} \upharpoonright \mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}}$  or  $\overline{\psi} \upharpoonright \mathbb{N}^{\mathbb{N}}_*$  is an extension of  $\phi$  to a closed continuous embedding. If (2a), (2b), (3a), or (3b) holds, then  $\overline{\psi}$  is an extension of  $\psi$  to a closed continuous embedding on  $\mathbb{N}^{\leq \mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}}$ ,  $\mathbb{N}^{\leq \mathbb{N}}_*$ , or  $\mathbb{N}^{\leq \mathbb{N}}$ .

**Proposition 5.7.** Suppose that X is an analytic metric space,  $\phi \colon \mathbb{N}_*^{\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}} \to X$ ,  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  is a  $\wedge$ -embedding, and  $\phi \circ \overline{\pi}$  is constant,  $\phi \circ \overline{\pi}$  extends to a closed continuous embedding on  $\mathbb{N}_*^{\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}}$  or  $\mathbb{N}_*^{\mathbb{N}}$ ,  $or \phi \circ \overline{\pi} \circ p^{-1}$  extends to a closed continuous embedding on  $\mathbb{N}^{<\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}}$ ,  $\mathbb{N}_*^{<\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}}$ ,  $\mathbb{N}^{<\mathbb{N}}$ ,  $\mathbb{N}^{<\mathbb{N}}$ ,  $\mathbb{N}^{<\mathbb{N}}$ ,  $\mathbb{N}^{<\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}}$ ,  $\mathbb{N}^{<\mathbb{N}}$ ,  $\mathbb{N}^{<\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}}$ ,  $\mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}}$ ,  $\mathbb{N}^{\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}}$ ,  $\mathbb{N}^{\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}}$ ,  $\mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}}$ ,  $\psi$  is a closed continuous embedding of  $\phi_0$  into  $\phi$ .

*Proof.* If  $\phi \circ \overline{\pi}$  is constant, then set  $\phi_0 = c_{\mathbb{N}^N \setminus \mathbb{N}^N}$  and let  $\psi$  be the unique function from  $c_{\mathbb{N}^N \setminus \mathbb{N}^N}(\mathbb{N}^N \setminus \mathbb{N}^N)$  to  $(\phi \circ \overline{\pi})(\mathbb{N}^N \setminus \mathbb{N}^N)$ . If  $\phi \circ \overline{\pi}$  extends to a closed continuous embedding  $\psi$  on  $Z \in {\mathbb{N}^N \setminus \mathbb{N}^N, \mathbb{N}^N}$ , then set  $\phi_0 = \iota_{\mathbb{N}^N \setminus \mathbb{N}^N, \mathbb{Z}}$ . And if  $\phi \circ \overline{\pi} \circ p^{-1}$  extends to a closed continuous embedding  $\psi$  on  $Z \in {\mathbb{N}^{<N} \setminus \mathbb{N}^N, \mathbb{N}^{<N}}$ , then set  $\phi_0 = \iota_{\mathbb{N}^{<N}, \mathbb{N}^{<N}}$ ,  $\mathbb{N}^{<N} \setminus \mathbb{N}^N, \mathbb{N}^{<N}$ ,  $\mathbb{N}^{<N} \setminus \mathbb{N}^N, \mathbb{N}^{<N}$ ,  $\mathbb{N}^{<N} \setminus \mathbb{N}^N, \mathbb{N}^{<N}$ , then set  $\phi_0 = \iota_{\mathbb{N}^{<N}, \mathbb{Z}} \circ p$ .  $\boxtimes$ 

### 6. Borel functions that are not Baire class one

Here we provide bases for the classes of non-Baire-class-one Borel functions and non- $\sigma$ -continuous-with-closed-witnesses Borel functions between analytic metric spaces.

**Proposition 6.1.** Suppose that X is a metric space and  $\phi \colon \mathbb{N}^{\mathbb{N}}_* \to X$ has the property that  $\phi \upharpoonright \mathbb{N}^{\mathbb{N}}$  is continuous and  $\phi(\mathbb{N}^{\mathbb{N}}) \nsubseteq \overline{\phi(\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}})}$ . Then there is a  $\wedge$ -embedding  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  with the property that  $(\overline{\phi \circ \pi})(\mathbb{N}^{\mathbb{N}}) \cap (\overline{\phi \circ \pi})(\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}}) = \emptyset$ .

Proof. Fix  $b \in \mathbb{N}^{\mathbb{N}}$  for which  $\phi(b)$  is not in the closure of  $\phi(\mathbb{N}_*^{\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}})$ . Then there is an open neighborhood U of  $\phi(b)$  disjoint from  $\phi(\mathbb{N}_*^{\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}})$ , as well as an open neighborhood V of  $\phi(b)$  whose closure is contained in U, in which case the continuity of  $\phi \upharpoonright \mathbb{N}^{\mathbb{N}}$  yields a proper initial segment s of b for which  $\phi(\mathcal{N}_s \cap \mathbb{N}^{\mathbb{N}}) \subseteq V$ . Then the  $\wedge$ -embedding  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  given by  $\pi(t) = s \frown t$  for all  $t \in \mathbb{N}^{<\mathbb{N}}$  is as desired.

Given  $\phi_{\mathbb{N}^{\mathbb{N}}} \colon \mathbb{N}^{\mathbb{N}} \to X$  and  $\phi_{\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}}} \colon \mathbb{N}^{\mathbb{N}}_* \to Y$ , let  $\phi_{\mathbb{N}^{\mathbb{N}}} \sqcup \phi_{\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}}}$ denote the corresponding function from  $\mathbb{N}^{\mathbb{N}}_*$  to the disjoint union  $X \sqcup Y$ . **Theorem 6.2.** Suppose that X and Y are analytic metric spaces and  $\phi: X \to Y$  is a Borel function that is not Baire class one. Then there exist  $\phi_{\mathbb{N}^{\mathbb{N}}} \in \{c_{\mathbb{N}^{\mathbb{N}}}\} \cup \{\iota_{\mathbb{N}^{\mathbb{N}},Z} \mid Z \in \{\mathbb{N}^{\mathbb{N}}, \mathbb{N}^{\mathbb{N}}_{*}\}\}$  and  $\phi_{\mathbb{N}^{\mathbb{N}},\mathbb{N}^{\mathbb{N}}} \in \{c_{\mathbb{N}^{\mathbb{N}},\mathbb{N}^{\mathbb{N}}}\} \cup \{\iota_{\mathbb{N}^{\mathbb{N}},\mathbb{N}^{\mathbb{N}}}\} \in \{c_{\mathbb{N}^{\mathbb{N}},\mathbb{N}^{\mathbb{N}}}\} \cup \{\iota_{\mathbb{N}^{\mathbb{N}},\mathbb{N},Z} \circ p \mid Z \in \{\mathbb{N}^{<\mathbb{N}}, \mathbb{N}^{<\mathbb{N}}_{*}, \mathbb{N}^{<\mathbb{N}}_{*} \setminus \mathbb{N}^{\mathbb{N}}, \mathbb{N}^{<\mathbb{N}}_{*}\}\}$  for which there is a closed continuous embedding of  $\phi_{\mathbb{N}^{\mathbb{N}}} \sqcup \phi_{\mathbb{N}^{\mathbb{N}},\mathbb{N}^{\mathbb{N}}}$  into  $\phi$ .

Proof. Hurewicz's dichotomy theorem for  $F_{\sigma}$  sets yields a closed continuous embedding  $\psi \colon \mathbb{N}^{\mathbb{N}}_* \to X$  with  $(\phi \circ \psi)(\mathbb{N}^{\mathbb{N}}) \cap \overline{(\phi \circ \psi)(\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}})} = \emptyset$ . As  $(\psi, \operatorname{id}_{\overline{(\phi \circ \psi)(\mathbb{N}^{\mathbb{N}}_*)}})$  is a closed continuous embedding of  $\phi \circ \psi$  into  $\phi$ , by replacing the latter with the former, we can assume that  $X = \mathbb{N}^{\mathbb{N}}_*$  and  $\phi(\mathbb{N}^{\mathbb{N}}) \cap \overline{\phi(\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}})} = \emptyset$ .

By Proposition 3.1, there is a  $\wedge$ -embedding  $\pi \colon \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$  for which  $(\phi \circ \overline{\pi}) \upharpoonright \mathbb{N}^{\mathbb{N}}$  is continuous. By composing  $\pi$  with the  $\wedge$ -embedding given by Proposition 6.1, we can assume that  $(\phi \circ \overline{\pi})(\mathbb{N}^{\mathbb{N}}) \cap (\phi \circ \overline{\pi})(\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}}) = \emptyset$ . By composing  $\pi$  with the  $\wedge$ -embedding given by Theorem 3.6, we can assume that  $(\phi \circ \overline{\pi}) \upharpoonright \mathbb{N}^{\mathbb{N}}$  is constant or extends to a closed continuous embedding on  $\mathbb{N}^{\mathbb{N}}$  or  $\mathbb{N}^{\mathbb{N}}_*$ . And by composing  $\pi$  with the  $\wedge$ -embedding given by Theorem 5.5, we can assume that  $(\phi \circ \overline{\pi}) \upharpoonright \mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}}$  is constant,  $(\phi \circ \overline{\pi}) \upharpoonright \mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}}$  extends to a closed continuous embedding on  $\mathbb{N}^{\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}}$  or  $\mathbb{N}^{\mathbb{N}}_*$ , or  $\phi \circ \overline{\pi} \circ p^{-1}$  extends to a closed continuous embedding on  $\mathbb{N}^{<\mathbb{N}}_*$ ,  $\mathbb{N}^{<\mathbb{N}}_* \setminus \mathbb{N}^{\mathbb{N}}_*$ .

By Proposition 3.7, there exist  $\phi_{\mathbb{N}^{\mathbb{N}}} \in \{c_{\mathbb{N}^{\mathbb{N}}}\} \cup \{\iota_{\mathbb{N}^{\mathbb{N}},Z} \mid Z \in \{\mathbb{N}^{\mathbb{N}}, \mathbb{N}_{*}^{\mathbb{N}}\}\}$ and  $\psi_{\mathbb{N}^{\mathbb{N}}} : \overline{\phi_{\mathbb{N}^{\mathbb{N}}}(\mathbb{N}^{\mathbb{N}})} \to \overline{\phi(\mathbb{N}^{\mathbb{N}})}$  for which  $(\overline{\pi} \upharpoonright \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}, \psi_{\mathbb{N}^{\mathbb{N}}})$  is a closed continuous embedding of  $\phi_{\mathbb{N}^{\mathbb{N}}}$  into  $\phi \upharpoonright \mathbb{N}^{\mathbb{N}}$ . By Proposition 5.7, there exist  $\phi_{\mathbb{N}_{*}^{\mathbb{N}}/\mathbb{N}^{\mathbb{N}}} \in \{c_{\mathbb{N}_{*}^{\mathbb{N}}/\mathbb{N}^{\mathbb{N}}}\} \cup \{\iota_{\mathbb{N}_{*}^{\mathbb{N}}/\mathbb{N}^{\mathbb{N}}, Z} \mid Z \in \{\mathbb{N}_{*}^{\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}}, \mathbb{N}_{*}^{\mathbb{N}}\}\} \cup \{\iota_{\mathbb{N}^{<\mathbb{N}}, Z} \circ p \mid Z \in \{\mathbb{N}^{<\mathbb{N}}, \mathbb{N}_{*}^{<\mathbb{N}}, \mathbb{N}_{*}^{<\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}}, \mathbb{N}_{*}^{<\mathbb{N}}\}\}$  and  $\psi_{\mathbb{N}_{*}^{\mathbb{N}}/\mathbb{N}^{\mathbb{N}}} : \overline{\phi_{\mathbb{N}_{*}^{\mathbb{N}}/\mathbb{N}^{\mathbb{N}}}(\mathbb{N}_{*}^{\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}})} \to \overline{\phi(\mathbb{N}_{*}^{\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}}, \mathbb{N}_{*}^{<\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}}, \mathbb{N}^{<\mathbb{N}})}$  for which  $(\overline{\pi} \upharpoonright \mathbb{N}_{*}^{\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}} \to \mathbb{N}_{*}^{\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}}, \psi_{\mathbb{N}_{*}/\mathbb{N}^{\mathbb{N}}})$  is a closed continuous embedding of  $\phi_{\mathbb{N}_{*}^{\mathbb{N}}/\mathbb{N}^{\mathbb{N}}}$  into  $\phi \upharpoonright \mathbb{N}_{*}^{\mathbb{N}} \setminus \mathbb{N}^{\mathbb{N}}$ . Then  $(\overline{\pi} \upharpoonright \mathbb{N}_{*}^{\mathbb{N}} \to \mathbb{N}_{*}^{\mathbb{N}}, \psi_{\mathbb{N}_{*}/\mathbb{N}^{\mathbb{N}}})$  is a closed continuous embedding of  $\phi_{\mathbb{N}_{*}/\mathbb{N}^{\mathbb{N}}}$  into  $\phi$ .

Theorems 4.5 and 6.2 together provide the promised twenty-seven element basis under closed continuous embeddability for the class of non- $\sigma$ -continuous-with-closed-witnesses Borel functions between analytic metric spaces.

#### References

[CMS] R. Carroy, B.D. Miller, and D.T. Soukup, *The open dihypergraph dichotomy* and the second level of the Borel hierarchy, to appear in Contemporary Mathematics.

- [JR82] J.E. Jayne and C.A. Rogers, First level Borel functions and isomorphisms, J. Math. Pures Appl. (9) 61 (1982), no. 2, 177–205. MR 673304 (84a:54072)
- [Kec95] A.S. Kechris, Classical descriptive set theory, Graduate Texts in Mathematics, vol. 156, Springer-Verlag, New York, 1995. MR 1321597 (96e:03057)
- [Sol98] S. Solecki, Decomposing Borel sets and functions and the structure of Baire class 1 functions, J. Amer. Math. Soc. 11 (1998), no. 3, 521–550. MR 1606843 (99h:26010)

RAPHAËL CARROY, KURT GÖDEL RESEARCH CENTER FOR MATHEMATICAL LOGIC, UNIVERSITÄT WIEN, WÄHRINGER STRASSE 25, 1090 WIEN, AUSTRIA *E-mail address*: raphael.carroy@univie.ac.at

URL: http://www.logique.jussieu.fr/~carroy/indexeng.html

BENJAMIN D. MILLER, KURT GÖDEL RESEARCH CENTER FOR MATHEMATI-CAL LOGIC, UNIVERSITÄT WIEN, WÄHRINGER STRASSE 25, 1090 WIEN, AUS-TRIA

*E-mail address*: benjamin.miller@univie.ac.at

URL: http://www.logic.univie.ac.at/benjamin.miller