

This is a repository copy of *The data-driven future of high-energy-density physics*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/182911/>

Version: Accepted Version

Article:

Hatfield, Peter W., Gaffney, Jim A., Anderson, Gemma J. et al. (20 more authors) (2021) The data-driven future of high-energy-density physics. *Nature*. pp. 351-361. ISSN 1476-4687

<https://doi.org/10.1038/s41586-021-03382-w>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

The data-driven future of high energy density physics

Peter W. Hatfield,^{1, a)} Jim A. Gaffney,^{2, b)} Gemma J. Anderson,^{2, c)} Suzanne Ali,² Luca Antonelli,³ Suzan Başğmez du Pree,⁴ Jonathan Citrin,⁵ Marta Fajardo,⁶ Patrick Knapp,⁷ Brendan Kettle,⁸ Bogdan Kustowski,² Michael J. MacDonald,² Derek Mariscal,² Madison E. Martin,² Taisuke Nagayama,⁷ Charlotte A.J. Palmer,⁹ J. L. Peterson,² Steven Rose,^{1, 8} JJ Ruby,¹⁰ Carl Shneider,¹¹ Matt J.V. Streeter,⁸ Will Trickey,³ and Ben Williams¹²

¹⁾Clarendon Laboratory, University of Oxford, Parks Road, Oxford OX1 3PU, UK

²⁾Lawrence Livermore National Laboratory, 7000 East Ave, Livermore, CA 94550, USA

³⁾York Plasma Institute, Department of Physics, University of York, Heslington, York YO10 5DD, UK

⁴⁾Nikhef, National Institute for Subatomic Physics, Science Park 105, 1098 XG, Amsterdam, Netherlands

⁵⁾DIFFER - Dutch Institute for Fundamental Energy Research, Eindhoven, Netherlands

⁶⁾Instituto de Plasmas e Fusão Nuclear, Instituto Superior Técnico, 1049-001 Lisboa, Portugal

⁷⁾Sandia National Laboratories, 1515 Eubank SE, Albuquerque, New Mexico 87185-1196, USA

⁸⁾Imperial College London, London, SW7 2AZ, UK

⁹⁾School of Mathematics and Physics, Queen's University Belfast, Belfast, BT7 1NN, UK

¹⁰⁾Laboratory for Laser Energetics, University of Rochester, Rochester, NY, USA

¹¹⁾Dutch National Center for Mathematics and Computer Science (CWI), Science Park 123, 1098 XG, Amsterdam, Netherlands

¹²⁾AWE Plc, Aldermaston, Reading, RG4 7PR, United Kingdom

(Dated: 19 March 2021)

PREFACE

The study of plasma physics under conditions of extreme temperatures, densities and electromagnetic field strengths is significant for our understanding of astrophysics, nuclear fusion and fundamental physics. These extreme physical systems are strongly non-linear and very difficult to understand theoretically or optimize experimentally. Here, we argue that machine learning models and data-driven methods are in the process of reshaping our exploration of these extreme systems that have hitherto proven far too non-linear for human researchers. From a fundamental perspective, our understanding can be helped by the way in which machine learning models can rapidly discover complex interactions in large data sets. From a practical point of view, the newest generation of extreme physics facilities can perform experiments multiple times a second (as opposed to \sim daily) – moving away from human-based control towards automatic control based on real-time interpretation of diagnostic data and updates of the physics model. To make the most of these emerging opportunities, we advance proposals for the community in terms of research design, training, best practices, and support for synthetic diagnostics and data analysis.

‘Set the controls for the heart of the Sun’ encouraged a 2004 paper¹ (riffing on the 1968 Pink Floyd song), describing the bright future of using Earth based experiments to create conditions similar to inside the Sun in the lab. Seventeen years later substantial advances have been made in this research programme. A question that is presently emerging however is *who* should be at the controls - humans, or artificial intelligences?

In the last few years plasma physics has been steadily beginning to explore the use of modern day data science and artificial intelligence methods to support research goals^{2,3}. In this article we will identify data science issues for physics specifically at the extreme⁴; extremely high temperatures, densities or electromagnetic field strengths - which have unique challenges. In particular phenomena at these conditions are highly *non-linear* - small parameter changes can lead to large changes in behaviour. Interpreting extreme physics data typically requires simultaneously comprehending large amounts of complex multi-modal data from multiple different sources. Optimising extreme physics systems requires fine-tuning over large numbers of (often highly correlated) parameters. Artificial Intelligence (AI) methods have proved highly successful at teasing out correlations in large data sets like these and we believe will be crucial for understanding and optimising systems that up to now have been inscrutable. These extreme conditions can be found in astrophysical scenarios, but can also be created using high-energy ‘drivers’ (often lasers) in the laboratory - millimetre sized plasmas with temperatures and pressures higher than the centre of the Sun. The field has seen an explosion of interest in machine learning techniques because new and future laser facilities have much higher shot (and corresponding data) rates than previous facilities. Data

^{a)}Electronic mail: peter.hatfield@physics.ox.ac.uk

^{b)}Electronic mail: gaffney3@llnl.gov

^{c)}Electronic mail: anderson276@llnl.gov

iments is moving beyond the limits of conventional methods. The quantity of experimental data is increasing both because shot rates on facilities are dramatically increasing (see Figure 1), but also because diagnostics are becoming more sophisticated; $\sim 150\text{GB}$ of data is taken on each NIF shot, and LCLS campaigns have reached $\sim 70\text{GB}$ per minute³⁵. Machine learning³⁶, Bayesian methods³⁷, and data-driven science³⁸ have been common in particle physics and astrophysics for many years, and are having large impacts on other multiscale, highly non-linear areas of physics e.g. climate science and Earth system science^{39,40}. Some AI solutions from other fields are likely to be applicable in plasma physics - but HEDP also has its own unique challenges. Specifically we typically want to (sometimes very rapidly) fine tune (either optimising or fitting a model) a large number of parameters for a desired outcome, based on a large number of multi-modal data sets. This is very difficult for humans, as it is very hard to simultaneously comprehend all the different sources and forms of data - but is achievable for AI. In HEDP, the highest performing experiments in the field are now increasingly data driven^{41,42}; the vision is to work towards a HEDP science where algorithms are at the centre of design, analysis of experiments, and discovery.

In the following sections we review key challenges and topics in extreme plasma physics, and highlight research areas where data science has dramatically impacted the field. We further lay out what the future might hold for data driven Extreme Physics and HEDP - and how the community must change and adapt how it practices research to make the most of these exciting new approaches.

CHALLENGES

The qualities that make HEDP an exciting area of research also contribute to causing the challenges inherent in any quantitative analysis of data. Table I summarises some key quantitative methods, comparing conventional and emerging approaches, and in the following sub-sections we discuss three primary challenges that data science techniques can help address in HEDP.

Experimental Design and Automation

A key component of HEDP is experimentation. Designing experiments is a hugely complex task, and researchers typically will have a wide range of overlapping goals during this phase. Researchers must consider what specific hypotheses are to be tested, and if the expected data would be sufficient to rule out alternatives. What design to shoot, diagnostic instruments to field, or astronomical observation to make will depend heavily on the specific science goal at hand. In current design approaches there is typically much intuition implicitly at play, and often experiments are built as an extension of what has been done before, limiting the regions of experimental space that are studied. Machine learning techniques offer a possible framework for the intuition of the computa-

tional scientists and experimental scientists to be explicitly included in a cohesive picture that considers both measurements that can be made, and which aspects of physics have the most leverage on those measurements. AI-aided design is beginning to be used in the creation of new HEDP experiments^{43,44}, and we foresee that becoming the norm in the coming years. However, machine learning methods have yet to demonstrate that they can “think outside the box” of pre-defined parameter spaces, therefore for the foreseeable future it will still be necessary to have substantial human input in the design process.

Experiments on state-of-the-art high repetition rate lasers, firing many times a second, cannot be done with a human in the loop - so in this case some algorithmic control is essential. This in principle could lead to huge savings of time, money, and human effort in the near future - allowing them to be redirected to aspects of research where they can be better used. Automating experiments combines control of experimental parameters and real-time analysis of experimental results into a single algorithm. The experimental goals are coded in to the automation, such that choices of how to vary the experimental inputs are made automatically in order to maximise the output.

Automation also allows for active feedback stabilisation of complex processes. This is of particular benefit to HEDP experiments, where many non-linear effects combine to determine the performance of what is a nominally unpredictable process. In this context, AI could be the solution to an otherwise intractable problem; the well-known ability of deep learning models to discover complex interactions⁴⁵ could be used to optimize systems that have proven far too non-linear for human researchers.

Data Synthesis

The measurements made in HEDP experiments are often highly integrated; experiments typically don’t measure the actual quantity of interest (QoI), and there are usually multiple confounding or nuisance variables that have to be controlled. Isolating a particular aspect of these systems is often not possible, resulting in a measurement of an evolving system subject to different conditions and physical processes. This complexity makes repeatability an issue. In order to isolate individual aspects of the underlying physics, experiments therefore typically require multiple, indirect observations, sometimes spread across several different experimental facilities. At most facilities researchers have developed multiple diagnostics for experiments; for example both x-ray and particle spectra may be measured on a single experiment, along with many other forms of experimental data - all of which might contribute to the determination of a single quantity. The analysis of such increasingly sophisticated interlinked data requires the use of more advanced modelling techniques to make the best use of the available data, and to sensibly quantify the uncertainty on any inferences. Looking forward, large quantities of data will require the development of streamlined automated data analysis tools to avoid read/write bottlenecks⁴⁶.

As well as combining data from multiple diagnostics, there

Analytical Task	Scientific Task	Conventional approaches	Limitations of conventional approaches	Emergent or potential approaches
Uncertainty Quantification	Quantifying uncertainty on estimate of some microphysics (e.g. opacity, equation-of-state), characterising uncertainty estimate on laser energy required for ignition	Simulation-based sensitivity studies, basic Poisson/Gaussian uncertainties, not including correlations	Local sensitivities only, relies on estimated uncertainties in underlying parameters, cannot account for simulation bias	MCMC, dropout, bootstrapping, quantification of ‘unknown unknowns’, invertible neural networks
Regression	Emulating an expensive simulation, building an empirical model of microphysics	Look-up tables, polynomial regression	Struggles in high dimension, incorporating known physics constraints, extrapolation is difficult, often not fast enough	Neural networks, Gaussian processes, autoencoders
Design	Selecting target design parameters, scheduling observation runs	Adjusting parameters by hand, using a combination of code output and Designer judgement	Hard in high dimensions, human intensive, cannot be done quickly, can miss optimal/novel designs	Bayesian optimisation, Genetic/Evolutionary algorithms
Pattern Recognition	Identifying target defects, characterising magnetic perturbations, image segmentation/featurization	Human inspection	Laborious and time consuming, subject to individual biases, uncontrolled approximation to true information content	Convolutional Neural Networks, Deep learning, random forest classifiers, Human-AI hybrid approaches, Application Specific Integrated Circuits incorporated into diagnostics, Generative adversarial networks
Data Synthesis	Combining data from multiple sources (e.g. different instruments)	Researcher guided inference from independently analysed diagnostic data	Difficult to do ‘by hand’, hard to take advantages of any degeneracies broken	Bayesian inference, data assimilation methods
Classification	Image classification, particle track classification, identify good/bad shots, anomaly detection	Simple analytic criteria, human inspection	Laborious and time consuming, potentially inaccurate	Random forest, decision trees, neural networks, deep learning, Generative adversarial networks
Model Calibration	Update physics models/parameters in the face of experimental data	Trial and error, single-point fitting to data, hand-tuning	Laborious and time consuming, inaccurate or missing treatment of uncertainties, miss multiple solutions, prone to overfitting	Bayesian inference, discrepancy modeling, transfer learning, multi-task learning, physics informed neural networks

TABLE I. Conventional and AI-enabled approaches to tasks in extreme plasma physics

are also challenges in combining data from multiple sources; ‘data synthesis’, where multiple forms of heterogeneous data are combined in a self-consistent manner in order to construct a more complete picture of the phenomenon of interest. Typically each diagnostic will be analysed separately, and overall conclusions reached heuristically by the researcher. However, combining all available data can help reduce error bars, break degeneracies, and cancel out uncorrelated systematics⁴⁷.

The long term vision for the best use of the physics data is to develop systems to combine data from multiple diagnostics on the same shot, multiple shots, shots on different facilities, and finally from different types of facility.

Physics models

The evolution of HEDP experiments is governed by multiple complex, non-linear physics models, each of which has

its own range of applicability and uncertainties. Solving these multiphysics computer models requires very expensive simulations, which are often not suited to next-generation HPC platforms. With the increasing desire to explore larger experimental parameter spaces, there is a shifting dynamic between single, incredibly computationally expensive, ‘hero’ simulations, and large-scale ensembles of simulations which only become meaningful when confronted with experimental data.

Due to numerical approximations, poorly known or unknown model parameters, and missing physics (*model discrepancy*), computer models often do not accurately represent the physical process under study. We can leverage real-world experiments to calibrate our computational models, enabling us to constrain some of the uncertain model parameters; ideally including an Uncertainty Quantification (UQ) analysis⁴⁸ and practicing ‘data assimilation’⁴⁹ that obeys physical laws. A useful approach to this problem is through *Bayesian inversion* (also known as model calibration)^{50,51}, which allows

prior knowledge to be included and for which convenient numerical tools exist. In a HEDP context, challenges with this approach include large parameter spaces, expensive models, and very sparse experimental data.

The computer models can often be prohibitively computationally expensive. In this case a *surrogate model* (or *emulator*) may be useful - running a moderate number of expensive simulations, and training a machine learning algorithm to reproduce what the simulation would have given as an output. Surrogate models are of course themselves only an approximation of the true model, introducing further uncertainty that needs to be accounted for.

Emulation can be done at the macro level (e.g. predicting outputs of a whole experiment), or at the level of individual modules run inline inside of a computer model. The use of emulators opens up an array of new inference methods that would not be practical with the full computational expense of a conventional simulation^{52–58}.

CASE STUDIES

Here we highlight three key areas where researchers are tackling the challenges described in Section II, and where data science is significantly impacting the practice of extreme plasma science.

Astrophysics

Plasmas are found throughout the Universe: in Solar physics (the centre of the Sun, the solar corona, solar wind); interplanetary, interstellar and intergalactic media; in Earth's and other planetary magnetospheres and ionospheres, and tails of comets; in compact astrophysical objects (white dwarfs, neutron stars, and black holes) and their accretion disks. After direct matter-antimatter annihilation, accretion onto compact objects is the most efficient energy source in the Universe⁵⁹; the Cosmos offers plenty of opportunity to probe extreme physics.

Understanding our closest star is of course of supreme practical importance - alongside curiosity-driven blue sky astrophysical motivation. Space weather refers to conditions on the Sun, in the heliosphere, in the solar wind, and in Earth's magnetosphere, ionosphere, and thermosphere, that can influence the performance and reliability of space-borne and ground-based technological systems and can endanger human life or health^{60,61}. Changes in the space environment, resulting mainly from changes on the Sun, include modification of the ambient plasma, particulate radiation (electrons, protons and ions), electromagnetic radiation (including radio, visible, UV, X-ray, and gamma radiation), and magnetic and electric fields.

As with many other areas of physics, the amount of data we have on the multi-scale complex physics experiment that is the Sun has massively increased in recent years from Solar space missions e.g. from both spacecraft, such as the Solar and Heliospheric Observatory (SOHO), the Hinode mis-

sion, the Solar Dynamics Observatory (SDO), the Parker Solar Probe (PSP) and Solar Orbiter (SolO), and CubeSats, such as the Miniature X-ray Solar Spectrometer (MinXSS). Machine learning has been employed successfully to both forecast and 'now-cast' space weather⁶². It has been used to make predictions and gain insight about: solar wind⁶³, solar flares^{64,65}, coronal mass ejections (CMEs)^{66,67}, Van Allen radiation belts⁶⁸, geomagnetically induced currents⁶⁹ and the role of auroras as proxy for ionospheric disturbances⁷⁰.

Outside our Solar System, large survey telescopes are taking huge amounts of data on astronomical bodies with extreme physics. A full range of data science based techniques are used to both a) identify objects of interest in the large data sets⁷¹, and b) understand their underlying physics. Machine learning and statistical methods have been used to: make a Bayesian constraint on supra-nuclear equations-of-state⁷², understand the interiors of exoplanets⁷³, constrain fundamental stellar parameters from asteroseismic observations⁷⁴, classify supernovae⁷⁵, identify and infer the properties of white dwarfs⁷⁶, classify states of black hole X-ray binaries⁷⁷, and emulate radiative transfer during the epoch of cosmological reionization⁷⁸.

HEDP experimental facilities can, as discussed, also probe the extreme plasma physics relevant in astrophysical bodies. For example, experiments have measured the equation of state at conditions relevant for the centres of gas giant exoplanets^{79,80}, tested theories on possible origins of magnetic fields on galactic scales⁸¹ and helped our understanding of white dwarf photosphere spectra⁸². There has recently been great success in applying data science methods to these experiments and making inferences with realistic uncertainties. Measurements of iron opacity at conditions relevant to solar physics on the Z facility for example have modelled a large number of possible sources of uncertainty and systematics, and have combined the data from multiple shots together to get realistic uncertainties. This measurement was statistically inconsistent with conventional opacity predictions and has potentially led to an adjustment of estimates of the metallicity (lithium and higher atomic number elements) of the Sun^{83,84}. Similarly, Bayesian estimates from experiment of nuclear reaction rate at conditions relevant for Big Bang nucleosynthesis were found to be 3% different from conventionally assumed; a level of precision needed for cosmological studies⁸⁵. As discussed in the introduction, future facilities will have shot rates that will make multi-shot studies like this the norm. Critically this will make it possible to consistently make realistic uncertainty quantifications of key parameters, and will also give the ability to probe large parts of parameter space (e.g. measure the equation-of-state or opacity at a large number of points in temperature-density space). With these vast quantities of observational and experimental data, the natural next step is to use experimentally calibrated models of microphysics like these in astrophysical models - with the potential to give improved predictions over theory-only models.

There is a huge amount of data on astrophysical plasmas of a wide variety of sources, taken in a huge variety of ways, but unfortunately they are currently held in different forms, by different communities. See Figure 2 for an infographic on




FIG. 2. **Integration of astrophysical information:** combining data from multiple sources. At the centre is a machine learning algorithm that is receiving data from multiple sources, and is able to update its beliefs based on this ('data assimilation'). The resulting data-driven results will take information from both theory, and data, to give more precise predictions, with realistic uncertainties.

how different astrophysical data sets could in future be combined in a data assimilation framework^{86,87}. Astronomers have long practiced '*multi-wavelength*' astronomy (astronomy observing at multiple electromagnetic wavelengths), and since ~ 2015 have practiced '*multi-messenger*' astronomy (astronomy taking data from multiple messengers of information, electromagnetic waves, gravitational waves, cosmic rays and neutrinos). It is now time for '*multi-provenance*' astronomy; integrating data from both observations and experiments into one coherent model of our understanding of astrophysical sources.

Inertial Confinement Fusion

The aim of igniting a self-propagating hydrogen fusion reaction in laboratory-scale plasmas has been a scientific grand challenge for over 50 years. The motivation is clear: a reliable fusion reaction could form the basis for an effectively limitless, clean, and safe energy source⁸⁸. There are good reasons the research has been long-running; generating densities and temperatures high enough to overcome the Coulomb repulsion between nuclei and Bremsstrahlung radiation losses from electrons, over timescales long enough to allow energy break-even, is extremely difficult. In nature, the required conditions are reached in the cores of stars as a result of gravitational collapse and confinement. On Earth, we require even more extreme conditions to account for the much shorter time scales.

In inertial confinement fusion (ICF) studies, millimetre-scale targets are driven to implode using high energy (100s kJ to MJ-class) drivers⁸⁹. For fusion ignition, the implosion and

subsequent stagnation needs to generate a high temperature (10^7 - 10^8 K) hydrogen plasma, surrounded by a compressed ($100\times - 1000\times$) fuel layer, confined by its own inward velocity for several hundred picoseconds⁹⁰. Experiments on current ICF facilities like the National Ignition Facility (NIF)^{91,92}, the Omega Laser Facility⁹³, and the Z pulsed-power machine^{94,95} routinely generate plasmas under solar core conditions and beyond.

ICF experiments present distinct data challenges due to their scale and complexity. Experimental facilities are expensive and are not expected to achieve high repetition-rate operation any time soon. Targets and drivers are very complex resulting in high-dimensional experimental design spaces. Experiments are also highly *integrated*, meaning direct measurement of any figure of merit beyond the raw energy yield is not possible. These factors mean that ICF datasets are always sparse, with multiple confounding factors and uncertain information content; researchers therefore place a very high value on theoretical studies undertaken using multiphysics simulation codes. While cheaper than experiments, the simulations are still expensive, requiring at least CPU-months to complete. They can also have significant bias⁹⁶, and therefore require calibration against the available experimental data (from both ICF, and smaller scale experiments focused on relevant phenomena). There is a significant need for new methods that can help with experimental design and optimization, interpretation of experimental data, linking experiments with physics models, as well as making reliable predictions of future experiments. As this work progresses, ICF is becoming a prototypical example of the difficulties associated with science in the data-poor regime.

As with the other examples in this perspective, the funda-

mental data problem is the synthesis of multiple sources of information. Here, the key aim is to efficiently use the sparse information available to update our physics understanding in order to make simulations more predictive of future experiments. Figure 3 shows a potential workflow that fully integrates data-driven and machine-learning methods to achieve this goal. There are three fundamental information sources; experimental data which may comprise ~ 10 ‘shots’ each producing multiple diagnostics with diverse data types; traditional high fidelity simulations which produce single best-physics predictions for each shot; and high-volume ensemble studies which use large numbers of (necessarily lower-fidelity) simulations to investigate competing physics hypothesis and provide protection against overfitting. The optimization and control of these information sources, combination of the resulting data, and updating of models to become more predictive present numerous opportunities for modern data science and machine learning approaches and each stage has seen recent active research.

Work to optimize the three information sources in Figure 3 include the acceleration of multiphysics simulations using deep learning^{55,97}, infrastructure for intelligent control of large-scale simulations⁹⁸, intelligent and data-informed design of experiments^{42,43,99}, as well as optimization of experimental facilities¹⁰⁰. These developments have enabled simulation studies of unprecedented size¹⁰¹ and the generation of open-source ICF datasets¹⁰² that motivate novel deep learning research^{54,57,103,104}. AI tools have been applied to the automatic analysis and featurization of complex data types like spectra, images^{54,105}, and line-of-sight dependent quantities. There has been significant interest in using Bayesian inference to improve diagnostics¹⁰⁶, and to synthesize observations in both focused HEDP experiments⁵¹ and full-scale ICF experiments^{50,107–109}. The ultimate aims of using these methods to improve physics understanding, and the reliability of simulations in extrapolating to new designs⁴⁸ or facilities, have been addressed through machine learning^{42,110}, Bayesian model calibration⁵⁰, and transfer learning^{58,111}.

Integrating the recent work we have described into a fully developed workflow similar to Figure 3 is still a significant challenge. Once achieved, however, we expect to gain an unprecedented view of the ICF design space, significantly better understanding of the conditions in current ICF experiments, and an improved understanding of the path to high yield and ignition.

Automation for High Repetition Rates

At the other end of the spectrum to NIF, which can only fire roughly once a day, are high-repetition rate lasers that can fire up to multiple times a second. This means huge amounts of data towards given science goals, but also means large aspects of the experimental process must be automated. To succeed, the automation of experiments requires both control of experimental parameters *and* real-time analysis of experimental results in one single algorithmic process; see Figure 4.

The first step in achieving this is for experimental goals to

be coded in to the automation algorithm, such that choices of how to vary the experimental inputs are made automatically in order to maximise the desired output. This approach enables huge increases in efficiency in optimisation and model learning experiments, especially important when these experiments are resource limited. In addition, automation also allows for active feedback stabilisation of complex processes; this is of particular benefit to high energy density and plasma physics experiments, where many non-linear (and also often non-equilibrium) effects combine to determine the performance of what is a nominally unpredictable process. Thought must also be given to other factors that will determine optimal data return; diagnostics used, signal-to-noise ratio achieved, and experimental parameter space covered. Many diagnostics have already been adapted for fast electronic readout¹¹², however, challenges still remain in adapting x-ray or charged particle imagers and similar detectors for both maximum flexibility, as well as robustness to the harsh radiation and debris environments of HEDP experiments - especially at multi-Hz repetition rates.

This approach may be used to perform the following tasks:

- *Optimisation*: A function of experimental diagnostics is used to calculate the ‘fitness’ which expresses how closely measurements reflect the desired performance. An iterative procedure is then performed to optimise this fitness value by controlling experiment input parameters. This can be done with any optimisation procedure e.g. an evolutionary approach (i.e. genetic algorithms^{41,113,114}), a numerical minimisation method (e.g. Nelder-Mead¹¹⁴) or by Bayesian optimisation using a machine learned surrogate model^{115,116}. This approach can also be used to limit the parameter search to satisfy some safety constraint, e.g. beam loss in a particle accelerator¹¹⁵. In general applying these approaches allows for rapid optimisation of experiments in a far more efficient manner than human controlled experiments - and produces much better results.
- *Stabilisation*: Active feedback can improve the stability of experimental performance by rapidly controlling input parameters to counteract oscillation or drifts in the apparatus¹¹⁷. This is routinely performed to stabilise component systems, such as alignment of laser beam transport, but can also be applied to highly complex and non-linear experimental phenomena, such as density limit disruptions in tokamaks using the predictions of neural network^{118–120}. A stable output source then allows for much better experimental or source application outcomes.
- *Model Inference*: A Bayesian approach to statistical inference and model validation requires incorporating experiment uncertainties from diagnostic data in a rigorous manner, accounting for correlations across all parameter spaces⁵⁰. This can lead not only to better estimates of the uncertainty, but the results of the inference can also dramatically change. Including this approach in real time in the data taking process can be




FIG. 3. **Integrating Information Sources in ICF Studies:** Our understanding of ICF implosion physics is based on a combination of high-volume, lower-fidelity simulation ensembles; sparse, difficult-to-diagnose experiments; and best-physics simulations that push the limits of high-performance computing technology. Creating and synthesizing these data into an improved understanding of the physics will require multiple new, complementary techniques from data science, uncertainty quantification (UQ) and artificial intelligence.

used to ensure that the experiment optimally constrains the physical models under examination - getting the most *information* per shot (as opposed to simply the most neutrons/x-rays etc.).

Encoding the scientific goals into the algorithm making the shot choice is just one of many challenges in automating high repetition rate facilities. Not only the physics of interest, but also the laser system and the target setup typically have inherent non-linearities that can make automating knowledge extraction extremely challenging. Small changes in system parameters (laser pulse width, shape, energy, focal spot conditions, target thickness etc.) can lead to large changes in experimental outcomes - requiring very fine control of the entire system. Thus the laser itself and the target must be modelled alongside the physics of interest. This complex multi-modal data must also be analysed as fast as the shot-rate to prevent another bottleneck in the experimental loop. In addition performance of diagnostics themselves might be impaired over time (for example if exposed to large radiation fluxes), requiring further modelling. The goal is for the AI to understand the effect of these diagnosed (and potentially undiagnosed) fluctuations in the system, rather than be confused by it. Human intuition risks misinterpreting evidence when many parameters are changing simultaneously. Finally, data archiving from experiments will rapidly become a challenge. Significant challenges in developing pipelines that can prevent data bottlenecks will become important, i.e. the operating algorithms may have to decide whether to record or destroy data

based on quality of inputs and outputs to avoid large amounts of spurious and insignificant data occupying many TB of storage systems. In summary there are two separate challenges that should not be conflated. There is both a) the technical challenge of delivering online feedback and real time data curation and b) the modelling problem of automating knowledge extraction from the complex HEDP data. Researchers should take care to identify what aspects of their specific scientific problem fit into these two categories, and seek appropriate solutions.

The computerised control of experimental parameters increases the convenience for the experimental operators, allows for automatic parameter scans and reduces the likelihood of experimental errors. Enabling experiment automation requires considerable additional investment and effort in the preparation of experimental apparatus and facilities. However, time and resources spent on this endeavour yield large returns once the experiment is fully operational due to the increase in efficiency and productivity.

POLICY PROPOSITIONS

To help the field take advantage of the aforementioned new ways of using data, we make a series of suggestions for how educational and research practices might be beneficially changed.




FIG. 4. **High repetition workflow:** the different components of a series of high repetition rate high powered laser. The AI system must have i) a model of its best estimate, with uncertainties, of the physics being probed, ii) a model of the current state of the laser (both so it can better achieve science goals, but also to avoid damaging itself), iii) a model of the target, iv) an algorithm to rapidly select what the next shot must be (depending on science goal), v) a system to actually fire the shot with no human intervention, vi) rapid automated data collection, vii) rapid physics, laser and target modelling, and viii) capacity to update a model of diagnostic performance (if needed).

Education

With the changes in HEDP data acquisition rates and analysis that have been described in this article, it is important to consider whether researchers who are new to the subject have sufficient training in the topics we have discussed.

There are courses in most universities that attempt to teach aspects of the subject; however they are not typically included in the HEDP curriculum. At the time of the writing, only 3 of the 40 Plasma Physics PhD Projects, Programs and Scholarships listed on www.findaphd.com contain a data science component.

While the subject is easily within reach for plasma physics graduate students, the unfamiliarity with the Bayesian perspective, and the necessary jargon, render it more difficult than necessary for a self-directed approach. We recommend that, *a minima*, a brief introduction^{36–38} be included in one of the Advanced Courses, with the opportunity to take an elective course on Bayesian analysis and Uncertainty Quantification. Ideally, a Bayesian analysis module should be included in future HEDP doctoral programmes, built from the cases presented in this Perspective. In addition to including data science in the general curriculum, existing certificate programs focused on both fundamental ML techniques and multi-disciplinary applications of data science could be leveraged by students and professionals alike. Conferences and workshops like the International Conference on Data Driven Plasma Science (which had its second meeting in 2019) and the 2018 APS Mini-Conference on Machine Learning, Data Science, and Artificial Intelligence in Plasma Research³ have flourished in the last few years, and these opportunities should be extended to those studying/training.

For all other graduate students and researchers, a complementary approach should be taken, through targeted work-

shops and schools. We believe that because the field is developing very quickly, it is preferable to consider teaching these topics as part of the existing HEDP workshops that have been established over the last few years for those new to the area. Topics should include Bayesian and frequentist statistics comparisons, UQ, Markov chain Monte Carlo (MCMC), surrogate building, deep neural nets, and optimization techniques for a range of dimensional spaces. Such workshops and meetings may also provide opportunities to engage the general ML community. The HEDP community could also entice ML researchers to collaborate in our field by presenting or organizing focused sessions at ML conferences, releasing datasets, exploring a ML challenge call focused on an application in our field, and pursuing direct research collaborations. Finally we would note that data science skill sets have broad application both in other research areas, as well as in industry. The proposed training is highly transferable and promises to be valuable to students regardless of specific career goals.

Research Practices

The changing nature of the field means practices within the field will also have to change. Researchers will need to become familiar with methods needed to run large numbers of simulations, and tools for storing larger amounts of data⁴⁶. In particular, the field will have to develop data standards so that data is easily compatible between different facilities. Adopting open data practices¹⁰² (like F.A.I.R., findability, accessibility, interoperability, and reusability¹²²) wherever possible is also likely to greatly aid collaboration and comparison of data sets, although this will not be possible for all researchers in this area. The importance of these approaches is already well understood by researchers in other fields, for example high en-

ergy physics¹²³ and astronomy¹²⁴; we foresee a similar level of data curation will soon be necessary in HEDP.

Many of the methods described in this paper require significant computational resources and good synthetic diagnostics, i.e., good simulations of what the data should look like through the actual pipeline the real data goes through. While most research efforts include a computational component, these are often disjoint from the analysis of experimental data. In future it is advisable for experimental time to also have associated funding for computation and analysis, and the development of synthetic diagnostics/data analysis. Diagnostics and experiments should be designed so that collected data can easily be used in conjunction with other shots (e.g. consistent pixel sizes). It may even be necessary that the commissioning of instruments comes with a corresponding budget to develop tools to simulate data as seen by the device. Data analysis is a core component of project commissioning and planning in many other areas of physics (e.g. the Euclid telescope¹²⁵).

High data rates are permitting the probing of low signal-to-noise phenomena e.g. possible beyond Standard Model physics like axions. If laser based accelerators are to play a role in the future of probing new physics in this way, then statistical analysis must be brought up to the same standards as are practiced in High Energy Physics (HEP). In particular we may wish to adopt stringent statistical significance requirements e.g. 5σ for any discovery of beyond Standard Model physics¹²⁶. Similarly using blinding methods are likely to become more necessary, where researchers deliberately hide some aspect of the labelling of the data from themselves to prevent subconscious bias or p-hacking¹²⁷. Using a blinding protocol does also present dangers and challenges; the complexity and bespoke properties common in many laser-plasma experiments make blinding difficult to implement. However, high-rep rates will make blinding strategies much more viable, and may help for analyses where particular outcomes have high psychological significance and/or there are lots of different potentially viable approaches to doing the analysis.

CONCLUSION

The volume of data on plasma physics at the extremes is rapidly growing and offers the potential for dramatically increased speed of scientific advance. These data are typically multi-modal and describe very non-linear systems, making interpretation challenging for humans, but tractable for AI algorithms. Plasma physics is unlikely to reach the extremely high data rates of HEP in the immediate future, and it is fully the case that AI will not solve all problems in the field. Nonetheless, this new approach in the field offers novel ways of working, and new ways to gain insight - we hope practitioners in the field will be able to find applicability for these methods in their research, see Box 1.

BOX 1

Key conclusions:

1. The application of machine learning and modern data science methods to extreme plasma physics and HEDP is rapidly growing and is aiding in producing realistic uncertainties on predictions
2. Higher repetition rate facilities open up a range of new ways of working; data driven discovery, blinding methods, greater reproducibility, automated data taking
3. Integrating machine learning based approaches into working practices can greatly save money, time, and human effort
4. AI based tools are now often more successful at optimising non-linear extreme physics systems and comprehending multi-modal data than humans

Key recommendations:

1. Researchers should think carefully about how to best use their data: what methods and diagnostics can they use to take the best data, get sensible uncertainties, and coherently combine with other data sets
2. Awards of experimental time and instrument construction should also include greater support for uncertainty quantification, building synthetic diagnostics and data analysis
3. Plasma physics graduate education and national lab training programmes begin to include basic data science courses
4. Researchers should try where possible to practice open science best practice; making code and data available publicly, using shared data standards between different facilities

In conclusion, modern data science has a lot to offer extreme plasma physics and HEDP science; the community must act now to identify which areas it will make the biggest impact in and make resources and training available to make the most of these novel approaches.

ACKNOWLEDGMENTS

This Perspective was the result of a meeting at the Lorentz Center, University of Leiden, 13th-17th January 2020. The Lorentz Centre is funded by the Dutch Research Council (NWO) and the University of Leiden. The meeting also had support from the John Fell Oxford University Press (OUP) Research Fund. The organisers are grateful to Tanja Uitbeij-

jerse (Lorentz Center) for facilitating a fruitful and productive meeting.

P.W.H. acknowledges funding from the Engineering and Physical Sciences Research Council. A portion of this work was performed under the auspices of the US Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344. J.A.G and G.J.A were supported by LLNL Laboratory Directed Research and Development project 18-SI-002. The paper has LLNL tracking number LLNL-JRNL-811857. This document was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor Lawrence Livermore National Security, LLC, nor any of their employees makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or Lawrence Livermore National Security, LLC. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or Lawrence Livermore National Security, LLC, and shall not be used for advertising or product endorsement purposes.

Sandia National Laboratories is a multimission laboratory managed and operated by National Technology & Engineering Solutions of Sandia, LLC, a wholly owned subsidiary of Honeywell International Inc., for the U.S. Department of Energy's National Nuclear Security Administration under contract DE-NA0003525. This paper describes objective technical results and analysis. Any subjective views or opinions that might be expressed in the paper do not necessarily represent the views of the U.S. Department of Energy or the United States Government.

AUTHOR CONTRIBUTIONS

P.W.H., J.A.G and G.J.A conceived the work and led the writing of the manuscript. All authors contributed to the manuscript and the ideas discussed at the Lorentz Center Meeting.

¹Rose, S. Set the controls for the heart of the Sun. *Contemporary Physics* **45**, 109–121 (2004).

²Spears, B. K. *et al.* Deep learning: A guide for practitioners in the physical sciences. *Physics of Plasmas* **25**, 080901 (2018).

This tutorial paper gives an introduction to scientific machine learning, with examples taken from ICF research

³Wang, Z., Peterson, J. L., Rea, C. & Humphreys, D. Special Issue on Machine Learning, Data Science, and Artificial Intelligence in Plasma Research. *IEEE Transactions on Plasma Science* **48**, 1–2 (2020).

⁴Colvin, J. & Larsen, J. *Extreme Physics* (Cambridge University Press, Cambridge, 2013).

⁵Graziani, F., Desjarlais, M. P., Redmer, R. & Trickey, S. B. (eds.) *Frontiers and Challenges in Warm Dense Matter*, vol. 96 of *Lecture Notes in*

Computational Science and Engineering (Springer International Publishing, Cham, 2014).

⁶Gould, O., Mangles, S., Rajantie, A., Rose, S. & Xie, C. Observing thermal Schwinger pair production. *Physical Review A* **99**, 052120 (2019).

⁷Millot, M. *et al.* Experimental evidence for superionic water ice using shock compression. *Nature Physics* **14**, 297–302 (2018).

⁸Celliers, P. M. *et al.* Insulator-metal transition in dense fluid deuterium. *Science* **361**, 677–682 (2018).

⁹Joshi, C. & Malka, V. Focus on Laser- and Beam-Driven Plasma Accelerators. *New Journal of Physics* **12**, 045003 (2010).

¹⁰Hidding, B., Foster, B., Hogan, M. J., Muggli, P. & Rosenzweig, J. B. Directions in plasma wakefield acceleration. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* **377**, 20190215 (2019).

¹¹Wang, W.-M. *et al.* Collimated ultrabright gamma rays from electron wiggle along a petawatt laser-irradiated wire in the QED regime. *Proceedings of the National Academy of Sciences* **115**, 9911–9916 (2018).

¹²Badziak, J. Laser-driven ion acceleration: methods, challenges and prospects. *Journal of Physics: Conference Series* **959**, 012001 (2018).

¹³Feng, J. *et al.* High-efficiency neutron source generation from photonuclear reactions driven by laser plasma accelerator. *High Energy Density Physics* **36**, 100753 (2020).

¹⁴Hidding, B. *et al.* Plasma wakefield accelerator research 2019 - 2040: A community-driven uk roadmap compiled by the plasma wakefield accelerator steering committee (pwasc). WorkingPaper (2019).

¹⁵Nuckolls, J., Wood, L., Thiessen, A. & Zimmerman, G. Laser compression of matter to super-high densities: Thermonuclear applications. *Nature* **239**, 139–142 (1972).

¹⁶Hurricane, O. A. & Program, I. Overview of Progress and Future Prospects in Indirect Drive Implosions on the National Ignition Facility. *Journal of Physics: Conference Series* **717**, 012005 (2016).

¹⁷Fournier, K. & K. NIF Monthly Highlights for July 2019. Tech. Rep., Lawrence Livermore National Laboratory (LLNL), Livermore, CA (United States) (2019). URL <http://www.osti.gov/servlets/purl/1548376/>.

¹⁸Remington, B. A., Rudd, R. E. & Wark, J. S. From microjoules to megajoules and kilobars to gigabars: Probing matter at extreme states of deformation. *Physics of Plasmas* **22**, 090501 (2015).

¹⁹Trines, R. CLF Annual Report 2017-2018. Tech. Rep., Science and Technology Facilities Council (2019). URL <https://www.clf.stfc.ac.uk/Pages/Annual-Report-2017-18.aspx>.

²⁰Sturm, C. & Stöcker, H. The facility for antiproton and ion research FAIR. *Physics of Particles and Nuclei Letters* **8**, 865–868 (2011).

²¹Roso, L. High repetition rate Petawatt lasers. *EPJ Web of Conferences* **167**, 01001 (2018).

²²Zheng, W. *et al.* Laser performance of the SG-III laser facility. *High Power Laser Science and Engineering* **4**, e21 (2016).

²³Danson, C. N. *et al.* Petawatt and exawatt class lasers worldwide. *High Power Laser Science and Engineering* **7**, e54 (2019).

This paper reviews some of the many high-powered lasers in use around the world.

²⁴*Opportunities in Intense Ultrafast Lasers* (National Academies Press, Washington, D.C., 2018).

²⁵Lin, Z. *et al.* SG-II laser elementary research and precision SG-II program. *Fusion Engineering and Design* **44**, 61–66 (1999).

²⁶LULI. LULI2000 User Guide. Tech. Rep., Intenses, Laboratoire pour l'Utilisation des Lasers (2019).

²⁷Kirillov, G. A., Murugov, V. M., Punin, V. T. & Shemyakin, V. I. High power laser system ISKRA V. *Laser and Particle Beams* **8**, 827–831 (1990).

²⁸Zhao, Z. & Wang, D. XFEL Projects in China (2019). Presented at LINAC2018 in Beijing, China, unpublished.

²⁹Zhang, Z. *et al.* The laser beamline in SULF facility. *High Power Laser Science and Engineering* **8**, e4 (2020).

³⁰Schramm, U. *et al.* First results with the novel petawatt laser acceleration facility in Dresden. In *Journal of Physics: Conference Series*, vol. 874, 12028 (Institute of Physics Publishing, 2017).

- ³¹XFEL. European XFEL Facts and Figures. URL https://www.xfel.eu/facility/overview/facts_amp_figures/index_eng.html.
- ³²LCLS-II. LCLS-II Design & Performance. URL <https://lcls.slac.stanford.edu/lcls-ii/design-and-performance>.
- ³³Yabashi, M., Tanaka, H., Tono, K. & Ishikawa, T. Status of the SACLA Facility. *Applied Sciences* **7**, 604 (2017).
- ³⁴CERN FAQ – LHC the guide. Tech. Rep., CERN-Brochure-2017-002-Eng, Education, Communications and Outreach Group, CERN (2017).
- ³⁵MacDonald, M. J. *et al.* Measurement of high-dynamic range x-ray Thomson scattering spectra for the characterization of nano-plasmas at LCLS. *Review of Scientific Instruments* **87**, 11E709 (2016).
- ³⁶Mitchell, T. *Machine Learning* (McGraw-Hill, New York, 1997).
- ³⁷Sivia, D. & Skilling, J. *Data analysis: a Bayesian tutorial* (Oxford University Press).
- ³⁸Brunton, S. L. & Kutz, J. N. *Data-Driven Science and Engineering* (Cambridge University Press, 2019).
- ³⁹Reichstein, M. *et al.* Deep learning and process understanding for data-driven Earth system science. *Nature* **566**, 195–204 (2019).
- ⁴⁰Fleming, S. W. & Gupta, H. V. The physics of river prediction. *Physics Today* **73**, 46–52 (2020).
- ⁴¹Streeter, M. J. V. *et al.* Temporal feedback control of high-intensity laser pulses to optimize ultrafast heating of atomic clusters. *Applied Physics Letters* **112**, 244101 (2018).
- ⁴²Gopalaswamy, V. *et al.* Tripled yield in direct-drive laser fusion through statistical modelling. *Nature* **565**, 581–586 (2019).
- This paper is the first to use data-driven approaches to motivate and actually carry out new ICF experiments.**
- ⁴³Hatfield, P. W., Rose, S. J. & Scott, R. H. H. The blind implosion-maker: Automated inertial confinement fusion experiment design. *Physics of Plasmas* **26**, 062706 (2019).
- ⁴⁴Martin, M., London, R., Goluoglu, S. & Whitley, H. An automated design process for short pulse laser driven opacity experiments. *High Energy Density Physics* **26**, 26–37 (2018).
- ⁴⁵Raghu, M. & Schmidt, E. A Survey of Deep Learning for Scientific Discovery (2020). 2003.11755.
- ⁴⁶Thayer, J. *et al.* Data systems for the Linac coherent light source. *Advanced Structural and Chemical Imaging* **3**, 3 (2017).
- ⁴⁷Bernal, J. L., Peacock, J. A., Bernal, J. L. & Peacock, J. A. Conservative cosmology: combining data with allowance for unknown systematics. *JCAP* **2018**, 002 (2018).
- This paper considers how to get realistic uncertainty estimates in the presence of unknown systematics.**
- ⁴⁸Osthus, D., Vander Wiel, S. A., Hoffman, N. M. & Wysocki, F. J. Prediction Uncertainties beyond the Range of Experience: A Case Study in Inertial Confinement Fusion Implosion Experiments. *SIAM/ASA Journal on Uncertainty Quantification* **7**, 604–633 (2019).
- ⁴⁹Lang, M. & Owens, M. J. A Variational Approach to Data Assimilation in the Solar Wind. *Space Weather* **17**, 59–83 (2019).
- ⁵⁰Gaffney, J. A. *et al.* Making inertial confinement fusion models more predictive. *Physics of Plasmas* **26**, 082704 (2019).
- ⁵¹Kasim, M. F., Galligan, T. P., Topp-Muggleston, J., Gregori, G. & Vinko, S. M. Inverse problem instabilities in large-scale modeling of matter in extreme conditions. *Physics of Plasmas* **26**, 112706 (2019).
- ⁵²van de Plassche, K. L. *et al.* Fast modeling of turbulent transport in fusion plasmas using neural networks. *Physics of Plasmas* **27**, 022310 (2020).
- ⁵³Meneghini, O. *et al.* Self-consistent core-pedestal transport simulations with neural network accelerated models. *Nuclear Fusion* **57**, 086034 (2017).
- ⁵⁴Anirudh, R., Thiagarajan, J. J., Bremer, P.-T. & Spears, B. K. Improved surrogates in inertial confinement fusion with manifold and cycle consistencies. *Proceedings of the National Academy of Sciences* **117**, 9741–9746 (2020).
- ⁵⁵Kluth, G. *et al.* Deep learning for NLTE spectral opacities. *Physics of Plasmas* **27**, 052707 (2020).
- This paper uses machine learning to emulate opacity calculations, in order to speed up ICF simulations.**
- ⁵⁶Humbird, K. D., Peterson, J. L. & McClarren, R. G. Deep neural network initialization with decision trees. *IEEE transactions on neural networks and learning systems* **30**, 1286–1295 (2018).
- This paper presents a data-driven method for selecting deep neural network architecture and hyperparameters**
- ⁵⁷Kasim, M. F. *et al.* Building high accuracy emulators for scientific simulations with deep neural architecture search (2020). arXiv:2001.08055.
- ⁵⁸Kustowski, B. *et al.* Transfer Learning as a Tool for Reducing Simulation Bias: Application to Inertial Confinement Fusion. *IEEE Transactions on Plasma Science* **48**, 46–53 (2020).
- Transfer learning updates a subset of a pre-trained deep neural network using experimental data. This paper explores its application to sparse ICF datasets**
- ⁵⁹Bambi, C. Astrophysical Black Holes: A Compact Pedagogical Review (2018). 1711.10256.
- ⁶⁰The National Space Weather Program Strategic Plan. Tech. Rep., CM-P30-1995, Washington D.C. (1995).
- ⁶¹Eastwood, J. P. *et al.* The Economic Impact of Space Weather: Where Do We Stand? *Risk Analysis* **37**, 206–218 (2017).
- ⁶²Camporeale, E. The Challenge of Machine Learning in Space Weather: Nowcasting and Forecasting. *Space Weather* **17**, 1166–1207 (2019).
- ⁶³Camporeale, E., Carè, A. & Borovsky, J. E. Classification of Solar Wind With Machine Learning. *Journal of Geophysical Research: Space Physics* **122**, 10,910–10,920 (2017).
- ⁶⁴Chen, Y. *et al.* Identifying Solar Flare Precursors Using Time Series of SDO/HMI Images and SHARP Parameters. *Space Weather* **17**, 1404–1426 (2019).
- ⁶⁵Campi, C. *et al.* Feature Ranking of Active Region Source Properties in Solar Flare Forecasting and the Uncompromised Stochasticity of Flare Occurrence. *The Astrophysical Journal* **883**, 150 (2019).
- ⁶⁶Inceoglu, F. *et al.* Using Machine Learning Methods to Forecast if Solar Flares Will Be Associated with CMEs and SEPs. *The Astrophysical Journal* **861**, 128 (2018).
- ⁶⁷Bobra, M. G. & Ilonidis, S. Predicting coronal mass ejections using machine learning methods. *The Astrophysical Journal* **821**, 127 (2016).
- ⁶⁸Sarma, R. *et al.* Bayesian Inference of Quasi-Linear Radial Diffusion Parameters using Van Allen Probes. *Journal of Geophysical Research: Space Physics* **125** (2020).
- ⁶⁹Camporeale, E. *et al.* A Gray-Box Model for a Probabilistic Estimate of Regional Ground Magnetic Perturbations: Enhancing the NOAA Operational Geospace Model With Machine Learning. *Journal of Geophysical Research: Space Physics* **125**, e27684 (2020). 1912.01038.
- ⁷⁰Lamb, K. E. Correlation of auroral dynamics and gnss scintillation with an autoencoder. In *Second Workshop on Machine Learning and the Physical Sciences (NeurIPS 2019)*, Vancouver, Canada (2019).
- ⁷¹Rowlinson, A. *et al.* Identifying transient and variable sources in radio images. *Astronomy and Computing* **27**, 111–129 (2019).
- ⁷²Coughlin, M. W., Dietrich, T., Margalit, B. & Metzger, B. D. Multimessenger Bayesian parameter inference of a binary neutron-star merger. *Monthly Notices of the Royal Astronomical Society: Letters* **489**, L91–L96 (2019).
- This paper uses both gravitational wave, and multi-wavelength electromagnetic data to simultaneously constrain stellar masses, orbital parameters, and supra-nuclear density equations-of-state.**
- ⁷³Dorn, C. *et al.* A generalized Bayesian inference method for constraining the interiors of super Earths and sub-Neptunes. *Astronomy & Astrophysics* **597**, A37 (2017).
- ⁷⁴Bellinger, E. P. *et al.* Fundamental parameters of main-sequence stars in an instant with machine learning. *The Astrophysical Journal* **830**, 31 (2016).
- ⁷⁵Lochner, M., McEwen, J. D., Peiris, H. V., Lahav, O. & Winter, M. K. Photometric supernova classification with machine learning. *The Astrophysical Journal Supplement Series* **225**, 31 (2016).

This paper uses machine learning to emulate opacity calculations,

- ⁷⁶Kong, X. *et al.* Spectral feature extraction for DB white dwarfs through machine learning applied to new discoveries in the SDSS DR12 and DR14. *PASP* **130**, 084203 (2018).
- ⁷⁷Huppenkothen, D., Heil, L. M., Hogg, D. W. & Mueller, A. Using machine learning to explore the long-term evolution of GRS 1915+105. *Monthly Notices of the Royal Astronomical Society* **466**, 2364–2377 (2017).
- ⁷⁸Chardin, J. *et al.* A deep learning model to emulate simulations of cosmic reionization. *MNRAS* **490**, 1055–1065 (2019).
- ⁷⁹Saumon, D. & Guillot, T. Shock Compression of Deuterium and the Interiors of Jupiter and Saturn. *The Astrophysical Journal* **609**, 1170–1180 (2004).
- ⁸⁰Smith, R. F. *et al.* Ramp compression of diamond to five terapascals. *Nature* **511**, 330–333 (2014).
- ⁸¹Tzeferacos, P. *et al.* Laboratory evidence of dynamo amplification of magnetic fields in a turbulent plasma. *Nature Communications* **9**, 591 (2018).
- ⁸²Falcon, R. E. *et al.* Laboratory measurements of white dwarf photospheric spectral lines: H β . *The Astrophysical Journal* **806**, 214 (2015).
- ⁸³Bailey, J. E. *et al.* A higher-than-predicted measurement of iron opacity at solar interior temperatures. *Nature* **517**, 56–59 (2015).
- ⁸⁴Nagayama, T. *et al.* Systematic Study of L-Shell Opacity at Stellar Interior Temperatures. *Physical Review Letters* **122**, 235001 (2019).
- ⁸⁵de Souza, R. S., Boston, S. R., Coc, A. & Iliadis, C. Thermonuclear fusion rates for tritium + deuterium using Bayesian methods. *Physical Review C* **99**, 014619 (2019).
- ⁸⁶Zhelavskaya, I. S., Shprits, Y. Y. & Spasojević, M. Empirical Modeling of the Plasmasphere Dynamics Using Neural Networks. *Journal of Geophysical Research: Space Physics* **122**, 11,227–11,244 (2017).
- ⁸⁷Zhelavskaya, I. S., Shprits, Y. Y. & Spasojevic, M. Reconstruction of Plasma Electron Density From Satellite Measurements Via Artificial Neural Networks. *Machine Learning Techniques for Space Weather* 301–327 (2018).
- ⁸⁸Freidberg, J. *Plasma Physics and Fusion Energy* (Cambridge University Press, 2008).
- ⁸⁹Atzeni, S., Meyer-ter Vehn, J. & Meyer-ter Vehn, J. *The Physics of Inertial Fusion: BeamPlasma Interaction, Hydrodynamics, Hot Dense Matter*. International Series of Monographs on Physics (Clarendon Press, 2004).
- ⁹⁰Lindl, J. *Inertial Confinement Fusion: The Quest for Ignition and Energy Gain Using Indirect Drive*. AIP-Press Series (AIP Press, 1998).
- ⁹¹Campbell, E. M. & Hogan, W. J. The National Ignition Facility - applications for inertial fusion energy and high-energy-density science. *Plasma Physics and Controlled Fusion* **41**, B39–B56 (1999).
- ⁹²Moses, E. I. Ignition on the national ignition facility. *Journal of Physics: Conference Series* **112**, 012003 (2008).
- ⁹³Boehly, T. *et al.* Initial performance results of the omega laser system. *Optics Communications* **133**, 495 – 506 (1997).
- ⁹⁴Slutz, S. A. High-gain magnetized inertial fusion. *Physical Review Letters* **108** (2012).
- ⁹⁵Gomez, M. R. *et al.* Experimental demonstration of fusion-relevant conditions in magnetized liner inertial fusion. *Phys. Rev. Lett.* **113**, 155003 (2014).
- ⁹⁶Dimonte, G. Quantitative metrics for evaluating thermonuclear design codes and physics models applied to the National Ignition Campaign. *Physics of Plasmas* **27**, 052709 (2020).
- ⁹⁷Yang, C. *et al.* Preparing dense net for automated hydra mesh management via reinforcement learning. Tech. Rep. LLNL-TR-799958, Lawrence Livermore National Laboratory (2019). URL <https://www.osti.gov/servlets/purl/1580017>.
- This report details the use of deep neural networks to automatically perform adaptive mesh refinement in radiation-hydrodynamics simulations**
- ⁹⁸Peterson, J. L. *et al.* Merlin: Enabling Machine Learning-Ready HPC Ensembles (2019). arXiv:1912.02892.
- ⁹⁹Peterson, J. L. *et al.* Zonal flow generation in inertial confinement fusion implosions. *Physics of Plasmas* **24**, 032702 (2017).
- This paper presents the first novel ICF design reached using machine learning.**
- ¹⁰⁰Amorin, C., Kegelmeyer, L. M. & Kegelmeyer, W. P. A hybrid deep learning architecture for classification of microscopic damage on National Ignition Facility laser optics. *Statistical Analysis and Data Mining: The ASA Data Science Journal* **12**, 505–513 (2019).
- Here, deep neural network-based image classification is used to detect and classify damage in the NIF laser system, allowing more reliable operation at high energy**
- ¹⁰¹Nora, R., Peterson, J. L., Spears, B. K., Field, J. E. & Brandon, S. Ensemble simulations of inertial confinement fusion implosions. *Statistical Analysis and Data Mining: The ASA Data Science Journal* **10**, 230–237 (2017).
- ¹⁰²Gaffney, J. A. *et al.* The JAG inertial confinement fusion simulation dataset for multi-modal scientific deep learning (2020). URL <https://library.ucsd.edu/dc/object/bb5534097t>.
- This dataset is an example of open data practices in ICF, and is one of the first multi-modal scientific datasets to be released by the HEDP community**
- ¹⁰³Thiagarajan, J. J. *et al.* Designing accurate emulators for scientific processes using calibration-driven deep models. *Nature Communications* **11**, 1–10 (2020). 2005.02328.
- ¹⁰⁴Hatfield, P. *et al.* Using sparse gaussian processes for predicting robust inertial confinement fusion implosion yields. *IEEE Transactions on Plasma Science* **48**, 14–21 (2020).
- ¹⁰⁵Glinisky, M. E. *et al.* Quantification of MagLIF Morphology using the Mallat Scattering Transformation. *Office of Scientific and Technical Information (OSTI)* (2019). URL <http://dx.doi.org/10.2172/1569785>.
- This paper uses a novel, physics-motivated deep neural network architecture to featurize images of ICF implosions driven on the Z pulsed-power machine**
- ¹⁰⁶Palaniyappan, S. *et al.* Hydro-scaling of direct-drive cylindrical implosions at the omega and the national ignition facility. *Physics of Plasmas* **27**, 042708 (2020).
- ¹⁰⁷Gaffney, J., Clark, D., Sonnad, V. & Libby, S. Development of a Bayesian method for the analysis of inertial confinement fusion experiments on the NIF. *Nuclear Fusion* **53**, 073032 (2013).
- ¹⁰⁸Gaffney, J., Clark, D., Sonnad, V. & Libby, S. Bayesian inference of inaccuracies in radiation transport physics from inertial confinement fusion experiments. *High Energy Density Physics* **9**, 457 – 461 (2013).
- ¹⁰⁹Knapp, P. A Bayesian Parameter Estimation Framework for Understanding Fusion Experiments on Z. Tech. Rep. (2018). URL <https://www.osti.gov/biblio/1525597>.
- ¹¹⁰Hsu, A., Cheng, B. & Bradley, P. A. Analysis of NIF scaling using physics informed machine learning. *Physics of Plasmas* **27**, 012703 (2020).
- ¹¹¹Humbird, K. D., Peterson, J. L., Spears, B. K. & McClarren, R. G. Transfer Learning to Model Inertial Confinement Fusion Experiments. *IEEE Transactions on Plasma Science* **48**, 61–70 (2020).
- ¹¹²Götzfried, J. *et al.* Research towards high-repetition rate laser-driven X-ray sources for imaging applications. *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment* **909**, 286–289 (2018). 1803.05415.
- ¹¹³He, Z.-H. *et al.* Coherent control of plasma dynamics. *Nature communications* **6**, 7156 (2015).
- ¹¹⁴Dann, S. J. *et al.* Laser wakefield acceleration with active feedback at 5 Hz. *Physical Review Accelerators and Beams* **22**, 041303 (2019).
- ¹¹⁵Kirschner, J., Mutny, M. M., Hiller, N., Ischebeck, R. & Krause, A. Adaptive and Safe Bayesian Optimization in High Dimensions via One-Dimensional Subspaces. *Proceedings of the 36th International Conference on Machine Learning* 3429–3438 (2019).
- This paper describes a Bayesian Optimisation method for a Free Electron Laser that prevents the laser from violating any safety constraints.**

- ¹¹⁶Shaloo, R. J. *et al.* Automation and control of laser wakefield accelerators using Bayesian optimization. *Nature Communications* **11**, 1–8 (2020). 2007.14340.
- ¹¹⁷Maier, A. R. *et al.* Decoding Sources of Energy Variability in a Laser-Plasma Accelerator. *Physical Review X* **10**, 031039 (2020).
- ¹¹⁸Team, J. T. *et al.* Hybrid neural network for density limit disruption prediction and avoidance on J-TEXT tokamak. *Nuclear Fusion* (2018).
- ¹¹⁹Fu, Y. *et al.* Machine learning control for disruption and tearing mode avoidance. *Physics of Plasmas* **27**, 022501 (2020).
- ¹²⁰Kates-Harbeck, J., Svyatkovskiy, A. & Tang, W. Predicting disruptive instabilities in controlled fusion plasmas through deep learning. *Nature* **568**, 526–531 (2019).
- ¹²¹Blanken, T., Felici, F., Rapson, C., de Baar, M. & Heemels, W. Control-oriented modeling of the plasma particle density in tokamaks and application to real-time density profile reconstruction. *Fusion Engineering and Design* **126**, 87–103 (2018).
- ¹²²Wilkinson, M. D. *et al.* Comment: The FAIR Guiding Principles for scientific data management and stewardship. *Scientific Data* **3**, 1–9 (2016).
- ¹²³Albertsson, K. *et al.* Machine Learning in High Energy Physics Community White Paper. *Journal of Physics: Conference Series* **1085**, 022008 (2018).
- ¹²⁴Borne, K. D. *et al.* The Revolution in Astronomy Education: Data Science for the Masses (2009). arXiv:0909.3895.
- ¹²⁵Pasian, F. *et al.* Science ground segment for the ESA Euclid Mission. In Radziwill, N. M. & Chiozzi, G. (eds.) *Software and Cyberinfrastructure for Astronomy II*, vol. 8451, 845104 (SPIE, 2012).
- ¹²⁶Lyons, L. Discovering the Significance of 5 sigma (2013). arXiv:1310.1284.
- This paper discusses the background, rationale, and advantages and disadvantages of the 5- σ criterion commonly used in particle physics.**
- ¹²⁷Roodman, A. Blind Analysis in Particle Physics. In *Proceedings of the Conference on Statistical Problems in Particle Physics, Astrophysics and Cosmology, Stanford Linear Accelerator Center, Stanford, California, September 8th – 11th, 2003* (2003).