Directive words of episturmian words: equivalences and normalization

Amy Glen[†]

Florence Levé[‡]

Gwénaël Richomme[‡]

February 26, 2008

Abstract

Episturmian morphisms constitute a powerful tool to study episturmian words. Indeed, any episturmian word can be infinitely decomposed over the set of pure episturmian morphisms. Thus, an episturmian word can be defined by one of its morphic decompositions or, equivalently, by a certain directive word. Here we characterize pairs of words directing a common episturmian word. We also propose a way to uniquely define any episturmian word through a normalization of its directive words. As a consequence of these results, we characterize episturmian words having a unique directive word.

Keywords: episturmian word; Sturmian word; Arnoux-Rauzy sequence; episturmian morphism; directive word.

MSC (2000): 68R15.

1 Introduction

Since the seminal works of Morse and Hedlund [21], Sturmian words have been widely studied and their beautiful properties are related to many fields like Number Theory, Geometry, Dynamical Systems, and Combinatorics on Words (see [1, 20, 23, 3] for recent surveys). These infinite words, which are defined on a binary alphabet, have numerous equivalent definitions and characterizations. Nowadays most works deal with generalizations of Sturmian words to arbitrary finite alphabets. Two very interesting generalizations are very close: the Arnoux-Rauzy sequences (e.g., see [2, 14, 23, 30]) and episturmian words (e.g., see [5, 13, 15]). The first of these two families is a particular subclass of the second one. More precisely, the family of episturmian words is composed of the Arnoux-Rauzy sequences, images of the Arnoux-Rauzy sequences by episturmian morphisms, and certain periodic infinite words. In the binary case, Arnoux-Rauzy sequences are exactly the Sturmian words whereas episturmian words include all recurrent balanced words, that is, periodic balanced words and Sturmian words (see [10, 22, 29] for recent results relating episturmian words to the balanced property). See also [9] for a recent survey on episturmian theory.

Epistumian morphisms play a central role in the study of these words. Introduced first as a generalization of Sturmian morphisms, Justin and Pirillo [13] showed that they are exactly the morphisms that preserve the aperiodic epistumian words. They also proved that any epistumian

[†]Amy Glen

LaCIM, Université du Québec à Montréal,

C.P. 8888, succursale Centre-ville, Montréal, Québec, CANADA, H3C 3P8

E-mail: amy.glen@gmail.com (with the support of CRM-ISM-LaCIM)

[‡]F. Levé, G. Richomme

Université de Picardie Jules Verne,

Laboratoire MIS (Modélisation, Information, Systèmes)

^{33,} Rue Saint Leu, F-80039 Amiens cedex 1, FRANCE

E-mail: {florence.leve, gwenael.richomme}@u-picardie.fr

word is the image of another episturmian word by some so-called *pure episturmian morphism*. Even more, any episturmian word can be infinitely decomposed over the set of pure episturmian morphisms. This last property allows an episturmian word to be defined by one of its morphic decompositions or, equivalently, by a certain *directive word*, which is an infinite sequence of rules for decomposing the given episturmian word by morphisms. In consequence, many properties of episturmian words can be deduced from properties of episturmian morphisms. This approach is used for instance in [4, 8, 16, 28, 29, 30] and of course in the papers of Justin *et al.* In Section 2, we recall useful results on episturmian words and explain the vision of morphic decompositions and directive words introduced by Justin and Pirillo in [13].

An episturmian word can have several directive words. The question: "When do two words direct a common episturmian word?" was considered in [15]. Using a block-equivalence notion for directive words, Justin and Pirillo provided several results to answer this question in most cases (see Section 3). In Section 4, we state a complete result characterizing the form of words directing a common episturmian word, without using block-equivalence.

In [4], Berthé, Holton, and Zamboni show that any Sturmian word has a unique directive word with some particular properties. In [18], the second and third authors rephrased this result and used it to characterize all quasiperiodic Sturmian words. In Section 5, we extend this result to all episturmian words by introducing a way to normalize the directive words of an episturmian word so that any episturmian word can be defined uniquely by its *normalized directive word*, defined by some factor avoidance (Theorem 5.2). This result was previously presented at the Sixth International Conference on Words [17] to characterize all quasiperiodic episturmian words (see also [11]).

As an application of the previous results, we end this paper with a characterization of episturmian words having a unique directive word.

2 Episturmian words and morphisms

We assume the reader is familiar with combinatorics on words and morphisms (e.g., see [19, 20]). In this section, we recall some basic definitions and properties relating to episturmian words which are needed later in the paper. For the most part, we follow the notation and terminology of [5, 13, 15, 10].

2.1 Notation and terminology

Let \mathcal{A} denote a finite *alphabet*. A finite *word* over \mathcal{A} is a finite sequence of letters from \mathcal{A} . The *empty word* ε is the empty sequence. Under the operation of concatenation, the set \mathcal{A}^* of all finite words over \mathcal{A} is a *free monoid* with identity element ε and set of generators \mathcal{A} . The set of *non-empty* words over \mathcal{A} is the *free semigroup* $\mathcal{A}^+ = \mathcal{A}^* \setminus \{\varepsilon\}$.

Given a finite word $w = x_1 x_2 \cdots x_m \in \mathcal{A}^+$ with each $x_i \in \mathcal{A}$, the *length* of w is |w| = m. The length of the empty word is 0. By $|w|_a$ we denote the number of occurrences of the letter a in the word w. If $|w|_a = 0$, then w is said to be *a-free*. For any integer $p \ge 1$, the *p*-th power of w is the word w^p obtained by concatenating p occurrences of w.

A (right) infinite word \mathbf{x} is a sequence indexed by \mathbb{N}^+ with values in \mathcal{A} , i.e., $\mathbf{x} = x_1 x_2 x_3 \cdots$ with each $x_i \in \mathcal{A}$. The set of all infinite words over \mathcal{A} is denoted by \mathcal{A}^{ω} . Given a non-empty finite word v, we denote by v^{ω} the infinite word obtained by concatenating v with itself infinitely many times. For easier reading, infinite words are hereafter typed in boldface to distinguish them from finite words.

Given a set X of words, X^* (resp. X^{ω}) is the set of all finite (resp. infinite) words that can be obtained by concatenating words of X. The empty word ε belongs to X^* .

A finite word w is a *factor* of a finite or infinite word z if z = uwv for some words u, v (where v is infinite iff z is infinite). Further, w is called a *prefix* (resp. *suffix*) of z if $u = \varepsilon$ (resp. $v = \varepsilon$). We use the notation $p^{-1}w$ (resp. ws^{-1}) to indicate the removal of a prefix p (resp. suffix s) of the

word w.

The *alphabet* of a word w, denoted by Alph(w) is the set of letters occurring in w, and if w is infinite, we denote by Ult(w) the set of all letters occurring infinitely often in w.

2.2 Episturmian words

In this paper, our vision of episturmian words will be the characteristic property stated in Theorem 2.1 below. Nevertheless, to give an idea of what an episturmian word is, let us give one of the equivalent definitions of an episturmian word provided in [5]. Before doing so, we recall that a factor u of an infinite word $\mathbf{w} \in \mathcal{A}^{\omega}$ is right (resp. left) special if ua, ub (resp. au, bu) are factors of \mathbf{w} for some letters $a, b \in \mathcal{A}, a \neq b$. We recall also that the reversal \tilde{w} of a finite word w is its mirror image: if $w = x_1 \dots x_{m-1} x_m$, then $\tilde{w} = x_m x_{m-1} \dots x_1$.

An infinite word $\mathbf{t} \in \mathcal{A}^{\omega}$ is *episturmian* if its set of factors is closed under reversal and \mathbf{t} has at most one right (or equivalently left) special factor of each length. Moreover, an episturmian word is *standard* if all of its left special factors are prefixes of it.

In the initiating paper [5], episturmian words were defined in two steps. Standard episturmian words were first introduced and studied as a generalization of standard Sturmian words. (Note that in the rest of this paper, we refer to a standard episturmian word as an *epistandard word*, for simplicity). Then an episturmian word was defined as an infinite word having exactly the same set of factors as some epistandard word.

Moreover, it was proved in [5] that any episturmian word is *recurrent*, that is, all of its factors occur infinitely often (actually episturmian words are uniformly recurrent but this will not be needed here). An *ultimately periodic* infinite word is a word that can be written as $uv^{\omega} = uvvv \cdots$, for some $u, v \in \mathcal{A}^*, v \neq \varepsilon$. If $u = \varepsilon$, then such a word is *periodic*. Since they are recurrent, all ultimately periodic episturmian words are periodic. Let us recall that an infinite word that is not ultimately periodic is said to be *aperiodic*.

2.3 Episturmian morphisms

To study episturmian words, Justin and Pirillo [13] introduced *episturmian morphisms*. In particular they proved that these morphisms (defined below) are precisely the morphisms that preserve the set of aperiodic episturmian words.

Let us recall that given an alphabet \mathcal{A} , a morphism f on \mathcal{A} is a map from \mathcal{A}^* to \mathcal{A}^* such that f(uv) = f(u)f(v) for any words u, v over \mathcal{A} . A morphism on \mathcal{A} is entirely defined by the images of letters in \mathcal{A} . All morphisms considered in this paper will be non-erasing: the image of any non-empty word is never empty. Hence the action of a morphism f on \mathcal{A}^* can be naturally extended to infinite words; that is, if $\mathbf{x} = x_1 x_2 x_3 \cdots \in \mathcal{A}^{\omega}$, then $f(\mathbf{x}) = f(x_1) f(x_2) f(x_3) \cdots$.

In what follows, we will denote the composition of morphisms by juxtaposition as for concatenation of words.

Epistumian morphisms are the compositions of the permutation morphisms (the morphisms f such that $f(\mathcal{A}) = \mathcal{A}$) and the morphisms L_a and R_a where, for all $a \in \mathcal{A}$:

$$L_a: \left\{ \begin{array}{ccc} a & \mapsto & a \\ b & \mapsto & ab \end{array} \right., \quad R_a: \left\{ \begin{array}{ccc} a & \mapsto & a \\ b & \mapsto & ba \end{array} \right. \quad \text{for all } b \neq a \text{ in } \mathcal{A}.$$

Here we will work only on *pure* episturmian morphisms, i.e., morphisms obtained by composition of elements of the sets:

$$\mathcal{L}_{\mathcal{A}} = \{ L_a \mid a \in \mathcal{A} \} \text{ and } \mathcal{R}_{\mathcal{A}} = \{ R_a \mid a \in \mathcal{A} \}.$$

Note. In [13], the morphism L_a (resp. R_a) is denoted by ψ_a (resp. $\overline{\psi}_a$). We adopt the current notation to emphasize the action of L_a (resp. R_a) when applied to a word, which consists in placing an occurrence of the letter a on the left (resp. right) of each occurrence of any letter different from a.

Epistandard morphisms are the morphisms obtained by concatenation of morphisms in $\mathcal{L}_{\mathcal{A}}$ and permutations on \mathcal{A} . Likewise, the *pure episturmian morphisms* (resp. *pure epistandard morphisms*) are the morphisms obtained by concatenation of morphisms in $\mathcal{L}_{\mathcal{A}} \cup \mathcal{R}_{\mathcal{A}}$ (resp. in $\mathcal{L}_{\mathcal{A}}$). Note that the episturmian morphisms are exactly the *Sturmian morphisms* when \mathcal{A} is a 2-letter alphabet.

All episturmian morphisms are injective on both finite and infinite words. The monoid of episturmian morphisms is *left cancellative* (see [26, Lem. 7.2]) which means that for any episturmian morphisms f, g, h, if fg = fh then g = h. Note that this fact, which is a by-product of the injectivity, can also be seen as a consequence of the invertibility of these morphisms (see [7, 12, 26, 32]).

2.4 Morphic decomposition of episturmian words

Justin and Pirillo [13] proved the following insightful characterizations of epistandard and episturmian words (see Theorem 2.1 below), which show that any episturmian word can be *infinitely decomposed* over the set of pure episturmian morphisms.

The statement of Theorem 2.1 needs some extra definitions and notation.

First we define the following new alphabet, $\overline{\mathcal{A}} = \{\overline{x} \mid x \in \mathcal{A}\}$. A letter \overline{x} is considered to be x with spin R, whilst x itself has spin L. A finite or infinite word over $\mathcal{A} \cup \overline{\mathcal{A}}$ is called a spinned word. To ease the reading, we sometimes call a letter with spin L (resp. spin R) an L-spinned (resp. R-spinned) letter. By extension, an L-spinned (resp. R-spinned) word is a word having only letters with spin L (resp. spin R).

The opposite \bar{w} of a finite or infinite spinned word w is obtained from w by exchanging all spins in w. For instance, if $v = ab\bar{a}$, then $\bar{v} = \bar{a}\bar{b}a$. When $v \in \mathcal{A}^+$, then its opposite $\bar{v} \in \bar{\mathcal{A}}^+$ is an R-spinned word and we set $\bar{\varepsilon} = \varepsilon$. Note that, given a finite or infinite word $w = w_1w_2...$ over \mathcal{A} , we sometimes denote $\bar{w} = \bar{w}_1\bar{w}_2\cdots$ any spinned word such that $\bar{w}_i = w_i$ if \bar{w}_i has spin L and $\bar{w}_i = \bar{w}_i$ if \bar{w}_i has spin R. Such a word \bar{w} is called a *spinned version* of w.

Note. In Justin and Pirillo's original papers, spins are 0 and 1 instead of L and R. It is convenient here to change this vision of the spins because of the relationship with episturmian morphisms, which we now recall.

For $a \in \mathcal{A}$, let $\mu_a = L_a$ and $\mu_{\bar{a}} = R_a$. This operator μ can be naturally extended (as done in [13]) to a morphism from the free monoid $(\mathcal{A} \cup \bar{\mathcal{A}})^*$ to a pure episturmian morphism: for a spinned finite word $\check{w} = \check{w}_1 \dots \check{w}_n$ over $\mathcal{A} \cup \bar{\mathcal{A}}, \mu_{\check{w}} = \mu_{\check{w}_1} \dots \mu_{\check{w}_n}$ (μ_{ε} is the identity morphism). We say that the word w directs or is a directive word of the morphism μ_w . The following result extends the notion of directive words to infinite episturmian words.

Theorem 2.1. [13]

- i) An infinite word $\mathbf{s} \in \mathcal{A}^{\omega}$ is epistandard if and only if there exist an infinite word $\Delta = x_1 x_2 x_3 \cdots$ over \mathcal{A} and an infinite sequence $(\mathbf{s}^{(n)})_{n\geq 0}$ of infinite words such that $\mathbf{s}^{(0)} = \mathbf{s}$ and for all $n \geq 1$, $\mathbf{s}^{(n-1)} = L_{x_n}(\mathbf{s}^{(n)})$.
- ii) An infinite word $\mathbf{t} \in \mathcal{A}^{\omega}$ is episturmian if and only if there exist a spinned infinite word $\check{\Delta} = \check{x}_1\check{x}_2\check{x}_3\cdots$ over $\mathcal{A}\cup\bar{\mathcal{A}}$ and an infinite sequence $(\mathbf{t}^{(n)})_{n\geq 0}$ of recurrent infinite words such that $\mathbf{t}^{(0)} = \mathbf{t}$ and for all $n \geq 1$, $\mathbf{t}^{(n-1)} = \mu_{\check{x}_n}(\mathbf{t}^{(n)})$.

For any epistandard word (resp. episturmian word) \mathbf{t} and *L*-spinned infinite word Δ (resp. spinned infinite word $\check{\Delta}$) satisfying the conditions of the above theorem, we say that Δ (resp. $\check{\Delta}$) is a *(spinned) directive word* for \mathbf{t} or that \mathbf{t} is *directed by* Δ (resp. $\check{\Delta}$). Notice that this directive word is exactly the one that arises from the equivalent definition of epistandard words that uses *palindromic closure* [5, 9, 13] and, in the binary case, it is related to the continued fraction of the slope of the straight line represented by a standard word (see [20]). It follows immediately from Theorem 2.1 that, with the notation of case *ii*), each $\mathbf{t}^{(n)}$ is an episturmian word directed by $\check{x}_{n+1}\check{x}_{n+2}\cdots$

The natural question: "Does any spinned infinite word direct a unique episturmian word?" is answered in [13]:

Proposition 2.2. [13, Prop. 3.11]

- 1. Any spinned infinite word Δ having infinitely many L-spinned letters directs a unique episturmian word beginning with the left-most letter having spin L in $\check{\Delta}$.
- 2. Any R-spinned infinite word $\check{\Delta}$ directs exactly $|\text{Ult}(\Delta)|$ episturmian words.
- 3. Let $\check{\Delta}$ be an *R*-spinned infinite word, and let *a* be a letter such that $\bar{a} \in \text{Ult}(\check{\Delta})$. Then $\check{\Delta}$ directs exactly one episturmian word starting with *a*.

Note. In [13], item 3 was stated in the more general case where $\check{\Delta}$ is ultimately *R*-spinned. In this case, $\check{\Delta}$ still directs exactly one episturmian word for each letter \bar{a} in Ult($\check{\Delta}$), but contrary to what is written in [13], nothing can be said on its first letter.

As a consequence of the previous proposition and part i) of Theorem 2.1, any L-spinned infinite word directs a unique epistandard word. The following important remark links the two parts of Theorem 2.1.

Remark 2.3. [13] If $\check{\Delta}$ is a spinned version of an *L*-spinned word Δ and if **t** is an epistumian word directed by $\check{\Delta}$, then the set of factors of **t** is exactly the set of factors of the epistandard word **s** directed by Δ .

Moreover (with the same notation as in the previous remark):

Remark 2.4. The episturmian word **t** is periodic if and only if the epistandard word **s** is periodic, and this holds if and only if there is only one letter occurring infinitely often in Δ , that is, $|\text{Ult}(\Delta)| =$ 1 (see [13, Prop. 2.9]). More precisely, a periodic episturmian word takes the form $(\mu_{\tilde{w}}(x))^{\omega}$ for some finite spinned word \tilde{w} and letter x.

Note. Sturmian words are precisely the aperiodic episturmian words on a 2-letter alphabet.

When an episturmian word is aperiodic, we have the following fundamental link between the words $(\mathbf{t}^{(n)})_{n\geq 0}$ and the spinned infinite word $\check{\Delta}$ occurring in Theorem 2.1: if a_n is the first letter of $\mathbf{t}^{(n)}$, then $\mu_{\check{x}_1...\check{x}_n}(a_n)$ is a prefix of \mathbf{t} and the sequence $(\mu_{\check{x}_1...\check{x}_n}(a_n))_{n\geq 1}$ is not ultimately constant (since $\check{\Delta}$ is not ultimately constant), then $\mathbf{t} = \lim_{n\to\infty} \mu_{\check{x}_1...\check{x}_n}(a_n)$. This fact is a slight generalization of a result of Risley and Zamboni [30, Prop. III.7] on *S-adic representations* for characteristic Arnoux-Rauzy sequences. See also the recent paper [4] for S-adic representations of Sturmian words. Note that *S-adic dynamical systems* were introduced by Ferenczi [6] as minimal dynamical systems (e.g., see [23]) generated by a finite number of substitutions. In the case of episturmian words, the notion itself is actually a reformulation of the well-known Rauzy rules, as studied in [25].

To anticipate next sections, let us also observe:

Remark 2.5. [13] If an aperiodic episturmian word is directed by two spinned words Δ_1 and Δ_2 , then Δ_1 and Δ_2 are spinned versions of a common *L*-spinned word.

This is no longer true for periodic episturmian words; for instance ab^{ω} and $\bar{b}a^{\omega}$ direct the same episturmian word $(ab)^{\omega} = ababab \cdots$.

3 Known results on directive-equivalent words

We have just seen an example of a periodic episturmian word that is directed by two different spinned infinite words. This situation holds also in the aperiodic case (see [13, 15]). For example,

the Tribonacci word (or Rauzy word [24]) is directed by $(abc)^{\omega}$ and also by $(abc)^n \bar{a} \bar{b} \bar{c} (a \bar{b} \bar{c})^{\omega}$ for each $n \geq 0$, as well as infinitely many other spinned words. More generally, by [13], any epistandard word has a unique L-spinned directive word but also has other directive words (see also [15] and Theorem 4.1).

We now consider in detail the following two questions: When do two finite spinned words direct a common episturmian morphism? When do two spinned infinite words direct a common unique episturmian word? We say that that two finite (resp. infinite) spinned words are *directive-equivalent* words if they direct a common episturmian morphism (resp. a common episturmian word).

In Section 3.1, we recall the characterizations of directive-equivalent finite spinned words. In Section 3.2, we recall known results about directive-equivalent infinite words. Section 4 will present a new characterization of these words.

3.1 Finite directive-equivalent words: presentation versus block-equivalence

Generalizing a study of the monoid of Sturmian morphisms by Séébold [31], the third author [26] answered the question: "When do two spinned finite words direct a common episturmian morphism?" by giving a presentation of the monoid of episturmian morphisms. This result was reformulated in [27] using another set of generators and it was independently and differently treated in [15]. As a direct consequence, one can see that the monoid of pure epistandard morphisms is a free monoid and one can obtain the following presentation of the monoid of pure episturmian morphisms:

Theorem 3.1. (direct consequence of [27, Prop. 6.5]; reformulation of [15, Th. 2.2])

The monoid of pure episturmian morphisms with $\{L_{\alpha}, R_{\alpha} \mid \alpha \in \mathcal{A}\}$ as a set of generators has the following presentation:

$$R_{a_1}R_{a_2}\ldots R_{a_k}L_{a_1}=L_{a_1}L_{a_2}\ldots L_{a_k}R_{a_1}$$

where $k \geq 1$ is an integer and $a_1, \ldots, a_k \in \mathcal{A}$ with $a_1 \neq a_i$ for all $i, 2 \leq i \leq k$.

This result means that two different compositions of morphisms in $\mathcal{L}_{\mathcal{A}} \cup \mathcal{R}_{\mathcal{A}}$ yield a common pure episturmian morphism if and only if one composition can be deduced from the other one in a rewriting system, called the *block-equivalence* in [15]. Although Theorem 3.1 allows us to show that many properties of episturmian words are linked to properties of episturmian morphisms, it will be convenient for us to have in mind the block-equivalence that we now recall.

A word of the form xvx, where $x \in \mathcal{A}$ and $v \in (\mathcal{A} \setminus \{x\})^*$, is called a (x-based) block. A (xbased) block-transformation is the replacement in a spinned word of an occurrence of $xv\bar{x}$ (where xvx is a block) by $\bar{x}\bar{v}x$ or vice-versa. Two finite spinned words w, w' are said to be block-equivalent if we can pass from one to the other by a (possibly empty) chain of block-transformations, in which case we write $w \equiv w'$. For example, $\bar{b}\bar{a}b\bar{c}b\bar{a}\bar{c}$ and $babc\bar{b}\bar{a}\bar{c}$ are block-equivalent because $\bar{b}\bar{a}b\bar{c}b\bar{a}\bar{c} \rightarrow ba\bar{b}c\bar{b}\bar{a}\bar{c}$ and vice-versa. The block-equivalence is an equivalence relation over spinned words, and moreover one can observe that if $w \equiv w'$ then w and w' are spinned versions of a common word over \mathcal{A} .

Theorem 3.1 can be reformulated in terms of block-equivalence:

Theorem 3.1. Let w, w' be two spinned words over $\mathcal{A} \cup \overline{\mathcal{A}}$. Then $\mu_w = \mu_{w'}$ if and only if $w \equiv w'$.

3.2 Infinite directive-equivalent words: previous results

The question: "When do two spinned infinite words direct a common unique episturmian word?" was tackled by Justin and Pirillo in [15] for *bi-infinite episturmian words*, that is, episturmian words with letters indexed by \mathbb{Z} (and not by \mathbb{N} as considered until now). Let us recall relations between right-infinite episturmian words and bi-infinite episturmian words (see [15, p. 332] and [9] for more details).

First we observe that a right-infinite episturmian word \mathbf{t} can be prolonged infinitely to the left with the same set of factors. Note also that the definition of episturmian words considered in Section 2.2 (using reversal and special factors) can be extended to bi-infinite words (see [15]). Furthermore, the characterization (Theorem 2.1) of right-infinite episturmian words by a sequence $(\mathbf{t}^{(i)})_{i\geq 0}$ extends to bi-infinite episturmian words, with all the $\mathbf{t}^{(i)}$ now bi-infinite episturmian words. That is, as for right-infinite episturmian words, we have bi-infinite words of the form $\mathbf{l}^{(i)}.\mathbf{r}^{(i)}$ where $\mathbf{l}^{(i)}$ is a left-infinite episturmian word and $\mathbf{r}^{(i)}$ is a right-infinite episturmian word. Moreover, if the bi-infinite episturmian word $\mathbf{b} = \mathbf{l}.\mathbf{r}$ is directed by $\check{\Delta}$ with associated bi-infinite episturmian words $\mathbf{r}^{(i)}$.

As a consequence of what precedes, Justin and Pirillo's results about spinned words directing a common bi-infinite episturmian word are still valid for words directing a common (right-infinite) episturmian word. We summarize now these results, which will be helpful for the proof of our main theorem (Theorem 4.1, to follow).

First of all, Justin and Pirillo characterized pairs of words directing a common episturmian word in the case of *wavy* directive words, that is, spinned infinite words containing infinitely many L-spinned letters and infinitely many R-spinned letters. This characterization uses the following extension of the block-equivalence \equiv for infinite words.

Let Δ_1, Δ_2 be spinned versions of Δ . We write $\Delta_1 \rightsquigarrow \Delta_2$ if there exist infinitely many prefixes f_i of Δ_1 and g_i of Δ_2 with the g_i of strictly increasing lengths, and such that, for all $i, |g_i| \leq |f_i|$ and $f_i \equiv g_i c_i$ for a suitable spinned word c_i . Infinite words Δ_1 and Δ_2 are said to be *block-equivalent* (denoted by $\Delta_1 \equiv \Delta_2$) if $\Delta_1 \rightsquigarrow \Delta_2$ and $\Delta_2 \rightsquigarrow \Delta_1$.

Theorem 3.2. [15, Th. 3.4, Cor. 3.5] Let Δ_1 and Δ_2 be wavy spinned versions of $\Delta \in A^{\omega}$ with $|\text{Ult}(\Delta)| > 1$. Then Δ_1 and Δ_2 direct a common (unique) episturmian word if and only if $\Delta_1 \equiv \Delta_2$.

Moreover when Δ_1 and Δ_2 do not have any common prefix modulo \equiv , and when there exists a letter x such that Δ_1 and Δ_2 begin with x and \bar{x} respectively, if $\Delta_1 \equiv \Delta_2$, then $\Delta_1 = x \prod_{n\geq 1} v_n \check{x}_n$, $\Delta_2 = \bar{x} \prod_{n\geq 1} \bar{v}_n \hat{x}_n$ for an L-spinned letter x, a sequence $(v_n)_{n\geq 1}$ of x-free L-spinned words, and sequences of spinned letters $(\check{x}_n)_{n\geq 1}$, $(\hat{x}_n)_{n\geq 1}$ in $\{x, \bar{x}\}$ such that $(\check{x}_n)_{n\geq 1}$ contains infinitely many times the R-spinned letter \bar{x} , and $(\hat{x}_n)_{n\geq 1}$ contains infinitely many times the L-spinned letter x.

The relation \rightsquigarrow (and hence the block-equivalence \equiv for infinite words) is rather intricate to understand. So in some way the forms of Δ_1 and Δ_2 at the end of Theorem 3.2 are, although technical, easier to understand. Theorem 4.1, which refines the end of the previous result and proves the converse, describes all possible forms for pairs of directive-equivalent words without any use of notations \rightsquigarrow and \equiv .

When one of the two considered directive words is not wavy, Justin and Pirillo established:

Proposition 3.3. [15, Prop. 3.6] Let Δ_1 and Δ_2 be spinned versions of a common word such that Δ_1 is wavy and letters of Δ_2 are ultimately of spin L (resp. ultimately of spin R). If Δ_1 and Δ_2 are directive-equivalent, then $\Delta_1 \rightsquigarrow \Delta_2$. Moreover there exist spinned words w_1 , w_2 , an L-spinned letter x, and L-spinned x-free words $(v_i)_{i\geq 1}$ such that $\mu_{w_1} = \mu_{w_2}$, $\Delta_1 = w_1 \bar{x} \prod_{i\geq 1} \bar{v}_i x$ and $\Delta_2 = w_2 x \prod_{i\geq 1} v_i x$ (resp. $\Delta_1 = w_1 x \prod_{i\geq 1} v_i \bar{x}$ and $\Delta_2 = w_2 \bar{x} \prod_{i\geq 1} v_i x$ (resp. $\Delta_1 = w_1 x \prod_{i\geq 1} v_i \bar{x}$ and $\Delta_2 = w_2 \bar{x} \prod_{i\geq 1} \bar{v}_i \bar{x}$).

With the next two results, they considered the remaining cases of words directing aperiodic epistumian words. In the first one, the spins of the letters in each of the two directive words are ultimately L or ultimately R. The second result shows that if one of the directive words has the spins of its letters ultimately L (resp. ultimately R), then the other directive word cannot have the spins of its letters ultimately R (resp. ultimately L).

Proposition 3.4. [15, Prop. 3.7] Let Δ_1 and Δ_2 be spinned versions of a common word $\Delta \in \mathcal{A}^{\omega}$ with $|\text{Ult}(\Delta)| > 1$. If there exist spinned words w_1, w_2 and an L-spinned infinite word Δ' such that $\Delta_1 = w_1 \Delta'$ and $\Delta_2 = w_2 \Delta'$ (resp. $\Delta_1 = w_1 \overline{\Delta}'$ and $\Delta_2 = w_2 \overline{\Delta}'$), then Δ_1, Δ_2 are directive-equivalent if and only if $\mu_{w_1} = \mu_{w_2}$. **Proposition 3.5.** [15, Prop. 3.9] Let Δ be an L-spinned infinite word. Then Δ and Δ do not direct a common right-infinite episturmian word.

Actually the previous statement is a corollary of Proposition 3.9 in [15] which considers more generally words directing episturmian words differing only by a shift.

Justin and Pirillo also discussed in [15] the periodic case and proved:

Proposition 3.6. [15, Prop. 3.10] Suppose that $\Delta_1 = \breve{w}\breve{y}a^{\omega}$ and $\Delta_2 = \hat{w}\hat{y}\bar{a}^{\omega}$, where \breve{w} and \hat{w} (resp. \breve{y} and \hat{y}) are spinned versions of a common word and a is an L-spinned letter. Then Δ_1 and Δ_2 are directive-equivalent if and only if there exist sequences of letters $(\breve{a}_n)_{n\geq 1}$ and $(\hat{a}_n)_{n\geq 1}$ such that $\breve{w}\breve{y}\prod_{n>1}\breve{a}_n \equiv \hat{w}\hat{y}\prod_{n>1}\hat{a}_n$.

We will see in Theorem 4.1 that other cases can occur for periodic episturmian words.

4 Directive-equivalent words: a characterization

As shown in the previous section, Justin and Pirillo provided quite complete results about directiveequivalent infinite words. Nevertheless they did not systematically provide the relative forms of two directive-equivalent words. The following characterization does it, moreover without the use of relations \rightsquigarrow and \equiv . This result also fully solves the periodic case, which was only partially solved in [15].

Theorem 4.1. Given two spinned infinite words Δ_1 and Δ_2 , the following assertions are equivalent.

- i) Δ_1 and Δ_2 direct a common right-infinite episturnian word;
- ii) Δ_1 and Δ_2 direct a common bi-infinite episturmian word;
- iii) One of the following cases holds for some i, j such that $\{i, j\} = \{1, 2\}$:
 - 1. $\Delta_i = \prod_{n \ge 1} v_n$, $\Delta_j = \prod_{n \ge 1} z_n$ where $(v_n)_{n \ge 1}$, $(z_n)_{n \ge 1}$ are spinned words such that $\mu_{v_n} = \mu_{z_n}$ for all $n \ge 1$;
 - 2. $\Delta_i = wx \prod_{n \ge 1} v_n \check{x}_n, \ \Delta_j = w' \bar{x} \prod_{n \ge 1} \bar{v}_n \hat{x}_n$ where w, w' are spinned words such that $\mu_w = \mu_{w'}, x \text{ is an } L$ -spinned letter, $(v_n)_{n \ge 1}$ is a sequence of non-empty x-free L-spinned words, and $(\check{x}_n)_{n \ge 1}, \ (\hat{x}_n)_{n \ge 1}$ are sequences of non-empty spinned words over $\{x, \bar{x}\}$ such that, for all $n \ge 1$, $|\check{x}_n| = |\hat{x}_n|$ and $|\check{x}_n|_x = |\hat{x}_n|_x$;
 - 3. $\Delta_1 = w\mathbf{x}$ and $\Delta_2 = w'\mathbf{y}$ where w, w' are spinned words, x and y are letters, and $\mathbf{x} \in \{x, \bar{x}\}^{\omega}, \mathbf{y} \in \{y, \bar{y}\}^{\omega}$ are spinned infinite words such that $\mu_w(x) = \mu_{w'}(y)$.

Note. For a, b, c three different letters in \mathcal{A} , the spinned infinite words $\Delta_1 = a(bc\bar{a})^{\omega}$ and $\Delta_2 = \bar{a}(\bar{b}c\bar{a})^{\omega}$ direct a common epistumian word that starts with the letter a. Indeed, these two directive words fulfill item 2 of Theorem 4.1 with $w = w' = \varepsilon$, x = a, and for all $n, v_n = bc$ and $\check{x}_n = \hat{x}_n = \bar{a}$. Moreover the fact that Δ_1 starts with the *L*-spinned letter a shows that the word it directs starts with a. Similarly $\Delta'_1 = \bar{a}b(ca\bar{b})^{\omega}$ and $\Delta'_2 = \bar{a}\bar{b}(c\bar{c}a\bar{b})^{\omega}$ direct a common epistumian word starting with the letter b. Since $\Delta_2 = \Delta'_2$, this shows that the relation "direct a common epistumian word" over spinned infinite word is not an equivalence relation.

Items 2 and 3 of Theorem 4.1 show that any episturmian word is directed by a spinned infinite word having infinitely many *L*-spinned letters, but also by a spinned word having both infinitely many *L*-spinned letters and infinitely many *R*-spinned letters (i.e., a wavy word). To emphasize the importance of these facts, let us recall from Proposition 2.2 that if $\check{\Delta}$ is a spinned infinite word over $\mathcal{A} \cup \bar{\mathcal{A}}$ with infinitely many *L*-spinned letters, then there exists a unique episturmian word **t** directed by $\check{\Delta}$. Unicity comes from the fact that the first letter of **t** is fixed by the first *L*-spinned letter in $\check{\Delta}$.

Before proving Theorem 4.1, let us make two more remarks.

Remark 4.2. In items 1 and 2 of Theorem 4.1, the two considered directive words are spinned versions of a common *L*-spinned word. This does not hold in item 3, which deals only with periodic episturmian words. This is consistent with Remark 2.5. As an example of item 3, one can consider the word $(ab)^{\omega} = L_a(b^{\omega}) = R_b(a^{\omega})$ which, as already said at the end of Section 2.4, is directed by ab^{ω} and by $\bar{b}a^{\omega}$ $(L_a(b) = ab = R_b(a))$. Note also that $(ab)^{\omega}$ is directed by $(a\bar{b})^{\omega}$, underlining the fact that x and y can be equal in item 3 of Theorem 4.1.

Remark 4.3. If an episturmian word **t** has two directive words satisfying items 2 or 3, then **t** has infinitely many directive words. Indeed, if item 2 is satisfied and \bar{x} occurs in \check{x}_p $(p \ge 1)$, then by Theorem 3.1, $x\left(\prod_{k=1}^{p-1} v_n \check{x}_n\right) v_p \check{x}'_p \bar{x} \equiv \bar{x} \left(\prod_{k=1}^{p-1} \bar{v}_n \check{x}_n\right) \bar{v}_p \check{x}'_p x$ where \check{x}'_p is such that $\check{x}_p \equiv \bar{x}\check{x}'_p$. Thus **t** is also directed by $w\bar{x}\left(\prod_{k=1}^{p-1} \bar{v}_n \check{x}_n\right) \bar{v}_p \check{x}'_p x \prod_{n\ge p+1} v_n \check{x}_n$. Similarly if item 2 is satisfied and x occurs in \check{x}_p $(p \ge 1)$, then **t** is also directed by $w'x\left(\prod_{k=1}^{p-1} v_n \hat{x}_n\right) v_p \check{x}'_p x \prod_{n\ge p+1} v_n \hat{x}_n$ where \hat{x}'_p is such that $\hat{x}_p \equiv x \hat{x}'_p$. If item 3 is satisfied, then **t** is periodic and directed by $w\mathbf{x}$ where **x** is any spinned version of x^{ω} .

The rest of this section is dedicated to the proof of Theorem 4.1.

Proof of Theorem 4.1. We have i \Leftrightarrow ii) by the remarks on bi-infinite words at the beginning of Section 3.2.

 $iii) \Rightarrow i$). Assume first that $\Delta_1 = \prod_{n \ge 1} v_n$ and $\Delta_2 = \prod_{n \ge 1} z_n$ for spinned words $(v_n)_{n \ge 1}, (z_n)_{n \ge 1}$ such that $\mu_{v_n} = \mu_{z_n}$ for all $n \ge 1$. From the latter equality and Theorem 3.1, Δ_1 has infinitely many *L*-spinned letters if and only if Δ_2 has infinitely many *L*-spinned letters.

Let us first consider the case when both Δ_1 and Δ_2 have infinitely many *L*-spinned letters. Without loss of generality we can assume that for all $n \geq 1$, v_n and z_n contain at least one *L*-spinned letter. Now we need to define some more notations. Let \mathbf{t}_1 and \mathbf{t}_2 be the episturmian words directed by Δ_1 and Δ_2 , respectively (these episturmian words exist and are unique by Proposition 2.2). For $n \geq 0$, let $\mathbf{t}_1^{(n)}$ and $\mathbf{t}_2^{(n)}$ be the episturmian words as in *ii*) of Theorem 2.1 and let a_n and b_n be their respective first letters. Finally, for $n \geq 1$, set $p_n = \prod_{i=1}^n v_i$ and $q_n = \prod_{i=1}^n z_i$. The words $\mu_{p_n}(a_{|p_n|})$ (resp. $\mu_{q_n}(b_{|q_n|})$) are prefixes of \mathbf{t}_1 (resp. of \mathbf{t}_2). The letter $a_{|p_n|}$ (resp. $b_{|q_n|}$) is the first letter of $\mu_{v_{n+1}}(\mathbf{t}_1^{(m)})$ (resp. $\mu_{z_{n+1}}(\mathbf{t}_2^{(m)})$) with $m = \sum_{i=1}^{n+1} |v_i| = \sum_{i=1}^{n+1} |z_i|$. Since v_{n+1} (resp. $z_{n+1}(w)$) for any word w. From $\mu_{v_{n+1}} = \mu_{z_{n+1}}$, we have $a_{|p_n|} = b_{|q_n|}$ and so $\mu_{p_n}(a_{|p_n|}) = \mu_{q_n}(b_{|q_n|})$ for all $n \geq 1$. If the sequence $(\mu_{p_n}(a_{|p_n|}))_{n\geq 1}$ is not ultimately constant, then from $\mathbf{t}_1 = \lim_{n\to\infty} \mu_{p_n}(a_{|p_n|})$ and $\mathbf{t}_2 = \lim_{n\to\infty} \mu_{q_n}(b_{|q_n|})$, we deduce that $\mathbf{t}_1 = \mathbf{t}_2$. If $(\mu_{p_n}(a_{|p_n|}))_{n\geq 1}$ is ultimately constant, then necessarily there exists a letter a and an integer m such that for all n > m, v_n and z_n belong to $\{a\}^*$. Then $\mathbf{t}_1 = \mu_{v_1...v_m}(a^\omega) = \mu_{z_1...z_m}(a^\omega) = \mathbf{t}_2$.

Now, with the same notations as in the above case, we consider the case when the letters of Δ_1 and Δ_2 are ultimately *R*-spinned. By Theorem 3.1, any equality $\mu_v = \mu_z$ (for some different spinned words v and z) implies that v and z both contain at least one *L*-spinned letter and one *R*-spinned letter. Hence, in our current case, there exists an integer m such that $v_n = z_n$ for all n > m. Let **t** be an episturmian word directed by $\prod_{n>m} v_n = \prod_{n>m} z_n$ (such an episturmian word exists by Proposition 2.2). Then $\mu_{p_m}(\mathbf{t}) = \mu_{q_m}(\mathbf{t})$ and Δ_1 and Δ_2 are directive-equivalent.

Now consider item 2 of part *iii*). We assume that $\Delta_1 = wx \prod_{n \ge 1} v_n \check{x}_n$ and $\Delta_2 = w' \bar{x} \prod_{n \ge 1} \bar{v}_n \hat{x}_n$ where w, w' are spinned words such that $\mu_w = \mu_{w'}, x$ is an *L*-spinned letter, $(v_n)_{n\ge 1}$ is a sequence of non-empty *x*-free *L*-spinned words, and $(\check{x}_n)_{n\ge 1}, (\hat{x}_n)_{n\ge 1}$ are non-empty spinned words over $\{x, \bar{x}\}$ such that, for all $n \ge 1$, $|\check{x}_n| = |\hat{x}_n|$ and $|\check{x}_n|_x = |\hat{x}_n|_x$. By injectivity of the morphisms $\mu_w = \mu_{w'},$ Δ_1 and Δ_2 are directive-equivalent if and only if $w^{-1}\Delta_1$ and $w'^{-1}\Delta_2$ are directive-equivalent. So, from now on, we assume without loss of generality that $w = w' = \varepsilon$.

By Proposition 2.2, there exist unique episturmian words \mathbf{t}_1 and \mathbf{t}_2 starting with x directed by the respective words Δ_1 and Δ_2 (observe that if $\hat{x}_n \in \bar{x}^+$ for all $n \ge 1$, then $\bar{x} \in \text{Ult}(\Delta_2)$). For $i \geq 1$, let $\Delta_1^{(i)} = x \prod_{n \geq i} v_n \breve{x}_n$ and $\Delta_2^{(i)} = \overline{x} \prod_{n \geq i} \overline{v}_n \widehat{x}_n$ and let $\mathbf{t}_1^{[i]}$ and $\mathbf{t}_2^{[i]}$ be the words beginning with x and directed by the respective words $\Delta_1^{(i)}$ and $\Delta_2^{(i)}$. (The episturmian words $\mathbf{t}_1^{[i]}$ and $\mathbf{t}_2^{[i]}$ exist by Proposition 2.2.) For $i \geq 1$ we also define $\alpha_i := |\breve{x}_i|_x = |\hat{x}_i|_x$ and $\beta_i := |\breve{x}_i|_{\overline{x}} = |\hat{x}_i|_{\overline{x}}$. Assume first that $\alpha_i \neq 0$. Then $\overline{x}\overline{v}_i \widehat{x}_i \equiv \overline{x}\overline{v}_i x x^{\alpha_i - 1} \overline{x}^{\beta_i} \equiv x v_i \overline{x} x^{\alpha_i - 1} \overline{x}^{\beta_i} \equiv x v_i \overline{x}^{\alpha_i - 1} \overline{x}^{\beta_i} \overline{x}$ and

Assume first that $\alpha_i \neq 0$. Then $\bar{x}\bar{v}_i\hat{x}_i \equiv \bar{x}\bar{v}_ixx^{\alpha_i-1}\bar{x}^{\beta_i} \equiv xv_i\bar{x}x^{\alpha_i-1}\bar{x}^{\beta_i} \equiv xv_ix^{\alpha_i-1}\bar{x}^{\beta_i}\bar{x}$ and $xv_i\check{x}_i \equiv xv_ix^{\alpha_i-1}\bar{x}^{\beta_i}x$. Let $p_i = xv_ix^{\alpha_i-1}\bar{x}^{\beta_i}$. From what precedes we deduce that $\Delta_1^{(i)}$ and $p_i\Delta_1^{(i+1)}$ are directive-equivalent. By the choice of words $\mathbf{t}_1^{[i]}$ and $\mathbf{t}_2^{[i]}$, we deduce that $\mathbf{t}_1^{[i]} = \mu_{p_i}(\mathbf{t}_1^{[i+1]})$ and $\mathbf{t}_2^{[i]} = \mu_{p_i}(\mathbf{t}_2^{[i+1]})$ and each of these words starts with $\mu_{p_i}(x)$.

Now let us consider the case when $\alpha_i = 0$. Then $\check{x}_i = \hat{x}_i = \bar{x}^{\beta_i}$. We have $xv_i\check{x}_i \equiv \bar{x}\bar{v}_i\bar{x}^{\beta_i-1}x$ and $\bar{x}\bar{v}_i\hat{x}_i = \bar{x}\bar{v}_i\bar{x}^{\beta_i-1}\bar{x}$. Taking $p_i = \bar{x}\bar{v}_i\bar{x}^{\beta_i-1}$, we reach the same conclusion as in the case when $\alpha_i \neq 0$.

It follows from what precedes that \mathbf{t}_1 and \mathbf{t}_2 both start with $\mu_{p_1\dots p_i}(x)$ for all $i \ge 1$. Since $v_i \ne \varepsilon$, the sequence $(\mu_{p_1\dots p_i}(x))_{i\ge 1}$ is not ultimately constant; whence $\mathbf{t}_1 = \mathbf{t}_2 = \lim_{i\to\infty} \mu_{p_1\dots p_i}(x)$.

Lastly, assume that $\Delta_1 = w\mathbf{x}$ and $\Delta_2 = w'\mathbf{y}$ for some spinned words w, w', some letters x and y, and some spinned infinite words $\mathbf{x} \in \{x, \bar{x}\}^{\omega}$, $\mathbf{y} \in \{y, \bar{y}\}^{\omega}$ such that $\mu_w(x) = \mu_{w'}(y)$. The word Δ_1 (resp. Δ_2) directs the episturmian word $\mu_w(x^{\omega}) = (\mu_w(x))^{\omega}$ (resp. $\mu_{w'}(y^{\omega}) = (\mu_{w'}(y))^{\omega}$). Hence Δ_1 and Δ_2 are directive-equivalent.

 $i) \Rightarrow iii$). Suppose Δ_1 and Δ_2 direct a common (right-infinite) episturmian word **t**. Let us first assume that **t** is aperiodic. Then, by Remark 2.5, Δ_1 and Δ_2 are spinned versions of a common infinite word $\Delta \in \mathcal{A}^{\omega}$. We now show that item 1 or item 2 holds using results of Justin and Pirillo in [15].

First consider the case when both Δ_1 and Δ_2 are wavy. Suppose there exist a sequence of prefixes $(p_n)_{n\geq 0}$ of Δ_1 and a sequence of prefixes $(p'_n)_{n\geq 0}$ of Δ_2 such that for all $n\geq 0$, $\mu_{p_n}=\mu_{p'_n}$. Without loss of generality we can assume that $p_0 = p'_0 = \varepsilon$ and the sequence $(|p_n|)_{n\geq 0}$ is strictly increasing. For $n\geq 1$, let v_n , z_n be such that $p_n = p_{n-1}v_n$, $p'_n = p'_{n-1}z_n$; that is $\Delta_1 = \prod_{n\geq 1} v_n$ and $\Delta_2 = \prod_{n\geq 1} z_n$. Let us prove by induction that $\mu_{v_n} = \mu_{z_n}$ for all $n\geq 1$. First $\mu_{v_1} = \mu_{p_1} = \mu_{p'_1} = \mu_{z_1}$. For $n\geq 2$, since $\mu_{p_n} = \mu_{p_{n-1}}\mu_{v_n}$, $\mu_{p'_n} = \mu_{p'_{n-1}}\mu_{z_n}$, $\mu_{p_n} = \mu_{p'_n}$ and $\mu_{p_{n-1}} = \mu_{p'_{n-1}}$, we have $\mu_{p_{n-1}}\mu_{v_n} = \mu_{p_{n-1}}\mu_{z_n}$ by left cancellativity of the monoid of episturmian morphisms. So item 1 is satisfied in this case.

Now assume that previous sequences $(p_n)_{n\geq 0}$ and $(p'_n)_{n\geq 0}$ do not exist. Let w and w' be the longest prefixes of the respective spinned words Δ_1 and Δ_2 such that $\mu_w = \mu_{w'}$. Further, let Δ'_1 and Δ'_2 be the spinned words such that $\Delta_1 = w\Delta'_1$ and $\Delta_2 = w'\Delta'_2$. Then, by injectivity of μ_w , the words Δ'_1 and Δ'_2 are directive-equivalent and have no prefixes with equal images by μ .

By Theorem 3.2, there exists a letter x in \mathcal{A} , a sequence of non-empty x-free words $(v_n)_{n\geq 1}$ over \mathcal{A} , and two sequences of non-empty words $(\check{x}_n)_{n\geq 1}$, $(\hat{x}_n)_{n\geq 1}$ over $\{x, \bar{x}\}$ such that $\Delta'_i = x \prod_{n\geq 1} v_n \check{x}_n$ and $\Delta'_j = \bar{x} \prod_{n\geq 1} \bar{v}_n \hat{x}_n$ for some integers i, j such that $\{i, j\} = \{1, 2\}$. We have to prove that for all $n \geq 1$, $|\check{x}_n| = |\hat{x}_n|$ and $|\check{x}_n|_x = |\hat{x}_n|_x$. We use induction on n and prove also that for all $n \geq 0$, the words $\Delta_i^{(n+1)} = x \prod_{m\geq n+1} v_m \check{x}_m$ and $\Delta_j^{(n+1)} = \bar{x} \prod_{m\geq n+1} \bar{v}_m \hat{x}_m$ are directive-equivalent.

Let $n \ge 1$ be an integer. By definition of $\Delta_i^{(1)} = \overline{\Delta}_i'$ and $\Delta_j^{(1)} = \Delta_j'$ (when n = 1) and by the induction hypothesis (when $n \ge 2$), we know that the words $\Delta_i^{(n)} = x \prod_{m \ge n} v_m \breve{x}_m$ and $\Delta_j^{(n)} = \overline{x} \prod_{m \ge n} \overline{v}_m \hat{x}_m$ are directive-equivalent.

Assume first that \hat{x}_n contains at least one occurrence of x. Then, with $\alpha_n = |\hat{x}_n|_x$ and $\beta_n = |\hat{x}_n|_{\bar{x}}$, we have $\bar{x}\bar{v}_n\hat{x}_n \equiv \bar{x}\bar{v}_nxx^{\alpha_n-1}\bar{x}^{\beta_n} \equiv xv_nx^{\alpha_n-1}\bar{x}^{\beta_n}\bar{x}$. By injectivity of the morphism μ_{xv_n} we deduce that the words $\check{x}_n \prod_{m \ge n+1} v_m \check{x}_m = \check{x}_n v_{n+1} \check{x}_{n+1} \prod_{m \ge n+2} v_m \check{x}_m$ and $x^{\alpha_n-1}\bar{x}^{\beta_n}\bar{x}\prod_{m\ge n+1}\bar{v}_m\hat{x}_m = x^{\alpha_n-1}\bar{x}^{\beta_n+1}\bar{v}_{n+1}\hat{x}_{n+1}\prod_{m\ge n+2}\bar{v}_m\hat{x}_m$ direct a common episturmian word \mathbf{t}_n . The word v_{n+1} is not empty. Let c be its first letter, let $D = c^{-1}v_{n+1}\check{x}_{n+1}\prod_{m\ge n+2}v_m\check{x}_m$ and let $D' = (\bar{c})^{-1}\bar{v}_{n+1}\hat{x}_{n+1}\prod_{m\ge n+2}\bar{v}_m\hat{x}_m$: the word \mathbf{t}_n is directed by \check{x}_ncD and by $x^{\alpha_n-1}\bar{x}^{\beta_n+1}\bar{c}D'$. Since Δ_j is wavy, D' is also wavy. So x occurs in D' (among the \hat{x}_n) and the word directed by D' starts with x. Consequently \mathbf{t}_n starts with $\mu_{x^{\alpha_n-1}\bar{x}^{\beta_n+1}\bar{c}(x) = x^{\alpha_n}cx^{\beta_{n+1}}$. The words

 v_m are non-empty, thus there exists a letter $d \neq x$ that occurs in the word directed by D'. Consequently $cx^{\alpha_n+\beta_n}d$ is the smallest factor of \mathbf{t}_n belonging to $c\{x\}^*d$. Since \mathbf{t}_n is also directed by $\check{x}_n cD$, it follows that \mathbf{t}_n starts with $x^{|\check{x}_n|_x}c$ and the smallest factor of \mathbf{t}_n belonging to $c\{x\}^*d$ is $cx^{|\check{x}_n|}d$. Hence $|\check{x}_n|_x = \alpha_n = |\hat{x}_n|_x$ and $|\check{x}_n| = \alpha_n + \beta_n = |\hat{x}_n|$. Consequently $\check{x}_n \equiv x^{\alpha_n} \bar{x}^{\beta_n} \equiv x^{\alpha_n - 1} \bar{x}^{\beta_n} x$. The injectivity of the morphism $\mu_{x^{\alpha_n - 1} \bar{x}^{\beta_n}}$ implies that $\hat{\Delta}_{i}^{(n+1)} = x \prod_{m \ge n+1} v_m \breve{x}_m \text{ and } \Delta_{j}^{(n+1)} = \overline{x} \prod_{m \ge n+1} \overline{v}_m \hat{x}_m \text{ are directive-equivalent.}$ When \breve{x}_n contains at least one occurrence of \overline{x} , we similarly reach the same conclusion.

Now we show that it is impossible that $\check{x}_n \in x^+$ and $\hat{x}_n \in \bar{x}^+$. Assume these relations hold and let k be the least integer strictly greater than n such that $x \in Alph(\hat{x}_k)$ (such an integer exists since Δ_i) is wavy). Let $\alpha_k = |\hat{x}_k|_x$ and $\beta_k = |\hat{x}_k|_{\bar{x}}$. Since all of the words $\hat{x}_n, \ldots, \hat{x}_{k-1}$ belong to \bar{x}^+ , we have $\bar{x}\bar{v}_n\hat{x}_n\bar{v}_{n+1}\ldots\bar{v}_k\hat{x}_k \equiv \bar{x}\bar{v}_n\hat{x}_n\bar{v}_{n+1}\ldots\bar{v}_kx^{\alpha_k}\bar{x}^{\beta_k} \equiv xv_n\hat{x}_nv_{n+1}\ldots v_k\bar{x}x^{\alpha_k-1}\bar{x}^{\beta_k}.$ Then by injectivity of the morphism μ_{xv_n} , there exists an episturmian word directed by both $\Delta = \breve{x}_n \prod_{m \ge n+1} v_m \breve{x}_m$ and $\Delta' = \hat{x}_n v_{n+1} \dots v_k x^{\alpha_k - 1} \bar{x}^{\beta_k} \bar{x} \prod_{m > k+1} \bar{v}_m \hat{x}_m$. But this is impossible since Δ directs a word starting with x (recall that $\check{x}_n \in x^+$) and Δ' directs a word starting with the first letter of v_{n+1} (recall that $\hat{x}_n \in \bar{x}^+$).

Let us now consider the case when one of the two words Δ_1 , Δ_2 is wavy and the other has all of its spins ultimately L or ultimately R. Then item 2 is verified by Proposition 3.3.

Suppose now that both Δ_1 and Δ_2 have all spins ultimately L (resp. ultimately R). Then by Remark 2.5, Δ_1 and Δ_2 are spinned versions of a common word. Hence $\Delta_1 = w\Delta$ and $\Delta_2 = w'\Delta$ (resp. $\Delta_1 = w\Delta$ and $\Delta_2 = w'\Delta$) for some spinned words w, w' of the same length and an infinite L-spinned word Δ (resp. R-spinned word Δ). Since Δ_1 and Δ_2 are directive-equivalent, $\mu_w = \mu_{w'}$ by Proposition 3.4, and furthermore Δ_1 and Δ_2 have infinitely many prefixes whose images are equal by μ . Therefore, as already seen, this situation satisfies item 1.

We have now ended the study of the aperiodic case, since by Proposition 3.5, Δ_1 and Δ_2 cannot direct a common aperiodic episturmian word if one of them has all spins ultimately L and the other has all spins ultimately R.

Finally we come to the periodic case: $\Delta_1 = w\mathbf{x}$ and $\Delta_2 = w'\mathbf{y}$ for some spinned words w, w', letters x and y, and spinned infinite words $\mathbf{x} \in \{x, \bar{x}\}^{\omega}$, $\mathbf{y} \in \{y, \bar{y}\}^{\omega}$. In this case, the episturmian word directed by Δ_1 and Δ_2 is $\mu_w(x)^{\omega} = \mu_{w'}(y)^{\omega}$, which implies that $\mu_w(x)^{|y|} = \mu_{w'}(y)^{|x|}$. Then (see [19] for instance) there exists a primitive word z such that $\mu_w(x)$ and $\mu_{w'}(y)$ are powers of z (let us recall that a word w is primitive if it is not an integer power of a shorter word, i.e., if $w = u^p$ with $p \in \mathbb{N}$, then p = 1 and w = u). One can quite easily verify that any epistumian morphism maps any primitive word to another primitive word (see also [13, Prop. 2.8, Prop. 3.15]). Since any letter constitutes a primitive word, both $\mu_w(x)$ and $\mu_{w'}(y)$ are primitive. Thus $\mu_w(x) = z = \mu_{w'}(y)$.

5 Normalized directive word of an episturmian word

In the previous section we have seen that any episturmian word t has a directive word with infinitely many L-spinned letters. To work on Sturmian words, Berthé, Holton, and Zamboni recently proved that it is always possible to choose a particular directive word:

Theorem 5.1. [4] Any Sturmian word w over $\{a, b\}$ has a unique representation of the form

$$\mathbf{w} = \lim_{n \to \infty} L_a^{d_1 - c_1} R_a^{c_1} L_b^{d_2 - c_2} R_b^{c_2} \dots L_a^{d_{2n-1} - c_{2n-1}} R_a^{c_{2n-1}} L_b^{d_{2n} - c_{2n}} R_b^{c_{2n}}(a)$$

where $d_k \ge c_k \ge 0$ for all integer $k \ge 1$, $d_k \ge 1$ for $k \ge 2$ and if $c_k = d_k$ then $c_{k-1} = 0$.

In other words, any Sturmian word has a unique directive word over $\{a, b, \bar{a}, \bar{b}\}$ containing infinitely many L-spinned letters but no factor of the form $\bar{a}\bar{b}^n a$ or $\bar{b}\bar{a}^n b$ with n an integer. Actually this result is quite natural if one thinks about the presentation of the monoid of Sturmian morphisms (see [31]). Using Theorems 3.1 and 4.1, we generalize Theorem 5.1 to episturnian words:

Theorem 5.2. Any episturmian word **t** has a spinned directive word containing infinitely many *L*-spinned letters, but no factor in $\bigcup_{a \in A} \bar{a} \bar{A}^* a$. Such a directive word is unique if **t** is aperiodic.

The example given in Remark 4.2 shows that unicity does not necessarily hold for periodic episturmian words. A directive word of an aperiodic episturmian word \mathbf{t} with the above property is called the *normalized directive word* of \mathbf{t} . We extend this definition to morphisms: a finite spinned word w is said to be a *normalized directive word* of the morphism μ_w if w has no factor in $\bigcup_{a \in A} \bar{a} \bar{A}^* a$.

One can observe that, by Theorem 3.1, for any morphism in $L_a \mathcal{L}^*_{\mathcal{A}} R_a$, we can find another decomposition of the morphism in the set $R_a \mathcal{R}^*_{\mathcal{A}} L_a$. Equivalently, for any spinned word in $a\mathcal{A}^*\bar{a}$, there exists a word w' in $\bar{a}\bar{\mathcal{A}}^*a$ such that $\mu_w = \mu_{w'}$. This is the main idea used in the proof of the lemma below. The proof of Theorem 5.2 is based on an extension of this lemma to infinite words.

Lemma 5.3. Any pure episturnian morphism has a unique normalized directive word.

Proof. Existence of the normalized directive word: Let $w = (w_i)_{1 \le i \le |w|}$ be a spinned word over $\mathcal{A} \cup \overline{\mathcal{A}}$. We construct by induction on |w| a normalized directive word of μ_w .

If |w| = 0, there is nothing to do: ε is a normalized directive word of the empty morphism. Assume we have constructed a normalized directive word $w' = (w'_i)_{1 \le i \le k}$ of the morphism $\mu_{w'} = \mu_w$.

Let \bar{x} be a letter in \mathcal{A} . Then, by normalization of w', the word $w'\bar{x}$ has no factor in $\bigcup_{a \in \mathcal{A}} \bar{a}\mathcal{A}^*a$. Moreover since $\mu_w = \mu_{w'}$, we have $\mu_{w\bar{x}} = \mu_{w'\bar{x}}$: the word $w'\bar{x}$ is a normalized directive word of $\mu_{w\bar{x}}$.

Now let x be a letter in \mathcal{A} . The word w'x can have factors in $\bigcup_{a \in \mathcal{A}} \bar{a} \bar{\mathcal{A}}^* a$, but only as suffixes. If this does not hold, as in the previous case, the word w'x is a normalized directive word of μ_{wx} . Else $w' = p\bar{x}\bar{u}_1\bar{x}\bar{u}_2\ldots\bar{x}\bar{u}_k$ for an integer $k \geq 1$, some L-spinned x-free words $(u_i)_{1\leq i\leq k}$ and a spinned word p having no suffix in $\bar{x}\bar{\mathcal{A}}^*$. The word $w''\bar{x}$ where $w'' = pxu_1\bar{x}u_2\ldots\bar{x}u_k$ contains no factor in $\bigcup_{a\in\mathcal{A}}\bar{a}\bar{\mathcal{A}}^*a$. Moreover Theorem 3.1 implies $\mu_{w'x} = \mu_{w''\bar{x}}$. Hence $w''\bar{x}$ is a normalized directive word of μ_{wx} .

Let us make a remark on the inductive construction presented in this proof:

Remark 5.4. Let u, v, u', v' be four spinned words such that u' (resp. v') is the normalized directive word obtained by the above construction from u (resp. v). If u is a prefix of v and if p is a prefix of u' ending by an L-spinned letter, then p is also a prefix of v'.

Unicity: Assume by way of contradiction that w and w' are two different spinned normalized words such that $\mu_w = \mu_{w'}$. By left cancellativity of the monoid of epistumian morphisms, we can assume that w and w' start with different letters. Moreover it follows from Theorem 3.1 that wand w' are spinned versions of a common word. Without loss of generality, we can assume that wbegins with a letter $a \in \mathcal{A}$ and w' begins with \bar{a} and so for any word z, $\mu_w(z) = \mu_{w'}(z)$ begins with a. Hence w' must start with $\bar{a}\bar{v}a$ for a word $v \in \mathcal{A}^*$. This contradicts its normalization.

Example 5.5. Let f be the pure episturmian morphism with directive word $\bar{a}\bar{b}c\bar{b}a\bar{b}\bar{a}c\bar{b}\bar{a}c\bar{a}$. By Theorem 3.1, $\mu_{\bar{a}c\bar{b}a\bar{c}a} = \mu_{\bar{a}c\bar{b}ac\bar{a}} = \mu_{acb\bar{a}c\bar{a}}$ and hence $\mu_{\bar{a}\bar{b}c\bar{b}a\bar{b}\bar{a}c\bar{b}\bar{a}c\bar{a}} = \mu_{\bar{a}\bar{b}c\bar{b}a\bar{b}acb\bar{a}c\bar{a}}$ and $\bar{a}\bar{b}c\bar{b}a\bar{b}acb\bar{a}c\bar{a}$ is the normalized directive word of f.

Now we provide the

Proof of Theorem 5.2.

Existence of the normalized directive word:

Let $\Delta = (w_i)_{i \geq 1}$ be a spinned directive word of an episturmian word **t** (with $w_i \in \mathcal{A} \cup \overline{\mathcal{A}}$). From Theorem 4.1, we can assume that Δ has infinitely many *L*-spinned letters.

By Lemma 5.3, for any $n \ge 1$, the morphism $\mu_{w_1...w_n}$ has a unique normalized directive word $(w_i^{(n)})_{1\le i\le n}$. (It follows from the proof of Lemma 5.3 that w_i and $w_i^{(n)}$ are spinned versions of a common letter).

Let p_n be the longest prefix of $w_1^{(n)} \dots w_n^{(n)}$ that belongs to $(\mathcal{A} \cup \overline{\mathcal{A}})^* \mathcal{A}$. Let $i_n \leq n$ be the integer such that $p_n = w_1^{(n)} \dots w_{i_n}^{(n)}$, and let π_n be the word $\pi_n = \mu_{w_1^{(n)} \dots w_{i_n-1}^{(n)}}(w_{i_n}^{(n)})$. Since the morphisms $\mu_{w_1 \dots w_n}$ and $\mu_{w_1^{(n)} \dots w_n^{(n)}}$ are equal, **t** has the directive word

 $(w_1^{(n)}, \ldots, w_n^{(n)}, w_{n+1}, w_{n+2}, \ldots)$, so π_n is a prefix of **t**.

By Remark 5.4, for any $n \ge 1$, p_n is a prefix of p_{n+1} , and since Δ contains infinitely many L-spinned letters, for any $n \ge 1$, there exists an m > n such that $|p_m| > |p_n|$.

If $|\text{Ult}(\Delta)| = 1$, then there exists a letter a and an integer m such that $\mathbf{t} = \mu_{p_m}(a^{\omega})$ and $p_m a^{\omega}$ is a normalized directive word of **t**. If $|\text{Ult}(\Delta)| > 1$, the sequence $(\pi_n)_{n \ge 1}$ is not ultimately constant, and $\lim_{n\to\infty} \pi_n = \mathbf{t}$. In this case \mathbf{t} is directed by the sequence $\lim_{n\to\infty} p_n$ which is normalized by construction (indeed otherwise one of the prefixes p_n would not be normalized).

Unicity of the normalized directive word:

Assume by way of contradiction that an aperiodic episturmian word \mathbf{t} has two different normalized spinned directive words $\Delta_1 = (w_n)_{n \geq 1}$ and $\Delta_2 = (w'_n)_{n \geq 1}$ (with w_n and $w'_n \in \mathcal{A} \cup \bar{\mathcal{A}}$ for all n). Let $i \geq 1$ be the smallest integer such that $w_i \neq w'_i$ (and for all $j < i, w_j = w'_j$). By Theorem 4.1, Δ_1 and Δ_2 are spinned versions of the same word (see Remark 4.2). Thus, without loss of generality, we can assume that $w_i = \bar{x}$ and $w'_i = x$ for some letter x.

Let $\mathbf{t}^{\prime(i)}$ be the epistumian word with (normalized) directive word $(w'_n)_{n>i}$ (by Proposition 2.2 this word is unique), then $\mathbf{t}^{\prime(i)}$ starts with x since $w_i = x$. Since the word $(w_n)_{n\geq 1}$ has infinitely many L-spinned letters, there exists an integer j > i such that $w_j = y$ for a letter $y \in \mathcal{A}$ and $w_\ell \in \overline{\mathcal{A}}$ for each ℓ , $i < \ell < j$. Let $\mathbf{t}^{(i)}$ be the word with normalized directive word $(w_n)_{n \ge i}$, then $\mathbf{t}^{(i)}$ has the word $\mu_{w_i...w_{j-1}}(y)$ as prefix since $w_j = y$ and so $\mathbf{t}^{(i)}$ starts with y since $w_i \ldots w_{j-1} \in \bar{\mathcal{A}}^*$. We have $\mathbf{t} = \mu_{w_1...w_{i-1}}(\mathbf{t}^{(i)}) = \mu_{w'_1...w'_{i-1}}(\mathbf{t}^{'(i)})$. By choice of $i, w_1...w_{i-1} = w'_1...w'_{i-1}$. Consequently, since episturmian morphisms are injective on infinite words, $\mathbf{t}^{(i)} = \mathbf{t}^{'(i)}$ and so x = y. But since $w_i = \bar{x}$, $w_{i+1} \dots w_{j-1} \in \overline{\mathcal{A}}^*$, and $w_j = x$, we reach a contradiction to the normalization of $(w_n)_{n \geq 1}$.

6 Episturmian words having a unique directive word

In Section 4 we have characterized pairs of words directing a common episturnian word. In Section 5 we have proposed a way to uniquely define any episturmian word through a normalization of its directives words (as mentioned in the introduction, see [4, 18, 17, 11] for some uses of this normalization). Using these results we now characterize episturnian words having a unique directive word.

Theorem 6.1. An episturnian word has a unique directive word if and only if its (normalized) directive word contains 1) infinitely many L-spinned letters, 2) infinitely many R-spinned letters, 3) no factor in $\bigcup_{a \in \mathcal{A}} \bar{a} \mathcal{A}^* a$, 4) no factor in $\bigcup_{a \in \mathcal{A}} a \mathcal{A}^* \bar{a}$.

Such an episturmian word is necessarily aperiodic.

Proof. Assume first that an episturnian word t has a unique spinned directive word Δ . By Theorem 5.2, Δ is normalized and so contains infinitely many L-spinned letters and no factor in $\bigcup_{a \in A} \bar{a} \bar{\mathcal{A}}^* a$. By item 3 of Theorem 4.1 and by Remark 2.4, t cannot be periodic. By item 2 of Theorem 4.1, Δ also contains infinitely many R-spinned letters, and hence is wavy (otherwise one can construct another directive word of \mathbf{t} – the fact that \mathbf{t} is aperiodic is important for having the $(v_n)_{n\geq 1}$ non-empty in this construction). Finally Theorem 3.1 implies the non-existence of a factor in $\bigcup_{a \in A} a \mathcal{A}^* \bar{a}$ (otherwise, one can again construct another directive word for t).

Let us now prove that the four conditions (given in the statement of the theorem) are sufficient. Arguing by contradiction, we assume that an episturmian word \mathbf{t} is directed by two spinned infinite words Δ_1 and Δ_2 , both fulfilling the four given conditions. We observe that if Δ_1 or Δ_2 is ultimately written over $\{x, \bar{x}\}$ for a letter x (which can occur only if t is periodic), then at least one of the conditions is not fulfilled. Thus the two words Δ_1 and Δ_2 should verify one of the two first items in part *iii*) of Theorem 4.1 (item 3 does not apply since **t** is aperiodic). But the hypotheses on Δ_1 and Δ_2 imply that only item 1 can be verified so that $\Delta_1 = \prod_{n \ge 1} v_n$, $\Delta_2 = \prod_{n \ge 1} z_n$ for spinned words $(v_n)_{n\ge 1}, (z_n)_{n\ge 1}$ such that $\mu_{v_n} = \mu_{z_n}$ for all $n \ge 1$. Now by Theorem 3.1 and by the fact that words $(v_n)_{n\ge 1}$ and $(z_n)_{n\ge 1}$ have no factor in $\bigcup_{a\in\mathcal{A}} \bar{a}\bar{\mathcal{A}}^*a$ nor $\bigcup_{a\in\mathcal{A}} a\mathcal{A}^*\bar{a}$, we must have $v_n = z_n$ for all $n \ge 1$. Thus $\Delta_1 = \Delta_2$.

As an example, a particular family of episturmian words having unique directive words consists of those directed by *regular wavy words*, i.e., spinned infinite words having both infinitely many *L*-spinned letters and infinitely many *R*-spinned letters such that each letter occurs with the same spin everywhere in the directive word. More formally, a spinned version \breve{w} of a finite or infinite word w is said to be *regular* if, for each letter $x \in Alph(w)$, all occurrences of \breve{x} in \breve{w} have the same spin (*L* or *R*). For example, $a\bar{b}aa\bar{c}\bar{b}$ and $(a\bar{b}c)^{\omega}$ are regular, whereas $a\bar{b}a\bar{a}\bar{c}b$ and $(a\bar{b}\bar{a})^{\omega}$ are not regular.

In the Sturmian case, we have:

Proposition 6.2. Any Sturmian word has a unique spinned directive word or infinitely many spinned directive words. Moreover, a Sturmian word has a unique directive word if and only if its (normalized) directive word is regular wavy.

Proof. Let Δ be the normalized directive word of a Sturmian word **t** over $\{a, b\}$. Then Δ contains no factor belonging to $\bar{a}\bar{b}^*a \cup \bar{b}\bar{a}^*b$ (where $\alpha^* = \{\alpha\}^*$ for any letter α).

Assume first that Δ contains infinitely many factors in $ab^*\bar{a} \cup ba^*\bar{b}$. Then $\Delta = p \bigcup_{n \ge 1} x_n y_n$ for some spinned words p and $(x_n, y_n)_{n\ge 1}$ such that, for all $n \ge 1$, $x_n \in ab^*\bar{a} \cup ba^*\bar{b}$ and $y_n \in \{a, b, \bar{a}, \bar{b}\}^*$. In this case, Δ has infinitely many directive words; indeed, the spinned words $(p[\bigcup_{n=1}^{k-1} x_n y_n]\bar{x}_k y_k \bigcup_{n>k+1} x_n y_n)_{k\ge 1}$ are (by Theorem 3.1) pairwise different directive words for t.

Now assume that Δ contains only finitely many factors in $ab^*\bar{a} \cup ba^*\bar{b}$. Since Δ contains no factor in $\bar{a}\bar{b}^*a \cup \bar{b}\bar{a}^*b$, it is ultimately regular wavy. More precisely Δ is regular wavy and either Δ belongs to $\{a, \bar{b}\}^{\omega} \cup \{\bar{a}, b\}^{\omega}$, or Δ belongs to one of the following sets of infinite words: $S_1 = \{a, b, \bar{a}, \bar{b}\}^*a\{\bar{a}, b\}^{\omega}$, $S_2 = \{a, b, \bar{a}, \bar{b}\}^*b\{a, \bar{b}\}^{\omega}$, $S_3 = \{a, b, \bar{a}, \bar{b}\}^*\bar{a}\{a, \bar{b}\}^{\omega}$ or $S_4 = \{a, b, \bar{a}, \bar{b}\}^*\bar{b}\{\bar{a}, b\}^{\omega}$. Assume $\Delta \in S_1$. Since any Sturmian word is aperiodic, Δ is not ultimately constant (see Remark 2.4). Thus $\Delta =$ $pa \bigcup_{n\geq 1} x_n \bar{a}$ with $x_n \in b^*$ for all $n \geq 1$. Once again in this case, **t** has infinitely many directive words since the words $(p[\bigcup_{n=1}^{k-1} \bar{a}\bar{x}_n]a \bigcup_{n\geq k} x_n \bar{a})_{k\geq 1}$ are pairwise different directive words for **t**. The cases when $\Delta \in S_2$ or $\Delta \in S_3$ or $\Delta \in S_4$ are similar.

We end with the case when Δ is regular wavy. In this case, Δ contains infinitely many *L*-spinned letters, infinitely many *R*-spinned letters, no factor in $ab^*\bar{a} \cup ba^*\bar{b}$, and no factor in $\bar{a}\bar{b}^*a \cup \bar{b}\bar{a}^*b$. Hence by Theorem 6.1, **t** has a unique directive word.

Proposition 6.2 shows a great difference between Sturmian words and episturmian words constructed over alphabets with at least three letters. Indeed, when considering words over a ternary alphabet, one can find episturmian words having exactly m directive words for any $m \ge 1$. For instance, the episturmian word \mathbf{t} directed by $\Delta = a(b\bar{a})^{m-1}b\bar{c}(ab\bar{c})^{\omega}$ has exactly m directive words, namely $(\bar{a}\bar{b})^i a(b\bar{a})^j b\bar{c}(ab\bar{c})^{\omega}$ with i+j=m-1. Notice that the suffix $b\bar{c}(ab\bar{c})^{\omega}$ of Δ is regular wavy, and the other m-1 spinned versions of Δ that also direct \mathbf{t} arise from the m-1 words that are block-equivalent to the prefix $a(b\bar{a})^{m-1}$.

Acknowledgement. The two last authors thank Eddy Godelle for his remarks and his suggestion for improvement of the proof of Theorem 5.2.

References

 J.-P. Allouche, J. Shallit, Automatic sequences: Theory, Applications, Generalizations, Cambridge University Press, 2003.

- [2] P. Arnoux, G. Rauzy, Représentation géométrique de suites de complexités 2n + 1, Bull. Soc. Math. France 119 (1991) 199–215.
- [3] J. Berstel, Sturmian and episturmian words (a survey of some recent results), in: Proceedings of CAI 2007, vol. 4728 of Lecture Notes in Computer Science, Springer-Verlag, 2007.
- [4] V. Berthé, C. Holton, L. Q. Zamboni, Initial powers of Sturmian sequences, Acta Arith. 122 (2006) 315–347.
- [5] X. Droubay, J. Justin, G. Pirillo, Episturmian words and some constructions of de Luca and Rauzy, Theoret. Comput. Sci. 255 (1-2) (2001) 539–553.
- [6] S. Ferenczi, Complexity of sequences and dynamical systems, Discrete Math. 206 (1999) 145–154.
- [7] A. Glen, On Sturmian and episturmian words, and related topics, Ph.D. thesis, The University of Adelaide, Australia (April 2006).
- [8] A. Glen, A characterization of fine words over a finite alphabet, Theoret. Comput. Sci. 391 (2008) 51–60.
- [9] A. Glen, J. Justin, Episturmian words: a survey, preprint, 2007.
- [10] A. Glen, J. Justin, G. Pirillo, Characterizations of finite and infinite episturmian words via lexicographic orderings, European J. Combin. 29 (2008) 45–58.
- [11] A. Glen, F. Levé, G. Richomme, Quasiperiodic and Lyndon episturnian words, preprint, 2008.
- [12] E. Godelle, Représentation par des transvections des groupes d'artin-tits, Group, Geometry and Dynamics 1 (2007) 111–133.
- [13] J. Justin, G. Pirillo, Episturmian words and episturmian morphisms, Theoret. Comput. Sci. 276 (1-2) (2002) 281–313.
- [14] J. Justin, G. Pirillo, On a characteristic property of Arnoux-Rauzy sequences, Theoret. Inform. Appl. 36 (4) (2003) 385–388.
- [15] J. Justin, G. Pirillo, Episturmian words: shifts, morphisms and numeration systems, Internat. J. Found. Comput. Sci. 15 (2) (2004) 329–348.
- [16] F. Levé, G. Richomme, Quasiperiodic infinite words: some answers, Bull. Eur. Assoc. Theor. Comput. Sci. 84 (2004) 128–138.
- [17] F. Levé, G. Richomme, Quasiperiodic episturmian words, in: Proceedings of the 6th International Conference on Words, Marseille, France, September 17-21, 2007.
- [18] F. Levé, G. Richomme, Quasiperiodic Sturmian words and morphisms, Theoret. Comput. Sci. 372 (2007) 15–25.
- [19] M. Lothaire, Combinatorics on Words, vol. 17 of Encyclopedia of Mathematics and its Applications, Addison-Wesley, 1983.
- [20] M. Lothaire, Algebraic Combinatorics on Words, vol. 90 of Encyclopedia of Mathematics and its Applications, Cambridge University Press, 2002.
- [21] M. Morse, G. Hedlund, Symbolic Dynamics II. Sturmian trajectories, Amer. J. Math. 61 (1940) 1–42.
- [22] G. Paquin, L. Vuillon, A characterization of balanced episturmian sequences, Electron. J. Combin. 14, #R33, pp. 12.
- [23] N. Pytheas Fogg, Substitutions in Dynamics, Arithmetics and Combinatorics, vol. 1794 of Lecture Notes in Mathematics, Springer, 2002.
- [24] G. Rauzy, Nombres algébriques et substitutions, Bull. Soc. Math. France 110 (1982) 147–178.
- [25] G. Rauzy, Mots infinis en arithmétique, in: M. Nivat, D. Perrin (eds.), Automata on Infinite words, vol. 192 of Lecture Notes in Computer Science, Springer-Verlag, Berlin, 1985.
- [26] G. Richomme, Conjugacy and episturmian morphisms, Theoret. Comput. Sci. 302 (1-3) (2003) 1–34.
- [27] G. Richomme, Lyndon morphisms, Bull. Belg. Math. Soc. Simon Stevin 10 (2003) 761–785.
- [28] G. Richomme, Conjugacy of morphisms and Lyndon decomposition of standard Sturmian words, Theoret. Comput. Sci. 380 (3) (2007) 393–400.
- [29] G. Richomme, A local balance property of episturmian words, in: Proc. DLT '07, vol. 4588 of Lecture Notes in Computer Science, Springer, Berlin, 2007, pp. 371–381.
- [30] R. Risley, L. Zamboni, A generalization of Sturmian sequences: combinatorial structure and transcendence, Acta Arith. 95 (2000) 167–184.
- [31] P. Séébold, Fibonacci morphisms and Sturmian words, Theoret. Comput. Sci. 88 (2) (1991) 365–384.
- [32] Z.-X. Wen, Y. Zhang, Some remarks on invertible substitutions on three letter alphabet, Chinese Sci. Bull. 44 (19) (1999) 1755–1760.