

Application of Human Factors Methods to Understand Missed Follow-up of Abnormal Test Results

Deevakar Rogith¹ Tyler Satterly^{2,3} Hardeep Singh^{2,3} Dean F. Sittig^{1,4} Elise Russo⁵
Michael W. Smith⁶ Don Roosan⁷ Viraj Bhise⁸ Daniel R. Murphy^{2,3}

¹The University of Texas Health Science Center at Houston School of Biomedical Informatics, Houston, Texas, United States

²Center for Innovations in Quality, Effectiveness and Safety, Michael E. DeBakey Veteran Affairs Medical Center, Houston, Texas, United States

³Department of Medicine, Section of Health Services Research, Baylor College of Medicine, Houston, Texas, United States

⁴UT-Memorial Hermann Center for Healthcare Quality and Safety, Houston, Texas, United States

⁵Department of Biomedical Informatics, Vanderbilt University Medical Center, Nashville, Tennessee, United States

⁶Department of Industrial and Mechanical Engineering, Universidad de las Americas Puebla, Cholula, Mexico

⁷Department of Pharmacy Practice and Administration, College of Pharmacy Western University of Health Sciences, Pomona, California, United States

⁸Department of Medicine, Geisel School of Medicine at Dartmouth, Hanover, New Hampshire, United States

Address for correspondence Deevakar Rogith, MBBS, PhD, 7000 Fannin Street Suite 600, Houston, TX 77030, United States (e-mail: Deevakar.Rogith@uth.tmc.edu).

Appl Clin Inform 2020;11:692–698.

Abstract

Objective This study demonstrates application of human factors methods for understanding causes for lack of timely follow-up of abnormal test results (“missed results”) in outpatient settings.

Methods We identified 30 cases of missed test results by querying electronic health record data, developed a critical decision method (CDM)-based interview guide to understand decision-making processes, and interviewed physicians who ordered these tests. We analyzed transcribed responses using a contextual inquiry (CI)-based methodology to identify contextual factors contributing to missed results. We then developed a CI-based flow model and conducted a fault tree analysis (FTA) to identify hierarchical relationships between factors that delayed action.

Results The flow model highlighted barriers in information flow and decision making, and the hierarchical model identified relationships between contributing factors for delayed action. Key findings including underdeveloped methods to track follow-up, as well as mismatches, in communication channels, timeframes, and expectations between patients and physicians.

Conclusion This case report illustrates how human factors-based approaches can enable analysis of contributing factors that lead to missed results, thus informing development of preventive strategies to address them.

Keywords

- ▶ contextual inquiry
- ▶ fault tree analysis
- ▶ abnormal laboratory follow-up
- ▶ human factors
- ▶ patient safety

received
March 4, 2020
accepted
August 1, 2020

© 2020 Georg Thieme Verlag KG
Stuttgart · New York

DOI <https://doi.org/10.1055/s-0040-1716537>.
ISSN 1869-0327.

Background and Significance

Lack of timely follow-up and missed follow-up of abnormal test results (henceforth, “missed results”) is a recognized patient safety concern.^{1,2} In ambulatory settings, the incidence of missed results can be up to 65% and can lead to delayed diagnosis or treatment.^{2,3} Knowledge on individual-level (vs. system-level) factors contributing to missed results is evolving. For example, it is imperative to understand individual's intent at the point of care and decision making that contribute to missed results especially within the complex sociotechnical context of electronic health records (EHRs)-enabled health systems. Human factors methods could advance understanding of physician decision-making processes and uncover factors related to individual decision-making processes.

Prior research on understanding causes of missed results is limited and has used retrospective chart reviews,² EHR activity logs,^{4,5} focus groups,^{6,7} cognitive task analyses,⁸ aggregated root cause analyses data,⁹ and safety event reporting.¹⁰ Our objective was to use human factors methods to further illustrate workflow and process issues related to missed results and focus on contributing factors. By using the human factors methods described in this study, we gathered additional detail about physician's decision-making process, illustrating factors that contribute to missed results.

We applied three human factors methods, including critical decision method (CDM)-based interviews, contextual inquiry (CI)-based analysis methodology, and fault tree analysis (FTA), to understand factors contributing to missed results. Missed follow-up of abnormal laboratory result involves complex human-system interaction factors; to uncover the related decision-making processes, we chose CDM-based interviews. Building on the decision-making process, we applied CI and FTA to illustrate contributing factors and interactions among the factors leading to missed follow-up. Such human factors methods can enhance understanding of where to focus strategies to reduce or mitigate negative outcomes.

Critical Decision Method–Based Interviews

CDM is a cognitive task analysis technique used to describe naturalistic decision making, and improves understanding of situational awareness, mental models, and decision points in particular situations.^{11–13} CDM involves gathering information about a personally experienced incident via focused interviews with task experts and identifying timelines, key decision points, and factors influencing decision making, such as clinical decision making in critical care.^{14–16} One of the limitations of the CDM method is the delay between the incident and the interview.¹⁷ By using near-real time detection of incidents, we reduce the problem of memory decay.

Contextual Inquiry–Based Analysis

The CI methodology helps in understanding the context of actions, such as physicians' responses to abnormal test results. CI is a structured methodology for modeling work domains and identifying user needs, guiding both interviews, and analysis. It focuses on four principles as follows: (1) identifying context of participants' work, (2) partnering

with participants to observe and discuss work, (3) interpreting insights and relaying them back to the participant, and (4) using the research question to guide the interactions. CI-based analysis helps generate models to represent different aspects of how work functions: communication flow and coordination, culture, task sequences, physical environment, and artifacts. CI has been used in developing tools and implementing new workflows in health care.^{18–20}

Fault Tree Analysis

FTA²¹ is a form of root cause analysis used to illustrate and analyze complex interacting pathways leading to process failures²¹ and is used for developing error prevention, monitoring, and intervention strategies.^{22–24} FTA models an outcome as a hierarchy of interacting contributing factors^{21,25} using Boolean logic operators (“AND” and “OR”). Construction of fault trees requires describing top-level outcomes and resolving them into basic (primary initiating events) and intermediate events (immediate causes for basic events).²⁶ This method enables visual analytics and probabilistic modeling of factors contributing to an outcome and has been applied in clinical use cases for studying factors related to adverse events.^{27–29}

A combination of these human factors methods could allow in-depth identification of causes for missed results and inform targeted solutions to improve decision-making processes.

Case Report

Setting

This study was performed at three large primary care clinics in Texas after Institutional Review Board approval. Each clinic used EHRs and included trainees.

Case Selection

We queried the clinical data repository at each site from January 1, 2015 to September 30, 2015 to identify abnormal imaging and laboratory results (►Table 1). A reviewer (V.B.) manually reviewed records to identify missed results, defined as lack of documented action (repeat or subsequent testing, referral placement, medication change, or patient

Table 1 List of abnormal laboratory result cases not followed-up

	Real	Vignette	Total
Chest X-ray	1		1
EKG	1	3	4
Hemoglobin	1		1
TSH		10	10
Urine albumin	2		2
Urine culture	2		2
Urine micro	8	2	10
Total	15	15	30

Abbreviations: EKG, electrocardiogram; TSH, thyroid stimulating hormone.

notification) within 14 days. We then invited 30 physicians who ordered the respective tests for interviews. We used maximum variation sampling techniques to maximize heterogeneity in clinic site and test types with each subsequent interview.

Interview

We created a CDM-based interview guide (►Appendix A) to understand follow-up in the context of a physician's own missed result cases including reasons for the miss. Questions identified factors delaying follow-up, not necessarily in the same case. Questions also both elicited specific factors contributing to missed results and identified relationships between work system and individual decision-making factors. Interviews were audio recorded and transcribed.

Three investigators (M.W.S., D.F.S., and D. Roosan) performed semistructured interviews with physicians using the CDM-based interview guide, and data were analyzed using the four CI principles. For the first 15 cases, interviewees were aware that they missed the follow-up (delay case interviews). However, this contributed to some reluctance in responding to questions about causes of missed results. To ensure responses were not constrained by recognition of their own potential oversight and enable more open discussion, we modified the method, so the remaining 15 cases were not traditional CDM interviews about the participants' own incidents, but instead vignettes similar to that experienced by their patient (vignette case interviews). The

vignettes were generated by removing identifying information about the patient, treating physician, and clinic.

Data Analysis

Two other independent reviewers (D. Rogith and T.S.) with human factors expertise analyzed data using CI-based and FTA methods. We adapted the CI to include only the flow diagram analysis. We adapted the FTA to consider all logical operators as "AND" operators.²⁶ The sociotechnical model³⁰ guided identification of factors contributing to missed results.

Reviewers first analyzed transcripts to identify underlying factors contributing to missed results. Because we aimed to identify information flow breakdowns related to missed results, reviewers used the CI-based analysis methodology to develop flow models of communication and coordination in result management decision making. Thus, the reviewers independently reviewed transcripts and identified discussion about information flow and workflow decisions related to managing the test result. These were represented in the model in terms of information flow for both people (e.g., physicians and patients) and data sources (e.g., laboratory results and EHR systems). Each reviewer then independently combined their 30 flow models into a single model before collaboratively reconciling into one final model (►Fig. 1).

Reviewers then performed FTAs to identify events leading to inaction for each case and used deductive reasoning to identify basic events from the interviews. To generate fault trees based on actual events, we chose only the 15 cases where




Fig. 1 Contextual inquiry flow model of follow-up of abnormal laboratory results. EHR, electronic health record.




Fig. 2 Fault tree analysis of events leading to no follow-up of abnormal laboratory results. EHR, electronic health record.

interviewees were aware that they had missed the follow-up, and both reviewers (D. Rogith and T.S.) independently conducted FTAs for each case to identify basic events. Basic events were then grouped into intermediate events based on the flow model described above. An interdisciplinary team discussed findings and consolidated intermediate events to generate a cumulative fault tree (► Fig. 2). Using this process, the basic and intermediate events were grouped into four categories: patient, clinical condition, physician, and EHR.

Results

Contributory causes identified from interview data are listed in ► Table 2. During delay case interviews, workflow issues were predominant (e.g., forgetting to notify patients about therapy changes based on results, diffusion of responsibility between referring physicians and residents for results follow-up, and language barriers). However, in vignette case interviews, EHR issues were more prominent. For example, limited patient portal usage led physicians to not send messages about results.

We developed a CI flow model describing physicians' processes for managing abnormal results (► Fig. 1). The flow model shows four different paths in physicians' action after test results as follows: (1) Identifying abnormal results, (2) tracking follow-up, (3) delegating follow-up, and (4) conducting follow-up. For each path, we identified barriers in the follow-up process. This

displays how physicians interact with abnormal results, their expectations for managing these results, and user requirements for completing the follow-up tasks. Key findings from the flow model included physicians' lack of methods to track follow-up and mismatch in communication channels, timeframes, and expectations between patients and physicians.

Several physicians described unwillingness to sending notifications to the EHR portals to communicate results because they felt patients may not use the portal. Some physicians reported that if the result was not acted on by a physician within a specific timeframe (e.g., 10 days), the EHR automatically released results without a physician interpretation. This removed the item from the physician's to-do list, limiting prompts to act. Furthermore, some physicians preferred only in-person communication of abnormal results at patients' next appointments, which may occur beyond the autorelease timeframe.

► Fig. 2 displays the FTA-based hierarchical model of factors contributing to missed test results, which displays the frequency of each occurrence among the 15 delay case interviews. The most common factors were physicians' assumption that ordering physicians are responsible for follow-up (5 of 15 cases). While most institutions designate responsibility for result follow-up to the ordering physician, physicians reported not being notified when resident-ordered tests returned, adding delays to follow-up. In specialty referrals, some physicians

Table 2 List of reasons for not following-up an abnormal laboratory result

Categories	Delay case interviews	Vignette case interviews
Test results	<ul style="list-style-type: none"> • Similar abnormal laboratory results in past • Results deemed abnormal but not clinically serious 	<ul style="list-style-type: none"> • Results arrive after patient's visit • Laboratory results are scanned or faxed so not available in structured data tabs • Noise in color coding of abnormal laboratory results
Physician actions	<ul style="list-style-type: none"> • Specialist expected to follow-up • Follow up deemed to be resident physician's responsibility • Communication breakdown during delegating follow-up action to be relayed by staff • No feedback from staff that abnormal results and follow-up actions communicated to patient 	<ul style="list-style-type: none"> • Forgetfulness • No dedicated staff for follow-up • Follow-up deemed an unbillable activity • Communication breakdown during delegating follow-up action to be relayed by staff • No feedback from staff that abnormal results and follow-up actions communicated to patient
Clinical actions	<ul style="list-style-type: none"> • Action taken in form of adding a clinical note or updating prescription without communication to patient • No treatment modifications necessary 	<ul style="list-style-type: none"> • Need to explain results in detail
EHR system	<ul style="list-style-type: none"> • Patient does not use portal • Unable to confirm whether patient accessed result via portal 	<ul style="list-style-type: none"> • Patient does not use portal • Unable to confirm whether patient accessed result via portal • Multiple clinics, multiple EHR systems—forgets to act on time • Time limit for auto notification is deemed to be too short (10 business days) • Not sure of difference between "communication" vs "release of abnormal laboratory result EHR features".
Other communications	<ul style="list-style-type: none"> • Inability to reach patient via phone • Physician prefers not to call patients • Language barriers • Mail—no feedback on status of mailing 	<ul style="list-style-type: none"> • Inability to reach patient via phone. No patient contact information available • Prefer direct communication (via SMS-like technology) • Mail—no feedback on status of mailing
Patient factors	<ul style="list-style-type: none"> • Patient deemed responsible for follow-up • Patient has another appointment within 2 weeks, so follow-up delayed 	<ul style="list-style-type: none"> • Patient has another appointment within 2 weeks, so follow-up delayed

Abbreviation: EHR, electronic health record.

Note: Factors common in identified cases and vignette cases are in bold.

reportedly assumed that referred specialist physicians will manage the results of the tests ordered by the referring physician because results would arrive at the time of the specialists' appointments. The FTA hierarchical model (→Fig. 2) also confirms mismatch in communication channels for follow-up (5 of 15 cases). We found that some physicians shifted responsibility for follow-up to patients, such as by releasing results electronically only when patients signed up to the organization's patient portal and/or expecting patients to schedule follow-up visits to discuss results.

These intermediate events provide a high-level cause for the individual basic events. Some intermediate events highlight professionalism issues, such as reliance on patient appointments and poor communication etiquette. One physician asserted that patients are responsible for scheduling follow-up visits within 2 weeks. In other cases, physicians preferred to wait on nonurgent results for patients' next visit if scheduled within 2 weeks. However, several physicians felt that follow-up discussion was often forgotten if patients missed or canceled visits.

Discussion

Using three complementary human factors methods, we identified causes for lack of abnormal laboratory results

follow-up. Reasons identified included physicians' expectations that patients are responsible for scheduling follow-up, mismatch in physician–patient communication preferences, and difficulties with managing abnormal results in EHR systems. These first-person accounts using human factors methods differ from prior studies in that individual decisionmaking, and workflow- and technology-related findings were prominent, allowing identification of contributing factors and barriers to action.

Combining CI-based and FTA analysis methods with CDM-based interviews permitted uncovering of contributing factors for missed results. For example, CDM-based interviews allow identification of multiple causes for inaction on abnormal results, while adding CI-based analysis enabled uncovering of contextual information such as information exchange between physician and staff (→Fig. 1). This helped categorize causes for missed results (→Table 2). Application of FTA then yielded a hierarchical model of missed results. Identifying basic contributory causes can assist in designing systems to manage abnormal test results, implementing results follow-up policies, and training clinicians to reduce breakdowns. Interestingly, we obtained richer detail with vignettes than delay cases, suggesting that physicians remain hesitant to discuss care breakdowns that they are involved in and providing guidance for future work in this area.

Limitations

Several limitations should be noted. First, our findings may be limited by socially desirability bias given the potentially sensitive topic of missed results. In vignette cases, physicians were unable to refer to their own experiences with the cases as they would in traditional CDM interviews. Additionally, findings from these three sites may not be generalizable to different practice settings and EHRs. Second, the CI-based analysis was performed by reviewers who did not perform the interviews; however, this offers a more independent assessment of findings that might not be apparent during interviews. The initial interview was collected to understand the decision making using CDM, and so for the secondary analysis using CI- and FTA-based methods, we used independent reviews. Third, CI methodology involves both observations and analysis. However, in clinical practice, it is impractical to directly observe rare events such as those under study in this care report. Nevertheless, CI-based analysis allowed useful information to be gleaned from postevent interview. Finally, we did not aim to identify specific actions to improve efficiency of test result management; however, our findings help inform future work to identify and test solutions.

Conclusion

We illustrate our application of diverse human factors methods, – CDM, CI, and FTA, to understand factors in abnormal test result follow-up. Our methods identified multiple factors contributing to missed follow-up, such as provider-patient communication channel mismatch and diffusion of responsibility. We focused on identifying barriers to successful follow-up and pathways leading to inaction. Future directions include expanding these methods to facilitate design information systems and implementation of preventive strategies to reduce missed test results.

Clinical Relevance Statement

Adverse events and care delays can occur when physicians miss taking actions on abnormal test results. However, individual decision-making factors surrounding such events are less understood. Using a combination of human factors methods described herein can identify key contributory factors that guide development of preventive interventions.

Multiple Choice Questions

1. Which method is useful in understanding information flow in decision making process?

- a. Critical decision method
- b. Contextual inquiry
- c. Process mining

Correct Answer: The correct answer is option b.

2. Which method is useful in understanding sequence of events leading to an adverse outcome?

- a. Fault tree analysis
- b. Process mining
- c. Critical decision method

Correct Answer: The correct answer is option a.

Protection of Human and Animal Subjects

The study was approved by Baylor College of Medicine Institutional Review board. Informed consent was obtained from the physicians prior to the interview.

Funding

This project is funded by the Agency for Health Care Research and Quality (R01HS022087) and partially funded by the Houston VA HSR&D Center for Innovations in Quality, Effectiveness and Safety (CIN 13–413). D.R.M. is additionally funded by an Agency for Healthcare Research and Quality Mentored Career Development Award (K08-HS022901) and H.S. is additionally supported by the VA Health Services Research and Development Service (CRE 12–033; Presidential Early Career Award for Scientists and Engineers USA 14–274), the VA National Center for Patient Safety, and the Agency for Health Care Research and Quality (R01HS022087). These funding sources had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; and preparation, review, or approval of the manuscript. There are no conflicts of interest for any authors.

Conflict of Interest

None declared.

References

- 1 Wynia MK, Classen DC. Improving ambulatory patient safety: learning from the last decade, moving ahead in the next. *JAMA* 2011;306(22):2504–2505
- 2 Callen JL, Westbrook JL, Georgiou A, Li J. Failure to follow-up test results for ambulatory patients: a systematic review. *J Gen Intern Med* 2012;27(10):1334–1348
- 3 Callen J, Georgiou A, Li J, Westbrook JL. The safety implications of missed test results for hospitalised patients: a systematic review. *BMJ Qual Saf* 2011;20(02):194–199
- 4 Singh H, Thomas EJ, Sittig DF, et al. Notification of abnormal lab test results in an electronic medical record: do any safety concerns remain? *Am J Med* 2010;123(03):238–244
- 5 Singh H, Thomas EJ, Mani S, et al. Timely follow-up of abnormal diagnostic imaging test results in an outpatient setting: are electronic medical records achieving their potential? *Arch Intern Med* 2009;169(17):1578–1586
- 6 Hysong SJ, Sawhney MK, Wilson L, et al. Understanding the management of electronic test result notifications in the outpatient setting. *BMC Med Inform Decis Mak* 2011;11(01):22
- 7 Singh H, Vij MS. Eight recommendations for policies for communicating abnormal test results. *Jt Comm J Qual Patient Saf* 2010;36(05):226–232
- 8 Hysong SJ, Sawhney MK, Wilson L, et al. Provider management strategies of abnormal test result alerts: a cognitive task analysis. *J Am Med Inform Assoc* 2010;17(01):71–77

- 9 Powell L, Sittig DF, Chrouser K, Singh H. Assessment of health information technology-related outpatient diagnostic delays in the US veterans affairs health care system: a qualitative study of aggregated root cause analysis data. *JAMA Netw Open* 2020;3(06):e206752
- 10 Lacson R, Cochon L, Ip I, et al. Classifying safety events related to diagnostic imaging from a safety reporting system using a human factors framework. *J Am Coll Radiol* 2019;16(03):282–288
- 11 Hoffman RR, Crandall B, Shadbolt N. Use of the critical decision method to elicit expert knowledge: a case study in the methodology of cognitive task analysis. *Hum Factors* 1998;40(02):254–276
- 12 Pascual R, Henderson S. Evidence of naturalistic decision making in military command and control. *Naturalistic Decision Making* 1997:217–26
- 13 Crandall B, Getchell-Reiter K. Critical decision method: a technique for eliciting concrete assessment indicators from the intuition of NICU nurses. *ANS Adv Nurs Sci* 1993;16(01):42–51
- 14 Patterson MD, Militello LG, Bunker A, et al. Leveraging the critical decision method to develop simulation-based training for early recognition of sepsis. *J Cogn Eng Decis Mak* 2016;10(01):36–56
- 15 Power N, Baldwin J, Plummer NR, Laha S. Critical care decision making: a pilot study to explore and compare the decision making processes used by critical care and non-critical care doctors when referring patients for admission. 2017:255. Presented at 13th International Conference on Naturalistic Decision; Bath, UK
- 16 Schnittker R, Marshall SD, Horberry T, Young K. Decision-centred design in healthcare: The process of identifying a decision support tool for airway management. *Appl Ergon* 2019;77:70–82
- 17 Klein GA, Calderwood R, Macgregor D. Critical decision method for eliciting knowledge. *IEEE Trans Syst Man Cybern* 1989;19(03):462–472
- 18 Coble J, Maffitt J, Orland M, Kahn M. Contextual inquiry: discovering physicians' true needs. *Proc Annu Symp Comput Appl Med Care* 1995:469–473
- 19 Ho J, Aridor O, Glinski DW, et al. Needs and workflow assessment prior to implementation of a digital pathology infrastructure for the US Air Force Medical Service. *J Pathol Inform* 2013;4:32
- 20 Liang J, Yan B, Yang C, Yang H. The design of hemiplegia rehabilitation equipment based on contextual inquiry. *IOP Conf. Ser.: Mater. Sci. Eng* 2019;520:012020. Doi: 10.1088/1757-899X/520/1/012020
- 21 Vesely WE, Roberts N. *Fault Tree Handbook*. Washington, D.C.: Systems and Reliability Research, Office of Nuclear Regulatory Research, U.S. Nuclear Regulatory Commission; 1981
- 22 Hessian R, Salter BB, Goodwin EF. Fault-tree analysis for system design, development, modification, and verification. *IEEE Trans Reliab* 1990;39(01):87–91
- 23 Rao KD, Gopika V, Rao VS, Kushwaha H, Verma AK, Srividya A. Dynamic fault tree analysis using Monte Carlo simulation in probabilistic safety assessment. *Reliab Eng Syst Saf* 2009;94(04):872–883
- 24 Hauptmanns U. Semi-quantitative fault tree analysis for process plant safety using frequency and probability ranges. *J Loss Prev Process Ind* 2004;17(05):339–345
- 25 Lee W-S, Grosh D, Tillman FA, Lie CH. Fault tree analysis, methods, and applications: a review. *reliability. IEEE Transactions on*. 1985; 34(03):194–203
- 26 Rogith D, Iyengar MS, Singh H. Using fault trees to advance understanding of diagnostic errors. *Jt Comm J Qual Patient Saf* 2017;43(11):598–605
- 27 Jonas JA, Devon EP, Ronan JC, et al. Determining preventability of pediatric readmissions using fault tree analysis. *J Hosp Med* 2016; 11(05):329–335
- 28 McElroy LM, Khorzad R, Rowe TA, et al. Fault tree analysis: assessing the adequacy of reporting efforts to reduce postoperative bloodstream infection. *Am J Med Qual* 2017;32(01):80–86
- 29 Li X-Z, Gao J-M, Zhao Y-Q, Chen Y-J. Infusion pump system safety analysis based on fault tree and Markov analysis. In: Wei J, ed. *Mechanical Engineering and Control Systems: Proceedings of the 2016 International Conference on Mechanical Engineering and Control System (MECS2016)*. New Jersey, NJ: World Scientific Publishing Co. Pte. Ltd.; 2017
- 30 Sittig DF, Singh H. A new socio-technical model for studying health information technology in complex adaptive healthcare systems. *Qual Saf Health Care* 2010;19(Suppl 3):i68–i74