Entropy of subordinate shift spaces

Marcin Kulczycki, Dominik Kwietniak, and Jian Li August 4, 2017

Abstract

We introduce a new family of shift spaces — the subordinate shifts. Using subordinate shifts we prove in an elementary way that for every nonnegative real number t there is a shift space with entropy t.

1 Introduction.

Let \mathcal{A} be a finite collection of *symbols* which we call an *alphabet*. Typical choices are $\mathcal{A} = \{0, 1, \dots, r-1\}$ for some integer r or a set of Roman letters, for example $\mathcal{A} = \{a, b, c, d\}$. The *full shift* over \mathcal{A} is the set $\mathcal{A}^{\mathbb{N}}$ of all infinite sequences of symbols. A *block* is any finite sequence of symbols.

A shift space over \mathcal{A} is a subset of $\mathcal{A}^{\mathbb{N}}$ defined by establishing some constraints on blocks, which are allowed to appear as subsequences. For example, the set of all infinite sequences of 0's and 1's that do not contain the block 11 is a shift space over $\{0,1\}$. A shift space becomes a dynamical system when we equip it with the *shift map*, which discards the first element of a sequence and shifts the remaining elements by one place to the left. The branch of mathematics concerning the study of shift spaces is known as *symbolic dynamics*.

Shift spaces are mathematical models for digitized information, arising frequently as a result of the discretization of dynamical processes. For example, imagine a point moving along a trajectory in space. Partition the space into finitely many pieces and assign a symbol to each piece. Now write down the sequence of symbols labelling the successive partition elements visited by the point as it follows the trajectory. We have encoded information about the trajectory in a sequence from the shift space over the set of labels of the partition cells.

As is often the case in mathematics, there is a notion of isomorphism for shift spaces known as *conjugacy* showing us when two seemingly different shift spaces are practically the same. In other words, conjugate shift spaces can be treated as two instances of the same underlying object (for more details we refer the reader to [7, Def. 1.5.9]). For example, there is neither loss nor gain of information when the full $\{0,1\}$ -shift is transformed into the full $\{4,7\}$ -shift by switching every 0 to 4 and every 1 to 7.

A common problem in symbolic dynamics is to decide whether two shift spaces are conjugate or not. Various invariants are devised to help us distinguish shift spaces which are *not* conjugate. A property of a shift space is a *conjugacy* invariant if whenever a shift space X possesses that property, then every shift space conjugate to X also possesses that property. Among the most important invariants is entropy, which is a measure of the complexity of a shift space. Entropy is a nonnegative real number equal to the asymptotic growth rate of the number of blocks that occur in a shift space.

Entropy is a conjugacy invariant [7, Cor. 4.1.10]. Therefore, if two shift spaces have different entropy, then they have different structure — they are not conjugate. For example, the full shift over $\{0,1\}$ is not conjugate to the full shift over $\{a,b,c\}$, as they have different entropies.

There is one problem with this useful tool: computing the entropy of an arbitrary shift space is often a difficult or even a hopeless task. For example, it is known that for every nonnegative real number t there is a shift space with entropy t, but this result is a part of nontrivial theory (see for example [10, pp. 178-9]).

We will show, however, that this can be proven in an elementary way. To this end, we will define a new class of shift spaces — the subordinate shifts — for some of which the calculation of entropy is straightforward and requires only basic combinatorics.

1.1 Notation and definitions.

Throughout this paper, the symbol \mathbb{N} denotes the set of *positive* integers. We denote the number of elements of a finite set A by |A|. Given any real number x, by $\lfloor x \rfloor$ we mean the largest integer not greater than x.

Recall that a sequence of real numbers $\{a_n\}_{n=1}^{\infty}$ is *subadditive* if $a_{m+n} \leq a_m + a_n$ for all $m, n \in \mathbb{N}$.

Fekete's Lemma. Let $\{a_n\}_{n=1}^{\infty}$ be a subadditive sequence of nonnegative real numbers. Then the sequence $\{a_n/n\}_{n=1}^{\infty}$ converges to a limit equal to the infimum of the terms of this sequence, that is

$$\lim_{n \to \infty} \frac{a_n}{n} = \inf_{n \in \mathbb{N}} \frac{a_n}{n}.$$

We follow the notation of Lind and Marcus [7] as close as possible. One notable exception is that we consider only one-sided shifts, while Lind and Marcus consider two-sided (invertible) shifts throughout most of their book.

Definitions (Full shifts). Let \mathcal{A} be a finite set, which we call the *alphabet*. We refer to the elements of \mathcal{A} as *symbols*. The *full* \mathcal{A} -shift is the collection of all infinite sequences of symbols from \mathcal{A} . The full \mathcal{A} -shift is denoted by

$$\mathcal{A}^{\mathbb{N}} = \{ x = (x_i)_{i=1}^{\infty} : x_i \in \mathcal{A} \text{ for all } i \in \mathbb{N} \}.$$

We usually write an element of $\mathcal{A}^{\mathbb{N}}$ as $x=(x_i)_{i=1}^{\infty}=x_1x_2x_3...$ Often we identify a finite set \mathcal{A} such that $|\mathcal{A}|=r$ with $\{0,1,\ldots,r-1\}$. A full r-shift is then the full shift over the alphabet $\{0,1,\ldots,r-1\}$ and a full binary shift is the full 2-shift.

Definitions (Blocks). A block over \mathcal{A} is a finite sequence of symbols from \mathcal{A} . We write blocks without separating their symbols, so a block over $\mathcal{A} = \{0, 1, 2\}$ might look like 01220120. The length of a block u is the number of symbols it contains. An n-block stands for a block of length n. We identify a symbol with the 1-block consisting of this symbol. An empty block is the unique block with no symbols and length zero that we denote ε . The set of all blocks over \mathcal{A} (including ε) is denoted by \mathcal{A}^* .

A concatenation of two blocks $u = a_1 \dots a_k$ and $v = b_1 \dots b_l$ is the block uv obtained by writing u first and then v, that is, $uv = a_1 \dots a_k b_1 \dots b_l$. The concatenation is an associative operation, because (uv)w = u(vw) for any blocks $u, v, w \in \mathcal{A}^*$. For this reason we may write uvw, or indeed concatenate any sequence of blocks (finite or not) without ambiguity. If $n \geq 1$, then u^n stands for the concatenation of n copies of u. Given a nonempty block $u \in \mathcal{A}^*$ we denote by u^{∞} the sequence $uuu \dots \in \mathcal{A}^{\mathbb{N}}$.

Let $x=(x_i)_{i=1}^{\infty}\in\mathcal{A}^{\mathbb{N}}$ and let $1\leq i\leq j$ be integers. We write $x_{[i,j]}=x_ix_{i+1}\dots x_j$ for the block of symbols in x starting from the i-th and ending at the j-th position. We say that a block $w\in A^*$ occurs in x and x contains w if $w=x_{[i,j]}$ for some integers $1\leq i\leq j$. Note that ε occurs in every sequence from $\mathcal{A}^{\mathbb{N}}$. Similarly, given an n-block $w=w_1\dots w_n\in\mathcal{A}^*$ we define $w_{[i,j]}=w_iw_{i+1}\dots w_j\in\mathcal{A}^*$ for each $1\leq i\leq j\leq n$.

A prefix of a block $z \in A^*$ is a block u such that z = uv for some $v \in A^*$.

Definition (Shift map). For every \mathcal{A} we define the *shift map* $\sigma: \mathcal{A}^{\mathbb{N}} \to \mathcal{A}^{\mathbb{N}}$. It maps a sequence $x = (x_i)_{i=1}^{\infty}$ to the sequence $\sigma(x) = (x_{i+1})_{i=1}^{\infty}$. Equivalently, $\sigma(x)$ is the sequence obtained by dropping the first symbol of x and moving the remaining symbols by one position to the left.

Definitions (Shift spaces). Given any collection \mathcal{F} of blocks over \mathcal{A} (i.e., a subset of \mathcal{A}^*) we define a *shift space specified by* \mathcal{F} , denoted by $X_{\mathcal{F}}$, as the set of all sequences from $\mathcal{A}^{\mathbb{N}}$ which do not contain any blocks from \mathcal{F} . We say that \mathcal{F} is a collection of *forbidden blocks for* $X_{\mathcal{F}}$ as blocks from \mathcal{F} are forbidden to occur in $X_{\mathcal{F}}$.

A shift space is a set $X \subset \mathcal{A}^{\mathbb{N}}$ such that $X = X_{\mathcal{F}}$ for some $\mathcal{F} \subset \mathcal{A}^*$. A binary shift space is a shift space over the alphabet $\{0,1\}$.

Exercise. Show that for every shift space X we have $\sigma(X) \subset X$. Find a shift space X for which $\sigma(X) \neq X$.

Definition (Language of a shift space). With each shift space X over \mathcal{A} , we may associate a set of blocks over \mathcal{A} which occur in some sequence $x \in X$. We call this set the *language of* X and denote it by $\mathcal{B}(X)$. We write $\mathcal{B}_n(X)$ for the set of all n-blocks contained in $\mathcal{B}(X)$.

Exercise. Show that if \mathcal{L} is a language of some shift space over \mathcal{A} , then \mathcal{L} is:

1. factorial, meaning that if $u \in \mathcal{L}$ and u = vw for some blocks $v, w \in \mathcal{A}^*$, then both v and w also belong to \mathcal{L} ,

2. prolongable, meaning that for every block u in \mathcal{L} there is a symbol $a \in \mathcal{A}$ such that ua also belongs to \mathcal{L} .

Actually, the converse is also true. Given a factorial and prolongable subset $\mathcal{L} \subset \mathcal{A}^*$ there is a shift space X such that \mathcal{L} is the language of X. A collection of forbidden blocks defining X is $\mathcal{F} = \mathcal{A}^* \setminus \mathcal{L}$.

We can also characterize points in a shift space X.

Lemma 1. Let $\mathcal{L} \subset \mathcal{A}^*$ be factorial and prolongable. Let X be a shift space such that $\mathcal{L} = \mathcal{B}(X)$. Then a point $x \in \mathcal{A}^{\mathbb{N}}$ is in X if and only if $x_{[i,j]} \in \mathcal{L}$ for all $i, j \in \mathbb{N}$ with i < j.

Shift spaces are determined by their language. In other words, two shift spaces are equal if and only if they have the same language [7, Proposition 1.3.4]. Hence there is a one-to-one correspondence between shift spaces over \mathcal{A} and factorial, prolongable subsets of \mathcal{A}^* . To define a shift space, one can either specify its set of forbidden blocks or its language.

Example. Let $x \in \mathcal{A}^{\mathbb{N}}$. Let \mathcal{B}_x be the collection of all blocks occurring in x. Since \mathcal{B}_x is a factorial and prolongable language, it defines a shift space, which we denote by Σ_x .

Example. Let $S \subset \mathbb{N} \cup \{0\}$. We define

$$\mathfrak{F}_S = \{1\underbrace{0\ldots 0}_p 1 : p \notin S\}.$$

The binary shift defined by forbidding blocks from \mathcal{F}_S is called the *S*-gap shift and is denoted by X_S . In particular, we call $X_{\mathbb{N}}$ the golden mean shift (a sequence belongs to it if and only if it does not contain the block 11).

The following fact merely states that we may construct inductively an *infinite* sequence starting from an infinite collection of *finite* sequences such that each sequence in it coincides with any shorter one as long as both are defined.

Lemma 2. Let $\{w^{(n)}\}_{n=1}^{\infty}$ be a sequence in A^* and for each $n \in \mathbb{N}$ let l(n) be the length of $w^{(n)}$. If for each $k \in \mathbb{N}$ the block $w^{(k)}$ is a prefix of $w^{(k+1)}$, then there is a point $x \in A^{\mathbb{N}}$ such that for each $n \in \mathbb{N}$ we have $x_{[1,l(n)]} = w^{(n)}$. Moreover, if $\lim_{n \to \infty} l(n) = \infty$, then x is unique.

We use superscripts in brackets as above to denote indices for sequences of blocks. That is, we write $\{w^{(n)}\}_{n=1}^{\infty}$ to denote a sequence of blocks. This way we may reserve subscripts for enumerating symbols within the block: $w^{(n)} = w_1^{(n)} w_2^{(n)} \dots w_k^{(n)}$.

Finally, we are ready to move to the entropy itself. In full generality, this concept was defined by Adler, Konheim and McAndrew [1] for an arbitrary compact topological space X and a continuous map $f: X \to X$. The definition below applies only to shift spaces but the resulting number is equal to the Adler, Konheim and McAndrew entropy of the shift treated as a dynamical system (see [7, Exercise 6.3.8]).

Definition (Entropy). Let log denote the logarithm to base 2 (choosing a different base would also yield a valid definition; it would change the value of entropy only by a multiplicative constant). Let $X \subset \mathcal{A}^{\mathbb{N}}$ be a nonempty shift space and let $m, n \in \mathbb{N}$. Observe that every block $w \in \mathcal{B}_{m+n}(X)$ can be written in a unique way as a concatenation w = uv, where $u \in \mathcal{B}_m(X)$ and $v \in \mathcal{B}_n(X)$. Therefore $|\mathcal{B}_{m+n}(X)| \leq |\mathcal{B}_m(X)| \cdot |\mathcal{B}_n(X)|$, and hence

$$\log |\mathfrak{B}_{m+n}(X)| \le \log |\mathfrak{B}_m(X)| + \log |\mathfrak{B}_n(X)|.$$

By applying Fekete's Lemma to the nonnegative sequence $\log |\mathcal{B}_n(X)|$ we may now define the *entropy of* X, denoted by h(X), as

$$h(X) = \lim_{n \to \infty} \frac{1}{n} \log |\mathcal{B}_n(X)| = \inf_{n \ge 1} \frac{1}{n} \log |\mathcal{B}_n(X)|.$$

Roughly speaking, the entropy measures the complexity of a shift space X in terms of the asymptotic growth rate of the number of n-blocks that appear in the language X. In other words, the number of n-blocks in a shift space of entropy $h \geq 0$ roughly equals 2^{nh} .

Example. Every finite shift space has entropy zero.

Example. The full A-shift has entropy $\log |A|$. In particular, the full r-shift has entropy $\log r$.

Observe that if $X, Y \subset \mathcal{A}^{\mathbb{N}}$ and $X \subset Y$, then $h(X) \leq h(Y)$. Since a shift space over \mathcal{A} is a subset of the full \mathcal{A} -shift, we may conclude that the entropy of any shift space over \mathcal{A} is a nonnegative real number bounded above by $\log |\mathcal{A}|$ (for systems that are not shifts it may well be infinite — see [2, Example 4.2.6]).

As mentioned in the introduction, computing the entropy of a shift space is a hard problem — try, for example to verify straight from the definition that the entropy of the golden mean shift is $\log((1+\sqrt{5})/2)$. There are (relatively rare) families of shift spaces (e.g. shifts of finite type, see [7]) for which we can actually provide a (theoretically) computable formula for entropy. Even in these special cases, one needs to apply some non-trivial tools. For the sake of illustration we recall some results from [7, Exercise 4.3.7].

Theorem 1. Let $S \subset \mathbb{N} \cup \{0\}$ and let X_S be the associated S-gap shift. Then $h(X_S) = \log \lambda$, where λ is the unique positive solution of the equation

$$\sum_{j \in S} x^{-j-1} = 1.$$

Theorem 2. For every $t \in [0,1]$ there is a set $S \subset \mathbb{N} \cup \{0\}$ (S depends on t) such that $h(X_S) = t$.

2 Subordinate shifts.

The goal of this section is to introduce a family of shift spaces with easily calculable entropy; a family rich enough to contain a shift space with every possible nonnegative entropy. For the remainder of the paper, we fix $A = \{0, 1, ..., r-1\}$.

Definition (Subordinate shift). We say that a block $w = w_1 \dots w_k \in \mathcal{A}^*$ dominates a block $v = v_1 \dots v_k \in \mathcal{A}^*$ if $v_i \leq w_i$ for $i = 1, \dots, k$. In an analogous way we define when one sequence from $\mathcal{A}^{\mathbb{N}}$ dominates another.

A subordinate of $\mathcal{L} \subset \mathcal{A}^*$ is the set \mathcal{L}^{\leq} of all blocks over \mathcal{A} that are dominated by some block in \mathcal{L} . Observe that if \mathcal{L} is factorial and prolongable, then the same holds for \mathcal{L}^{\leq} . In particular, given a point $x \in \mathcal{A}^{\mathbb{N}}$, we may define a subordinate shift of x, denoted by $X^{\leq x}$, as a shift space given by the language \mathcal{B}_x^{\leq} , where \mathcal{B}_x is the language of blocks occurring in x.

Example. All binary blocks of length 3 are dominated by 111. The blocks 0000, 0001, 0100, and 0101 are the only blocks dominated by 0101.

Subordinate shifts are *hereditary* (this is a notion introduced in [4] and examined in [6]). It can be shown that a hereditary shift is subordinate if and only if it is irreducible in the sense of [7, Definition 1.3.6]. It turns out that a shift space that has been recently extensively studied is an example of a subordinate shift.

Example. Recall that a positive integer n is square-free if there is no prime number p such that p^2 divides n. Let η be a point in $\{0,1\}^{\mathbb{N}}$ given by

$$\eta_n = \begin{cases} 1 & \text{if } n \text{ is square-free,} \\ 0 & \text{otherwise.} \end{cases}$$

In other words, $\eta_n = (\mu(n))^2$, where $\mu \colon \mathbb{N} \to \mathbb{N}$ is the famous Möbius function. It can be shown that $S = X^{\leq \eta}$ is the *square-free flow*; that is a shift space, whose structure is strongly tied to the statistical properties of square-free numbers. For more details see [8, 9]. The study of the square-free flow has been recently extended to the more general context of \mathcal{B} -free integers; that is to say integers with no factor in a given family \mathcal{B} of pairwise relatively prime integers, the sum of whose reciprocals is finite, see [3, 5].

We aim to show that if given $t \in [0,1]$, then we are able to choose a point x(t) from the full binary shift such that $h(X^{\leq x(t)}) = t$. First we tackle rational entropies.

Lemma 3. If $w \in \{0,1\}^*$ is a block of length q with p occurrences of the symbol 1 and $x = w^{\infty}$, then $h(X^{\leq x}) = p/q$.

Proof. Since replacing in w any subset of 1's with 0's leads to a block dominated by w which is in $\mathcal{B}(X^{\leq x})$, we know that there are at least 2^p blocks in $\mathcal{B}_q(X^{\leq x})$. It follows that $h(X^{\leq x}) \geq p/q$.

On the other hand, the periodicity of $x = w^{\infty}$ implies that for each $j \in \mathbb{N}$ there are at most q different blocks of length qj in \mathcal{B}_x . Each such block dominates exactly 2^{pj} blocks from $\{0,1\}^*$. Therefore there are at most $q \cdot 2^{pj}$ blocks in $\mathcal{B}_{qj}(X^{\leq x})$. Consequently,

$$h(X^{\leq x}) = \lim_{n \to \infty} \frac{1}{n} \log |\mathcal{B}_n(X^{\leq x})| = \lim_{j \to \infty} \frac{1}{qj} \log |\mathcal{B}_{qj}(X^{\leq x})|$$

$$\leq \lim_{j \to \infty} \frac{1}{qj} \log(q \cdot 2^{pj}) = p/q.$$

We now show how to construct a shift space with entropy $\pi/8$. We hope that this example will make the general construction that comes after it much clearer.

Example. Let $t = \pi/8 = 0.3926990816...$ We consider the following sequence of rational approximations of t from the above:

```
0.4, 0.4, 0.393, 0.3927, 0.3927, 0.3927, 0.3926991, 0.39269909, \dots
```

We obtain our n-th approximation by rounding up t to the nearest number with no more than n digits after the decimal point. Let p_n be this n-th approximation times 10^n .

We now inductively build a sequence of blocks $\{w^{(n)}\}_{n=1}^{\infty}$ from $\{0,1\}^*$ such that for every $n \in \mathbb{N}$:

- 1. $w^{(n)}$ has length 10^n ,
- 2. the symbol 1 appears exactly p_n times in $w^{(n)}$,
- 3. $w^{(n)}$ is a prefix of $w^{(n+1)}$
- 4. $w^{(n+1)}$ is dominated by $(w^{(n)})^{10}$.

We can visualize the construction of $w^{(n+1)}$ as a process with two steps. In the first step we concatenate ten copies of $w^{(n)}$. In the second step we keep the first p_{n+1} occurrences of the symbol 1 in $w^{(n+1)}$ and replace the rest by 0's.

In our example, we first put $w^{(1)} = 1111000000$. We want the symbol 1 to appear $p_2 = 40$ times in $w^{(2)}$, so we define $w^{(2)} = (w^{(1)})^{10}$ — there is no need to remove any 1's. Then in $w^{(3)}$ we need to see 393 appearances of 1, so we define $w^{(3)}$ as a concatenation of nine copies of $w^{(2)}$ and a block v of length 100 which agrees with $w^{(2)}$ except that the last seven 1's appearing in $w^{(2)}$ are replaced by 0's in v. The construction continues on inductively.

Applying Lemma 2 to the sequence of binary blocks $\{w^{(n)}\}_{n=1}^{\infty}$, we obtain a point $x \in \{0,1\}^{\mathbb{N}}$. The entropy of the subordinate shift $X^{\leq x}$ is $t = \pi/8$. The proof of this fact is contained in the general result below.

We are now equipped to tackle the main theorem of this paper.

Theorem 3. For every $t \in [0,1]$ there is a binary subordinate shift with entropy t.

Proof. For every $n \in \mathbb{N}$ let s_n be the rational approximation of t obtained by rounding up the decimal expansion of t to the nearest number with no more than n digits after the decimal point. Let $p_n = s_n \cdot 10^n$.

We now inductively build a sequence of blocks $\{w^{(n)}\}_{n=1}^{\infty}$ from $\{0,1\}^*$ such that for every $n \in \mathbb{N}$:

- 1. $w^{(n)}$ has length 10^n ,
- 2. the symbol 1 appears exactly p_n times in $w^{(n)}$,
- 3. $w^{(n)}$ is a prefix of $w^{(n+1)}$,
- 4. $w^{(n+1)}$ is dominated by $(w^{(n)})^{10}$.

We start with

$$w^{(1)} = \underbrace{1 \dots 1}_{p_1} \underbrace{0 \dots 0}_{10 - p_1}.$$

Assume now that we have defined $w^{(1)}, \ldots, w^{(n)}$ so that the four conditions above are satisfied to the extent to which they apply to $w^{(1)}, \ldots, w^{(n)}$. We now concatenate ten copies of $w^{(n)}$. We keep the first p_{n+1} occurrences of the symbol 1 unaltered in the sequence, replace the rest by 0, and call the result $w^{(n+1)}$. It is easy to verify that the conditions above now hold to the extent to which they apply to $w^{(1)}, \ldots, w^{(n+1)}$, so the inductive construction is complete.

Applying Lemma 2 to the sequence $\{w^{(n)}\}_{n=1}^{\infty}$ we obtain a point $x \in \{0,1\}^{\mathbb{N}}$. Observe that for every $n \in \mathbb{N}$ the point x is dominated by $(w^{(n)})^{\infty}$. It follows that $X^{\leq x} \subset X^{\leq (w^{(n)})^{\infty}}$, and so $h(X^{\leq x}) \leq h(X^{\leq (w^{(n)})^{\infty}})$. By Lemma 3 we have $h(X^{\leq (w^{(n)})^{\infty}}) = s_n$, and therefore

$$h(X^{\leq x}) \leq \lim_{n \to \infty} h(X^{\leq (w^{(n)})^{\infty}}) = \lim_{n \to \infty} s_n = t.$$

On the other hand, observe that for every $n \in \mathbb{N}$ we have $w^{(n)} \in \mathcal{B}_{10^n}(X^{\leq x})$, so all 2^{p_n} blocks dominated by $w^{(n)}$ are also in $\mathcal{B}_{10^n}(X^{\leq x})$. Therefore $\log |\mathcal{B}_{10^n}(X^{\leq x})| \geq \log 2^{p_n} = p_n$, and so

$$t = \lim_{n \to \infty} s_n = \lim_{n \to \infty} \frac{p_n}{10^n} \le \lim_{n \to \infty} \frac{1}{10^n} \log |\mathfrak{B}_{10^n}(X^{\le x})| = h(X^{\le x}),$$

which completes the proof of $h(X^{\leq x}) = t$.

Exercise. Prove that the shift space $X^{\leq x}$ constructed in the proof of Theorem 3 is *irreducible* (see [7, Definition 1.3.6]), meaning that given any pair of blocks $u, v \in \mathcal{B}(X^{\leq x})$ there is a block $y \in \mathcal{B}(X^{\leq x})$ such that $uyv \in \mathcal{B}(X^{\leq x})$.

It remains to show that the conclusion of Theorem 3 is true for every t > 1.

Theorem 4. For every $t \in (1, \infty)$ there is a shift space with entropy t.

Proof. Let $t \in (1, \infty)$ be given. Pick $k \in \mathbb{N}$ and $s \in (0, 1)$ such that ks = t. Use Theorem 3 to obtain a binary subordinate shift X with entropy s. Using X we now construct a shift space $Y \subset \{0, 1, \ldots, 2^k - 1\}^{\mathbb{N}}$.

We start with $Y = \emptyset$. For every point $w \in X$ we perform the following procedure:

- 1. express w as a concatenation $w_1w_2...$, where each block w_n has length k,
- 2. for every w_n we determine the number $a_n \in \{0, 1, ..., 2^k 1\}$ such that w_n is the binary notation for a_n (for example, for k = 2, we get $00 \mapsto 0$, $01 \mapsto 1$, $10 \mapsto 2$, and $11 \mapsto 3$),
- 3. we add the sequence $a_1 a_2 \dots$ to Y (note that it is the a_n 's that are symbols here, not their digits).

It is elementary to check that Y is a shift space. It suffices to analyze how the language of Y is created from the language of X.

Observe that for every $n \in \mathbb{N}$ we have $|\mathcal{B}_n(Y)| = |\mathcal{B}_{kn}(X)|$. Therefore

$$h(Y) = \lim_{n \to \infty} \frac{1}{n} \log |\mathcal{B}_n(Y)| = k \cdot \lim_{n \to \infty} \frac{1}{kn} \log |\mathcal{B}_{kn}(X)| = k \cdot h(X) = t. \quad \Box$$

Acknowledgment.

The authors would like to thank the referees for their thorough and careful work. Preparing this article we asked our students and colleagues to comment on it. We are grateful to: Jakub Byszewski, Vaughn Climenhaga, Jakub Konieczny, Marcin Lara, Simon Lunn, Martha Łącka, Dariusz Matlak, Samuel Roth, and Maciej Ulas for their remarks and suggestions. The research of Dominik Kwietniak was supported by the National Science Centre (NCN) under grant Maestro 2013/08/A/ST1/00275. The research of Jian Li was supported by Scientific Research Fund of Shantou University (YR13001).

References

- R. L. Adler, A. G. Konheim, M. H. McAndrew, Topological entropy, Trans. Amer. Math. Soc. 114 (1965) 309–319.
- [2] L. Alsedá, J. Llibre, M. Misiurewicz, Combinatorial Dynamics and Entropy in Dimension One. Second edition. World Scientific, River Edge, NJ, 2000.
- [3] E.-H. El Abdalaoui, M. Lemańczyk, T. De La Rue, A dynamical point of view on the set of B-free integers (2013), available at http://arxiv.org/abs/1311.3752.
- [4] D. Kerr, H. Li, Independence in topological and C^* -dynamics, Math.~Ann. 338 (2007) 869–926.

- [5] J. Kułaga-Przymus, M. Lemańczyk, B. Weiss, On invariant measures for B-free systems (2014), available at http://arxiv.org/abs/1406.3745.
- [6] D. Kwietniak, Topological entropy and distributional chaos in hereditary shifts with applications to spacing shifts and beta shifts, *Discrete Contin. Dyn. Syst.* 33 (2013) 2451–2467.
- [7] D. Lind, B. Marcus, An Introduction to Symbolic Dynamics and Coding. Cambridge University Press, Cambridge, 1995.
- [8] R. Peckner, Uniqueness of the measure of maximal entropy for the square-free flow (2014), available at http://arxiv.org/abs/1205.2905.
- [9] P. Sarnak, Three lectures the Möbius funcon tion. randomness. and dynamics (2011),available at http://publications.ias.edu/sarnak/paper/512.
- [10] P. Walters, An Introduction to Ergodic Theory. Graduate Texts in Mathematics, Vol. 79, Springer-Verlag, New York-Berlin, 1982.
- (M. Kulczycki) Institute of Mathematics, Faculty of Mathematics and Computer Science, Jagiellonian University in Kraków, ul. Łojasiewicza 6, 30-348 Kraków, Poland

E-mail address, M. Kulczycki: Marcin.Kulczycki@im.uj.edu.pl

(D. Kwietniak) Institute of Mathematics, Faculty of Mathematics and Computer Science, Jagiellonian University in Kraków, ul. Łojasiewicza 6, 30-348 Kraków, Poland

E-mail address, D. Kwietniak: dominik.kwietniak@uj.edu.pl

(J. Li) Department of Mathematics, Shantou University, Shantou, Guangdong 515063, P.R. China

E-mail address, J. Li: lijian09@mail.ustc.edu.cn