

MCMC Bayesian Estimation in FIEGARCH Models

Taiane S. Prass^{(a)*}, Sílvia R.C. Lopes^(a) and Jorge A. Achcar^(b)

(a) Mathematics Institute - UFRGS, Porto Alegre, RS, Brazil

(b) Medical School - USP, Ribeirão Preto, SP, Brazil

March 2, 2022

Abstract

Bayesian inference for fractionally integrated exponential generalized autoregressive conditional heteroskedastic (FIEGARCH) models using Markov Chain Monte Carlo (MCMC) methods is described. A simulation study is presented to assess the performance of the procedure, under the presence of long-memory in the volatility. Samples from FIEGARCH processes are obtained upon considering the generalized error distribution (GED) for the innovation process. Different values for the tail-thickness parameter ν are considered covering both scenarios, innovation processes with lighter ($\nu < 2$) and heavier ($\nu > 2$) tails than the Gaussian distribution ($\nu = 2$). A sensitivity analysis is performed by considering different prior density functions and by integrating (or not) the knowledge on the true parameter values to select the hyperparameter values.

Key words: Bayesian inference, MCMC, FIEGARCH processes, Long-range dependence.

1 Introduction

ARCH-type (Autoregressive Conditional Heteroskedasticity) and stochastic volatility (Breidt et al., 1998) models are commonly used in financial time series modeling to represent the dynamic evolution of volatilities. By ARCH-type models we mean not only the ARCH model proposed by Engle (1982) but also several generalizations that were lately proposed.

Among the most popular generalizations of the ARCH model is the generalized ARCH (GARCH) model, introduced by Bollerslev (1986), for which the conditional variance depends not only on the p past values of the process (as in the ARCH model), but also on the q past values of the conditional variance. Although the ARCH and GARCH models are widely used in practice, they do not take into account the asymmetry in the volatility, that is, the fact that volatility tends to rise in response to “bad” news and to fall in response to “good” news. As an alternative, Nelson (1991) introduces the exponential GARCH (EGARCH) model. This model not only describes the asymmetry on the volatility, but also have the advantage that the positivity of the conditional variance is always attained since it is defined in terms of the logarithm function.

*Corresponding author: taianeprass@gmail.com

The fractionally integrated EGARCH (FIEGARCH) and fractionally integrated GARCH (FIGARCH) models proposed, respectively, by [Bollerslev and Mikkelsen \(1996\)](#) and [Baillie et al. \(1996\)](#), generalize the EGARCH ([Nelson, 1991](#)) and the GARCH ([Bollerslev, 1986](#)) models, respectively. FIEGARCH models have not only the capability of modeling clusters of volatility (as ARCH and GARCH models do) and capturing its asymmetry (as the EGARCH model does) but they also take into account the characteristic of long memory in the volatility (as the FIGARCH model does). The non-stationarity of FIGARCH models (in the weak sense) makes this class of models less attractive for practical applications. Another drawback of the FIGARCH models is that we must have $d \geq 0$ and the polynomial coefficients in its definition must satisfy some restrictions so the conditional variance will be positive. FIEGARCH models do not have this problem since the variance is defined in terms of the logarithm function, moreover, they are weak stationary whenever the long memory parameter d is smaller than 0.5 ([Lopes and Prass, 2013](#)).

A complete study on the theoretical properties of FIEGARCH processes is presented in [Lopes and Prass \(2013\)](#). The authors also conduct a simulation study to analyze the finite sample performance of the quasi-maximum likelihood (QML) procedure on parameter estimation. The QML procedure became popular for two main reasons. First, the expression for the quasi-likelihood function is simpler for the Gaussian case than when considering, for example, the Student's t or the generalized error distribution (GED). Second, since the parameters of the distribution function are not estimated, the dimension of the optimization problem is reduced. On the other hand, the results in [Lopes and Prass \(2013\)](#) indicate that, although the QML presents a relatively good performance when the sample size is 2000 and the estimation improves as the sample size increases, it does so very slowly.

In this work we propose the use of Bayesian methods using Monte Carlo simulation techniques on the estimation of the FIEGARCH model parameters. This procedure is usually considered to analyze financial time series assuming stochastic volatility models (see, for example, [Meyer and Yu, 2000](#)), mostly because of the difficulty on applying traditional statistical techniques due to the complexity of the likelihood function. To generate samples from the joint posterior distribution for the parameters of interest we use MCMC (Markov Chain Monte Carlo) methods as the Gibbs Sampling algorithm (see, for example, [Gelfand and Smith, 1990](#); [Casela and George, 1992](#)) or the Metropolis-Hastings algorithm (see, for example, [Smith and Roberts, 1993](#); [Chib and Greenberg, 1995](#)). These samples are generated from all conditional posterior distributions for each parameter given all the other parameters and the data.

A simulation study is conducted to assess the finite sample performance of the procedure proposed here, under the presence of long-memory in the volatility. The samples from FIEGARCH processes are obtained upon considering the GED for the innovation process. Taking into account that financial time series are usually characterized by heavy tailed distributions, different values for the tail-thickness parameter ν are considered covering both scenarios: innovation processes with lighter and heavier tails than the Gaussian distribution. A sensitivity analysis is performed by considering different prior density functions and by integrating (or not) the knowledge on the true parameter values to select the hyperparameter values.

The paper is organized as follows. In [Section 2](#) a review on the definition and main properties of FIEGARCH processes is presented. [Section 3](#) describes the parameter estimation procedure when Bayesian inference using MCMC is considered. [Section 4](#) describes the steps used in the simulation study, such as the data generating process, the prior selection procedure and the performance measures considered. This section also reports the simulation results. [Section 5](#) concludes the paper.

2 FIEGARCH Processes

Let $(1 - \mathcal{B})^{-d}$ be the operator defined by its Maclaurin series expansion, namely,

$$(1 - \mathcal{B})^{-d} = \sum_{k=0}^{\infty} \frac{\Gamma(k+d)}{\Gamma(k+1)\Gamma(d)} := \sum_{k=0}^{\infty} \tau_{d,k} \mathcal{B}^k, \quad (1)$$

where $\tau_{d,k} := \frac{\Gamma(k+d)}{\Gamma(k+1)\Gamma(d)}$, for all $k \geq 1$, $\Gamma(\cdot)$ is the gamma function and \mathcal{B} is the backward shift operator defined by $\mathcal{B}^k(X_t) = X_{t-k}$, for all $k \in \mathbb{N}$.

Assume that $\alpha(\cdot)$ and $\beta(\cdot)$ are polynomials of order p and q , respectively, defined by

$$\alpha(z) = \sum_{i=0}^p (-\alpha_i) z^i \quad \text{and} \quad \beta(z) = \sum_{j=0}^q (-\beta_j) z^j, \quad (2)$$

with $\alpha_0 = \beta_0 = -1$. If $\alpha(\cdot)$ and $\beta(\cdot)$ have no common roots and $\beta(z) \neq 0$ in the closed disk $\{z : |z| \leq 1\}$, then the function $\lambda(\cdot)$, defined by

$$\lambda(z) = \frac{\alpha(z)}{\beta(z)} (1 - z)^{-d} := \sum_{k=0}^{\infty} \lambda_{d,k} z^k, \quad \text{for all } |z| < 1, \quad (3)$$

is analytic in the open disc $\{z : |z| < 1\}$, for any $d > 0$, and in the closed disk $\{z : |z| \leq 1\}$, whenever $d \leq 0$. Therefore, $\lambda(\cdot)$ is well defined and the power series representation in (3) is unique. More specifically, the coefficients $\lambda_{d,k}$, for all $k \in \mathbb{N}$, are given by (see [Lopes and Prass, 2013](#))

$$\lambda_{d,0} = 1 \quad \text{and} \quad \lambda_{d,k} = -\alpha_k^* + \sum_{i=0}^{k-1} \lambda_i \sum_{j=0}^{k-i} \beta_j^* \delta_{d,k-i-j}, \quad \text{for all } k \geq 1, \quad (4)$$

where

$$\alpha_m^* := \begin{cases} \alpha_m, & \text{if } 0 \leq m \leq p; \\ 0, & \text{if } m > p; \end{cases} \quad \beta_m^* := \begin{cases} \beta_m, & \text{if } 0 \leq m \leq q; \\ 0, & \text{if } m > q; \end{cases} \quad (5)$$

and $\delta_{d,j} := \tau_{-d,j}$, for all $j \in \mathbb{N}$, are the coefficients obtained upon replacing $-d$ by d in (1), that is

$$\sum_{k=0}^{\infty} \delta_{d,k} \mathcal{B}^k := \sum_{j=0}^{\infty} \tau_{-d,j} \mathcal{B}^j = (1 - \mathcal{B})^d.$$

Let $\theta, \gamma \in \mathbb{R}$ and $\{Z_t\}_{t \in \mathbb{Z}}$ be a sequence of independent and identically distributed (i.i.d.) random variables, with zero mean and variance equal to one. Assume that θ and γ are not both equal to zero and define $\{g(Z_t)\}_{t \in \mathbb{Z}}$ by

$$g(Z_t) = \theta Z_t + \gamma[|Z_t| - \mathbb{E}(|Z_t|)], \quad \text{for all } t \in \mathbb{Z}. \quad (6)$$

It follows that (see [Lopes and Prass, 2013](#)) $\{g(Z_t)\}_{t \in \mathbb{Z}}$ is a strictly stationary and ergodic process. Moreover, since $\mathbb{E}(Z_0^2) < \infty$, then $\{g(Z_t)\}_{t \in \mathbb{Z}}$ is also weakly stationary with mean zero (therefore a white noise process) and variance σ_g^2 given by

$$\sigma_g^2 = \theta^2 + \gamma^2 - [\gamma \mathbb{E}(|Z_0|)]^2 + 2\theta\gamma \mathbb{E}(Z_0|Z_0|). \quad (7)$$

Now, for any $d < 0.5$ and $\omega \in \mathbb{R}$, let $\{X_t\}_{t \in \mathbb{Z}}$ be the stochastic process defined by

$$X_t = \sigma_t Z_t, \quad (8)$$

$$\begin{aligned} \ln(\sigma_t^2) &= \omega + \frac{\alpha(\mathcal{B})}{\beta(\mathcal{B})} (1 - \mathcal{B})^{-d} g(Z_{t-1}) \\ &= \omega + \sum_{k=0}^{\infty} \lambda_{d,k} g(Z_{t-1-k}), \quad \text{for all } t \in \mathbb{Z}. \end{aligned} \quad (9)$$

Then $\{X_t\}_{t \in \mathbb{Z}}$ is a *Fractionally Integrated EGARCH process*, denoted by FIEGARCH(p, d, q) (Bollerslev and Mikkelsen, 1996).

The properties of FIEGARCH(p, d, q) processes, with $d < 0.5$, are given below (the proofs of these properties can be found in Lopes and Prass, 2013). Henceforth $GED(\nu)$ denotes the generalized error distribution with tail thickness parameter ν .

Proposition 1. *Let $\{X_t\}_{t \in \mathbb{Z}}$ FIEGARCH(p, d, a) process. Then the following properties hold:*

1. $\{\ln(\sigma_t^2)\}_{t \in \mathbb{Z}}$ is a stationary (weakly and strictly) and an ergodic process and the random variable $\ln(\sigma_t^2)$ is almost surely finite, for all $t \in \mathbb{Z}$;
2. if $d \in (-1, 0.5)$ and $\alpha(z) \neq 0$, for $|z| \leq 1$, the process $\{\ln(\sigma_t^2)\}_{t \in \mathbb{Z}}$ is invertible;
3. $\{X_t\}_{t \in \mathbb{Z}}$ and $\{\sigma_t^2\}_{t \in \mathbb{Z}}$ are strictly stationary and ergodic processes;
4. if $\{Z_t\}_{t \in \mathbb{Z}}$ is a sequence of i.i.d. $GED(\nu)$ random variables, with $\nu > 1$, zero mean and variance equal to one, then $\mathbb{E}(X_t^r) < \infty$ and $\mathbb{E}(\sigma_t^{2r}) < \infty$, for all $t \in \mathbb{Z}$ and $r > 0$.

3 Parameter Estimation: Bayesian Inference using MCMC

Let ν be the parameter (or vector of parameters) associated to the probability density function of Z_0 and denote by

- $\boldsymbol{\eta} = (\nu, d, \theta, \gamma, \omega, \alpha_1, \dots, \alpha_p, \beta_1, \dots, \beta_q)' := (\eta_1, \eta_2, \dots, \eta_{5+p+q})'$ the vector of unknown parameters in (9);
- $\boldsymbol{\eta}_{(-i)}$ the vector containing all parameters in $\boldsymbol{\eta}$ except η_i , for each $i \in \{1, \dots, 5 + p + q\}$;
- $p_Z(\cdot | \nu)$ the probability density function of Z_0 given ν ;
- \mathcal{F}_t the σ -algebra generated by $\{Z_s\}_{s \leq t}$;
- $p_{X_t}(\cdot | \boldsymbol{\eta}, \mathcal{F}_{t-1})$ the probability density function of X_t given $\boldsymbol{\eta}$ and \mathcal{F}_{t-1} , for all $t \in \mathbb{Z}$.

From (9) it is evident that, given $\boldsymbol{\eta}$, σ_t is a \mathcal{F}_{t-1} -measurable random variable. Moreover, since $X_t = \sigma_t Z_t$ and $p_Z(\cdot | \nu, \mathcal{F}_{t-1}) = p_Z(\cdot | \nu)$, the following equality holds

$$p_{X_t}(x_t | \boldsymbol{\eta}, \mathcal{F}_{t-1}) = \frac{1}{\sigma_t} p_Z(x_t \sigma_t^{-1} | \nu), \quad \text{with} \quad \sigma_t = \exp \left\{ \frac{1}{2} \left[\omega + \sum_{k=0}^{\infty} \lambda_{d,k} g(z_{t-1-k}) \right] \right\}, \quad (10)$$

Metropolis-Hastings (Metropolis et al., 1953; Hastings, 1970) algorithms. This method is usually referred to as Gibbs sampler with Metropolis steps. In this case, to draw the random variate η_i , one shall follow the same steps 1-3 just described. However, instead of sampling directly from $p(\eta_i|\mathbf{X}, \boldsymbol{\eta}_{(-i)}, \mathcal{F}_0)$, one shall consider the Metropolis-Hastings algorithm with $p(\eta_i|\mathbf{X}, \boldsymbol{\eta}_{(-i)}, \mathcal{F}_0)$ as the invariant (target) distribution.

Metropolis-Hastings algorithm is easy to implement since it does not require knowing the normalization constant $p_{(-i)}(\mathbf{X}, \boldsymbol{\eta}_{(-i)}, I_0)$, defined in (12). For simplicity of notation, in what follows $p_*(\cdot)$ shall denote any one of the non-normalized probability density function which corresponds to $p(\eta_i|\mathbf{X}, \boldsymbol{\eta}_{(-i)}, \mathcal{F}_0)$, for $i \in \{1, \dots, 5 + p + q\}$. The Metropolis-Hastings sampling procedure consists of the following steps.

Step 1. Select a transition kernel²(also called proposal distribution) $q(\cdot|\cdot)$ for which the sampling procedure is known.

Step 2. Set an arbitrary initial value y_0 for the chain. Let $m = 0$.

Step 3. Generate a draw ξ from $q(\cdot|y_m)$.

Step 4. Calculate $\alpha(y_m, \xi) = \min \left\{ 1, \frac{p_*(\xi)q(y_m|\xi)}{p_*(y_m)q(\xi|y_m)} \right\}$.

Step 5. Draw $u \sim \mathcal{U}[0, 1]$.

Step 6. Define $y_{m+1} = \begin{cases} \xi, & \text{if } u < \alpha(y_m, \xi); \\ y_m, & \text{otherwise.} \end{cases}$

Step 7. If $m + 1 < N$ (where N is the desired sample size), let $m = m + 1$ and go to Step 3.

Remark 1.

1. When considering Gibbs sampler with Metropolis steps only one iteration of Metropolis-Hastings algorithm is performed for each Gibbs sampler iteration.
2. In both cases, Gibbs sampler and Metropolis-Hastings algorithm, it is advised to discard the first B (for some $B < N$) observations (that is, the burn-in sample) to assure the chain convergence.
3. The sample obtained from the algorithm described above is not independent. An alternative is to run parallel chains instead. Another common strategy to reduce sample autocorrelations is thinning the Markov chain, that is, to keep only every k -th simulated draw from each sequence. There is some controversy surrounding the question of whether or not it is better to run one long chain or several shorter ones (Gelman and Rubin, 1992; Geyer, 1992). Also, MacEachern and Berliner (1994) show that one always get more precise posterior estimates if the entire Markov chain is used instead of the thinned one.

4 Simulation Study

This simulation study considers FIEGARCH(0, d , 0) processes. Under this scenario, the vector of unknown parameters is $\boldsymbol{\eta} = (\nu, d, \theta, \gamma, \omega)' := (\eta_1, \dots, \eta_5)'$. The Bayesian inference approach, using MCMC to obtain posterior density functions, is used to estimate the parameters of the model.

²A transition kernel is a function $q(x|y)$ which is a probability measure with respect to x , so $\int q(x|y)dx = 1$.

4.1 Data Generating Process

The samples from FIEGARCH(0, d , 0) processes are obtained by setting the following.

- $Z_0 \sim \text{GED}(\nu)$, with zero mean and variance equal to one. Thus,

$$p_z(z|\nu) = \frac{\nu \exp\left\{-\frac{1}{2}|z\lambda_\nu^{-1}|^\nu\right\}}{\lambda_\nu 2^{1+1/\nu}\Gamma(1/\nu)}, \quad \lambda_\nu = \left[2^{-2/\nu} \frac{\Gamma(1/\nu)}{\Gamma(3/\nu)}\right]^{1/2} \quad \text{for all } z \in \mathbb{R};$$

- $d \in \{0.10, 0.25, 0.35, 0.45\}$ and $\nu \in \{1.1, 1.5, 1.9, 2.5, 5\}$;
- for all models, $\omega = -5.40$, $\theta = -0.15$ and $\gamma = 0.24$. These values are close to the ones already observed in practical applications (see, for instance, Nelson, 1991; Bollerslev and Mikkelsen, 1996; Ruiz and Veiga, 2008; Lopes and Prass, 2013).
- the infinite sum in (9) is truncated at $m^* = 50,000$.

For each combination of d and ν , a sample $\{z_t\}_{t=-m^*}^n$, of size $m^* + n + 1$, is drawn from the $\text{GED}(\nu)$ distribution and then the sample $\{x_t\}_{t=1}^n$, from the FIEGARCH(0, d , 0) process, is obtained through the relation

$$\ln(\sigma_t^2) = \omega + \sum_{k=0}^{m^*} \lambda_{d,k} g(z_{t-1-k}) \quad \text{and} \quad x_t = \sigma_t z_t, \quad \text{for all } t = 1, \dots, n.$$

4.2 Parameter Estimation Settings

The samples from the posterior distributions are obtained by considering the Gibbs sampler algorithm with Metropolis steps as described in Section 3. The transition kernel $q(\cdot|\cdot)$ considered in the Metropolis-Hastings algorithm is the function defined as

$$q(x|y) = f(x; y, \sigma, a, b),$$

where $f(\cdot; \cdot, \cdot, \cdot, \cdot)$ is the truncated normal density function, defined as

$$f(x; \mu, \sigma, a, b) = \begin{cases} \frac{1}{\sigma} \frac{\phi\left(\frac{x-\mu}{\sigma}\right)}{\Phi\left(\frac{b-\mu}{\sigma}\right) - \Phi\left(\frac{a-\mu}{\sigma}\right)}, & \text{if } a \leq x \leq b, \\ 0, & \text{otherwise,} \end{cases}$$

where $\phi(\cdot)$ and $\Phi(\cdot)$ are, respectively, the probability density and cumulative distribution functions of the standard normal distribution; $a, b \in \mathbb{R}$ are, respectively, the lower and upper limits of the distribution's support; μ and σ denote, respectively, the distribution's (non-truncated version).

To select a reasonable $\boldsymbol{\eta}^{(0)}$, $p_{\mathbf{X}}(\mathbf{X}|\boldsymbol{\eta}, \mathcal{F}_0)$ is calculated for different combinations of ν, d, θ, γ and ω . Then $\boldsymbol{\eta}^{(0)}$ is defined as the vector $\boldsymbol{\eta} = (\nu, d, \theta, \gamma, \omega)'$ with higher likelihood function value. To eliminate any dependence on the initial $\boldsymbol{\eta}^{(0)}$ a burn-in of size 1000 is considered.

A sample obtained by the method being described will probably present significant correlation³. However, due to the ergodicity property of the Markov chain, the estimation of the

³In fact, for the parameter d , this correlation could only be removed when the thinning parameter t was set to 200.

mean is not affected by the correlation in the sample. Therefore, to avoid unnecessary computational work, which ultimately would not lead to improvement in terms of parameter estimation, thinning is not implemented. Nevertheless, an example showing the influence of using the entire chain, the thinned chain or only the first 1000 observations of the entire chain (after burn-in) is provided in the following.

Example 1. Let $\boldsymbol{\eta} = (\nu, d, \theta, \gamma, \omega)'$ $:= (\eta_1, \dots, \eta_5)'$ and assume that

$$\pi_i(\eta_i) = \begin{cases} c_i, & \text{if } \eta_i \in I_i; \\ 0, & \text{otherwise;} \end{cases} \quad \text{for each } i \in \{1, \dots, 5\}, \quad (13)$$

with $c_1 = 1$, $c_2 = c_3 = c_4 = 2$, $c_5 = 1/30$, $I_1 = (0, \infty)$, $I_2 = [0, 0.5]$, $I_3[-0.5, 0]$, $I_4 = I_2$ and $I_5 = [-15, 15]$.

In the sequel, $\{\eta_i^{(k)}\}_{k=1}^n$ denotes the chain of size n obtained from the posterior distribution of η_i , upon considering the prior $\pi_i(\eta_i)$ defined in (13), for each $i \in \{1, \dots, 5\}$. Also, \mathfrak{b} , \mathfrak{t} and N denote, respectively, the burn-in size, the thinning parameter and the sample size of the thinned chain⁴ obtained from $\{\eta_i^{(k)}\}_{k=1}^n$, for any $i \in \{1, \dots, 5\}$.

Figure 1 presents the graph of $\{\eta_i^{(k)}\}_{k=1}^n$, for each $i \in \{1, \dots, 5\}$, with $n = 200,801$. Figure 1 also shows the thinned chain of size $N = 1000$ obtained by considering $\mathfrak{b} = 1000$ and $\mathfrak{t} = 200$. Furthermore, Figure 1 gives the sample of size 1000, obtained from $\{\eta_i^{(k)}\}_{k=1}^n$ by considering a burn-in equal to 1000 and no thinning, for each $i \in \{1, \dots, 5\}$. The true parameter values of the FIEGARCH(0, d , 0) model corresponding to these graphs are $\nu_0 = 1.9$, $d_0 = 0.25$, $\theta_0 = -0.15$, $\gamma_0 = 0.24$ and $\omega_0 = -5.4$. Figure 2 gives the histogram and kernel density functions corresponding to each sample in Figure 1. The graphs of the prior $\pi_i(\eta_i)$ defined in (13), for $i \in \{1, \dots, 5\}$, are represented in Figure 2 by the dashed lines. For a better visualization of the posterior distributions, in Figure 2, the range for the x -axis was restricted to the intervals $[-1.5, 2.5]$, $[-0.5, 0]$, $[0, 0.5]$ and $[-5.6, -5.1]$, respectively, for the parameters ν , θ , γ and ω .




Figure 1: Original chain with sample size 200801 (top row). Thinned chain with sample size 1000 and thinning parameter equal to 200 (middle row). Unthinned chain with sample size 1000 (bottom row). For the middle and bottom rows the burn-in size is equal to 1000. The true parameter values of the FIEGARCH(0, d , 0) model corresponding to these graphs are $\nu_0 = 1.9$, $d_0 = 0.25$, $\theta_0 = -0.15$, $\gamma_0 = 0.24$ and $\omega_0 = -5.4$.

⁴Observe that, by setting $\mathfrak{b} = 1000$ and $\mathfrak{t} = 200$, then a thinned chain of size $N = 1000$ can only be obtained from $\{\eta_i^{(k)}\}_{k=1}^n$ when $n \geq \mathfrak{b} + 1 + \mathfrak{t}(N - 1) = 200,801$.




Figure 2: Histogram and kernel density functions for the original chain with sample size 200801 (top row); the thinned chain with sample size 1000 and thinning parameter equal to 200 (middle row) and the unthinned chain with sample size 1000 (bottom row). For the middle and bottom rows the burn-in size is equal to 1000. The dashed lines correspond to the graphs of the priors $\pi_i(\eta_i)$ defined in (13), for $i \in \{1, \dots, 5\}$. The range for the x -axis was restricted to the intervals $[-1.5, 2.5]$, $[-0.5, 0]$, $[0, 0.5]$ and $[-5.6, -5.1]$, respectively, for the parameters ν, θ, γ and ω . The true parameter values of the FIEGARCH(0, d , 0) model corresponding to these graphs are $\nu_0 = 1.9$, $d_0 = 0.25$, $\theta_0 = -0.15$, $\gamma_0 = 0.24$ and $\omega_0 = -5.4$.

As shown in Figure 1 (see also Table 1), the mean of the posterior distribution does not change significantly when the entire sample or the thinned chain is considered instead of the unthinned one. On the other hand, Figure 2 reinforces the idea that the entire chain gives better estimates for the density function (notice that the curves in the graphs are smoother). Although the thinned chain is not as efficient as the entire chain, it still provides better estimates for the density function than the unthinned one.

Table 1 presents the summary statistics for the samples obtained from the posterior distribution for each parameter of the FIEGARCH(0, d , 0) model. This table considers the entire, thinned and unthinned chains. The statistics reported in this table are the sample mean ($\bar{\eta}_i$), the sample standard deviation (sd_{η_i}) and the 95% credibility interval $CI_{0.95}(\eta_i)$ for the parameter η_i in $\boldsymbol{\eta} = (\nu, d, \theta, \gamma, \omega)' := (\eta_1, \dots, \eta_5)'$, for each $i \in \{1, \dots, 5\}$. The true parameter values considered for this illustration are $\nu_0 \in \{1.1, 1.5, 2.5, 5.0\}$, $d_0 = 0.25$, $\theta_0 = -0.15$, $\gamma_0 = 0.24$ and $\omega_0 = -5.4$.

From Table 1 it is clear that, for any η_i , with $i \in \{1, \dots, 5\}$, the use of the entire or the thinned (thinning parameter $t = 200$) does not yield significant improvement in terms of parameter estimation. Not even the differences in the sample standard deviations or in the credibility intervals, which are the statistics affected by the sample correlation, justify the computational effort to obtain a sample of size 200,801. The same conclusions are obtained when considering $d_0 \in \{0.10, 0.35, 0.45\}$. This concludes the example.

Different prior distributions are tested as explained in the sequel. Since the conditional probability density function of I_0 given $\boldsymbol{\eta}$ is difficult to obtain, in all scenarios, it is assumed that $g(Z_s) = 0$, for all $s \leq 1$, and it is fixed $p_{I_0}(\cdot | \boldsymbol{\eta}) = 1$. Moreover, since (9) is well defined regardless the relation among the parameters of the model, it is assumed that

$$p_{(-i)}(\boldsymbol{\eta}_{(-i)} | \eta_i) \propto \prod_{j \neq i} \pi_j(\eta_j), \quad \text{for any } i \in \{1, \dots, 5\}.$$

Table 1: Summary for the entire, thinned (thinning parameter $t = 200$) and unthinned sample from posterior distributions considering all prior uniforms: mean $\bar{\eta}_i$, standard deviation sd_{η_i} and the 95% credibility interval $CI_{0.95}(\eta_i)$ for the parameter η_i in $\boldsymbol{\eta} = (\nu, d, \theta, \gamma, \omega)'$:= $(\eta_1, \dots, \eta_5)'$, for each $i \in \{1, \dots, 5\}$. The true parameter values considered in this simulation are $\nu_0 \in \{1.1, 1.5, 2.5, 5.0\}$, $d_0 = 0.25$, $\theta_0 = -0.15$, $\gamma_0 = -0.24$ and $\omega_0 = -5.4$. For the thinned and unthinned samples, the burn-in size is $b = 1000$.

Chain	ν_0	$\bar{\nu}$ (sd $_{\nu}$) $CI_{0.95}(\nu)$	\bar{d} (sd $_d$) $CI_{0.95}(d)$	$\bar{\theta}$ (sd $_{\theta}$) $CI_{0.95}(\theta)$	$\bar{\gamma}$ (sd $_{\gamma}$) $CI_{0.95}(\gamma)$	$\bar{\omega}$ (sd $_{\omega}$) $CI_{0.95}(\omega)$
Entire	1.1	1.095 (0.047) [1.006; 1.188]	0.263 (0.112) [0.035; 0.464]	-0.087 (0.038) [-0.164; -0.016]	0.233 (0.062) [0.114; 0.358]	-5.469 (0.078) [-5.612; -5.299]
	1.5	1.478 (0.067) [1.351; 1.612]	0.220 (0.081) [0.056; 0.375]	-0.184 (0.037) [-0.257; -0.111]	0.240 (0.057) [0.130; 0.355]	-5.408 (0.058) [-5.520; -5.291]
	1.9	2.077 (0.107) [1.874; 2.297]	0.252 (0.075) [0.092; 0.392]	-0.209 (0.032) [-0.272; -0.148]	0.220 (0.051) [0.122; 0.322]	-5.386 (0.058) [-5.487; -5.261]
	2.5	2.727 (0.153) [2.441; 3.040]	0.298 (0.056) [0.184; 0.405]	-0.203 (0.025) [-0.253; -0.153]	0.205 (0.045) [0.118; 0.296]	-5.361 (0.053) [-5.463; -5.252]
	5.0	5.227 (0.366) [4.548; 5.978]	0.220 (0.051) [0.115; 0.317]	-0.173 (0.019) [-0.211; -0.135]	0.294 (0.036) [0.224; 0.366]	-5.303 (0.039) [-5.384; -5.230]
Thinned	1.1	1.095 (0.047) [1.006; 1.193]	0.264 (0.110) [0.038; 0.462]	-0.087 (0.038) [-0.161; -0.015]	0.233 (0.062) [0.116; 0.362]	-5.469 (0.080) [-5.609; -5.288]
	1.5	1.480 (0.067) [1.353; 1.613]	0.221 (0.077) [0.070; 0.367]	-0.184 (0.037) [-0.257; -0.104]	0.240 (0.057) [0.129; 0.352]	-5.405 (0.060) [-5.524; -5.291]
	1.9	2.079 (0.103) [1.888; 2.292]	0.253 (0.075) [0.098; 0.387]	-0.210 (0.031) [-0.269; -0.152]	0.218 (0.052) [0.120; 0.327]	-5.388 (0.056) [-5.486; -5.261]
	2.5	2.718 (0.154) [2.427; 3.064]	0.298 (0.058) [0.181; 0.410]	-0.202 (0.026) [-0.255; -0.153]	0.205 (0.044) [0.121; 0.291]	-5.361 (0.053) [-5.466; -5.254]
	5.0	5.224 (0.363) [4.534; 5.991]	0.218 (0.052) [0.112; 0.316]	-0.173 (0.019) [-0.211; -0.137]	0.295 (0.037) [0.224; 0.370]	-5.302 (0.039) [-5.391; -5.230]
Unthinned	1.1	1.108 (0.039) [1.028; 1.198]	0.265 (0.129) [0.016; 0.467]	-0.089 (0.038) [-0.176; -0.019]	0.230 (0.060) [0.108; 0.345]	-5.476 (0.052) [-5.553; -5.366]
	1.5	1.474 (0.067) [1.352; 1.644]	0.250 (0.071) [0.123; 0.393]	-0.175 (0.035) [-0.248; -0.100]	0.240 (0.053) [0.135; 0.353]	-5.426 (0.057) [-5.535; -5.344]
	1.9	2.072 (0.109) [1.885; 2.303]	0.245 (0.074) [0.068; 0.400]	-0.210 (0.032) [-0.273; -0.152]	0.223 (0.059) [0.116; 0.349]	-5.400 (0.062) [-5.537; -5.244]
	2.5	2.720 (0.148) [2.412; 3.016]	0.308 (0.060) [0.192; 0.426]	-0.200 (0.026) [-0.257; -0.155]	0.198 (0.042) [0.119; 0.281]	-5.367 (0.058) [-5.462; -5.274]
	5.0	5.311 (0.356) [4.676; 6.070]	0.226 (0.045) [0.137; 0.316]	-0.176 (0.019) [-0.215; -0.140]	0.293 (0.036) [0.225; 0.368]	-5.291 (0.035) [-5.346; -5.246]

In a first moment the prior distributions for ν, d, θ, γ and ω are selected by considering only the basic set of information usually available in practice. The information on each parameter and the corresponding prior selected are given in Table 2. This scenario shall be referred to as Case 1. Table 3 presents the mean, standard deviation, lower and upper limits for the transition kernel considered at iteration m of the Gibbs sampler with Metropolis steps, when the prior for η_i in $\boldsymbol{\eta} = (\nu, d, \theta, \gamma, \omega)'$:= $(\eta_1, \dots, \eta_5)'$, for each $i \in \{1, \dots, 5\}$, is defined according to Case 1.

In a second moment the knowledge on the true parameter values is gradually incorporated to provide more informative priors for d, θ and/or γ . This analysis, combined with the first scenario, provides information on the sensitivity of the estimates with respect to the priors functions and hyperparameters. In all cases, the priors for ν and ω are the same and are the

Table 2: Information available in practice for the parameter η_i in $\boldsymbol{\eta} = (\nu, d, \theta, \gamma, \omega)'$:= $(\eta_1, \dots, \eta_5)'$ and the corresponding prior considered, for each $i \in \{1, \dots, 5\}$.

Information Available	Prior
The generalized error distribution is well defined for any $\nu > 0$.	$\nu \sim \mathbb{I}_{(0, \infty)}(\nu)$ *
Long-memory in volatility is observed if and only if $d \in (0, 0.5)$. This characteristic can be detected, for example, through the periodogram function of the time series $\{\ln(X_t^2)\}_{t=1}^n$ (see Lopes and Prass, 2013).	$d \sim \mathcal{U}(0, 0.5)$
Empirical evidence suggests that $\theta \in [-1, 0]$. **	$\theta \sim \mathcal{U}(-1, 0)$
Empirical evidence suggests that $\gamma \in [0, 1]$. **	$\gamma \sim \mathcal{U}(0, 1)$
$\omega = \mathbb{E}(\ln(h_t^2)) = \mathbb{E}(\ln(X_t^2)) + \mathbb{E}(\ln(Z_t^2))$. The choice of the interval for ω will depend on the magnitude of the data. The sample mean of $\{\ln(X_t^2)\}_{t=1}^n$ or $\ln(\hat{\sigma}_X^2)$, where $\hat{\sigma}_X^2$ is the sample variance of $\{X_t\}_{t=1}^n$, can be used to obtain a rough approximation for ω	$\omega \sim \mathcal{U}(-15, 15)$.

Notes: * Given $A \subset \mathbb{R}$, the symbol $\mathbb{I}_A(x)$ denotes the improper prior defined as 1, if $x \in A$, and 0, if $x \notin A$.
** See, for instance, Nelson (1991); Bollerslev and Mikkelsen (1996); Ruiz and Veiga (2008); Lopes and Prass (2013). To the best of our knowledge, a FIEGARCH model for which θ or γ are not in the intervals, respectively, $[-1, 0]$ and $[0, 1]$ has never been reported in the literature.

Table 3: Parameters of the truncated normal distribution (transition kernel) considered, at iteration m of the Gibbs sampler, to obtain the sample from the posterior distribution of the parameter η_i in $\boldsymbol{\eta} = (\nu, d, \theta, \gamma, \omega)'$:= $(\eta_1, \dots, \eta_5)'$, for each $i \in \{1, \dots, 5\}$.

Parameter	ν	d	θ	γ	ω
Mean (y)	$\nu^{(m-1)}$	$d^{(m-1)}$	$\theta^{(m-1)}$	$\gamma^{(m-1)}$	$\omega^{(m-1)}$
Standard Deviation (σ)	0.500	0.025	0.050	0.050	1.500
Lower Limit (a)	0.000	0.000	-1.000	0.000	-15.000
Upper Limit (b)	10.000	0.500	0.000	1.000	15.000

Note: $\eta_i^{(m-1)}$, for any $i \in \{1, \dots, 5\}$, denotes the parameter value obtained in the $(m - 1)$ th iteration. Different combinations of standard deviation, lower and upper limits were tested for the parameter η_i in $\boldsymbol{\eta} = (\nu, d, \theta, \gamma, \omega)'$, for each $i \in \{1, \dots, 5\}$. The values presented here correspond to the final choice.

ones defined in Table 2. The scenarios considered in this second step are described in the following and shall be referred to as Case 2 - Case 5.

Case 2: Gaussian Prior for $x = \phi^{-1}(d)$ and Uniform Priors for θ and γ .

In this case θ and γ remain with the same priors as in Case 1. For the parameter d it is assumed that $x \sim \mathcal{N}(\mu_\phi, \sigma_\phi^2)$ and $d = \phi(x)$, where $\phi : \mathbb{R} \rightarrow (0, 0.5)$ is given by

$$\phi(x) = \frac{e^x}{2(1 + e^x)}, \quad \text{for all } x \in \mathbb{R}. \tag{14}$$

First, the knowledge of d_0 is applied to set $\mu_\phi = \phi^{-1}(d_0)$, so $\mu_\phi \in \{-1.386, 0.000, 0.847, 2.197\}$,

respectively, for $d_0 \in \{0.10, 0.25, 0.35, 0.45\}$. This scenario shall be referred to as C2.1. Second, the knowledge on d_0 is ignored and the parameter μ_ϕ is assumed to be equal to zero. This scenario shall be referred to as C2.2. For both, C2.1 and C2.2, different values of σ_ϕ are tested. Third, the approaches considered in C2.1 and C2.2 are combined by setting $\mu_\phi = \phi^{-1}(\bar{d})$, where \bar{d} is the estimate of d obtained in C2.2. This scenario shall be referred to as C2.3. The value of σ_ϕ considered in C2.3 is the one which provides better estimates for d in C2.1.

The kernel parameter values for ν, θ, γ and ω are the same as in the Case 1. For $x = \phi^{-1}(d)$, at iteration m of the Gibbs sampler, the kernels mean (y), standard deviation (sd), lower (a) and upper limits (b) are set, respectively, as $x^{(m-1)}$, 1, -10 and 10, where $x^{(m-1)}$ is the parameter value obtained at iteration $m - 1$.

Case 3: Gaussian Prior for $x = \phi^{-1}(d)$, Beta Prior for $-\theta$ and Uniform Prior for γ .

In this case, the priors of γ and d are the same ones considered, respectively, in Case 1 and in scenario C2.1 of Case 2. It is also assumed that $-\theta \sim \text{Beta}(a_1, b_1)$, which is equivalent to set

$$\pi_3(\theta) = (-\theta)^{a_1-1}(1+\theta)^{b_1-1}B(a_1, b_1), \quad \theta \in [-1, 0],$$

where $B(\cdot, \cdot)$ is the beta function.

First, the fact that $X \sim \text{Beta}(a, b)$ implies $\mathbb{E}(X) = a(a+b)^{-1}$, is applied to set $b_1 = a_1(1+\theta_0)(-\theta_0)^{-1}$, where $\theta_0 = -0.15$ is the true parameter value considered in this simulation study. Different values of a_1 are tested. This scenario shall be referred to as C3.1. Second, the knowledge on θ_0 is ignored and different combinations of a_1 and b_1 are tested. This scenario shall be referred to as C3.2. Third, the approaches considered in C3.1 and C3.2 are combined by setting $b_1 = a_1(1+\bar{\theta}_0)(-\bar{\theta}_0)^{-1}$, where $-\bar{\theta}_0$ is the estimate of θ obtained in C3.2. The value of a_1 considered in this case is the one which provides better estimates for θ in C3.1. This scenario shall be referred to as C3.3.

The kernel parameter values are the same as in Case 2.

Case 4: Gaussian Prior for $x = \phi^{-1}(d)$, Beta Priors for $-\theta$ and γ .

In this case, the priors of d and $-\theta$ are the same ones considered, respectively, in scenario C2.1 of Case 2 and in scenario C3.1 of Case 3. It is also assumed that $\gamma \sim \text{Beta}(a_2, b_2)$. Two scenarios, denoted by C4.1 and C4.2 are considered. With the obvious identifications, the construction of C4.1 and C4.2 is analogous, respectively, to the construction of scenarios C3.1 and C3.2 in Case 3.

The kernel parameter values are the same as in Case 2.

Case 5: Beta Priors for $2d$, $-\theta$ and γ .

In this case, the priors of $-\theta$ and γ are the same ones considered, respectively, in scenario C3.1 of Case 3 and in scenario C4.1 of Case 4. Moreover, for each $d_0 \in \{0.10, 0.25, 0.35, 0.45\}$ considered in this simulation study, it is assumed that $2d \sim \text{Beta}(a_3, b_3)$, which is equivalent to set

$$\pi_2(d) = 2(2d)^{a_3-1}(1-2d)^{b_3-1}B(a_3, b_3), \quad d \in [0, 0.5],$$

where $B(\cdot, \cdot)$ is the beta function.

In this case, only two scenarios are considered. First, it is assumed that $b_3 = a_3(1 - 2d_0)(2d_0)^{-1}$ and different values of a_3 are tested. This scenario shall be referred to as C5.1. Second, an approach similar to scenarios C3.3 and C4.3, respectively, in Case 3 and Case 4, is considered. However, in this case, it is assumed that $b_3 = a_3(1 - 2\bar{d})(2\bar{d})^{-1}$, with \bar{d} obtained in Case 1. The value of a_3 considered in this case is the one which provides better estimates for d in C5.1. This scenario shall be referred to as C5.2.

The kernel parameter values are the same as in Case 1.

4.3 Estimates and Performance Measures

Let $\{\eta_i^{(k)}\}_{k=1}^M$ be a sample of size M from the posteriori distribution of η_i in $\boldsymbol{\eta} = (\nu, d, \theta, \gamma, \omega)' := (\eta_1, \dots, \eta_5)'$, for any $i \in \{1, \dots, 5\}$. Denote by $\bar{\eta}_i$ and sd_{η_i} , respectively, the sample mean and standard deviation of $\{\eta_i^{(k)}\}_{k=1}^M$, namely,

$$\bar{\eta}_i = \frac{1}{M} \sum_{k=1}^M \eta_i^{(k)} \quad \text{and} \quad \text{sd}_{\eta_i} = \sqrt{\frac{1}{M} \sum_{k=1}^M (\eta_i^{(k)} - \bar{\eta}_i)^2}, \quad \text{for any } i \in \{1, \dots, 5\}.$$

Then the estimate $\hat{\eta}_i$ of η_i is defined as $\hat{\eta}_i := \bar{\eta}_i$.

Moreover, let $\hat{q}_i(\alpha)$ denote the α quantile⁵ for the posterior sample distribution of η_i , for any $\alpha \in [0, 1]$ and $i \in \{1, \dots, 5\}$. Then a $100(1 - \alpha)\%$ credibility interval for η_i is given by

$$CI_{1-\alpha}(\eta_i) = \left[\hat{q}_i\left(\frac{\alpha}{2}\right), \hat{q}_i\left(\frac{1-\alpha}{2}\right) \right], \quad \text{for any } i \in \{1, \dots, 5\}.$$

Furthermore, the estimation bias and the absolute percentage error (ape) of estimation are given, respectively, by

$$\text{bias}_{\eta_i} = \bar{\eta}_i - \eta_i \quad \text{and} \quad \text{ape}_{\eta_i} = \left| \frac{\text{bias}_{\eta_i}}{\eta_i} \right|, \quad \text{for any } i \in \{1, \dots, 5\}.$$

4.4 Results

The results obtained in this simulation study, by considering the scenarios described in Section 4.2, are the following.

Case 1: The Priors as Defined in Table 2.

Table 4 present the summary statistics for the samples obtained from the posterior distribution for each parameter of the FIEGARCH(0, d , 0) model. The statistics reported in this table (the same applies to Table 5) are the sample mean ($\bar{\eta}_i$), the sample standard deviation (sd_{η_i}) and the 95% credibility interval $CI_{0.95}(\eta_i)$ for the parameter η_i in $\boldsymbol{\eta} = (\nu, d, \theta, \gamma, \omega)' := (\eta_1, \dots, \eta_5)'$, for each $i \in \{1, \dots, 5\}$. The bold-face font for the mean indicates that the absolute percentage error of estimation (ape_{η_i}) in the corresponding case is higher than 0.10 (that is, 10%). The bold-face font for the credibility interval indicates that the true parameter value is not contained in the interval.

⁵In this work, the following definition is adopted (Brockwell and Davis, 1991). Given any $0 \leq \alpha \leq 1$, the number $q(\alpha)$ satisfying $\mathbb{P}(X \leq q(\alpha)) \geq \alpha$ and $\mathbb{P}(X \geq q(\alpha)) \geq 1 - \alpha$, is called a quantile of order α (or α quantile) for the random variable X (or for the distribution function of X).

Table 4: Summary for the sample obtained from posterior distributions considering all prior uniforms: mean $\bar{\eta}_i$, standard deviation sd_{η_i} and the 95% credibility interval $CI_{0.95}(\eta_i)$ for the parameter η_i in $\boldsymbol{\eta} = (\nu, d, \theta, \gamma, \omega)' := (\eta_1, \dots, \eta_5)'$, for each $i \in \{1, \dots, 5\}$. The true parameter values considered in this simulation are $d_0 \in \{0.10, 0.25, 0.35, 0.45\}$, $\nu_0 \in \{1.1, 1.5, 2.5, 5.0\}$, $\theta_0 = -0.15$, $\gamma_0 = -0.24$ and $\omega_0 = -5.4$.

d_0	ν_0	$\bar{\nu}$ (sd $_{\nu}$) $CI_{0.95}(\nu)$	\bar{d} (sd $_d$) $CI_{0.95}(d)$	$\bar{\theta}$ (sd $_{\theta}$) $CI_{0.95}(\theta)$	$\bar{\gamma}$ (sd $_{\gamma}$) $CI_{0.95}(\gamma)$	$\bar{\omega}$ (sd $_{\omega}$) $CI_{0.95}(\omega)$
0.10	1.1	1.093 (0.044) [0.989; 1.195]	0.181 (0.123) [0.005; 0.458]	-0.084 (0.041) [-0.171; -0.013]	0.236 (0.066) [0.093; 0.357]	-5.438 (0.058) [-5.551; -5.337]
	1.5	1.480 (0.069) [1.353; 1.635]	0.147 (0.079) [0.020; 0.330]	-0.177 (0.038) [-0.258; -0.106]	0.232 (0.052) [0.122; 0.340]	-5.420 (0.036) [-5.510; -5.338]
	1.9	2.088 (0.111) [1.908; 2.296]	0.093 (0.055) [0.004; 0.201]	-0.220 (0.032) [-0.286; -0.154]	0.216 (0.060) [0.105; 0.337]	-5.410 (0.035) [-5.486; -5.340]
	2.5	2.724 (0.140) [2.491; 3.027]	0.192 (0.076) [0.038; 0.330]	-0.201 (0.025) [-0.256; -0.153]	0.198 (0.045) [0.116; 0.287]	-5.388 (0.031) [-5.448; -5.333]
	5.0	5.297 (0.364) [4.641; 6.014]	0.101 (0.051) [0.015; 0.217]	-0.174 (0.020) [-0.215; -0.133]	0.297 (0.036) [0.232; 0.374]	-5.336 (0.028) [-5.383; -5.287]
0.25	1.1	1.108 (0.039) [1.028; 1.198]	0.265 (0.129) [0.016; 0.467]	-0.089 (0.038) [-0.176; -0.019]	0.230 (0.060) [0.108; 0.345]	-5.476 (0.052) [-5.553; -5.366]
	1.5	1.474 (0.067) [1.352; 1.644]	0.250 (0.071) [0.123; 0.393]	-0.175 (0.035) [-0.248; -0.100]	0.240 (0.053) [0.135; 0.353]	-5.426 (0.057) [-5.535; -5.344]
	1.9	2.072 (0.109) [1.885; 2.303]	0.245 (0.074) [0.068; 0.400]	-0.210 (0.032) [-0.273; -0.152]	0.223 (0.059) [0.116; 0.349]	-5.400 (0.062) [-5.537; -5.244]
	2.5	2.720 (0.148) [2.412; 3.016]	0.308 (0.060) [0.192; 0.426]	-0.200 (0.026) [-0.257; -0.155]	0.198 (0.042) [0.119; 0.281]	-5.367 (0.058) [-5.462; -5.274]
	5.0	5.311 (0.356) [4.676; 6.070]	0.226 (0.045) [0.137; 0.316]	-0.176 (0.019) [-0.215; -0.140]	0.293 (0.036) [0.225; 0.368]	-5.291 (0.035) [-5.346; -5.246]
0.35	1.1	1.099 (0.040) [1.009; 1.194]	0.349 (0.108) [0.093; 0.492]	-0.097 (0.038) [-0.178; -0.027]	0.230 (0.056) [0.121; 0.330]	-5.495 (0.090) [-5.674; -5.318]
	1.5	1.479 (0.065) [1.352; 1.639]	0.329 (0.065) [0.204; 0.461]	-0.178 (0.036) [-0.246; -0.106]	0.246 (0.052) [0.143; 0.340]	-5.423 (0.076) [-5.561; -5.302]
	1.9	2.064 (0.110) [1.843; 2.299]	0.364 (0.054) [0.227; 0.461]	-0.199 (0.030) [-0.265; -0.139]	0.233 (0.050) [0.139; 0.330]	-5.377 (0.090) [-5.535; -5.199]
	2.5	2.732 (0.150) [2.481; 3.031]	0.380 (0.052) [0.283; 0.479]	-0.201 (0.024) [-0.254; -0.161]	0.200 (0.043) [0.110; 0.285]	-5.307 (0.066) [-5.410; -5.149]
	5.0	5.229 (0.321) [4.603; 5.864]	0.318 (0.040) [0.243; 0.409]	-0.177 (0.019) [-0.216; -0.140]	0.289 (0.036) [0.227; 0.366]	-5.227 (0.046) [-5.298; -5.127]
0.45	1.1	1.096 (0.039) [1.010; 1.161]	0.436 (0.053) [0.313; 0.499]	-0.115 (0.034) [-0.187; -0.047]	0.241 (0.054) [0.136; 0.338]	-5.453 (0.128) [-5.716; -5.174]
	1.5	1.475 (0.073) [1.353; 1.627]	0.411 (0.048) [0.311; 0.494]	-0.179 (0.034) [-0.246; -0.110]	0.257 (0.048) [0.158; 0.353]	-5.411 (0.123) [-5.600; -5.133]
	1.9	2.052 (0.111) [1.846; 2.279]	0.450 (0.032) [0.385; 0.497]	-0.191 (0.026) [-0.238; -0.141]	0.243 (0.043) [0.165; 0.320]	-5.367 (0.130) [-5.614; -5.125]
	2.5	2.725 (0.152) [2.461; 3.021]	0.447 (0.032) [0.381; 0.495]	-0.206 (0.021) [-0.247; -0.167]	0.211 (0.041) [0.133; 0.296]	-5.150 (0.081) [-5.310; -4.985]
	5.0	5.177 (0.322) [4.553; 5.832]	0.417 (0.032) [0.348; 0.480]	-0.177 (0.019) [-0.220; -0.140]	0.286 (0.032) [0.228; 0.350]	-5.041 (0.068) [-5.182; -4.929]

Note: The bold-face font for the estimated **mean** indicates that the absolute percentage error is higher than 10%. The bold-face font for the **credibility interval** indicates that the interval does not contain the true parameter value.

From Table 4 one observes that the parameters ν and ω are always well estimated, in terms of absolute percentage error (ape), regardless the combination of $d_0 \in \{0.10, 0.25, 0.35, 0.45\}$ and $\nu_0 \in \{1.1, 1.5, 2.5, 5.0\}$ considered (the error is less than 10% in all cases). The credibility interval $CI_{0.95}(\nu)$ contains the true parameter value ν_0 in all cases, except when $d_0 = 0.10$ and $\nu_0 = 1.9$. Also, the estimation bias for ν is always negative when $\nu_0 < 1.9$ and positive when $\nu_0 \geq 1.9$, except for the combination $(\nu_0, d_0) = (1.1, 0.25)$. For the parameter ω , the credibility interval $CI_{0.95}(\omega)$ does not contain the true parameter value ($\omega_0 = -5.4$) in 5 out of 20 combinations of ν_0 and d_0 (see $\nu_0 = 5$ and all d_0 ; $\nu_0 = 2.5$ and $d_0 = 0.45$). Moreover, the estimation bias for ω is always negative when $\nu_0 \leq 1.5$ and always positive when $\nu_0 \geq 2.5$.

Table 4 also reports that $\text{ape}_\theta > 10\%$ for all combinations of d_0 and ν_0 . On the other hand, in most cases (14 out of 20), the credibility interval $CI_{0.95}(\theta)$ contains the true parameter value $\theta_0 = -0.15$. The cases for which $\theta_0 \notin CI_{0.95}(\theta)$ are $\nu = 1.9$ and $d_0 \in \{0.10, 0.25\}$ and $\nu_0 = 2.5$ and any d_0 . The bias for θ is always positive when $\nu_0 = 1.1$ (for any d_0) and negative in all other cases.

Furthermore, Table 4 shows that the parameter γ seems to be better estimated when the GED distribution presents heavy tails ($\nu_0 < 2$), except when $d = 0.10$, in which case $\text{ape}_\gamma > 10\%$ when $\nu_0 = 1.9$. Also, with exception of four cases ($d_0 = 0.10$ and $\nu_0 \in \{1.1, 1.5, 2.5\}$; $d_0 = 0.25$ and $\nu_0 = 2.5$), the parameter d is always well estimated. The bias for parameters d and γ does not seem to follow any pattern and both, $d_0 \in CI_{0.95}(d)$ and $\gamma_0 \in CI_{0.95}(\gamma)$, for any combination of ν_0 and d_0 .

Case 2: Gaussian Prior for $x = \phi^{-1}(d)$ and Uniform Priors for θ and γ .

Changing the prior for d does not yield significant difference on the estimation of ν , θ , γ and ω .

When the true value of d_0 is used to set $\mu_\phi = \phi^{-1}(d_0)$ (scenario C2.1), the best performance is observed by letting $\sigma_\phi = 0.15$. In this case, the absolute percentage error of estimation (ape_d) is smaller than 10% for all combinations of ν_0 and d_0 , with $d_0 \in \{0.10, 0.25, 0.35, 0.45\}$ and $\nu_0 \in \{1.1, 1.5, 2.5, 5.0\}$. If $\sigma_\phi = 0.10$ the chain takes too long to move from the initial point when $d_0 = 0.10$. When $\sigma_\phi = 0.25$, there is only one case for which $\text{ape}_d > 10\%$ ($d_0 = 0.10$ and $\nu_0 = 2.5$). In fact, in this case, $\text{ape}_d = 0.103$, which is still acceptable ($\sigma_\phi = 0.15$ still seems to be the best choice). Furthermore, as σ_ϕ increases, the number of cases for which $\text{ape}_d > 10\%$ also increases. For instance, when $\sigma_\phi \in \{0.50, 1.00, 3.00\}$, $\text{ape}_d > 10\%$ in 2, 4 and 10 cases, respectively.

When d_0 is assumed unknown and μ_ϕ is set to zero (scenario C2.2), $\sigma_\phi = 3$ seems to provide better results than smaller values of σ_ϕ . Under this scenario, $\text{ape}_d > 10\%$ for 8 out of 20 combinations of ν_0 and d_0 . Therefore, $d \sim \mathcal{U}(0, 0.5)$ still provides better estimates for the parameter d (see Table 4). Higher values of σ_ϕ do not improve the estimation of d . Too high values of σ_ϕ actually make the estimation worst. In particular, when $\sigma_\phi = 4$ the results are similar to $\sigma_\phi = 3$, if $d_0 > 0.1$. If $d_0 = 0.1$ then $\sigma_\phi = 3$ is slightly better than $\sigma_\phi = 4$. When $\sigma_\phi = 5$, ape_d is, in most cases, higher than when $\sigma_\phi = 3$. For σ_ϕ smaller than 3 the estimation bias is much higher. For instance, when $\sigma_\phi = 0.15$, $\text{ape}_d \leq 10\%$ only for $d_0 = 0.25$ (for all ν_0). For all other combinations of d_0 and ν_0 $\text{ape}_d > 20\%$. Also, when $b = 1$, $\text{ape}_d > 20\%$ in 12 out of 20 cases. In particular, $\text{ape}_d > 20\%$ for $d_0 = 0.10$ and all ν_0 . As it should be expected, C2.1 performs much better than C2.2.

Upon considering a two step estimator (scenario C2.3), no improvement is observed, when compared to scenario C2.2. In fact, the estimates obtained by letting $\mu_\phi = \phi^{-1}(\bar{d})$ (where \bar{d} is the estimate of d obtained in C2.2) and $\sigma_\phi = 0.15$ (the parameter which leads to the best performance in C2.1) are very close to \bar{d} itself.

Case 3: Gaussian Prior for $x = \phi^{-1}(d)$, Beta Prior for $-\theta$ and Uniform Prior for γ .

The estimation of ν , d , γ and ω is not significantly affected by the change in the prior for $-\theta$.

When the knowledge on the true parameter values d_0 and θ_0 is applied to set $\mu_\phi = \phi^{-1}(d_0)$, $\sigma_\phi = 0.15$, for each $d_0 \in \{0.10, 0.25, 0.35, 0.45\}$ (the best scenario in Case 2), and $b_1 = a_1(1 + \theta_0)(-\theta_0)^{-1}$ (scenario C3.1), it is observed that larger values of a_1 lead to better estimates for θ . Although any $a_1 \in \{110, 150, 200\}$ leads to $\text{ape}_\theta \leq 10\%$, for all combinations of $d_0 \in \{0.10, 0.25, 0.35, 0.45\}$ and $\nu_0 \in \{1.1, 1.5, 2.5, 5.0\}$, the best performance is obtained by setting $a_1 = 110$ ($b_1 \approx 623.333$). As a_1 decreases, the estimation performance decreases. For instance, when $a_1 = 100$, one case for which $\text{ape}_\theta > 10\%$ is observed. When $a_1 = 20$ the number of cases for which $\text{ape}_\theta > 10\%$ increases to 10 and no case for which $\text{ape}_\theta \leq 10\%$ is observed if $a_1 \in \{2.0, 0.1\}$. More specifically, for any $a_1 \in \{2.0, 0.1\}$, $10\% < \text{ape}_\theta \leq 20\%$ for $\nu_0 \in \{1.5, 5.0\}$ and all values of d_0 (8 out of 20 cases) and, in the remaining 12 cases, $\text{ape}_\theta > 20\%$.

By assuming θ_0 unknown (scenario C3.2) or by considering a two step estimator (scenario C3.3), no case for which $\text{ape}_\theta < 10\%$ is observed. The combinations of a_1 and b_1 tested in scenario C3.2 are: $(a_1, b_1) \in \{(2, 3), (2, 5), (2, 9), (4, 7), (5, 7), (10, 40), (10, 60), (10, 70), (100, 500), (100, 600)\}$. Among these values, the best performance is obtained when $a_1 = 10$ and $b_1 = 50$. In this case, $10\% < \text{ape}_\theta \leq 20\%$ in 12 out of 20 cases, which is slightly better than the performance obtained assuming $\theta \sim \mathcal{U}(0, 1)$ (in this case, $10\% < \text{ape}_\theta \leq 20\%$ in 8 out of 20 cases).

Case 4: Gaussian Prior for $x = \phi^{-1}(d)$, Beta Priors for $-\theta$ and γ .

Analogously to Case 2 and Case 3, the estimation of ν , d , θ and ω is not significantly affected by the change in the prior for γ .

By considering the true parameter values d_0 , θ_0 and γ_0 and setting $\mu_\phi = \phi^{-1}(d_0)$, $\sigma_\phi = 0.15$, for each $d_0 \in \{0.10, 0.25, 0.35, 0.45\}$ (the best scenario in Case 2), $a_1 = 110$, $b_1 = a_1(1 + \theta_0)(-\theta_0)^{-1}$ (the best scenario in Case 3) and $b_2 = a_2(1 - \gamma_0)\gamma_0^{-1}$ (scenario C4.1), it is observed the following: larger values of a_2 (smaller than a_1 , however) lead to better estimates for γ and as a_2 decreases, the estimation performance decays. For instance, when $a_2 = 40$ only one case for which $\text{ape}_\theta > 10\%$ is observed and when $a_2 \in \{10, 25, 30\}$, the number of cases increases to 5 ($d_0 = 0.45$ and all ν_0). On the other hand, any $a_2 \in \{50, 100\}$ gives $\text{ape}_\theta \leq 10\%$, for all combinations of $d_0 \in \{0.10, 0.25, 0.35, 0.45\}$ and $\nu_0 \in \{1.1, 1.5, 2.5, 5.0\}$. The simulation results for $a_2 = 50$ ($b_2 \approx 158.333$) are illustrated in Figure 3.

Figure 3 shows the sample mean (solid circle) and the 95% credibility interval (solid line) for the sample obtained from the posterior distribution of ν, d, θ, γ and ω (respectively, from top to bottom), for each combination of d_0 and ν_0 . The true parameter values $\nu_0, d_0, \theta_0, \gamma_0$ and ω_0 are represented in the corresponding row by the dashed line. The graphs related to θ, γ and ω (respectively, the third, fourth and fifth rows, from top to bottom) consider the same scale for all $d_0 \in \{0.10, 0.25, 0.35, 0.45\}$. Also, for the parameters θ, γ and ω , there is one graph for each d_0 and, for each one of these graphs, the true value of ν_0 is indicated in the x -axis.

From Figure 3 one observes that, for ν and ω , the conclusion regarding the estimation bias and the credibility intervals are basically the same as in Case 1 (see Table 4). On the other hand, under C4.1 of Case 4), $\text{ape}_{\eta_i} < 10\%$, for all $i \in \{1, \dots, 5\}$ and any combination of $d_0 \in \{0.10, 0.25, 0.35, 0.45\}$ and $\nu_0 \in \{1.1, 1.5, 2.5, 5.0\}$ (compare the parameter θ in Table 4 and Figure 3). As in Case 1, the bias for θ is always positive when $\nu_0 = 1.1$ (for any d_0) and negative when $\nu_0 > 1.1$ and the bias for the parameters d and γ does not seem to follow any pattern. Under the current scenario, d_0, θ_0 and γ_0 are all contained in the respective credibility




Figure 3: Posterior mean (solid circle), the true parameter value (dashed line) and the 95% credibility interval (solid line) for the parameters ν, d, θ, γ and ω (from top to bottom), for each combination of d_0 and ν_0 . The posterior distributions were obtained by considering an improper prior for ν , a Gaussian prior for $\phi^{-1}(d)$, Beta priors for $-\theta$ and γ and a uniform prior for ω . The true parameters values considered in this simulation are $d_0 \in \{0.10, 0.25, 0.35, 0.45\}$, $\nu_0 \in \{1.1, 1.5, 2.5, 5.0\}$, $\theta_0 = -0.15$, $\gamma_0 = -0.24$ and $\omega_0 = -5.4$.

intervals, for any combination of $d_0 \in \{0.10, 0.25, 0.35, 0.45\}$ and $\nu_0 \in \{1.1, 1.5, 2.5, 5.0\}$.

When the true value of γ_0 is not used to choose b_2 (scenario C4.2) similar results to the ones in Figure 3 are still obtained for some combinations of (a_2, b_2) . Not surprisingly, the pairs (a_2, b_2) which lead to good estimates are such that $a_2(a_2 + b_2)^{-1}$ (the mean μ_B of the prior distribution) is close to γ_0 . For instance, when $(a_2, b_2) \in \{(100, 300), (100, 350)\}$ (μ_B is, respectively, equal to 0.25 and 0.22, while $\gamma_0 = 0.24$) it is obtained $\text{ape}_\gamma < 10\%$ for all combinations of $d_0 \in \{0.10, 0.25, 0.35, 0.45\}$ and $\nu_0 \in \{1.1, 1.5, 2.5, 5.0\}$. The pair $(a_2, b_2) = (100, 350)$ provides slightly better results than $(a_2, b_2) = (100, 300)$ only when $\nu_0 = 5$. When $(a_2, b_2) = (100, 280)$ (so $\mu_B \approx 0.26$), $\text{ape}_\gamma \leq 10\%$ in 16 out of 20 cases (in the remaining 4 cases,

ape_γ does not exceed 13.4%).

On the other hand, choosing a_2 and b_2 such that $a_2(a_2 + b_2)^{-1}$ is close to the true γ_0 does not necessarily lead to good estimates. For instance, if $(a_2, b_2) = (5, 15)$ then $\mu_B = 0.25$, as it is when $(a_2, b_2) = (100, 300)$, but $\text{ape}_\gamma > 10\%$ in 6 out of 20 cases. Also, it is not evident that the more distant $a_2(a_2 + b_2)^{-1}$ is from γ_0 , the worst is the estimation. For instance, by letting $(a_2, b_2) \in \{(3, 15), (100, 440), (10, 30), (10, 40), (20, 80), (100, 400), (100, 270), (5, 10)\}$ then, respectively, $\mu_B \in \{0.167, 0.185, 0.200, 0.200, 0.200, 0.200, 0.270, 0.330\}$ and it is observed that $\text{ape}_\gamma > 10\%$ in 14, 20, 4, 10, 12, 16, 13 and 5 out of 20 cases, respectively.

Case 5: Beta Priors for $2d$, $-\theta$ and γ .

Analogously to all other cases, the estimation of ν , θ , γ and ω is not significantly affected by the change in the prior for d .

Upon assuming $-\theta \sim \text{Beta}(a_1, b_1)$ and $\gamma \sim \text{Beta}(a_2, b_2)$, with the same a_1 , a_2 , b_1 and b_2 as in scenario C4.1 of Case 4, and letting $2d \sim \text{Beta}(a_3, b_3)$, with $b_3 = a_3(1 - 2d_0)(2d_0)^{-1}$, for each $d_0 \in \{0.10, 0.25, 0.35, 0.45\}$ (scenario C5.1 of Case 5), the following is concluded. If $a_3 \in \{25, 50\}$ then $\text{ape}_d < 10\%$ for all $d_0 \in \{0.10, 0.25, 0.35, 0.45\}$. By increasing or decreasing too much the a_3 values the estimation performance decays. For instance, $a_3 \in \{0.10, 0.20, 2.00\}$ yields $\text{ape}_d > 10\%$ in 3, 1 and 7 cases, respectively.

Table 5 reports the simulation results for $a_3 = 25$ and $b_3 = a_3(1 - 2d_0)(2d_0)^{-1}$, which gives $b_1 \in \{100.000, 25.000, 10.714, 2.778\}$, respectively, for $d_0 \in \{0.10, 0.25, 0.35, 0.45\}$. The conclusions on the results presented in this table are the same as in Figure 3. Although the credibility intervals for d are slightly wider in Table 5 than in Figure 3, in both tables $d_0 \in CI_{0.95}(d)$ for any combination of $d_0 \in \{0.10, 0.25, 0.35, 0.45\}$ and $\nu_0 \in \{1.1, 1.5, 2.5, 5.0\}$.

As in Case 2, when considering a two step estimator (scenario C5.2), no improvement is observed, when compared to Case 1. In fact, once again, the estimates obtained by letting $a_3 = 25$ and $b_3 = a_3(1 - 2\bar{d})(2\bar{d})^{-1}$, where \bar{d} is the estimate of d obtained in Case 1, are very close to \bar{d} itself.

5 Conclusions

The Bayesian inference approach for parameter estimation on FIEGARCH models was described and a Monte Carlo simulation study was conducted to analyze the performance of the method under the presence of long-memory in volatility. The samples from FIEGARCH processes were obtained by considering the infinite sum representation for the logarithm of the volatility. A recurrence formula was used to obtain the coefficients for this representation. The generalized error distribution, with different tail-thickness parameters was considered so both innovation processes with lighter and heavier tails than the Gaussian distribution, were covered.

Markov Chain Monte Carlo (MCMC) methods were used to obtain samples from the posterior distribution of the parameters. A sensitivity analysis was performed by considering the following steps. First, an improper prior for ν and uniform priors d , θ , γ and ω were selected. In this case, only the basic set of information usually available in practice was considered. Second, non-uniform priors were selected for one or more parameters in $\{d, \theta, \gamma\}$. A Gaussian prior for $\phi^{-1}(d)$, with $\phi(\cdot)$ defined in (14), combined with uniform or Beta priors for θ ($-\theta$ in the Beta case) and γ was considered. In the sequel, a comparison was made by assuming Beta priors for $2d$, $-\theta$ and γ . The sensitivity analysis was completed by integrating (or not) the knowledge on

Table 5: Summary for the sample obtained from posterior distributions considering Beta priors for $2d, -\theta$ and γ : mean $\bar{\eta}_i$, standard deviation sd_{η_i} and the 95% credibility interval $CI_{0.95}(\eta_i)$ for the parameter η_i in $\boldsymbol{\eta} = (\nu, d, \theta, \gamma, \omega)' := (\eta_1, \dots, \eta_5)'$, for each $i \in \{1, \dots, 5\}$. The true parameter values considered in this simulation are $d_0 \in \{0.10, 0.25, 0.35, 0.45\}$, $\nu_0 \in \{1.1, 1.5, 2.5, 5.0\}$, $\theta_0 = -0.15$, $\gamma_0 = -0.24$ and $\omega_0 = -5.4$.

d_0	ν_0	$\bar{\nu}$ (sd $_{\nu}$) $CI_{0.95}(\nu)$	\bar{d} (sd $_d$) $CI_{0.95}(d)$	$\bar{\theta}$ (sd $_{\theta}$) $CI_{0.95}(\theta)$	$\bar{\gamma}$ (sd $_{\gamma}$) $CI_{0.95}(\gamma)$	$\bar{\omega}$ (sd $_{\omega}$) $CI_{0.95}(\omega)$
0.10	1.1	1.102 (0.045) [1.012; 1.198]	0.100 (0.018) [0.069; 0.139]	-0.144 (0.011) [-0.170; -0.124]	0.243 (0.026) [0.199; 0.294]	-5.472 (0.063) [-5.569; -5.347]
	1.5	1.483 (0.063) [1.364; 1.607]	0.102 (0.019) [0.069; 0.141]	-0.154 (0.013) [-0.178; -0.132]	0.239 (0.024) [0.190; 0.285]	-5.412 (0.047) [-5.524; -5.333]
	1.9	2.067 (0.108) [1.879; 2.309]	0.101 (0.016) [0.070; 0.134]	-0.160 (0.012) [-0.186; -0.137]	0.234 (0.027) [0.184; 0.291]	-5.422 (0.034) [-5.498; -5.360]
	2.5	2.702 (0.143) [2.442; 2.977]	0.106 (0.019) [0.074; 0.150]	-0.160 (0.012) [-0.186; -0.136]	0.231 (0.023) [0.187; 0.277]	-5.407 (0.022) [-5.470; -5.375]
	5.0	5.251 (0.391) [4.514; 6.080]	0.099 (0.017) [0.070; 0.132]	-0.158 (0.012) [-0.186; -0.134]	0.263 (0.024) [0.220; 0.312]	-5.343 (0.036) [-5.390; -5.265]
0.25	1.1	1.095 (0.044) [0.986; 1.190]	0.255 (0.031) [0.197; 0.315]	-0.145 (0.012) [-0.173; -0.123]	0.242 (0.028) [0.191; 0.295]	-5.455 (0.057) [-5.559; -5.348]
	1.5	1.485 (0.066) [1.357; 1.628]	0.252 (0.030) [0.193; 0.313]	-0.154 (0.013) [-0.180; -0.132]	0.240 (0.022) [0.195; 0.281]	-5.421 (0.046) [-5.529; -5.324]
	1.9	2.058 (0.113) [1.864; 2.316]	0.260 (0.030) [0.194; 0.309]	-0.159 (0.012) [-0.185; -0.138]	0.236 (0.028) [0.189; 0.293]	-5.431 (0.048) [-5.525; -5.340]
	2.5	2.719 (0.146) [2.452; 3.008]	0.271 (0.029) [0.216; 0.326]	-0.163 (0.012) [-0.188; -0.137]	0.229 (0.021) [0.186; 0.275]	-5.386 (0.038) [-5.469; -5.304]
	5.0	5.208 (0.326) [4.548; 5.871]	0.244 (0.028) [0.190; 0.299]	-0.159 (0.013) [-0.185; -0.134]	0.260 (0.023) [0.213; 0.309]	-5.313 (0.034) [-5.381; -5.256]
0.35	1.1	1.097 (0.038) [1.014; 1.175]	0.355 (0.034) [0.283; 0.413]	-0.145 (0.012) [-0.169; -0.126]	0.241 (0.027) [0.186; 0.298]	-5.469 (0.080) [-5.630; -5.289]
	1.5	1.481 (0.064) [1.359; 1.628]	0.349 (0.030) [0.285; 0.403]	-0.154 (0.012) [-0.178; -0.131]	0.239 (0.024) [0.192; 0.285]	-5.447 (0.077) [-5.587; -5.310]
	1.9	2.070 (0.102) [1.870; 2.288]	0.370 (0.029) [0.306; 0.421]	-0.160 (0.012) [-0.183; -0.136]	0.236 (0.027) [0.186; 0.293]	-5.414 (0.082) [-5.587; -5.237]
	2.5	2.720 (0.147) [2.449; 3.009]	0.375 (0.028) [0.321; 0.426]	-0.163 (0.011) [-0.185; -0.143]	0.228 (0.023) [0.186; 0.274]	-5.337 (0.063) [-5.440; -5.221]
	5.0	5.147 (0.346) [4.598; 5.965]	0.344 (0.027) [0.287; 0.395]	-0.159 (0.012) [-0.185; -0.137]	0.258 (0.023) [0.214; 0.305]	-5.255 (0.048) [-5.321; -5.171]
0.45	1.1	1.101 (0.042) [1.024; 1.191]	0.454 (0.024) [0.402; 0.489]	-0.145 (0.012) [-0.169; -0.122]	0.238 (0.026) [0.189; 0.286]	-5.424 (0.128) [-5.682; -5.160]
	1.5	1.493 (0.073) [1.362; 1.645]	0.450 (0.024) [0.395; 0.488]	-0.154 (0.012) [-0.177; -0.132]	0.243 (0.024) [0.195; 0.291]	-5.414 (0.132) [-5.681; -5.139]
	1.9	2.045 (0.109) [1.844; 2.241]	0.464 (0.019) [0.419; 0.491]	-0.158 (0.011) [-0.181; -0.134]	0.239 (0.026) [0.193; 0.291]	-5.424 (0.126) [-5.717; -5.216]
	2.5	2.742 (0.142) [2.507; 3.010]	0.466 (0.017) [0.431; 0.493]	-0.164 (0.012) [-0.191; -0.141]	0.228 (0.022) [0.189; 0.275]	-5.170 (0.084) [-5.308; -5.002]
	5.0	5.164 (0.346) [4.558; 5.883]	0.447 (0.022) [0.396; 0.486]	-0.158 (0.011) [-0.182; -0.137]	0.256 (0.022) [0.214; 0.300]	-5.070 (0.076) [-5.227; -4.942]

Note: The bold-face font for the **credibility interval** indicates that the interval does not contain the true parameter value.

the true parameter values to select the hyperparameter values.

An example was presented to illustrate the similarities or differences on the mean, standard deviation and credibility intervals estimated by considering a chain of size $N = 200801$, a thinned chain (thinning parameter 200 and burn-in size 1000) and a sample of size 1000 (obtained from the larger chain, after the burn-in of size 1000). Given the ergodicity of the Markov chain, the posterior means for all three chains were very close. The differences on the standard deviations and credibility intervals are not significant enough to justify the use of the entire or thinned chains. Although the example only presents the case $d_0 = 0.25$, the same conclusions apply to $d_0 \in \{0.10, 0.35, 0.45\}$.

The simulation study showed that if the prior of one or more parameters is changed, the estimation of the other parameters is not significantly affected. The parameters ν and ω are always well estimated, in terms of absolute percentage error, regardless priors considered for d, θ and γ , for any combination of $d_0 \in \{0.10, 0.25, 0.35, 0.45\}$ and $\nu_0 \in \{1.1, 1.5, 2.5, 5.0\}$. With a few exceptions, the true parameter value ν_0 was contained in the 95% credibility interval, for any combination of $\nu_0 \in \{1.1, 1.5, 1.9, 2.5, 5.0\}$ and $d_0 \in \{0.10, 0.25, 0.35, 0.45\}$ considered. The true parameter value ω_0 was not contained in any credibility interval when $\nu_0 = 5$.

Regardless the prior considered, the parameter d is usually better estimated when $d \in \{0.35, 0.45\}$. The Gaussian prior for $\phi^{-1}(d)$ only provided better results (globally) when the knowledge on the true parameter value d_0 was used to set $\mu_\phi = \phi^{-1}(d_0)$ and σ_ϕ was set to some value smaller or equal than 1. In particular, only when $b = 0.15$ the absolute percentage error of estimation (ape) became smaller than 10% for all $d_0 \in \{0.10, 0.25, 0.35, 0.45\}$. Although the credibility intervals for d are slightly wider when a Beta prior is considered, the use of the Beta prior for $2d$ neither improves nor degrades the estimation performance, compared to the Gaussian prior for $\phi^{-1}(d)$.

The absolute percentage error of estimation for θ (ape_θ) only became smaller than 10% when the Beta prior was considered and the true value of the parameter was used to select the hyperparameter. When θ_0 was assumed unknown the ape_θ was always between 10% and 38.1%. The parameter γ is always better estimated than θ , for any priors considered. Similar to d and θ , the best performance is obtained when the true parameter value is used to select the hyperparameters. On the other hand, γ is the only parameter for which there are hyperparameter values that do not yield $\mu_B = \gamma_0$ (μ_B is the mean of the prior distribution and γ_0 is the true parameter value) while still providing good estimates.

Acknowledgments

T.S. Prass was supported by CNPq-Brazil. S.R.C. Lopes was partially supported by CNPq-Brazil, by CAPES-Brazil, by INCT em *Matemática* and by Pronex *Probabilidade e Processos Estocásticos* - E-26/170.008/2008 -APQ1. The authors are grateful to the (Brazilian) National Center of Super Computing (CESUP-UFRGS) for the computational resources.

References

- Baillie, R.; T. Bollerslev and H.O. Mikkelsen (1996). "Fractionally Integrated Generalized Autoregressive Conditional Heteroskedasticity". *Journal of Econometrics*, vol. 74, 3-30.
- Bollerslev, T. (1986). "Generalized Autoregressive Conditional Heteroskedasticity". *Journal of Econometrics*, vol. 31, 307-327.

- Bollerslev, T. and H.O. Mikkelsen (1996). "Modeling and Pricing Long Memory in Stock Market Volatility". *Journal of Econometrics*, vol. 73, 151-184.
- Breidt, F.; N. Crato and P.J.F. de Lima (1998). "On the Detection and Estimation of Long Memory in Stochastic Volatility". *Journal of Econometrics*, vol. 83, 325-348.
- Brockwell, P.J. and R.A. Davis (1991). *Time Series: Theory and Methods*. Second Edition. New York: Springer-Verlag.
- Casella, G. and E.I. George (1992). "Explaining the Gibbs Sampler", *The American Statistician*, vol 46(3), 167-174.
- Chib, S. and E. Greenberg (1995). "Understanding the Metropolis-Hastings Algorithm", *The American Statistician*, vol. 49(4), 327-335.
- Engle, R.F. (1982). "Autoregressive Conditional Heteroskedasticity with Estimates of Variance of U.K. Inflation". *Econometrica*, vol. 50, 987-1008.
- Gelfand, A.E. and A.F.M Smith (1990). "Sampling-Based Approaches to Calculating Marginal Densities". *Journal of the American Statistical Association*, vol. 85(410), 398-409.
- Geman, S. and D. Geman (1984). "Stochastic Relaxation, Gibbs Distributions and the Bayesian Restoration of Images". *IEEE Transactions on Pattern Analysis and Machine Intelligence*, vol. 12, 609-628.
- Gelman, A. and D. Rubin (1992). "Inference from Iterative Simulation Using Multiple Sequences". *Statistical Science*, vol. 7, 457-511.
- Geyer, C. J. (1992). "Practical Markov chain Monte Carlo". *Statistical Science*, vol. 7, 473-511.
- Hastings, W.K. (1970). "Monte Carlo Sampling Methods Using Markov Chains and Their Applications". *Biometrika*, vol. 57(1), 97-109.
- Lopes, S.R.C and T.S. Prass (2013). "Theoretical Results on Fractionally Integrated Exponential Generalized Autoregressive Conditional Heteroskedastic Processes." Working Paper.
- MacEachern, S.N. and L.M. Berliner (1994). "Subsampling the Gibbs Sampler". *The American Statistician*, vol. 48, 188-190.
- Metropolis, N.; A.W. Rosenbluth; M.N. Rosenbluth; A.H. Teller and E. Teller (1953). "Equations of State Calculations by Fast Computing Machines". *Journal of Chemical Physics*, vol. 21(6), 1087-1092.
- Meyer, R. and T. Yu (2000). "Bugs for a Bayesian Analysis of Stochastic Volatility Models". *Econometrics Journal*, vol. 3, 198-215.
- Nelson, D.B. (1991). "Conditional Heteroskedasticity in Asset Returns: A New Approach". *Econometrica*, vol. 59, 347-370.
- Ruiz E. and H. Veiga (2008). "Modelling long-memory volatilities with leverage effect: A-LMSV versus FIEGARCH". *Computational Statistics and Data Analysis*, vol. 52(6), 2846-2862.
- Smith, A.F.M. and G.O. Roberts (1993). "Bayesian computation via the Gibbs sampler and related Markov chain Monte Carlo methods". *Journal of the Royal Statistical Society, Ser. B*, vol. 55, 3-23.