EVIDENCE FOR THE DYNAMICAL BRAUER-MANIN CRITERION

EKATERINA AMERIK, PÄR KURLBERG, KHOA NGUYEN, ADAM TOWSLEY, BIANCA VIRAY, AND JOSÉ FELIPE VOLOCH

ABSTRACT. Let $\varphi \colon X \to X$ be a morphism of a variety over a number field K. We consider local conditions and a "Brauer-Manin" condition, defined by Hsia and Silverman, for the orbit of a point $P \in X(K)$ to be disjoint from a subvariety $V \subseteq X$, i.e., for $\mathcal{O}_{\varphi}(P) \cap V = \emptyset$. We provide evidence that the dynamical Brauer-Manin condition is sufficient to explain the lack of points in the intersection $\mathcal{O}_{\varphi}(P) \cap V$; this evidence stems from a probabilistic argument as well as unconditional results in the case of étale maps.

1. INTRODUCTION

In recent work, Hsia and Silverman (Hsia and Silverman, 2009) ask a dynamical question in analogy with a question of Scharaschkin (Scharaschkin, 1999); the dynamical question is as follows. Let $\varphi \colon X \to X$ be a self-morphism of a variety over a number field K, let $V \subseteq X$ be a subvariety, and let $P \in X(K)$. Hsia and Silverman ask whether the closure of the intersection of the orbit

$$\mathcal{O}_{\varphi}(P) := \{P, \varphi(P), \varphi(\varphi(P)), \ldots\}$$

with the subvariety V is equal to the intersection of $V(\mathbb{A}_K)$ with the closure of $\mathcal{O}_{\varphi}(P)$ in the adelic topology, i.e. whether

$$\mathcal{C}(V(K) \cap \mathcal{O}_{\varphi}(P)) = V(\mathbb{A}_K) \cap \mathcal{C}(\mathcal{O}_{\varphi}(P)),$$

where $\mathcal{C}(-)$ denotes the closure in the adelic topology.

The purpose of this paper is to give some justification to the assertion that a closely related question has a positive answer. In particular, when $K = \mathbb{Q}$, (and a choice of an integral model for V is made) we give evidence for the assertion that

$$V(K) \cap \mathcal{O}_{\varphi}(P) = \emptyset \Rightarrow V(\mathbb{Z}/m\mathbb{Z}) \cap (\mathcal{O}_{\varphi}(P) \bmod m) = \emptyset$$
(1.1)

for some integer m, as long as the map φ is sufficiently random.

Our evidence is twofold: a probabilistic argument that holds under certain randomness assumptions, and unconditional results in the case that φ is étale and V is φ^k -invariant or φ -preperiodic.

The probabilistic argument is given in Section 2 and is based on a probabilistic argument of Poonen (Poonen, 2006) for the original question of Scharaschkin. Precisely, we show the following:

E.A. was partially supported by AG Laboratory NRU-HSE, RF government grant, ag. 11.G34.31.0023. Her research was carried out within the NRU HSE Academic Fund Program for 2013-2014, research grant No. 12-01-0107. P.K. was partially supported by grants from the Göran Gustafsson Foundation and the Swedish Research Council. K.N. was partially supported by NSF grant DMS-0901149. B.V. was supported by ICERM, the Centre Interfacultaire Bernoulli, and NSF grant DMS-1002933. J.F.V. was supported by the Simons Foundation (grant #234591) and by the Centre Interfacultaire Bernoulli.

Theorem 1.1. Assume that φ is sufficiently random (see Section 2.1 for the precise statement) and that the lengths of the cyclic part of the orbit $\mathcal{O}_{\varphi}(P)$ modulo p are integers with at least the same probability of being smooth as that of a random integer of similar size. Then there exists a sequence of squarefree integers m such that

$$\operatorname{Prob}(C_m \cap V(\mathbb{Z}/m\mathbb{Z}) = \emptyset) = 1 - o(1)$$

as $m \to \infty$. Here C_m denotes the cyclic part of $\mathcal{O}_{\varphi}(P)$ in $X(\mathbb{Z}/m)$.

Remark 1.2. If the intersection of the orbit $\mathcal{O}_{\varphi}(P)$ modulo m with $V(\mathbb{Z}/m\mathbb{Z})$ is contained only in the tail of $\mathcal{O}_{\varphi}(P) \mod m$, then the intersection $(\mathcal{O}_{\varphi}(P) \mod m) \cap V(\mathbb{Z}/m\mathbb{Z})$ consists of finitely many *iterates* of P under φ . More specifically, $\varphi^n(P) \mod m \notin V(\mathbb{Z}/m\mathbb{Z})$ for all $n > N_0$, where φ^n denotes the composition of φ with itself n times. Therefore $V(K) \cap \mathcal{O}_{\varphi}(P)$ is contained in

$$\{P, \varphi(P), \varphi^2(P), \dots, \varphi^{N_0}(P)\},$$

and so can be computed by a finite computation. Therefore, it is reasonable to restrict to the cyclic part of the orbit in the above theorem.

In Section 3, we provide numerical evidence for the randomness assumptions needed in the heuristic argument from Section 2. We also describe experiments on randomly generated morphisms of \mathbb{A}^5 which support the argument that (1.1) holds.

The unconditional results are the focus of Section 4. Assume that X is quasi-projective, that φ is étale, and that $\varphi^k(V) \subseteq V$, i.e., that V is φ^k -invariant, for some positive integer k. Under these assumptions we show that if $\mathcal{O}_{\varphi}(P) \cap V = \emptyset$, then for all but finitely many primes p, there exists an n = n(p) such that $\mathcal{O}_{\varphi}(P) \mod p^n \cap V(\mathbb{Z}/p^n\mathbb{Z}) = \emptyset$. If every irreducible component of V is preperiodic, i.e., $\varphi^{k_0+k}(V) = \varphi^{k_0}(V)$ for some integers k_0, k with k > 0, Xis quasi-projective, and φ is étale and closed, then we obtain the same result. More generally, we prove

Theorem 1.3. Let X be a quasi-projective variety over a global field K. Assume that φ is étale, and either (1) that V is φ^m -invariant or (2) that φ is closed and every irreducible component of V is φ -preperiodic. If $P \in X(K)$ is such that $V(K) \cap \mathcal{O}_{\varphi}(P) = \emptyset$ then, for all but finitely many primes v,

$$V(K_v) \cap \mathcal{C}_v(\mathcal{O}_\varphi(P)) = \emptyset,$$

where $C_v(-)$ denotes closure in the v-adic topology.

In 4.6, we discuss whether the assumption that V is preperiodic or φ^k -invariant can be weakened in any way.

Acknowledgements

This project was started during the workshop on Global Arithmetic Dynamics, part of the semester program on Complex and Arithmetic Dynamics at the Institute for Computational and Experimental Research Mathematics (ICERM). We thank Zoé Chatzidakis, Alice Medvedev, and Thomas Scanlon for helpful discussions during the workshop, and we are grateful to the ICERM staff and the organizers of the workshop, Xander Faber, Michelle Manes, Lucien Szpiro, Thomas Tucker, and Michael Zieve, for their support. We thank Serge Cantat for pointing out a construction in Section 4.6.

2. Probabilistic Proof

Let X be a projective variety over \mathbb{Q} , let $V \subset X$ be a closed \mathbb{Q} -subvariety, and let $\varphi \colon X \to X$ be a \mathbb{Q} -morphism. We write

$$d_1 := \dim(X),$$
 and $d_2 := \dim(V).$

The goal of this section is to prove, under some randomness assumptions, that if $\mathcal{O}_{\varphi}(P) \cap V = \emptyset$, then with probability 1 there exists a positive integer m such that, roughly speaking, the cyclic part of $\mathcal{O}_{\varphi}(P) \mod m$ is disjoint from $V(\mathbb{Z}/m\mathbb{Z})$. To make this precise, we need some further notation. Fix a finite set of primes S such that X, V and P extend to flat projective models \mathscr{X}, \mathscr{V} and \mathscr{P} , respectively, over \mathbb{Z}_S . After possibly enlarging S, we also assume that φ extends to a morphism $\tilde{\varphi} \colon \mathscr{X} \to \mathscr{X}$. Then for any integer m which is relatively prime to all elements of S, we may consider the base change $X_m := \mathscr{X} \times_{\mathbb{Z}_S} \mathbb{Z}/m\mathbb{Z}$. Similarly, we have $P_m := \mathscr{P} \times_{\mathbb{Z}_S} \mathbb{Z}/m\mathbb{Z}, V_m := \mathscr{V} \times_{\mathbb{Z}_S} \mathbb{Z}/m\mathbb{Z}$ and $\varphi_m := \tilde{\varphi}|_{X_m} \colon X_m \to X_m$. We will often abuse notation and write $V(\mathbb{Z}/m)$ for $V_m(\mathbb{Z}/m)$.

We write

$$O_m := \{P_m, \varphi_m(P_m), \varphi_m(\varphi_m(P_m)), \ldots\}.$$

As O_m is contained in the finite set $X_m(\mathbb{Z}/m)$, there is some pair of non-negative integers $k_0 < k_1$ such that $\varphi_m^{k_0}(P_m) = \varphi_m^{k_1}(P_m)$. Let k_0 be the minimal such integer; then we define the cyclic part of O_m as

$$C_m := \{\varphi_m^{k_0}(P_m), \varphi_m^{k_0+1}(P_m), \varphi_m^{k_0+2}(P_m), \ldots\}.$$

2.1. Assumptions on φ and V. With the above notation, our randomness assumptions on φ and V are then based on the following heuristics:

- (i) The reduction of morphisms $\varphi \colon X \to X$ modulo p behave like random maps on a finite set $S = X(\mathbb{F}_p)$.
- (ii) For any $y \in X(\mathbb{F}_p)$, the condition that $y \in O_p$ is independent from the condition that $y \in V(\mathbb{F}_p)$.
- (iii) For any $y \in X(\mathbb{Q}) \setminus V(\mathbb{Q})$, the condition that $\bar{y} \in V(\mathbb{F}_p)$ is independent from the condition that $\bar{y} \in V(\mathbb{F}_q)$ for primes $p \neq q$.

These heuristics are made precise as follows.

(i) Orbit size assumption: For all primes $p \notin S$,

$$|O_p| = p^{d_1/2 + o(1)}, \text{ and } |C_p| = p^{d_1/2 + o(1)}.$$
 (2.1)

(ii) Independence assumption on $\mathcal{O}_{\varphi}(P)$ and V modulo p: For all primes $p \notin S$

$$\operatorname{Prob}(x_1, \dots, x_k \notin V(\mathbb{F}_p) \mid x_1, \dots, x_k \in O_p) = (1 - 1/p^{d_1 - d_2 + o(1)})^k, \text{ and} \\ \operatorname{Prob}(x_1, \dots, x_k \notin V(\mathbb{F}_p) \mid x_1, \dots, x_k \in C_p) = (1 - 1/p^{d_1 - d_2 + o(1)})^k,$$

$$(2.2)$$

for $k = O(p^{d_1/2 + o(1)})$, and x_1, \ldots, x_k distinct points modulo p.

(iii) Asymptotic independence modulo (large) primes: if $\mathcal{O}_{\varphi}(P) \cap V(\mathbb{Q}) = \emptyset$, then

$$\operatorname{Prob}(O_p \cap V(\mathbb{F}_p) \neq \emptyset \ \forall p > T) = (1 + o(1)) \cdot \prod_{p > T} \operatorname{Prob}(O_p \cap V(\mathbb{F}_p) \neq \emptyset),$$

$$\operatorname{Prob}(C_p \cap V(\mathbb{F}_p) \neq \emptyset \ \forall p > T) = (1 + o(1)) \cdot \prod_{p > T} \operatorname{Prob}(C_p \cap V(\mathbb{F}_p) \neq \emptyset).$$
(2.3)

Remark 2.1. Any two different models will differ at only finitely many primes. Therefore, since assumptions (2.1)–(2.3) say something about behavior at sufficiently large primes, these assumptions are really assumptions on ϕ, X, V and P and not $\tilde{\phi}, \mathcal{X}, \mathcal{V}, \mathcal{P}$.

If A is a (large) finite set, and $f : A \to A$ is a sufficiently random map, then cardinality of the forward orbit of a random starting point is likely to be of size $|A|^{1/2+o(1)}$ (Flajolet and Odlyzko, 1990). This, together with the Weil conjectures which give that $|X(\mathbb{F}_p)| = p^{d_1+o(1)}$ and $|V(\mathbb{F}_p)| = p^{d_2+o(1)}$, shows that the above heuristics imply the precise assumptions (2.1)– (2.3).

Remark 2.2. We warn the reader that there are maps of special type for which the random map heuristic does not apply. For example, for linear automorphisms on \mathbb{P}^n , orbits can be as large as $p^{d_1+o(1)}$.

2.2. Nonempty intersections modulo p. Under assumptions (2.1)-(2.3) and if V has sufficiently small codimension, then, with probability 1, we will have $\mathcal{O}_{\varphi}(P) \cap V(\mathbb{F}_p) \neq \emptyset$ for all but finitely many p even if $\mathcal{O}_{\varphi}(P)$ and V have empty intersection. The intuition is as follows. If V is, say, of codimension 1 in X, then the probability that the reduction of $\varphi^n(P)$ modulo p is contained in $V(\mathbb{F}_p)$ should equal 1/p. Furthermore, if the orbit is of length $p^{d_1/2}$, as we would expect, then the likelihood of $V(\mathbb{F}_p) \cap O_p = \emptyset$ should be given by $(1 - 1/p)^{|O_p|} = \exp(-|O_p|/p)$, assuming p is sufficiently large. Thus, if the orbits are long, there will be "accidental" intersections modulo p.

Proposition 2.3. Assume that assumptions (2.1)–(2.3) hold, and that $d_2 > d_1/2$. Then, as $T \to \infty$,

$$\operatorname{Prob}(O_p \cap V(\mathbb{F}_p) \neq \emptyset \ \forall p > T) = 1 - e^{-T^{d_2 - d_1/2 + o(1)}}$$

Remark 2.4. The proposition still holds under a relaxed orbit size assumption: it is sufficient to assume $|O_p| = p^{d'_1/2 + o(1)}$ for some $d'_1 \in \mathbb{Z}_{\geq 0}$ such that $d_2 > d'_1/2$.

Proof. By assumption (2.1), $|O_p| = p^{d_1/2 + o(1)}$, and by the Weil conjectures, $|V(\mathbb{F}_p)| = p^{d_2 + o(1)}$, $|X(\mathbb{F}_p)| = p^{d_1 + o(1)}$. Therefore, assumption (2.2) gives that

$$\operatorname{Prob}(O_p \cap V(\mathbb{F}_p) = \emptyset) = \left(1 - p^{d_2 - d_1 + o(1)}\right)^{|O_p|} = \left(1 - p^{d_2 - d_1 + o(1)}\right)^{p^{d_1/2 + o(1)}} = e^{-p^{d_1/2 - (d_1 - d_2) + o(1)}} = e^{-p^{d_2 - d_1/2 + o(1)}}.$$

In particular,

$$\operatorname{Prob}(O_p \cap V(\mathbb{F}_p) \neq \emptyset) = 1 - e^{-p^{d_2 - d_1/2 + o(1)}}$$

and therefore, by assumption (2.3),

$$\operatorname{Prob}(O_p \cap V(\mathbb{F}_p) \neq \emptyset \ \forall p > T) = \prod_{p > T} (1 - e^{-p^{d_2 - d_1/2 + o(1)}}) = 1 - e^{-T^{d_2 - d_1/2 + o(1)}}.$$

2.3. Empty intersections for some composite m. The situation is quite different over composite integers m. Indeed, the main result of this section is that, assuming some further randomness properties, there exist integers m such that the probability that C_m and $V(\mathbb{Z}/m\mathbb{Z})$ are disjoint is arbitrarily close to 1.

We begin by recalling some background on smooth numbers.

Definition 2.5. An integer n is y-smooth if all primes p dividing n are bounded above by y. Define

$$\psi(x, y) := |\{n \le x : n \text{ is } y - \text{smooth}\}|.$$

Smooth integers have the following well-known distribution (Tenenbaum and Mendès France, 2000, Thm. 10, p. 97) for $\alpha \in (0, 1)$ and x tending to infinity,

$$\psi(x, x^{\alpha}) = (1 + o(1)) \cdot x\rho(u)$$

where $u := \log x / \log x^{\alpha} = 1/\alpha$, and $\rho(u) \in (0, 1)$ for $u \in (1, \infty)$.

Our analysis will be based on the following heuristic: that $|C_p|$ has the same "likelihood" of being smooth as a random integer of the same size. By (2.1), $|C_p| = p^{d_1/2+o(1)}$, so the heuristic implies that the density of primes p for which $|C_p| = p^{d_1/2+o(1)}$ is p^{α} -smooth equals $\rho(u)$ where $u = \log p^{d_1/2+o(1)}/\log p^{\alpha} = d_1(1+o(1))/(2\alpha)$, and hence that

$$\prod_{\substack{p \le x \\ C_p \mid \text{ is } x^{\alpha}-\text{smooth}}} p = \exp\left(\sum_{\substack{p \le x \\ |C_p| \text{ is } x^{\alpha}-\text{smooth}}} \log p\right) = \exp\left(x \cdot \rho\left(\frac{d_1(1+o(1))}{2\alpha}\right)\right).$$

This heuristic leads us to the following precise cycle length smoothness assumption: For α and d_1 fixed and $x \to \infty$,

$$\prod_{\substack{p \le x \\ |C_p| \text{ is } x^{\alpha} - \text{smooth}}} p = \exp\left(x \cdot \rho\left(\frac{d_1(1+o(1))}{2\alpha}\right)\right).$$
(2.4)

Proposition 2.6. Assume (2.1)–(2.4). Then there exists a sequence of squarefree integers m such that

$$\operatorname{Prob}(C_m \cap V(\mathbb{Z}/m\mathbb{Z}) = \emptyset) = 1 - o(1)$$

as $m \to \infty$.

Proof. Define

$$m_{x,\alpha} := \prod_{\substack{p \le x \\ |C_p| \text{ is } x^{\alpha} \text{-smooth}}} p.$$

Since for any square free integer M, $|C_M| = \lim_{p|M} |C_p|$, we have, as $x \to \infty$,

$$|C_{m_{x,\alpha}}| \le \prod_{p \le x^{\alpha}} p^{\log x^{d_1/2 + o(1)/\log(p)}} = \exp\left(\sum_{p \le x^{\alpha}} (d_1/2 + o(1)) \log x\right)$$
$$= \exp\left((d_1/2 + o(1)) \cdot \log x \cdot \pi(x^{\alpha})\right) = \exp\left(x^{\alpha} \cdot \frac{d_1 \cdot (1 + o(1))}{2\alpha}\right).$$

In particular, $|C_{m_{x,1/3}}| = \exp(O(x^{1/3}))$. Therefore, if we take

$$m = m_{x,1/3} = \prod_{\substack{p \le x \\ |C_p| \text{ is } x^{1/3} \text{-smooth}}} p$$

we find that $m = \exp(\Theta(x))$, thus $|C_m| = \exp(O(x^{1/3})) = m^{o(1)}$, and hence

$$\operatorname{Prob}(C_m \cap V(\mathbb{Z}/m\mathbb{Z}) = \emptyset) = \left(1 - \frac{1}{m^{d_1 - d_2}}\right)^{|C_m|} = \left(1 - \frac{1}{m^{d_1 - d_2}}\right)^{m^{d_1 - d_2} \frac{|C_m|}{m^{d_1 - d_2}}} = e^{-\frac{|C_m|}{m^{d_1 - d_2}}} = e^{-o(1)} = 1 - o(1).$$

s $x \to \infty$.

a

3. Computations

3.1. Cycle length smoothness assumption. We ran experiments, detailed below, to justify the assumption (2.4) on the smoothness of the cycle lengths. Our experiments do not confirm this assumption. Fortunately, it is clear from the proof of proposition 2.6 that we only need that the cycle lengths be at least as smooth as the prediction (2.4) and this is what we see in the experiments. We also found some maps with special properties for which the cycle lengths are even smoother. We conjecture that cycle lengths be at least as smooth as the prediction (2.4) in all cases but we don't know how to explain the extra smoothness shown in the experiments.

We will consider three maps:

$$\begin{split} \varphi \colon \mathbb{P}^1 \to \mathbb{P}^1, \quad & (x:y) \mapsto (x^2 + 5y^2 : y^2), \\ \psi \colon \mathbb{P}^2 \to \mathbb{P}^2, \quad & (x:y:z) \mapsto \left(x^2 + y^2 : x^2 + 3y^2 - 2xy + z^2 : z^2\right), \text{ and} \\ \sigma \colon \mathbb{P}^3 \to \mathbb{P}^3, \quad & (x:y:z:w) \mapsto \left(x^2 + y^2 - z^2 + yw + w^2 : x^2 - xy + xz + 2w^2 : z^2 - yz + xz + 3w^2 : w^2\right), \end{split}$$

For each prime less than 100,000, 500,000 and 1,000,000 respectively we compute C_p , the length of the periodic cycle length of [1:1], [1:1:1], or [1:1:1:1] under φ , ψ , and σ

respectively in \mathbb{F}_p . Setting $\alpha = 1/3$, we compute $S(x) := \left[\prod_{\substack{p \le x \\ |C_p| \text{ is } x^{\alpha} - \text{smooth}} p\right]$ at each prime

in the range specified. We then create the graphs below which compare $\log S(x)$ to the predicted value of $x \cdot \rho(u)$ where $u = \frac{d_1}{2\alpha}$ and ρ is the Dickman ρ -function. Recall that assumption (2.4) states that $\log S(x)$ should behave linearly; the graphs (figures 1, 2, and 3) below) support this assumption. The data appears linear for large enough x, with slope at least as big as predicted.

All computations were performed using C and Sage 4.8 (Stein, 2013).




FIGURE 1. $\log (S(x))$ for $\varphi(x)$ and x < 100,000




FIGURE 2. $\log (S(x))$: for $\psi(x)$ and x < 500,000

3.2. Experiments. In this section, we let $X = \mathbb{A}^5$ and let $V = V(1 - x^2 - y^2 - z^2 - w^2 - v^2)$. Fix a point $P \in X(\mathbb{Z})$ and consider a morphism $\phi: X \to X$ with integer coefficients; then $\mathcal{O}_{\phi}(P) \subseteq X(\mathbb{Z})$. As V contains few integral points, namely only those points with exactly one coordinate equal to ± 1 and the remaining coordinates 0, one expects the intersection




FIGURE 3. log (S(x)) for $\sigma(x)$ and x < 1,000,000

 $\mathcal{O}_{\phi}(P) \cap V$ to be empty. Thus, by the arguments in §2, we expect to find an integer m such that $O_m(P) \cap V(\mathbb{Z}/m\mathbb{Z}) = \emptyset$.

We considered a fixed integer starting point P and 500 randomly generated morphisms from $X \to X$ with integer coefficients. For each of these morphisms, we computed whether the intersection $O_m(P) \cap V(\mathbb{Z}/m\mathbb{Z})$ was empty for some positive integer $m \leq 2000$. The results are as follows:

- (i) For 86.8% of these maps, i.e., 434 out of 500, there exists a prime power $q \leq 2000$ such that $O_q(P) \cap V(\mathbb{Z}/q\mathbb{Z}) = \emptyset$.
- (ii) For 96.2% of these maps, i.e., 481 out of 500, there exists a positive integer $m \leq 2000$ such that $O_m(P) \cap V(\mathbb{Z}/m\mathbb{Z}) = \emptyset$.
- (iii) For 11 of the 19 remaining maps, we conclude that $V \cap \mathcal{O}_{\varphi}(P) = \emptyset$ by finding an integer 2000 < m < 11,500 such that $O_m(P) \cap V(\mathbb{Z}/m\mathbb{Z}) = \emptyset$. In each of these cases, the smallest such m is supported at more than 1 prime.
- (iv) For the remaining 8 maps, $O_m(P) \cap V(\mathbb{Z}/m\mathbb{Z}) \neq \emptyset$ for all m < 11,500. However, we are still able to conclude that $V \cap \mathcal{O}_{\varphi}(P) = \emptyset$ by showing that the image of $V(\mathbb{Z})$ modulo 7 is disjoint from $O_7(P)$.

All computations were performed using Magma (Bosma et al., 1997).

4. A DYNAMICAL HASSE PRINCIPLE FOR ÉTALE MORPHISMS

4.1. Notation. Let K be a global field, and let M_K denote its set of places. For a finite set S of places of K containing all the archimedean places, we write $\mathcal{O}_{K,S}$ to denote the ring of S-integers. For all $v \in M_K$, we use K_v to denote the v-adic completion. If v is nonarchimedean, we write \mathcal{O}_v , \mathfrak{m}_v , and k_v for the valuation ring, maximal ideal, and residue field of v, respectively. Let X denote a K-variety, i.e., a reduced separated scheme of finite type over K, let $V \subseteq X$ denote a closed K-subvariety, and let $\varphi \colon X \to X$ denote a K-endomorphism. For any K-variety Y, define

$$Y(K,S) := \prod_{v \notin S} Y(K_v).$$
(4.1)

We equip $Y(K_v)$ with the v-adic topology and Y(K, S) with the product topology. We view Y(K) as a subset of Y(K, S) via the diagonal embedding. For every subset T of Y(K, S) or $Y(K_v)$, we write $\mathcal{C}(T)$ or $\mathcal{C}_v(T)$ for the closure of T in the product topology or v-adic topology, respectively.

Since Y is separated, $Y(K_v)$, and hence Y(K, S), is Hausdorff. We note that as Y is not assumed to be projective, Y(K, S) need not agree with the adelic points of Y. For basic terminologies and properties of scheme theory, we refer the readers to (Hartshorne, 1977), for properties of smooth and étale morphisms used throughout this section, we refer the readers to (Grothendieck, 1964).

4.2. The dynamical Hasse principle for étale maps and preperiodic subvarieties. For any point $P \in X(K)$, we have the following containments:

$$V(K) \cap \mathcal{O}_{\varphi}(P) \subseteq V(K,S) \cap \mathcal{C}(\mathcal{O}_{\varphi}(P)) \subseteq \prod_{v \notin S} V(K_v) \cap \mathcal{C}_v(\mathcal{O}_{\varphi}(P)).$$

Recall the definition by Hsia-Silverman (Hsia and Silverman, 2009, p.237–238) that (X, V, φ) is said to be dynamical Brauer-Manin S-unobstructed if the leftmost containment is an equality for every $P \in X(K)$ satisfying $\mathcal{O}_{\varphi}(P) \cap V^{\text{pp}} = \emptyset$, where V^{pp} is the union of all positive dimensional preperiodic subvarieties of V. In analogy, we define the dynamical Hasse principle:

Definition 4.1. The triple (X, V, φ) is said to satisfy the dynamical Hasse principle (over K) if for every $P \in X(K)$ such that $\mathcal{O}_{\varphi}(P) \cap V(K) = \emptyset$, there exists a place v (depending on P) such that $\mathcal{C}_{v}(\mathcal{O}_{\varphi}(P)) \cap V(K_{v}) = \emptyset$. If there are infinitely many such places v, we say that (X, V, φ) satisfies the strong dynamical Hasse principle.

When $V = V^{\text{pp}}$, if (X, V, φ) satisfies the strong dynamical Hasse principle then it is immediate that (X, V, S) is Brauer-Manin S-unobstructed for every S. The reason is that for every $P \in X(K)$ such that $\mathcal{O}_{\varphi}(P) \cap V(K) = \emptyset$, both containments above are equalities since all the three sets are the empty set. Our main results in this section are the following:

Theorem 4.2. Assume that X is quasi-projective, that φ is étale, and that V is φ^m -invariant, i.e., that $\varphi^m(V) \subseteq V$, for some $m \in \mathbb{Z}_{>0}$. If $P \in X(K)$ is such that $V(K) \cap \mathcal{O}_{\varphi}(P) = \emptyset$ then, for all but finitely many primes v,

$$V(K_v) \cap \mathcal{C}_v(\mathcal{O}_\varphi(P)) = \emptyset$$

Consequently, (X, V, φ) satisfies the strong dynamical Hasse principle.

We obtain a similar result when V is φ -preperiodic, under the mild additional assumption that φ is closed:

Theorem 4.3. Assume that X is quasi-projective and that φ is étale and closed. Let V be a preperiodic subvariety of X, which means that there exist $k \ge 0$ and m > 0 such that

 $\varphi^{k+m}(V) \subset \varphi^k(V)$. For every $P \in X(K)$, if $V(K) \cap \mathcal{O}_{\varphi}(P) = \emptyset$ then, for all but finitely many primes v,

$$V(K_v) \cap \mathcal{C}_v(\mathcal{O}_\varphi(P)) = \emptyset$$

Consequently, (X, V, φ) satisfies the strong dynamical Hasse principle.

Notice that in the theorems, we do not suppose that V is irreducible. But it is clear that the general case reduces to this. Indeed, for instance in 4.3, let C be an irreducible component of V such that the closure of the orbit of P meets C. Though the component C does not have to be preperiodic in general, C is preperiodic when it is of maximal dimension (thanks to the étaleness of φ). Moreover, if C is not preperiodic, then some power of φ sends it to a component of V of greater dimension. Thus, if the closure of the orbit of P meets C, it necessarily meets some preperiodic component of V, so we may replace V by this component.

In §4.3 we prove a local version of Theorem 4.2. Next in §4.4, we show how Theorem 4.2 follows from the local version, Theorem 4.4. In §4.5 we prove Theorem 4.3 and in §4.6 we give some closing remarks.

4.3. The local statement. Throughout this section, we work locally. Let A denote a complete discrete valuation ring, \mathfrak{m} its maximal ideal, and k its residue field; we will assume that k is perfect. We write F for the fraction field of A.

The goal of this section is to prove the following:

Theorem 4.4. Let \mathscr{X} be a smooth quasi-projective scheme over A, let φ be an étale endomorphism of \mathscr{X} , and let \mathscr{V} be a reduced and closed subscheme of \mathscr{X} such that $\varphi^{M}(\mathscr{V}) \subseteq \mathscr{V}$ for some $M \geq 1$. Let $P \in \mathscr{X}(A)$ be such that the reduction \overline{P} of P modulo \mathfrak{m} is preperiodic under φ . If $\mathscr{V}(A) \cap \mathcal{O}_{\varphi}(P) = \emptyset$, then $\mathscr{V}(A)$ does not intersect the \mathfrak{m} -adic closure of $\mathcal{O}_{\varphi}(P)$.

We beging with a few preliminary lemmas.

Remark 4.5. Lemmas 4.6–4.9 below are similar to results in (Bell et al., 2010, Sec. 2) which were proved for $A = \mathbb{Z}_p$.

Lemma 4.6. Let \mathscr{X} be a separated A-scheme of finite type. Then we have the following.

- (a) $\mathscr{X}(A)$ is an open and closed subset of $\mathscr{X}(F)$, and it is compact if k is finite.
- (b) For each $x \in \mathscr{X}(A)$, and each positive integer n,

 $C(x,n) := \{ x' \in \mathscr{X}(A) : x' = x \text{ in } \mathscr{X}(A/\mathfrak{m}^n) \}$

is open and closed and the collection $\{C(x,n): n > 0\}$ forms a basis of neighborhoods of x. Furthermore, for any $n \in \mathbb{Z}_{>0}$, the set $\mathscr{X}(A)$ is a disjoint union of subsets of the form C(x,n); if k is finite, then $\mathscr{X}(A)$ is a disjoint union of finitely many such subsets.

Proof. Note that the \mathfrak{m} -adic topology on $\mathscr{X}(F)$ is defined as follows. Let $\{U_{\alpha}\}$ be an affine open cover of \mathscr{X} ; assume that each U_{α} is of the form $\operatorname{Spec}(A[T_1, \ldots, T_m]/I_{\alpha})$. Then $U_{\alpha}(F)$ is identified with the zero locus of I_{α} in F^m , and we equip it with the usual \mathfrak{m} -adic topology. This topology is independent of the choice of an affine covering and independent of the choice of presentation $\operatorname{Spec}(A[T_1, \ldots, T_m]/I_{\alpha})$ of an open in the covering. Since, by assumption, \mathscr{X} is of finite type, we assume that the covering is finite. All assertions in the lemma follow immediately from the above description of the \mathfrak{m} -adic topology on $\mathscr{X}(F)$. \Box Now we will prove a series of lemmas that give an explicit presentation of the local ring at points of \mathscr{X} and \mathscr{V} in terms of power series rings with coefficients in A.

Lemma 4.7. Let \mathscr{X} be a smooth quasi-projective scheme over A, let P be an A-point of \mathscr{X} , and let \overline{P} denote its reduction in $\mathscr{X}(k)$. Then the $\mathfrak{m}_{\overline{P}}$ -adic completion $\hat{\mathcal{O}}_{\mathscr{X},\overline{P}}$ is isomorphic to $A[[T_1,\ldots,T_g]]$, where g is the relative dimension of \mathscr{X} at P over A. Moreover, a choice of an isomorphism

$$\tau \colon \hat{\mathcal{O}}_{\mathscr{X},\bar{P}} \xrightarrow{\sim} A[[T_1,\ldots,T_g]]$$

gives the corresponding homeomorphism

$$h_{\tau}: \ C(P,1) \cong \mathfrak{m}^g. \tag{4.2}$$

Proof. If F is characteristic 0 and k is characteristic p, then the ring $\hat{\mathcal{O}}_{\mathscr{X},\bar{P}}$ is finite (as a module) over $\mathcal{W}(k)[[T_1,\ldots,T_g]]$ where $\mathcal{W}(k)$ is the ring of Witt vectors over k (Matsumura, 1986, Ch. 10). Since the point P gives rise to a section of the structural morphism:

$$A \longrightarrow \hat{\mathcal{O}}_{\mathscr{X},\bar{P}}$$

we have $\hat{\mathcal{O}}_{\mathscr{X},\bar{P}}$ is isomorphic to $A[[T_1,\ldots,T_g]]$.

If F and k have equal characteristic, we have $\hat{\mathcal{O}}_{\mathscr{X},\bar{P}}$ is isomorphic to $k[[T_1,\ldots,T_{g+1}]]$, and A is isomorphic to k[[T]]. Again, the point P makes sure that the image of T in $\hat{\mathcal{O}}_{\mathscr{X},\bar{P}}$ lies in $\hat{\mathfrak{m}} - \hat{\mathfrak{m}}^2$ where $\hat{\mathfrak{m}}$ is the maximal ideal of $\hat{\mathcal{O}}_{\mathscr{X},\bar{P}}$. Thus the structural morphism makes $\hat{\mathcal{O}}_{\mathscr{X},\bar{P}}$ a power series in g variables with coefficients in A.

Every point u in C(P, 1) is equivalent to an A-morphism:

$$\tilde{u}: \operatorname{Spec} A \to \operatorname{Spec} \mathcal{O}_{\mathscr{X},\bar{P}},$$

which is in turn equivalent to specifying an A-algebra homomorphism:

$$\tilde{u}^{\sharp}: \mathcal{O}_{\mathscr{X},\bar{P}} \to A.$$

Thus, for a fixed isomorphism $\tau : \hat{\mathcal{O}}_{\mathscr{X},\bar{P}} \xrightarrow{\sim} A[[T_1,\ldots,T_g]]$, we have that giving \tilde{u}^{\sharp} is equivalent to giving an element:

$$h_{\tau}(u) := (\tilde{u}^{\sharp}(T_i))_{i=1}^g \in \mathfrak{m}^g.$$

Lemma 4.8. Retain the notation from Lemma 4.7. Assume that \mathscr{V} is a closed subscheme of \mathscr{X} such that $\bar{P} \in \mathscr{V}(k)$; write \mathscr{I} for the ideal sheaf of \mathscr{V} . After fixing an isomorphism $\tau: \hat{\mathcal{O}}_{\mathscr{X},\bar{P}} \xrightarrow{\sim} A[[T_1,\ldots,T_g]]$ (which exists by Lemma 4.7), we may view $\mathscr{I}_{\bar{P}}\hat{\mathcal{O}}_{\mathscr{X},\bar{P}}$ as an ideal of $A[[T_1,\ldots,T_g]]$. Then the homeomorphism h_{τ} in (4.2) maps $\mathscr{V}(A) \cap C(P,1)$ onto

$$\{\beta \in \mathfrak{m}^g : f(\beta) = 0 \ \forall f \in \mathscr{I}_{\bar{P}}\mathcal{O}_{\mathscr{X},\bar{P}}\}.$$

Furthermore, the $\mathfrak{m}_{\bar{P}}$ -adic completion

$$\hat{\mathcal{O}}_{\mathscr{V},\bar{P}} \cong \hat{\mathcal{O}}_{\mathscr{X},\bar{P}} / \mathscr{I}_{\bar{P}} \hat{\mathcal{O}}_{\mathscr{X},\bar{P}}$$

is reduced if \mathscr{V} is reduced at \overline{P} .

Proof. The first assertion is immediate. The fact that $\hat{\mathcal{O}}_{\mathscr{V},\bar{P}} \cong \hat{\mathcal{O}}_{\mathscr{X},\bar{P}}/\mathscr{I}_{\bar{P}}\hat{\mathcal{O}}_{\mathscr{X},\bar{P}}$ is a wellknown property: taking completion preserves exact sequences of finitely generated modules over a Noetherian ring (see, for example (Atiyah and Macdonald, 1969, p. 108)). Since the local ring $\mathcal{O}_{\mathscr{V},\bar{P}}$ is reduced and excellent, its completion is reduced (Matsumura, 1986, Ch. 11).

Lemma 4.9. Retain the notation from Lemma 4.7 and fix an isomorphism $\tau: \hat{\mathcal{O}}_{\mathscr{X},\bar{P}} \xrightarrow{\sim} A[[T_1,\ldots,T_g]]$. Assume that φ is an endomorphism of \mathscr{X} such that $\overline{\varphi(P)} = \bar{P}$. Then there are power series f_1,\ldots,f_g in $A[[T_1,\ldots,T_g]]$ such that $f = (f_1,\ldots,f_g)$ fits into the following commutative diagram:

$$C(P,1) \xrightarrow{\varphi} C(P,1) \tag{4.3}$$

$$\downarrow^{h_{\tau}} \qquad \qquad \downarrow^{h_{\tau}} \qquad \qquad \downarrow^{h_{\tau}}$$

$$\mathfrak{m}^{g} \xrightarrow{f} \qquad \mathfrak{m}^{g}$$

Proof. Using τ to identify $\hat{\mathcal{O}}_{\mathscr{X},\bar{P}}$ with $A[[T_1,\ldots,T_g]]$, we define:

$$f_i = \widehat{\varphi^*} T_i \quad \forall 1 \le i \le g,$$

where $\widehat{\varphi^*}$: $\hat{\mathcal{O}}_{\mathscr{X},\bar{P}} \to \hat{\mathcal{O}}_{\mathscr{X},\bar{P}}$. Then the commutativity is immediate.

We will also use the following easy lemma:

Lemma 4.10. Let $R = A[[T_1, \ldots, T_g]]$, and $\varphi \colon R \to R$ a ring isomorphism. If I is a radical ideal of R such that $\varphi(I) \subseteq I$, then $\varphi(I) = I$.

Proof. Although the lemma is stated for a particular ring R, it will be clear from the proof that we require only that R is a Noetherian equidimensional catenary ring (see (Eisenbud, 1995, pp. 452, 453) for the definition). Let P_1, \ldots, P_n be the minimal primes containing I; then $I = \bigcap_{j=1}^{n} P_j$. Let $D = \max_{1 \le j \le n} \dim R/P_j$. We will prove that for any integer d, φ fixes the set

$$\mathcal{S}_d := \{ P_j : \dim R / P_j = d \};$$

the proof uses backwards induction on $0 \le d \le D$. From this statement, the lemma follows immediately.

Since, for each $1 \le j \le n$

$$\bigcap_{i=1}^{n} \varphi(P_i) = \varphi(I) \subseteq I \subseteq P_j,$$

there exists some *i* (depending on *j*) such that $\varphi(P_i) \subseteq P_j$. Let *P* be an arbitrary prime in \mathcal{S}_D . Then there is some P_i such that $\varphi(P_i) \subseteq P$. We have:

$$D = \dim R/P \le \dim R/\varphi(P_i) = \dim R/P_i.$$

Since D is the maximal dimension, we must have equality and so $\varphi(P_i) = P$. Then a counting argument shows that φ fixes S_D .

Now fix $1 \leq D' < D$ and assume that φ fixes S_d for all $D' < d \leq D$. Let P be an arbitrary prime in $S_{D'}$, and let $1 \leq i \leq n$ be such that $\varphi(P_i) \subseteq P$. If $d := \dim R/P_i$ is strictly greater

that D', then, by the induction hypothesis, $\varphi(P_i) = P_j$ for some j. Since P_j and P are two distinct minimal primes containing I and $P_j \subseteq P$, this yields a contradiction. Thus, we have

$$D' = \dim R/P \le \dim R/\varphi(P_i) = \dim R/P_i = d \le D'_i$$

and so $\varphi(P_i) = P$. Again, a counting argument shows that φ fixes $S_{D'}$, as desired.

Now we are equipped to prove Theorem 4.4.

Proof of Theorem 4.4. Assume that $\mathscr{V} \cap \mathcal{O}_{\varphi}(P) = \emptyset$. We first perform a few reductions. By assumption, $\varphi^{M}(\mathscr{V}) \subseteq \mathscr{V}$ for some positive integer M; after replacing the data (φ, P) with $(\varphi^{M}, \varphi^{i}(P))$ for each $0 \leq i < M$, we may assume that M = 1. We have also assumed that P is φ -preperiodic modulo \mathfrak{m} , i.e., that $\varphi^{N_{0}+N}(P) \equiv \varphi^{N_{0}}(P) \pmod{\mathfrak{m}}$. Since $\mathscr{V} \cap \mathcal{O}_{\varphi}(P) = \emptyset$, the \mathfrak{m} -adic closure of $\mathcal{O}_{\varphi}(P)$ intersects \mathscr{V} if and only if \mathfrak{m} -adic closure of $\mathcal{O}_{\varphi}(\varphi^{N_{0}}(P))$ intersects \mathscr{V} if and only if \mathfrak{m} -adic closure of $\mathcal{O}_{\varphi}(\varphi^{N_{0}}(P))$ intersects \mathscr{V} . Therefore, by replacing P with $\varphi^{N_{0}}(P)$, we may assume that \overline{P} is φ -periodic of period N. For each $0 \leq i \leq N-1$, by replacing the data (φ, P) with the data $(\varphi^{N}, \varphi^{i}(P))$, we may assume that φ fixes \overline{P} . By Lemma 4.7, we may identify $\hat{\mathcal{O}}_{\mathscr{X},\overline{P}}$ and C(P,1) with $A[[T_{1},\ldots,T_{g}]]$ and \mathfrak{m}^{g} , respectively. By Lemma 4.9, there exist power series $f = (f_{1},\ldots,f_{g})$ such that φ acts by f in the chart \mathfrak{m}^{g} .

If $\bar{P} \notin \mathscr{V}(k)$, then C(P) := C(P, 1) contains the **m**-adic closure of $\mathcal{O}_{\varphi}(P)$ and is disjoint from $\mathscr{V}(A)$, so we are done. Now suppose that $\bar{P} \in \mathscr{V}(k)$. As in Lemma 4.8, let \mathscr{I} denote the ideal sheaf of \mathscr{V} . Let H_1, \ldots, H_m denote a choice of generators of the ideal $\mathscr{I}_{\bar{P}} \hat{\mathcal{O}}_{\mathscr{X},\bar{P}}$ of $A[[T_1, \ldots, T_g]]$. Because $\varphi(\mathscr{V}) \subseteq \mathscr{V}$, we have $\varphi^*(\mathscr{I}_{\bar{P}}) \subseteq \mathscr{I}_{\bar{P}}$ where $\varphi^* : \mathcal{O}_{\mathscr{X},\bar{P}} \to \mathcal{O}_{\mathscr{X},\bar{P}}$ is the induced map on the stalk at \bar{P} . Therefore $\widehat{\varphi^*}(\mathscr{I}_{\bar{P}}) \hat{\mathcal{O}}_{\mathscr{X},\bar{P}} \subseteq \mathscr{I}_{\bar{P}} \hat{\mathcal{O}}_{\mathscr{X},\bar{P}}$. By Lemma 4.8, $\mathscr{I}_{\bar{P}} \hat{\mathcal{O}}_{\mathscr{X},\bar{P}}$ is a radical ideal of $\hat{\mathcal{O}}_{\mathscr{X},\bar{P}}$, and, since φ is étale, $\widehat{\varphi^*} : \hat{\mathcal{O}}_{\mathscr{X},\bar{P}} \to \hat{\mathcal{O}}_{\mathscr{X},\bar{P}}$ is a ring isomorphism. Thus, we may apply Lemma 4.10 to conclude that $\widehat{\varphi^*}(\mathscr{I}_{\bar{P}}) \hat{\mathcal{O}}_{\mathscr{X},\bar{P}} = \mathscr{I}_{\bar{P}} \hat{\mathcal{O}}_{\mathscr{X},\bar{P}}$. Therefore, there exist power series $c_{ij} \in A[[T_1, \ldots, T_g]]$, where $1 \leq i, j \leq m$, such that

$$H_{i} = \sum_{j=1}^{m} c_{ij} H_{j}(f_{1}, \dots, f_{g}) \quad \forall 1 \le i \le m.$$
(4.4)

For a vector $v = (v_i)$ in $(F)^m$, define:

$$||v|| = \max_{1 \le i \le m} |v_i|,$$

and for $u \in \mathfrak{m}^{g}$, write $H(u) = (H_{i}(u)) \in A^{m}$. By (4.4), we now have:

$$||H(f(u))|| \ge ||H(u)|| \quad \forall u = (u_i) \in \mathfrak{m}^g.$$

$$(4.5)$$

Thus, the closure of $\mathcal{O}_{\varphi}(P)$ is contained in

$$\{u\in\mathfrak{m}^g:\ ||H(u)||\geq ||H(P)||\}$$

Since ||H(P)|| > 0 and H vanishes on $\mathscr{V}(A)$, this completes the proof.

Remark 4.11. If $\varphi(\mathscr{V}) \subseteq \mathscr{V}$, then we can prove a stronger statement than (4.5), namely:

$$||H(F(u))|| = ||H(u)||.$$
(4.6)

The proof proceeds in the same way as the proof of (4.5). More explicitly, we may write each $H_i(f_1, \ldots, f_g)$ as a linear combination of the H_j 's and obtain $||H(f(u))|| \leq ||H(u)||$.

Morally speaking, if we consider ||H(P)|| as the "distance" from P to \mathscr{V} then equality (4.6) means that when P comes close to \mathscr{V} (i.e. $\bar{P} \in \mathscr{V}(k)$) and φ fixes P modulo \mathfrak{m} , then all

elements in $\mathcal{O}_{\varphi}(P)$ are equidistant to \mathscr{V} . Thus P becomes a "satellite" of \mathscr{V} by moving around \mathscr{V} under φ .

If F is a finite extension of \mathbb{Q}_p for some prime p, then there is another proof of Theorem 4.4 using a uniformization of $\mathcal{O}_{\omega}(P)$ defined as follows.

Definition 4.12. Retain the notation from Theorem 4.4. Assume that F is a finite extension of \mathbb{Q}_p and that φ fixes P modulo \mathfrak{m} . Using Lemma 4.7, identify $\hat{\mathcal{O}}_{\mathscr{X},\bar{P}}$ with $A[[T_1,\ldots,T_g]]$, and C(P) := C(P,1) with \mathfrak{m}^g . The orbit $\mathcal{O}_{\varphi}(P)$ has a uniformization if there exist power series G_1,\ldots,G_g in F[[T]], convergent on \mathbb{Z}_p such that:

- (i) $(G_1(0), \ldots, G_q(0)) = P$, which equals 0 in \mathfrak{m}^g , and
- (ii) $\varphi(G_1(z), \dots, G_g(z)) = (G_1(z+1), \dots, G_g(z+1)) \ \forall z \in \mathbb{Z}_p.$

Proof of Theorem 4.4 using uniformization. This proof is similar to arguments developed in (Bell et al., 2010).

As above, we reduce to the case that M = 1 (i.e. \mathscr{V} is invariant) and assume that $\overline{P} \in \mathscr{V}(k)$. By (Bell et al., 2010, Theorem 3.3) or (Amerik, 2011, Theorem 7), there is a uniformization of $\mathscr{O}_{\varphi}(P)$ (possibly after replacing φ by an iterate). Let $G = (G_1, \ldots, G_g)$ be such a uniformization. By Definition 4.12, the m-adic closure of $\mathcal{O}_{\varphi}(P)$ is contained in $G(\mathbb{Z}_p)$. If there is some $u \in \mathbb{Z}_p$ such that $G(u) \in \mathscr{V}(A)$, then $G(u+n) \in \mathscr{V}(A)$ for every natural number n. Let H = 0 be any of the equations defining \mathscr{V} in \mathscr{X} . Then we have $H \circ G(u+n) = 0$ for every natural number n. Since a nonzero p-adic analytic function on \mathbb{Z}_p can have only finitely many zeros, the analytic function $H \circ G$ must be identically zero on \mathbb{Z}_p . Therefore $G(\mathbb{Z}_p) \subseteq \mathscr{V}(A)$ and so the whole orbit of P is contained in $\mathscr{V}(A)$, contradicting our assumption that $\mathcal{O}_{\varphi}(P) \cap \mathscr{V}(A) = \emptyset$.

While the argument using uniformization is simpler and might be applicable to some cases where φ is not étale, our approach still has a number of advantages.

- The proof of Theorem 4.4 that does not require on uniformization can be translated into a simple and effective algorithm, hence is suitable for computational purposes.
- Lemma 4.10 and results of the same type may be of independent interest. Indeed, results of a similar spirit are studied in a recent preprint of Bell and Lagarias (Bell and Lagarias).
- Our proof requires no assumption on the characteristic of F, whereas uniformization of orbits does not hold if char(F) > 0 (for example, it is impossible to have an "exponential function" since dividing by n! is prohibited).

4.4. **Proof of Theorem 4.2.** In this section, we present the proof of Theorem 4.2. First we show that for all but finitely many places v, the assumptions of Theorem 4.4 hold.

Lemma 4.13. Assume that φ is étale and that $\varphi(V) \subseteq V$. Fix a point $P \in X(K)$, then there exists a finite set $S \subseteq M_K$ containing all the archimedean places such that X, V, φ and P extend to models $\mathscr{X}, \mathscr{V}, \tilde{\varphi}$ and \mathscr{P} , respectively, over $\mathcal{O}_{K,S}$ with the following properties:

- $\tilde{\varphi}$ is étale,
- \mathscr{X} is quasi-projective and smooth, and
- \mathscr{V} is a reduced, closed subscheme of \mathscr{X} , is flat over $\mathcal{O}_{K,S}$, and $\tilde{\varphi}(\mathscr{V}) \subseteq \mathscr{V}$.

Proof. We first find models for X, V, φ and P over, say, some $\mathcal{O}_{K,S}$. As the locus in $\operatorname{Spec}(\mathcal{O}_{K,S})$ over which \mathscr{X} is not smooth, \mathscr{V} is not flat, or $\tilde{\varphi}$ is not étale is closed, by

enlarging S, we may assume that this locus is empty. Since \mathscr{V} is flat over $\mathcal{O}_{K,S}$ and its generic fiber is reduced, \mathscr{V} is itself reduced.

It remains to ensure that $\tilde{\varphi}(\mathscr{V}) \subseteq \mathscr{V}$. By enlarging S again, we may assume that every irreducible component of \mathscr{V} contains some point in the generic fiber. Then since $\varphi(V) \subseteq V$ and V is dense in \mathscr{V} , we have $\tilde{\varphi}(\mathscr{V}) \subseteq \mathscr{V}$.

Remark 4.14. The proof of Theorem 4.2 is simpler in the case when X is smooth. For clarity, we present the proof under this additional assumption first, and then give the general case.

Proof of Theorem 4.2 under the assumption that X is smooth. Fix $S \subseteq M_K$ as in Lemma 4.13. For $v \notin S$, let $\mathcal{O}_{(v)} = K \cap \mathcal{O}_v$ be the valuation ring of v in K. By Lemma 4.13, there exist models \mathscr{X} and \mathscr{V} of X and V, respectively, over $\mathcal{O}_{(v)}$. We take the formally smooth base change from $\mathcal{O}_{(v)}$ to \mathcal{O}_v , and abuse notation by (still) using \mathscr{V} and \mathscr{X} to denote the resulting models over \mathcal{O}_v .

Now we apply Theorem 4.4 to show that $\mathscr{V}(\mathcal{O}_v)$ does not intersect the closure of $\mathcal{O}_{\varphi}(P)$ in $\mathscr{X}(\mathcal{O}_v)$. Since $\mathscr{X}(\mathcal{O}_v)$ is closed in $\mathscr{X}(K_v) = X(K_v)$, $V(K_v) = \mathscr{V}(K_v)$ does not intersect the closure of $\mathcal{O}_{\varphi}(P)$ in $X(K_v)$. This completes the proof in the case when X is smooth. \Box

Proof of Theorem 4.2 in the general case. Let M > 0 be such that $\varphi^M(V) \subseteq V$. As in the proof of Theorem 4.4, by replacing the data (φ, P) with the data $(\varphi^M, \varphi^i(P))$ for $0 \leq i < M - 1$, we may assume that M = 1. We prove the theorem by Noetherian induction.

If $\dim(X) = 0$, there is nothing to prove. Assume that $\dim(X) > 0$; let X' be the smooth locus of X, and let X'' = X - X' with the reduced closed subscheme structure. Since φ is étale, φ preserves both X' and X''. If $P \in X''(K)$ then we use the induction hypothesis on X''.

Now suppose that $P \in X'(K)$. Write $V' = V \cap X'$. By Lemma 4.13, there exists a finite set of places S such that X and V, respectively, have flat (hence reduced) models \mathscr{X} and \mathscr{V} over $\mathcal{O}_{K,S}$. In addition, φ and P both extend to $\mathcal{O}_{K,S}$ (by an abuse of notation, we denote these extensions by φ and P, respectively), and $\varphi(\mathscr{V}) \subseteq \mathscr{V}$. Since φ is étale at P generically, by enlarging S, we may assume that φ is étale at every point in P.

Let \mathscr{X}' denote the open subscheme of \mathscr{X} consisting of smooth points over $\mathcal{O}_{K,S}$, and let $\mathscr{V}' = \mathscr{V} \cap \mathscr{X}'$ be a model of V' over $\mathcal{O}_{K,S}$. Now let v be a place outside S and pull back all models to \mathcal{O}_v . By Theorem 4.4, $\mathscr{V}'(\mathcal{O}_v)$ does not intersect the v-adic closure of $\mathcal{O}_{\varphi}(P)$ in $\mathscr{X}'(\mathcal{O}_v)$. Since, by Lemma 4.6, $\mathscr{X}'(\mathcal{O}_v)$ is quasi-compact and $X(K_v)$ is Hausdorff, the set $\mathscr{X}'(\mathcal{O}_v)$ is closed in $X(K_v)$. Therefore $V(K_v)$ does not intersect the v-adic closure of $\mathcal{O}_{\varphi}(P)$ in $X(K_v)$. This completes the proof.

4.5. **Proof of Theorem 4.3.** We proceed by induction on the dimension of V. The case $\dim(V) = 0$ is easy, as follows. Let U be the union of all iterates of V. If $\mathcal{O}_{\varphi}(P)$ intersects U then P is preperiodic and there is nothing to prove. Otherwise, we apply Theorem 4.2 for U instead of V. To carry out the induction step, we need the following lemma which might be of independent interest.

Lemma 4.15. Let X and φ be as in Theorem 4.3. Let Y be an irreducible φ -preperiodic subvariety of X. Let Y_1 be a periodic iterate of Y. Then every irreducible component of $Y \cap Y_1$ is preperiodic.

Proof. Replacing φ by an iterate, we reduce to the case $Y_1 = \varphi(Y) = \varphi^2(Y) \neq Y$. Let $\nu_1: \tilde{Y}_1 \to Y_1$ and $\nu: \varphi^{-1}(Y_1) \to \varphi^{-1}(Y_1)$ denote the normalizations of Y_1 and $\varphi^{-1}(Y_1)$, respectively. For every integer $n \geq 1$, define:

$$Z_n := \left\{ x \in Y_1 : |\nu_1^{-1}(x)| \ge n \right\}$$

where $|\nu_1^{-1}(x)|$ is counted with multiplicity. By the semicontinuity theorem, Z_n is closed in Y_1 .

Now let W be an irreducible component of $Y \cap Y_1$ having dimensional d and generic point w. Let $s = |\nu^{-1}(\varphi(w))|$, the main observation is that $\varphi(W)$ is an irreducible component of Z_s . Assume otherwise and let C_1 be an irreducible component of Z_s strictly containing $\varphi(W)$. Let c_1 denote the generic point of C_1 . Since $c_1 \in Z_s$, we have $|\nu_1^{-1}(c_1)| \ge s$. On the other hand, by semicontinuity $|\nu_1^{-1}(c_1)| \le |\nu_1^{-1}(\varphi(w))| = s$. Hence we must have $|\nu_1^{-1}(c_1)| = s$.

By faithful flatness, the map

$$\operatorname{Spec}(\mathcal{O}_{V_1},\varphi(w))\longrightarrow \operatorname{Spec}(\mathcal{O}_{\varphi^{-1}(V_1)},w)$$

induced by φ is surjective (Atiyah and Macdonald, 1969, p.68). Therefore, there exists c in $\varphi^{-1}(V_1)$ such that $\varphi(c) = c_1$ and the Zariski closure C of c contains W. From the last paragraph and the fact that taking normalization commutes with étale base changes, we have:

$$|\nu^{-1}(w)| = |\nu_1^{-1}(\varphi(w))| = s = |\nu_1^{-1}(c_1)| = |\nu^{-1}(c)|$$
(4.7)

By comparing dimensions, we see that V and V_1 are two irreducible components of $\varphi^{-1}(V_1)$ containing w. Since C_1 strictly contains $\varphi(W)$, we have that C strictly contains W. Note that it is impossible for both V and V_1 to contain C since W is an irreducible component of $V \cap V_1$. Therefore the set of irreducible components of $\varphi^{-1}(V_1)$ containing c is strictly smaller than the set of irreducible components of $\varphi^{-1}(V)$ containing w. Together with the semicontinuity theorem, we have $|\nu^{-1}(c)| < |\nu^{-1}(w)|$ contradicting (4.7).

Now let \mathcal{Z} denote the union of all *d*-dimensional irreducible components of all the Z_n 's for $n \geq 1$. This is a finite union since $Z_n = \emptyset$ for all sufficiently large *n* thanks to finiteness of ν_1 . We have proved that $\varphi(W)$ is an irreducible component of \mathcal{Z} . For each $m \geq 1$, repeat the same arguments for φ^m instead of φ , we have that $\varphi^m(W)$ is an irreducible component of \mathcal{Z} . This proves that W is preperiodic.

We now return to the proof of Theorem 4.3. We may assume that V is irreducible and not periodic. Replacing φ by an iterate, we may assume $V_1 = \varphi(V) = \varphi^2(V) \neq V$.

If $\mathcal{O}_{\varphi}(P) \cap V_1(K) = \emptyset$ then by Theorem 4.2, there is a finite set of places S_1 such that $\mathcal{C}_v(\mathcal{O}_{\varphi}(P)) \cap V_1(K_v) = \emptyset$ for $v \notin S_1$. This implies $\mathcal{C}_v(\mathcal{O}_{\varphi}(P)) \cap V(K_v) = \emptyset$ for every $v \notin S_1$. Now assume that $\varphi^m(P) \in V_1(K)$ for some $m \ge 0$. For every v, we have

$$\mathcal{C}_v(\mathcal{O}_\varphi(\varphi^m(P))) \subseteq V_1(K_v)$$

and

$$\mathcal{C}_{v}(\mathcal{O}_{\varphi}(P)) = \mathcal{C}_{v}(\mathcal{O}_{\varphi}(\varphi^{m}(P))) \cup \{P, \dots, \varphi^{m-1}(P)\}$$

If
$$V(K_v) \cap \mathcal{C}_v(\mathcal{O}_{\varphi}(P)) \neq \emptyset$$
, then the above identities together with $V(K) \cap \mathcal{O}_{\varphi}(P) = \emptyset$ imply:

 $\mathcal{C}_v(\mathcal{O}_\varphi(P)) \cap (V \cap V_1)(K_v) \neq \emptyset.$

This can only happen for finitely many v's by applying the induction hypothesis for $V \cap V_1$, whose irreducible components are preperiodic by Lemma 4.15.

4.6. Closing remarks. It is natural to ask whether the assumption in Theorem 4.2 that V is preperiodic is necessary. We show that if K is a number field and V = Q is a single K-point, then (X, Q, φ) fails the strong Hasse principle if and only if Q is not periodic but has almost everywhere periodic reduction, i.e. for all but finitely many primes \mathfrak{p} , the reduction of Q modulo \mathfrak{p} is periodic.

Proposition 4.16. Let K be a number field, let $\varphi : X \to X$ be an étale K-endomorphism of X and let $Q \in X(K)$.

- (a) Let p be a prime such that we have models over O_p. If the reduction of Q modulo p is periodic then the p-adic closure of the orbit of φ(Q) contains Q. Consequently, if Q is not periodic but has almost everywhere periodic reduction then (X, Q, φ) does not satisfy the strong Hasse principle.
- (b) Conversely, if Q is either periodic or does not have almost everywhere periodic reduction, then (X, Q, φ) satisfies the strong dynamical Hasse principle.

Proof. (a) The first assertion follows immediately from the \mathfrak{p} -adic uniformization of the φ^{N} -orbit of Q (for some N >> 0 depending on \mathfrak{p}) and the fact that for every analytic function G from $\mathcal{O}_{\mathfrak{p}}$ to $\mathcal{O}_{\mathfrak{p}}^{g}$, the point G(0) lies in the closure of $\{G(1), G(2), \ldots\}$. Such uniformization exists by (Bell et al., 2010, Theorem 3.3), or more precisely by its generalization in (Amerik, 2011, Theorem 7). For the second assertion, note that the orbit of $P = \varphi(Q)$ does not contain Q but the \mathfrak{p} -adic closure of this orbit contains Q for almost all \mathfrak{p} .

(b) The case that Q is periodic follows from Theorem 4.2. Hence we assume that Q is non-periodic. Let $P \in X(K)$ such that $Q \notin \mathcal{O}_{\varphi}(P)$. For every \mathfrak{p} such that X, φ, P and Qhave models over $\mathcal{O}_{\mathfrak{p}}$, if $Q \in \mathcal{C}_{\mathfrak{p}}(\mathcal{O}_{\varphi}(P))$ then Q and $\varphi^m(P)$ have the same reduction modulo \mathfrak{p} for m as large as we like. This implies that Q is periodic modulo \mathfrak{p} . But there are infinitely many primes such that this conclusion does not hold thanks to our assumption on Q, hence (X, Q, φ) satisfies the strong dynamical Hasse principle. \Box

Some results in the literature suggest that the examples of non-periodic points with almost everywhere periodic reduction must be very special, and so the strong dynamical Hasse principle mostly holds when the endomorphism is étale and the subvariety is a single point. For instance, by a result of Pink (Pink, 2004, Corollary 4.3), such points cannot exist for the multiplication-by-d map on an abelian variety. Furthermore, by (Benedetto et al., 2013, Corollary 1.2), such points also cannot exist for a self-map of \mathbb{P}^1 of degree at least two (though such a map is not étale, and thus 4.16 does not apply directly).

On the other hand, the translation-by-one map on \mathbb{P}^1 is an obvious example where nonperiodic points become periodic modulo almost all primes, so that the strong dynamical Hasse principle fails; more generally, an automorphism of \mathbb{P}^n of infinite order, given by an integer-valued matrix, has the same property. As it was pointed out to us by Serge Cantat, one can use this to construct other, though somewhat artificial, examples: take an automorphism φ of a smooth variety X with a fixed point x, inducing an infinite-order automorphism on the tangent space at x. Then φ lifts to the blow-up of X at x, inducing an automorphism of the exceptional divisor, and points of that exceptional divisor are periodic modulo almost all primes.

It seems reasonable to conjecture that non-periodic points with almost everywhere periodic reduction do not exist for polarized morphisms φ (that is, morphisms such that $\varphi^* \mathcal{L} = \mathcal{L}^{\otimes k}$ for some integer k > 1 and some ample line bundle \mathcal{L}), so that the strong dynamical Hasse

principle holds for number fields K, étale polarized morphisms φ and $V \in X(K)$. Notice however that étale polarized endomorphisms are extremely rare, cf. (Fakhruddin, 2003).

For the sake of completeness, we note that for curves over number fields, the only counterexamples to the dynamical Brauer-Manin criterion are automorphisms φ of a very special kind.

Proposition 4.17. Let X be a smooth geometrically integral projective curve of genus g over a number field K, let φ be a nonconstant self-map of X over K, and V a finite subset of X(K). We make the following additional assumptions.

- (a) If g = 0, assume that φ is not conjugate to $z \mapsto z + 1$,
- (b) If g = 1, assume that φ has a preperiodic point in $X(\overline{K})$. (If we regard X as an elliptic curve, this condition is equivalent to the condition that φ is not a translate by a non-torsion point.)

Then we have the following equality:

$$V(K) \cap \mathcal{O}_{\varphi}(P) = V(K, S) \cap \mathcal{C}(\mathcal{O}_{\varphi}(P)).$$

Remark 4.18. If $X = \mathbb{P}^1$, V is a finite set of points and $\deg(\varphi) \geq 2$, then Silverman and Voloch have shown that $\mathcal{O}_{\varphi}(P) \cap V(K) = \mathcal{C}(\mathcal{O}_{\varphi}(P)) \cap V(K, S)$ (Silverman and Voloch, 2009, Theorem 1).

Remark 4.19. When X is an abelian variety, V is an arbitrary subvariety and φ is a K-endomorphism of X such that $\mathbb{Z}[\varphi]$ is an integral domain, Hsia and Silverman show the equality:

$$\mathcal{O}_{\varphi}(P) \cap V(K) = \mathcal{C}(\mathcal{O}_{\varphi}(P)) \cap V(K,S)$$

under certain strong conditions. We refer the readers to (Hsia and Silverman, 2009, Theorem 11) for more details. Our proof of Proposition 4.17 gives an unconditional proof of their result when X is an elliptic curve.

Proof. The case $g \ge 2$ is trivial since all endomorphisms of curves of genus at least two are of finite order. If g = 0, this follows from (Silverman and Voloch, 2009) as mentioned above (see Remark 9 from (Silverman and Voloch, 2009) if deg(φ) = 1).

Now consider the case when g = 1. If P is φ -preperiodic then there is nothing to prove, so we assume that P is wandering. There exists a non-negative integer N such that $\varphi^M(P) \notin V(K)$ for all M > N. After replacing P by $\varphi^{N+1}(P)$, we may assume the φ -orbit of P does not intersect V(K). It remains to show that $V(K, S) \cap \mathcal{C}(\mathcal{O}_{\varphi}(P)) = \emptyset$. By assumption (b), there is some M > 0 such that φ^M has a fixed point. By replacing the data (φ, P) with $(\varphi^M, \varphi^i(P))$ for $0 \leq i < M$, we may assume that φ has a fixed point.

Note that if we can prove

$$V(L, S_L) \cap \mathcal{C}(\mathcal{O}_{\varphi}(P)) = \emptyset$$

for L a finite extension of K, and $S_L \subset M_L$ the set of places of L lying above places in S, then this implies that $V(K, S) \cap \mathcal{C}(\mathcal{O}_{\varphi}(P)) = \emptyset$. Thus, we may assume that there is a fixed point $O \in X(K)$. Using O as the point at infinity we have the following Weierstrass equation for X:

$$y^2 = x^3 + Ax + B.$$

Let U(z)/V(z) be the Lattès map associated to φ . Since O is fixed by φ , we have deg $(V) < \deg(U) = \deg(\varphi)$. Let S' be a finite set of places such that $U, V \in \mathcal{O}_{K,S'}[z]$ and such that

the leading coefficients of U and V are in $\mathcal{O}_{K,S'}^{\times}$. Then

$$\forall v \notin S' \; \forall Q \in X(K), \quad \text{if } v(x(Q)) < 0, \text{ then } v(x(\varphi(Q))) < 0.$$

$$(4.8)$$

Assume that $V(K, S) \cap \mathcal{C}(\mathcal{O}_{\varphi}(P)) \neq \emptyset$ and let $a = (a_{\mathfrak{p}})_{\mathfrak{p}\notin S}$ be an element of the intersection. Note that since V is a finite set of points, $V(K_{\mathfrak{p}}) = V(K)$ and so $a_{\mathfrak{p}} \in V(K)$ for every $\mathfrak{p} \notin S$. By Theorem 4.3, for almost all \mathfrak{p} , the \mathfrak{p} -adic closure of $\mathcal{O}_{\varphi}(P)$ does not contain O. Therefore $a_{\mathfrak{p}} \neq O$ for almost all \mathfrak{p} . For such \mathfrak{p} , we can write $a_{\mathfrak{p}}$ with affine coordinates $a_{\mathfrak{p}} = (x_{\mathfrak{p}}, y_{\mathfrak{p}})$. We now enlarge S' so that:

- (i) $S \subseteq S'$.
- (ii) $a_{\mathfrak{p}} \neq O$ for $\mathfrak{p} \notin S'$.
- (iii) All points in $V(K) \{O\}$ have $\mathcal{O}_{K,S'}$ -integral affine coordinates. This means $x_{\mathfrak{p}}, y_{\mathfrak{p}} \in \mathcal{O}_{\mathfrak{p}}$ for $\mathfrak{p} \notin S'$.
- (iv) $A, B \in \mathcal{O}_{K,S'}$.

Since $a \in \mathcal{C}(\mathcal{O}_{\varphi}(P))$, properties (i), (iii) and (4.8) imply that $v(x(\varphi^n(P))) \geq 0$ for every n, and every $v \notin S'$. By property (iv) and the Weierstrass equation for X, we have $v(y(\varphi^n(P))) \geq 0$ for every n, and every $v \notin S'$. Therefore $\varphi^n(P)$ has $\mathcal{O}_{K,S'}$ -integral affine coordinates for every n. This contradicts Siegel's theorem asserting finiteness of integral points.

References

Amerik, E. 2011. Existence of non-preperiodic algebraic points for a rational self-map of infinite order, Math. Research Letters 18, 251-256. $\uparrow 4.3$, 4.6

Atiyah, M. F. and I. G. Macdonald. 1969. Introduction to commutative algebra, Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont. MR0242802 (39 #4129) ↑4.3, 4.5

Bell, J. P., D. Ghioca, and T. J. Tucker. 2010. *The dynamical Mordell-Lang problem for étale maps*, Amer. J. Math. **132**, no. 6, 1655–1675. MR2766180 (2012a:37202) ↑4.5, 4.3, 4.6

Benedetto, Robert L., Dragos Ghioca, Benjamin Hutz, Pär Kurlberg, Thomas Scanlon, and Thomas J. Tucker. 2013. *Periods of rational maps modulo primes*, Math. Ann. **355**, no. 2, 637–660, DOI 10.1007/s00208-012-0799-8. MR3010142 ↑4.6

Bell, J. and J. Lagarias. A Skolem-Mahler-Lech Theorem For Ideal Inclusions. preprint, 2012. ↑4.3

Bosma, Wieb, John Cannon, and Catherine Playoust. 1997. The Magma algebra system. I. The user language, J. Symbolic Comput. 24, no. 3-4, 235–265, DOI 10.1006/jsco.1996.0125. Computational algebra and number theory (London, 1993). MR1484478 \uparrow 3.2

Eisenbud, David. 1995. Commutative algebra, Graduate Texts in Mathematics, vol. 150, Springer-Verlag, New York. With a view toward algebraic geometry. MR1322960 (97a:13001) ↑4.3

Fakhruddin, Najmuddin. 2003. Questions on self maps of algebraic varieties, J. Ramanujan Math. Soc. 18, no. 2, 109–122. MR1995861 ↑4.6

Flajolet, Philippe and Andrew M. Odlyzko. 1990. *Random mapping statistics*, Advances in cryptology— EUROCRYPT '89 (Houthalen, 1989), Lecture Notes in Comput. Sci., vol. 434, Springer, Berlin, pp. 329–354. MR1083961 ↑2.1

Grothendieck, A. 1964. Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes des schémas. I, Inst. Hautes Études Sci. Publ. Math. 20, 259 pp. ↑4.1

Hartshorne, Robin. 1977. Algebraic geometry, Springer-Verlag, New York. Graduate Texts in Mathematics, No. 52. MR0463157 (57 #3116) $\uparrow 4.1$

Hsia, Liang-Chung and Joseph Silverman. 2009. On a dynamical Brauer-Manin obstruction, J. Théor. Nombres Bordeaux **21**, no. 1, 235–250 (English, with English and French summaries). MR2537714 (2010f:37157) \uparrow 1, 4.2, 4.19

Matsumura, Hideyuki. 1986. Commutative ring theory, Cambridge Studies in Advanced Mathematics, vol. 8, Cambridge University Press, Cambridge. Translated from the Japanese by M. Reid. MR879273 (88h:13001) \uparrow 4.3, 4.3

Pink, Richard. 2004. On the order of the reduction of a point on an abelian variety, Math. Ann. **330**, no. 2, 275–291, DOI 10.1007/s00208-004-0548-8. MR2089426 (2005g:11101) ↑4.6

Poonen, Bjorn. 2006. Heuristics for the Brauer-Manin obstruction for curves, Experiment. Math. 15, no. 4, 415–420. MR2293593 (2008d:11062) $\uparrow 1$

Scharaschkin, Victor. 1999. Local-global problems and the Brauer-Manin obstruction, ProQuest LLC, Ann Arbor, MI. Thesis (Ph.D.)–University of Michigan. MR2700328 ↑1

Silverman, Joseph H. and José Felipe Voloch. 2009. A local-global criterion for dynamics on \mathbb{P}^1 , Acta Arith. **137**, no. 3, 285–294, DOI 10.4064/aa137-3-8. MR2496466 (2010g:37173) \uparrow 4.18, 4.6

Stein, W. A. 2013. Sage Mathematics Software (Version 4.8), The Sage Development Team. http://www.sagemath.org. \uparrow 3.1

Tenenbaum, Gérald and Michel Mendès France. 2000. *The prime numbers and their distribution*, Student Mathematical Library, vol. 6, American Mathematical Society, Providence, RI. Translated from the 1997 French original by Philip G. Spain. MR1756233 (2001j:11086) ↑2.3

LABORATORY OF ALGEBRAIC GEOMETRY, NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF ECONOMICS, 7 VAVILOVA STR., MOSCOW, RUSSIA, 117312; AND LABORATOIRE DE MATH., UNIVERSITÉ PARIS-SUD, CAMPUS D'ORSAY, BÂTIMENT 425, 91405 ORSAY, FRANCE

E-mail address: Ekaterina.Amerik@math.u-psud.fr *URL*: www.math.u-psud.fr/~amerik

Department of Mathematics, KTH Royal Institute of Technology, SE-100 44 Stockholm, Sweden

E-mail address: kurlberg@math.kth.se *URL*: www.math.kth.se/~kurlberg

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720 *E-mail address*: khoanguyen2511@gmail.com *URL*: http://math.berkeley.edu/~khoa

DEPARTMENT OF MATHEMATICS, CUNY GRADUATE CENTER, NEW YORK, NY, 10016, USA *E-mail address*: atowsley@gc.cuny.edu *URL*: https://sites.google.com/site/atowsley42/

DEPARTMENT OF MATHEMATICS, BOX 1917, BROWN UNIVERSITY, PROVIDENCE, RI 02912, USA *E-mail address:* bviray@math.brown.edu *URL:* http://math.brown.edu/~bviray

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEXAS, AUSTIN, TX 78712, USA *E-mail address*: voloch@math.utexas.edu *URL*: http://www.ma.utexas.edu/users/voloch/