

## COMPUTING ISOGENY COVARIANT DIFFERENTIAL MODULAR FORMS

CHRIS HURLBURT

ABSTRACT. We present the computation modulo  $p^2$  and explicit formulas for the unique isogeny covariant differential modular form of order one and weight  $\chi_{-p-1,-p}$  called  $f_{\text{jet}}$ , an isogeny covariant differential modular form of order two and weight  $\chi_{-p^2-p,-1,-1}$  denoted by  $f_{\text{jet}}h_{\text{jet}}$ , and an isogeny covariant differential modular form  $h_{\text{jet}}$  of order two and weight  $\chi_{1-p^2,0,-1}$ .

### 1. INTRODUCTION

In this paper we introduce explicit formulas modulo  $p^2$  for various differential modular forms discussed by Buium in [3], [2], [4]. The central modular form discussed is the unique, up to multiplication by an element in  $\mathbb{Z}_p^*$ , isogeny covariant differential modular form of order one and weight  $\chi_{-p-1,-1}$  called  $f_{\text{jet}}$ ,  $f_p^1$ , and  $f_{\text{jet}}^1$ , respectively, in [3], [2], [4]. For the rest of this paper we mean “unique up to multiplication by an element in  $\mathbb{Z}_p^*$ ” when we say “unique”, and we will refer to the unique isogeny covariant differential modular form of order one and weight  $\chi_{-p-1,-1}$  by  $f_{\text{jet}}$ . This modular form has many interesting connections detailed in [1], [3], [2], and [4]. We compute  $f_{\text{jet}}$  in a  $p$ -adic fashion following the construction of  $f_{\text{jet}}$  detailed in [3] which allows us to compute modulo  $p^n$  or specifically modulo  $p^2$ . Then we use the explicit formula from this computation to provide modulo  $p^2$  formulas for order two differential modular forms. The specific order two isogeny covariant differential modular forms we describe are  $f_{\text{jet}}h_{\text{jet}}$  from [3] also referred to as  $k_p^2$  in [2] or  $f_{\text{jet}}^{1,2}$  in [4] and  $h_{\text{jet}}$  from [3] also referred to as  $k_p^2/f_p^1$  in [2]. We note that modulo  $p$  neither of these order two modular forms contain any second order terms.

The strategy is simple. We know that the isogeny covariant differential modular forms  $f_{\text{jet}}h_{\text{jet}}$  and  $h_{\text{jet}}$  of order two and weights  $\chi_{-p^2-p,-1,-1}$  and  $\chi_{1-p^2,0,-1}$ , respectively, are  $f_{\text{jet}}h_{\text{jet}} = \phi(f_{\text{jet}})$ , where  $\phi$  is the lifting of the Frobenius morphism, and outside the locus, where  $f_{\text{jet}}$  modulo  $p$  is zero  $h_{\text{jet}} = \frac{\phi(f_{\text{jet}})}{f_{\text{jet}}}$  [4]. We should note that  $h_{\text{jet}}$  is defined only outside this zero locus of  $f_{\text{jet}}$  modulo  $p$ . In [5] we have the explicit computation of  $\bar{\mathcal{F}}_{\text{def}}$  (the  $p$ -derivation analog of the Kodaira-Spencer class) which is the reduction modulo  $p$  of the unique isogeny covariant differential modular form of weight  $\chi_{-p-1,-1}$ . By uniqueness,  $f_{\text{jet}} \equiv c\bar{\mathcal{F}}_{\text{def}}$  modulo  $p$  for some  $c \in \mathbb{Z}_p^*$ . Here we compute  $f_{\text{jet}}$  directly allowing us to give a formula for the unique

---

Received by the editor January 14, 2004 and, in revised form, April 16, 2004.

2000 *Mathematics Subject Classification*. Primary 11F11; Secondary 12H05.

This research was supported in part by NSA grant MDA904-03-1-0031.

isogeny covariant differential modular form modulo  $p^2$  and not just modulo  $p$ . We also then are able to describe the order two terms that occur in  $f_{\text{jet}}h_{\text{jet}}$  and  $h_{\text{jet}}$  modulo  $p^2$  but not modulo  $p$ .

For both context and notation we give the relevant definitions of differential modular forms. Let  $p > 3$  be a prime number. Let  $M^0 = \mathbb{Z}_p[a_4, a_6, \Delta^{-1}]^\wedge$ ,  $M^1 = \mathbb{Z}_p[a_4, a_6, \delta a_4, \delta a_6, \Delta^{-1}]^\wedge$ , and  $M^2 = \mathbb{Z}_p[a_4, a_6, \delta a_4, \delta a_6, \delta^2 a_4, \delta^2 a_6, \Delta^{-1}]^\wedge$ , where  $\Delta = -2^4(4a_4^3 + 27a_6^2)$  and  $\mathbb{Z}_p$  is the  $p$ -adic integer. We note that  $a_4, a_6, \delta a_4, \delta a_6, \delta^2 a_4, \delta^2 a_6$  are variables over  $\mathbb{Z}_p$  and that  $^\wedge$  represents the  $p$ -adic completion. Then the elements of  $M^1$  are called  $\delta$  modular forms of order one and elements of  $M^2$  are called  $\delta$  modular forms of order two as defined by Buium in [3].

Recall now that a  $p$ -derivation is a set theoretic map,  $\delta : A \rightarrow B$ , from a ring  $A$  to an  $A$ -algebra  $B$  such that

$$(1.1) \quad \delta(x + y) = \delta x + \delta y + C_p(x, y),$$

$$(1.2) \quad \delta(xy) = y^p \delta x + x^p \delta y + p \delta x \delta y$$

for all  $x, y \in A$ , where  $C_p(X, Y) = \frac{X^p + Y^p - (X+Y)^p}{p}$ . In Section 2 we will expand these axioms into a more complete list of properties of  $p$ -derivations. For now, if  $A$  is a complete discrete valuation ring  $R$ , where  $R$  has maximal ideal generated by  $p$  and an algebraically closed residue field  $k$ , and if  $\phi$  is the unique lifting of the Frobenius morphism to  $A$ , then the  $p$ -derivation given by  $\delta(x) = (\phi(x) - x^p)/p$  is unique on  $R$ .

Now we use the  $R$  and  $\delta$  from our example and set

$$M(R) = \{(a, b) \in R^2 \mid 4a^3 + 27b^2 \in R^*\}.$$

Then the set  $M(R)$  is in one-to-one correspondence with the set of pairs consisting of an elliptic curve over  $R$  and an invertible 1-form; namely, each  $(\bar{a}_4, \bar{a}_6) \in M(R)$  corresponds to  $(E, dx/2y)$ , where  $E$  is the projective closure of the affine plane curve  $y^2 = x^3 + \bar{a}_4 x + \bar{a}_6$ . For any  $f \in M^1$ , if we substitute  $a, b, \delta a, \delta b$  in for  $a_4, a_6, \delta a_4, \delta a_6$ , then  $f$  defines a map (still denoted by  $f$ ) from  $M(R)$  to  $R$ . This element in  $M^1$  is in fact uniquely determined by the map from  $M(R)$  to  $R$ . Similar statements are true for  $f \in M^2$ .

We define a  $\delta$ -character of order  $\leq 1$  to be a group homomorphism  $\chi : R^* \rightarrow R^*$  of the form  $\chi = \chi_{m,n}$ , where

$$\chi_{m,n}(\lambda) = \lambda^m \left( \frac{\phi(\lambda)}{\lambda^p} \right)^n.$$

Then a  $\delta$ -modular function of order one has weight  $\chi$  if for any  $\lambda \in R^*$

$$f(\lambda^4 a, \lambda^6 b) = \chi(\lambda) f(a, b)$$

for all  $(a, b) \in M(R)$ . We can easily extend the definition of  $\delta$ -characters to higher orders. Namely, a  $\delta$ -character of order  $\leq 2$  is a group homomorphism  $\chi : R^* \rightarrow R^*$  of the form  $\chi = \chi_{m,n,r}$  where

$$\chi_{m,n,r}(\lambda) = \lambda^m \left( \frac{\phi(\lambda)}{\lambda^p} \right)^n \left( \frac{\phi^2(\lambda)}{\lambda^{p^2}} \right)^r.$$

The criterion for a  $\delta$ -modular function of order two to have weight  $\chi$  is exactly the same as the criterion for a  $\delta$ -modular function of order one to have weight  $\chi$ . A  $\delta$ -modular form is a  $\delta$ -modular function with a weight.

A  $\delta$ -modular form is *isogeny covariant* if for any two pairs  $(a, b)$  and  $(\tilde{a}, \tilde{b})$  with an etale isogeny of degree  $N$  between the corresponding elliptic curves that pulls back  $\frac{dx}{y}$  to  $\frac{d\tilde{x}}{\tilde{y}}$

$$f(a, b) = N^{-k/2} f(\tilde{a}, \tilde{b}),$$

where  $k$  is a constant that depends solely on the weight. Note that for  $\chi = \chi_{m,n}$  the constant is  $k = m + n(1 - p)$  and for  $\chi = \chi_{m,n,r}$  the constant is  $k = m + n(1 - p) + r(1 - p^2)$ .

**Theorem 1.1.** *The isogeny covariant differential modular form of order one and weight  $\chi_{-p-1,-1}$  modulo  $p^2$  is*

$$f_{\text{jet}} = \left[ \frac{-72a_6^p \delta a_4 + 48a_4^p \delta a_6}{\Delta^p} \right] \gamma_{2p,p} + h + pH,$$

where  $\gamma_{2p,p}$  and  $h$  are polynomials in  $M_1^0 := M^0 \otimes \mathbb{Z}_p/(p^2)$ ,  $H$  is a polynomial in  $M_0^1 := M^1 \otimes \mathbb{Z}_p/(p)$ , and  $H$  is a nonhomogeneous quadratic in  $\delta a_4$  and  $\delta a_6$ .

Explicit formulas for  $h$  and  $H$  are given in Theorem 6.11 and an explicit formula for  $\gamma_{2p,p}$  is given in Proposition 6.2.

**Theorem 1.2.** *The isogeny covariant differential modular form  $f_{\text{jet}} h_{\text{jet}}$  of order two and weight  $\chi_{-p^2-p,-1,-1}$  modulo  $p^2$  is*

$$\begin{aligned} f_{\text{jet}} h_{\text{jet}} = & \left[ \frac{-72a_6^{p^2} (\delta a_4)^p + 48a_4^{p^2} (\delta a_6)^p}{\Delta^{p^2}} \right] \gamma_{2p,p}(a_4^p, a_6^p) + h^* \\ & + p \left[ \frac{-72a_6^{p^2} \delta^2 a_4 + 48a_4^{p^2} \delta^2 a_6}{\Delta^{p^2}} \right] \gamma_{2p,p}(a_4^p, a_6^p) + pJ, \end{aligned}$$

where  $h^*$  is a polynomial in  $M_1^1 := M^1 \otimes \mathbb{Z}_p/(p^2)$  and  $J$  is a polynomial in  $M_0^1$ .

**Corollary 1.3.** *The isogeny covariant differential modular form  $h_{\text{jet}}$  of order two and weight  $\chi_{1-p^2,0,-1}$  modulo  $p^2$  is*

$$\begin{aligned} h_{\text{jet}} = & \frac{\left[ -72a_6^{p^2} (\delta a_4)^p + 48a_4^{p^2} (\delta a_6)^p \right] \gamma_{2p,p}(a_4^p, a_6^p) + \Delta^{p^2} h^*}{\Delta^{p^2-p} ([-72a_6^p \delta a_4 + 48a_4^p \delta a_6] \gamma_{2p,p} + \Delta^p h)} \\ & + p \left( \frac{\left( \left[ -72a_6^{p^2} (\delta a_4)^p + 48a_4^{p^2} (\delta a_6)^p \right] \gamma_{2p,p}(a_4^p, a_6^p) + \Delta^{p^2} h^* \right) H}{\Delta^{p^2-2p} ([-72a_6^p \delta a_4 + 48a_4^p \delta a_6] \gamma_{2p,p} + \Delta^p h)^2} \right. \\ & \left. + \frac{\left[ 72a_6^{p^2} \delta^2 a_4 + 48a_4^{p^2} \delta^2 a_6 \right] \gamma_{2p,p}(a_4^p, a_6^p) + \Delta^{p^2} J}{\Delta^{p^2-p} ([-72a_6^p \delta a_4 + 48a_4^p \delta a_6] \gamma_{2p,p} + \Delta^p h)} \right), \end{aligned}$$

where  $h$ ,  $h^*$ ,  $H$ , and  $J$  are the same as in Theorems 1.1 and 1.2.

What follows is preliminary information to the calculation of  $f_{\text{jet}}$ . Let  $E$  be the elliptic curve in Weierstrass form over  $M^0$  defined by the homogeneous equation

$$f(X, Y, W) = WY^2 - X^3 - a_4 XW^2 - a_6 W^3.$$

Let  $U$  and  $V$  be the affine open subsets of  $E$  given by the equations  $f(x, y, 1)$  and  $f(z, 1, w)$ , respectively. So

$$U = \text{Spec } M^0[X, Y]/(f(X, Y, 1)) = \text{Spec } M^0[x, y],$$

$$V = \text{Spec } M^0[Z, W]/(f(Z, 1, W)) = \text{Spec } M^0[z, w],$$

and on  $U \cap V$

$$z = -x/y,$$

$$w = -1/y,$$

whence  $E = U \cup V$ . Next we define the first jets of  $U$  and  $V$  to be the sets

$$U^1 = \operatorname{Spec} M^1[X, Y, \delta X, \delta Y]/(f(X, Y, 1), \delta f(X, Y, 1)) = \operatorname{Spec} M^1[x, y, \delta x, \delta y],$$

$$V^1 = \operatorname{Spec} M^1[Z, W, \delta Z, \delta W]/(f(Z, 1, W), \delta f(Z, 1, W)) = \operatorname{Spec} M^1[z, w, \delta z, \delta w].$$

Then  $E^1$ , the first jet space of  $E$ , is the gluing of  $U^1$  and  $V^1$  by the maps

$$(1.3) \quad \begin{aligned} z &= -x/y, \\ w &= -1/y, \\ \delta z &= \frac{x^p \delta y - y^p \delta x}{y^p(y^p + p\delta y)}, \\ \delta w &= \frac{\delta y}{y^p(y^p + p\delta y)}. \end{aligned}$$

We can extend the group law on  $E$  to a group law on  $E^1$ . The group law arises naturally by construction from the group law on  $E$  just as  $E^1$  arises naturally by construction from  $E$ . This will be detailed explicitly in Section 3.

From now on we will also use the following notation. First by  $M_n^i$  we mean  $M^i \otimes \mathbb{Z}_p/(p^{n+1})$ . For example  $M_0^1 = \mathbb{F}_p[a_4, a_6, \delta a_4, \delta a_6, \Delta^{-1}]$ , where  $\mathbb{F}_p$  is the finite field of  $p$  elements. Second by  $E_m^1$  we mean  $E^i \otimes M_m^1$ , and by  $E_m = E_m^0$  we mean  $E \otimes M_m^0$ . Also we will use  $\delta(a_4)$  interchangeably for  $\delta a_4$ ,  $\delta(a_6)$  interchangeably for  $\delta a_6$ , etc.

To compute  $f_{\text{jet}}$ , the isogeny covariant  $\delta$  modular form of weight  $\chi_{-p-1, -1}$ , we work from [2, Construction 4.1]. The same construction is also described in [4] and [3]. First we find two sections  $s_U$  and  $s_V$  of the morphisms  $U^1 \rightarrow U \otimes M^1$  and  $V^1 \rightarrow V \otimes M^1$ , respectively, such that  $s_U$  defines a morphism from  $U \otimes M^1$  to  $U^1$  and  $s_V$  defines a morphism from  $V \otimes M^1$  to  $V^1$ . Then the difference of the sections under the group law induces a morphism  $s_U - s_V : U \cap V \otimes M^1 \rightarrow E^1$ . Let  $\zeta$  be the  $\delta z$  coordinate in the difference  $s_U - s_V$ . By the  $\delta z$  coordinate, we mean the image of  $\delta z \in U^1 \cap V^1$  under the ring homomorphism induced by the morphism  $s_U - s_V : U \cap V \otimes M^1 \rightarrow E^1$ . Let  $\log_{\mathcal{F}_1^{\phi^1}}(\xi)$  be the formal logarithm of the Frobenius twist of the formal group of the elliptic curve, namely

$$\log_{\mathcal{F}_1^{\phi^1}}(\xi) = \xi + \frac{p\phi(c_1)}{2}\xi^2 + \frac{p^2\phi(c_2)}{3}\xi^3 + \cdots,$$

where the  $c_i$  are the coefficients of the power series expansion of the invariant differential [3]. Then  $\log_{\mathcal{F}_1^{\phi^1}}(\zeta)$  is a cohomology class in  $H^1(E \otimes M^1, \mathcal{O}) \simeq H^1(E, \mathcal{O}) \otimes M^1$ , and this resulting class has a representative of the form  $\sum a_n y^n + x \sum b_n y^n + x^2 \sum e_n y^n$ . The modular form  $f_{\text{jet}}$  is the coefficient  $e_{-1}$  of  $x^2/y$  in this representative which is the residue of the cohomology class, namely the image of the cohomology class under the Serre duality pairing.

We will actually work modulo  $p^2$  which means that our end result will be  $f_{\text{jet}}$  modulo  $p^2$ . In fact  $f_{\text{jet}} \in M^1$  is a restricted power series whose coefficients expand exponentially in the number of terms in each coefficient of a power of  $p$ . Therefore, the formulas necessary to express  $f_{\text{jet}}$  modulo  $p^n$  for  $n > 2$  are prohibitive in length. At this point we note that the formal logarithm of the Frobenius twist of the formal

group of the elliptic curve modulo  $p^2$  is in fact the identity. This certainly simplifies one step of the computation modulo  $p^2$ ; however, for  $n > 4$  this formal logarithm is no longer trivial modulo  $p^n$ , meaning this step is not trivial for large  $n$ . As a preliminary step to computing  $f_{\text{jet}}$  we detail some computation guidelines for  $p$ -derivations and the group law for  $E^1$  modulo  $p^2$ .

## 2. PROPERTIES OF $p$ -DERIVATIONS

Recall that a  $p$ -derivation is a set theoretic map,  $\delta : A \rightarrow B$ , from a ring  $A$  to an  $A$ -algebra  $B$  with  $\delta(1) = 0$  such that

$$\delta(x + y) = \delta x + \delta y + C_p(x, y), \quad \delta(xy) = y^p \delta x + x^p \delta y + p \delta x \delta y$$

for all  $x, y \in A$ , where  $C_p(X, Y) = \frac{X^p + Y^p - (X+Y)^p}{p}$ . In the case when  $A = B = R$ , where  $R$  is a complete discrete valuation ring with maximal ideal generated by  $p$  and has an algebraically closed residue field, there is a unique  $p$ -derivation given by  $\delta(x) = (\phi(x) - x^p)/p$ , where  $\phi$  is the unique lifting of the Frobenius morphism to  $R$ .

This definition implies that if  $\varphi : A \rightarrow B$  is the ring homomorphism associated to  $B$  being an  $A$ -algebra, then

$$(\varphi, \delta) : A \rightarrow W_2(B)$$

is a ring homomorphism, where  $W_2(B)$  is the ring of *Witt vectors* of length two on  $B$ . With  $(\varphi, \delta)$  as above,  $\phi : A \rightarrow B$  defined by  $\phi(x) = \varphi(x)^p + p\delta(x)$  is a ring homomorphism. In case  $B = A$ , this is a lifting of the Frobenius endomorphism  $F(x) = x^p$  of  $A/pA$ .

While no further axioms for  $p$ -derivations beyond those in the definition are necessary for computation, the following  $p$ -derivation rules are very convenient for computation. Before introducing these rules we must define an extension of  $C_p(X, Y)$ .

**Definition 2.1.** For any  $\sum q$ , let

$$C_p^{\text{ext}}(\sum q) = \frac{\sum q^p - (\sum q)^p}{p}.$$

Note that  $C_p^{\text{ext}}(X + Y) = \frac{X^p + Y^p - (X+Y)^p}{p} = C_p(X, Y)$ ; thus, this is a very natural definition.

**Lemma 2.2.** Let  $\delta : A \rightarrow B$  be a  $p$ -derivation, let  $g = \sum q, x, y \in A$ , and let  $n > 0$  be an integer. Then the following are true.

- (1)  $\delta(\sum q) = \sum \delta q + C_p^{\text{ext}}(\sum q)$ .
- (2)  $\delta(-1) = 0$ .
- (3)  $\delta(-x) = -\delta x$ .
- (4)  $\delta(x^n) = \sum_{k=1}^n \binom{n}{k} p^{k-1} x^{(n-k)p} (\delta x)^k = \frac{-x^{np} + (x^p + p\delta x)^n}{p}$ .
- (5)  $\delta\left(\frac{1}{x}\right) = \frac{-\delta x}{x^p(x^p + p\delta x)}$ .
- (6)  $\delta\left(\frac{y}{x}\right) = \frac{x^p \delta y - y^p \delta x}{x^p(x^p + p\delta x)}$ .

### 3. THE GROUP LAW FOR THE FIRST $p$ -JET SPACE OF $E$

We now want to make the group law on  $E^1$  explicit. This is necessary since the main result requires us to subtract two sections using the group law. The group law on the first  $p$ -jet is induced by the group law on the elliptic curve  $E$ , so we start by giving the group law on  $E$ . Let  $\rho$  and  $\psi$  be the equations that define the group law on  $E$ . So if  $(z_1, w_1) \oplus (z_2, w_2) = (z_3, w_3)$ , then

$$\begin{aligned} z_3 &= \rho(z_1, w_1, z_2, w_2), \\ w_3 &= \psi(z_1, w_1, z_2, w_2). \end{aligned}$$

Then the group law on  $E^1$  is an extension of the group law on  $E$  such that if  $(z_1, w_1, \delta z_1, \delta w_1) \oplus (z_2, w_2, \delta z_2, \delta w_2) = (z_3, w_3, \delta z_3, \delta w_3)$ , then

$$\begin{aligned} z_3 &= \rho(z_1, w_1, z_2, w_2), \\ w_3 &= \psi(z_1, w_1, z_2, w_2), \\ \delta z_3 &= \delta(\rho(z_1, w_1, z_2, w_2)), \\ \delta w_3 &= \delta(\psi(z_1, w_1, z_2, w_2)). \end{aligned}$$

To find appropriate  $\rho$  and  $\psi$ , we must consider actual formulas for the group law. On the elliptic curve  $E$ , the group law can be explicitly formulated using the chord-tangent approach. In this approach we consider that every line intersects the elliptic curve at exactly three points counting multiplicity. We choose a specific point,  $O$ , to be the origin; in this case the point we choose to be the origin is the point at infinity,  $(0, 1, 0)$ . We then define the inverse of a point  $P$  to be the third point on the line that intersects  $P$  and the origin. We denote this point by  $-P$ . So if we want to add  $P \oplus Q$ , we take the line through  $P$  and  $Q$  and let  $R$  be the third point on the line. Then we define  $P \oplus Q = -R$ . This definition arises naturally from the theory of Weil divisors. We refer to the case when  $P = Q$  as the tangent case and  $P \neq Q$  as the chord case. From now on we will focus on the chord case of the chord-tangent approach since that is the most general case and the case used when computing the group law for a  $p$ -jet space.

We use the standard procedure for finding explicit formulas for group law in the  $z$  and  $w$  coordinates. In these coordinates our origin is  $(0, 0)$ . To start with, we recall data on  $V$ ; namely, that  $f(z, 1, w) = w - z^3 - a_4zw^2 - a_6w^3$  is the curve we will be using and that any line through  $P = (z_0, w_0)$  and the origin of  $(0, 0)$  will intersect  $f(z, 1, w)$  at the third point  $(-z_0, -w_0)$ . Whence  $-P = (-z_0, -w_0)$ .

Now consider two points  $P_1$  and  $P_2$  denoted by  $(z_i, w_i)$  for  $i = 1, 2$ , respectively. If we assume that  $z_1 \neq z_2$ , the line connecting these two points is

$$w = \frac{w_2 - w_1}{z_2 - z_1}(z - z_1) + w_1.$$

To find the sum  $P_1 \oplus P_2$ , we must find the three points counting multiplicity of the intersection of this line with the curve  $f(z, 1, w)$ . If we substitute the line into the curve  $f(z, 1, w)$  we get a cubic equation in terms of  $z$ . Finding these three points becomes a matter of finding the roots of the resulting cubic equation. On the other hand, we already know two of the roots, namely  $z = z_1$  and  $z = z_2$ . The third root

is

$$z = \frac{-2w_2z_1 - w_1z_1 + 2w_1z_2 - 3a_6w_2w_1^2z_2 - a_4w_1^2z_2^2 + w_2z_2 + 3w_2^2z_1a_6w_1 + w_2^2z_1^2a_4}{3a_6w_2w_1(w_2 - w_1) + 3z_2z_1(z_2 - z_1) + a_4(w_2^2z_1 - w_1^2z_2) + w_1 - w_2 + 2a_4w_2w_1(z_2 - z_1)}.$$

So for  $z_1 \neq z_2$ , the  $P_3 = P_1 \oplus P_2$  has coordinates

$$z_3 = -\frac{-2w_2z_1 - w_1z_1 + 2w_1z_2 - 3a_6w_2w_1^2z_2 - a_4w_1^2z_2^2 + w_2z_2 + 3w_2^2z_1a_6w_1 + w_2^2z_1^2a_4}{3a_6w_2w_1(w_2 - w_1) + 3z_2z_1(z_2 - z_1) + a_4(w_2^2z_1 - w_1^2z_2) + w_1 - w_2 + 2a_4w_2w_1(z_2 - z_1)},$$

$$w_3 = -\frac{3w_2z_2z_1^2 + z_1w_2^2w_1a_4 - 3z_1w_1z_2^2 + w_1^2 - w_2w_1^2z_2a_4 - w_2^2}{3a_6w_2w_1(w_2 - w_1) + 3z_2z_1(z_2 - z_1) + a_4(w_2^2z_1 - w_1^2z_2) + w_1 - w_2 + 2a_4w_2w_1(z_2 - z_1)}.$$

From this information, if we want the formulation of group law on  $E^1$  we must simply take the  $p$ -derivation of these equations. The resulting group law for  $P_i = (z_i, w_i, \delta z_i, \delta w_i) \in V^1$  is

$$z_3 = -\frac{\alpha}{\mu},$$

$$w_3 = -\frac{\beta}{\mu},$$

$$\delta z_3 = -\frac{\mu^p \delta \alpha - \alpha^p \delta \mu}{\mu^p (\mu^p + p \delta \mu)},$$

$$\delta w_3 = -\frac{\mu^p \delta \beta - \beta^p \delta \mu}{\mu^p (\mu^p + p \delta \mu)},$$

where  $P_3 = P_1 \oplus P_2$ ,

$$\alpha = -2w_2z_1 - w_1z_1 + 2w_1z_2 - 3a_6w_2w_1^2z_2 - a_4w_1^2z_2^2 + w_2z_2 + 3w_2^2z_1a_6w_1 + w_2^2z_1^2a_4,$$

$$\beta = 3w_2z_2z_1^2 + z_1w_2^2w_1a_4 - 3z_1w_1z_2^2 + w_1^2 - w_2w_1^2z_2a_4 - w_2^2,$$

$$\mu = 3a_6w_2w_1(w_2 - w_1) + 3z_2z_1(z_2 - z_1) + a_4(w_2^2z_1 - w_1^2z_2) + w_1 - w_2 + 2a_4w_2w_1(z_2 - z_1),$$

and  $\delta\alpha, \delta\beta, \delta\mu$  are the respective  $p$ -derivatives which are not included here because of their lengthy nature. On the other hand, this group law also describes the group law on  $E_m^1$ . For example, if  $m = 1$ , then we consider this same group law modulo  $p^2$ . This does shorten the expressions of  $\delta\alpha, \delta\beta$ , and  $\delta\mu$  for  $m \leq 5$  to lengths that are possible to work with in computer algebra systems.

Besides shortening the expressions for  $\alpha, \beta$ , and  $\mu$  one other advantage of explicitly detailing the group law on  $E_1^1$  rather than  $E^1$  is that we may write  $\delta z_3$  and  $\delta w_3$  in terms of polynomials in  $\delta\alpha, \delta\beta$ , and  $\delta\mu$  by using their series expansions. Hence

we have the following description for the group law on  $E_1^1$ :

$$\begin{aligned} z_3 &= -\frac{\alpha}{\mu}, \\ w_3 &= -\frac{\beta}{\mu}, \\ \delta z_3 &= \frac{1}{\mu^{3p}}(-\mu^p \delta \alpha + \alpha^p \delta \mu)(\mu^p - p \delta \mu), \\ \delta w_3 &= \frac{1}{\mu^{3p}}(-\mu^p \delta \beta + \beta^p \delta \mu)(\mu^p - p \delta \mu), \end{aligned}$$

where  $\delta \alpha$ ,  $\delta \beta$ , and  $\delta \mu$  are now expressions modulo  $p^2$ .

4. THE SECTION ON  $U$  THAT DEFINES A MAP FROM  $U \otimes M^1$  TO  $U^1$  AND THE SECTION ON  $V$  THAT DEFINES A MAP FROM  $V \otimes M^1$  TO  $V^1$

We in fact want a specific map from  $U$  to  $U^1$ ; namely, the morphism which takes  $\delta x$  and  $\delta y$  to elements such that  $\delta f(x, y, 1)$  is mapped to 0. To do this we use a variant of Hensel's Lemma involving two variables which will be illuminated as we go along. To find the appropriate  $\delta x$  and  $\delta y$ , we first consider the explicit expression of  $\delta f(x, y, 1)$ ,

$$\begin{aligned} (4.1) \quad & -p^2 \delta x^3 + (-3x^p \delta x^2 - \delta a_4 \delta x + \delta y^2)p \\ & + (-3x^{2p} - a_4^p) \delta x + 2y^p \delta y - \delta a_4 x^p - \delta a_6 + C_p^{\text{ext}}(y^2 - x^3 - a_4 x - a_6), \end{aligned}$$

and from this polynomial define  $P_{U,0} = -\delta a_4 x^p - \delta a_6 + C_p^{\text{ext}}(y^2 - x^3 - a_4 x - a_6)$ . From now on for convenience of notation we will denote  $f(x, y, 1)$  simply by  $f$ . We will also denote  $\frac{\partial f}{\partial x}$  and  $\frac{\partial f}{\partial y}$  by  $f_x$  and  $f_y$ , respectively.

Now we let  $A$  and  $B$  be polynomials in  $M^0[x, y]$  such that  $Af_x + Bf_y = 1$ . Specifically

$$\begin{aligned} A &= \frac{2^4(4a_4^2 + 6x^2 a_4 - 9xa_6)}{\Delta}, \\ B &= \frac{2^3(9y)(2xa_4 - 3a_6)}{\Delta}. \end{aligned}$$

By simple arithmetic,  $A^p f_x^p + B^p f_y^p = 1 + A^p f_x^p + B^p f_y^p - (Af_x + Bf_y)^p = 1 + pC_p^{\text{ext}}(Af_x + Bf_y)$ . Now we consider the relationship between  $f_x^p$ ,  $f_y^p$  and the coefficients of  $\delta x$  and  $\delta y$ , respectively. First recall that  $n = n^p + p\delta(n)$  for any positive integer  $n$ . So we can write the coefficients of  $\delta x$  and  $\delta y$  from equation (4.1) as

$$\begin{aligned} \text{Coefficient of } \delta x &= -3x^{2p} - a_4^p = -(3^p + p\delta(3))x^{2p} - a_4^p \\ &= -p\delta(3)x^{2p} + f_x^p + pC_p^{\text{ext}}(-3x^2 - a_4) \\ &= f_x^p + p(-\delta(3)x^{2p} + C_p^{\text{ext}}(-3x^2 - a_4)), \\ \text{Coefficient of } \delta y &= 2y^p = (2^p + p\delta(2))y^p \\ &= f_y^p + p\delta(2)y^p. \end{aligned}$$



Then combining these with the equation  $A^p f_x^p + B^p f_y^p = 1 + pC_p^{\text{ext}}(Af_x + Bf_y)$ ,

$$\begin{aligned} & A^p(-3x^{2p} - a_4^p) + B^p(2y^p) \\ &= A^p(f_x^p + p(-\delta(3)x^{2p} + C_p^{\text{ext}}(-3x^2 - a_4))) + B^p(f_y^p + p\delta(2)y^p) \\ &= 1 + p(C_p^{\text{ext}}(Af_x + Bf_y) + A^p(-\delta(3)x^{2p} + C_p^{\text{ext}}(-3x^2 - a_4)) + B^p\delta(2)y^p). \end{aligned}$$

Now if we let  $R_{U,0} = C_p^{\text{ext}}(Af_x + Bf_y) + A^p(-\delta(3)x^{2p} + C_p^{\text{ext}}(-3x^2 - a_4)) + B^p\delta(2)y^p$ , then  $A^p(-3x^{2p} - a_4^p) + B^p(2y^p) = 1 + pR_{U,0}$ .

With the computations in the previous paragraph we now have enough tools to perform the iteration step in Hensel's Lemma. We assume that

$$\begin{aligned} \delta x &= -P_{U,0}A^p + p\eta, \\ \delta y &= -P_{U,0}B^p + p\sigma, \end{aligned}$$

and plug these assumptions into equation (4.1). Then we solve the resulting equation for  $\eta$  and  $\sigma$ , keeping in mind that we are working modulo  $p^2$ . (Note: The procedure is the same working modulo  $p^3$  etc., but in that case one must assume a  $p^2$  term for the  $\delta x$  and  $\delta y$  and then perform the iteration twice.)

$$\begin{aligned} & (-3x^p\delta x^2 - \delta a_4\delta x + \delta y^2)p + (-3x^{2p} - a_4^p)\delta x + 2y^p\delta y + P_{U,0} \\ &= p(-3x^pP_{U,0}^2A^{2p} + \delta a_4P_{U,0}A^p + P_{U,0}^2B^{2p} \\ &\quad - P_{U,0}R_{U,0} + (-3x^{2p} - a_4^p)\eta + 2y^p\sigma). \end{aligned}$$

Now if we let  $P_{U,1} = -3x^pP_{U,0}^2A^{2p} + \delta a_4P_{U,0}A^p + P_{U,0}^2B^{2p} - P_{U,0}R_{U,0}$ , then

$$\begin{aligned} \eta &= -P_{U,1}A^p, \\ \sigma &= -P_{U,1}B^p. \end{aligned}$$

So the morphism that takes  $x$  to  $x$ ,  $y$  to  $y$ ,  $\delta x$  to  $-P_{U,0}A^p - pP_{U,1}A^p$ , and  $\delta y$  to  $-P_{U,0}B^p - pP_{U,1}B^p$  will map  $\delta f$  to 0. Our corresponding section,  $s_U$ , is

$$(x, y, -A^p(P_{U,0} + pP_{U,1}), -B^p(P_{U,0} + pP_{U,1})).$$

Next we find the section  $s_V$  that defines a specific map from  $V$  to  $V^1$  such that under this map  $\delta f(z, 1, w)$  is taken to 0. Since the techniques used are identical to those used to find  $s_U$ , we will omit most of the details. From now on for convenience of notation we will refer to  $f(z, 1, w)$  as  $g$ , and  $\frac{\partial g}{\partial z}$ ,  $\frac{\partial g}{\partial w}$  will be referred to as  $g_z$  and  $g_w$ , respectively.

Let  $P_{V,0} = -\delta a_6w^{3p} - \delta a_4z^pw^{2p} + C_p^{\text{ext}}(w - z^3 - a_4zw^2 - a_6w^3)$  and let  $C$  and  $D$  be polynomials in  $M^0[z, w]$  such that  $Cg_z + Dg_w = 1$ . Specifically

$$C = z(-\frac{3}{2}a_6w - a_4z), \quad D = -\frac{3}{2}a_6w^2 - wa_4z + 1.$$

Next let

$$R_{V,0} = C_p^{\text{ext}}(Cg_z + Dg_w) + C^p(-\delta(3)z^{2p} + C_p^{\text{ext}}(-3z^2 - a_4w^2)) \\ + D^p(-\delta(3)a_6^p w^{2p} - \delta(2)a_4^p z^p w^p + C_p^{\text{ext}}(1 - 3a_6w^2 - 2a_4zw))$$

and let

$$P_{V,1} = -3z^p(P_{V,0}C^p)^2 + (2a_4^p w^p(-P_{V,0}D^p) + \delta a_4 w^{2p})(P_{V,0}C^p) \\ - (a_4^p z^p + 3a_6^p w^p)(P_{V,0}D^p)^2 - (3\delta a_6 w^{2p} + 2\delta a_4 z^p w^p)(-P_{V,0}D^p) - P_{V,0}R_{V,0}.$$

Then the section  $s_V$  defining a map from  $V$  to  $V^1$  is

$$(z, w, -C^p(P_{V,0} + pP_{V,1}), -D^p(P_{V,0} + pP_{V,1})).$$

### 5. $s_U - s_V$ UNDER THE GROUP LAW

We now need the  $\delta z$  coordinate also referred to as  $\zeta$  in the difference,  $s_U - s_V$ , of these two sections under the group law. We will work with the element  $(z, w, z', w')$  where  $z' = \delta z$  and  $w' = \delta w$ . Recall from the Introduction that our gluing maps on the intersection  $U^1 \cap V^1$  are

$$z = -x/y, \\ w = -1/y, \\ \delta z = \frac{x^p \delta y - y^p \delta x}{y^p(y^p + p\delta y)}, \\ \delta w = \frac{\delta y}{y^p(y^p + p\delta y)}.$$

So if we let  $x' = \delta x$  and  $y' = \delta y$ , then in terms of the coordinates on  $U^1$ , our element is  $(-x/y, -1/y, \frac{x^p y' - y^p x'}{y^p(y^p + p y')}, \frac{y'}{y^p(y^p + p y')})$ , which modulo  $p^2$  is the same as  $(-x/y, -1/y, \frac{1}{y^{3p}}(x^p y' - y^p x')(y^p - p y'), \frac{1}{y^{3p}}(y')(y^p - p y'))$ . Then under the map  $s_U$ , this element is mapped to

$$\left( -x/y, -1/y, \frac{-x^p B^p(P_{U,0} + pP_{U,1}) + y^p A^p(P_{U,0} + pP_{U,1})}{y^p(y^p - pB^p(P_{U,0} + pP_{U,1}))}, \frac{-B^p(P_{U,0} + pP_{U,1})}{y^p(y^p - pB^p(P_{U,0} + pP_{U,1}))} \right)$$

which simplifies modulo  $p^2$  to

$$\left( -x/y, -1/y, \frac{(-x^p B^p + y^p A^p)(y^p P_{U,0} + p(B^p P_{U,0}^2 + y^p P_{U,1}))}{y^{3p}}, \frac{-B^p(y^p P_{U,0} + p(B^p P_{U,0}^2 + y^p P_{U,1}))}{y^{3p}} \right).$$

Under the map  $s_V$  the element  $(z, w, z', w')$  is mapped to

$$(z, w, -C^p(P_{V,0} + pP_{V,1}), -D^p(P_{V,0} + pP_{V,1})).$$

The image of the element  $(z, w, z', w')$  under the difference map  $s_U - s_V$  is the difference under the group law on  $E^1$  of the image of  $(z, w, z', w')$  under  $s_U$  and the image of  $(z, w, z', w')$  under  $s_V$ . In order to take the difference we must first take the inverse under the group law of the image of  $(z, w, z', w')$  under  $s_V$ , which is

$$(-z, -w, C^p(P_{V,0} + pP_{V,1}), D^p(P_{V,0} + pP_{V,1}))$$

and then add this to the image of  $(z, w, z', w')$  under  $s_U$ . Specifically we will let

$$\begin{aligned}
z_1 &= -x/y = z, \\
w_1 &= -1/y = w, \\
\delta z_1 &= \frac{(-x^p B^p + y^p A^p)(y^p P_{U,0} + p(B^p P_{U,0}^2 + y^p P_{U,1}))}{y^{3p}} \\
&= (w^p z^p B^p + w^p A^p)(-P_{U,0} + p(w^p B^p P_{U,0}^2 - P_{U,1})), \\
\delta w_1 &= \frac{-B^p(y^p P_{U,0} + p(B^p P_{U,0}^2 + y^p P_{U,1}))}{y^{3p}} \\
&= w^{2p} B^p(-P_{U,0} + p(w^p B^p P_{U,0}^2 - P_{U,1})), \\
z_2 &= -z, \\
w_2 &= -w, \\
\delta z_2 &= C^p(P_{V,0} + pP_{V,1}), \\
\delta w_2 &= D^p(P_{V,0} + pP_{V,1}),
\end{aligned}$$

and apply the explicit formulation of the group law detailed in Section 3. Also since for the purpose of our computation we only need the  $\delta z_3$  term, this is the only one we will formulate in detail.

We are going to be analyzing  $\frac{1}{\mu^{3p}}(-\mu^p \delta \alpha + \alpha^p \delta \mu)(\mu^p - p \delta \mu)$  with the above terms substituted in for  $z_1$ ,  $w_1$ , etc. When we do this,

$$\begin{aligned}
\alpha &= 0, \\
\mu &= 2w + 6a_6 w^3 + 6a_4 z w^2 + 6z^3,
\end{aligned}$$

which, if we then add  $6f(z, 1, w)$ , we have  $\mu = 8w = -8/y$ .

**Proposition 5.1.** *The  $\zeta = \delta z$  coordinate of  $s_U - s_V$  is*

$$\zeta = \frac{-\delta \alpha}{(8w)^p} + p \frac{\delta \alpha \delta \mu}{(8w)^{2p}},$$

where the above expressions are used for  $z_1$ ,  $z_2$ ,  $\delta z_1$ ,  $\delta z_2$ ,  $\delta w_1$ , etc. in  $\delta \alpha$  and  $\delta \mu$ .

The next step in the computation of  $f_{\text{def}}$  is to apply the formal logarithm of the Frobenius twist of the formal group of the elliptic curve to  $\zeta$ . This is a triviality as mentioned in the Introduction by the following proposition.

**Proposition 5.2.** *Let  $\log_{\mathcal{F}_1^{\phi^1}}(\xi)$  be the formal logarithm of the Frobenius twist of the formal group of the elliptic curve. Then*

$$\log_{\mathcal{F}_1^{\phi^1}}(\zeta) = \zeta \text{ modulo } p^2.$$

*Proof.* Recall

$$\log_{\mathcal{F}_1^{\phi^1}}(\xi) := \xi + \frac{p\phi(c_1)}{2}\xi^2 + \frac{p^2\phi(c_2)}{3}\xi^3 + \dots,$$

where the  $c_i$  are the coefficients of the power series expansion of the invariant differential [3, p. 127]. From [6, p. 113] we know that the invariant differential

$$\omega(z) = (1 + 2a_4 z^4 + \dots)dz$$

and so  $c_1 = 0$ ,  $c_2 = 0$ ,  $c_3 = 0$ ,  $c_4 = 2a_4 \dots$ . However, the power of  $p$  in  $p^n/n$  is at least 2 for all  $n \geq 4$ . Hence, modulo  $p^2$  the power series  $\log_{\mathcal{F}_1^{\phi^1}}(\xi)$  is the identity.  $\square$

## 6. RESIDUE OF THE COHOMOLOGY CLASS

Recall that any cohomology class in  $H^1(E \otimes M^1, \mathcal{O}) \simeq H^1(E, \mathcal{O}) \otimes M^1$  has a representative of the form  $\sum a_n y^n + x \sum b_n y^n + x^2 \sum e_n y^n$ . Let us refer to the coefficient  $e_{-1}$  in a sum  $\sum a_n y^n + x \sum b_n y^n + x^2 \sum e_n y^n$  as the residue of the sum. The final step in the computation of  $f_{\text{jet}}$  modulo  $p^2$  is to take the residue of  $\zeta = \delta z$  coordinate of  $s_U - s_V$ , which is a cohomology class as a result of the above proposition.

While the idea behind taking the residue is simple, namely write  $\zeta$  as  $\sum a_n y^n + x \sum b_n y^n + x^2 \sum e_n y^n$  and take the coefficient of  $x^2/y$  in this sum which is  $e_{-1}$ , the practice is computationally unfeasible. Instead we break the process of finding the residue of  $\zeta$  into parts. The residue map has some useful properties; namely, it is linear and the residue of any function that is regular on  $U$  or any function that is regular on  $V$  is zero. So we can take the residue of the terms in  $\zeta$  and then add together the result to get the residue of  $\zeta$ .

As a preliminary step to the task of analyzing the residue of the terms of  $\zeta = \delta z$ , we write the following expressions in both coordinates of  $U$  and coordinates of  $V$ :

$$\begin{aligned} A &= \frac{2^4(4a_4^2 + 6x^2a_4 - 9xa_6)}{\Delta} = \frac{2^4(4a_4^2w^2 + 6z^2a_4 - 9zwa_6)}{w^2\Delta}, \\ B &= \frac{2^3(9y)(2xa_4 - 3a_6)}{\Delta} = -\frac{2^3(9)(2za_4 - 3wa_6)}{w^2\Delta}, \\ C &= z\left(-\frac{3}{2}a_6w - a_4z\right) = \frac{x}{y^2}\left(-\frac{3}{2}a_6 - a_4x\right), \\ D &= -\frac{3}{2}a_6w^2 - wa_4z + 1 = \frac{1}{y^2}\left(x^3 - \frac{1}{2}a_6\right), \\ P_{U,0} &= \frac{P_{V,0}}{w^{3p}}. \end{aligned}$$

We also note that  $P_{U,0}$  and  $P_{U,1}$  are regular on  $U$ , and that  $P_{V,0}$ ,  $P_{V,1}$ , and  $C_p^{\text{ext}}(3a_6w^3 + 3a_6w^3 + 3z^3 + 3z^3 + a_4w^2z + a_4w^2z + w + w + 2a_4w^2z + 2a_4w^2z)$  are regular on  $V$ . So as an example the following combinations of taken from  $\frac{-\delta\alpha}{(8w)^p}$  are regular on  $V$ :

$$\frac{-p^3 D^p C^p a_6^p w^p P_{V,0}^2}{8^p}, \quad \frac{p^2 w^p a_4^p z^p C^p P_{V,1}}{8^p}, \quad \frac{-C^p P_{V,0}}{8^p}.$$

More examples of regular combinations on  $V$ , this time taken from  $\frac{\delta\alpha\delta\mu}{(8w)^{2p}}$ , are

$$\frac{(3^p w^{3p} C^p a_6^p P_{V,0})(3(3^p) a_6^p w^{4p} B^p P_{U,0})}{(8w)^{2p}} \quad \text{and} \quad \frac{(-3^p w^{4p} a_6^p A^p P_{U,0})(2(3^p) \delta(a_6) w^{3p})}{(8w)^{2p}}.$$

Since the residue of terms that are regular on either  $U$  or  $V$  is zero, we can exclude these terms from consideration in computing the residue class of  $\zeta$ . This leads to the following proposition in which for brevity's sake we let

$$\Upsilon = C_p^{\text{ext}}(y^2 - x^3 - a_4x - a_6).$$

**Proposition 6.1.** *The residue of  $\zeta$  is equal to the residue of*

$$\begin{aligned} & \left( \frac{(1-2^p)A^p}{y^p} + \frac{-2x^p B^p}{y^{2p}} + \frac{(-3^p a_6^p - 2a_4^p x^p)A^p}{y^{3p}} + \frac{(-1-2^p)x^p D^p}{y^{3p}} \right) \frac{P_{U,0}}{8^p} \\ & + p \left( F_1 C_p^{\text{ext}} (3a_6 w^3 + 3a_6 w^3 + 3z^3 + 3z^3 + a_4 w^2 z \right. \\ & \quad \left. + a_4 w^2 z + w + w + 2a_4 w^2 z + 2a_4 w^2 z) \right. \\ & \quad \left. + F_2 C_p^{\text{ext}} (A f_x + B f_y) + F_3 C_p^{\text{ext}} (-3x^2 - a_4) \right. \\ & \quad \left. + F_4 \Upsilon^2 + (F_5 + F_6 + F_7) \Upsilon + F_8 \right), \end{aligned}$$

where  $F_i$  are polynomials in  $M_1^1[x^p, y^p, \Upsilon]$ .

*Proof.* This is proved by the very precise removal of almost all regular terms using a computer algebra system.  $\square$

It is now necessary to compute residues of terms whose residue may be nontrivial. Namely, we provide a formula for the residue of  $\frac{x^a}{y^b}$  which we will call  $\gamma_{a,b}$ . We let  $\binom{n}{k}$  denote the binomial coefficient with the convention that  $\binom{n}{k} = 0$  if  $k > n$ . Then from [5] we know

**Proposition 6.2.** *Let  $a$  and  $b$  be positive integers. Let  $m$  and  $n \in \{0, 1, 2\}$  be integers such that  $a = 3m + n$ . Then the residue of  $\frac{x^a}{y^b}$  is*

$$\gamma_{a,b} = \begin{cases} 0 & \text{if } b \text{ is even,} \\ \sum_{k=0}^{\infty} \binom{m+k}{3k+2-n} \binom{m-2k-2+n}{\frac{b-1}{2}} (-1)^{m+k-\frac{b-1}{2}} (a_4)^{3k+2-n} (a_6)^{m-2k-2+n-\frac{b-1}{2}} & \text{if } b \text{ is odd.} \end{cases}$$

Obviously, because of the convention for binomial coefficients, there will be integers  $a$  and  $b$  with  $b$  odd for which  $\gamma_{a,b}$  is 0. In fact, if  $\frac{3b}{2} > a$ ,  $\gamma_{a,b} = 0$  because of the binomial coefficient  $\binom{m-2k-2+n}{\frac{b-1}{2}}$ . We now introduce a series of propositions that are just expanded formulas for expressions found in Proposition 6.1.

**Proposition 6.3.**

$$\begin{aligned} \Upsilon &= C_p^{\text{ext}} (y^2 - x^3 - a_4 x - a_6) \\ &= \frac{1}{p} \left[ \left( \sum_{k=1}^{p-1} \binom{p}{k} (-1)^k \right) y^{2p} - \sum_{k=1}^{p-1} \binom{p}{k} x^{3k} (a_4 x + a_6)^{p-k} - \sum_{k=1}^{p-1} \binom{p}{k} a_4^k a_6^{p-k} x^k \right]. \end{aligned}$$

**Proposition 6.4.**

$$\begin{aligned} A^p &= \frac{2^{4p}(4a_4^2 + 6x^2 a_4 - 9x a_6)^p}{\Delta^p} \\ &= \frac{2^{4p}}{\Delta^p} \left[ 4^p a_4^{2p} + 6^p x^{2p} a_4^p - 9^p x^p a_6^p \right. \\ & \quad \left. + \sum_{k=1}^{p-1} \binom{p}{k} (4a_4^2)^{p-k} (6x^2 a_4 - 9x a_6)^k + \sum_{k=1}^{p-1} \binom{p}{k} (6x^2 a_4)^k (-9x a_6)^{p-k} \right]. \end{aligned}$$

**Proposition 6.5.**

$$\begin{aligned} B^p &= \frac{2^{3p}(9y)^p(2xa_4 - 3a_6)^p}{\Delta^p} \\ &= \frac{2^{3p}(9y)^p}{\Delta^p} \left[ 2^p x^p a_4^p - 3^p a_6^p + \sum_{k=1}^{p-1} \binom{p}{k} (2xa_4)^k (-3a_6)^{p-k} \right]. \end{aligned}$$

**Proposition 6.6.**

$$\begin{aligned} C_p^{\text{ext}}(3a_6w^3 + 3a_6w^3 + 3z^3 + 3z^3 + a_4w^2z + a_4w^2z + w + w + 2a_4w^2z + 2a_4w^2z) \\ = \frac{1}{p} [2(3a_6)^p + 2(3x^3)^p + 2(1 + 2^p)(a_4x)^p + (2 - 8^p)y^{2p}] \left( \frac{-1}{y^{3p}} \right). \end{aligned}$$

**Proposition 6.7.**

$$\begin{aligned} C_p^{\text{ext}}(Af_x + Bf_y) \\ = \frac{1}{p} \sum_{k=1}^{p-1} \sum_{i=0}^k \sum_{j=0}^{p-i} \binom{p}{k} \binom{k}{i} \binom{p-i}{j} (-1)^{k-i} \\ \times \left( \frac{9(2^4)}{\Delta} \right)^{p-i} (2a_4)^j (-3a_6)^{p-i-j} x^j y^{2(p-i)}. \end{aligned}$$

**Proposition 6.8.**

$$C_p^{\text{ext}}(-3x^2 - a_4) = -\frac{1}{p} \sum_{k=1}^{p-1} \binom{p}{k} 3^k a_4^{p-k} x^{2k}.$$

Using this series of propositions, we can now explicitly write down the residue of  $\zeta$  by computing the residue of the formula in Proposition 6.1. First we introduce some more notation.

**Definition 6.9.** Define  $\mu_{a,b}$  to be the residue of  $\frac{x^a \Upsilon}{y^b}$  where

$$\Upsilon = C_p^{\text{ext}}(y^2 - x^3 - a_4x - a_6).$$

**Definition 6.10.** Define  $\tau_{a,b}$  to be the residue of  $\frac{x^a \Upsilon^2}{y^b}$  where

$$\Upsilon = C_p^{\text{ext}}(y^2 - x^3 - a_4x - a_6).$$

Using the formulas above, it is easy to check that for some, but not all, values,  $\mu_{a,b}$  will be zero. Similarly there are values of  $a$  and  $b$  for which  $\tau_{a,b}$  is nonzero and for which it is zero. Some examples of  $\mu_{a,b}$  are

$$\begin{aligned} \mu_{p,3p} &= 0, \\ \mu_{p,p} &= \frac{1}{p} \left[ -\sum_{k=1}^{p-1} \binom{p}{k} \sum_{i=0}^{p-k} \binom{p-k}{i} a_4^i a_6^{p-k-i} \gamma_{p+3k+i,p} - \sum_{k=1}^{p-1} \binom{p}{k} a_4^k a_6^{p-k} \gamma_{p+k,p} \right]. \end{aligned}$$

Note that both  $\gamma_{a,b}$  and  $\mu_{a,b}$  are in  $M_1^0$ . We can now prove the following theorem.

**Theorem 6.11.** *The reduction modulo  $p^2$  of  $f_{\text{jet}}$  is*

$$\begin{aligned} & \left[ \frac{9^p(2^p - 4^p - 2(3^p))a_6^p \delta(a_4)}{\Delta^p} + \frac{2^p(-6^p + 12^p + 2(9^p))a_4^p \delta(a_6)}{\Delta^p} \right] \gamma_{2p,p} \\ & + \frac{1}{\Delta^p} \left[ 2^p(1 - 2^p)4^p a_4^{2p} \mu_{0,p} + (-18^p(1 - 2^p) + 2(27^p))a_6^p \mu_{p,p} \right. \\ & + (12^p(1 - 2^p) - 2(18^p))a_4^p \mu_{2p,p} + (2(18^p) - 36^p)a_4^p a_6^p \mu_{2p,3p} - 2(12^p)a_4^{2p} \mu_{3p,3p} \left. \right] \\ & + p(H_0 + H_1 + H_2 + H_3 + H_4 + H_5 + H_6 + H_7 + H_8) \end{aligned}$$

where  $H_0$  is

$$\begin{aligned} & \frac{1}{p} \left( (1 - 2^p) \frac{2^p}{\Delta^p} \left[ -\delta(a_4)6^p a_4^p \gamma_{3p,p} + \sum_{k=1}^{p-1} \binom{p}{k} (4a_4^2)^{p-k} \sum_{i=0}^k \binom{k}{i} (6a_4)^i (-9a_6)^{k-i} \right. \right. \\ & \quad \times (-\delta(a_4)\gamma_{p+k+i,p} - \delta(a_6)\gamma_{k+i,p} + \mu_{k+i,p}) \\ & \quad \left. + \sum_{k=1}^{p-1} \binom{p}{k} (6a_4)^k (-9a_6)^{p-k} (-\delta(a_4)\gamma_{2p+k,p} - \delta(a_6)\gamma_{p+k,p} + \mu_{p+k,p}) \right] \\ & - 2 \frac{9^p}{\Delta^p} \left[ -\delta(a_4)2^p a_4^p \gamma_{3p,p} \right. \\ & \quad \left. + \sum_{k=1}^{p-1} \binom{p}{k} (2a_4)^k (-3a_6)^{p-k} (-\delta(a_4)\gamma_{2p+k,p} - \delta(a_6)\gamma_{p+k,p} + \mu_{p+k,p}) \right] \\ & - (3^p a_6^p) \frac{2^p}{\Delta^p} \left[ \sum_{k=1}^{p-1} \binom{p}{k} (4a_4^2)^{p-k} \sum_{i=0}^k \binom{k}{i} (6a_4)^i (-9a_6)^{k-i} \mu_{k+i,3p} \right. \\ & \quad \left. + \sum_{k=1}^{p-1} \binom{p}{k} (6a_4)^k (-9a_6)^{p-k} \mu_{p+k,3p} \right] \\ & - (2a_4^p) \frac{2^p}{\Delta^p} \left[ \sum_{k=1}^{p-1} \binom{p}{k} (4a_4^2)^{p-k} \sum_{i=0}^k \binom{k}{i} (6a_4)^i (-9a_6)^{k-i} \mu_{p+k+i,3p} \right. \\ & \quad \left. + \sum_{k=1}^{p-1} \binom{p}{k} (6a_4)^k (-9a_6)^{p-k} \mu_{2p+k,3p} \right] \Bigg), \end{aligned}$$

$H_1$  is

$$\begin{aligned} & \frac{-1}{p} \left( 9\delta(a_4) a_4^p (3^p) \gamma_{3p,p} \right. \\ & - \frac{9}{2} a_4^p \left( 2(3a_6)^p \mu_{2p,3p} + 2(3^p) \mu_{5p,3p} + 2(1+2^p) a_4^p \mu_{3p,3p} + (2-8^p) \mu_{2p,p} \right) \\ & + (9a_6^p - \frac{3}{2} a_4^p) \left( 2(3^p) \mu_{4p,3p} + 2(1+2^p) a_4^p \mu_{2p,3p} + (2-8^p) \mu_{p,p} \right) \\ & - a_4^{2p} \left( 2(3^p) \mu_{3p,3p} + (2-8^p) \mu_{0,p} \right) \\ & + \left( \left( \frac{3}{2} a_4^p - 9a_6^p \right) \delta(a_4) + \frac{9}{2} a_4^p \delta(a_6) \right) \left( 2(3^p) \gamma_{5p,3p} + (2-8^p) \gamma_{2p,p} \right) \\ & \quad \left. + \left( -3a_4^{2p} + \frac{9}{2} a_6^p a_4^p \right) \left( 2(3^p) \mu_{5p,5p} + (2-8^p) \mu_{2p,3p} \right) \right) / \Delta^p, \end{aligned}$$

$H_2$  is

$$\begin{aligned}
& \frac{1}{p} \sum_{k=1}^{p-1} \sum_{i=0}^k \sum_{j=0}^{p-i} \binom{p}{k} \binom{k}{i} \binom{p-i}{j} (-1)^{k-i} \left( \frac{9(2^4)}{\Delta} \right)^{p-i} (2a_4)^j (-3a_6)^{p-i-j} \\
& \times \left[ \left( (24a_4^{2p} - 36a_6^p a_4^p) \mu_{2p+j, 3p+2(p-i)} + (36a_6^p a_4^p + 16a_4^{3p} - 54a_6^{2p}) \mu_{p+j, 3p+2(p-i)} \right. \right. \\
& \quad + 24a_6^p a_4^{2p} \mu_{j, 3p+2(p-i)} + 36a_4^p \mu_{2p+j, p+2(p-i)} \\
& \quad \left. \left. + (12a_4^p - 72a_6^p) \mu_{p+j, p+2(p-i)} + 8a_4^{2p} \mu_{j, p+2(p-i)} \right) / \Delta^p \right. \\
& + \left( 36a_4^p \delta(a_4) \gamma_{3p+j, p+2(p-i)} \right. \\
& \quad + ((72a_6^p - 12a_4^p) \delta(a_4) - 36a_4^p \delta(a_6)) \gamma_{2p+j, p+2(p-i)} \\
& \quad + (28a_4^{2p} \delta(a_4) + (72a_6^p - 12a_4^p) \delta(a_6)) \gamma_{p+j, p+2(p-i)} \\
& \quad \left. \left. + ((72a_4^p a_6^p - 24a_4^{2p}) \delta(a_4) - 8a_4^{2p} \delta(a_6)) \gamma_{j, p+2(p-i)} \right) / \Delta^p \right. \\
& + \left( ((54a_6^{2p} - 16a_4^{3p} - 36a_4^p a_6^p) \delta(a_4) + (36a_4^p a_6^p - 24a_4^{2p}) \delta(a_6)) \gamma_{2p+j, 3p+2(p-i)} \right. \\
& \quad + ((24a_4^{3p} - 60a_4^{2p} a_6^p) \delta(a_4) \\
& \quad \left. \left. + (54a_6^{2p} - 16a_4^{3p} - 36a_4^p a_6^p) \delta(a_6)) \gamma_{p+j, 3p+2(p-i)} \right) / \Delta^p \right],
\end{aligned}$$

$H_3$  is

$$\begin{aligned}
& -\frac{1}{p} \sum_{k=1}^{p-1} \binom{p}{k} 3^k a_4^{p-k} \\
& \times \left( \left( (2304a_4^{3p} + 10368a_6^{2p} - 8640a_6^p a_4^p + 1152a_4^{2p}) \mu_{2p+2k, p} \right. \right. \\
& \quad + (-1920a_4^{3p} - 576a_6^p a_4^{2p}) \mu_{p+2k, p} \\
& \quad + (512a_4^{4p} + 10368a_6^{2p} a_4^p - 10368a_6^p a_4^{2p} + 2304a_4^{3p}) \mu_{2k, p} \\
& \quad + (-10368a_6^{2p} a_4^p + 3456a_6^p a_4^{2p} \\
& \quad \quad + 3072a_4^{4p} + 7776a_6^{3p} - 4608a_6^p a_4^{3p}) \mu_{2p+2k, 3p} \\
& \quad + (11520a_6^p a_4^{3p} + 1024a_4^{5p} - 12096a_6^{2p} a_4^{2p} - 2304a_4^{4p}) \mu_{p+2k, 3p} \\
& \quad \left. \left. + (-5184a_6^{3p} a_4^p - 2304a_6^p a_4^{3p} + 1536a_6^p a_4^{4p} + 6912a_6^{2p} a_4^{2p}) \mu_{2k, 3p} \right) / \Delta^{2p} \right. \\
& + \left( ((1920a_4^{3p} + 576a_6^p a_4^{2p}) \delta(a_4) \right. \\
& \quad + (8640a_6^p a_4^p - 1152a_4^{2p} - 2304a_4^{3p} - 10368a_6^{2p}) \delta(a_6)) \gamma_{2p+2k, p} \\
& \quad + ((1792a_4^{4p} - 1152a_4^{3p} + 1728a_6^p a_4^{2p}) \delta(a_4) \\
& \quad \quad + (1920a_4^{3p} + 576a_6^p a_4^{2p}) \delta(a_6)) \gamma_{p+2k, p} \\
& \quad + ((6912a_6^p a_4^{3p} + 1728a_6^{2p} a_4^p \\
& \quad \quad + 2592a_6^{3p} - 3072a_4^{4p} - 2304a_6^p a_4^{2p}) \delta(a_4) \\
& \quad \left. \left. + (-512a_4^{4p} - 10368a_6^{2p} a_4^p + 10368a_6^p a_4^{2p} - 2304a_4^{3p}) \delta(a_6)) \gamma_{2k, p} \right) / \Delta^{2p} \right),
\end{aligned}$$



$H_4$  is

$$\begin{aligned}
& \left( -\frac{1}{2} \left( -3732480 a_6^{2p} a_4^{2p} - 829440 a_6^p a_4^{4p} \right. \right. \\
& \quad + 3359232 a_6^{3p} a_4^p + 3456 a_4^{2p} \Delta^p + 774144 a_4^{5p} \\
& \quad + 1990656 a_6^p a_4^{3p} - 12960 a_4^p a_6^p \Delta^p - 663552 a_4^{4p} \left. \right) \tau_{2p,p} \\
& - \frac{1}{2} \left( 2592 a_4^{2p} \Delta^p - 3317760 a_4^{5p} - 7713792 a_6^{2p} a_4^{3p} \right. \\
& \quad + 19440 a_6^{2p} \Delta^p + 9123840 a_6^p a_4^{4p} - 7278336 a_6^{4p} \\
& \quad + 2985984 a_6^{2p} a_4^{2p} - 1327104 a_6^p a_4^{3p} + 3359232 a_6^{3p} a_4^p \\
& \quad + 24576 a_4^{6p} - 12960 a_4^p a_6^p \Delta^p - 288 a_4^{3p} \Delta^p \left. \right) \tau_{p,p} \\
& - \frac{1}{2} \left( -1769472 a_4^{6p} - 8957952 a_6^{3p} a_4^{2p} - 11232 a_4^{2p} a_6^p \Delta^p \right. \\
& \quad + 2875392 a_4^{5p} a_6^p + 3456 a_4^{3p} \Delta^p + 1327104 a_4^{5p} \\
& \quad \left. + 20404224 a_6^{2p} a_4^{3p} - 10616832 a_6^p a_4^{4p} \right) \tau_{0,p} \Big) / (\Delta^{3p}) \\
& + \left( -\frac{1}{2} \left( -393216 a_4^{7p} + 22394880 a_6^{3p} a_4^{2p} - 3456 a_4^{2p} a_6^p \Delta^p \right. \right. \\
& \quad - 11197440 a_6^{4p} a_4^p - 10616832 a_4^{5p} a_6^p + 2592 a_4^p a_6^{2p} \Delta^p \\
& \quad + 3317760 a_6^p a_4^{4p} + 1152 a_4^{4p} \Delta^p + 2654208 a_4^{6p} \\
& \quad - 14929920 a_6^{2p} a_4^{3p} + 9953280 a_6^{2p} a_4^{4p} + 1152 a_4^{3p} \Delta^p \left. \right) \tau_{2p,3p} \\
& - \frac{1}{2} \left( 12607488 a_4^{5p} a_6^p - 8957952 a_6^{3p} a_4^{2p} - 3244032 a_6^p a_4^{6p} \right. \\
& \quad + 19408896 a_6^{3p} a_4^{3p} + 1536 a_4^{4p} \Delta^p + 13436928 a_6^{4p} a_4^p \\
& \quad - 28864512 a_6^{2p} a_4^{4p} + 5184 a_4^{2p} a_6^p \Delta^p + 11664 a_6^{3p} \Delta^p \\
& \quad - 1327104 a_4^{6p} - 15552 a_4^p a_6^{2p} \Delta^p + 1769472 a_4^{7p} \\
& \quad - 2304 a_4^{3p} a_6^p \Delta^p + 1990656 a_6^{2p} a_4^{3p} - 6718464 a_6^{5p} \left. \right) \tau_{p,3p} \\
& - \frac{1}{2} \left( -20901888 a_6^{3p} a_4^{3p} - 18144 a_4^{2p} a_6^{2p} \Delta^p - 3456 a_4^{4p} \Delta^p \right. \\
& \quad + 9953280 a_6^{2p} a_4^{4p} + 13436928 a_6^{4p} a_4^{2p} + 17280 a_4^{3p} a_6^p \Delta^p \\
& \quad + 1769472 a_6^p a_4^{6p} + 1664 a_4^{5p} \Delta^p \\
& \quad \left. - 2654208 a_6^{2p} a_4^{5p} - 1327104 a_4^{5p} a_6^p \right) \tau_{0,3p} \Big) / (\Delta^{3p}) \\
& + \left( -\frac{1}{2} \left( -10368 a_4^{3p} a_6^p \Delta^p + 31104 a_4^{2p} a_6^{2p} \Delta^p - 54 \Delta^{2p} a_4^p \right. \right. \\
& \quad \left. + 10368 \Delta^p a_6^p a_4^{4p} - 23328 \Delta^p a_6^{3p} a_4^p - 4608 a_4^{5p} \Delta^p \right) \tau_{2p,5p} \Big) / (\Delta^{3p}),
\end{aligned}$$

$H_5$  is

$$\begin{aligned} & \left( \left( \frac{1}{2} \left( -14556672 a_6^{4p} - 6635520 a_4^{5p} + 6718464 a_6^{3p} a_4^p + 5971968 a_6^{2p} a_4^{2p} \right. \right. \right. \\ & \quad + 2880 a_4^{2p} \Delta^p + 18144 a_6^{2p} \Delta^p - 5184 a_4^{3p} \Delta^p - 15427584 a_6^{2p} a_4^{3p} \\ & \quad - 2654208 a_6^p a_4^{3p} - 8640 a_4^p a_6^p \Delta^p + 49152 a_4^{6p} + 18247680 a_6^p a_4^{4p} \Big) \delta(a_4) \\ & + \frac{1}{2} \left( 1548288 a_4^{5p} + 6912 a_4^{2p} \Delta^p - 25920 a_4^p a_6^p \Delta^p - 1658880 a_6^p a_4^{4p} \right. \\ & \quad + 6718464 a_6^{3p} a_4^p - 7464960 a_6^{2p} a_4^{2p} + 3981312 a_6^p a_4^{3p} - 1327104 a_4^{4p} \Big) \delta(a_6) \\ & + \frac{1}{2} \left( 3584 a_4^{4p} \delta(3) \Delta^p + 3456 \delta(3) a_4^{2p} a_6^p \Delta^p + 4608 \delta(2) a_4^{4p} \Delta^p \right. \\ & \quad \left. \left. + 18 \delta(3) a_4^p \Delta^{2p} - 2304 a_4^{3p} \delta(3) \Delta^p \right) \mu_{2p,p} \right) / (\Delta^{3p}), \end{aligned}$$

$H_6$  is

$$\begin{aligned} & \left( \left( \frac{1}{2} \left( 3840 a_4^{3p} \Delta^p - 25214976 a_6^p a_4^{4p} \right. \right. \right. \\ & \quad + 3981312 a_4^{5p} + 7409664 a_4^{5p} a_6^p + 4608 a_4^{2p} a_6^p \Delta^p \\ & \quad - 5087232 a_4^{6p} + 48273408 a_6^{2p} a_4^{3p} - 24634368 a_6^{3p} a_4^{2p} \Big) \delta(a_4) \\ & + \frac{1}{2} \left( -14556672 a_6^{4p} - 15427584 a_6^{2p} a_4^{3p} - 25920 a_4^p a_6^p \Delta^p \right. \\ & \quad - 2654208 a_6^p a_4^{3p} - 6635520 a_4^{5p} + 38880 a_6^{2p} \Delta^p \\ & \quad + 18247680 a_6^p a_4^{4p} + 5184 a_4^{2p} \Delta^p + 6718464 a_6^{3p} a_4^p \\ & \quad \left. + 5971968 a_6^{2p} a_4^{2p} - 576 a_4^{3p} \Delta^p + 49152 a_4^{6p} \right) \delta(a_6) \\ & + \frac{1}{2} \left( 5184 \delta(3) a_6^{3p} \Delta^p - 24 \delta(2) a_4^p \Delta^{2p} - 36 \delta(3) a_6^p \Delta^{2p} + 23328 \delta(2) a_6^{3p} \Delta^p \right. \\ & \quad + 36 \delta(2) a_6^p \Delta^{2p} - 6912 \delta(2) a_4^{4p} \Delta^p + 12672 a_4^{3p} \delta(3) a_6^p \Delta^p \\ & \quad - 15552 \delta(2) a_6^{2p} a_4^p \Delta^p + 3456 \delta(3) a_6^{2p} a_4^p \Delta^p + 10368 \delta(2) a_4^{3p} a_6^p \Delta^p \\ & \quad \left. - 4608 \delta(3) a_4^{2p} a_6^p \Delta^p - 9984 a_4^{4p} \delta(3) \Delta^p + 6 \delta(3) a_4^p \Delta^{2p} \right) \mu_{p,p} \\ & + \left( \frac{1}{2} \left( 7962624 a_6^p a_4^{4p} - 33841152 a_6^{2p} a_4^{3p} - 22781952 a_4^{5p} a_6^p \right. \right. \\ & \quad + 5308416 a_4^{6p} + 52254720 a_6^{3p} a_4^{2p} - 29113344 a_6^{4p} a_4^p \\ & \quad + 54 \Delta^{2p} a_4^p - 786432 a_4^{7p} + 6912 a_4^{2p} a_6^p \Delta^p + 10368 a_4^p a_6^{2p} \Delta^p \\ & \quad \left. - 2304 a_4^{3p} \Delta^p + 1280 a_4^{4p} \Delta^p + 21565440 a_6^{2p} a_4^{4p} \right) \delta(a_4) \\ & + \frac{1}{2} \left( 40808448 a_6^{2p} a_4^{3p} - 22464 a_4^{2p} a_6^p \Delta^p \right. \\ & \quad - 17915904 a_6^{3p} a_4^{2p} + 5750784 a_4^{5p} a_6^p + 2654208 a_4^{5p} \\ & \quad \left. - 21233664 a_6^p a_4^{4p} - 3538944 a_4^{6p} + 6912 a_4^{3p} \Delta^p \right) \delta(a_6) \\ & + \frac{1}{2} \left( -6912 \delta(2) a_4^{3p} a_6^p \Delta^p - 2048 a_4^{5p} \delta(3) \Delta^p \right. \\ & \quad + 4608 a_4^{4p} \delta(3) \Delta^p - 14 \delta(3) a_4^{2p} \Delta^{2p} + 23040 \delta(3) a_6^{2p} a_4^{2p} \Delta^p \\ & \quad \left. - 26880 a_4^{3p} \delta(3) a_6^p \Delta^p + 2 \delta(2) a_4^{2p} \Delta^{2p} \right) \mu_{0,p} \Big) / (\Delta^{3p}), \end{aligned}$$

$H_7$  is

$$\begin{aligned}
& \left( \frac{1}{2} \left( -17915904 a_6^{3p} a_4^{2p} + 25214976 a_4^{5p} a_6^p + 7776 a_6^{3p} \Delta^p \right. \right. \\
& \quad + 38817792 a_6^{3p} a_4^{3p} + 4608 a_4^{3p} a_6^p \Delta^p + 3456 a_4^{2p} a_6^p \Delta^p \\
& \quad + 3981312 a_6^{2p} a_4^{3p} + 26873856 a_6^{4p} a_4^p - 2654208 a_4^{6p} \\
& \quad - 13436928 a_6^{5p} - 30 \Delta^{2p} a_4^p - 6488064 a_6^p a_4^{6p} + 3538944 a_4^{7p} \\
& \quad \left. - 36 \Delta^{2p} a_6^p - 10368 a_4^p a_6^{2p} \Delta^p - 3072 a_4^{4p} \Delta^p - 57729024 a_6^{2p} a_4^{4p} \right) \delta(a_4) \\
& + \frac{1}{2} \left( 54 \Delta^{2p} a_4^p + 5184 a_4^p a_6^{2p} \Delta^p - 21233664 a_4^{5p} a_6^p \right. \\
& \quad + 19906560 a_6^{2p} a_4^{4p} + 2304 a_4^{3p} \Delta^p + 6635520 a_6^p a_4^{4p} \\
& \quad - 29859840 a_6^{2p} a_4^{3p} + 2304 a_4^{4p} \Delta^p - 6912 a_4^{2p} a_6^p \Delta^p \\
& \quad + 44789760 a_6^{3p} a_4^{2p} + 5308416 a_4^{6p} - 22394880 a_6^{4p} a_4^p - 786432 a_4^{7p} \left. \right) \delta(a_6) \\
& + \frac{1}{2} \left( 25920 a_6^{3p} \delta(3) a_4^p \Delta^p + 6144 a_4^{5p} \delta(3) \Delta^p - 36 \delta(2) a_4^p a_6^p \Delta^{2p} \right. \\
& \quad + 11520 a_4^{3p} \delta(3) a_6^p \Delta^p + 6 \delta(2) a_4^{2p} \Delta^{2p} + 6 \delta(3) a_4^{2p} \Delta^{2p} \\
& \quad \left. + 18 \delta(3) a_6^p a_4^p \Delta^{2p} - 34560 \delta(3) a_6^{2p} a_4^{2p} \Delta^p - 12288 a_4^{4p} \delta(3) a_6^p \Delta^p \right) \mu_{2p,3p} / (\Delta^{3p}),
\end{aligned}$$

and  $H_8$  is

$$\begin{aligned}
& \left( - \left( -1271808 a_4^{6p} + 995328 a_4^{5p} + 12068352 a_6^{2p} a_4^{3p} \right. \right. \\
& \quad - 6158592 a_6^{3p} a_4^{2p} + 1852416 a_4^{5p} a_6^p \\
& \quad - 6303744 a_6^p a_4^{4p} + 1920 a_4^{3p} \Delta^p + 1440 a_4^{2p} a_6^p \Delta^p \left. \right) \delta(a_4)^2 \\
& - \left( 9123840 a_6^p a_4^{4p} - 7278336 a_6^{4p} + 24576 a_4^{6p} \right. \\
& \quad - 3317760 a_4^{5p} - 7713792 a_6^{2p} a_4^{3p} - 2592 a_4^{3p} \Delta^p \\
& \quad + 9072 a_6^{2p} \Delta^p - 4320 a_4^p a_6^p \Delta^p - 1327104 a_6^p a_4^{3p} \\
& \quad + 1440 a_4^{2p} \Delta^p + 2985984 a_6^{2p} a_4^{2p} + 3359232 a_6^{3p} a_4^p \left. \right) \delta(a_4) \delta(a_6) \\
& - \left( 387072 a_4^{5p} + 1728 a_4^{2p} \Delta^p - 1866240 a_6^{2p} a_4^{2p} \right. \\
& \quad + 995328 a_6^p a_4^{3p} - 331776 a_4^{4p} + 1679616 a_6^{3p} a_4^p \\
& \quad \left. - 6480 a_4^p a_6^p \Delta^p - 414720 a_6^p a_4^{4p} \right) \delta(a_6)^2 \\
& - \left( 18 \delta(2) a_6^p \Delta^{2p} - 18 \delta(3) a_6^p \Delta^{2p} + 1728 \delta(3) a_6^{2p} a_4^p \Delta^p \right. \\
& \quad - 7776 \delta(2) a_6^{2p} a_4^p \Delta^p + 6336 a_4^{3p} \delta(3) a_6^p \Delta^p \\
& \quad - 2304 \delta(3) a_4^{2p} a_6^p \Delta^p + 5184 \delta(2) a_4^{3p} a_6^p \Delta^p \\
& \quad + 3 \delta(3) a_4^p \Delta^{2p} - 12 \delta(2) a_4^p \Delta^{2p} - 3456 \delta(2) a_4^{4p} \Delta^p \\
& \quad + 11664 \delta(2) a_6^{3p} \Delta^p + 2592 \delta(3) a_6^{3p} \Delta^p - 4992 a_4^{4p} \delta(3) \Delta^p \left. \right) \delta(a_4) \\
& - \left( 1792 a_4^{4p} \delta(3) \Delta^p - 1152 a_4^{3p} \delta(3) \Delta^p + 1728 \delta(3) a_4^{2p} a_6^p \Delta^p \right. \\
& \quad \left. + 9 \delta(3) a_4^p \Delta^{2p} + 2304 \delta(2) a_4^{4p} \Delta^p \right) \delta(a_6) \Big) \frac{\gamma_{2p,p}}{\Delta^{3p}}.
\end{aligned}$$

*Proof.* We simply apply the most recent propositions to the actual formulas from Proposition 6.1. It should be noted that the  $H_i$  correspond to the  $F_i$  in Proposition 6.1 and that upon further analysis certain terms like  $\frac{(-1-2^p)x^p D^p}{y^{3p}}$  have zero residue even though it is not immediately obvious that the term is regular.  $\square$

From now on, when we refer to  $H_0, H_1, H_2, H_3, H_4, H_5, H_6, H_7$ , and  $H_8$  we will mean the polynomials in this theorem. We note that  $H_0, H_1, H_2, H_3, H_5, H_6, H_7$  are in  $M_1^1$  and are linear in  $\delta(a_4)$  and  $\delta(a_6)$ ,  $H_4 \in M_1^0$ , and  $H_8 \in M_1^1$  is quadratic in  $\delta(a_4)$  and  $\delta(a_6)$ .

## 7. ORDER TWO MODULAR FORMS

We remind ourselves that  $\phi$ , the unique lifting of the Frobenius morphism to  $R$ , extends to a homomorphism from  $M_1^1 \rightarrow M_1^2$  by taking, e.g.,  $a_4 \mapsto a_4^p + p\delta(a_4)$  and  $\delta(a_6) \mapsto \delta(a_6)^p + p\delta(a_6)$ . Hence if we start with a polynomial in  $M^0$  like  $\gamma_{a,b}$ , then  $\phi(\gamma_{a,b}) = \gamma_{a,b}(a_4^p + p\delta(a_4), a_6^p + p\delta(a_6)) \in M^1$ , where by this notation we mean substitute  $a_4^p + p\delta(a_4)$  in for  $a_4$  and  $a_6^p + p\delta(a_6)$  in for  $a_6$ .

**Definition 7.1.** Let  $\tilde{\gamma}_{a,b}$  be the polynomial in  $M^1$  such that  $\phi(\gamma_{a,b}) = \gamma_{a,b}(a_4^p, a_6^p) + p\tilde{\gamma}_{a,b}$ .

An explicit formula for  $\tilde{\gamma}_{a,b}$  is simple to compute by expanding the formula

$$\phi(\gamma_{a,b}) = \begin{cases} 0 & \text{if } b \text{ is even,} \\ \sum_{k=0}^{\infty} \binom{m+k}{3k+2-n} \binom{m-2k-2+n}{\frac{b-1}{2}} & \\ \quad (-1)^{m+k-\frac{b-1}{2}} (a_4^p + p\delta(a_4))^{3k+2-n} (a_6^p + p\delta(a_6))^{m-2k-2+n-\frac{b-1}{2}} & \text{if } b \text{ is odd.} \end{cases}$$

and modulo  $p^2$ ,  $\tilde{\gamma}_{a,b}$  is linear in  $\delta(a_4), \delta(a_6)$ . In addition  $\phi(\gamma_{a,b}) = \gamma_{a,b}^p + p\delta(\gamma_{a,b})$ ; however, note that  $\tilde{\gamma}_{a,b}$  does not equal  $p\delta(\gamma_{a,b})$  because the latter is missing the terms from  $\gamma_{a,b}^p$  whose coefficients are divisible by  $p$ .

**Definition 7.2.** Let  $\tilde{\mu}_{a,b}$  be the polynomial in  $M^1$  such that  $\phi(\mu_{a,b}) = \mu_{a,b}(a_4^p, a_6^p) + p\tilde{\mu}_{a,b}$ .

We recall that the isogeny covariant differential modular form  $f_{\text{jet}} h_{\text{jet}}$  is  $\phi(f_{\text{jet}})$ .

**Theorem 7.3.** The reduction modulo  $p^2$  of  $f_{\text{jet}} h_{\text{jet}}$  is

$$\begin{aligned} & \left[ \frac{-72a_6^{p^2} \delta(a_4)^p + 48a_4^{p^2} \delta(a_6)^p}{\Delta^{p^2}} \right] \gamma_{2p,p}(a_4^p, a_6^p) \\ & + \frac{1}{\Delta^{p^2}} \left[ -8a_4^{2p^2} \mu_{0,p}(a_4^p, a_6^p) + 72a_6^{p^2} \mu_{p,p}(a_4^p, a_6^p) \right. \\ & \quad \left. - 48a_4^{p^2} \mu_{2p,p}(a_4^p, a_6^p) - 24a_4^{2p^2} \mu_{3p,3p}(a_4^p, a_6^p) \right] \\ & + p \left[ \frac{-72a_6^{p^2} \delta^2(a_4) + 48a_4^{p^2} \delta^2(a_6)}{\Delta^{p^2}} \right] \gamma_{2p,p}(a_4^p, a_6^p) + pJ_0, \end{aligned}$$

where  $J_0$  is

$$\begin{aligned} & \left( \frac{1}{\Delta^{p^2}} \left[ (-72a_6^{p^2} \delta(a_4)^p + 48a_4^{p^2} \delta(a_6)^p) \tilde{\gamma}_{2p,p} - 8a_4^{2p^2} \tilde{\mu}_{0,p} \right. \right. \\ & \quad + 72a_6^{p^2} \tilde{\mu}_{p,p} - 48a_4^{p^2} \tilde{\mu}_{2p,p} - 24a_4^{2p^2} \tilde{\mu}_{3p,3p} \\ & \quad + ((27\delta(2) + 66\delta(3))a_6^{p^2} \delta(a_4)^p \\ & \quad + (-28\delta(3) - 42\delta(2))a_4^{p^2} \delta(a_6)^p) \gamma_{2p,p}^p + 20\delta(2)a_4^{2p^2} \mu_{0,p}^p \\ & \quad + (-27\delta(2) - 66\delta(3))a_6^{p^2} \mu_{p,p}^p + (42\delta(2) + 28\delta(3))a_4^{p^2} \mu_{2p,p}^p \\ & \quad \left. + 18\delta(2)a_4^{p^2} a_6^{p^2} \mu_{2p,3p}^p + (8\delta(3) + 24\delta(2))a_4^{2p^2} \mu_{3p,3p}^p \right] \\ & \quad \left. + H_0^p + H_1^p + H_2^p + H_3^p + H_4^p + H_5^p + H_6^p + H_7^p + H_8^p \right). \end{aligned}$$

*Proof.* Let  $H_i$  be the polynomials from Theorem 6.11. The formula follows immediately from the fact that  $f_{\text{jet}} h_{\text{jet}} = \phi(f_{\text{jet}})$ .  $\square$

Next working modulo  $p^2$ ,  $h_{\text{jet}} = (f_{\text{jet}} h_{\text{jet}})/f_{\text{jet}}$  is  $\frac{h_0}{f_0} + p \left( \frac{-h_0 f_1}{f_0^2} + \frac{h_1}{f_0} \right)$ , where  $f_0$  is the coefficient of  $p^0$  in  $f_{\text{jet}}$ ,  $f_1$  is the coefficient of  $p$ ,  $h_0$  is the coefficient of  $p^0$  in  $f_{\text{jet}} h_{\text{jet}}$ , and  $h_1$  is the coefficient of  $p$ . In particular

$$\begin{aligned} f_0 &= \left[ \frac{-72a_6^p \delta(a_4) + 48a_4^p \delta(a_6)}{\Delta^p} \right] \gamma_{2p,p} \\ & \quad + \frac{1}{\Delta^p} \left[ -8a_4^{2p} \mu_{0,p} + 72a_6^p \mu_{p,p} - 48a_4^p \mu_{2p,p} - 24a_4^{2p} \mu_{3p,3p} \right], \\ f_1 &= \frac{1}{\Delta^p} \left[ ((27\delta(2) + 66\delta(3))a_6^p \delta(a_4) \right. \\ & \quad + (-28\delta(3) - 42\delta(2))a_4^p \delta(a_6)) \gamma_{2p,p} + 20\delta(2)a_4^{2p} \mu_{0,p} \\ & \quad + (-27\delta(2) - 66\delta(3))a_6^p \mu_{p,p} + (42\delta(2) + 28\delta(3))a_4^p \mu_{2p,p} \\ & \quad \left. + 18\delta(2)a_4^p a_6^p \mu_{2p,3p} + (8\delta(3) + 24\delta(2))a_4^{2p} \mu_{3p,3p} \right] \\ & \quad + (H_0 + H_1 + H_2 + H_3 + H_4 + H_5 + H_6 + H_7 + H_8), \\ h_0 &= \left[ \frac{-72a_6^{p^2} \delta(a_4)^p + 48a_4^{p^2} \delta(a_6)^p}{\Delta^{p^2}} \right] \gamma_{2p,p}(a_4^p, a_6^p) \\ & \quad + \frac{1}{\Delta^{p^2}} \left[ -8a_4^{2p^2} \mu_{0,p}(a_4^p, a_6^p) + 72a_6^{p^2} \mu_{p,p}(a_4^p, a_6^p) \right. \\ & \quad \left. - 48a_4^{p^2} \mu_{2p,p}(a_4^p, a_6^p) - 24a_4^{2p^2} \mu_{3p,3p}(a_4^p, a_6^p) \right], \\ h_1 &= \left[ \frac{-72a_6^{p^2} \delta^2(a_4) + 48a_4^{p^2} \delta^2(a_6)}{\Delta^{p^2}} \right] \gamma_{2p,p}(a_4^p, a_6^p) + J_0. \end{aligned}$$

Therefore we have the following explicit formulation for  $h_{\text{jet}}$  where  $J_0$  is the polynomial from Theorem 7.3.

**Theorem 7.4.** *The reduction modulo  $p^2$  of  $h_{\text{jet}}$  is*

$$\begin{aligned} & \left( (-72a_6^{p^2}\delta(a_4)^p + 48a_4^{p^2}\delta(a_6)^p)\gamma_{2p,p}(a_4^p, a_6^p) - 8a_4^{2p^2}\mu_{0,p}(a_4^p, a_6^p) \right. \\ & \quad \left. + 72a_6^{p^2}\mu_{p,p}(a_4^p, a_6^p) - 48a_4^{p^2}\mu_{2p,p}(a_4^p, a_6^p) - 24a_4^{2p^2}\mu_{3p,3p}(a_4^p, a_6^p) \right) / \\ & \quad \left( \Delta^{p^2-p} \left( (-72a_6^p\delta(a_4) + 48a_4^p\delta(a_6))\gamma_{2p,p} \right. \right. \\ & \quad \left. \left. - 8a_4^{2p}\mu_{0,p} + 72a_6^p\mu_{p,p} - 48a_4^p\mu_{2p,p} - 24a_4^{2p}\mu_{3p,3p} \right) \right) + pK_0 \\ & + \frac{p(-72a_6^{p^2}\delta^2(a_4) + 48a_4^{p^2}\delta^2(a_6))\gamma_{2p,p}(a_4^p, a_6^p)}{\Delta^{p^2-p}((-72a_6^p\delta(a_4) + 48a_4^p\delta(a_6))\gamma_{2p,p} - 8a_4^{2p}\mu_{0,p} + 72a_6^p\mu_{p,p} - 48a_4^p\mu_{2p,p} - 24a_4^{2p}\mu_{3p,3p})}, \end{aligned}$$

where  $K_0$  is

$$\begin{aligned} & \frac{\Delta^p J_0}{\left( (-72a_6^p\delta(a_4) + 48a_4^p\delta(a_6))\gamma_{2p,p} - 8a_4^{2p}\mu_{0,p} + 72a_6^p\mu_{p,p} - 48a_4^p\mu_{2p,p} - 24a_4^{2p}\mu_{3p,3p} \right)} \\ & - \left( (-72a_6^{p^2}\delta(a_4)^p + 48a_4^{p^2}\delta(a_6)^p)\gamma_{2p,p}(a_4^p, a_6^p) - 8a_4^{2p^2}\mu_{0,p}(a_4^p, a_6^p) \right. \\ & \quad \left. + 72a_6^{p^2}\mu_{p,p}(a_4^p, a_6^p) - 48a_4^{p^2}\mu_{2p,p}(a_4^p, a_6^p) - 24a_4^{2p^2}\mu_{3p,3p}(a_4^p, a_6^p) \right) \\ & \times \left( \frac{1}{\Delta^p} \left[ ((27\delta(2) + 66\delta(3))a_6^p\delta(a_4) + (-28\delta(3) - 42\delta(2))a_4^p\delta(a_6))\gamma_{2p,p} \right. \right. \\ & \quad \left. \left. + 20\delta(2)a_4^{2p}\mu_{0,p} + (-27\delta(2) - 66\delta(3))a_6^p\mu_{p,p} + (42\delta(2) + 28\delta(3))a_4^p\mu_{2p,p} \right. \right. \\ & \quad \left. \left. + 18\delta(2)a_4^p a_6^p \mu_{2p,3p} + (8\delta(3) + 24\delta(2))a_4^{2p}\mu_{3p,3p} \right] + \sum_{i=0}^8 H_i \right) / \\ & \left( \left( (-72a_6^p\delta(a_4) + 48a_4^p\delta(a_6))\gamma_{2p,p} \right. \right. \\ & \quad \left. \left. - 8a_4^{2p}\mu_{0,p} + 72a_6^p\mu_{p,p} - 48a_4^p\mu_{2p,p} - 24a_4^{2p}\mu_{3p,3p} \right)^2 \Delta^{p^2-2p} \right). \end{aligned}$$

#### REFERENCES

1. M. Barcan and A. Buium, *Siegel Differential Modular Forms*, Int. Math. Res. Not. (2002), no. 28, 1457–1503. MR1908022 (2003g:11044)
2. A. Buium, *Geometry of Fermat Adeles*, Preprint, 1999.
3. ———, *Differential Modular Forms*, J. Reine Angew. Math. (2000), no. 520, 95–167. MR1748272 (2002d:11042)
4. ———, *Arithmetic Differential Invariants*, In preparation, 2003.
5. C. Hurlburt, *Isogeny Covariant Differential Modular Forms modulo  $p$* , Compositio Mathematica **128** (2001), no. 1, 17–34. MR1847663 (2002i:11053)
6. J. Silverman, *The Arithmetic of Elliptic Curves*, Graduate Texts in Mathematics, vol. 106, Springer Verlag, 1986. MR0817210 (87g:11070)

DEPARTMENT OF MATHEMATICS, NORTHERN ILLINOIS UNIVERSITY, DEKALB, ILLINOIS 60115  
E-mail address: [hurlburt@math.niu.edu](mailto:hurlburt@math.niu.edu)