

**THE FREQUENCY DECOMPOSITION MULTILEVEL METHOD:
A ROBUST ADDITIVE HIERARCHICAL BASIS
PRECONDITIONER**

ROB STEVENSON

ABSTRACT. Hackbusch's frequency decomposition multilevel method is characterized by the application of three additional coarse-grid corrections in parallel to the standard one. Each coarse-grid correction was designed to damp errors from a different part of the frequency spectrum. In this paper, we introduce a cheap variant of this method, partly based on semicoarsening, which demands fewer recursive calls than the original version. Using the theory of the additive Schwarz methods, we will prove robustness of our method as a preconditioner applied to anisotropic equations.

1. INTRODUCTION

As is well known, the rate of convergence of a multilevel method applied to a discretized elliptic boundary value problem is less than one uniformly in the top level. Yet, without a special choice of the components of the method, the rate of convergence tends to one as the problem becomes less elliptic (singularly perturbed problems), that is, the method is not *robust*. This paper concentrates on the question of robustness for so-called anisotropic problems. The classical way to obtain a robust multilevel method is to choose a smoother adapted to the problem. A disadvantage of this approach is that the resulting smoothers are often expensive, not well parallelizable or, in three dimensions, hard to find.

An alternative approach is to add more coarse-grid corrections to the multilevel method. Representatives of this class of methods are Hackbusch' Frequency Decomposition Multilevel Method (FDMLM) ([2, 3, 5, 6]), which is the subject of this paper, and the Multiple Semi-Coarsened Grids Method ([8, 9, 10]) introduced by Mulder.

In two dimensions, the FD *Two-Level* Method consists of four coarse-grid corrections, that can be performed in parallel, each of them designed to reduce errors in a (non-overlapping) part of the frequency spectrum. To speed up convergence, smoothers can be added to the algorithm but we shall not consider this option. In the (V-cycle) FDMLM, each of the four coarse-grid problems is solved by means of a recursive call, thus involving four coarse-grid corrections on the next coarser level. For a complete explanation of the ideas behind this method, we refer to the papers of Hackbusch. In [5], it has been proved that the *FD TLM* yields a

Received by the editor August 1, 1994.

1991 *Mathematics Subject Classification.* Primary 65N55, 65N30.

Key words and phrases. Frequency decomposition, multilevel method, semicoarsening, finite elements, hierarchical basis, additive Schwarz method, subspace decomposition, robustness.

robust preconditioner, that is, the condition number of the preconditioned system is bounded uniformly in the top level *and* the anisotropy. Until now, robustness of the FDMLM is an open problem.

In this paper, we study a cheap variant of the FDMLM. As was already noted in [3], one of the coarse-grid problems generated by the FDTLM has a bounded condition number ($:= \lambda_{\max}/\lambda_{\min}$) uniformly with respect to the level and the anisotropy. Therefore, instead of applying a recursive call, this system can better be solved using a cheap iterative solver as, e.g., Jacobi's method. Apart from this, with our FDMLM, we solve two of the three remaining coarse-grid problems by means of only *two* instead of four coarse-grid corrections on the next coarser level by using semicoarsening. It will appear then that also one of these two corrections yields a system with bounded condition number, which therefore can be solved cheaply. On the other system we apply the semicoarsening idea recursively.

For any dimension d , the complexity of the resulting algorithm is equivalent to the number of unknowns, even if one would apply more than one recursive calls at certain places in the algorithm. Considered as an additive Schwarz method or, in the terminology of [12], a Parallel Subspace Correction method, it consists of $(\#\text{levels})^d$ subspace corrections compared to $\sim (2^d)^{\#\text{levels}}$ subspace corrections for the FDMLM in its original form.

Using the theory of the additive Schwarz methods, we will prove robustness of our FDMLM as a preconditioner. To do that, we first reformulate the method in an abstract finite element context. This kind of formulation of a multilevel method was introduced in [1]. Then with the help of tensor products, the question of robustness will be reduced to the question of convergence of the method in one dimension applied to the identity and the Laplace operator.

In one dimension, the subspace decomposition that defines our method appears to be very similar to the decomposition of the finite element space into the differences of successive L^2 -orthogonal projections onto the finite element spaces corresponding to coarser grids. In particular, we will show that also our decomposition induces an L^2 -equivalent norm, which means convergence for the identity. The fact that the decomposition using L^2 -orthogonal projections yields an H^1 -equivalent norm plays a crucial role in the modern regularity-free convergence proofs of standard multilevel methods (cf. [12, 13]). By adapting Xu's proof of this result, we will prove the same for our decomposition and with that, convergence for the Laplace operator.

Our FDMLM can be seen as block Jacobi's method after a basis transformation to a certain hierarchical basis. Our convergence result means that independent of the dimension, the stiffness matrix after this transformation has a bounded condition number uniformly in the level and the anisotropy.

In a forthcoming paper ([11]), we will discuss an efficient implementation of the method and present numerical results.

Following [12], we shall use the notations \lesssim , \gtrsim and $\bar{\approx}$. When we write

$$x_1 \lesssim y_1, x_2 \gtrsim y_2 \text{ and } x_3 \bar{\approx} y_3,$$

there exist constants C_1 , c_2 , c_3 and C_3 *independent* of relevant parameters such as the level or the anisotropy, such that

$$x_1 \leq C_1 y_1, x_2 \geq c_2 y_2 \text{ and } c_3 x_3 \leq y_3 \leq C_3 x_3.$$

2. DESCRIPTION OF THE METHOD

2.1. **Basic definitions.** We start by giving some definitions for the *one-dimensional* case. Let $\Omega = (0, 1)$, $h_k = 2^{-(k+1)}$ ($k \in \mathbf{N}_0 = \{0, 1, 2, \dots\}$) and let $\Omega_k^\iota = \Omega \cap h_k(\mathbf{Z} + \frac{1}{2}\iota)$ ($\iota \in \{0, 1\}$). Note that $\Omega_k^0 = \Omega_{k-1}^0 \cup \Omega_{k-1}^1$ (cf. Figure 1). We equip




FIGURE 1. “Standard” and “shifted” coarse-grids Ω_{k-1}^0 and Ω_{k-1}^1 respectively

the space of grid functions on Ω_k^ι , denoted by $\ell^2(\Omega_k^\iota)$, with the Euclidean scalar product

$$\langle \mu, \nu \rangle = \sum_{x \in \Omega_k^\iota} \mu(x) \overline{\nu(x)}$$

and norm $\| \cdot \| = \langle \cdot, \cdot \rangle^{\frac{1}{2}}$.

The prolongations $p^\iota : \ell^2(\Omega_{k-1}^\iota) \rightarrow \ell^2(\Omega_k^0)$ are defined in difference stencil notation as $p^0 = \frac{1}{2} \begin{bmatrix} 1 & 2 & 1 \end{bmatrix}$ (linear interpolation) and $p^1 = \frac{1}{2} \begin{bmatrix} -1 & 2 & -1 \end{bmatrix}$. They satisfy

$$(1) \quad \text{range } p^0 \oplus \text{range } p^1 = \ell^2(\Omega_k^0).$$

The restrictions $r^\iota : \ell^2(\Omega_k^0) \rightarrow \ell^2(\Omega_{k-1}^\iota)$ are defined as adjoints of the corresponding prolongations, that is, $r^0 = \frac{1}{2} \begin{bmatrix} 1 & 2 & 1 \end{bmatrix}$ and $r^1 = \frac{1}{2} \begin{bmatrix} -1 & 2 & -1 \end{bmatrix}$.

For the general *d-dimensional* case, we define the grids $\Omega_k^\iota = \Omega_{k_1}^{\iota_1} \times \dots \times \Omega_{k_d}^{\iota_d}$ ($k \in \mathbf{N}_0^d$, $\iota \in \{0, 1\}^d$). We equip the space $\ell^2(\Omega_k^\iota)$ of grid functions on Ω_k^ι also with the Euclidean scalar product $\langle \mu, \nu \rangle = \sum_{x \in \Omega_k^\iota} \mu(x) \overline{\nu(x)}$.

Since we exploit tensor products quite often, note that $\ell^2(\Omega_k^\iota) = \bigotimes_{j=1}^d \ell^2(\Omega_{k_j}^{\iota_j})$, i.e., $\ell^2(\Omega_k^\iota) = \text{span} \{ \bigotimes_{j=1}^d \mu_j : \mu_j \in \ell^2(\Omega_{k_j}^{\iota_j}) \}$, where $(\bigotimes_{j=1}^d \mu_j)(x) := \prod_{j=1}^d \mu_j(x_j)$. Furthermore, we have $\langle \bigotimes_{j=1}^d \mu_j, \bigotimes_{j=1}^d \nu_j \rangle = \prod_{j=1}^d \langle \mu_j, \nu_j \rangle$.

2.2. **Derivation of the (modified) FDMLM.** First, we consider the two-dimensional case. In [2], the FD *Two-Level Method* to solve a system $\mathcal{A}\mu = \beta$ on $\Omega_{J,J}^{0,0}$ was defined by

$$\mu \leftarrow \mu - \sum_{\iota, j \in \{0, 1\}} p_x^\iota \otimes p_y^j (r_x^\iota \otimes r_y^j \mathcal{A} p_x^\iota \otimes p_y^j)^{-1} r_x^\iota \otimes r_y^j (\mathcal{A}\mu - \beta),$$

where thus $r_x^\iota \otimes r_y^j \mathcal{A} p_x^\iota \otimes p_y^j$ acts on the space of grid functions on $\Omega_{j-1, j-1}^{\iota j}$ ($= \Omega_{j-1}^\iota \times \Omega_{j-1}^j$). [It will be clear why we avoid the term *two-grid method*.] Using the abbreviations $p^{\iota j}$ and $r^{\iota j}$ for $p_x^\iota \otimes p_y^j$ and $r_x^\iota \otimes r_y^j$, respectively, we have

$$p^{\iota j} = \frac{1}{4} \begin{bmatrix} (-1)^{\iota+j} & (-1)^j 2 & (-1)^{\iota+j} \\ (-1)^{\iota} 2 & 4 & (-1)^{\iota} 2 \\ (-1)^{\iota+j} & (-1)^j 2 & (-1)^{\iota+j} \end{bmatrix}$$

and

$$r^{ij}(= (p^{ij})^*) = \frac{1}{4} \begin{bmatrix} (-1)^{i+j} & (-1)^j 2 & (-1)^{i+j} \\ (-1)^i 2 & 4 & (-1)^i 2 \\ (-1)^{i+j} & (-1)^j 2 & (-1)^{i+j} \end{bmatrix}.$$

We consider only $\mathcal{A} > 0$. Then, because of the Galerkin approach, the error amplification operator of the method is given by

$$\mu^* - \mu^{\text{new}} = (I - \sum_{i,j \in \{0,1\}} \mathcal{P}^{ij})(\mu^* - \mu^{\text{old}}),$$

where μ^* is the exact solution and \mathcal{P}^{ij} is the projection from $\ell^2(\Omega_{JJ}^{00})$ onto the range p^{ij} orthogonal with respect to $\langle \mathcal{A}, \cdot \rangle$. The range of the standard prolongation p^{00} contains the “smooth” functions. The one-dimensional prolongation p^1 was chosen such that the ranges of the p^{ij} for $(i, j) \neq (0, 0)$ contain the different types of oscillating functions, so that also errors of that kind are corrected.

We consider systems that arise from the application of the bilinear finite element method to

$$\begin{cases} -(a_1 \partial_1^2 + a_2 \partial_2^2)u = f & \text{on } \Omega^2, \\ u = 0 & \text{on } \partial\Omega^2, \end{cases}$$

that is,

$$\mathcal{A} = \frac{1}{6}a_1 \begin{bmatrix} -1 & 2 & -1 \end{bmatrix} \begin{bmatrix} 1 \\ 4 \\ 1 \end{bmatrix} + \frac{1}{6}a_2 \begin{bmatrix} -1 \\ 2 \\ -1 \end{bmatrix} \begin{bmatrix} 1 & 4 & 1 \end{bmatrix} : \ell^2(\Omega_{JJ}^{00}) \rightarrow \ell^2(\Omega_{JJ}^{00}),$$

where $a_1, a_2 \geq 0$ and $a_1 + a_2 > 0$. This kind of problem is called *anisotropic* when $a_1 \ll a_2$ or $a_1 \gg a_2$. In [5], it was proved that the FDTLM yields a *robust* preconditioner, that is, the condition number $\kappa(\sum_{i,j \in \{0,1\}} p^{ij} (r^{ij} \mathcal{A} p^{ij})^{-1} r^{ij} \mathcal{A}) \lesssim 1$ (uniformly in J and a_i). Our aim is to prove the same for a multilevel version.

In its original form, the multilevel version consisted of recursive calls for each of the four coarse-grid problems on the grids Ω_{J-1J-1}^{00} , Ω_{J-1J-1}^{01} , Ω_{J-1J-1}^{10} and Ω_{J-1J-1}^{11} . By $r^0 \begin{bmatrix} -1 & 2 & -1 \end{bmatrix} p^0 = \frac{1}{2} \begin{bmatrix} -1 & 2 & -1 \end{bmatrix}$, $r^0 \begin{bmatrix} 1 & 4 & 1 \end{bmatrix} p^0 = 2 \begin{bmatrix} 1 & 4 & 1 \end{bmatrix}$, $r^1 \begin{bmatrix} 1 & 4 & 1 \end{bmatrix} p^1 = 4I$ and

$$r^1 \begin{bmatrix} -1 & 2 & -1 \end{bmatrix} p^1 = \frac{1}{2} \begin{bmatrix} 7 & 3 & & & \\ 3 & 10 & 3 & & \\ & & \ddots & & \\ & & & 3 & 10 & 3 \\ & & & & 7 & 3 \end{bmatrix} =: \frac{1}{2} \begin{bmatrix} 3 & 10 & 3 \end{bmatrix}^{\sim},$$

the operators on the spaces of grid functions on these grids are

$$r^{00} \mathcal{A} p^{00} = \frac{1}{6}a_1 \begin{bmatrix} -1 & 2 & -1 \end{bmatrix} \begin{bmatrix} 1 \\ 4 \\ 1 \end{bmatrix} + \frac{1}{6}a_2 \begin{bmatrix} -1 \\ 2 \\ -1 \end{bmatrix} \begin{bmatrix} 1 & 4 & 1 \end{bmatrix} \quad \text{on } \ell^2(\Omega_{J-1J-1}^{00}),$$

$$r^{01} \mathcal{A} p^{01} = \frac{1}{3}a_1 \begin{bmatrix} -1 & 2 & -1 \end{bmatrix} + \frac{1}{6}a_2 \begin{bmatrix} 3 \\ 10 \\ 3 \end{bmatrix}^{\sim} \begin{bmatrix} 1 & 4 & 1 \end{bmatrix} \quad \text{on } \ell^2(\Omega_{J-1J-1}^{01}),$$

$$r^{10}\mathcal{A}p^{10} = \frac{1}{6}a_1 [3 \quad 10 \quad 3] \sim \begin{bmatrix} 1 \\ 4 \\ 1 \end{bmatrix} + \frac{1}{3}a_2 \begin{bmatrix} -1 \\ 2 \\ -1 \end{bmatrix} \quad \text{on } \ell^2(\Omega_{J-1J-1}^{10})$$

and

$$r^{11}\mathcal{A}p^{11} = \frac{1}{3}a_1 [3 \quad 10 \quad 3] \sim + \frac{1}{3}a_2 \begin{bmatrix} 3 \\ 10 \\ 3 \end{bmatrix} \sim \quad \text{on } \ell^2(\Omega_{J-1J-1}^{11}).$$

As noted in [3], there holds $\kappa(r^{11}\mathcal{A}p^{11}) \lesssim 1$ (uniformly in the grid sizes and a_i). The argument is that for $\mathcal{B}_1, \mathcal{B}_2 > 0$, we have $\kappa(\mathcal{B}_1 + \mathcal{B}_2) \leq \max\{\kappa(\mathcal{B}_1), \kappa(\mathcal{B}_2)\}$. So instead of applying a recursive call, the corresponding system can be solved using a cheap iterative solver. Furthermore, in [3] it was argued that in the cases $a_1 \leq a_2$ or $a_1 \geq a_2$ also one of the two operators $r^{01}\mathcal{A}p^{01}$ and $r^{10}\mathcal{A}p^{10}$ has a bounded condition number. Yet, this argument cannot be applied to construct a method that is robust for the general variable-coefficient case. Therefore, we will use another idea to further reduce the number of recursive calls.

Consider the following two operators that arise from $\tilde{\mathcal{A}} := r^{01}\mathcal{A}p^{01}$ by means of *semicoarsening* Ω_{J-1J-1}^{01} in the x -direction, that is, *in the direction where we have not applied p^1 so far*,

$$r_x^0\tilde{\mathcal{A}}p_x^0 = \frac{1}{6}a_1 [-1 \quad 2 \quad -1] + \frac{1}{3}a_2 \begin{bmatrix} 3 \\ 10 \\ 3 \end{bmatrix} \sim [1 \quad 4 \quad 1] \quad \text{on } \ell^2(\Omega_{J-2J-1}^{01})$$

and

$$r_x^1\tilde{\mathcal{A}}p_x^1 = \frac{1}{6}a_1 [3 \quad 10 \quad 3] \sim + \frac{2}{3}a_2 \begin{bmatrix} 3 \\ 10 \\ 3 \end{bmatrix} \sim \quad \text{on } \ell^2(\Omega_{J-2J-1}^{11}).$$

Then the first operator is of the same type as $\tilde{\mathcal{A}}$, and so we can apply (x -)semicoarsening recursively or, if $J - 2 = 0$, the operator has a bounded condition number and therefore the system can be solved using a cheap iterative solver. The second operator always has a bounded condition number.

Analogously to the above procedure, we can solve the system on Ω_{J-1J-1}^{10} using semicoarsening in the y -direction. Finally, as with the original version, the system on Ω_{J-1J-1}^{00} is solved with a recursive call of the entire method, with which this informal description of the modified FDMMLM is completed (see Figure 2).

In view of the following, note that since, e.g., on Ω_{J-1J-1}^{01} no system is solved (unless $J - 1 = 0$), but only coarse-grid corrections are invoked, this (intermediate) grid and the operators defined on it are only important for an efficient implementation. Because of the Galerkin approach, the mathematical properties of the resulting method are determined by the (sequence of) prolongations from the grids on which systems are (approximately) solved (leaves in the tree of Figure 2) onto the finest grid Ω_{JJ}^{00} . For example, for $J = 2$ these prolongations are $p_x^0p_x^0 \otimes p_y^0p_y^0$, $p_x^0p_x^0 \otimes p_y^0p_y^1$, $p_x^0p_x^1 \otimes p_y^1$, $p_x^0p_x^1 \otimes p_y^0p_y^0$, $p_x^0p_x^1 \otimes p_y^0p_y^1$, $p_x^0p_x^1 \otimes p_y^1$, $p_x^1 \otimes p_y^0p_y^0$, $p_x^1 \otimes p_y^0p_y^1$ and $p_x^1 \otimes p_y^1$, i.e., tensor products of all possible combinations of $p_x^0p_x^0$, $p_x^0p_x^1$, p_x^1 and $p_y^0p_y^0$, $p_y^0p_y^1$, p_y^1 .

We are now ready to give a formal description of the modified FDMMLM.




FIGURE 2. Grids and prolongations defining the modified FDMLM in two dimensions. [We dotted the lines at the bottom of the figure since the pictures of the grids correspond to $J = 2$, i.e., the three-level case.]

Algorithm 2.1. Let $\mathcal{A}\mu = \beta$ be a system on the d -dimensional grid $\Omega_{J,\dots,J}^{0,\dots,0}$. For $0 \leq k \leq J$, define the one-dimensional prolongation $p_k = p_k^{(J)}$ by

$$p_k = \begin{cases} \underbrace{p^0 \cdots p^0}_{(J-k) \times} p^1 & : \ell^2(\Omega_{k-1}^1) \rightarrow \ell^2(\Omega_J^0), \quad k \geq 1, \\ \underbrace{p^0 \cdots p^0}_{J \times} & : \ell^2(\Omega_0^0) \rightarrow \ell^2(\Omega_J^0), \quad k = 0. \end{cases}$$

Note that from (1) we have $\ell^2(\Omega_J^0) = \bigoplus_{k=0}^J \text{range } p_k$. For $k \in I := \{0, \dots, J\}^d$, we define

$$p_k = \bigotimes_{j=1}^d p_{k_j, x_j}, \quad r_k = p_k^* (= \bigotimes_{j=1}^d r_{k_d, x_d}) \quad \text{and} \quad \mathcal{A}_k = r_k \mathcal{A} p_k.$$

Now let \mathcal{B}_k be such that \mathcal{B}_k^{-1} is a cheap approximation of \mathcal{A}_k^{-1} . Then the (modified) FDMLM is defined by

$$(2) \quad \mu \leftarrow \mu - \sum_{k \in I} p_k \mathcal{B}_k^{-1} r_k (\mathcal{A}\mu - \beta).$$

Our FDMLM is an example of an *additive Schwarz method* or *Parallel Subspace Correction method* with subspace $\text{range } p_k$ satisfying $\ell^2(\Omega_{J,\dots,J}^{0,\dots,0}) = \bigoplus_{k \in I} \text{range } p_k$.

Remark 2.2. Since, e.g., $\Omega_{J-1,J-1}^{01}$ ($d = 2$) is coarsened only in the x -direction, the elementary one-dimensional prolongations p^0 and p^1 , which are the building blocks

of all prolongations in the algorithm, always map onto the space of grid functions on a non-shifted grid, that is, a grid Ω_k^0 for some $k \in \mathbf{N}$. So in contrast to the original FDMLM, we do not have to construct boundary adaptations for p^0 and p^1 in order to maintain property (1).

We want to prove robustness of this method applied to anisotropic problems. As a consequence of the following lemma it is then sufficient to analyze the FDMLM with exact subspace corrections ($\mathcal{B}_k = \mathcal{A}_k$). The straightforward proof of this lemma is left to the reader.

Lemma 2.3. *Let $\mathcal{A} > 0$ and $\mathcal{B}_k > 0$ ($k \in I$). Define $\Lambda = \max_{k \in I} \lambda_{\max}(\mathcal{B}_k^{-1} \mathcal{A}_k)$ and $\lambda = \min_{k \in I} \lambda_{\min}(\mathcal{B}_k^{-1} \mathcal{A}_k)$. Then*

$$\kappa \left(\sum_{k \in I} p_k \mathcal{B}_k^{-1} r_k \mathcal{A} \right) \leq \frac{\Lambda}{\lambda} \kappa \left(\sum_{k \in I} p_k \mathcal{A}_k^{-1} r_k \mathcal{A} \right).$$

Analogously to the two-dimensional case, for d -dimensional anisotropic problems we have that $\kappa(\mathcal{A}_k) \lesssim 1$ ($k \in I$). So already the simple Richardson iteration, that is $\mathcal{B}_k = \rho(\mathcal{A}_k)$, gives $\frac{\Lambda}{\lambda} = \max_{k \in I} \kappa(\mathcal{A}_k) \lesssim 1$.

2.3. Computational complexity. For ease of presentation we consider the two-dimensional case. The general case can be handled by induction.

Assume that the application of \mathcal{B}_k^{-1} ($k \in I$) costs a number of operations that is equivalent to the number of points of the grid in question. For $k_1 \leq k_2$, let $W_{k_1 k_2}^{01}$ ($W_{k_2 k_1}^{10}$) be the number of arithmetic operations necessary to treat a system on $\Omega_{k_1 k_2}^{01}$ ($\Omega_{k_2 k_1}^{10}$) using the recursive application of semicoarsening in the x - (y -) direction. Then we have $W_{k_1 k_2}^{01} \approx \#\Omega_{k_1 k_2}^{01} + W_{k_1-1 k_2}^{01}$, which gives $W_{k_1 k_2}^{01} \approx \#\Omega_{k_1 k_2}^{01}$ and analogously $W_{k_2 k_1}^{10} \approx \#\Omega_{k_2 k_1}^{10}$. Finally, let W_{kk}^{00} be the number of arithmetic operations necessary for an entire FDMLM call on Ω_{kk}^{00} . We conclude that

$$\begin{aligned} W_{kk}^{00} &\approx W_{k-1 k-1}^{00} + W_{k-1 k-1}^{01} + W_{k-1 k-1}^{10} + \#\Omega_{kk}^{00} \\ &\approx W_{k-1 k-1}^{00} + \#\Omega_{kk}^{00}, \end{aligned}$$

which implies $W_{kk}^{00} \approx \#\Omega_{kk}^{00}$. Note that since $\#\Omega_{k-1 k-1}^{00} / \#\Omega_{kk}^{00} = \frac{1}{4}$, more than one recursive call on $\Omega_{k-1 k-1}^{00}$ can be applied. The number of recursive calls involving semicoarsening should be restricted to one.

3. PROOF OF ROBUSTNESS OF THE FDMLM

3.1. Coordinate-free finite element formulation. To facilitate the analysis, we reformulate the algorithm in a more abstract context. We start by giving some definitions for the *one-dimensional* case.

For $i \in \{0, 1\}$, $k \in \mathbf{N}_0$, $x \in \Omega_k^i$, define $\delta_{k,x}^i \in \ell^2(\Omega_k^i)$ by $\delta_{k,x}^i(y) = \begin{cases} 1, & x = y, \\ 0, & x \neq y. \end{cases}$

Define $P_k : \ell^2(\Omega_k^0) \rightarrow H_0^1(\Omega) \subset L^2(\Omega)$ as the linear interpolation operator using zero boundary values. Put $\mathcal{M}_k = \text{range} P_k$, that is, \mathcal{M}_k is the linear finite element space corresponding to the grid Ω_k^0 . The basis $\{\phi_{k,x}^0 := P_k \delta_{k,x}^0 : x \in \Omega_k^0\}$ is the standard (nodal) basis of \mathcal{M}_k . We equip \mathcal{M}_k with scalar product

$$(3) \quad \langle \cdot, \cdot \rangle_{\mathcal{M}_k} = h_k \langle P_k^{-1} \cdot, P_k^{-1} \cdot \rangle$$

and norm $\| \cdot \|_{\mathcal{M}_k} = \langle \cdot, \cdot \rangle_{\mathcal{M}_k}^{\frac{1}{2}}$. It is well known that $\| \cdot \|_{\mathcal{M}_k} \approx \| \cdot \|_{L^2}$ (uniformly in k).

For $k > 1$, we define $\mathcal{V}_k = \text{range}(P_k p^1 : \ell^2(\Omega_{k-1}^1) \rightarrow H_0^1(\Omega))$. We will call the basis $\{\phi_{k,x}^1 := P_k p^1 \delta_{k-1,x}^1 : x \in \Omega_{k-1}^1\}$ the standard basis of \mathcal{V}_k . Using $P_k p^0 = P_{k-1}$, we find that (1) is equivalent to

$$(4) \quad \mathcal{M}_k = \mathcal{M}_{k-1} \oplus^{\perp \mathcal{M}_k} \mathcal{V}_k.$$

So, with the definition $\mathcal{V}_0 = \mathcal{M}_0$, the union of the bases of $\mathcal{V}_0, \dots, \mathcal{V}_k$ forms a basis of \mathcal{M}_k , which is therefore called a *hierarchical basis* (cf. Figure 3). Note that (4)




FIGURE 3. Hierarchical basis of \mathcal{M}_2

does not imply that $\mathcal{V}_0, \dots, \mathcal{V}_k$ are mutually orthogonal with respect to some scalar product.

For $k \in \mathbb{N}_0$, let $I_k : \mathcal{V}_k \rightarrow L^2(\Omega)$ be the inclusion operator. Since

$$P_J p_k = \begin{cases} P_J p^0 \cdots p^0 p^1 = P_k p^1, & 1 \leq k \leq J, \\ P_J \underbrace{p^0 \cdots p^0}_{(J-k) \times} = P_0, & k = 0, \end{cases}$$

we find that p_k is the representation of I_k with respect to the standard bases on \mathcal{V}_k and \mathcal{M}_J .

As usual, for some basis $\{\psi_i : i \in I_{\mathcal{W}}\}$ of a subspace $\mathcal{W} \subset L^2(\Omega)$, we define the dual basis $\{\tilde{\psi}_i : i \in I_{\mathcal{W}}\}$ of \mathcal{W} by $(\tilde{\psi}_i, \psi_j)_{L^2} = \begin{cases} 1, & i = j, \\ 0, & i \neq j. \end{cases}$ Let $V_k : L^2(\Omega) \rightarrow \mathcal{V}_k$

be the adjoint of I_k with respect to the L^2 -scalar product on both spaces, that is, V_k is the L^2 -orthogonal projection onto \mathcal{V}_k . Then we find that the representation of $V_k|_{\mathcal{M}_J}$ with respect to the dual bases on \mathcal{M}_J and \mathcal{V}_k is equal to the adjoint of p_k , that is, r_k .

In the *multidimensional* case everything is defined using tensor products, that is,

$$\mathcal{M}_k = \bigotimes_{j=1}^d \mathcal{M}_{k_j}, \quad \mathcal{V}_k = \bigotimes_{j=1}^d \mathcal{V}_{k_j}, \quad I_k = \bigotimes_{j=1}^d I_{k_j} \quad \text{and} \quad V_k (= I_k^*) = \bigotimes_{j=1}^d V_{k_j}.$$

So, $I_k : \mathcal{V}_k \rightarrow L^2(\Omega^d)$ is the inclusion operator and $V_k : L^2(\Omega^d) \rightarrow \mathcal{V}_k$ is the L^2 -orthogonal projection onto \mathcal{V}_k . We equip \mathcal{M}_k and \mathcal{V}_k with standard bases obtained by making tensor products of the standard basis functions of its factors, that is, the standard bases consist of functions of the form $\phi_{k,x}^i = \bigotimes_{j=1}^d \phi_{k_j,x_j}^{i_j}$. Concerning dual basis functions, note that $\tilde{\phi}_{k,x}^i = \bigotimes_{j=1}^d \tilde{\phi}_{k_j,x_j}^{i_j}$. Using the abbreviation \mathbf{m} for multi-indices $(m, \dots, m) \in \mathbf{N}_0^d$, we conclude that for $k \in I$, p_k is the representation of I_k with respect to the standard bases on \mathcal{V}_k and $\mathcal{M}_{\mathbf{J}}$, and r_k is the representation of $V_k|_{\mathcal{M}_{\mathbf{J}}}$ with respect to the dual bases.

Finally, for a given system $\mathcal{A}\mu = \beta$ on $\Omega_{\mathbf{J}}^0$, define $A : \mathcal{M}_{\mathbf{J}} \rightarrow \mathcal{M}_{\mathbf{J}}$ by

$$(5) \quad (A\phi_{\mathbf{J},y}^0, \phi_{\mathbf{J},x}^0)_{L^2} = \mathcal{A}_{xy} \quad (x, y \in \Omega_{\mathbf{J}}^0).$$

Then \mathcal{A} is the representation of A with respect to the standard basis on its domain and dual basis on its image. With the definition $A_k = V_k A I_k$, we arrive at the conclusion that (2), with $\mathcal{B}_k = \mathcal{A}_k$, is a matrix formulation of the iteration

$$(6) \quad u \leftarrow u - \sum_{k \in I} I_k A_k^{-1} V_k (A u - f),$$

where $u = \sum_{x \in \Omega_{\mathbf{J}}^0} \mu(x) \phi_{\mathbf{J},x}^0$, $f = \sum_{x \in \Omega_{\mathbf{J}}^0} \beta(x) \tilde{\phi}_{\mathbf{J},x}^0$ (that is, $\beta(x) = (f, \phi_{\mathbf{J},x}^0)_{L^2}$). Because the condition number κ was defined as the quotient of the largest and smallest eigenvalue, clearly we have $\kappa(\sum_{k \in I} p_k A_k^{-1} r_k \mathcal{A}) = \kappa(\sum_{k \in I} I_k A_k^{-1} V_k A)$.

Remark 3.1. We defined the operator A using the matrix \mathcal{A} . Of course, the usual procedure is the other way around. If a is a bilinear form on $H_0^1(\Omega^d)$ and $A : \mathcal{M}_{\mathbf{J}} \rightarrow \mathcal{M}_{\mathbf{J}}$ is defined by $(Au, v)_{L^2} = a(u, v)$ ($u, v \in \mathcal{M}_{\mathbf{J}}$), then \mathcal{A} defined by (5) is called the *stiffness matrix* with respect to the (multilinear) basis $\{\phi_{\mathbf{J},x}^0 : x \in \Omega_{\mathbf{J}}^0\}$.

Remark 3.2. Consider the hierarchical basis of $\mathcal{M}_{\mathbf{J}} = \bigoplus_{k \in I} \mathcal{V}_k$ that is obtained by taking the union of the standard bases of the \mathcal{V}_k . The iteration (6) with respect to this basis, that is, the hierarchical basis for the solution and its dual for the right-hand side, is just block Jacobi's method with a partitioning into blocks corresponding to the spaces \mathcal{V}_k . As we have seen, for anisotropic problems, the diagonal blocks have bounded condition number and so robustness of (6) implies that, properly scaled, the stiffness matrix with respect to this hierarchical basis has a bounded condition number uniformly in the level and the anisotropy.

3.2. Main theorem; reduction to one-dimensional cases. The fact that $\mathcal{M}_{\mathbf{J}} = \bigoplus_{k \in I} \mathcal{V}_k$ is a *direct* sum decomposition implies that there exist projections $Z_k = Z_k^{(\mathbf{J})} : \mathcal{M}_{\mathbf{J}} \rightarrow \mathcal{V}_k$ such that $\sum_{k \in I} Z_k = I$ on $\mathcal{M}_{\mathbf{J}}$. Note that $Z_k I_{k'} = 0$ if $k \neq k'$ and that $Z_k I_k$ is the identity on \mathcal{V}_k .

Lemma 3.3. *Define $W = \sum_{k \in I} Z_k^* A_k Z_k : \mathcal{M}_{\mathbf{J}} \rightarrow \mathcal{M}_{\mathbf{J}}$. Then W^{-1} exists and is equal to $\sum_{k \in I} I_k A_k^{-1} V_k$.*

Proof. $\sum_{k \in I} Z_k^* A_k Z_k \sum_{k' \in I} I_{k'} A_{k'}^{-1} V_{k'} = \sum_{k \in I} Z_k^* V_k = (\sum_{k \in I} I_k Z_k)^* = I$. \square

This lemma shows that for $A > 0$, the condition number $\kappa(\sum_{k \in I} I_k A_k^{-1} V_k A)$ is the quotient $\frac{\Gamma}{\gamma}$ of the optimum constants in the inequalities $\gamma W \leq A \leq \Gamma W$ or, by $A_k = V_k A I_k$,

$$(7) \quad \gamma \sum_{k \in I} (AZ_k u, Z_k u)_{L^2} \leq (Au, u)_{L^2} \leq \Gamma \sum_{k \in I} (AZ_k u, Z_k u)_{L^2} \quad (u \in \mathcal{M}_{\mathbf{J}}).$$

We are now ready to formulate our main theorem.

Theorem 3.4. *For nonnegative constants a_j and b with $\sum_{j=1}^d a_j + b > 0$, let*

$$a(u, v) = \sum_{j=1}^d \int_{\Omega^d} a_j \partial_j u \overline{\partial_j v} + \int_{\Omega^d} bu\overline{v} \quad (u, v \in H_0^1(\Omega^d))$$

and let $A : \mathcal{M}_J \rightarrow \mathcal{M}_J$ be defined by $(Au, v)_{L^2} = a(u, v)$ ($u, v \in \mathcal{M}_J$). Then we have $\kappa(\sum_{k \in I} I_k A_k^{-1} V_k A) \lesssim 1$ (uniformly in J , a_j and b).

Remark 3.5. From (7), we immediately see that the theorem can be extended to all operators \tilde{A} for which there exist $c, C > 0$ such that

$$c(Au, u)_{L^2} \leq (\tilde{A}u, u)_{L^2} \leq C(Au, u)_{L^2}$$

for some A as described in the theorem. Examples are *linear* finite element discretizations or discretizations of elliptic boundary value problems with *nonconstant* coefficients. With a view to the nonconstant-coefficient case, we note that clearly the theorem does not yield boundedness of the condition number that is uniform in C/c .

To prove Theorem 3.4, we first note that if (7) is satisfied by $A^{(1)}$ and $A^{(2)}$, then it is satisfied by $c_1 A^{(1)} + c_2 A^{(2)}$ for any $c_1, c_2 \geq 0$. As a consequence, we only have to consider tensor product operators $A = \otimes_{j=1}^d A_j$, where $(A_j u, u)_{L^2} = (u, u)_{L^2}$ or $(A_j u, u)_{L^2} = (u', u')_{L^2}$ ($u \in \mathcal{M}_J$). Secondly, we observe that (7) is equivalent to

$$(8) \quad \sigma\left(\sum_{k \in I} A^{-\frac{1}{2}} Z_k^* A Z_k A^{-\frac{1}{2}}\right) \subset \left[\frac{1}{\Gamma}, \frac{1}{\gamma}\right].$$

As a special case of the general definition, the one-dimensional $Z_k^{(J)} : \mathcal{M}_J \rightarrow \mathcal{V}_k$ were defined by $\sum_{k=0}^J Z_k^{(J)} = I$. For $k \in I$, we have $Z_k (= Z_k^{(J)}) = \otimes_{j=1}^d Z_{k_j}^{(J)}$. From

$$\begin{aligned} \sigma\left(\sum_{k \in I} A^{-\frac{1}{2}} Z_k^* A Z_k A^{-\frac{1}{2}}\right) &= \sigma\left(\sum_{k \in I} \otimes_{j=1}^d A_j^{-\frac{1}{2}} Z_{k_j}^{(J)*} A_j Z_{k_j}^{(J)} A_j^{-\frac{1}{2}}\right) \\ &= \sigma\left(\otimes_{j=1}^d \sum_{k=0}^J A_j^{-\frac{1}{2}} Z_{k_j}^{(J)*} A_j Z_{k_j}^{(J)} A_j^{-\frac{1}{2}}\right) = \prod_{j=1}^d \sigma\left(\sum_{k=0}^J A_j^{-\frac{1}{2}} Z_{k_j}^{(J)*} A_j Z_{k_j}^{(J)} A_j^{-\frac{1}{2}}\right), \end{aligned}$$

and again by using the equivalence of (8) and (7) but now for the one-dimensional case, we conclude that it suffices to prove the following norm equivalences in *one dimension*:

$$(9) \quad \sum_{k=0}^J \|Z_k^{(J)} u\|_{L^2}^2 \approx \|u\|_{L^2}^2$$

and

$$(10) \quad \sum_{k=0}^J \|Z_k^{(J)} u\|_{H^1}^2 \approx \|u\|_{H^1}^2$$

($u \in \mathcal{M}_J$), where $\|\cdot\|_{H^1} := (\cdot, \cdot)_{H^1}^{\frac{1}{2}}$ and $(u, v)_{H^1} := (u', v')_{L^2}$.

We start with constructing an explicit formula for the one-dimensional projections $Z_k^{(J)}$:

Lemma 3.6. *Let $Y_k : \mathcal{M}_{k+1} \rightarrow \mathcal{M}_k$ be the projection on \mathcal{M}_k orthogonal with respect to $\langle \cdot, \cdot \rangle_{\mathcal{M}_{k+1}}$. Then*

$$(11) \quad Z_k^{(J)} = \begin{cases} Y_0 \cdots Y_{J-1}, & k = 0, \\ (I - Y_{k-1})Y_k \cdots Y_{J-1}, & 1 \leq k \leq J. \end{cases}$$

Proof. From $\mathcal{M}_{J-1} \oplus^{\perp \mathcal{M}_J} \mathcal{V}_J = M_J$ and $M_{J-1} = \bigoplus_{k=1}^{J-1} \mathcal{V}_k$, it follows that $Z_J^{(J)} = (I - Y_{J-1})$ and $Z_k^{(J)} = Z_k^{(J-1)}Y_{J-1}$ ($0 \leq k \leq J-1$). \square

Remark 3.7. As noted before, $\|\cdot\|_{\mathcal{M}_k} \approx \|\cdot\|_{L^2}$ on \mathcal{M}_k (uniformly in k). If $\|\cdot\|_{\mathcal{M}_k}$ would be equal to $\|\cdot\|_{L^2}$ (which is not the case), then $Z_k^{(J)}$ defined by (11) would be equal to $(Q_k - Q_{k-1})|_{\mathcal{M}_J}$, where $Q_k : L^2(\Omega) \rightarrow \mathcal{M}_k$ is the L^2 -orthogonal projection onto \mathcal{M}_k and $Q_{-1} := 0$. For the decomposition $u = \sum_{k=0}^J (Q_k - Q_{k-1})u$, (9) is trivially true. The corresponding relation (10) is famous and is the key to the modern regularity-free convergence proofs of standard multilevel methods (cf. [12, 13]). Unlike the decomposition $u = \sum_{k=0}^J (Q_k - Q_{k-1})u$, our decomposition was not introduced as a clever trick for the analysis of an overlapping subspace correction method, but it was yielded by the method itself.

In the next two subsections, §§3.3 and 3.4, we will prove the norm equivalences (9) and (10), respectively. We will prove (9) by estimating the angles between the spaces \mathcal{V}_k with respect to the L^2 -scalar product for our one-dimensional regular grid case. For the same case, an alternative proof exploiting the standard bases of the \mathcal{V}_k appeared in [7]. We will prove (10) in an abstract framework using (9), $\sum_{k=0}^J \|(Q_k - Q_{k-1})u\|_{H^1}^2 \approx \|u\|_{H^1}^2$ ($u \in \mathcal{M}_J$) and $\|\cdot\|_{\mathcal{M}_k} \approx \|\cdot\|_{L^2}$ on \mathcal{M}_k .

3.3. An L^2 -equivalent norm on \mathcal{M}_J . Since $Z_k^{(J)}$ is the projection from \mathcal{M}_J onto \mathcal{V}_k satisfying $\sum_{k=0}^J Z_k^{(J)} = I$, the validity of (9) depends on the angle between the spaces \mathcal{V}_k .

Lemma 3.8. *Let θ_{kl} be the smallest constant satisfying*

$$|(u, v)_{L^2}| \leq \theta_{kl} \|u\|_{L^2} \|v\|_{L^2} \quad \text{for all } u \in \mathcal{V}_k, v \in \mathcal{V}_l$$

(strengthened Cauchy-Schwarz inequality). Put $\Theta = (\theta_{kl})_{k,l}$ and let $\rho(\Theta - I) < 1$. Then

$$1 - \rho(\Theta - I) \leq \|u\|_{L^2}^2 / \sum_{k=0}^J \|Z_k^{(J)}u\|_{L^2}^2 \leq 1 + \rho(\Theta - I) \quad (u \in \mathcal{M}_J).$$

Proof. Use

$$\|u\|_{L^2}^2 = \left(\sum_{k=0}^J Z_k^{(J)}u, \sum_{l=0}^J Z_l^{(J)}u \right)_{L^2} = \sum_{k=0}^J \|Z_k^{(J)}u\|_{L^2}^2 + \sum_{0 \leq k \neq l \leq J} (Z_k^{(J)}u, Z_l^{(J)}u)_{L^2}$$

and

$$\begin{aligned} \left| \sum_{0 \leq k \neq l \leq J} (Z_k^{(J)}u, Z_l^{(J)}u)_{L^2} \right| &\leq \sum_{0 \leq k \neq l \leq J} \theta_{kl} \|Z_k^{(J)}u\|_{L^2} \|Z_l^{(J)}u\|_{L^2} \\ &\leq \rho(\Theta - I) \sum_{k=0}^J \|Z_k^{(J)}u\|_{L^2}^2. \quad \square \end{aligned}$$

Proposition 3.9. *For $k > l$, there holds*

$$\theta_{kl} \leq \|(r^1 M_k p^1)^{-\frac{1}{2}} r^1 M_k \underbrace{p^0 \cdots p^0}_{(k-l) \times} M_l^{-\frac{1}{2}}\|,$$

where $M_m := R_m P_m$ (mass matrix) and R_m is the adjoint of $P_m : \ell^2(\Omega_m^0) \rightarrow L^2(\Omega)$.

Proof. The result follows from

$$\begin{aligned} \theta_{kl} &= \sup_{\substack{0 \neq u \in \mathcal{V}_k \\ 0 \neq v \in \mathcal{V}_l}} \frac{|(u, v)_{L^2}|}{\|u\|_{L^2} \|v\|_{L^2}} \leq \sup_{\substack{0 \neq u \in \mathcal{V}_k \\ 0 \neq v \in \mathcal{M}_l}} \frac{|(u, v)_{L^2}|}{\|u\|_{L^2} \|v\|_{L^2}} \\ &= \sup_{\substack{0 \neq \mu \in \ell^2(\Omega_{k-1}^1) \\ 0 \neq \nu \in \ell^2(\Omega_l^0)}} \frac{|(P_k p^1 \mu, P_l \nu)_{L^2}|}{(P_k p^1 \mu, P_k p^1 \mu)_{L^2}^{\frac{1}{2}} (P_l \nu, P_l \nu)_{L^2}^{\frac{1}{2}}} \\ &= \sup_{\substack{0 \neq \mu \in \ell^2(\Omega_{k-1}^1) \\ 0 \neq \nu \in \ell^2(\Omega_l^0)}} \frac{|(P_k p^1 \mu, P_k \underbrace{p^0 \cdots p^0}_{(k-l) \times} \nu)_{L^2}|}{\langle r^1 M_k p^1 \mu, \mu \rangle^{\frac{1}{2}} \langle M_l \nu, \nu \rangle^{\frac{1}{2}}} \\ &= \sup_{\substack{0 \neq \tilde{\mu} \in \ell^2(\Omega_{k-1}^1) \\ 0 \neq \tilde{\nu} \in \ell^2(\Omega_l^0)}} \frac{|(\tilde{\mu}, (r^1 M_k p^1)^{-\frac{1}{2}} r^1 M_k \underbrace{p^0 \cdots p^0}_{(k-l) \times} M_l^{-\frac{1}{2}} \tilde{\nu})|}{\|\tilde{\mu}\| \|\tilde{\nu}\|}. \quad \square \end{aligned}$$

We will now estimate the upper bound for θ_{kl} from Proposition 3.9 in our one-dimensional regular grid case resulting in (9). Define $\bar{p}^i : \ell^2(\Omega_{m-1}^i) \rightarrow \ell^2(\Omega_m^0)$ by

$$(\bar{p}^i u)(x) = \begin{cases} u(x), & x \in \Omega_{m-1}^i, \\ 0, & x \in \Omega_m^0 \setminus \Omega_{m-1}^i \end{cases} \text{ and } \bar{r}^i : \ell^2(\Omega_m^0) \rightarrow \ell^2(\Omega_{m-1}^i) \text{ by } (\bar{r}^i u)(x) = u(x).$$

Then $\bar{r}^i = (\bar{p}^i)^*$, $p^i = \frac{1}{2} [(-1)^i \quad 2 \quad (-1)^i] \bar{p}^i$ and $r^i = \bar{r}^i \frac{1}{2} [(-1)^i \quad 2 \quad (-1)^i]$.

The mass matrix M_m is given by the difference stencil $\frac{1}{6} h_m [1 \quad 4 \quad 1]$. It satisfies the relation

$$(12) \quad (M_m - h_m I) p^0 = \frac{1}{4} \bar{p}^0 (M_{m-1} - h_{m-1} I).$$

Since $r^1 p^0 = 0$ (use (1)), we obtain

$$r^1 M_k \underbrace{p^0 \cdots p^0}_{(k-l) \times} = r^1 (M_k - h_k I) \underbrace{p^0 \cdots p^0}_{(k-l) \times} = r^1 \left(\frac{1}{4}\right)^{k-l} \underbrace{\bar{p}^0 \cdots \bar{p}^0}_{(k-l) \times} (M_l - h_l I).$$

The set $\{\psi_m^{(j)} : x \mapsto \sqrt{2h_m} \sin(\pi j x)\}_{j \in \{1, \dots, n_m\}}$, where $n_m := h_m^{-1} - 1$, forms an orthonormal basis of $\ell^2(\Omega_m^0)$ that consists of eigenvectors $\psi_m^{(j)}$ of M_m and $\frac{1}{2} [(-1)^i \quad 2 \quad (-1)^i]$ with eigenvalues $\frac{1}{3} h_m (2 + \cos(\pi j h_m))$ and $(1 + (-1)^i \cos(\pi j h_m))$, respectively. There holds

$$\bar{p}^0 \psi_{m-1}^{(j)} = \frac{1}{2} \sqrt{2} (\psi_m^{(j)} - \psi_m^{(n_m+1-j)}) \quad (\{j \in \{1, \dots, n_{m-1}\}\}).$$

For r^1, p^1 , we have

$$\bar{r}^1 \psi_m^{(j)} = \bar{r}^1 \psi_m^{(n_m+1-j)} = \psi_m^{(j)}|_{\Omega_{m-1}^1} \text{ and } \bar{p}^1 (\psi_m^{(j)}|_{\Omega_{m-1}^1}) = \frac{1}{2} (\psi_m^{(j)} + \psi_m^{(n_m+1-j)})$$

($j \in \{1, \dots, n_{m-1} + 1\}$). From

$$\begin{aligned} \langle \psi_m^{(j)}|_{\Omega_{m-1}^1}, \psi_m^{(i)}|_{\Omega_{m-1}^1} \rangle &= \langle \bar{r}^1 \psi_m^{(j)}, \bar{r}^1 \psi_m^{(i)} \rangle \\ &= \langle \frac{1}{2}(\psi_m^{(j)} + \psi_m^{(n_{m-1}+1-j)}), \psi_m^{(i)} \rangle = \begin{cases} 0, & i \neq j \in \{1, \dots, n_{m-1} + 1\}, \\ \frac{1}{2}, & i = j \in \{1, \dots, n_{m-1}\}, \\ 1, & i = j = n_{m-1} + 1, \end{cases} \end{aligned}$$

we have that $\{\sqrt{2}\psi_m^{(j)}|_{\Omega_{m-1}^1} : j \in \{1, \dots, n_{m-1}\}\} \cup \{\psi_m^{(n_{m-1}+1)}|_{\Omega_{m-1}^1}\}$ is an orthonormal basis of $\ell^2(\Omega_{m-1}^1)$. Note that $\text{span}\{\psi_m^{(n_{m-1}+1)}\} \perp \text{range } p^0 : \ell^2(\Omega_{m-1}^0) \rightarrow \ell^2(\Omega_m^0)$, which means that the different scaling of $\psi_m^{(n_{m-1}+1)}|_{\Omega_{m-1}^1}$ does not enter our computations.

After a basis transformation to the orthonormal bases of $\ell^2(\Omega_l^0)$ and $\ell^2(\Omega_k^1)$, straightforward computations using the relations above now show that for $l = k - 1$,

$$\begin{aligned} \theta_{kk-1} &\leq \| (r^1 M_k p^1)^{-\frac{1}{2}} r^1 \frac{1}{4} \bar{p}^0 (M_{k-1} - h_{k-1} I) M_{k-1}^{-\frac{1}{2}} \| \\ &= \max \left\{ \frac{x(1-x^2)}{2\sqrt{1+2x^2}} : x = \cos(\pi j h_k), j \in \{1, \dots, n_{k-1}\} \right\} \\ &\leq \eta := \max \left\{ \frac{x(1-x^2)}{2\sqrt{1+2x^2}} : x \in [0, 1] \right\} \approx .153 \end{aligned}$$

and that for $l < k - 1$,

$$\begin{aligned} \theta_{kl} &\leq \| (r^1 M_k p^1)^{-\frac{1}{2}} r^1 \frac{1}{4} \bar{p}^0 \| \| \frac{1}{4} \underbrace{\bar{p}^0 \dots \bar{p}^0}_{(k-l-1) \times} \| \| (M_l - h_l I) M_l^{-\frac{1}{2}} \| \\ &= \max \left\{ \frac{\sqrt{6}|x|}{8\sqrt{h_k}} : x = \cos(\pi j h_k), j \in \{1, \dots, n_{k-1}\} \right\} \\ &\quad \cdot \frac{1}{4}^{k-1-l} \cdot \max \left\{ \frac{|1-x|\sqrt{h_l}}{(2+x)\sqrt{3}} : x = \cos(\pi j h_l), j \in \{1, \dots, n_l\} \right\} \\ &\leq \frac{\sqrt{6}}{8\sqrt{h_k}} \cdot \frac{1}{4}^{k-1-l} \cdot \frac{2\sqrt{h_l}}{\sqrt{3}} = \sqrt{2} \left(\frac{1}{4}\sqrt{2}\right)^{k-l}. \end{aligned}$$

By using Lemma 3.8 and $\rho(\Theta - I) \leq \|\Theta - I\|_\infty \leq 2(\eta + \sqrt{2} \sum_{n \geq 2} (\frac{1}{4}\sqrt{2})^n) \approx .85 < 1$, we conclude that (9) is valid.

3.4. An H^1 -equivalent norm on \mathcal{M}_J . A consequence of (9) is that

$$(13) \quad \|Z_l^{(k)}\|_{L^2 \leftarrow L^2} \lesssim 1 \quad (\text{uniformly in } k \geq l).$$

Let $Q_k : L^2(\Omega) \rightarrow \mathcal{M}_k$ be the L^2 -orthogonal projection on \mathcal{M}_k . Then it is well known that

$$(14) \quad \|I - Q_k\|_{L^2 \leftarrow H^1} \lesssim h_k.$$

From the *inverse estimate*

$$(15) \quad \|\cdot\|_{H^1} \lesssim h_k^{-1} \|\cdot\|_{L^2} \quad \text{on } \mathcal{M}_k,$$

we have

$$(16) \quad \|Q_k\|_{H^1 \leftarrow H^1} \lesssim 1.$$

The estimate (14) also implies that for $u \in \mathcal{V}_k$,

$$(17) \quad \|u\|_{L^2} \approx \|u\|_{\mathcal{M}_k} = \|(I - Y_{k-1})u\|_{\mathcal{M}_k} \\ \leq \|(I - Q_{k-1})u\|_{\mathcal{M}_k} \approx \|(I - Q_{k-1})u\|_{L^2} \lesssim h_{k-1}\|u\|_{H^1}.$$

In [12, Appendix], a compact proof is given of

$$(18) \quad \sum_{k=1}^J \|(Q_k - Q_{k-1})u\|_{H^1}^2 \approx \|u\|_{H^1}^2 \quad (u \in \mathcal{M}_J)$$

(where $Q_{-1} := 0$). As we will see, this proof with $Z_k^{(J)}$ playing the role of $Q_k - Q_{k-1}$ will also yield (10).

Theorem 3.10. *We have $\sum_{k=0}^J \|Z_k^{(J)}u\|_{H^1}^2 \approx \|u\|_{H^1}^2$ ($u \in \mathcal{M}_J$).*

Proof (based on [12, Appendix]). Let $u \in \mathcal{M}_J$ and $u_l = (Q_l - Q_{l-1})u$. Then $u_l \in \mathcal{M}_l$, and so for $l < k \leq J$, we have $Z_k^{(J)}u_l = 0$. For $k \leq l$, it follows from (15), (13) and (14) that

$$\|Z_k^{(J)}u_l\|_{H^1} \lesssim h_k^{-1}\|Z_k^{(J)}u_l\|_{L^2} \lesssim h_k^{-1}\|u_l\|_{L^2} \lesssim h_k^{-1}h_l\|u_l\|_{H^1}.$$

Let $l \wedge m = \min\{l, m\}$. Writing $u = \sum_{l=0}^J u_l$, we get

$$\begin{aligned} \sum_{k=0}^J \|Z_k^{(J)}u\|_{H^1}^2 &= \sum_{k=0}^J \sum_{l,m=k}^J (Z_k^{(J)}u_l, Z_k^{(J)}u_m)_{H^1} = \sum_{l,m=0}^J \sum_{k=0}^{l \wedge m} (Z_k^{(J)}u_l, Z_k^{(J)}u_m)_{H^1} \\ &\leq \sum_{l,m=0}^J \sum_{k=0}^{l \wedge m} h_k^{-2}h_lh_m\|u_l\|_{H^1}\|u_m\|_{H^1} \lesssim \sum_{l,m=0}^J h_{l \wedge m}^{-2}h_lh_m\|u_l\|_{H^1}\|u_m\|_{H^1} \\ &\lesssim \sum_{l=0}^J \|u_l\|_{H^1}^2 \approx \|u\|_{H^1}^2 \end{aligned}$$

by (18).

In [12, Lemma 6.1], it was proved that for $k \leq l$, $u \in \mathcal{M}_k$ and $v \in \mathcal{M}_l$,

$$|(u, v)_{H^1}| \lesssim (h_k h_l)^{-\frac{1}{2}} \|u\|_{H^1} \|v\|_{L^2}.$$

Using this and (17), we obtain for $u \in \mathcal{M}_J$

$$\begin{aligned} \|u\|_{H^1}^2 &= \sum_{k,l=0}^J (Z_k^{(J)}u, Z_l^{(J)}u)_{H^1} \\ &\lesssim \sum_{k,l=0}^J (h_k h_l)^{-\frac{1}{2}} \min\{h_{k-1}, h_{l-1}\} \|Z_k^{(J)}u\|_{H^1} \|Z_l^{(J)}u\|_{H^1} \lesssim \sum_{k=0}^J \|Z_k^{(J)}u\|_{H^1}^2, \end{aligned}$$

which completes the proof.

ACKNOWLEDGEMENTS

The author wishes to acknowledge the hospitality of the Institut für Informatik und Praktische Mathematik, Universität Kiel, where much of this work was performed. Furthermore, he would like to thank Professors W. Hackbusch (Kiel) and M. Dryja (Warsaw) and Dr. A. Reusken (Eindhoven) for fruitful discussions.

REFERENCES

1. J.H. Bramble and J.E. Pasciak, *New convergence estimates for multigrid algorithms*, Math. Comp. **49** (1987), 311–329. MR **89b**:65234
2. W. Hackbusch, *A new approach to robust multi-grid methods*, ICIAM '87: *Proceedings of the First International Conference on Industrial and Applied Mathematics* (Philadelphia) (J. McKenna and R. Temam, eds.), SIAM, 1988. MR **90f**:65193
3. ———, *The frequency decomposition multi-grid method I. Application to anisotropic equations*, Numer. Math. **56** (1989), 229–245. MR **90i**:65212
4. ———, *Iterative lösung großer schwachbesetzter gleichungssysteme*, B.G. Teubner, Stuttgart, 1991. MR **92i**:65002
5. ———, *The frequency decomposition multi-grid method II. Convergence analysis based on the additive Schwarz method*, Numer. Math. **63** (1992), 433–453. MR **93j**:65203
6. ———, *The frequency decomposition multi-grid method*, Multigrid Methods IV, *Proceedings of the Fourth European Multigrid Conference, Amsterdam, July 6-9, 1993* (Basel) (P.W. Hemker and P. Wesseling, eds.), Birkhäuser, 1994, pp. 43–56. MR **95h**:65095
7. J. Junkherr, *Multigrid methods for weakly singular integral equations of the first kind*, Ph.D. thesis, Christian-Albrechts-Universität Kiel, 1994.
8. W. Mulder, *A new multigrid approach to convection problems*, J. Comp. Phys. **83** (1989), 303–323. MR **90f**:76013
9. N.H. Naik and J. Van Rosendale, *The improved robustness of multigrid elliptic solvers based on multiple semicoarsened grids*, SIAM J. Numer. Anal. **30** (1993), 215–229. MR **94b**:65159
10. C.W. Oosterlee and P. Wesseling, *On the robustness of a multiple semi-coarsened grid method*, ZAMM **75** (1995), no. 4, 251–257. CMP 95:12
11. R.P. Stevenson, *Robustness of the additive and multiplicative frequency decomposition multi-level method*, Computing **54** (1995), 331–346. CMP 95:13
12. J. Xu, *Iterative methods by space decomposition and subspace correction*, SIAM Rev. **34** (1992), 581–613. MR **93k**:65029
13. H. Yserentant, *Old and new convergence proofs for multigrid methods*, Acta Numerica (1993), 285–326. MR **94i**:65128

DEPARTMENT OF MATHEMATICS, NIJMEGEN UNIVERSITY, TOERNOOIVELD 1, 6525 ED NIJMEGEN, THE NETHERLANDS

E-mail address: `stevens@sci.kun.nl`