
Logics of formal inconsistency based on distributive involutive residuated lattices

F. ESTEVA, *Artificial Intelligence Research Institute (IIIA), Spanish National Research Council (CSIC), Campus UAB s/n, Bellaterra 08193, Spain.*
E-mail: esteva@iiia.csic.es

A. FIGALLO-ORELLANO, *Departamento de Matemática, Universidad Nacional del Sur (UNS), Bahía Blanca 8000, Argentina and Centre for Logic, Epistemology, and The History of Science (CLE), University of Campinas (UNICAMP), Campinas, SP, 13083-859, Brazil.*
E-mail: aldofigallo@gmail.com

T. FLAMINIO, *Artificial Intelligence Research Institute (IIIA), Spanish National Research Council (CSIC), Campus UAB s/n, Bellaterra 08193, Spain.*
E-mail: tommaso@iiia.csic.es

L. GODO, *Artificial Intelligence Research Institute (IIIA), Spanish National Research Council (CSIC), Campus UAB s/n, Bellaterra 08193, Spain.*
E-mail: godo@iiia.csic.es

Abstract

The aim of this paper is to develop an algebraic and logical study of certain paraconsistent systems, from the family of the logics of formal inconsistency (LFIs), which are definable from the degree-preserving companions of logics of distributive involutive residuated lattices (dIRLs) with a consistency operator, the latter including as particular cases, Nelson logic (NL), involutive monoidal t-norm based logic (IMTL) or nilpotent minimum (NM) logic. To this end, we first algebraically study enriched dIRLs with suitable consistency operators. In fact, we consider three classes of consistency operators, leading respectively to three subquasivarieties of such expanded residuated lattices. We characterize the simple and subdirectly irreducible members of these quasivarieties, and we extend Sendlewski's representation results for the case of Nelson lattices with consistency operators. Finally, we define and axiomatize the logics of three quasivarieties of dIRLs and their corresponding degree-preserving companions that belong to the family of LFIs.

Keywords: Logics of formal inconsistency, paraconsistent logics, degree-preserving logics, distributive involutive residuated lattices, Nelson lattices.

1 Introduction

The aim of this paper is to develop an algebraic and logical study of paraconsistent systems definable from the degree-preserving companions of logics of distributive involutive residuated lattices (dIRLs) with a consistency operator. The initial motivation comes from different considerations relating paraconsistency and Nelson's constructive logic with strong negation.

In the 1950's, constructive logic with strong negation, nowadays commonly known under the name of *Nelson logic* (even also called N3), was formulated by Nelson and Markov as a result of

2 Logics of formal inconsistency on *dIRL*

certain philosophical objections to the intuitionistic negation pointed out by Rasiowa [43], see also her celebrated book [44, Ch. XII]. The criticism concerned its disadvantageous non-constructive property, namely that the derivability of the formula $\neg(\alpha \wedge \beta)$ in an intuitionistic propositional calculus does not imply that at least one of the formulas $\neg\alpha, \neg\beta$ is derivable.

Although *Nelson algebras*, the algebraic semantics of Nelson logic developed by Rasiowa [43, 44], were not originally presented as a subclass of residuated lattices, in 2008 Spinks and Veroff [47, 48] have shown that Nelson logic is indeed a substructural logic. More precisely, they show that Nelson algebras are termwise equivalent to certain involutive, bounded, commutative and integral residuated lattices, called *Nelson (residuated) lattices*. Busaniche and Cignoli [7] have further contributed to the algebraic study of Nelson lattices.

A paraconsistent version of Nelson logic was first introduced in [1], where the authors observe that the weaker system obtained from Nelson logic by deleting the axiom schema $\varphi \rightarrow (\sim\varphi \rightarrow \psi)$ could be used to reason under inconsistency without incurring in a trivial logic. Semantics for this paraconsistent version of Nelson logic have been studied by Odintsov in [38, 39], who calls the logic **N4**, in terms of what is known in the literature as *Fidel structures* [25] and *twist-structures* [24, 51]. Later on, Busaniche and Cignoli [8] provided another algebraic semantics under the umbrella of non-integral commutative residuated lattices with involution. More recently, Carnielli and Rodrigues [15] show that **N4** is in fact equivalent to a paraconsistent logic of evidence, called BLE (for basic logic of evidence), for which they provide a semantics based on non truth-functional evaluations.

Our initial interest also was on paraconsistent variants of Nelson logic, but taking a different road. Indeed, our idea was to consider paraconsistent logics from the family of logics of formal inconsistency (LFIs), introduced by Carnielli and Marcos in 2000 (see e.g. [13]), and also studied e.g. by Avron [2, 3]. LFIs constitute a generalization of da Costa's C-systems [20, 21]. The main characteristic of these logics is that they internalize in the object language a notion of *consistency* by means of a specific connective \circ (primitive or definable) in the following sense: although LFIs are not explosive in general, meaning that for at least a formula φ the theory $\{\varphi, \sim\varphi\}$ is consistent, the connective \circ allows to recover the explosion property from a formula ψ and its negation $\sim\psi$ whenever they are retained to be *consistent*, that is to say, whenever ψ falls under the scope of \circ . In other words, even if $\{\varphi, \sim\varphi\}$ is not explosive, $\{\varphi, \sim\varphi, \circ\varphi\}$ trivializes because $\circ\varphi$ states that φ is consistent. It is worth noticing that, in fact, Carnielli and Rodrigues have already introduced in [15] an LFI based on paraconsistent Nelson logic **N4**, called **LET_J** (for logic of evidence and truth), by adding a consistency-like operation to BLE, their paraconsistent and paracomplete basic logic of evidence.

There is however another approach to define LFIs based on Nelson logic, and more generally, on logics of involutive residuated lattices. In general, given a quasivariety of bounded residuated lattices \mathbb{Q} , its corresponding logic is given by the usual (non-paraconsistent) truth-preserving notion of logical consequence $\models_{\mathbb{Q}}$, i.e. a formula φ follows from a set of formulas Γ , written $\Gamma \models_{\mathbb{Q}} \varphi$, if $e(\varphi) = 1$ whenever $e(\psi) = 1$ for all $\psi \in \Gamma$ and for all evaluations e on every algebra \mathbf{A} in the quasivariety \mathbb{Q} . A weaker notion of consequence companion of \models is the one called *degree-preserving* logical consequence, where a formula φ follows from a set of formulas Γ , written $\Gamma \models_{\mathbb{Q}}^{\vee} \varphi$ if $e(\varphi) \geq a$ whenever $e(\psi) \geq a$ for all $\psi \in \Gamma$ and for all evaluations e on every algebra $\mathbf{A} \in \mathbb{Q}$ and every element $a \in A$. This weaker notion of logical consequence, firstly introduced in [52], has been further investigated in e.g. [6, 26, 27]. The point is that, as observed in [22] and unlike the truth-preserving logics, the degree-preserving companions of a large class of fuzzy logics (i.e. logics of varieties of prelinear residuated lattices), in particular those with an involutive negation, are paraconsistent. Interestingly, this is also the case of Nelson logic and, more generally, the logics of varieties of involutive residuated lattices. However, although the degree-preserving companions of these logics are paraconsistent, they are not expressive enough to define a consistency connective

\circ in its own language (see [22, Corollary 2 and Example 2] and Remark 3.6 below), and hence they are not LFIs. A further step was made in [19] where the authors introduce a wide class of LFIs by first expanding (non pseudo-complemented) fuzzy logics with a consistency operator while preserving the semi-linearity of the logics, and then considering their corresponding degree-preserving companions.

Given all these antecedents, our initial aim was to follow a similar approach to [19] to expand the language of Nelson logic by a primitive consistency connective \circ and to add axioms and rules encoding suitable postulates in such a way that the corresponding degree-preserving companion of the logic arising in this way be an LFI. However, in doing so, we realized that, essentially, all the definitions and results we got for Nelson logic and lattices, as underlying logical and algebraic framework, remain valid for the more general framework of involutive, distributive (bounded, commutative and integral) residuated lattices and their logics. Therefore, in this paper, we have finally chosen to present our algebraic and logical results in this more general setting and only particularize to Nelson logic when necessary.

The organization of this paper is as follows. In Section 2, the basic notions about varieties of involutive residuated lattices and their logics are recalled. Section 3 contains the main algebraic results and it is divided into three subsections, in each of which we will introduce a specific type of consistency operator and we will present results concerning the so arising quasivarieties. In Section 4 we consider the particular case of Nelson lattices with a consistency operator, for which we prove structural results and their relation to Heyting algebras with dual pseudocomplement. In particular, in Subsection 4.2 we also present additional results on subdirectly irreducible and simple Nelson lattices with a consistency operator. Section 5 considers the logical counterparts of the classes of algebras studied in Section 3 and their associated LFI over the degree-preserving companions of the former. In the last subsection of Section 5, we will point out that, besides being LFIs, our logics are logics of formal undeterminedness¹ (LFUs) as well, and we analyse the relationships between these two notions. We conclude in Section 6 collecting some final remarks and our future work on the topics of this paper.

2 Preliminary notions

Recall that a *commutative, integral, bounded residuated lattice*, which we will simply call *residuated lattice*, is an algebra $\mathbf{A} = \langle A, \wedge, \vee, *, \rightarrow, 0, 1 \rangle$ of type $(2, 2, 2, 2, 0, 0)$ such that $\langle A, *, 1 \rangle$ is a commutative monoid, $\langle A, \wedge, \vee, 0, 1 \rangle$ is a bounded lattice with least element 0 and greatest element 1 and such that the following residuation condition holds: $x * y \leq z$ iff $x \leq y \rightarrow z$, where x, y, z denote arbitrary elements of A and \leq is the order given by the lattice structure. Since we assume the neutral element of the monoid reduct coincides with the greatest element of its lattice reduct, we have that $x \leq y$ iff $x \rightarrow y = 1$.

It is well known that the class \mathbb{RL} of residuated lattices forms a variety, which is related to different and well-known varieties studied in substructural and fuzzy logics literature. In particular, \mathbb{RL} coincides with the variety of \mathbf{FL}_{ew} algebras of [28]. According to the denotational conventions of [28], \mathbf{FL} refers to the ‘Full Lambek calculus’, which is the base system and associated algebras, and subindices indicate several axiomatic extensions with properties such as exchange (e) or weakening (w).

¹A logic of formal undeterminedness is a paracomplete logic with a unary *determinedness* operator that controls the law of the excluded middle, it is a sort of dual notion of an LFI [35].

4 Logics of formal inconsistency on dIRL

A residuated lattice is called *distributive* if its lattice reduct is a distributive lattice, that is to say, if it satisfies the distributivity equations:

$$x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z) \text{ and } x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z). \quad (\text{Dist})$$

The variety of distributive residuated lattices will be denoted by $d\mathbb{R}\mathbb{L}$.

A residuated lattice is called *involutive* if it satisfies the double negation equation:

$$\sim\sim x = x, \quad (\text{Inv})$$

where $\sim x$ is $x \rightarrow 0$. In every involutive residuated lattice, it is possible to prove that $x * y = \sim(x \rightarrow \sim y)$ and $x \rightarrow y = \sim(x * \sim y)$. Clearly involutive residuated lattices form a variety that will be denoted by $\mathbb{I}\mathbb{R}\mathbb{L}$.²

The variety of distributive and involutive residuated lattices (dIRL algebras for short) will be henceforth denoted by $d\mathbb{I}\mathbb{R}\mathbb{L}$.

In this paper we will also consider some proper subvarieties of $d\mathbb{I}\mathbb{R}\mathbb{L}$, in particular:

- The variety $\mathbb{N}\mathbb{L}$ of *Nelson residuated lattices*, defined within $\mathbb{I}\mathbb{R}\mathbb{L}$ by the so-called Nelson equation:

$$(((x * x) \rightarrow y) \wedge ((\sim y * \sim y) \rightarrow \sim x)) \rightarrow (x \rightarrow y) = 1. \quad (\text{Nel})$$

- The variety $\mathbb{I}\mathbb{M}\mathbb{T}\mathbb{L}$ of *involutive monoidal t-norm based algebras* (or IMTL-algebras for short), which can be defined as the proper subvariety of $\mathbb{I}\mathbb{R}\mathbb{L}$ (and of $d\mathbb{I}\mathbb{R}\mathbb{L}$) of those algebras satisfying the *prelinearity equation*:

$$(x \rightarrow y) \vee (y \rightarrow x) = 1. \quad (\text{Pre})$$

- The variety $\mathbb{N}\mathbb{M}$ of *nilpotent minimum algebras* (or NM-algebras for short), which is identified as proper subvariety of $\mathbb{I}\mathbb{M}\mathbb{T}\mathbb{L}$ by the following equation:

$$(x * y \rightarrow 0) \vee (x \wedge y \rightarrow x * y) = 1, \quad (\text{NM})$$

or equivalently, as shown in [7], the subvariety of $\mathbb{N}\mathbb{L}$ of those algebras satisfying the prelinearity equation.

Notice that Nelson, IMTL and NM algebras can be axiomatized directly within $\mathbb{I}\mathbb{R}\mathbb{L}$ without explicitly requiring the distributivity equations to hold.

In Figure 1 we represent the graph of the above considered subvarieties of $\mathbb{I}\mathbb{R}\mathbb{L}$ together with their characteristic axioms.

2.1 Truth-preserving and degree-preserving logics of residuated lattices

The substructural logic that is complete with respect to the variety of (bounded, commutative, integral) residuated lattices is the so-called Full Lambek calculus extended with exchange (commutativity) and weakening (integrality), FL_{ew} , see e.g. [28].³ The language of FL_{ew} consists of denumerably many propositional variables p_1, p_2, \dots , binary connectives $\wedge, \vee, \&, \rightarrow$ and the truth constant \perp . Formulas, which will be denoted by lower case greek letters ϕ, ψ, \dots , are defined by induction as usual. Further connectives and constants are definable; in particular, $\neg\phi$ stands for

²In the literature, involutive residuated lattices have been called involutive FL_{ew} algebras (IFL_{ew} algebras), see e.g. [32], and they are also known as the algebras of the *affine linear logic without exponentials*, see e.g. [5]. In this paper we shall adopt the notation $\mathbb{I}\mathbb{R}\mathbb{L}$ algebras to denote them without danger of confusion.

³An equivalent system, called monoidal logic, was previously defined and studied by Höhle in [30].

6 Logics of formal inconsistency on dIRL

This paper will be mainly concerned with the logic dIRL and some of its axiomatic extensions. Hence, distributivity will always be assumed to hold. In particular we will consider *Nelson logic* NL obtained by extending dIRL by the Nelson axiom:

$$\text{Ax16. } (((\psi \& \psi) \rightarrow \phi) \wedge ((\neg \phi \& \neg \phi) \rightarrow \neg \psi)) \rightarrow (\psi \rightarrow \phi).$$

The logic IMTL, as introduced in [23], is the axiomatic extension of the logic MTL (the logic of the variety of prelinear residuated lattices) with the axiom (Ax14) or, equivalently, obtained by extending IRL by the prelinearity axiom:

$$\text{Ax17. } (\phi \rightarrow \psi) \vee (\psi \rightarrow \phi).$$

Finally, the *nilpotent minimum* logic NM, introduced also in [23], is the axiomatic extension of IMTL by the axiom

$$\text{Ax18. } (\phi \& \psi \rightarrow \perp) \vee (\phi \wedge \psi \rightarrow \phi \& \psi).$$

Equivalently, NM can be obtained as the axiomatic extension of Nelson logic NL by the prelinearity axiom.

We will denote by \vdash_L the notion of proof for each logic $L \in \{\text{IRL}, \text{dIRL}, \text{IMTL}, \text{NL}, \text{NM}\}$ defined as usual from the corresponding sets of axioms described above and the inference rule of Modus Ponens (MP).

Each of these logics L is algebraizable (as all of them are axiomatic extensions of FL_{ew}), and thus it has an equivalent algebraic semantics given by the corresponding variety of L algebras introduced before, and which brings the same name. This means that the truth-preserving (finitary) consequence relation \models_L induced by the variety of L algebras, defined as

$$\Gamma \models_L \varphi \quad \text{iff} \quad \text{for every } L \text{ algebra } \mathcal{A} \text{ and every } \mathcal{A} \text{ evaluation } e, \\ \text{if } e(\psi) = 1 \text{ for every } \psi \in \Gamma, \text{ then } e(\varphi) = 1,$$

is such that \vdash_L is sound and complete w.r.t. \models_L .

Moreover, for each such a logic L , in [6, 27] the authors introduce a companion logic denoted L^\leq , whose associated consequence relation, which will be denoted as \models_L^\leq , has the following semantics: for every set of formulas $\Gamma \cup \{\varphi\}$,

$$\Gamma \models_L^\leq \varphi \quad \text{iff} \quad \text{for every } L \text{ algebra } \mathcal{A}, \text{ every } a \in \mathcal{A}, \text{ and every } \mathcal{A} \text{ evaluation } e, \\ \text{if } a \leq e(\psi) \text{ for every } \psi \in \Gamma, \text{ then } a \leq e(\varphi).$$

By obvious reasons, L^\leq is known as the companion logic of L *preserving degrees of truth*, or the *degree-preserving companion* of L . It is not difficult to show that L and L^\leq have the same valid formulas (i.e. $\vdash_L \varphi$ iff $\vdash_{L^\leq} \varphi$), and that, for every finite set of formulas $\Gamma \cup \{\varphi\}$, the following property holds:

$$\Gamma \models_L^\leq \varphi \quad \text{iff} \quad \vdash_L \Gamma^\wedge \rightarrow \varphi,$$

where Γ^\wedge means $\gamma_1 \wedge \dots \wedge \gamma_k$ if $\Gamma = \{\gamma_1, \dots, \gamma_k\}$ (when Γ is empty then Γ^\wedge is taken as \top).

As it regards to axiomatization, if L is an axiomatic extension of FL_{ew} , then the logic L^\leq admits a Hilbert-style axiomatization having the same axioms as L and the following deduction rules [6]:

- (Adj- \wedge) from φ and ψ derive $\varphi \wedge \psi$
- (MP- r) if $\vdash_L \varphi \rightarrow \psi$, then from φ and $\varphi \rightarrow \psi$, derive ψ

Note that (MP-*r*) is a *restricted form* of the Modus Ponens rule, as it is only applicable when $\varphi \rightarrow \psi$ is a theorem of L . Thus, although L and L^{\leq} share the same theorems, L^{\leq} is in fact a weaker logic than L .

If the set of theorems of L is decidable, then the above systems of axioms and rules provide a recursive Hilbert-style axiomatization of L^{\leq} .

In the more general case of L being not an axiomatic extension but a finitary Rasiowa-implicative expansion of FL_{ew} (i.e. L may have new inference rules and hence its algebraic semantics may be a sub-quasivariety of \mathbb{RL}), the definition of the degree-preserving companion L^{\leq} keeps being the same as above. However, the axiomatization needs to be tuned. Assume the new inference rules of L are as follows:

(R_i) from Γ_i derive φ_i ,

for $i \in I$. Then, following the same idea of [6, Th. 2.12], one can show the following generalized result about the axiomatization of L^{\leq} .

PROPOSITION 2.1

Let L be an expansion of FL_{ew} , with the above set of new inference rules $\{(R_i)\}_{i \in I}$. Then L^{\leq} is axiomatized by the axioms of L , the inference rules (Adj- \wedge) and (MP-*r*) and the following restricted inference rules:

(R_i -*r*) If $\vdash_L \Gamma_i$, then from Γ_i derive φ_i

for each $i \in I$.

The proof is completely analogous to the one in [6] in the context of expansions of MTL and it is omitted.

3 Distributive involutive residuated lattices expanded by a consistency operator

Paraconsistency is the study of logics having a negation operator \sim that are *not explosive* with respect to that negation; that is to say, logics for which there exists at least a formula ϕ such that the theory $\{\phi, \sim\phi\}$ does not entail any other formula. Therefore, a paraconsistent logic is a logic having at least a contradictory but non-trivial theory.

Among the plethora of paraconsistent logics proposed in the literature, the LFIs (see, for instance, [12, 13]) play an important role, since they internalize in the object language the very notion of *consistency*⁴ by means of a specific connective, primitive or not. This generalizes the strategy of da Costa, who introduced in [21] the well-known hierarchy of systems C_n , for $n > 0$. Briefly said, LFIs have a non-explosive negation \sim , as well as a (primitive or derived) *consistency connective* \circ , which allows to recover the explosion law in a controlled way.

Let Σ be a propositional signature, which contains a negation \sim and a primitive or defined unary connective \circ , let $\mathcal{V} = \{p_1, p_2, \dots\}$ be a denumerable set of propositional variables and let $L = (\Sigma, \vdash)$ be a Tarskian, finitary and structural logic defined over Σ and \mathcal{V} . Then, according to e.g. [12], we have the following definition.

DEFINITION 3.1

L is said to be a LFI with respect to \sim and \circ if the following holds:

- (i) $\varphi, \sim\varphi \not\vdash \psi$ for some φ and ψ ;

⁴A formula ϕ is named consistent in a paraconsistent logic when $\{\phi, \sim\phi\}$ is an explosive theory.

8 Logics of formal inconsistency on dIRL

- (ii) there are two formulas α and β such that
 - (ii.a) $\circ\alpha, \alpha \not\vdash \beta$;
 - (ii.b) $\circ\alpha, \sim\alpha \not\vdash \beta$;
- (iii) $\circ\varphi, \varphi, \sim\varphi \vdash \psi$ for every φ and ψ .

Moreover, in [12], the authors also consider the following stronger notion of LFIs.

DEFINITION 3.2

\mathbf{L} is said to be a *strong LFI* with respect to \sim and \circ if the following holds:

- (i) if p and q are two different propositional variables then
 - (i.a) $p, \sim p \not\vdash q$
 - (i.b) $\circ p, p \not\vdash q$
 - (i.c) $\circ p, \sim p \not\vdash q$
- (ii) $\circ\varphi, \varphi, \sim\varphi \vdash \psi$ for every φ and ψ .

Our aim is to consider different possibilities of defining (strong) LFIs over the degree-preserving logic companion of the logic dIRL. To this end, we will first study suitable expansions of dIRL algebras by a new consistency operator \circ in such a way that their degree-preserving companions are LFIs. Actually, we will consider three different axiomatic definitions of varying strength. Similar ideas of having LFIs over a degree-preserving logic companion of other class of algebras can be seen in [19] and [11].

Let \mathbf{A} be an involutive residuated lattice and let $\circ : A \rightarrow A$ be a unary operation on \mathbf{A} . With the above goal in mind, we consider different properties we may ask to the \circ operation in order to be a suitable consistency operator.

It is clear that the minimal properties to require to \circ to be a consistency operator are:

- ($\circ 0$) $\circ(1) = \circ(0) = 1$
- ($\circ 1$) $x \wedge \sim x \wedge \circ(x) = 0$

However, these properties turn out to be a weak specification in the sense that, for a given dIRL \mathbf{A} , one can define many operations satisfying the above sets of properties, in particular, one can always define a minimal operation by letting $\circ(x) = 0$ for all $x \in A \setminus \{0, 1\}$. A natural way out is to require that \circ provides the maximum value in A such that ($\circ 1$) is satisfied and hence we can define $\circ(x)$ to be the $\max\{z \in A \mid x \wedge \sim x \wedge z = 0\}$. In a sense, such an operator, if it exists, can be considered as the least committed one satisfying ($\circ 1$). This is formally achieved by considering the following additional requirement:

- ($\circ 2$) if $x \wedge \sim x \wedge y = 0$ then $y \leq \circ(x)$

Since Boolean elements are the prototypical examples of consistent and explosive elements, another reasonable property one can further require to \circ is to be a *Boolean operator*, that is to say, that for each $x \in A$ to require $\circ(x) \vee \sim\circ(x) = 1$. This can be achieved in at least two ways.

A first possibility is to define $\circ(x)$ as the maximum, among the set $B(\mathbf{A}) = \{x \in A \mid x \wedge \sim x = 0\}$ of Boolean elements of \mathbf{A} , satisfying the condition ($\circ 1$) above. In other words, to take $\circ(x) = \max\{z \in B(\mathbf{A}) \mid x \wedge \sim x \wedge z = 0\}$. This amounts to replace ($\circ 2$) by the following two new conditions:

- ($\circ 3$) if $x \wedge \sim x \wedge y = 0$ and $y \wedge \sim y = 0$ then $y \leq \circ(x)$
- ($\circ 4$) $\circ(x) \vee \sim\circ(x) = 1$

A second possibility is to consider an operator \circ such that $\circ(x)$ is, at the same time, both the $\max\{z \in A \mid x \wedge \sim x \wedge z = 0\}$ and a Boolean element. In fact this latter requirement differs from $\max\{z \in B(\mathbf{A}) \mid x \wedge \sim x \wedge z = 0\}$. This is achieved by asking \circ to satisfy the above conditions ($\circ 0$), ($\circ 1$), ($\circ 2$) and ($\circ 4$).

Actually, in this paper, we will study expansions of involutive residuated lattices with these three types of consistency operators \circ . Namely, besides satisfying ($\circ 0$) and ($\circ 1$), we will consider operators:

- (i) additionally satisfying ($\circ 2$), which we will call *maximal consistency operators* (**max**-consistency operators for short), in Subsection 3.1;
- (ii) additionally satisfying ($\circ 3$) and ($\circ 4$), which we will call *maximal Boolean consistency operators* (**maxB**-consistency operators for short), in Subsection 3.2; and
- (iii) additionally satisfying ($\circ 2$) and ($\circ 4$), which we will call *Boolean and maximal consistency operators* (**Bmax**-consistency operators, for short), in Subsection 3.3.

As one can already anticipate, **Bmax**-consistency operators are those operators that are both **max**- and **maxB**-consistency operators. We will formally show this in Subsection 3.3.

The following observations on congruences, filters and subdirect product decompositions will be useful in the rest of the paper.

As is well known, an *implicative filter* of a (bounded) residuated lattice \mathbf{A} is a subset $F \subseteq A$ such that $1 \in F$ and it is closed under modus ponens: $x \in F$ and $x \rightarrow y \in F$ imply $y \in F$.⁵ For each implicative filter F , the binary relation $\theta(F)$ defined by $(x, y) \in \theta(F)$ if and only if $x \rightarrow y, y \rightarrow x \in F$ is a congruence of the residuated lattice \mathbf{A} , and $F = \{z \in A : (z, 1) \in \theta(F)\}$. This is actually a one-to-one correspondence between the lattice of congruences and the lattice of implicative filters for the variety of bounded residuated lattices.

Moreover, since the classes of expansions of dIRLs with the above types of consistency operators involve not only equations but also quasi-equations, we will also deal with quasivarieties. In a quasivariety, congruences that allow for the decomposition of an algebra as a subdirect product of subdirectly irreducible components are required to satisfy an additional condition: the quotient of an algebra by a congruence has to belong to the quasivariety, see e.g. [42]. This condition is automatically satisfied in varieties but not in quasivarieties. Congruences satisfying this condition are usually called *Q-congruences*. Similarly, filters that are in a one-to-one correspondence between *Q*-congruences are implicative filters ‘closed’ by the quasiequations of the quasivariety and are called *Q-filters*.

3.1 Distributive involutive residuated lattices with **max**-consistency operators

We start by formally defining the first class of dIRLs with a consistency operator, namely with a **max**-consistency operator.

DEFINITION 3.3

A *dIRL with a **max**-consistency operator* (a *dIRL_c algebra* for short) is a pair (\mathbf{A}, \circ) where \mathbf{A} is a dIRL and $\circ : A \rightarrow A$ satisfies the following two conditions: for all $x, y, z \in A$,

- ($\circ 1$) $x \wedge \sim x \wedge \circ(x) = 0$
- ($\circ 2$) if $x \wedge \sim x \wedge y = 0$ then $y \leq \circ(x)$.

⁵Recall that an implicative filter of a residuated lattice \mathbf{A} can be equivalently defined as a subset $F \subseteq A$ such that $1 \in F$ and is closed by the monoidal operation $*$ of the residuated lattice.

10 Logics of formal inconsistency on dIRL

From the definition, it is clear that the class of dIRL_c algebras is a quasivariety.

It is easy to check that conditions (o1) and (o2) faithfully capture the expected behavior max-consistency operators as described above.

LEMMA 3.4

Let \mathbf{A} be a dIRL algebra, and let \circ be a unary operation on A . Then (\mathbf{A}, \circ) is a dIRL_c algebra iff, for any $x \in A$, $\circ(x) = \max\{z \in A \mid x \wedge \sim x \wedge z = 0\}$.

PROOF. Clearly, for a given $x \in A$, the set $\{z \in A \mid x \wedge \sim x \wedge z = 0\}$ is closed by \vee , it has $\circ(x)$ as an upper bound by (o2) and moreover $\circ(x)$ belongs to that set by (o1). \square

Conditions (o1) and (o2) are also enough to ensure that condition (o0) also holds in a dIRL_c algebra. This and other easy properties of dIRL_c algebras are displayed in the next lemma.

LEMMA 3.5

The following properties hold in a dIRL_c algebra (\mathbf{A}, \circ) :

- (i) $\circ(x) = \circ(\sim x) = \circ(x \wedge \sim x) = \circ(x \vee \sim x)$
- (ii) $\circ(x) = 1$ iff x is Boolean, in particular $\circ(1) = \circ(0) = 1$.

PROOF. (i) follows from Lemma 3.4 just noticing that $\sim \sim x = x$, and $\circ(1) = 1$ follows from (o2) by taking $x = y = 1$. (ii) also follows from Lemma 3.4 by noticing that $x \wedge \sim x = 0$ for any Boolean element x . \square

As an example of a finite dIRL_c algebra, let $L = \{0, a, b, c, d, e, f, 1\}$ and consider the lattice (L, \wedge, \vee) represented in the upper part of Figure 2. Then the algebra $\mathbf{L} = (L, \wedge, \vee, *, \rightarrow, \sim, 0, 1)$, where $*$ and \sim are those specified in the tables of Figure 2 and where $x \rightarrow y = \sim(x * \sim y)$, is a dIRL (it is indeed a finite *nilpotent minimum* algebra) and (\mathbf{L}, \circ) is a dIRL_c algebra.

REMARK 3.6


It follows from Lemma 3.4 that if a max-consistency operator is definable in an involutive residuated lattice it is uniquely determined. Moreover, it also follows that the max-consistency operator is always definable in every finite involutive residuated lattice. However, it is not always the case in infinite involutive residuated lattices. Actually, also according to Lemma 3.4 above, such an operator is definable if and only if all elements of the involutive residuated lattice of the form $x \wedge \sim x$ admit a pseudo-complement. The following are examples of infinite dIRLs in which the consistency operator cannot be defined.

(1) Let \mathbf{A} be the an algebra over $[0, 1] \setminus \{\frac{1}{2}\}$ with the nilpotent minimum operations: $x * y = \min(x, y)$ if $x + y > 1$ and $x * y = 0$ otherwise; and $\sim x = 1 - x$. Let $A^+ = \{x \in A \mid x > 1/2\}$ and $A^- = \{x \in A \mid x < 1/2\}$ be the sets of positive and negative elements of \mathbf{A} respectively. Further, let \mathbf{B} be the nilpotent minimum subalgebra of $\mathbf{A} \times \mathbf{A}$ defined on the sublattice $B = (A^+ \times A^+) \cup (A^- \times A^-)$. Take an element $(x, 1) \in B$ such that $x \in A^+$. An easy computation shows that $\circ((x, 1))$ does not exist.

(2) Let \mathbf{A} be the 1-generated free MV algebra [17], i.e. up to isomorphism, the algebra of continuous piecewise linear functions with integer coefficients from $[0, 1]$ to $[0, 1]$ and with operations defined as follows: for all $f, g \in A$ and for all $x \in [0, 1]$,

$$(f \oplus g)(x) = \min\{1, f(x) + g(x)\} \text{ and } \sim f(x) = 1 - f(x).$$

Consider the function $f : [0, 1] \rightarrow [0, 1]$ defined by the stipulation $f(x) = 0$ for all $x \in [0, 1/3) \cup (2/3, 1]$, while $f(x) = \min\{3x - 1, -3x + 2\}$ on $[1/3, 2/3]$. A direct computation shows that



*	0	a	b	c	d	e	f	1	$\sim x$	$\circ x$
0	0	0	0	0	0	0	0	0	1	1
a	0	0	0	0	0	a	0	a	f	b
b	0	0	0	0	0	0	b	b	e	a
c	0	0	0	0	0	a	e	c	d	0
d	0	0	0	0	d	e	d	d	c	0
e	0	a	0	a	e	e	d	e	b	a
f	0	0	b	e	d	d	a	f	a	b
1	0	a	b	c	d	e	f	1	1	1

FIGURE 2 A finite lattice of eight elements together with three operations $*$, \sim and \circ .

$\sim f(x) = 1$ for all $x \in [0, 1/3) \cup (2/3, 1]$ and $f(x) = \max\{3x - 1, -3x + 2\}$ on $[1/3, 2/3]$ so that $f \leq \sim f$ and hence $f \wedge \sim f = f$. Therefore the set $\{g \in A \mid f \wedge \sim f \wedge g = 0\} = \{g \in A \mid f \wedge g = 0\}$. However, notice that $\sup\{g \in A \mid f \wedge g = 0\}$ is a function $h : [0, 1] \rightarrow [0, 1]$ such that $h(x) = 0$ for all $x \in [0, 1/3) \cup (2/3, 1]$ and $h(x) = 1$ for all $x \in [1/3, 2/3]$ and hence it is not continuous and hence it does not belong to A . As a consequence $\circ(f)$ is not definable.

Taking into account the observation above on filters in quasivarieties, a Q -filter F of a dIRL $_c$ algebra (\mathbf{A}, \circ) , besides being implicative, has to additionally satisfy the following two conditions:

- (F1) if $x \rightarrow y, y \rightarrow x \in F$ then $\circ x \rightarrow \circ y, \circ y \rightarrow \circ x \in F$,
- (F2) if $x \vee \sim x \vee \sim y \in F$ then $y \rightarrow \circ x \in F$.

We shall call such a filter a \circ -filter. Note that from (F1) it follows in particular that \circ -filters are closed by \circ : if $x \in F$ then $\circ x \in F$ as well. Since in every dIRL $_c$ algebra (\mathbf{A}, \circ) , \circ -filters bijectively correspond to Q -congruences by the maps

$$F \mapsto \Theta_F = \{(a, b) \in A \times A \mid (a \rightarrow b) \wedge (b \rightarrow a) \in F\}$$

and

$$\Theta \mapsto F_\Theta = \{a \in A \mid a\Theta 1\},$$

we will henceforth say that a dIRL_c algebra (\mathbf{A}, \circ) is *simple* if it only has two \mathcal{Q} -congruences, and hence if it only has two (trivial) \circ -filters: $\{1\}$ and A . Furthermore, a quasivariety of dIRL_c algebras will be said to be *semisimple* provided that all its subdirectly irreducible elements are simple in the above sense.

LEMMA 3.7

Let \mathbf{A} be a subdirectly irreducible dIRL and define $\circ : A \rightarrow A$ as $\circ(1) = \circ(0) = 1$ and $\circ(x) = 0$ otherwise. Then (\mathbf{A}, \circ) is a simple dIRL_c algebra.

PROOF. Let us start by showing that \circ defined as in the statement is a \max -consistency operator. First of all it is immediate to see that $(\circ 1)$ holds, then let us hence show $(\circ 2)$. To this end assume that $x \wedge \sim x \wedge y = 0$, or equivalently that $\sim(x \wedge \sim x \wedge y) = \sim x \vee x \vee \sim y = 1$. Since \mathbf{A} is subdirectly irreducible, by [41, Theorem 4.1] one has that $(\sim x \vee x) \vee \sim y = 1$ if either $x \vee \sim x = 1$ or $\sim y = 1$. If the former is the case, then x is Boolean. Since \mathbf{A} is subdirectly irreducible, it is hence directly indecomposable. Therefore, [33, Proposition 1.5] implies $x \in \{0, 1\}$. Then, by definition $\circ(x) = 1$, whence necessarily $\circ(x) \geq y$. Conversely, if $\sim y = 1$, then $y = 0 \leq \circ(x)$.

Finally assume, without loss of generality, that A contains at least an element x distinct from 0 and 1, for otherwise \mathbf{A} would be the two element Boolean algebra and (\mathbf{A}, \circ) would be obviously simple. Furthermore, let $F \neq \{1\}$ be a \circ -filter of (\mathbf{A}, \circ) . Then F must contain x and since F is closed under \circ , by definition of \circ , $\circ(x) = 0 \in F$. Thus, $F = A$ and (\mathbf{A}, \circ) is simple. \square

The next result provides a generalization to the case of dIRL_c algebras of the well-known result showing that a residuated lattice \mathbf{A} is directly indecomposable iff $B(\mathbf{A}) = \{0, 1\}$, see [33, Proposition 1.5].

THEOREM 3.8

Let (\mathbf{A}, \circ) be a dIRL_c algebra. Then (\mathbf{A}, \circ) is directly indecomposable iff $B(\mathbf{A}) = \{0, 1\}$.

PROOF. The right-to-left direction is clear. Indeed, if (\mathbf{A}, \circ) were product of two (or more) dIRL_c algebras, say (\mathbf{A}_1, \circ) and (\mathbf{A}_2, \circ) , denoting by $1_1, 0_1$ the top and the bottom elements of the first and by $1_2, 0_2$ the top and the bottom elements of the second, then $(1_1, 1_2)$, $(0_1, 0_2)$, $(1_1, 0_2)$ and $(0_1, 1_2)$ would be four distinct Boolean elements of \mathbf{A} contradicting the fact that $B(\mathbf{A}) = \{0, 1\}$.

Hence, let us prove the left-to-right direction. First of all observe that if $z \in B(\mathbf{A})$, for all $x \in A$ $x \wedge z = x * z$. The proof is again by reduction ad absurdum. Assume there is $z \in B(\mathbf{A}) \setminus \{0, 1\}$, and let us see that $[z]$ is a \circ -filter. It is clear that it is an implicative filter. Let us check that $[z]$ satisfies the conditions of a \circ -filter:

- First we prove that $z \leq (x \rightarrow y), (y \rightarrow x)$ implies $z \leq (\circ x \rightarrow \circ y), (\circ y \rightarrow \circ x)$. Since $z \leq (x \rightarrow y)$ we have $x \wedge z = x * z \leq y$ since z is Boolean. Hence $\sim y \leq \sim(x \wedge z) = \sim x \vee \sim z$. Also, since $z \leq y \rightarrow x$, we have $y \leq z \rightarrow x$. Then we have the following inequalities:

$$y \wedge \sim y \wedge (z \wedge \circ x) \leq y \wedge (\sim x \vee \sim z) \wedge z \wedge \circ x = (y \wedge \sim x \wedge z \wedge \circ x) \vee (y \wedge \sim z \wedge z \wedge \circ x) \leq ((z \rightarrow x) \wedge \sim x \wedge z \wedge \circ x) \vee 0 = ((z * (z \rightarrow x)) \wedge \sim x \wedge \circ x) \leq x \wedge \sim x \wedge \circ x = 0.$$
Therefore, we have $z \wedge \circ x \leq \circ y$, i.e. $z \leq \circ x \rightarrow \circ y$. Analogously, one can prove $z \leq \circ y \rightarrow \circ x$.
- Second, we prove that $z \leq x \vee \sim x \vee \sim y$ implies $z \leq y \rightarrow \circ x$. Note that $x \vee \sim x \vee \sim y = \sim(x \wedge \sim x \wedge y)$. Then we have:

$$x \wedge \sim x \wedge (z \wedge y) = (x \wedge \sim x \wedge y) \wedge z = (x \wedge \sim x \wedge y) * z \leq (x \wedge \sim x \wedge y) * \sim(x \wedge \sim x \wedge y) = 0.$$
Therefore, $z * y = z \wedge y \leq \circ x$, hence, $z \leq y \rightarrow \circ x$.

Analogously, we can prove that $[\sim z]$ is a \circ -filter. Then if $B(\mathbf{A}) \setminus \{0, 1\}$ is non-empty, there exists $z \in B(\mathbf{A}) \setminus \{0, 1\}$ and the \circ -filters $[z]$ and $[\sim z]$ are such that $[z] \cap [\sim z] = \{1\}$, while the filter generated by $[z] \cup [\sim z]$ contains z and contains $\sim z$ and hence it contains $z \wedge \sim z = 0$. Thus it coincides with the

entire A . Clearly, their associated Q -congruences, say Θ_z and $\Theta_{\sim z}$, permute $\Theta_{\sim z}(\Theta_z) = \Theta_z(\Theta_{\sim z})$. Indeed $(a, b) \in \Theta_{\sim z}(\Theta_z)$ iff there exists $c \in A$ such that $(a \rightarrow c) \wedge (c \rightarrow a) \geq z$ and $(c \rightarrow b) \wedge (b \rightarrow c) \geq \sim z$ iff, by the commutativity of \wedge and the reflexive property of congruences, $(b, a) = (a, b) \in \Theta_z(\Theta_{\sim z})$. Thus, they form a nontrivial pair of complementary factor congruences and (\mathbf{A}, \circ) is not directly indecomposable. \square

Thus, since every subdirectly irreducible algebra is also directly indecomposable, we also have the following result.

COROLLARY 3.9

Let (\mathbf{A}, \circ) be a subdirectly irreducible dIRL $_c$ algebra. Then $B(\mathbf{A}) = \{0, 1\}$.

When the algebra \mathbf{A} is finite or is a *connected or disconnected rotation* of a residuated lattice, we can prove more. Connected and disconnected rotations of residuated lattices were studied by Jenei in [31]. The paper [10] (see also [50]) studies varieties of algebras obtained as connected or disconnected rotations of residuated lattices from a purely algebraic perspective. There, the authors introduce a variety of algebras, denoted by IMVR3, whose directly indecomposable elements are obtained as *generalized 3-rotations* (in the terminology of [10]) of residuated lattices. Within directly indecomposable IMVR3 algebras, we can hence identify structures with a negation fixpoint (corresponding to Jenei’s connected rotations) and structures without a negation fixpoint (corresponding to Jenei’s disconnected rotations). As for distributivity, in [10, Remark 3.9], it is observed that any IMVR3 algebra obtained as a generalized 3-rotation of a residuated lattice \mathbf{R} is distributive iff so is \mathbf{R} . We will henceforth say that an algebra is a *dIMVR3 algebra* if it is a distributive IMVR3 algebra and the corresponding variety will be denoted by dIMVR3. Thus, every algebra of dIMVR3 is a commutative, integral, bounded, involutive and distributive residuated lattice, i.e. dIMVR3 as a subvariety of dIRL.

By [10, Theorem 4.6], the domain of every directly indecomposable IMVR3 algebra \mathbf{A} can be partitioned in two sets,

$$A^+ = \{x : x \geq \sim x\} \text{ and } A^- = \{x : x \leq \sim x\}.$$

Furthermore, every directly indecomposable IMVR3 algebra satisfies the following conditions: if $x \in A^+$ and $y \in A^-$, then (1) $x \geq y$ and (2) $y^2 = y * y = 0$.

In what follows, dIRL $_c$ algebras (\mathbf{A}, \circ) where $\mathbf{A} \in \text{dIMVR3}$ will be called a dIMVR3 $_c$ algebra.

The next theorem characterizes, in particular, those subdirectly irreducible dIRL $_c$ algebras (\mathbf{A}, \circ) in which \mathbf{A} is either a finite or a dIMVR3 $_c$ algebra. It is worth pointing out that the following characterization does not hold in general for infinite structures. In subsection 4.2, we will give an example of an infinite dIRL $_c$ algebra, which is subdirectly irreducible but not simple.

THEOREM 3.10

Let (\mathbf{A}, \circ) be a finite dIRL $_c$ algebra or an arbitrary dIMVR3 $_c$ algebra. Then the following conditions are equivalent:

- (i) $B(\mathbf{A}) = \{0, 1\}$,
- (ii) (\mathbf{A}, \circ) is a directly indecomposable dIRL $_c$ algebra,
- (iii) (\mathbf{A}, \circ) is a subdirectly irreducible dIRL $_c$ algebra,
- (iv) (\mathbf{A}, \circ) is a simple dIRL $_c$ algebra.

As a consequence, the quasivariety of dIMVR3 $_c$ algebras is semisimple.

PROOF. Due to Theorem 3.8 and Corollary 3.9, we are left to prove only that (iii) implies (iv), since (iv) implies (iii) in general.

In the case of \mathbf{A} being finite, towards a contradiction, assume (\mathbf{A}, \circ) is not simple. Hence, there is a \circ -filter F such that $F \neq A$ and $F \neq \{1\}$. Since A is finite, there is $0 < a \in A \setminus \{1\}$ such that $F = [a]$. Since F is a \circ -filter, $\circ(a) \in F$ and thus it must be $a \leq \circ(a)$. But then we have $0 = a \wedge \sim a \wedge \circ(a) = a \wedge \sim a$, i.e. a is Boolean, and hence $B(\mathbf{A}) \neq \{0, 1\}$ and hence (\mathbf{A}, \circ) is not subdirectly irreducible by Corollary 3.9.

Assume now (\mathbf{A}, \circ) is a subdirectly irreducible, and hence a directly indecomposable, dIMVR3_c algebra. We have to show that (\mathbf{A}, \circ) is simple. By Theorem 3.8, (\mathbf{A}, \circ) is directly indecomposable iff $B(\mathbf{A}) = \{0, 1\}$. Therefore, by [33, Proposition 1.5], \mathbf{A} is directly indecomposable as a dIMVR3 algebra. Moreover, [10, Theorems 4.6] shows that if a dIMVR3 algebra \mathbf{A} is directly indecomposable, its domain, as mentioned above, can be expressed as $A = A^+ \cup A^-$, where $A^+ = \{x : x \geq \sim x\}$ and $A^- = \{x : x \leq \sim x\}$, and satisfying $x \geq y$ whenever $x \in A^+$ and $y \in A^-$, and $y^2 = 0$ for every $y \in A^-$. Now, if $x \in A^+$ then $\sim x \in A^-$ and thus, $0 = x \wedge \sim x \wedge \circ(x) = \sim x \wedge \circ(x)$, from which it follows that $\circ(x) \in A^-$ as well, and hence $(\circ(x))^2 = 0$. Since any \circ -filter F of (\mathbf{A}, \circ) containing an element $x \neq 1$ has to contain $\circ(x)$ as well, it has to contain $(\circ(x))^2$ as well, i.e. F must be such that $0 \in F$. Therefore, F must be the whole algebra domain A . Thus, we have shown that (\mathbf{A}, \circ) is a simple dIRL_c algebra. \square

In the light of the Theorem 3.10 above, it is clear that the lattice of Figure 2 together with the operations of the table of Figure 2 is an example of a finite dIRL_c algebra whose Boolean elements are 0 and 1, and hence it is directly indecomposable, subdirectly irreducible and simple.

3.2 *Distributive involutive residuated lattices with a maxB-consistency operator*

We now start considering consistency operators that map the elements of a dIRL algebra into Boolean elements of the same. The next definition introduces dIRL s with a maxB -consistency operator.

DEFINITION 3.11

A *dIRL with a maxB-consistency operator* (or $\text{dIRL}_c^{\text{mB}}$ algebra for short) is a pair (\mathbf{A}, \circ) where \mathbf{A} is a dIRL algebra and $\circ : A \rightarrow A$ satisfies the following conditions: for all $x, y, z \in A$,

- (o1) $x \wedge \sim x \wedge \circ(x) = 0$
- (o3) if $x \wedge \sim x \wedge y = 0$ and $y \wedge \sim y = 0$, then $y \leq \circ(x)$
- (o4) $\circ(x) \wedge \sim \circ(x) = 0$

Again, from this definition, it follows that the class of $\text{dIRL}_c^{\text{mB}}$ algebras is a quasivariety. Moreover, and similarly to the case of max -consistency operators, also now the conditions (o1), (o3) and (o4) capture the expected behavior of maxB -consistency operators as described at the beginning of this Section 3.

LEMMA 3.12

Let \mathbf{A} be a dIRL , and let \circ be unary operation on A . Then (\mathbf{A}, \circ) is an $\text{dIRL}_c^{\text{mB}}$ algebra iff, for any $x \in A$,

$$\circ(x) = \max\{z \in B(\mathbf{A}) \mid x \wedge \sim x \wedge z = 0\}.$$

PROOF. Clearly, for a given $x \in A$, the set $\{z \in B(\mathbf{A}) \mid x \wedge \sim x \wedge z = 0\}$ is closed by \vee , it has $\circ(x)$ as an upper bound by (o3) and moreover $\circ(x)$ belongs to that set by (o1) and (o4). \square

From the definition it is easy to prove the following properties of maxB -consistency operators.

TABLE 1 Two operators for the Nelson lattice of Figure 2.

x	$\diamond x$	$\circ x$
0	1	1
a	0	b
b	0	a
c	0	0
d	0	0
e	0	a
f	0	b
1	1	1

PROPOSITION 3.13

The following properties hold in a $\text{dIRL}_c^{\text{mB}}$ algebra (\mathbf{A}, \circ) :

- (i) $\circ(x) = \max\{z \in B(\mathbf{A}) \mid z \leq x \vee \sim x\}$
- (ii) $\circ(x) = \circ(\sim x) = \circ(x \wedge \sim x) = \circ(x \vee \sim x)$
- (iii) $\circ(x) = 1$ iff $x \in B(\mathbf{A})$, in particular $\circ(1) = \circ(0) = 1$
- (iv) $\circ \circ(x) = 1$.

Notice that if (\mathbf{A}, \circ) is a dIRL_c algebra and (\mathbf{A}, \diamond) is $\text{dIRL}_c^{\text{mB}}$ algebra, then from Lemmas 3.4 and 3.12 it is clear that $\diamond(x) \leq \circ(x)$ for all $x \in A$.

In the nilpotent minimum algebra of Figure 2, the corresponding mB-consistency operator \diamond comes defined as $\diamond(0) = \diamond(1) = 1$ and $\diamond(x) = 0$ otherwise. Since this is a different operation from the operation \circ in the table of Figure 2, it readily follows that the classes of dIRL_c algebras and $\text{dIRL}_c^{\text{mB}}$ algebras are different. Indeed, consider again the two unary operations in Table 1. It turns out that (\mathbf{L}, \circ) is a dIRL_c algebra but not a $\text{dIRL}_c^{\text{mB}}$ algebra, while (\mathbf{L}, \diamond) is a $\text{dIRL}_c^{\text{mB}}$ algebra but not a dIRL_c algebra.

REMARK 3.14

Like in the case of a max-consistency operator over a dIRL algebra, a maxB-consistency operator, if it exists, is unique. Actually, according to Lemma 3.12, the maxB-consistency operator \circ is definable on a dIRL if and only if all the elements of the form $x \wedge \sim x$ have a minimum Boolean element above them. Since the value of $\circ(x)$ is a maximum of Boolean elements, \circ is always definable in any dIRL \mathbf{A} such that $B(\mathbf{A})$ is finite. However, the definability of the \circ operator is not guaranteed when $B(\mathbf{A})$ is infinite.

In the quasivariety of $\text{dIRL}_c^{\text{mB}}$ algebras, the corresponding notion of Q -filter is that of an implicative filter F further satisfying the following two conditions:

- (F1) if $x \rightarrow y, y \rightarrow x \in F$ then $\circ x \rightarrow \circ y, \circ y \rightarrow \circ x \in F$
- (F3) if $x \vee \sim x \vee \sim y \in F$ and $y \vee \sim y \in F$ then $y \rightarrow \circ x \in F$.

We will call them \circ_b -filters.

From this we can provide a characterization of subdirectly irreducible dIRL_c algebras. To begin with, we can observe that the same proof of Lemma 3.7 also applies to the case of $\text{dIRL}_c^{\text{mB}}$

algebras with a unique atom and hence to all finite and subdirectly irreducible dIRL algebras with a mB -consistency operator as well.

LEMMA 3.15

Let \mathbf{A} be a subdirectly irreducible dIRL with a unique atom and define $\circ : A \rightarrow A$ as $\circ(1) = \circ(0) = 1$ and $\circ(x) = 0$ otherwise. Then (\mathbf{A}, \circ) is a simple $\text{dIRL}_c^{\text{mB}}$ algebra.

Moving from max - to maxB -consistency operators allows us to improve the results shown in Theorem 3.8 and Theorem 3.10 and, as anticipated, to characterize all subdirectly irreducible $\text{dIRL}_c^{\text{mB}}$ algebras.

THEOREM 3.16

Let (\mathbf{A}, \circ) be a $\text{dIRL}_c^{\text{mB}}$ algebra, then the following conditions are equivalent:

- (i) $B(\mathbf{A}) = \{0, 1\}$
- (ii) (\mathbf{A}, \circ) is a directly indecomposable $\text{dIRL}_c^{\text{mB}}$ algebra
- (iii) (\mathbf{A}, \circ) is a subdirectly irreducible $\text{dIRL}_c^{\text{mB}}$ algebra
- (iv) (\mathbf{A}, \circ) is a simple $\text{dIRL}_c^{\text{mB}}$ algebra.

PROOF. That (iv) implies (iii) and (iii) implies (ii) is clear. It is hence left to show that (ii) implies (i) and (i) implies (iv).

(ii) implies (i). The same proof of Theorem 3.8 applies.

(i) implies (iv). If $B(\mathbf{A}) = \{0, 1\}$ then a simple computation shows that $\circ(0) = \circ(1) = 1$ and $\circ(x) = 0$ otherwise since for all $x \in A \setminus \{0, 1\}$, $x \wedge \sim x \neq 0$. Then every \circ_b -filter F containing an element $x \notin \{0, 1\}$ is equal to A and thus A is simple. \square

As a direct consequence of this result we have the following corollary.

COROLLARY 3.17

The quasi-variety of $\text{dIRL}_c^{\text{mB}}$ algebras is semisimple.

3.3 *Distributive involutive residuated lattices with a Bmax-consistency operator*

Finally, in this subsection, we consider dIRLs expanded with a Bmax -consistency operator.

DEFINITION 3.18

A dIRL with a Bmax -consistency operator (or $\text{dIRL}_c^{\text{Bm}}$ algebra for short) is a dIRL_c algebra (\mathbf{A}, \circ) satisfying the additional condition: for all $x \in A$,

$$(\circ 4) \quad \circ(x) \vee \sim \circ(x) = 1.$$

From the very definition, it is clear that the class of $\text{dIRL}_c^{\text{Bm}}$ algebras constitutes a subquasivariety of the quasivariety of dIRL_c algebras.

We also have a characterization of Bmax -consistency operators similar to the case of max -consistency operators with the obvious modification.

LEMMA 3.19

Let \mathbf{A} a dIRL, and let \circ be a unary operation on A . Then (\mathbf{A}, \circ) is a $\text{dIRL}_c^{\text{Bm}}$ algebra iff, for any $x \in A$,

$$\circ(x) = \max\{z \in A \mid x \wedge \sim x \wedge z = 0\} \text{ and } \circ(x) \wedge \sim \circ(x) = 0.$$

Actually, as anticipated, Bmax-consistency operators are just max-consistency operators that are maxB-consistency operators as well.

LEMMA 3.20

The quasivariety of $\text{dIRL}_c^{\text{Bm}}$ algebras is the intersection of the quasivariety of dIRL_c algebras and the quasivariety of $\text{dIRL}_c^{\text{mB}}$ algebras.

PROOF. Let (\mathbf{A}, \circ) be a $\text{dIRL}_c^{\text{Bm}}$ algebra. We have to show that \circ is a maxB-consistency operator as well. Since $\circ(x)$ is Boolean, $\circ(x) \in \{z \in B(\mathbf{A}) \mid x \wedge \sim x \wedge z = 0\}$, i.e. we also have $\circ(x) = \max\{z \in B(\mathbf{A}) \mid x \wedge \sim x \wedge z = 0\}$. Conversely, if \circ is both a max- and maxB-consistency operator, then it is clear that $\circ(x) = \max\{z \in A \mid x \wedge \sim x \wedge z = 0\}$ and that $\circ(x)$ is Boolean, i.e. \circ is a Bmax-consistency operator. \square

The following proposition collects basic properties of $\text{dIRL}_c^{\text{Bm}}$ algebras.

PROPOSITION 3.21

For a given $\text{dIRL}_c^{\text{Bm}}$ algebra (\mathbf{A}, \circ) and $x \in A$, we have

- (i) $\circ\circ(x) = 1$.
- (ii) $\circ(x) = 1$ iff $x \wedge \sim x = 0$ iff x is a Boolean element.

Again, an analogous result to Lemma 3.7 and Lemma 3.15 for dIRL_c algebras and $\text{dIRL}_c^{\text{mB}}$ algebras respectively also holds for $\text{dIRL}_c^{\text{Bm}}$.

LEMMA 3.22

Let \mathbf{A} be a subdirectly irreducible dIRL algebra with a unique atom and let \circ the unary operation on A defined as $\circ(1) = \circ(0) = 1$ and $\circ(x) = 0$ otherwise. Then (\mathbf{A}, \circ) is a simple $\text{dIRL}_c^{\text{Bm}}$ algebra.

Since $\text{dIRL}_c^{\text{Bm}}$ algebras are dIRL_c algebras fulfilling the additional equation ($\circ 4$), the corresponding notions of Q-filters and Q-congruences are the same, and we can obtain similar results as those obtained for dIRL_c algebras and $\text{dIRL}_c^{\text{mB}}$ algebras. In particular, the following result characterizes subdirectly irreducible $\text{dIRL}_c^{\text{Bm}}$ algebras.

THEOREM 3.23

Let (\mathbf{A}, \circ) be a $\text{dIRL}_c^{\text{Bm}}$ algebra, then the following conditions are equivalent:

- (i) $B(\mathbf{A}) = \{0, 1\}$
- (ii) (\mathbf{A}, \circ) is a directly indecomposable $\text{dIRL}_c^{\text{Bm}}$ algebra
- (iii) (\mathbf{A}, \circ) is a subdirectly irreducible $\text{dIRL}_c^{\text{Bm}}$ algebra
- (iv) (\mathbf{A}, \circ) is a simple $\text{dIRL}_c^{\text{Bm}}$ algebra

PROOF. The same proof of Theorem 3.16 applies. \square

3.4 The prelinear case: IMTL algebras

We now turn the attention to the particular case of prelinear dIRL algebras with a consistency operator \circ . As shown in [23], involutive residuated lattices satisfying the prelinearity equation

$$(x \rightarrow y) \vee (y \rightarrow x) = 1$$

are precisely the so-called *involutive MTL algebras* (or *IMTL algebras* for short). Let us start showing a first result concerning IMTL algebras expanded by a Bmax-consistency operator.

LEMMA 3.24

Let \mathbf{A} be a IMTL algebra that is not a chain. Then there are two elements $x, y \neq 1$ such that $x \vee y = 1$.

PROOF. If \mathbf{A} is not a chain, let a, b two incomparable elements. Then $x = a \rightarrow b$ and $y = b \rightarrow a$ must be different from 1, and by prelinearity, $x \vee y = (a \rightarrow b) \vee (b \rightarrow a) = 1$. \square

Let us call $\text{IMTL}_c^{\text{Bm}}$ algebras those $\text{dIRL}_c^{\text{Bm}}$ algebras (\mathbf{A}, \circ) such that \mathbf{A} is a IMTL algebra.

PROPOSITION 3.25

Let (\mathbf{A}, \circ) be a $\text{IMTL}_c^{\text{Bm}}$ algebra. Then the following statements are equivalent:

- (i) $B(\mathbf{A}) = \{0, 1\}$
- (ii) \mathbf{A} is an IMTL chain
- (iii) (\mathbf{A}, \circ) is a directly indecomposable $\text{IMTL}_c^{\text{Bm}}$ algebra
- (iv) (\mathbf{A}, \circ) is a subdirectly irreducible $\text{IMTL}_c^{\text{Bm}}$ algebra
- (v) (\mathbf{A}, \circ) is a simple $\text{IMTL}_c^{\text{Bm}}$ algebra

PROOF. Thanks to Theorem 3.23 it is sufficient to prove the equivalence between (i) and (ii).

(i) implies (ii). By Theorem 3.23, $B(\mathbf{A}) = \{0, 1\}$. By Lemma 3.24 if \mathbf{A} is not a chain there exist elements $x, y \neq 1$ such that $x \vee y = 1$. Moreover, $x \wedge \sim x \neq 0$, since if $x \wedge \sim x = 0$ then x would be a Boolean element, a contradiction with the assumption that $x \neq 1, 0$. Moreover, $x \wedge \sim x \wedge \sim y \leq \sim x \wedge \sim y = \sim(x \vee y) = 0$, and thus $\circ(x) \geq \sim y$. Since $B(\mathbf{A}) = \{0, 1\}$ and $\sim y \neq 0$, it follows that $\circ(x) = 1$, which contradicts the definition of \circ .

In order to prove that (ii) implies (i), let us show that (ii) implies (v) whence the result will follow from Theorem 3.23. If \mathbf{A} is an IMTL chain, then (v) obviously follows because, over a chain, \circ is defined as $\circ(0) = \circ(1) = 1$ and $\circ(x) = 0$ otherwise, and thus, a \circ -filter F that contains an element $x \neq 1, 0$ must contain $\circ(x) = 0$, hence $F = A$, whence (\mathbf{A}, \circ) is a simple. \square

Let us remark that Proposition 3.25 above does not apply to IMTL algebras expanded by \max and $\max\mathbf{B}$ -consistency operators. Indeed, consider the IMTL algebra \mathbf{A} of Figure 2 expanded by either its (unique) \max or $\max\mathbf{B}$ -consistency operator \circ . By Theorem 3.10 and Theorem 3.16, (\mathbf{A}, \circ) is directly indecomposable, subdirectly irreducible, simple and $B(\mathbf{A}) = \{0, 1\}$; however, its lattice reduct is not totally ordered. Moreover, it is clear that the same finite algebra \mathbf{A} does not admit a $\mathbf{B}\max$ -consistency operator, while on the other hand, it is always definable on a chain.

It is worth pointing out that Theorem 3.10 applies to this peculiar case and, in particular, to those IMTL_c algebra whose underlying IMTL algebra is either finite or is obtained as rotation of a prelinear semihoop. The latter class of structures has been largely studied in [37].

Now, we turn our attention to the variety of IMTL_* algebras that was defined in [19] and let us analyse analogies and differences between them and IMTL algebras with a consistency operator \circ . Indeed, in [19], the authors consider expansions of IMTL algebras with a consistency operator that, to avoid a notational clash, we will denote here by $*$. The equations for $*$ are stronger than the ones we introduced in this paper and in fact they allow us to prove that the variety of IMTL_* is *semilinear*, that is to say, it is generated by its totally ordered members.

More precisely, a pair $(\mathbf{A}, *)$ is an IMTL_* algebra if \mathbf{A} is an IMTL algebra and the following equations for $*$ hold:

- (B1) $x \wedge \neg x \wedge *(x) = 0$
- (B2) $*(x \leftrightarrow y) \leq (*x \leftrightarrow *y)$
- (B3) $*(x \vee y) \leq *(x) \vee *y$
- (B4) $*(0) = *(1) = 1$

The variety of IMTL_* algebras is shown in [19] to be semisimple, and the simple IMTL_* algebras are those defined over IMTL chains, i.e. all IMTL_* algebras are subdirect products of chains, and chains are simple. As also shown in [19], if \mathbf{A} is a IMTL chain, the unique $*$ operator that makes $(\mathbf{A}, *)$ an IMTL_* algebra is the one defined as $*(0) = *(1) = 1$ and $*(x) = 0$ otherwise. Hence IMTL_* , dIRL_c , $\text{dIRL}_c^{\text{mB}}$ and $\text{dIRL}_c^{\text{Bm}}$ algebras over IMTL chains share the same consistency operators.

The following is a consequence of Proposition 3.25.

COROLLARY 3.26

The class of $\text{IMTL}_c^{\text{Bm}}$ algebras is a variety that coincides with the variety of IMTL_* algebras. Thus, both are semilinear and semisimple.

PROOF. Let us start showing that the quasivariety of $\text{IMTL}_c^{\text{Bm}}$ algebras is indeed a variety. To this end notice that by Proposition 3.25, the class of $\text{IMTL}_c^{\text{Bm}}$ algebras is generated by its totally ordered members. Moreover, following [19], for every $\text{IMTL}_c^{\text{Bm}}$ algebra (\mathbf{A}, \circ) , the unary operation $\Delta : A \rightarrow A$ defined by $\Delta(x) = x \wedge \circ(x)$ coincides with the Baaz–Monteiro projection operator [4, 36]. These facts show that the algebraic logic of $\text{IMTL}_c^{\text{Bm}}$ algebras is a Δ -core fuzzy logic in the sense of [18] and hence, for every finite set of formulas $\{\psi_1, \dots, \psi_k, \phi\}$, it holds $\{\psi_1, \dots, \psi_k\} \vdash \phi$ iff $\vdash (\Delta\psi_1 \wedge \dots \wedge \Delta\psi_k) \rightarrow \phi$ (see [29, Theorem 2.4.14]). Algebraically, this means that every quasi-equation describing $\text{IMTL}_c^{\text{Bm}}$ algebras can be written equationally, whence it is indeed a variety.

Finally, the claim follows from Proposition 3.25 above. Indeed, both the variety of $\text{IMTL}_c^{\text{Bm}}$ algebras and the variety of IMTL_* algebras are generated by its totally ordered members. We already observed that, on IMTL chains, the unique Bmax -consistency operator \circ coincides with the $*$ -operator: it maps all non-Boolean elements to 0, and 0 and 1 to 1. Thus the two varieties are generated by the same structures and hence they coincide. \square

In fact the previous result can be proved in a slightly more general setting. To this end consider the *semilinear extension* of the quasi-varieties of dIRL_c , $\text{dIRL}_c^{\text{mB}}$ and $\text{dIRL}_c^{\text{Bm}}$ algebras. That is to say the quasi-varieties obtained from the above ones by replacing $(\circ 2)$ in Definition 3.3 by the quasi-equation

$$(x \wedge \sim x \wedge y) \vee z = 0 \text{ implies } (y \rightarrow \circ(x)) \vee z = 1$$

and $(\circ 3)$ in Definition 3.11 by

$$(x \wedge \sim x \wedge y) \vee z = 0 \text{ and } (y \wedge \sim y) \vee z = 0 \text{ implies } (y \rightarrow \circ(x)) \vee z = 1.$$

By general results it follows that every algebra in the semilinear extension of the quasi-varieties of dIRL_c , $\text{dIRL}_c^{\text{mB}}$ and $\text{dIRL}_c^{\text{Bm}}$ algebras can be represented as a subdirect product of totally ordered structures in the same classes [16]. Thus, the following corollary easily holds.

COROLLARY 3.27

The semilinear extensions of dIRL_c , $\text{dIRL}_c^{\text{mB}}$ and $\text{dIRL}_c^{\text{Bm}}$ algebras form varieties and they all coincide with the variety of IMTL_* algebras.

PROOF. Over a dIRL -chain, max , Bmax and maxB consistency operators coincide and therefore the sub-quasivariety generated by them are the same and coincide with the variety of IMTL_* algebras. \square

Nevertheless, there are IMTL_c algebras that are not subdirect product of chains as the one in Figure 2 that is simple, hence subdirectly irreducible, but it is not totally ordered. Therefore, the quasi-varieties of IMTL_c and $\text{IMTL}_c^{\text{mb}}$ algebras are not semilinear.




FIGURE 3 Diagram of main classes of algebras with a consistency operator we consider in this paper and their relationships as expansions of dIRL.

Of course, the quasivariety of $\text{IMTL}_c^{\text{mB}}$ algebras is semisimple, but it is an open problem whether the quasivariety of IMTL_c algebras is semisimple or whether there exist subdirectly irreducible algebras that are not simple, as in the general case of dIRL or NL algebras.

As a sort of summary, Figure 3 provides a graphical representation of the main quasivarieties of algebras expanded with a consistency operator that we have considered in this section, where dashed arrows stand for expansions and solid arrows denote extensions.

4 The particular case of Nelson lattices with a max-consistency operator

In this section we will focus on expansions of Nelson lattices by means of a max-consistency operator. In particular we will take advantage of well-developed structural properties for these algebras in terms of *twist-products* to both investigate more in details subdirectly irreducible structures and also to show that Theorem 3.10 does not hold in the general setting of infinite structures.⁶

Nelson lattices can be represented by means of Heyting algebras and their *Boolean filters*, that is to say, filters F of a Heyting algebra \mathbf{H} such that the quotient \mathbf{H}/F is a Boolean algebra. This relation has been investigated by several authors and main contributions in this sense have been provided by Fidel [24], Vakarelov [51] and Sendlewski [46].

In the following two subsections, after some needed brief preliminaries, we are going to prove similar representations for Nelson lattices expanded by consistency operators (Subsection 4.1) and,

⁶Twist-products have been developed also for, in general unbounded, involutive residuated lattices (see [9]). However, these results, although being more general, do not lead to a complete representation of all (distributive) involutive residuated lattices. Thus, here we preferred to restrict to Nelson lattices since in this setting we can obtain stronger results.

as anticipated in Subsection 3.1, we will also show more results on subdirectly irreducible algebras (Subsection 4.2).

It is worth to notice that prelinear Nelson lattices precisely correspond to NM algebras and moreover the twist-product preserves prelinearity. In other words, each NM algebra \mathbf{A} can be uniquely represented by a prelinear Heyting algebra \mathbf{G} (aka a Gödel algebra) and a Boolean filter of \mathbf{G} [7, Theorem 6.19]. As a consequence, all the results of this section (with the clear exception of the counterexample we show in Subsection 4.2) apply to NM algebras with a consistency operator.

Conforming to the previous notation, we will call NL_c , NL_c^{mB} and NL_c^{Bm} algebras to the Nelson lattices expanded, respectively, with \max -, $\max B$ - and $B\max$ -consistency operators.

To start with, let us recall the following result, whose formulation is taken from [8].

THEOREM 4.1 ([46]).

Given a Heyting algebra $\mathbf{H} = (H, \wedge, \vee, \rightarrow_H, 0, 1)$ and a Boolean filter F of \mathbf{H} let

$$N(\mathbf{H}, F) := \{(x, y) \in H \times H : x \wedge y = 0 \text{ and } x \vee y \in F\}.$$

Then we have the following:

- (i) $\mathbf{N}(\mathbf{H}, F) = (N(\mathbf{H}, F), \vee, \wedge, *, \rightarrow, \sim, 0, 1)$ is a Nelson lattice, where operations are defined as follows:

$$\begin{aligned} (x, y) \vee (s, t) &= (x \vee s, y \wedge t), \\ (x, y) \wedge (s, t) &= (x \wedge s, y \vee t), \\ (x, y) * (s, t) &= (x \wedge s, (x \rightarrow_H t) \wedge (s \rightarrow_H y)), \\ (x, y) \rightarrow (s, t) &= ((x \rightarrow_H s) \wedge (t \rightarrow_H y), x \wedge t), \\ \sim(x, y) &= (y, x), \\ 1 &= (1, 0), \\ 0 &= (0, 1). \end{aligned}$$

- (ii) If F_1, F_2 are Boolean filters of \mathbf{H} , then $\mathbf{N}(\mathbf{H}, F_1)$ is a subalgebra of $\mathbf{N}(\mathbf{H}, F_2)$ if and only if $F_1 \subseteq F_2$.

Following the tradition (see [8] for instance), we will include, for every Heyting algebra \mathbf{H} , the improper filter H among the Boolean filters of \mathbf{H} . In what follows, the Nelson lattice $\mathbf{N}(\mathbf{H}, H)$ will be simply denoted by $\mathbf{N}(\mathbf{H})$.

The following result, independently proved by Fidel in [24] and by Vakarelov in [51], can hence be stated as a corollary of Theorem 4.1 above.

COROLLARY 4.2

For every Nelson lattice \mathbf{A} there is a Heyting algebra \mathbf{H} such that \mathbf{A} is isomorphic to a subalgebra of $\mathbf{N}(\mathbf{H})$.

In what follows we will need a further requirement for the Boolean filter of a Heyting algebra in order to extend the previous representation to Nelson lattices with consistency operators. Let us hence recall from [45] that a unary operation \ulcorner on a Heyting algebra \mathbf{H} is called a *dual pseudocomplement*, if the following equations are satisfied:

- (D1) $x \vee \ulcorner(x \vee y) = x \vee \ulcorner y$,
 (D2) $x \vee \ulcorner 1 = x$,
 (D3) $\ulcorner \ulcorner 1 = 1$.

In any Heyting algebra \mathbf{H} , if the dual pseudocomplement of $x \in H$ exists, then it is defined as

$$\neg x = \min\{z \in H \mid z \vee x = 1\}. \quad (1)$$

Recall from [45] that in every Heyting algebra with dual pseudocomplement, congruences bijectively correspond to *normal filters*, that is to say, implicative filters that satisfy the following further requirement: if $x \in F$, then $\neg\neg x \in F$ as well.

We will make use of the following easy result that is known in the literature, but we provide the proof for the sake of self-containedness.

LEMMA 4.3

A filter F of a Heyting algebra \mathbf{H} is Boolean iff every $a \in H$, $a \vee \neg a \in F$.

PROOF. Assume that F is Boolean, and hence \mathbf{H}/F is a Boolean algebra. Thus for all $[a]_F \in H/F$, $[a]_F \vee \neg[a]_F = [1]_F$, i.e. $[a \vee \neg a]_F = [1]_F$ and hence $a \vee \neg a \in F$. Conversely, if $a \vee \neg a \in F$ for all $a \in H$, the quotient \mathbf{H}/F satisfies $[1]_F = [a \vee \neg a]_F = [a]_F \vee \neg[a]_F$ for all $a \in H$. Therefore, \mathbf{H}/F is a Boolean algebra whence F is Boolean. \square

4.1 Representation of Nelson lattices with consistency operators

Thanks to the representation of Nelson lattices in terms of Heyting algebras and Boolean filters, if $\mathbf{A} = \mathbf{N}(\mathbf{H}, F)$ is a Nelson lattice and $\circ : A \rightarrow A$ is a \max -consistency operator, by virtue of Lemma 3.4, \circ can be equivalently reformulated in the following way: for all $(a, b) \in A$,

$$\circ(a, b) = \max\{(z, z') \in A \mid a \vee b \vee z' = 1\}. \quad (2)$$

Indeed, taking into account how the operations are defined in $\mathbf{N}(\mathbf{H}, F)$ and that $a \wedge b = 0$, we have the following chain of equalities:

$$\begin{aligned} \circ(a, b) &= \max\{(z, z') \in A \mid (a, b) \wedge \neg(a, b) \wedge (z, z') = (0, 1)\} \\ &= \max\{(z, z') \in A \mid (a, b) \wedge (b, a) \wedge (z, z') = (0, 1)\} \\ &= \max\{(z, z') \in A \mid (a \wedge b \wedge z, b \vee a \vee z') = (0, 1)\} \\ &= \max\{(z, z') \in A \mid (0, a \vee b \vee z') = (0, 1)\} \\ &= \max\{(z, z') \in A \mid a \vee b \vee z' = 1\}. \end{aligned}$$

We are now going to show a representation for NL_c algebras in terms of Heyting algebras, Boolean filters and the dual-pseudocomplement. First, we need to prove the following.

LEMMA 4.4

Let \mathbf{H} be a Heyting algebra and let F be a Boolean filter of \mathbf{H} . If the dual pseudo-complement of $a \vee b$, $\neg(a \vee b)$, exists for all those a, b in H such that $a \wedge b = 0$ and $a \vee b \in F$, then the \max -consistency operator $\circ(a, b)$ exists in the Nelson lattice $\mathbf{N}(\mathbf{H}, F)$ and $\circ(a, b) = (\neg d, d)$, where $d = \neg(a \vee b)$.

PROOF. Recalling the above equation (2), consider the set $D = \{(z, z') \in N(H, F) \mid a \vee b \vee z' = 1\}$. It is clear that, by definition, if $\max D$ exists, then $\circ(a, b) = \max D$. Besides, we can write $D = \{(z, z') \in H \times H \mid z \wedge z' = 0, z \vee z' \in F, a \vee b \vee z' = 1\}$.

Now, for each $z' \in H$ such that $a \vee b \vee z' = 1$, let us define $D_{z'} = \{(z, z') \mid z \in H, z \wedge z' = 0, z \vee z' \in F\}$. Hence, it is clear that

$$D = \bigcup \{D_{z'} \mid z' \in H, a \vee b \vee z' = 1\}$$

and $D_{z'} \neq \emptyset$ for each $z' \in H$ such that $a \vee b \vee z' = 1$. Moreover, it is possible to prove that $\max D_{z'}$ exists and $\max D_{z'} = (\neg z', z')$. Indeed, if $(z, z') \in D_{z'}$ then $(z, z') \leq (\neg z', z')$ because, if $(z, z') \in D_{z'}$

then $z \wedge z' = 0$ and then, $z \leq \neg z'$. Therefore, $\max D_{z'} = (\neg z', z')$. On the other hand, we can see that $\max\{\max D_{z'} \mid z' \in H, a \vee b \vee z' = 1\}$ exists and equals $(\neg d, d)$, where, recalling Equation (1), $d = \neg(a \vee b)$. Indeed, $\max\{\max D_{z'} \mid z' \in H, a \vee b \vee z' = 1\} = \max\{(\neg z', z') \mid z' \in H, a \vee b \vee z' = 1\} = (\neg \min\{z' \mid a \vee b \vee z' = 1\}, \min\{z' \mid a \vee b \vee z' = 1\}) = (\neg d, d)$, where $d = \neg(a \vee b)$.

Finally, we are going to show that $\max D$ exists and in fact $\max D = (\neg d, d)$. We have to prove the following two conditions:

(I) $(\neg d, d) \in D$ and (II) $(f, g) \leq (\neg d, d)$ for any $(f, g) \in D$.

As for condition (I), we have to check that (i) $\neg d \wedge d = 0$, which is obvious, (ii) $\neg d \vee d \in F$, which follows from Lemma 4.3 and (iii) $a \vee b \vee d = 1$, which also follows because $d = \neg(a \vee b)$. As for condition (II), assume $(f, g) \in D$. Then $f \wedge g = 0, f \vee g \in F$ and $a \vee b \vee g = 1$. From $f \wedge g = 0$ it follows that $f \leq \neg g$, and from $a \vee b \vee g = 1$ and (1) it follows that $g \geq \neg(a \vee b)$ and $f \leq \neg g \leq \neg(\neg(a \vee b))$. Therefore, $(f, g) \leq (\neg(\neg(a \vee b)), \neg(a \vee b))$. \square

Therefore, every NL_c algebra can be represented as follows.

THEOREM 4.5

Every NL_c algebra is of the form $(\mathbf{N}(\mathbf{H}, F), \circ)$ where \mathbf{H} is a Heyting algebra and F is a Boolean filter of H such that the dual pseudo-complement exists in \mathbf{H} for all the elements of F , and for all $(a, b) \in N(H, F)$, $\circ(a, b) = (\neg\neg(a \vee b), \neg(a \vee b))$.

PROOF. Let us define $G = \{a \vee b \mid a, b \in H, (a, b) \in A\} = \{a \vee b \mid a, b \in H, a \wedge b = 0, a \vee b \in F\}$ and let us prove that $G = F$. It is clear that, by definition of G , $G \subseteq F$. Now, let $y \in F$ and write it as $y = y \vee 0$. Then, $y \wedge 0 = 0$ and that $y \vee 0 = y \in F$, hence $y \in G$. Hence, $F \subseteq G$.

Therefore, by Lemma 4.4, if the dual pseudocomplement \neg exists for all $x \in F$, the max-consistency operator \circ exists in $\mathbf{N}(\mathbf{H}, F)$ and it is in the form $\circ(a, b) = (\neg\neg(a \vee b), \neg(a \vee b))$. Conversely, if \circ exists in $\mathbf{N}(\mathbf{H}, F)$, then \neg exists for all $a \vee b \in G$ and hence it exists for all the elements of the Boolean filter F by the previous argument. \square

The last theorem shows under which conditions a Heyting algebra \mathbf{H} and a Boolean filter F of \mathbf{H} induces an NL_c algebra over the Nelson lattice $\mathbf{N}(\mathbf{H}, F)$. It is clear that in every finite Heyting algebra all the elements have a dual pseudocomplement, whence all the elements of each of its Boolean filters also have. However, this might not be the case for infinite algebras. In order to see an example, consider the infinite Heyting algebra \mathbf{H} depicted in Figure 4. It is clear that the element x_1 does not have a dual pseudocomplement, while all other elements do have. Then, take the implicative filter $F = \{y_i \mid i \in N\}$. It is easy to see that F is Boolean since the quotient algebra \mathbf{H}/F is the Boolean algebra $\{0, 1\}$. Thus, although the dual pseudocomplement is not defined on the whole domain H , Theorem 4.5 ensures that $(\mathbf{N}(\mathbf{H}, F), \circ)$ is a NL_c algebra.

The final result of this subsection is a direct consequence of Theorem 4.5 above and it extends Sendewski's Theorem (see Theorem 4.1 (ii)) to Nelson lattices expanded by a consistency operators.

THEOREM 4.6

Let \mathbf{H} be a Heyting algebra and let F_1, F_2 be two Boolean filters of \mathbf{H} . Then $(\mathbf{N}(\mathbf{H}, F_1), \circ)$ is a NL_c subalgebra of $(\mathbf{N}(\mathbf{H}, F_2), \circ)$ iff $F_1 \subseteq F_2$.

PROOF. It is an immediate consequence of Theorem 4.5 and of the fact that the value of $\circ(a, b)$ only depends on a and b . \square

Let us close this subsection with a brief comment aimed at clarifying what is the situation for the two remaining cases: NL_c^{mB} and NL_c^{Bm} algebras.




FIGURE 4 A Heyting algebra that is not dually pseudo-complemented.

(1) As for NL_c^{mB} algebras, it is easy to see that the Boolean range of the $\max B$ -consistency operator is characterized, in terms of twist product, by requiring in Theorem 4.5 that the dual pseudocomplement \neg exists for all the elements of the Boolean filter F and, in addition, it maps all elements of F into a Boolean element of H .

(2) As for NL_c^{Bm} algebras, on the other hand, we need to define a new unary operator on a Heyting algebra \mathbf{H} , which we denote by \neg^b , called a *Boolean dual pseudocomplement*, defined by the following condition: for all $x \in H$,

$$\neg^b x = \min\{z \in B(\mathbf{H}) \mid x \vee z = 1\}.$$

Notice that, similarly to a dual pseudocomplement that assigns a Boolean value to an element of a Heyting algebra, \neg^b also ranges on the Boolean subalgebra $B(\mathbf{H})$ of \mathbf{H} . But notice that the operator \neg^b need not always exist since, for a fixed x in a Heyting algebra \mathbf{H} , the set $\{z \in B(\mathbf{H}) \mid x \vee z = 1\}$ might not have a minimum in $B(\mathbf{H})$. Again, a variant of Theorem 4.5 holds for NL_c^{Bm} algebras stating that if \mathbf{H} is a Heyting algebra and is F a Boolean filter of \mathbf{H} , $\neg^b x$ exists for all $x \in F$ iff the $\max B$ -consistency operator \circ exists in $N(\mathbf{H}, F)$. In case it exists, the $B\max$ -consistency operator \circ is definable in $N(\mathbf{H}, F)$ as follows: for every (a, b) , $\circ(a, b) = (\neg^b(a \vee b), \neg^b(a \vee b))$.

4.2 More on subdirectly irreducible and simple NL_c algebras

In the light of the results of the previous subsection, we are now in position to add some results on subdirectly irreducible NL_c algebras. In particular this subsection is devoted to show that Theorem 3.10 cannot be extended to infinite Nelson lattices with a \max -consistency operator. To this end, consider a Heyting algebra with dual pseudocomplement $\mathbf{H} = (H, \wedge, \vee, \rightarrow_H, \neg, 0, 1)$ and a normal Boolean filter F_B of \mathbf{H} (recall how normal filters are defined at the beginning this section). Since \mathbf{H} has a dual pseudocomplement, also F_B has it as well, and hence the

max-consistency operator \circ can be defined on $\mathbf{N}(\mathbf{H}, F_B)$ as in Lemma 4.4: for all $(a, b) \in N(\mathbf{H}, F_B)$, $\circ(a, b) = (\neg\neg(a \vee b), \neg(a \vee b))$. Now define the following subset of $N(\mathbf{H}, F)$:

$$G = \{(a, b) \in N(\mathbf{H}, F_B) \mid a \in F_B\}. \quad (3)$$

Then the following holds.

LEMMA 4.7

The set G is a \circ -filter of $(\mathbf{N}(\mathbf{H}, F_B), \circ)$. Furthermore if F_B is proper then so is G .

PROOF. Let us start showing that G is an implicative filter, and to this end let us prove that G is closed under $*$. If $(a, b), (a', b') \in G$, then $a, a' \in F_B$, whence $a \wedge a' \in F_B$ as well. Therefore, $(a, b) * (a', b') = (a \wedge a', (a \rightarrow_H b') \wedge (a' \rightarrow_H b)) \in G$.

Second, let us prove that G satisfies conditions (F1) and (F2). As for the former, assume that $(a, b) \rightarrow (a', b'), (a', b') \rightarrow (a, b) \in G$ and let us show that $\circ((a, b) \rightarrow (a', b')), \circ((a', b') \rightarrow (a, b)) \in G$. For all $(x, y), (z, k) \in N(\mathbf{H}, F_B)$, $(x, y) \rightarrow (z, k) \in G$ iff $(x \rightarrow_H x'), (y' \rightarrow_H y) \in F_B$, therefore

$$a \rightarrow_H a', b' \rightarrow_H b, a' \rightarrow_H b, b \rightarrow_H b' \in F_B.$$

In addition $a \vee b, a' \vee b' \in F_B$ because $(a, b), (a', b') \in N(\mathbf{H}, F_B)$, whence $\neg\neg(a \vee b), \neg\neg(a' \vee b') \in F_B$ since F_B is normal. Therefore one immediately has (i) $\neg\neg(a \vee b) \rightarrow_H \neg\neg(a' \vee b') \in F_B$ and (ii) $\neg\neg(a' \vee b') \rightarrow_H \neg\neg(a \vee b) \in F_B$. Furthermore, by hypothesis $a \leftrightarrow_H a', b \leftrightarrow_H b' \in F_B$, whence (iii) $\neg(a \vee b) \leftrightarrow_H \neg(a' \vee b') \in F_B$. Therefore, $\circ(a, b) \rightarrow \circ(a', b') = (\neg\neg(a \vee b), \neg(a \vee b)) \rightarrow_H (\neg\neg(a' \vee b'), \neg(a' \vee b')) = (\neg\neg(a \vee b) \rightarrow \neg\neg(a' \vee b'), \neg\neg(a \vee b) \wedge \neg(a' \vee b')) \in G$ by (i) and by (ii) $\circ(a', b') \rightarrow \circ(a, b) \in G$ as well. Thus (F1) holds.

As for (F2), assume that $(a, b) \vee \neg(a, b) \vee \neg(c, d) \in G$, or equivalently by definition of G , that $a \vee b \vee d \in F_B$. Thus, in order to prove that $(c, d) \rightarrow \circ(a, b) \in G$, we need to show (i) $c \rightarrow_H \neg\neg(a \vee b) \in F_B$ and (ii) $\neg(a \vee b) \rightarrow_H d \in F_B$. Now, since $(a, b) \in N(\mathbf{H}, F_B)$, $a \vee b \in F_B$ and $\neg\neg(a \vee b) \in F_B$ because F_B is normal. Therefore, $c \rightarrow_H \neg\neg(a \vee b) \in F_B$. As to prove (ii), notice that every Heyting algebra satisfies $\neg x \vee y \leq x \rightarrow_H y$, and hence $\neg(a \vee b) \rightarrow_H d \geq \neg\neg(a \vee b) \vee d \geq \neg\neg(a \vee b) \in F_B$.

Finally, let us observe that if F_B is proper, i.e. if $\{1\} \neq F_B \neq H$, then by definition $\{1\} \neq G \neq N(\mathbf{H}, F_B)$ and hence G is proper as well. \square

LEMMA 4.8

Let \mathbf{H} be a Heyting algebra with dual pseudocomplement and F_B a Boolean filter of \mathbf{H} . Then, for every proper \circ -filter G of $\mathbf{N}(\mathbf{H}, F_B)$, the set $F(G) = \{a \in F_B \mid \exists b \in H, (a, b) \in G\}$ is a proper normal filter of \mathbf{H} .

PROOF. The fact that $F(G)$ is an implicative filter is immediate. Thus, let us assume that $a \in F(G)$. By definition, it means that $(a, b) \in G$ for some $b \in H$. Since $(a, b) \leq (a, 0)$ and $(a, 0)$ clearly belongs to $N(\mathbf{H}, F_B)$, $(a, 0) \in F(G)$ as well. Therefore, $\circ(a, 0) = (\neg\neg a, \neg a) \in G$ because G is a \circ -filter, and hence $\neg\neg a \in F(G)$ proving that $F(G)$ is normal.

It is easy to see that if G is proper, then $F(G)$ is proper as well. \square

THEOREM 4.9


Let \mathbf{H} be a Heyting algebra with dual pseudocomplement and F_B a Boolean filter of \mathbf{H} . Then $G = \{(a, b) \in N(\mathbf{H}, F_B) \mid a \in F_B\}$ is a proper \circ -filter of $(\mathbf{N}(\mathbf{H}, F_B))$ iff F_B is proper and normal. Furthermore, if F_B is the minimal filter of \mathbf{H} , then G is the minimal filter of $\mathbf{N}(\mathbf{H}, F_B)$.

PROOF. In the light of Lemma 4.7 and Lemma 4.8, in order to prove the first part of the claim, it is enough to observe that if G is defined as in (3) and $F(G)$ is as in Lemma 4.8, then indeed $F(G) = F_B$. Therefore, let us prove the last claim and assume that G is not minimal. Thus, let R be a \circ -filter of $\mathbf{N}(\mathbf{H}, F_B)$ such that $R \subset G$. Then $F(R) \subset F(G) = F_B$ and F_B would not be minimal, a contradiction. \square

As an application of this last result, let us now show that Theorem 3.10 fails for general infinite structures, i.e. there exist NL_c algebras that are subdirectly irreducible but not simple. To this end, consider the following example that we have elaborated from an insight provided by Taylor in a personal communication and that can be found in [49, Figure 8.1 of page 99].

EXAMPLE 4.10

Let $Z = \{\omega\} \cup X \cup Y$ being $X = \{x_i : i \in \mathbb{N}\}$ and $Y = \{y_i : i \in \mathbb{N}\}$ and the following diagram of the partial order on Z :



Let us topologize X in the following manner: $U \subseteq X$ is open iff either $\omega \notin U$ or $\omega \in U$ and $X \setminus U$ is finite. Topologize Y similarly. Therefore, U is open in Z if only if $U \cap X$ and $U \cap Y$ are open in X and Y , respectively. Let H be the set of coplen upsets of Z . A direct computation shows those are the following subsets of Z :

Type (1): subsets U not containing ω and containing finite subsets of elements of X and Y and satisfying the condition that if $y_i \in U$ then $x_i, x_{i-1} \in U$ (in particular if $y_1 \in U$ then $x_1 \in U$ as the particular case of $i = 1$).

Type (2): subsets U containing ω and containing all elements of X and Y except, at most, a finite subset of them and satisfying the condition that if $y_i \in U$ then $x_i, x_{i-1} \in U$.

Using the standard construction for Priestley duality is possible to see that $\mathbf{H} = (H, \cap, \cup, \rightarrow_H, \neg, \emptyset, Z)$ is a Heyting algebra in which the dual pseudocomplementation \neg is indeed definable. Moreover, let us observe the following facts:

- the pseudocomplement is defined as $\neg D = D \rightarrow_H \emptyset$.
- the atoms of \mathbf{H} are the subsets $\{x_i\}$ for $i \in \mathbb{N}$.
- the antiatoms of \mathbf{H} are the subsets $\{\omega\} \cup X \cup (Y \setminus \{y_i\})$ for every $i \in \mathbb{N}$.

As proved in [45], the congruences of \mathbf{H} are in one-to-one correspondence with normal filters, i.e. lattice filters F closed by a combination of the two pseudocomplementations, i.e. if $U \in F$, then $\neg\neg(U) \in F$.

By the result about Priestley duality we can prove that the normal filters of \mathbf{H} are (i) the one containing only the maximum $\{Z\}$, (ii) the one containing all subsets of type (2) and (iii) the full algebra \mathbf{H} . Then \mathbf{H} has only three congruences that form a three element chain and thus it is subdirectly irreducible but not simple.

In order to be self-contained let us sketch the proof of this fact:

- (1) Compute the normal filter F generated by the antiatom $D = \{\omega\} \cup X \cup (Y \setminus \{y_1\})$. Since $\neg D = \{y_1, x_1\}$ and $\neg\neg(D) = \{\omega\} \cup X \cup (Y \setminus \{y_1, y_2\})$, we deduce that $\{\omega\} \cup X \cup (Y \setminus \{y_2\}) \in F$ and, recursively, we obtain that all antiatoms belong to F , and so F consists of all subsets of type (2)
- (2) Compute the normal filter generated by any subset of type (1). It is easy to see that it is the full Heyting algebra.
- (3) Then the set of congruences of \mathbf{H} has exactly three congruences: the identity, the one corresponding to the normal filter F and the full algebra.

Also notice that the normal filter F defined in (1) above is indeed Boolean, because an easy computation shows that the quotient \mathbf{H}/\equiv_F is the two-element Boolean algebra.

Then let us consider the NL_c algebra $(\mathbf{N}(\mathbf{H}, F), \circ)$. Notice that $N(\mathbf{H}, F) = \{(D, E) \in H \times H : D \cap E = \emptyset, D \cup E \in F\}$ is indeed the set of pairs (D, E) such that D is of type (1) and $E \subseteq \neg D$. By Theorem 4.9, the \circ -filters of $(\mathbf{N}(\mathbf{H}, F), \circ)$ are the singleton $\{(Z, \emptyset)\}$, the set $\mathcal{F} = \{(D, E) \in R(H, F) : D \in F\}$ and the full Nelson lattice $N(\mathbf{H}, F)$. Thus the \circ -congruences of $(\mathbf{N}(\mathbf{H}, F), \circ)$ form a three-element chain and hence our NL_c algebra is subdirectly irreducible but not simple.

REMARK 4.11

The residuated lattice of the last example is not an IMTL algebra, i.e. is not prelinear, it does not satisfy the equation $(x \rightarrow y) \vee (y \rightarrow x) = 1$ as the following example shows: take $A = \{x_i\}$ and $B = \{x_{i-1}\}$, two atoms of the Nelson lattice. Then an easy computation shows that

- $A \rightarrow B = \{w\} \cup X \cup Y \setminus \{x_i, y_i, y_{i+1}\}$,
- $A \rightarrow B = \{w\} \cup X \cup Y \setminus \{x_{i-1}, y_i, y_{i-1}\}$.

Therefore, $(A \rightarrow B) \vee (B \rightarrow A) = \{w\} \cup X \cup Y \setminus \{y_i\}$, which is not the maximum of the lattice, and thus the residuated lattice is not prelinear.

5 Adding consistency operators to the logic dIRL and its paraconsistent companions

Let us recall from Section 2.1 the logic dIRL, the logic corresponding to the variety of dIRLs, that can be presented as the axiomatic extension of FL_{ew} with the following axioms:

- (Inv) $\phi \rightarrow \neg\neg\phi$,
 (Dist) $\phi \wedge (\psi \vee \chi) \rightarrow (\phi \wedge \psi) \vee (\phi \wedge \chi)$

and its degree-preserving companion dIRL^{\leq} . Since the residual negation \neg in dIRL is involutive, and hence it does not prove the pseudo-complementation axiom, i.e. $\not\vdash_{\text{NL}} \phi \wedge \neg\phi \rightarrow \perp$, the logic dIRL^{\leq} satisfies

$$\phi, \neg\phi \not\vdash_{\text{dIRL}^{\leq}} \perp$$

and hence it is paraconsistent with respect to the residual negation \neg . However, dIRL^{\leq} is not an LFI since we cannot define, in its language, a consistency connective \circ satisfying

$$\circ\phi, \phi, \neg\phi \vdash_{\text{dIRL}^{\leq}} \psi$$

for every ϕ and ψ . This has been the main motivation in the previous sections for the algebraic study of expansions of dIRL algebras with suitable consistency operators. With this background, in this section, we will consider the logical counterparts of those expansions, namely the expansion of the

logic dIRL with three classes of consistency operators \circ whose degree-preserving companions will turn out to be LFIs.

5.1 dIRL logic with a max-consistency operator

In this section we start by defining the expansion of the logic dIRL with a new connective \circ whose algebraic interpretation corresponds to a max-consistency operator.

DEFINITION 5.1

The logic dIRL_c is the expansion of the logic dIRL in a language that incorporates a new unary connective \circ with the following additional axiom:

$$(A1) \quad \neg(\varphi \wedge \neg\varphi \wedge \circ\varphi)$$

and inference rules:

$$(CNG) \quad \frac{\varphi \leftrightarrow \psi}{\circ\varphi \leftrightarrow \circ\psi} \quad (\text{Max}) \quad \frac{\varphi \vee \neg\varphi \vee \neg\psi}{\psi \rightarrow \circ\varphi} .$$

Clearly, the axiom (A1) and the rule (Max) are the logical counterparts of conditions ($\circ 1$) and ($\circ 2$) in dIRL_c algebras, respectively, while the rule (CNG) enforces \circ to be congruent w.r.t. logical equivalence.

Some observations follow:

- (i) Both $\circ\top$ and $\circ\perp$ are derivable in dIRL_c , it suffices in (Max) to take $\psi = \top$ and then $\varphi = \top$ (resp. $\varphi = \perp$).
- (ii) The rule of necessitation for \circ :

$$\frac{\varphi}{\circ\varphi}$$

is also derivable. Indeed, assuming φ , we can derive $\varphi \rightarrow \top$ and $\top \rightarrow \varphi$ in dIRL_c. As a matter of fact, $\varphi \rightarrow \top$ and $\varphi \rightarrow (\psi \rightarrow \varphi)$ both are theorems of RL (and hence a theorem of dIRL_c a fortiori). Thus, $\varphi \vdash_{\text{dIRL}_c} \varphi \leftrightarrow \top$.

- (ii) Now, by (CNG) with $\psi = \top$, $\varphi \leftrightarrow \top \vdash_{\text{dIRL}_c} \circ\varphi \leftrightarrow \circ\top$ and since by the above (i) $\circ\top$ is a theorem of dIRL_c, by $\circ\varphi \leftrightarrow \circ\top$ and $\circ\top$ we get $\circ\varphi$ by modus ponens.

LEMMA 5.2

The following derivabilities hold in dIRL_c :

- (i) $\varphi \vee \neg\varphi \vdash_{\text{dIRL}_c} \circ\varphi$
- (ii) $\vdash_{\text{dIRL}_c} \circ\varphi \rightarrow \neg\varphi \vee \varphi$
- (iii) $\varphi \vee \neg\varphi \dashv\vdash_{\text{dIRL}_c} \circ\varphi$

PROOF.

- (i) It follows by using the rule (Max) with $\psi = \top$.
- (ii) Notice that in dIRLs the inequality $\sim x \vee y \leq x \rightarrow y$ holds true, and hence, since dIRL is sound and complete with respect to dIRL algebras, $(\neg\varphi \vee \psi) \rightarrow (\varphi \rightarrow \psi)$ is a theorem in dIRL. Therefore, since $\neg(\varphi \wedge \neg\varphi \wedge \circ\varphi)$ is logically equivalent to $\neg\varphi \vee \varphi \vee \neg\circ\varphi$ and, by the previous observation, in dIRL_c the latter logically implies $\circ\varphi \rightarrow \neg\varphi \vee \varphi$. Hence $\text{dIRL}_c \vdash \circ\varphi \rightarrow \neg\varphi \vee \varphi$.
- (iii) Direct from (i) and (ii). □

It is easy to check that, due to the presence of the rule (CNG) for \circ , dIRL_c is a Rasiowa implicative logic, hence it is algebraizable, and its equivalent algebraic semantics is given by the quasi-variety of dIRL_c algebras studied in Section 3.1.

PROPOSITION 5.3

dIRL_c is strongly complete w.r.t. the class of dIRL_c algebras.

Now, thanks to results on dIRL_c algebras in Section 3.1, we can prove that dIRL_c is a conservative expansion of dIRL .

PROPOSITION 5.4

dIRL_c is a conservative expansion of dIRL , i.e. if $\Gamma \cup \{\varphi\}$ is a set of formulas in the language of dIRL , i.e. without the connective \circ , then $\Gamma \vdash_{\text{dIRL}_c} \varphi$ iff $\Gamma \vdash_{\text{dIRL}} \varphi$.

PROOF. The right-to-left direction is obvious. Thus, assume $\Gamma \not\vdash_{\text{dIRL}} \varphi$, then there is a subdirectly irreducible dIRL algebra \mathbf{A} and an \mathbf{A} evaluation of formulas e such that $e(\psi) = 1$ for all $\psi \in \Gamma$ and $e(\varphi) < 1$. By Lemma 3.7, the unary operation \circ on A defined as $\circ(1) = \circ(0) = 1$ and $\circ(x) = 0$ otherwise makes (\mathbf{A}, \circ) a simple dIRL_c algebra. Then e can be extended in the obvious way to an (\mathbf{A}, \circ) evaluation e' agreeing with e on the formulas not containing the connective \circ . Therefore, we have $e'(\psi) = 1$ for all $\psi \in \Gamma$ and $e'(\varphi) < 1$, i.e. $\Gamma \not\vdash_{\text{dIRL}_c} \varphi$. \square

Now we move to the logic dIRL_c^{\leq} , the degree-preserving companion of the logic dIRL_c .

DEFINITION 5.5

The degree-preserving companion of logic dIRL_c is the logic dIRL_c^{\leq} defined by the following axioms and rules:

- Axioms of dIRL_c^{\leq} are those of dIRL_c
- Rules of dIRL_c^{\leq} are the following:
 - (Adj- \wedge) from φ and ψ derive $\varphi \wedge \psi$
 - (MP- r) if $\vdash_{\text{dIRL}_c} \varphi \rightarrow \psi$, then from φ and $\varphi \rightarrow \psi$, derive ψ
 - (CNG- r) if $\vdash_{\text{dIRL}_c} \varphi \leftrightarrow \psi$, then from $\varphi \leftrightarrow \psi$ derive $\varphi \circ \leftrightarrow \varphi \circ \psi$
 - (Max- r) if $\vdash_{\text{dIRL}_c} \varphi \vee \neg\varphi \vee \neg\psi$, then from $\varphi \vee \neg\varphi \vee \neg\psi$ derive $\psi \rightarrow \varphi \circ$.

We will denote by $\vdash_{\text{dIRL}_c}^{\leq}$ its corresponding notion of proof. By Proposition 2.1, this axiomatization is sound and complete w.r.t. the semantical consequence relation $\models_{\text{dIRL}_c}^{\leq}$, defined in the obvious way.

Observe that the rules (MP- r), (CNG- r) and (Max- r) are *restricted* in the sense that the premises are required to satisfy an extra condition, to be theorems of the logic.

Now we can check that dIRL_c^{\leq} is in fact a strong LFI in the sense of Definition 3.2.

PROPOSITION 5.6

The logic dIRL_c^{\leq} is a strong LFI w.r.t. the negation \neg and the consistency operator \circ .

PROOF. By the above-mentioned completeness of dIRL_c^{\leq} w.r.t. $\models_{\text{dIRL}_c}^{\leq}$, it is enough to check the conditions (i.a), (i.b), (i.c) and (ii) in Definition 3.2 for the semantical consequence relation $\models_{\text{dIRL}_c}^{\leq}$. If p, q are two different propositional variables, then we have the following:

- (i.a) any evaluation e such that $e(p) > 0$ and $e(q) = 0$ is such that $e(p \wedge \neg p) > e(q)$ and hence $p, \neg p \not\models_{\text{dIRL}_c}^{\leq} q$;

(i.b) any evaluation e such that $e(p) = e(op) = 1$ and $e(q) = 0$ is such that $e(p \wedge op) > e(q)$, and hence $p, op \not\models_{\text{dIRL}_c}^{\leq} q$;

(i.c) any evaluation e such that $e(\neg p) = e(op) = 1$ and $e(q) = 0$ is such that $e(\neg p \wedge op) > e(q)$, and hence $\neg p, op \not\models_{\text{dIRL}_c}^{\leq} q$; and

(ii) any evaluation e is such that $e(p \wedge \neg p \wedge op) = 0$ and hence $p, \neg p, op \models_{\text{dIRL}_c}^{\leq} \perp$. \square

In the context of LFIs, it is a desirable property to recover the classical reasoning by means of the consistency connective \circ (see [13]). Specifically, let CPL be classical propositional logic. If L is a given LFI such that its reduct to the language of CPL is a sublogic of CPL, then a derivability adjustment theorem (DAT) for L with respect to CPL is as follows: for every finite set of formulas $\Gamma \cup \{\varphi\}$ in the language of CPL, there exists a finite set of formulas Θ in the language of L, whose variables occur in formulas of $\Gamma \cup \{\varphi\}$, such that

$$\text{(DAT)} \quad \Gamma \vdash_{\text{CPL}} \varphi \text{ iff } \circ(\Theta) \cup \Gamma \vdash_{\text{L}} \varphi.$$

When the operator \circ enjoys the *propagation property* in the logic L with respect to a set X of classical connectives, i.e. when

$$\circ \varphi_1, \dots, \circ \varphi_n \vdash_{\text{L}} \circ \#(\varphi_1, \dots, \varphi_n),$$

for every n -ary connective $\# \in X$ and formulas $\varphi_1, \dots, \varphi_n$ built with connectives from X , then the DAT takes the following, simplified form: for every finite set of formulas $\Gamma \cup \{\varphi\}$ in the language of CPL,

$$\text{(PDAT)} \quad \Gamma \vdash_{\text{CPL}} \varphi \text{ iff } \{\circ p_1, \dots, \circ p_m\} \cup \Gamma \vdash_{\text{L}} \varphi$$

where $\{p_1, \dots, p_m\}$ is the set of propositional variables occurring in $\Gamma \cup \{\varphi\}$.

In particular, checking whether dIRL_c^{\leq} satisfies the propagation property for the connectives $X = \{\perp, \wedge, \&, \rightarrow\}$ amounts to check the following conditions:

$$\left\{ \begin{array}{l} \vdash_{\text{dIRL}_c} \circ \perp \\ \vdash_{\text{dIRL}_c} (\circ \varphi \wedge \circ \psi) \rightarrow \circ(\varphi \# \psi), \quad \text{for each binary } \# \in X. \end{array} \right.$$

In Proposition 5.8 below we will show that the operator \circ of dIRL_c satisfies the propagation property with respect to all connectives of logic dIRL .⁷ Our proof is algebraic and for that we will need the claims proved in the next lemma.

LEMMA 5.7

The following equations hold in the quasivariety of dIRL_c algebras:

- (i) $x \wedge \sim x \wedge \sim y \wedge \circ(x \vee (y \wedge z)) = 0$
- (ii) $(x * y) \vee ((\sim x \vee \sim \circ(x)) \wedge y) = y$.

PROOF.

- (i) By $(\circ 1)$, we know that the equation $(x \vee y) \wedge \sim(x \vee y) \wedge \circ(x \vee y) = 0$ holds, i.e. we have $(x \vee y) \wedge \sim x \wedge \sim y \wedge \circ(x \vee y) = 0$. But since $x \leq (x \vee y)$ we also have that $x \wedge \sim x \wedge \sim y \wedge \circ(x \vee y) = 0$. Now, by replacing y by $y \wedge z$ in the former equation, we get $x \wedge \sim x \wedge \sim (y \wedge z) \wedge \circ(x \vee (y \wedge z)) = 0$, and since $\sim(y \wedge z) \geq \sim y$, we finally get $x \wedge \sim x \wedge \sim y \wedge \circ(x \vee (y \wedge z)) = 0$.

⁷The proof of the propagation property for the monoidal conjunction and is in fact a re-elaboration of a proof found by the automated theorem-prover Prover9 [34].

(ii) We start from the equation $x \vee \sim x \vee \sim \circ(x) = 1$, we multiply by y in both sides and get the following equations:

$$\begin{aligned} y * (x \vee \sim x \vee \sim \circ(x)) &= y, (x * y) \vee y * (\sim x \vee \sim \circ(x)) = y, y = (x * y) \vee (y * (\sim x \vee \sim \circ(x))) \leq \\ &(x * y) \vee (y \wedge (\sim x \vee \sim \circ(x))) \leq y \vee y = y. \\ \text{Hence } (x * y) \vee (y \wedge (\sim x \vee \sim \circ(x))) &= y. \quad \square \end{aligned}$$

PROPOSITION 5.8

The logic dIRL_c satisfies the following conditions:

- $\vdash_{\text{dIRL}_c} \circ \perp$, and
- $\vdash_{\text{dIRL}_c} (\circ \varphi \wedge \circ \psi) \rightarrow \circ(\varphi \# \psi)$, for $\# \in \{\wedge, \vee, \&, \rightarrow\}$,

and thus, in dIRL_c^{\leq} the consistency connective \circ satisfies the propagation properties w.r.t. all connectives of the logic dIRL .

PROOF. The first condition is satisfied since $\circ \perp$ is a theorem of the logic, as observed above after Def. 5.1. The proof is algebraic, so we have to prove that, for any dIRL_c algebra (\mathbf{A}, \circ) , the following condition holds: $\circ(x) \wedge \circ(y) \leq \circ(x \# y)$, for all $\# \in \{\wedge, \vee, *\}$. As for the lattice operations \wedge and \vee , we have

$$\begin{aligned} \circ(x \wedge y) &= \max\{z \mid (x \wedge y) \wedge \sim(x \wedge y) \wedge z = 0\} \\ &= \max(\max\{z \mid x \wedge y \wedge \sim x \wedge z = 0\}, \max\{z \mid x \wedge y \wedge \sim y \wedge z = 0\}) \\ &\geq \max(\max\{z \mid x \wedge \sim x \wedge z = 0\}, \max\{z \mid y \wedge \sim y \wedge z = 0\}) \\ &= \circ(x) \vee \circ(y) \geq \circ(x) \wedge \circ(y). \\ \\ \circ(x \vee y) &= \max\{z \mid (x \vee y) \wedge \sim(x \vee y) \wedge z = 0\} \\ &= \max(\max\{z \mid x \wedge \sim x \wedge \sim y \wedge z = 0\}, \max\{z \mid y \wedge \sim x \wedge \sim y \wedge z = 0\}) \\ &\geq \max(\max\{z \mid x \wedge \sim x \wedge z = 0\}, \max\{z \mid y \wedge \sim y \wedge z = 0\}) \\ &= \circ(x) \vee \circ(y) \geq \circ(x) \wedge \circ(y). \end{aligned}$$

Next, let us show that $\circ(x) \wedge \circ(y) \leq \circ(x * y)$. By (o2), it is enough to prove that $(x * y) \wedge \sim(x * y) \wedge \circ(x) \wedge \circ(y) = 0$. To do so, by (i) of Lemma 5.7, we can start from the equation

$$u \wedge \sim u \wedge \sim v \wedge \circ(u \vee (v \wedge t)) = 0,$$

and make the following substitutions: replace u by $x * y$, v by $\sim(x \wedge \circ(x))$ and t by y , and get

$$x * y \wedge \sim(x * y) \wedge \sim(\sim(x \wedge \circ(x))) \wedge \circ(x * y \vee (\sim(x \wedge \circ(x)) \wedge y)) = 0,$$

i.e. $x * y \wedge \sim(x * y) \wedge x \wedge \circ(x) \wedge \circ(x * y \vee (\sim(x \wedge \circ(x)) \wedge y)) = 0$. But $x * y \leq x$ and by (ii) of Lemma 5.7, $x * y \vee (\sim(x \wedge \circ(x)) \wedge y) = y$, and hence we finally get $x * y \wedge \sim(x * y) \wedge \circ(x) \wedge \circ(y) = 0$, as claimed.

Finally, since by (ii) of Proposition 3.13, we have $\circ(x) = \circ(\sim x)$, and $x \rightarrow y = \sim(x * \sim y)$, the propagation property for \rightarrow , i.e. $\circ(x) \wedge \circ(y) \leq \circ(x \rightarrow y)$, directly follows as well. \square

Finally, we are interested in investigating whether we can expect some form close to (PDAT) for the logic dIRL_c^{\leq} . Next theorem provides a result in this direction showing that it is possible to recover classical logic derivations inside the LFI logic dIRL_c^{\leq} .

THEOREM 5.9 (PDAT like for dIRL_c^{\leq}).

Let $\Gamma \cup \{\varphi\}$ be a finite set of formulas in the language of CPL and let $\{p_1, \dots, p_m\}$ the set of

propositional variables appearing in $\Gamma \cup \{\varphi\}$. Then, there is a natural $k > 0$ such that

$$\Gamma \vdash_{\text{CPL}} \varphi \text{ iff } \Gamma \vdash_{\text{dIRL}_c}^{\leq} \left(\bigwedge_{i=1}^m \circ(p_i) \right)^k \rightarrow \varphi.$$

PROOF. Assume $\Gamma \vdash_{\text{CPL}} \varphi$, or equivalently, $\vdash_{\text{CPL}} \Gamma^\wedge \rightarrow \varphi$. First of all, observe that we also have

$$\{p_i \vee \neg p_i \mid i = 1, 2, \dots, m\} \vdash_{\text{dIRL}} \Gamma^\wedge \rightarrow \varphi,$$

since the class of Boolean subalgebras of dIRLs is in fact the whole class of Boolean algebras. Then by the local deduction-detachment theorem of dIRL, there is a natural $k > 0$ such that $\vdash_{\text{dIRL}} (\bigwedge_{i=1}^m (p_i \vee \neg p_i))^k \rightarrow (\Gamma^\wedge \rightarrow \varphi)$, and thus this theorem is also valid in dIRL_c . But this implies, by (iii) of Lemma 5.2, $\vdash_{\text{dIRL}_c} (\bigwedge_{i=1}^m (\circ p_i))^k \rightarrow (\Gamma^\wedge \rightarrow \varphi)$ and hence $\Gamma \vdash_{\text{dIRL}_c} (\bigwedge_{i=1}^m \circ(p_i))^k \rightarrow \varphi$ as well.

Conversely, assume $\vdash_{\text{dIRL}_c} (\bigwedge_{i=1}^m (\circ p_i))^k \rightarrow (\Gamma^\wedge \rightarrow \varphi)$, and let e be any evaluation on the two-element Boolean algebra $\mathbf{2}$. Since $\mathbf{2}$ is a dIRL_c algebra with $\circ(0) = \circ(1) = 1$, we have $e((\bigwedge_{i=1}^m \circ p_i)^k \rightarrow (\Gamma^\wedge \rightarrow \varphi)) = 1$. But then we necessarily have $e(\Gamma^\wedge \rightarrow \varphi) = 1$, because $e(\bigwedge_{i=1}^m \circ p_i) = 1$. Therefore, $\Gamma^\wedge \rightarrow \varphi$ is a CPL tautology and so $\Gamma \vdash_{\text{CPL}} \varphi$. \square

Note that, in the above formulation of the PDAT theorem for dIRL_c , the natural number k depends on the formulas $\Gamma \cup \{\varphi\}$ involved, and hence it is not possible to fix such exponent k in advance. However, if we replace the logic dIRL_c by any of its axiomatic extensions L validating the n -potency axiom

$$\varphi^n \rightarrow \varphi^{n+1}$$

for some n , then one can be more precise and state the PDAT theorem for L_c^{\leq} : for any set of formulas $\Gamma \cup \{\varphi\}$, the following condition holds:

$$\Gamma \vdash_{\text{CPL}} \varphi \text{ iff } \Gamma \vdash_{\text{dIRL}_c}^{\leq} \left(\bigwedge_{i=1}^m \circ(p_i) \right)^n \rightarrow \varphi.$$

In particular, when L is Nelson logic or its prelineal extension NM logic, which are 2-potent logics, we can take $n = 2$ in the previous expression.

5.2 *dIRL logic with a maxB-consistency operator*

In this section we consider the logic $\text{dIRL}_c^{\text{mB}}$ corresponding to the quasivariety of $\text{dIRL}_c^{\text{mB}}$ algebras and its paraconsistent degree preserving companion $(\text{dIRL}_c^{\text{mB}})^{\leq}$. The content is mostly parallel to the previous section, only some details change. In particular we get a simpler PDAT theorem due to the fact that the consistency connective is Boolean in this case.

DEFINITION 5.10

The logic $\text{dIRL}_c^{\text{mB}}$ is the expansion of the logic dIRL, in a language that incorporates a new unary connective \circ , with the following additional axioms:

- (A1) $\neg(\varphi \wedge \neg\varphi \wedge \circ\varphi)$
- (A2) $\circ\varphi \vee \neg\circ\varphi$

and inference rules:

$$(\text{CNG}) \frac{\varphi \leftrightarrow \psi}{\circ\varphi \leftrightarrow \circ\psi} \quad (\text{MaxB}) \frac{\neg(\varphi \wedge \neg\varphi \wedge \psi), \quad \psi \vee \neg\psi}{\psi \rightarrow \circ\varphi}.$$

Analogously to the case of $d\text{IRL}_c$, thanks to Lemma 3.7, we also have that $d\text{IRL}_c^{\text{mB}}$ is a conservative expansion of $d\text{IRL}$.

We can check now that $d\text{IRL}_c^{\text{mB}}$ keeps validating the same properties in Lemma 5.2 for $d\text{IRL}_c$, plus a new one.

LEMMA 5.11

The logic $d\text{IRL}_c^{\text{mB}}$ enjoys the following properties:

- (i) $\varphi \vee \neg\varphi \vdash_{d\text{IRL}_c^{\text{mB}}} \circ\varphi$
- (ii) $d\text{IRL}_c^{\text{mB}} \vdash \circ\varphi \rightarrow \neg\varphi \vee \varphi$
- (iii) $\varphi \vee \neg\varphi \not\vdash_{d\text{IRL}_c^{\text{mB}}} \circ\varphi$
- (iv) $d\text{IRL}_c^{\text{mB}} \vdash \circ\circ\varphi$.

PROOF. (i), (ii) and (iii) are as in Lemma 5.2. As for (iv), using (Max) with $\psi = \top$, we have $\neg(\circ\varphi \wedge \neg\circ\varphi) \vdash \circ\circ\varphi$, but the premise is in fact Axiom (A2). \square

Again, as in the case of the logic $d\text{IRL}_c$, it is easy to check that, due to the (Cong) rule for \circ , $d\text{IRL}_c^{\text{mB}}$ is a Rasiowa implicative logic and hence it is algebraizable. The equivalent algebraic semantics is now given by the quasi-variety of $d\text{IRL}_c^{\text{mB}}$ algebras.

We now introduce the degree-preserving companion of $d\text{IRL}_c^{\text{mB}}$.

DEFINITION 5.12

The degree-preserving companion of logic $d\text{IRL}_c^{\text{mB}}$ is the logic $(d\text{IRL}_c^{\text{mB}})^{\leq}$ defined by the following axioms and rules:

- Axioms of $(d\text{IRL}_c^{\text{mB}})^{\leq}$ are those of $d\text{IRL}_c^{\text{mB}}$
- Rules of $(d\text{IRL}_c^{\text{mB}})^{\leq}$ are those of $d\text{IRL}_c^{\leq}$ but replacing the rule (Max-r) by the following rule:

(MaxB-r) if $\vdash_{d\text{IRL}_c^{\text{mB}}} \neg(\varphi \wedge \neg\varphi \wedge \psi)$ and $\vdash_{d\text{IRL}_c^{\text{mB}}} \psi \vee \neg\psi$, then from $\neg(\varphi \wedge \neg\varphi \wedge \psi)$ and $\psi \vee \neg\psi$ derive $\psi \rightarrow \circ\varphi$

We will denote by $\vdash_{d\text{IRL}_c^{\text{mB}}}^{\leq}$ its corresponding notion of proof. As in the previous subsection, by Proposition 2.1, this axiomatization of $(d\text{IRL}_c^{\text{mB}})^{\leq}$ is sound and complete w.r.t. the semantical consequence relation $\models_{d\text{IRL}_c^{\text{mB}}}^{\leq}$, defined in the obvious way, and moreover $(d\text{IRL}_c^{\text{mB}})^{\leq}$ is a strong LFI as well. The proof is the same of Proposition 5.6 and it is omitted.

PROPOSITION 5.13

The logic $(d\text{IRL}_c^{\text{mB}})^{\leq}$ is a strong LFI w.r.t. to the negation \neg and the consistency operator \circ .

As in the case of the logic $d\text{IRL}_c$, the \circ connective in the logic $d\text{IRL}_c^{\text{mB}}$ nicely propagates through the rest of connectives $X = \{\perp, \wedge, \vee, \&, \rightarrow\}$. This is shown next, after a previous lemma, formally similar to Lemma 5.8, but this one is much easier to prove.

LEMMA 5.14

The following provability holds in $\text{dIRL}_c^{\text{mB}}$:

$$\varphi \vee \neg\varphi, \psi \vee \neg\psi \vdash_{\text{dIRL}_c^{\text{mB}}} (\varphi\#\psi) \vee \neg(\varphi\#\psi)$$

for every binary connective $\# \in X$.PROOF. Using a semantic argument, it holds since the binary connectives $\wedge, \vee, \&, \rightarrow$ are closed under Boolean values in every evaluation over any $\text{dIRL}_c^{\text{mB}}$ algebra. \square

PROPOSITION 5.15

 $\text{dIRL}_c^{\text{mB}}$ satisfies the following conditions:

- $\vdash_{\text{dIRL}_c^{\text{mB}}} \circ\perp$, and
- $\vdash_{\text{dIRL}_c^{\text{mB}}} (\circ\varphi \wedge \circ\psi) \rightarrow \circ(\varphi\#\psi)$, for every $\# \in \{\wedge, \vee, \&, \rightarrow\}$,

and thus, in $(\text{dIRL}_c^{\text{mB}})^{\leq}$ the consistency connective \circ satisfies the propagation property w.r.t. all the connectives in $X = \{\perp, \wedge, \vee, \&, \rightarrow\}$.PROOF. Straightforward from the above lemma, taking into account that, by (iii) of Lemma 5.11, $\varphi \vee \neg\varphi \dashv\vdash_{\text{dIRL}_c^{\text{mB}}} \circ\varphi$. \square This propagation property for \circ , together with the fact that $\circ\varphi$ is a Boolean formula for any φ , allows us to formulate the following PDAT theorem for $(\text{dIRL}_c^{\text{mB}})^{\leq}$ in simpler terms.THEOREM 5.16 (PDAT for $(\text{NL}_c^{\text{mB}})^{\leq}$).Let $\Gamma \cup \{\varphi\}$ be a finite set of formulas in the language of CPL and let $\{p_1, \dots, p_m\}$ the set of propositional variables appearing in $\Gamma \cup \{\varphi\}$. Then

$$\Gamma \vdash_{\text{CPL}} \varphi \text{ iff } \{\circ p_1, \dots, \circ p_m\} \cup \Gamma \vdash_{\text{dIRL}_c^{\text{mB}}}^{\leq} \varphi.$$

PROOF. The proof is essentially the same as in Theorem 5.9 for dIRL_c^{\leq} . The only difference is that now $(\bigwedge_{i=1}^m \circ(p_i))^n$ turns out to be equivalent to just $\bigwedge_{i=1}^n \circ(p_i)$ since this is a Boolean formula, and this conjunction can be moved back to the premise as the set of formulas $\{\circ p_1, \dots, \circ p_m\}$. \square

5.3 dIRL logic with a Bmax-consistency operator

Finally, we consider in this section the logic resulting from expanding the logic of dIRLs with a Bmax-consistency connective and its paraconsistent degree-preserving companion. Both of them are very similar to the case of maxB-consistency connective of previous section.

DEFINITION 5.17

The logic $\text{dIRL}_c^{\text{Bm}}$ is the axiomatic extension of dIRL_c with the following additional axiom:

$$(A4) \quad \circ\varphi \vee \neg\circ\varphi.$$

It is clear that $\text{dIRL}_c^{\text{Bm}}$ is thus algebraizable, with equivalent algebraic semantics given by the quasivariety of $\text{dIRL}_c^{\text{Bm}}$ algebras.

PROPOSITION 5.18

 $\text{dIRL}_c^{\text{Bm}}$ is strongly complete w.r.t the class of $\text{dIRL}_c^{\text{Bm}}$ algebras.

Actually, by definition, the logic $d\text{IRL}_c^{\text{Bm}}$ is an extension of $d\text{IRL}_c$, but it is also an extension of $d\text{IRL}_c^{\text{mB}}$, as the $d\text{IRL}_c$'s rule (Max) is stronger than the $d\text{IRL}_c^{\text{mB}}$'s rule (MaxB), and hence the latter is derivable in $d\text{IRL}_c^{\text{Bm}}$. As a consequence, all the properties that have been shown for $d\text{IRL}_c$ and $d\text{IRL}_c^{\text{mB}}$ also hold for $d\text{IRL}_c^{\text{Bm}}$.

This has also impact on the properties of the corresponding degree-preserving companion of $d\text{IRL}_c^{\text{Bm}}$.

DEFINITION 5.19

The degree-preserving companion of logic $d\text{IRL}_c^{\text{Bm}}$ is the logic $(d\text{IRL}_c^{\text{Bm}})^{\leq}$ defined as the axiomatic extension of $d\text{IRL}_c^{\leq}$ with the axiom (A4).

Of course, as in the previous cases, the logic $(d\text{IRL}_c^{\text{Bm}})^{\leq}$ is a strong LFI (details are omitted). Moreover, it follows that the consistency connective \circ in $(d\text{IRL}_c^{\text{Bm}})^{\leq}$ also satisfies the same propagation properties and the same PDAT theorem as for $(d\text{IRL}_c^{\text{mB}})^{\leq}$.

THEOREM 5.20 (PDAT for $(d\text{IRL}_c^{\text{Bm}})^{\leq}$).

Let $\Gamma \cup \{\varphi\}$ be a finite set of formulas in the language of CPL and let $\{p_1, \dots, p_m\}$ be the set of propositional variables appearing in $\Gamma \cup \{\varphi\}$. Then, the following condition holds:

$$\Gamma \vdash_{\text{CPL}} \varphi \text{ iff } \{\circ p_1, \dots, \circ p_m\} \cup \Gamma \vdash_{d\text{IRL}_c^{\text{Bm}}}^{\leq} \varphi.$$

5.4 Logics of formal undeterminedness

In this final subsection, we would like to emphasize that the logics $d\text{IRL}_c^{\leq}$, $(d\text{IRL}_c^{\text{mB}})^{\leq}$ and $(d\text{IRL}_c^{\text{Bm}})^{\leq}$ are not only LFIs but also Logics of Formal Undeterminedness (*LFUs*), the latter being introduced in [35].

LFUs are in a sense dual to LFIs, since in the same way that a consistency operator in an LFI controls the explosion in the presence of a contradiction, a ‘determinedness’ operator \circ in an LFU controls the law of the excluded middle, namely: given a paracomplete logic L , i.e. such that $\not\vdash_L \varphi \vee \sim\varphi$, if \circ is a determinedness operator in L , then $\circ\varphi \vdash_L \varphi \vee \sim\varphi$. More formally, we have the following definition.

DEFINITION 5.21

Let $L = \langle \Sigma, \vdash \rangle$ be a Tarskian, finitary and structural logic defined over a signature Σ with a disjunction \vee , a negation \sim and a primitive or defined unary connective \circ . Then L is said to be a *LFU* with respect to \sim and \circ if the following holds:

- (i) $\not\vdash \varphi \vee \sim\varphi$, for some φ ;
- (ii) there is a formula φ such that
 - (ii.a) $\circ\varphi \not\vdash \varphi$;
 - (ii.b) $\circ\varphi \not\vdash \sim\varphi$;
- (iii) $\circ\varphi \vdash \varphi \vee \sim\varphi$, for every φ .

It turns out that, as we will formally show below, the consistency operator \circ of any of the logics $d\text{IRL}_c$, $d\text{IRL}_c^{\text{mB}}$ and $d\text{IRL}_c^{\text{Bm}}$ is a determinedness operator as well, and hence all these explosive logics are LFUs. Moreover, this is also the case for the paraconsistent logics $d\text{IRL}_c^{\leq}$, $(d\text{IRL}_c^{\text{mB}})^{\leq}$ and $(d\text{IRL}_c^{\text{Bm}})^{\leq}$. This is clear for the explosive logics $d\text{IRL}_c$, $d\text{IRL}_c^{\text{mB}}$ and $d\text{IRL}_c^{\text{Bm}}$ due to (i) of

Lemma 5.2, but also for the logics $(\text{dIRL}_c^{\text{mB}})^{\leq}$ and $(\text{dIRL}_c^{\text{Bm}})^{\leq}$ since in these logics $\circ\varphi$ is a Boolean formula. It is not so obvious in the case of the logic dIRL_c^{\leq} , where checking condition (iii) above

$$\circ\varphi \vdash_{\text{dIRL}_c}^{\leq} \varphi \vee \sim\varphi,$$

amounts to check the validity of the inference

$$\vdash_{\text{dIRL}_c} \circ\varphi \rightarrow \varphi \vee \sim\varphi.$$

But this follows from observing that the formula $(\sim\delta \vee \gamma) \rightarrow (\delta \rightarrow \gamma)$ is a theorem in dIRL , and that $\sim\circ\varphi \vee (\varphi \vee \sim\varphi)$ is a theorem of dIRL_c , as it is an equivalent rewriting of the axiom (A1).

In algebraic terms, we can analyse the above observations as follows: if a operator \circ on a dIRL algebra satisfies the basic equation we imposed for a consistency operator, i.e.

$$(\circ 1) \quad x \wedge \sim x \wedge \circ(x) = 0$$

then the following inequality is valid as well:

$$(\circ d) \quad \circ(x) \leq x \vee \sim x.$$

As a matter of fact, we have the following lemma.

LEMMA 5.22

Let \mathbf{A} be a dIRL algebra and let $\circ : A \rightarrow A$ be a unary operation. Then $(\circ 1)$ implies $(\circ d)$.

PROOF. In every residuated lattice $x * y \leq x \wedge y$ and hence, by $(\circ 1)$, one has $\circ(x) * (x \wedge \sim x) \leq \circ(x) \wedge (x \wedge \sim x) = 0$ and hence $\circ(x) \leq \sim(x \wedge \sim x)$. By the involutive property of \sim , then, $\circ(x) \leq \sim(x \wedge \sim x) = x \vee \sim x$. Therefore, it is clear that $(\circ 1)$ implies $(\circ d)$. \square

This means in particular that all the three logics dIRL_c^{\leq} , $(\text{dIRL}_c^{\text{mB}})^{\leq}$ and $(\text{dIRL}_c^{\text{Bm}})^{\leq}$, besides being LFIs, are LFUs as well. In other words, these logics are *strict LFIUs*, as coined in [14]. Therefore, in these logics, the operator \circ is in fact a *classicality* operator in the sense of Omori in his recent paper [40].

On the other hand, the other implication $(\circ d)$ implies $(\circ 1)$ does not hold in general, since $(\circ d)$ is equivalent to $\circ(x) * (x \wedge \sim x) = 0$, which, in general, is strictly weaker than $(\circ 1)$. However, it indeed holds true if we require $\circ(x)$ be Boolean.

LEMMA 5.23

Let \mathbf{A} be a dIRL algebra and let $\circ : A \rightarrow A$ be a unary operation such that the following equation holds:

$$\circ(x) \vee \sim\circ(x) = 1$$

Then $(\circ d)$ implies $(\circ 1)$.

PROOF. It follows by easily checking that the following equations hold in any dIRL algebra: $x \rightarrow (x \wedge y \rightarrow x * y) = 1$ and $\neg x \rightarrow (x \wedge y \rightarrow x * y) = 1$, and hence the equation $(x \vee \neg x) \rightarrow (x \wedge y \rightarrow x * y) = 1$ holds as well. Therefore, if $\circ(x) \vee \sim\circ(x) = 1$, then we have $\circ(x) \wedge y \rightarrow \circ(x) * y = 1$, for every y . In particular, taking $y = x \wedge \sim x$, we get $\circ(x) \wedge (x \wedge \sim x) \rightarrow \circ(x) * (x \wedge \sim x) = 1$, i.e. $\circ(x) \wedge (x \wedge \sim x) \leq \circ(x) * (x \wedge \sim x)$, as desired. \square

Therefore, being LFI turns out to be equivalent to being LFU if we replace $(\circ 1)$ by $(\circ d)$ in the logics $(\text{dIRL}_c^{\text{mB}})^{\leq}$ and $(\text{dIRL}_c^{\text{Bm}})^{\leq}$, but not in $(\text{dIRL}_c)^{\leq}$.

6 Concluding remarks

In this paper we have been concerned with introducing LFIs upon the class of substructural logics having subvarieties of dIRLs as algebraic semantics. To do so, we have first introduced and studied, from an algebraic point of view, dIRLs expanded with three suitable types of consistency operators. Particular attention has been paid to the subvariety of Nelson lattices. Then, the corresponding truth-preserving and degree-preserving logics have been axiomatized, the latter being paraconsistent and falling within the class of LFIs.

At this point there are several observations we deem interesting to discuss. The first one concerns with our request for the involutive lattices to be distributive. In fact, although distributivity is quite often a necessary request (see for instance the proofs of Theorem 3.8 and Theorem 5.9) it is questionable if similar results could be obtained getting rid of that property. Besides being more general, avoiding distributivity might frame the results of this paper in the setting of affine linear logic without exponentials of which (not necessarily distributive) involutive residuated lattices provide an algebraic semantics.

The second one is in fact an open problem. In Section 4.2 we have shown an example of dIRL_c algebra (in fact a NL_c algebra) that is subdirectly irreducible but not simple. It would be interesting to know whether there exists a subdirectly irreducible IMTL_c algebra that is not simple as well.

The third observation concerns the status of the modus ponens rule in the logics dIRL_c^{\leq} , $(\text{dIRL}_c^{\text{mB}})^{\leq}$ and $(\text{dIRL}_c^{\text{Bm}})^{\leq}$. By the very definition of degree-preserving logics, modus ponens is not a valid rule in any of them, only a restricted version where the implication is a theorem holds. However, in the logics $(\text{dIRL}_c^{\text{mB}})^{\leq}$ and $(\text{dIRL}_c^{\text{Bm}})^{\leq}$, due to the PDAT theorems for them (Theorems 5.16 and 5.20 resp.), it is clear that assuming the propositions involved are *consistent* guarantee the validity of the following modus ponens-like inference:

$$\{\varphi, \varphi \rightarrow \psi, \circ\varphi, \circ\psi\} \vdash \psi.$$

The situation in the logic dIRL_c^{\leq} is different, as even this weaker form of modus ponens is not valid. Indeed, take the dIRL_c algebra \mathbf{L} depicted in Figure 2 and an evaluation e on L such that $e(\varphi) = e$ and $e(\psi) = a$, and hence $e(\circ\varphi) = \circ(e) = b$ and $e(\circ\psi) = \circ(a) = b$. A simple computation also shows that $e(\varphi \rightarrow \psi) = c$. Then we have $e(\varphi) \wedge e(\varphi \rightarrow \psi) \wedge e(\circ\varphi) \wedge e(\circ\psi) = e \wedge c \wedge b \wedge b = b \not\leq a = e(\psi)$. Therefore,

$$\{\varphi, \varphi \rightarrow \psi, \circ\varphi, \circ\psi\} \not\vdash_{\text{dIRL}_c^{\leq}} \psi.$$

Nevertheless, such a ‘witnessed form’ of modus ponens holds in dIRL_c^{\leq} if we replace the residual implication $\varphi \rightarrow \psi$ by the material implication $\neg\varphi \vee \psi$. In fact, using axiom (A1), it is not difficult to check that $\varphi \wedge (\neg\varphi \vee \psi) \wedge \circ\varphi \wedge \circ\psi$ is logically equivalent to $\varphi \wedge \psi \wedge \circ\varphi \wedge \circ\psi$, which logically implies ψ . Therefore, the following pattern of inference

$$\{\varphi, \neg\varphi \vee \psi, \circ\varphi, \circ\psi\} \vdash_{\text{dIRL}_c^{\leq}} \psi$$

is indeed valid in dIRL_c^{\leq} .

We finish with two further questions for future research. We have seen that, in the prelinear case, the quasivarieties of $\text{IMTL}_c^{\text{mB}}$ and $\text{IMTL}_c^{\text{Bm}}$ algebras are in fact varieties, since in these algebras the Baaz–Delta connective Δ is definable as $\Delta(x) = x \wedge \circ(x)$. Thus, a first question to be investigated is whether the quasivarieties of dIRL_c , $\text{dIRL}_c^{\text{mB}}$ and $\text{dIRL}_c^{\text{Bm}}$ algebras are actually varieties or, otherwise, proper quasivarieties. A second question is to study an alternative way of defining LFIs related to dIRL logics without relying on the degree-preserving companions. The idea would be to start from the work by Busaniche and Cignoli [8] on the algebraic characterization of

the paraconsistent Nelson logic **N4** in terms of a certain variety of non-integral involutive residuated lattices with a constant, and try to generalize it to the logic *dIRL*, and then adding a suitable consistency operator.

Acknowledgements

The authors are indebted to the anonymous reviewers for their remarks and suggestions that have helped to significantly improve the final layout of this paper. Esteva, Flaminio and Godo acknowledge partial support by the Spanish project PID2019-111544GB-C21. Flaminio also acknowledges partial support by the Spanish Ramón y Cajal research program RYC-2016-19799. Figallo-Orellano acknowledges the support of a post-doctoral grant 2016/21928-0 from São Paulo Research Foundation (FAPESP), Brazil.

References

- [1] A. Almkudad and D. Nelson. Constructible falsity and inexact predicates. *Journal of Symbolic Logic*, **49**, 231–233, 1984.
- [2] A. Avron. Non-deterministic semantics for logics with a consistency operator. *International Journal of Approximate Reasoning*, **45**, 271–287, 2007.
- [3] A. Avron. Modular semantics for some basic logics of formal inconsistency. In *The Many Sides of Logic*, W. A. Carnielli, M. E. Coniglio and I. M. L. D. Ottaviano., eds, pp. 15–26. Studies in Logic. vol. **21**. College Publications, 2009.
- [4] M. Baaz. Infinite-valued Gödel logics with 0-1-projections and relativizations. In *GÖDEL 96, LNL 6*, P. Hájek., ed, pp. 23–33. Springer, 1996.
- [5] P. Baldi, A. Ciabattoni and F. Gulisano. Standard completeness for extensions of IMTL. In *Proceedings of the 2017 IEEE International Conference on Fuzzy Systems (FUZZ-IEEE)*, pp. 1–6, Napoli, Italy, 2017.
- [6] F. Bou, F. Esteva, J. M. Font, A. Gil, L. Godo, A. Torrents and V. Verdú. Logics preserving degrees of truth from varieties of residuated lattices. *Journal of Logic and Computation*, **19**, 1031–1069, 2009.
- [7] M. Busaniche and R. Cignoli. Constructive logic with strong negation as a substructural logic. *Journal of Logic and Computation*, **20**, 761–793, 2010.
- [8] M. Busaniche and R. Cignoli. Residuated lattices as an algebraic semantics for paraconsistent Nelson’s logic. *Journal of Logic and Computation*, **19**, 1019–1029, 2009.
- [9] M. Busaniche and R. Cignoli. The subvariety of commutative residuated lattices represented by twist-products. *Algebra Universalis*, **71**, 5–22, 2014.
- [10] M. Busaniche, M. A. Marcos and S. Ugolini. Representation by triples of algebras with an MV-retract. *Fuzzy Sets and Systems*, **369**, 82–102, 2019.
- [11] L. Cantú and M. Figallo. On the logic that preserves degrees of truth associated to involutive Stone algebras. *Logic Journal of the IGPL*, **28**, 1000–1020, 2020.
- [12] W. Carnielli and M. E. Coniglio. *Paraconsistent Logic: Consistency, Contradiction and Negation*. Logic, Epistemology, and the Unity of Science, vol. **40**. Springer, 2016.
- [13] W. Carnielli, M. E. Coniglio and J. Marcos. Logics of formal inconsistency. In *Handbook of Philosophical Logic*, D. Gabbay and F. Guenther., eds, vol. **14**, 2nd. edn., pp. 1–93. Springer, 2007.
- [14] W. Carnielli, M. E. Coniglio and A. Rodrigues. Recovery operators, paraconsistency and duality. *Logic Journal of the IGPL*, **28**, 624–656, 2020.

- [15] W. Carnielli and A. Rodrigues. An epistemic approach to paraconsistency: a logic of evidence and truth. *Synthese*, **196**, 3789–3813, 2019.
- [16] P. Cintula and C. Noguera. A general framework for mathematical fuzzy logic. In *Handbook of Mathematical Fuzzy Logic—Volume 1*, vol. **37** of Studies in Logic, Mathematical Logic and Foundations, P. Cintula, P. Hájek and C. Noguera., eds, pp. 103–207, London, 2011.
- [17] R. Cignoli, I. D’Ottaviano and D. Mundici. *Algebraic Foundations of Many-Valued Reasoning*. Trends in Logic—Studia Logica Library, vol. **7**. Kluwer, 2000.
- [18] P. Cintula, F. Esteva, J. Gispert, L. Godo, F. Montagna and C. Noguera. Distinguished algebraic semantics for t-norm based fuzzy logics: methods and algebraic equivalencies. *Annals of Pure and Applied Logic*, **160**, 53–81, 2009.
- [19] M. E. Coniglio, F. Esteva and L. Godo. Logics of formal inconsistency arising from systems of fuzzy logic. *Logic Journal of the IGPL*, **22**, 880–904, 2014.
- [20] N. da Costa. On the theory of inconsistent formal systems. *Notre Dame Journal of Formal Logic*, **15**, 497–510, 1974.
- [21] N. da Costa. *Sistemas Formais Inconsistentes*. Habilitation Thesis, Universidade Federal do Paraná, Curitiba, Brazil, 1963. Republished by Editora UFPR, Curitiba, Brazil, 1993.
- [22] R. Ertola, F. Esteva, T. Flaminio, L. Godo and C. Noguera. Paraconsistency properties in degree-preserving fuzzy logics. *Soft Computing*, **19**, 531–546, 2015.
- [23] F. Esteva and L. Godo. Monoidal t-norm based logic: towards a logic for left-continuous t-norms. *Fuzzy Sets and Systems*, **124**, 271–288, 2001.
- [24] M. M. Fidel. An algebraic study of a propositional system of Nelson. In *Mathematical Logic. Proceedings of the First Brazilian Conference, Lectures in Pure and Applied Mathematics*, A. I. Arruda, N. C. A. da Costa and R. Chuaqui., eds, vol. **39**, pp. 99–117. Marcel Dekker, New York and Basel, 1978.
- [25] M. M. Fidel. An algebraic study of logic with constructive negation. A. I. Arruda, N. C. A. da Costa, and A. M. A. Sette, editors, *Proceedings of the Third Brazilian Conference on Mathematical Logic, Recife, 1979*, Sociedade Brasileira de Logica, Campinas, pp. 119–129. 1980.
- [26] J. M. Font. Taking degrees of truth seriously. *Studia Logica*, **91**, 383–406, 2009.
- [27] J. M. Font, A. Gil, A. Torrens and V. Verdú. On the infinite-valued Łukasiewicz logic that preserves degrees of truth. *Archive for Mathematical Logic*, **45**, 839–868, 2006.
- [28] N. Galatos, P. Jipsen, T. Kowalski and H. Ono. *Residuated Lattices: an Algebraic Glimpse at Substructural Logics*. Studies in Logic and the Foundations of Mathematics, vol. **151**. Elsevier, 2007.
- [29] P. Hájek. *Metamathematics of Fuzzy Logic*. Trends in Logic—Studia Logica Library, vol. **4**. Kluwer, 1998.
- [30] U. Höhle. Commutative, residuated ℓ -monoids. In *Non-Classical Logics and Their Applications to Fuzzy Subsets*, U. Höhle and E. P. Klement., eds, pp. 53–106. Kluwer Academic Publishers, 1995.
- [31] S. Jenei. On the structure of rotation-invariant semigroups. *Archive for Mathematical Logic*, **42**, 489–514, 2003.
- [32] Y. Katoh, T. Kowalski and M. Ueda. Almost minimal varieties related to fuzzy logic. *Reports on Mathematical Logic*, **41**, 173–194, 2006.
- [33] T. Kowalski and H. Ono. *Residuated Lattices: An Algebraic Glimpse at Logic Without Contraction*. Monograph, Japan Advanced Institute of Science and Technology, 2001.
- [34] W. McCune. Prover9 and Mace4. <http://www.cs.unm.edu/mccune/prover9/>, 2005–2010.

- [35] J. Marcos. Nearly every normal modal logic is paranormal. *Logique et Analyse*, **48**, 279–300, 2005.
- [36] A. Monteiro. Sur les algèbres de Heyting simetriques. *Portugaliae Math.*, **39**, 1–237, 1980.
- [37] C. Noguera. *Algebraic Study of Axiomatic Extensions of Triangular Norm Based Fuzzy Logics*. PhD Thesis, University of Barcelona, 2006.
- [38] S. Odintsov. Algebraic semantics for paraconsistent Nelson’s logic. *Journal of Logic and Computation*, **13**(4):453–468, 2003.
- [39] S. Odintsov. On the representation of N4-lattices. *Studia Logica*, **76**, 385–405, 2002.
- [40] H. Omori. From logics of formal inconsistency to logics of formal classicality. *Logic Journal of the IGPL*, **28**, 684–711, 2020.
- [41] H. Ono. Logics without the contraction rule and residuated lattices. *Australasian Journal of Logic*, **8**, 50–81, 2011.
- [42] D. Pigozzi. Finite basis theorems for relatively congruence-distributive quasivarieties. *Transactions of the American Mathematical Society*, **310**, 499–533, 1988.
- [43] H. Rasiowa. N-lattices and constructive logic with strong negation. *Fundamenta Mathematicae*, **46**, 61–80, 1958.
- [44] H. Rasiowa. *An Algebraic Approach to Non-Classical Logics*. Studies in Logic and the Foundations of Mathematics, vol. **78**. North-Holland Publishing Company, Amsterdam and London, and American Elsevier Publishing Company, Inc, New York, 1974.
- [45] H. P. Sankappanavar. Heyting algebras with dual pseudocomplementation. *Pacific Journal of Mathematics*, **117**, 405–415, 1985.
- [46] A. Sendlewski. Nelson algebras through Heyting ones. I. *Studia Logica*, **49**, 105–126, 1990.
- [47] M. Spinks and R. Veroff. Constructive logic with strong negation is a substructural logic over FL_{ew} . I. *Studia Logica*, **88**, 325–348, 2008.
- [48] M. Spinks and R. Veroff. Constructive logic with strong negation is a substructural logic over FL_{ew} . II. *Studia Logica*, **89**, 401–425, 2008.
- [49] C. J. Taylor. *Double Heyting Algebras*. PhD Thesis, La Trobe University, 2017.
- [50] S. Ugolini. *Varieties of Residuated Lattices with an MV-Retract and an Investigation Into State Theory*. PhD Thesis, University of Pisa, 2018.
- [51] D. Vakarelov. Notes on N-lattices and constructive logic with strong negation. *Studia Logica*, **34**, 109–125, 1977.
- [52] R. Wójcicki. *Theory of Logical Calculi: Basic Theory of Consequence Operations*. Synthese Library, vol. **199**. Springer Netherlands, 1988.

Received 20 August 2020