

Preferential and Preferential-discriminative Consequence Relations

Jonathan Ben-Naim
LIF, CNRS
CMI 39, rue Joliot-Curie
F-13453 Marseille Cedex 13, France
jbennaim@lif.univ-mrs.fr

November 23, 2018

Abstract

The present paper follows the line of research which consists in investigating consequence relations that are both paraconsistent and plausible (generally non-monotonic too). More precisely, we lay the focus on preferential consequence relations, i.e. those relations that can be defined by a binary preference relation on states labelled by valuations (such as in Non-monotonic Reasoning). The first purpose of the paper is to provide characterizations for families of them, in a general framework that covers e.g. the ones of the well-known paraconsistent logics J_3 and $FOUR$. The second and main purpose is to provide, again in a general framework, characterizations for families of preferential-discriminative consequence relations. The latter are defined exactly as the plain versions, except that among the conclusions, a formula is rejected if its negation is also present.

Remark 1 A previous version of this paper has been published: *The Journal of Logic and Computation*, 15(3):263-294, 2005. The present version contains improvements about the presentation. Consequently, section ordering, definition labels, proposition labels, etc. are different according to whether the present or the previous version is considered.

1 Introduction

In many situations, an agent is confronted with incomplete and/or inconsistent information and then the classical consequence relation proves to be insufficient. Indeed, in case of inconsistent information, it leads to accept every formula as a conclusion, which amounts to loose the whole information. Therefore, the agent needs another relation leading to non-trivial conclusions in spite of the presence of contradictions. So, several paraconsistent consequence relations have been developed. In the present paper, we will pay attention in particular to certain many-valued ones [Bel77b, Bel77a, DdC70, CMdA00, dACM02, AA94, AA96, AA98]. They are defined in frameworks where valuations can assign more than two different truth values to formulas. In fact, they tolerate contradictions within the conclusions, but reject the principle of explosion according to which a single contradiction entails the deduction of every formula.

In case of incomplete information, the classical consequence relation also shows its limits. Indeed, no risk is taken, the conclusions are sure, but too few. The agent often needs another relation, more daring, leading to accept as conclusions formulas that are not necessarily sure, but still plausible. Eventually, some “hasty” conclusions will be rejected later, in the presence of additional information. So, a lot of plausible (generally, non-monotonic) consequence relations have been developed. Gabbay, Makinson, Kraus, Lehmann, and Magidor investigated extensively properties that should be satisfied by such relations [Gab85, Mak89, Mak94, KLM90, LM92]. In addition, central tools to define plausible relations are *choice functions* [Che54, Arr59, Sen70, AM81, Leh02, Leh01, Sch92, Sch04]. Indeed, suppose we have at our disposal a function μ , called a choice function, which chooses in any set of valuations V , those elements that are preferred, not necessarily in the absolute sense, but when the valuations in V are the only ones under consideration. Then, it is natural to conclude α (a formula) from Γ (a set of formulas) iff every model for Γ chosen by μ is a model for α . This constitutes a plausible (generally, non-monotonic) consequence relation.

In the present paper, we will lay the focus on a particular family of choice functions. Let us present it. Suppose we are given a binary preference relation \prec on states labelled by valuations (in the style of e.g. [KLM90, Sch04]). This defines naturally a choice function. Indeed, choose in any set of valuations V , each element which labels a state which is \prec -preferred among all the states labelled by an element of V . Those choice functions which can be defined in this manner constitute the aforementioned family. The consequence relations defined by this family will be called *preferential consequence relations*.

For a long time, research efforts on paraconsistent relations and plausible relations were separated. However, in many applications, the information is both incomplete and inconsistent. For instance, the semantic web or big databases inevitably contain inconsistencies. This can be due to human or material imperfections as well as contradictory sources of information. On the other hand, neither the web nor big databases can contain “all” information. Indeed, there are rules of which the exceptions cannot be enumerated. Also, some information might be left voluntarily vague or in concise form. Consequently, consequence relations that are both paraconsistent and plausible are useful to reason in such applications.

Such relations first appear in e.g. [Pri91, Bat98, KL92, AA00, KM02]. The idea begins by taking a many-valued framework to get paraconsistency. Then, only those models that are most preferred according to some particular binary preference relation on valuations (in the style of [Sho88, Sho87]) are relevant for making inference, which provides plausibility (and in fact also non-monotonicity). In [AL01b, AL01a], A. Avron and I. Lev generalized the study to families of binary preference relations which compare two valuations using, for each of them, this part of a certain set of formulas it satisfies. The present paper follows this line of research by combining many-valued frameworks

and choice functions.

More explicitly, we will investigate preferential consequence relations in a general framework. According to the different assumptions which will be made about the latter, it will cover various kinds of frameworks, including e.g. the classical propositional one as well as some many-valued ones. Moreover, in the many-valued frameworks, preferential relations lead to rational and non-trivial conclusions in spite of the presence of contradictions and are thus useful to deal with both incomplete and inconsistent information. However, they will not satisfy the Disjunctive Syllogism (from α and $\neg\alpha \vee \beta$ we can conclude β), whilst they satisfy it in classical frameworks.

In addition, it is in the many-valued frameworks that new relations, which we will investigate in detail, are really interesting: *preferential-discriminative consequence relations*. They are defined exactly as the plain versions, except that among the conclusions, a formula is rejected if its negation is also present. In classical frameworks, they do not bring something really new. Indeed, instead of concluding everything in the face of inconsistent information, we will simply conclude nothing. On the other hand, in many-valued frameworks, where the conclusions are rational even from inconsistent information, the discriminative versions will reject the contradictions among them, rendering them all the more rational.

The contribution of the present paper can now be summarized in one sentence: we characterized, in a general framework, several (sub)families of preferential(-discriminative) consequence relations. In many cases, our characterizations are purely syntactic. This has a lot of advantages, let us quote some important ones. Take some syntactic conditions that characterize a family of those consequence relations. This gives a syntactic point of view on this family defined semantically, which enables us to compare it to conditions known on the “market”, and thus to other consequence relations. This can also give rise to questions like: if we modified the conditions in such and such a natural-looking way, what would happen on the semantic side? More generally, this can open the door to questions that would not easily come to mind otherwise or to techniques of proof that could not have been employed in the semantic approach. Finally, this can help to find or improve proof systems based on the family, like a Gentzen proof system for instance.

Several characterizations can be found in the literature for preferential relations (e.g. [Gab85, Mak89, Mak94, KLM90, LM92, Leh02, Leh01, Sch92, Sch96, Sch00, Sch04]). We will provide some new ones, though, to do so, we have been inspired by techniques of K. Schlechta [Sch04]. In fact, our innovation is rather related to the discriminative version. To the author knowledge, the present paper is the first systematic work of characterization for preferential-discriminative consequence relations.

The rest of the paper is organized as follows. In Section 2.1, we introduce our general framework and the different assumptions which sometimes will be made about it. We will see that it covers in particular the many-valued frameworks of the well-known paraconsistent logics $\mathcal{F}\mathcal{O}\mathcal{U}\mathcal{R}$ and J_3 . In Section 2.2, we present choice functions and some of their well-known properties. We will see which properties characterize those choice functions that can be defined by a binary preference relation on states labelled by valuations. In Section 2.3, we define preferential(-discriminative) consequence relations and give examples in both the classical and the many-valued frameworks. We will also recall a characterization which involves the well-known system \mathbf{P} of Kraus, Lehmann, and Magidor. In section 3, we provide our characterizations. Finally, we conclude in Section 4.

2 Background

2.1 Semantic structures

2.1.1 Definitions and properties

We will work with general formulas, valuations, and satisfaction. A similar approach has been taken in two well-known papers [Mak05, Leh01].

Definition 2 We say that \mathcal{S} is a *semantic structure* iff $\mathcal{S} = \langle \mathcal{F}, \mathcal{V}, \models \rangle$ where \mathcal{F} is a set, \mathcal{V} is a set, and \models is a relation on $\mathcal{V} \times \mathcal{F}$.

Intuitively, \mathcal{F} is a set of formulas, \mathcal{V} a set of valuations for these formulas, and \models a satisfaction relation for these objects (i.e. $v \models \alpha$ means the formula α is satisfied in the valuation v , i.e. v is a model for α).

Notation 3 Let $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ be a semantic structure, $\Gamma \subseteq \mathcal{F}$, and $V \subseteq \mathcal{V}$. Then,

$$M_\Gamma := \{v \in \mathcal{V} : \forall \alpha \in \Gamma, v \models \alpha\},$$

$$T(V) := \{\alpha \in \mathcal{F} : V \subseteq M_\alpha\},$$

$$\mathbf{D} := \{V \subseteq \mathcal{V} : \exists \Gamma \subseteq \mathcal{F}, M_\Gamma = V\}.$$

Suppose \mathcal{L} is a language, \neg a unary connective of \mathcal{L} , and \mathcal{F} the set of all wffs of \mathcal{L} . Then,

$$T_d(V) := \{\alpha \in \mathcal{F} : V \subseteq M_\alpha \text{ and } V \not\subseteq M_{\neg\alpha}\},$$

$$T_c(V) := \{\alpha \in \mathcal{F} : V \subseteq M_\alpha \text{ and } V \subseteq M_{\neg\alpha}\},$$

$$\mathbf{C} := \{V \subseteq \mathcal{V} : \forall \alpha \in \mathcal{F}, V \not\subseteq M_\alpha \text{ or } V \not\subseteq M_{\neg\alpha}\}.$$

Intuitively, M_Γ is the set of all models for Γ and $T(V)$ the set of all formulas satisfied in V . Every element of $T(V)$ belongs either to $T_d(V)$ or $T_c(V)$, according to whether its negation is also in $T(V)$. \mathbf{D} is the set of all those sets of valuations that are definable by a set of formulas and \mathbf{C} the set of all those sets of valuations that do not satisfy both a formula and its negation. As usual, $M_{\Gamma, \alpha}$, $T(V, v)$ stand for respectively $M_{\Gamma \cup \{\alpha\}}$, $T(V \cup \{v\})$, etc.

Remark 4 The notations M_Γ , $T(V)$, etc. should contain the semantic structure on which they are based. To increase readability, we will omit it. There will never be any ambiguity. We will omit similar things with other notations in the sequel, for the same reason.

A semantic structure defines a basic consequence relation:

Notation 5 We denote by \mathcal{P} the power set operator.

Let $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ be a semantic structure.

We denote by \vdash the relation on $\mathcal{P}(\mathcal{F}) \times \mathcal{F}$ such that $\forall \Gamma \subseteq \mathcal{F}, \forall \alpha \in \mathcal{F}$,

$$\Gamma \vdash \alpha \text{ iff } M_\Gamma \subseteq M_\alpha.$$

Let \sim be a relation on $\mathcal{P}(\mathcal{F}) \times \mathcal{F}$. Then,

$$\sim(\Gamma) := \{\alpha \in \mathcal{F} : \Gamma \sim \alpha\}.$$

Suppose \mathcal{L} is a language, \neg a unary connective of \mathcal{L} , \mathcal{F} the set of all wffs of \mathcal{L} , and $\Gamma \subseteq \mathcal{F}$.

Then, we say that Γ is *consistent* iff $\forall \alpha \in \mathcal{F}, \Gamma \not\vdash \alpha$ or $\Gamma \not\vdash \neg\alpha$.

The following trivial facts hold, we will use them implicitly in the sequel:

Remark 6 Let $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ be a semantic structure and $\Gamma, \Delta \subseteq \mathcal{F}$. Then:

$$M_{\Gamma, \Delta} = M_{\Gamma} \cap M_{\Delta};$$

$$\vdash(\Gamma) = T(M_{\Gamma});$$

$$M_{\Gamma} = M_{\vdash(\Gamma)};$$

$$\Gamma \subseteq \vdash(\Delta) \text{ iff } \vdash(\Gamma) \subseteq \vdash(\Delta) \text{ iff } M_{\Delta} \subseteq M_{\Gamma}.$$

Sometimes, we will need some of the following assumptions about a semantic structure:

Definition 7 Suppose $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ is a semantic structure.

Then, define the following assumptions about it:

(A1) \mathcal{V} is finite.

Suppose \mathcal{L} is a language, \neg a unary connective of \mathcal{L} , and \mathcal{F} the set of all wffs of \mathcal{L} . Then, define:

(A2) $\forall \Gamma \subseteq \mathcal{F}, \forall \alpha \in \mathcal{F}$, if $\alpha \notin T(M_{\Gamma})$ and $\neg\alpha \notin T(M_{\Gamma})$, then $M_{\Gamma} \cap M_{\alpha} \not\subseteq M_{\neg\alpha}$.

Suppose \vee and \wedge are binary connectives of \mathcal{L} . Then, define:

(A3) $\forall \alpha, \beta \in \mathcal{F}$, we have:

$$M_{\alpha \vee \beta} = M_{\alpha} \cup M_{\beta};$$

$$M_{\alpha \wedge \beta} = M_{\alpha} \cap M_{\beta};$$

$$M_{\neg\neg\alpha} = M_{\alpha};$$

$$M_{\neg(\alpha \vee \beta)} = M_{\neg\alpha \wedge \neg\beta};$$

$$M_{\neg(\alpha \wedge \beta)} = M_{\neg\alpha \vee \neg\beta}.$$

Clearly, those assumptions are satisfied by classical semantic structures, i.e. structures where \mathcal{F} , \mathcal{V} , and \models are classical. In addition, we will see, in Sections 2.1.2 and 2.1.3, that they are satisfied also by certain many-valued semantic structures.

2.1.2 The semantic structure defined by *FOUR*

The logic *FOUR* was introduced by N. Belnap in [Bel77a, Bel77b]. This logic is useful to deal with inconsistent information. Several presentations are possible, depending on the language under consideration. For the needs of the present paper, a classical propositional language will be sufficient. The logic has been investigated intensively in e.g. [AA94, AA96, AA98], where richer languages, containing an implication connective \supset (first introduced by A. Avron [Avr91]), were considered.

Notation 8 We denote by \mathcal{A} a set of propositional symbols (or atoms).

We denote by \mathcal{L}_c the classical propositional language containing \mathcal{A} , the usual constants *false* and *true*, and the usual connectives \neg , \vee , and \wedge .

We denote by \mathcal{F}_c the set of all wffs of \mathcal{L}_c .

We recall a possible meaning for the logic *FOUR* (more details can be found in [CLM99, Bel77a, Bel77b]). Consider a system in which there are, on the one hand, sources of information and, on the other hand, a processor that listens to them. The sources provide information about the atoms only, not about the compound formulas. For each atom p , there are exactly four possibilities: either the processor is informed (by the sources, taken as a whole) that p is true; or he is informed that p is false; or he is informed of both; or he has no information about p .

Notation 9 Denote by 0 and 1 the classical truth values and define:

$$\mathbf{f} := \{0\}; \quad \mathbf{t} := \{1\}; \quad \top := \{0, 1\}; \quad \perp := \emptyset.$$

The global information given by the sources to the processor can be modelled by a function s from \mathcal{A} to $\{\mathbf{f}, \mathbf{t}, \top, \perp\}$. Intuitively, $1 \in s(p)$ means the processor is informed that p is true, whilst $0 \in s(p)$ means he is informed that p is false.

Then, the processor naturally builds information about the compound formulas from s . Before he starts to do so, the situation can be modelled by a function v from \mathcal{F}_c to $\{\mathbf{f}, \mathbf{t}, \top, \perp\}$ which agrees with s about the atoms and which assigns \perp to all compound formulas. Now, take p and q in \mathcal{A} and suppose $1 \in v(p)$ or $1 \in v(q)$. Then, the processor naturally adds 1 to $v(p \vee q)$. Similarly, if $0 \in v(p)$ and $0 \in v(q)$, then he adds 0 in $v(p \vee q)$. Of course, such rules hold for \neg and \wedge too.

Suppose all those rules are applied recursively to all compound formulas. Then, v represents the “full” (or developed) information given by the sources to the processor. Now, the valuations of the logic *FOUR* can be defined as exactly those functions that can be built in this manner (i.e. like v) from some of these sources-processor systems. More formally,

Definition 10 We say that v is a *four-valued valuation* iff v is a function from \mathcal{F}_c to $\{\mathbf{f}, \mathbf{t}, \top, \perp\}$ such that $v(\text{true}) = \mathbf{t}$, $v(\text{false}) = \mathbf{f}$ and $\forall \alpha, \beta \in \mathcal{F}_c$,

- $1 \in v(\neg\alpha)$ iff $0 \in v(\alpha)$;
- $0 \in v(\neg\alpha)$ iff $1 \in v(\alpha)$;
- $1 \in v(\alpha \vee \beta)$ iff $1 \in v(\alpha)$ or $1 \in v(\beta)$;
- $0 \in v(\alpha \vee \beta)$ iff $0 \in v(\alpha)$ and $0 \in v(\beta)$;
- $1 \in v(\alpha \wedge \beta)$ iff $1 \in v(\alpha)$ and $1 \in v(\beta)$;
- $0 \in v(\alpha \wedge \beta)$ iff $0 \in v(\alpha)$ or $0 \in v(\beta)$.

We denote by \mathcal{V}_4 the set of all four-valued valuations.

The definition may become more accessible if we see the four-valued valuations as those functions that satisfy Tables 1, 2, and 3 below:

$v(\alpha)$	$v(\neg\alpha)$
\mathbf{f}	\mathbf{t}
\mathbf{t}	\mathbf{f}
\top	\top
\perp	\perp

Table 1.

$v(\alpha)$	$v(\beta)$			
\mathbf{f}	\mathbf{f}	\mathbf{t}	\top	\perp
\mathbf{t}	\mathbf{t}	\mathbf{t}	\top	\mathbf{t}
\top	\top	\mathbf{t}	\top	\mathbf{t}
\perp	\perp	\mathbf{t}	\mathbf{t}	\perp

Table 2.

$v(\alpha)$	$v(\beta)$			
\mathbf{f}	\mathbf{f}	\mathbf{f}	\mathbf{f}	\mathbf{f}
\mathbf{t}	\mathbf{f}	\mathbf{t}	\top	\perp
\top	\mathbf{f}	\top	\top	\mathbf{f}
\perp	\mathbf{f}	\perp	\mathbf{f}	\perp

Table 3.

In the logic *FOUR*, a formula α is considered to be satisfied iff the processor is informed that it is true (it does not matter whether he is also informed that α is false).

Notation 11 We denote by \models_4 the relation on $\mathcal{V}_4 \times \mathcal{F}_c$ such that $\forall v \in \mathcal{V}_4, \forall \alpha \in \mathcal{F}_c$, we have $v \models_4 \alpha$ iff $1 \in v(\alpha)$.

When the *FOUR* semantic structure $\langle \mathcal{F}_c, \mathcal{V}_4, \models_4 \rangle$ is under consideration, proof systems for \vdash are available. For instance, A. Avron and O. Arieli provided several ones in e.g. [AA94, AA96, AA98].

Note that the *FOUR* semantic structure satisfies (A3). In addition, if \mathcal{A} is finite, then (A1) is also satisfied. However, (A2) is not satisfied by this structure. In Section 2.1.3, we turn to a many-valued semantic structure which satisfies (A2).

2.1.3 The semantic structure defined by J_3

The logic J_3 was introduced in [DdC70] to answer a question posed in 1948 by S. Jaśkowski, who was interested in systematizing theories capable of containing contradictions, especially if they occur in dialectical reasoning. The step from informal reasoning under contradictions and formal reasoning with databases and information was done in [CMdA00] (also specialized for real database models in [dACM02]), where another formulation of J_3 called **LFII** was introduced, and its first-order version, semantics and proof theory were studied in detail. Investigations of J_3 have also been made in e.g. [Avr91], where richer languages than our \mathcal{L}_c were considered.

The valuations of the logic J_3 can be given the same meaning as those of the logic \mathcal{FOUR} , except that the consideration is restricted to those systems where the sources, taken as a whole, always give some information about an atom. More formally,

Definition 12 We say that v is a *three-valued valuation* iff v is a function from \mathcal{F}_c to $\{\mathbf{f}, \mathbf{t}, \top\}$ such that $v(\text{true}) = \mathbf{t}$, $v(\text{false}) = \mathbf{f}$ and $\forall \alpha, \beta \in \mathcal{F}_c$,

- $1 \in v(\neg\alpha)$ iff $0 \in v(\alpha)$;
- $0 \in v(\neg\alpha)$ iff $1 \in v(\alpha)$;
- $1 \in v(\alpha \vee \beta)$ iff $1 \in v(\alpha)$ or $1 \in v(\beta)$;
- $0 \in v(\alpha \vee \beta)$ iff $0 \in v(\alpha)$ and $0 \in v(\beta)$;
- $1 \in v(\alpha \wedge \beta)$ iff $1 \in v(\alpha)$ and $1 \in v(\beta)$;
- $0 \in v(\alpha \wedge \beta)$ iff $0 \in v(\alpha)$ or $0 \in v(\beta)$.

We denote by \mathcal{V}_3 the set of all three-valued valuations.

As previously, the definition may become more accessible if we see the three-valued valuations as those functions that satisfy Tables 4, 5, and 6 below:

$v(\alpha)$	$v(\neg\alpha)$
\mathbf{f}	\mathbf{t}
\mathbf{t}	\mathbf{f}
\top	\top

Table 4.

$v(\alpha)$	$v(\beta)$		
\mathbf{f}	\mathbf{f}	\mathbf{t}	\top
\mathbf{t}	\mathbf{t}	\mathbf{t}	\mathbf{t}
\top	\top	\mathbf{t}	\top

Table 5.

$v(\alpha)$	$v(\beta)$		
\mathbf{f}	\mathbf{f}	\mathbf{t}	\top
\mathbf{t}	\mathbf{f}	\mathbf{t}	\top
\top	\mathbf{f}	\top	\top

Table 6.

We turn to the satisfaction relation.

Notation 13 We denote by \models_3 the relation on $\mathcal{V}_3 \times \mathcal{F}_c$ such that $\forall v \in \mathcal{V}_3, \forall \alpha \in \mathcal{F}_c$, we have $v \models_3 \alpha$ iff $1 \in v(\alpha)$.

When the J_3 semantic structure $\langle \mathcal{F}_c, \mathcal{V}_3, \models_3 \rangle$ is considered, proof systems for \vdash are available. Some have been provided in e.g. [Avr91, DdC70] and in chapter IX of [Eps90].

The J_3 structure satisfies (A3) and (A2). In addition, if \mathcal{A} is finite, then it satisfies (A1) too.

2.2 Choice functions

2.2.1 Definitions and properties

In many situations, an agent has some way to choose in any set of valuations V , those elements that are preferred (the bests, the more normal, etc.), not necessarily in the absolute sense, but when the valuations in V are the only ones under consideration. In Social Choice, this is modelled by choice functions [Che54, Arr59, Sen70, AM81, Leh02, Leh01].

Definition 14 Let \mathcal{V} be a set, $\mathbf{V} \subseteq \mathcal{P}(\mathcal{V})$, $\mathbf{W} \subseteq \mathcal{P}(\mathcal{V})$, and μ a function from \mathbf{V} to \mathbf{W} . We say that μ is a *choice function* iff $\forall V \in \mathbf{V}, \mu(V) \subseteq V$.

Several properties for choice functions have been put in evidence by researchers in Social Choice. Let us present two important ones (a better presentation can be found in [Leh01]). Suppose W is a set of valuations, V is a subset of W , and $v \in V$ is a preferred valuation of W . Then, a natural requirement is that v is a preferred valuation of V . Indeed, in many situations, the larger a set is, the harder it is to be a preferred element of it, and he who can do the most can do the least. This property appears in [Che54] and has been given the name Coherence in [Mou85].

We turn to the second property. Suppose W is a set of valuations, V is a subset of W , and suppose all the preferred valuations of W belong to V . Then, they are expected to include all the preferred valuations of V . The importance of this property has been put in evidence by [Aiz85, AM81] and has been given the name Local Monotonicity in e.g. [Leh01].

Definition 15 Let \mathcal{V} be a set, $\mathbf{V} \subseteq \mathcal{P}(\mathcal{V})$, $\mathbf{W} \subseteq \mathcal{P}(\mathcal{V})$, and μ a choice function from \mathbf{V} to \mathbf{W} . We say that μ is *coherent* iff $\forall V, W \in \mathbf{V}$,

$$\text{if } V \subseteq W, \text{ then } \mu(W) \cap V \subseteq \mu(V).$$

We say that μ is *locally monotonic* (LM) iff $\forall V, W \in \mathbf{V}$,

$$\text{if } \mu(W) \subseteq V \subseteq W, \text{ then } \mu(V) \subseteq \mu(W).$$

In addition to their intuitive meanings, these properties are important because, as was shown by K. Schlechta in [Sch00], they characterize those choice functions that can be defined by a binary preference relation on states labelled by valuations (in the style of e.g. [KLM90]). We will take a closer look at this in Section 2.2.2.

When a semantic structure is under consideration, two new properties can be defined. Each of them conveys a simple and natural meaning.

Definition 16 Let $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ be a semantic structure, $\mathbf{V} \subseteq \mathcal{P}(\mathcal{V})$, $\mathbf{W} \subseteq \mathcal{P}(\mathcal{V})$, and μ a choice function from \mathbf{V} to \mathbf{W} .

We say that μ is *definability preserving* (DP) iff

$$\forall V \in \mathbf{V} \cap \mathbf{D}, \mu(V) \in \mathbf{D}.$$

Suppose \mathcal{L} is a language, \neg a unary connective of \mathcal{L} , and \mathcal{F} the set of all wffs of \mathcal{L} .

We say that μ is *coherency preserving* (CP) iff

$$\forall V \in \mathbf{V} \cap \mathbf{C}, \mu(V) \in \mathbf{C}.$$

Definability Preservation has been put in evidence first in [Sch92]. One of its advantages is that when the choice functions under consideration satisfy it, we will provide characterizations with purely syntactic conditions. To the author knowledge, the present paper is the first to introduce Coherency Preservation. An advantage of this property is that when the choice functions under consideration satisfy it, we will not need to assume (A2) to show our characterizations (in the discriminative case).

2.2.2 Preference structures

Binary preference relations on valuations have been investigated by e.g. B. Hansson to give semantics for deontic logics [Han69]. Y. Shoham rediscovered them to give semantics for plausible non-monotonic logics [Sho88, Sho87]. Then, it seems that Imielinski is one of the first persons to introduce binary preference relations on states labelled by valuations [Imi87]. They have been used to give more general semantics for plausible non-monotonic logics, see e.g. [KLM90, LM92, Sch92, Sch96, Sch00, Sch04]. Let us present them.

Definition 17 We say that \mathcal{R} is a preference structure on a set \mathcal{V} iff $\mathcal{R} = \langle \mathcal{S}, l, \prec \rangle$ where \mathcal{S} is a set, l is a function from \mathcal{S} to \mathcal{V} , and \prec is a relation on $\mathcal{S} \times \mathcal{S}$.

In fact, preference structures are essentially Kripke structures. The difference lies in the interpretation of \prec . In a Kripke structure, it is seen as an accessibility relation, whilst, in a preference structure, it is seen as a preference relation. We recall a possible meaning for preference structures (see e.g. [KLM90, Sch04] for details about meaning). Intuitively, \mathcal{V} is a set of valuations for some language \mathcal{L} and \mathcal{S} a set of valuations for some language \mathcal{L}' richer than \mathcal{L} . The elements of \mathcal{S} are called states. $l(s)$ corresponds precisely to this part of s that is about the formulas of \mathcal{L} only. We call l a labelling function. Finally, \prec is a preference relation, i.e. $s \prec s'$ means s is preferred to s' .

We turn to well-known properties for preference structures.

Definition 18 Suppose \mathcal{V} is a set, $\mathcal{R} = \langle \mathcal{S}, l, \prec \rangle$ is a preference structure on \mathcal{V} , $S \subseteq \mathcal{S}$, $s \in S$, $V \subseteq \mathcal{V}$, and $\mathbf{V} \subseteq \mathcal{P}(\mathcal{V})$.

We say that \mathcal{R} is *transitive* (resp. *irreflexive*) iff \prec is transitive (resp. irreflexive).

We say that s is *preferred* in S iff $\forall s' \in S, s' \not\prec s$.

$L(V) := \{s \in \mathcal{S} : l(s) \in V\}$ (intuitively, $L(V)$ contains the states labelled by the elements of V).

We say that \mathcal{R} is *\mathbf{V} -smooth* (alias *\mathbf{V} -stoppered*) iff $\forall V \in \mathbf{V}, \forall s \in L(V)$,

either s is preferred in $L(V)$ or there exists s' preferred in $L(V)$ such that $s' \prec s$.

A preference structure defines naturally a choice function. The idea is to choose in any set of valuations V , each element which labels a state which is preferred among all the states labelled by the elements of V .

Definition 19 Suppose $\mathcal{R} = \langle \mathcal{S}, l, \prec \rangle$ is a preference structure on a set \mathcal{V} .

We denote by $\mu_{\mathcal{R}}$ the function from $\mathcal{P}(\mathcal{V})$ to $\mathcal{P}(\mathcal{V})$ such that $\forall V \subseteq \mathcal{V}$,

$$\mu_{\mathcal{R}}(V) = \{v \in V : \exists s \in L(v), s \text{ is preferred in } L(V)\}.$$

In [Sch00], Schlechta showed that Coherence and Local Monotonicity characterize those choice functions that can be defined by a preference structure. Details are given in the proposition just below. It is an immediate corollary of Proposition 2.4, Proposition 2.15, and Fact 1.3 of [Sch00].

Proposition 20 Taken from [Sch00].

Let \mathcal{V} be a set, \mathbf{V} and \mathbf{W} subsets of $\mathcal{P}(\mathcal{V})$, and μ a choice function from \mathbf{V} to \mathbf{W} . Then,

- (0) μ is coherent iff there exists a transitive and irreflexive preference structure \mathcal{R} on \mathcal{V} such that $\forall V \in \mathbf{V}$, we have $\mu(V) = \mu_{\mathcal{R}}(V)$.

Suppose $\forall V, W \in \mathbf{V}$, we have $V \cup W \in \mathbf{V}$ and $V \cap W \in \mathbf{V}$. Then,

- (1) μ is coherent and LM iff there exists a \mathbf{V} -smooth, transitive, and irreflexive preference structure \mathcal{R} on \mathcal{V} such that $\forall V \in \mathbf{V}$, we have $\mu(V) = \mu_{\mathcal{R}}(V)$.

In fact, in [Sch00], the codomain of μ is required to be its domain: \mathbf{V} . However, this plays no role in the proofs. Therefore, verbatim the same proofs are valid when the codomain of μ is an arbitrary subset \mathbf{W} of $\mathcal{P}(\mathcal{V})$. Both myself and Schlechta checked it.

2.3 Preferential(-discriminative) consequence relations

2.3.1 Definitions

Suppose we are given a semantic structure and a choice function μ on the valuations. Then, it is natural to conclude a formula α from a set of formulas Γ iff every model for Γ chosen by μ is a model for α . More formally:

Definition 21 Suppose $\mathcal{S} = \langle \mathcal{F}, \mathcal{V}, \models \rangle$ is a semantic structure and \sim a relation on $\mathcal{P}(\mathcal{F}) \times \mathcal{F}$. We say that \sim is a *preferential consequence relation* iff there exists a coherent choice function μ from \mathbf{D} to $\mathcal{P}(\mathcal{V})$ such that $\forall \Gamma \subseteq \mathcal{F}, \forall \alpha \in \mathcal{F}$,

$$\Gamma \sim \alpha \text{ iff } \mu(M_{\Gamma}) \subseteq M_{\alpha}.$$

In addition, if μ is LM, DP, etc., then so is \sim .

These consequence relations are called “preferential” because, in the light of Proposition 20, they can be defined equivalently with preference structures, instead of coherent choice functions. They lead to “jump” to plausible conclusions which will eventually be withdrawn later, in the presence of additional information. Therefore, they are useful to deal with incomplete information. We will give an example with a classical semantic structure in Section 2.4.1.

In addition, if a many-valued semantic structure is considered, they lead to rational and non-trivial conclusions in spite of the presence of contradictions and are thus useful to treat both incomplete and inconsistent information. However, they will not satisfy the Disjunctive Syllogism. We will give an example with the *FOUR* semantic structure in Section 2.4.2.

Now, we turn to a qualified version of preferential consequence. It captures the idea that the contradictions in the conclusions should be rejected.

Definition 22 Suppose \mathcal{L} is a language, \neg a unary connective of \mathcal{L} , \mathcal{F} the set of all wffs of \mathcal{L} , $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ a semantic structure, and \sim a relation on $\mathcal{P}(\mathcal{F}) \times \mathcal{F}$.

We say that \sim is a *preferential-discriminative consequence relation* iff there is a coherent choice function μ from \mathbf{D} to $\mathcal{P}(\mathcal{V})$ such that $\forall \Gamma \subseteq \mathcal{F}, \forall \alpha \in \mathcal{F}$,

$$\Gamma \sim \alpha \text{ iff } \mu(M_{\Gamma}) \subseteq M_{\alpha} \text{ and } \mu(M_{\Gamma}) \not\subseteq M_{\neg\alpha}.$$

In addition, if μ is LM, DP, etc., then so is \sim .

If a classical semantic structure is considered, the discriminative version does not bring something really new. Indeed, the only difference will be to conclude nothing instead of everything in the face of inconsistent information. On the other hand, with a many-valued structure, the conclusions are rational even from inconsistent information. The discriminative version will then reject the contradictions in the conclusions, rendering the latter all the more rational.

In Definitions 21 and 22, the domain of the choice function is \mathbf{D} . This is natural as only the elements of \mathbf{D} play a role in the definition of a preferential(-discriminative) consequence relation. This point of view has been adopted in e.g. [Leh01] (see Section 6). Now, one might want a definition with choice functions of which the domain is $\mathcal{P}(\mathcal{V})$. In fact, some families of relations can be defined equivalently with \mathbf{D} or $\mathcal{P}(\mathcal{V})$. For instance, as is noted in [Leh01], if μ is a coherent choice function from \mathbf{D} to $\mathcal{P}(\mathcal{V})$, then the function μ' from $\mathcal{P}(\mathcal{V})$ to $\mathcal{P}(\mathcal{V})$ defined by $\mu'(V) = V \cap \mu(M_{T(V)})$ is a coherent choice function which agrees with μ on \mathbf{D} .

Several characterizations for preferential consequence relations can be found in the literature (e.g. [KLM90, LM92, Leh02, Leh01, Sch92, Sch96, Sch00, Sch04]). In particular, we will recall (in Section 2.4) a characterization that involves the well-known system \mathbf{P} of [KLM90].

As said previously, in the light of Proposition 20, preferential(-discriminative) consequence relations could have been introduced equivalently with preference structures. We opted for coherent choice functions for two reasons. First, they give a clearer meaning. Indeed, properties like Coherence have simple intuitive justifications, whilst preference structures contain “states”, but it is not perfectly clear what a state is in daily life. By the way, in [KLM90], Kraus, Lehmann, and Magidor did not consider preference structures to be ontological justifications for their interest in the formal systems investigated, but to be technical tools to study those systems and in particular settle questions of interderivability and find efficient decision procedures (see the end of Section 1.2 of [KLM90]).

Second, in the proofs, we will work directly with choice functions and their properties, not with preference structures. By the way, the techniques developed in the present paper (especially in the discriminative case) can certainly be adapted to new properties.

2.4 The system \mathbf{P}

Gabbay, Makinson, Kraus, Lehmann, and Magidor investigated extensively properties which should be satisfied by plausible non-monotonic consequence relations [Gab85, Mak89, Mak94, KLM90, LM92]. A certain set of properties, called the system \mathbf{P} , plays a central role in this area. It is essentially due to Kraus, Lehmann, and Magidor [KLM90] and has been investigated further in [LM92]. Let's present it.

Definition 23 Suppose \mathcal{L} is a language containing the usual connectives \neg and \vee , \mathcal{F} the set of all wffs of \mathcal{L} , $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ a semantic structure, and \vdash a relation on $\mathcal{F} \times \mathcal{F}$.

Then, the system \mathbf{P} is the set of the six following conditions: $\forall \alpha, \beta, \gamma \in \mathcal{F}$,

Reflexivity $\alpha \vdash \alpha$

Left Logical Equivalence $\frac{\vdash \alpha \leftrightarrow \beta \quad \alpha \vdash \gamma}{\beta \vdash \gamma}$

Right Weakening $\frac{\vdash \alpha \rightarrow \beta \quad \gamma \vdash \alpha}{\gamma \vdash \beta}$

Cut $\frac{\alpha \wedge \beta \vdash \gamma \quad \alpha \vdash \beta}{\alpha \vdash \gamma}$

Cautious Monotonicity $\frac{\alpha \vdash \beta \quad \alpha \vdash \gamma}{\alpha \wedge \beta \vdash \gamma}$

$$\text{Or } \frac{\alpha \sim \gamma \quad \beta \sim \gamma}{\alpha \vee \beta \sim \gamma}$$

Note that $\alpha \wedge \beta$ is a shorthand for $\neg(\neg\alpha \vee \neg\beta)$. Similarly, $\alpha \rightarrow \beta$ and $\alpha \leftrightarrow \beta$ are shorthands. Note again that **P** without **Or** is called **C**. The system **C** is closely related to the cumulative inference which was investigated by Makinson in [Mak89]. In addition, it seems to correspond to what Gabbay proposed in [Gab85]. Concerning the rule **Or**, it corresponds to the axiom CA of conditional logic.

All the properties in **P** are sound if we read $\alpha \sim \beta$ as “ β is a plausible consequence of α ”. In addition, **P** is complete in the sense that it characterizes those consequence relations that can be defined by a smooth transitive irreflexive preference structure. This is what makes **P** central. More formally:

Definition 24 Suppose $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ is a semantic structure.

Then, $\mathbf{D}_f := \{V \subseteq \mathcal{V} : \exists \alpha \in \mathcal{F}, V = M_\alpha\}$.

Suppose \mathcal{L} is a language containing the usual connectives \neg and \vee , and \mathcal{F} the set of all wffs of \mathcal{L} .

Then define the following condition: $\forall v \in \mathcal{V}, \forall \alpha, \beta \in \mathcal{F}, \forall \Gamma \subseteq \mathcal{F}$,

(KLM0) $v \models \neg\alpha$ iff $v \not\models \alpha$;

(KLM1) $v \models \alpha \vee \beta$ iff $v \models \alpha$ or $v \models \beta$.

(KLM2) if for every finite subset Δ of Γ , $M_\Delta \neq \emptyset$, then $M_\Gamma \neq \emptyset$.

Note that (KLM2) is called “assumption of compactness” in [KLM90].

Proposition 25 [KLM90] Suppose \mathcal{L} is a language containing the usual connectives \neg and \vee , \mathcal{F} the set of all wffs of \mathcal{L} , $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ a semantic structure satisfying (KLM0)–(KLM2), and \sim a relation of $\mathcal{F} \times \mathcal{F}$.

Then, \sim satisfies all the properties of **P** iff there exists a \mathbf{D}_f -smooth transitive irreflexive preference structure \mathcal{R} on \mathcal{V} such that $\forall \alpha, \beta \in \mathcal{F}, \alpha \sim \beta$ iff $\mu_{\mathcal{R}}(M_\alpha) \subseteq M_\beta$.

Note that \sim is a relation on $\mathcal{F} \times \mathcal{F}$, not $\mathcal{P}(\mathcal{F}) \times \mathcal{F}$. This difference is crucial. Indeed, if we adapt the conditions of **P** in the obvious way to relations on $\mathcal{P}(\mathcal{F}) \times \mathcal{F}$ and if we replace \mathbf{D}_f by **D** in Proposition 25, then the latter does no longer hold. This negative result was shown by Schlechta in [Sch92].

Now, by Propositions 20 and 25, we immediately get the following representation theorem:

Proposition 26 Suppose Definition 21 (of preferential consequence relations) is adapted in the obvious way to relations on $\mathcal{F} \times \mathcal{F}$ (essentially, replace **D** by \mathbf{D}_f), \mathcal{L} is a language containing the usual connectives \neg and \vee , \mathcal{F} the set of all wffs of \mathcal{L} , \sim a relation on $\mathcal{F} \times \mathcal{F}$, and $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ a semantic structure such that (KLM0)–(KLM2) hold and $\forall V, W \in \mathbf{D}_f, V \cup W \in \mathbf{D}_f$ and $V \cap W \in \mathbf{D}_f$. Then, LM preferential consequence relations are precisely those relations that satisfy the system **P**.

2.4.1 Example with a classical semantic structure

Let \mathcal{L} be a classical propositional language of which the atoms are r , q , and p . Intuitively, r means Nixon is a republican, q means Nixon is a quaker, and p means Nixon is a pacifist. Let \mathcal{F} be the set of all wffs of \mathcal{L} , \mathcal{V} the set of all classical two-valued valuations of \mathcal{L} , and \models the classical satisfaction


relation for these objects. Then, \mathcal{V} is the set of the 8 following valuations: $v_0, v_1, v_2, v_3, v_4, v_5, v_6,$ and v_7 , which are defined in the obvious way by the following table:

	r	q	p
v_0	0	0	0
v_1	0	0	1
v_2	0	1	0
v_3	0	1	1
v_4	1	0	0
v_5	1	0	1
v_6	1	1	0
v_7	1	1	1

Now, consider the class of all republicans and the class of all quakers. Consider that a republican is normal iff he is not a pacifist and that a quaker is normal iff he is a pacifist. And, consider that a valuation v is more normal than a valuation w from the point of view of a class C iff

- Nixon is an individual of C in both v and w ;
- Nixon is normal in v ;
- Nixon is not normal in w .

In the following graph, there is an arrow from a valuation v to a valuation w iff v is more normal than w from the point of view of some class:



Given those considerations a natural preference structure on \mathcal{V} is $\mathcal{R} = \langle \mathcal{V}, l, \prec \rangle$, where l is identity and \prec is the relation such that $\forall v, w \in \mathcal{V}$, we have $v \prec w$ iff (1) or (2) below holds (i.e. there is an arrow from v to w):

- (1) $v \models r$ and $v \models \neg p$ and $w \models r$ and $w \not\models \neg p$;
- (2) $v \models q$ and $v \models p$ and $w \models q$ and $w \not\models p$.

Finally, let \sim be the preferential consequence relation defined by the coherent choice function $\mu_{\mathcal{R}}$.

Then, \sim leads us to “jump” to plausible conclusions from incomplete information and to revise previous “hasty” conclusions in the face of new and fuller information. For instance, $r \sim \neg p$ and $\{r, p\} \not\sim \neg p$ and $q \sim p$ and $\{q, \neg p\} \not\sim p$.

However, \sim is not paraconsistent. In addition, some sets of formulas are rendered useless, because there is no preferred model for them, though there are models for them. For instance, $\{q, r\} \sim \alpha, \forall \alpha \in \mathcal{F}$.

2.4.2 Example with the *FOUR* semantic structure

Consider the *FOUR* semantic structure $\langle \mathcal{F}_c, \mathcal{V}_4, \models_4 \rangle$ and suppose $\mathcal{A} = \{r, q, p\}$ (these objects have been defined in Section 2.1.2). In addition, make the same considerations about Nixon, the classes, normality, etc., as in Section 2.4.1, except that this time a valuation v is considered to be more normal than a valuation w from the point of view of a class C iff

- in both v and w , the processor is informed that Nixon is an individual of C ;
- in v , he is informed that Nixon is normal and not informed of the contrary;
- in w , he is not informed that Nixon is normal.

See Section 2.1.2 for recalls about the sources-processor systems. Given those considerations a natural preference structure on \mathcal{V}_4 is $\mathcal{R} = \langle \mathcal{V}_4, l, \prec \rangle$, where l is identity and \prec is the relation such that $\forall v, w \in \mathcal{V}_4$, we have $v \prec w$ iff (1) or (2) below holds (i.e. v is more normal than w from the point of view of some class):

- (1) $v \models r$ and $v \models \neg p$ and $v \not\models p$ and $w \models r$ and $w \not\models \neg p$;
- (2) $v \models q$ and $v \models p$ and $v \not\models \neg p$ and $w \models q$ and $w \not\models p$.

Let \sim be the preferential consequence relation defined by the coherent choice function $\mu_{\mathcal{R}}$.

Then, again we “jump” to plausible conclusions and revise previous “hasty” conclusions. For instance, $r \sim \neg p$ and $\{r, p\} \not\sim \neg p$ and $q \sim p$ and $\{q, \neg p\} \not\sim p$.

In addition, \sim is paraconsistent. For instance, $\{p, \neg p, q\} \sim p$ and $\{p, \neg p, q\} \sim \neg p$ and $\{p, \neg p, q\} \sim q$ and $\{p, \neg p, q\} \not\sim \neg q$. And, it happens less often that a set of formulas is rendered useless because there is no preferred model for it, though there are models for it. For instance, this time, $\{q, r\} \sim p$ and $\{q, r\} \sim \neg p$ and $\{q, r\} \sim q$ and $\{q, r\} \not\sim \neg q$ and $\{q, r\} \sim r$ and $\{q, r\} \not\sim \neg r$.

However, \sim does not satisfy the Disjunctive Syllogism. Indeed, for instance, $\{\neg r, r \vee q\} \not\sim q$.

3 Contributions

The main contributions of the present paper are summarized below. We characterized (in many cases, by purely syntactic conditions) families of preferential and preferential-discriminative consequence relations. Sometimes, we will need to make some assumptions about the semantic structure under consideration. However, no assumption will be needed for the three following families:

- the preferential consequence relations (Section 3.2);
- the DP preferential consequence relations (Section 3.1);
- the DP LM preferential consequence relations (Section 3.1).

We will assume (A1) and (A3) for:

- the CP preferential-discriminative consequence relations (Section 3.4);
- the CP DP preferential-discriminative consequence relations (Section 3.3);
- the CP DP LM preferential-discriminative consequence relations (Section 3.3).

And, we will need (A1), (A2), and (A3) for:

- the preferential-discriminative consequence relations (Section 3.4);
- the DP preferential-discriminative consequence relations (Section 3.3);
- the DP LM preferential-discriminative consequence relations (Section 3.3).

3.1 The non-discriminative and definability preserving case

The characterizations in this section have already been given in Proposition 3.1 of [Sch00], under the assumption that a classical propositional semantic structure is considered. Using the same techniques as those of Schlechta, we show that his characterizations hold with any semantic structure.

Notation 27 Let $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ be a semantic structure and \sim a relation on $\mathcal{P}(\mathcal{F}) \times \mathcal{F}$. Then, consider the following conditions: $\forall \Gamma, \Delta \subseteq \mathcal{F}$,

- (\sim 0) if $\vdash(\Gamma) = \vdash(\Delta)$, then $\sim(\Gamma) = \sim(\Delta)$;
- (\sim 1) $\vdash(\sim(\Gamma)) = \sim(\Gamma)$;
- (\sim 2) $\Gamma \subseteq \sim(\Gamma)$;
- (\sim 3) $\sim(\Gamma, \Delta) \subseteq \vdash(\sim(\Gamma), \Delta)$;
- (\sim 4) if $\Gamma \subseteq \vdash(\Delta) \subseteq \sim(\Gamma)$, then $\sim(\Gamma) \subseteq \sim(\Delta)$.

Note that those conditions are purely syntactic when there is a proof system available for \vdash (which is the case with e.g. the classical, *FOUR*, and *J₃* semantic structures).

Proposition 28 Let $\mathcal{S} = \langle \mathcal{F}, \mathcal{V}, \models \rangle$ be a semantic structure and \sim a relation on $\mathcal{P}(\mathcal{F}) \times \mathcal{F}$. Then,

- (0) \sim is a DP preferential consequence relation iff (\sim 0), (\sim 1), (\sim 2), and (\sim 3) hold;
- (1) \sim is a DP LM preferential consequence relation iff (\sim 0), (\sim 1), (\sim 2), (\sim 3), and (\sim 4) hold.

Proof *Proof of (0).* Direction: “ \rightarrow ”.

By hypothesis, there exists a DP coherent choice function μ from \mathbf{D} to $\mathcal{P}(\mathcal{V})$ such that $\forall \Gamma \subseteq \mathcal{F}$, $\sim(\Gamma) = T(\mu(M_\Gamma))$. We will show:

- (0.0) \sim satisfies (\sim 0);
- (0.1) \sim satisfies (\sim 1);
- (0.2) \sim satisfies (\sim 2).

Before turning to (\sim 3), we need a preliminary result:

- (0.3) $\forall \Gamma \subseteq \mathcal{F}$, we have $\mu(M_\Gamma) = M_{\sim(\Gamma)}$;
- (0.4) \sim satisfies (\sim 3).

Direction: “ \leftarrow ”.

Suppose \sim satisfies (\sim 0), (\sim 1), (\sim 2), and (\sim 3).

Let μ be the function from \mathbf{D} to $\mathcal{P}(\mathcal{V})$ such that $\forall \Gamma \subseteq \mathcal{F}$, $\mu(M_\Gamma) = M_{\sim(\Gamma)}$.

Then, μ is well-defined.

Indeed, If $\Gamma, \Delta \subseteq \mathcal{F}$ and $M_\Gamma = M_\Delta$, then $\vdash(\Gamma) = \vdash(\Delta)$, thus, by (\sim 0), $\sim(\Gamma) = \sim(\Delta)$.

In addition, μ is obviously DP. We show the following which ends the proof:

- (0.5) μ is a choice function;

(0.6) μ is coherent;

(0.7) $\forall \Gamma \subseteq \mathcal{F}$, we have $\sim(\Gamma) = T(\mu(M_\Gamma))$.

Proof of (0.0). Let $\Gamma, \Delta \subseteq \mathcal{F}$ and suppose $\vdash(\Gamma) = \vdash(\Delta)$.

Then, $M_\Gamma = M_\Delta$. Thus, $\sim(\Gamma) = T(\mu(M_\Gamma)) = T(\mu(M_\Delta)) = \sim(\Delta)$.

Proof of (0.1). Let $\Gamma \subseteq \mathcal{F}$. Then, $\vdash(\sim(\Gamma)) = \vdash(T(\mu(M_\Gamma))) = T(M_{T(\mu(M_\Gamma))}) = \sim(\Gamma)$.

Proof of (0.2). Let $\Gamma \subseteq \mathcal{F}$. Then, $\Gamma \subseteq T(M_\Gamma) \subseteq T(\mu(M_\Gamma)) = \sim(\Gamma)$.

Proof of (0.3). Let $\Gamma \subseteq \mathcal{F}$. As, μ is DP, $\mu(M_\Gamma) \in \mathbf{D}$.

Thus, $\exists \Gamma' \subseteq \mathcal{F}$, $\mu(M_\Gamma) = M_{\Gamma'}$. Therefore, $\mu(M_\Gamma) = M_{\Gamma'} = M_{T(M_{\Gamma'})} = M_{T(\mu(M_\Gamma))} = M_{\sim(\Gamma)}$.

Proof of (0.4). Let $\Gamma, \Delta \subseteq \mathcal{F}$. As, $M_{\Gamma, \Delta} \subseteq M_\Gamma$ and μ is coherent, $\mu(M_\Gamma) \cap M_{\Gamma, \Delta} \subseteq \mu(M_{\Gamma, \Delta})$.

Therefore, $\sim(\Gamma, \Delta) = T(\mu(M_{\Gamma, \Delta})) \subseteq T(\mu(M_\Gamma) \cap M_{\Gamma, \Delta}) = T(\mu(M_\Gamma) \cap M_\Delta)$.

Thus, by (0.0), $\sim(\Gamma, \Delta) \subseteq T(M_{\sim(\Gamma)} \cap M_\Delta) = T(M_{\sim(\Gamma), \Delta}) = \vdash(\sim(\Gamma), \Delta)$.

Proof of (0.5). Let $\Gamma \subseteq \mathcal{F}$. Then, $\mu(M_\Gamma) = M_{\sim(\Gamma)}$, which is, by (\sim 2), a subset of M_Γ .

Proof of (0.6). Let $\Gamma, \Delta \subseteq \mathcal{F}$ and suppose $M_\Gamma \subseteq M_\Delta$.

Then, $\mu(M_\Delta) \cap M_\Gamma = M_{\sim(\Delta)} \cap M_\Gamma = M_{\sim(\Delta), \Gamma}$.

But, by (\sim 3), $M_{\sim(\Delta), \Gamma} \subseteq M_{\sim(\Delta, \Gamma)} = \mu(M_{\Delta, \Gamma}) = \mu(M_\Gamma)$.

Proof of (0.7). Let $\Gamma \subseteq \mathcal{F}$. Then, by (\sim 1), $\sim(\Gamma) = \vdash(\sim(\Gamma)) = T(M_{\sim(\Gamma)}) = T(\mu(M_\Gamma))$.

Proof of (1). Direction: “ \rightarrow ”.

Verbatim the same proof as for (0), except that in addition μ is LM.

We use it to show that \sim satisfies (\sim 4).

Let $\Gamma, \Delta \subseteq \mathcal{F}$ and suppose $\Gamma \subseteq \vdash(\Delta) \subseteq \sim(\Gamma)$.

Then, by (0.3), $\mu(M_\Gamma) = M_{\sim(\Gamma)} \subseteq M_{\vdash(\Delta)} = M_\Delta \subseteq M_\Gamma$.

Therefore, as μ is locally monotonic, $\mu(M_\Delta) \subseteq \mu(M_\Gamma)$.

Thus, $\sim(\Gamma) = T(\mu(M_\Gamma)) \subseteq T(\mu(M_\Delta)) = \sim(\Delta)$.

Direction: “ \leftarrow ”.

Verbatim the same proof as for (0), except that in addition (\sim 4) is satisfied.

We use it to show that μ is locally monotonic.

Let $\Gamma, \Delta \subseteq \mathcal{F}$ and suppose $\mu(M_\Gamma) \subseteq M_\Delta \subseteq M_\Gamma$.

Then, $M_{\sim(\Gamma)} \subseteq M_\Delta \subseteq M_\Gamma$. Therefore, $\Gamma \subseteq T(M_\Gamma) \subseteq T(M_\Delta) = \vdash(\Delta)$.

On the other hand, $\vdash(\Delta) = T(M_\Delta) \subseteq T(M_{\sim(\Gamma)}) = \vdash(\sim(\Gamma))$ which is, by (\sim 1), equal to $\sim(\Gamma)$.

Thus, by (\sim 4), we have $\sim(\Gamma) \subseteq \sim(\Delta)$. Therefore, $\mu(M_\Delta) = M_{\sim(\Delta)} \subseteq M_{\sim(\Gamma)} = \mu(M_\Gamma)$. ■

3.2 The non-discriminative and not necessarily definability preserving case

In this section, we will characterize the family of all preferential consequence relations. Unlike in Section 3.1, our conditions will not be purely syntactic (i.e. using only \vdash , \sim , etc.). In fact, properties like Coherence cannot be translated in syntactic terms because the choice functions under consideration are not necessarily definability preserving. Indeed, we do no longer have at our disposal the remarkable equality: $\mu(M_\Gamma) = M_{\sim(\Gamma)}$, which is of great help to perform the translation and which

holds precisely because of Definability Preservation.

In Proposition 5.2.11 of [Sch04], K. Schlechta provided a characterization of the aforementioned family, under the assumption that a classical propositional semantic structure is considered. Note that most of his work is done in a very general, in fact algebraic, framework. Only at the end, he applied his general lemmas in a classical framework to get the characterization. The conditions he gave, as ours, are not purely syntactic (e.g. they involve the notion of model, etc.). Moreover, some limits of what can be done in this area have been put in evidence by Schlechta. Approximatively, he showed in Proposition 5.2.15 of the same book that, in an infinite classical framework, there does not exist a characterization containing only conditions which are universally quantified, of limited size, and using only simple operations (like e.g. \cup , \cap , \setminus).

The purpose of the present section is to provide a new characterization, more elegant than the one of Schlechta and that hold with any semantic structure. To do so, we have been inspired by the algebraic part of the work of Schlechta (see Proposition 5.2.5 of [Sch04]). Technically, the idea begins by building from any function f , a coherent choice function μ_f such that whenever f “covers” some coherent choice function, it necessarily covers μ_f .

Definition 29 Let \mathcal{V} be a set, \mathbf{V} and \mathbf{W} subsets of $\mathcal{P}(\mathcal{V})$, and f a function from \mathbf{V} to \mathbf{W} . We denote by μ_f the function from \mathbf{V} to $\mathcal{P}(\mathcal{V})$ such that $\forall V \in \mathbf{V}$,

$$\mu_f(V) = \{v \in V : \forall W \in \mathbf{V}, \text{ if } v \in W \subseteq V, \text{ then } v \in f(W)\}.$$

Lemma 30 Let \mathcal{V} be a set, \mathbf{V} and \mathbf{W} subsets of $\mathcal{P}(\mathcal{V})$, and f a function from \mathbf{V} to \mathbf{W} . Then, μ_f is a coherent choice function.

Proof μ_f is obviously a choice function. It remains to show that it is coherent.

Suppose $V, W \in \mathbf{V}$, $V \subseteq W$, and $v \in \mu_f(W) \cap V$. We show $v \in \mu_f(V)$.

To do so, suppose the contrary, i.e. suppose $v \notin \mu_f(V)$.

Then, as $v \in V$, we have $\exists Z \in \mathbf{V}$, $Z \subseteq V$, $v \in Z$, and $v \notin f(Z)$.

But, $V \subseteq W$, thus $Z \subseteq W$. Therefore, by definition of μ_f , $v \notin \mu_f(W)$, which is impossible. ■

Lemma 31 Let \mathcal{V} be a set, \mathbf{V} , \mathbf{W} , and \mathbf{X} subsets of $\mathcal{P}(\mathcal{V})$, f a function from \mathbf{V} to \mathbf{W} , and μ a coherent choice function from \mathbf{V} to \mathbf{X} such that $\forall V \in \mathbf{V}$, $f(V) = M_{T(\mu(V))}$.

Then, $\forall V \in \mathbf{V}$, $f(V) = M_{T(\mu_f(V))}$.

Proof Let $V \in \mathbf{V}$. We show $f(V) = M_{T(\mu_f(V))}$.

Case 1: $\exists v \in \mu(V)$, $v \notin \mu_f(V)$.

As $\mu(V) \subseteq V$, we have $v \in V$.

Thus, by definition of μ_f , $\exists W \in \mathbf{V}$, $W \subseteq V$, $v \in W$, and $v \notin f(W) = M_{T(\mu(W))} \supseteq \mu(W)$.

On the other hand, as μ is coherent, $\mu(V) \cap W \subseteq \mu(W)$. Thus, $v \in \mu(W)$, which is impossible.

Case 2: $\mu(V) \subseteq \mu_f(V)$.

Case 2.1: $\exists v \in \mu_f(V)$, $v \notin f(V)$.

Then, $\exists W \in \mathbf{V}$, $W \subseteq V$, $v \in W$, and $v \notin f(W)$. Indeed, just take V itself for the choice of W .

Therefore, $v \notin \mu_f(V)$, which is impossible.

Case 2.2: $\mu_f(V) \subseteq f(V)$.

Then, $f(V) = M_{T(\mu(V))} \subseteq M_{T(\mu_f(V))} \subseteq M_{T(f(V))} = M_{T(M_{T(\mu(V))})} = M_{T(\mu(V))} = f(V)$. ■

Now, everything is ready to show the representation result.

Notation 32 Let $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ be a semantic structure and \sim a relation on $\mathcal{P}(\mathcal{F}) \times \mathcal{F}$.

Then, consider the following condition: $\forall \Gamma \subseteq \mathcal{F}$,

(~ 5) $\vdash(\Gamma) = T(\{v \in M_\Gamma : \forall \Delta \subseteq \mathcal{F}, \text{ if } v \in M_\Delta \subseteq M_\Gamma, \text{ then } v \in M_{\vdash(\Delta)}\})$.

Proposition 33 Let $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ be a semantic structure and \vdash a relation on $\mathcal{P}(\mathcal{F}) \times \mathcal{F}$. Then, \vdash is a preferential consequence relation iff (~ 5) holds.

Proof *Direction: “ \rightarrow ”.*

There exists a coherent choice function μ from \mathbf{D} to $\mathcal{P}(\mathcal{V})$ such that $\forall \Gamma \subseteq \mathcal{F}, \vdash(\Gamma) = T(\mu(M_\Gamma))$.

Let f be the function from \mathbf{D} to \mathbf{D} such that $\forall V \in \mathbf{D}$, we have $f(V) = M_{T(\mu(V))}$.

By Lemma 31, $\forall V \in \mathbf{D}$, we have $f(V) = M_{T(\mu_f(V))}$.

Note that $\forall \Gamma \subseteq \mathcal{F}, f(M_\Gamma) = M_{T(\mu(M_\Gamma))} = M_{\vdash(\Gamma)}$.

We show that (~ 5) holds. Let $\Gamma \subseteq \mathcal{F}$.

Then, $\vdash(\Gamma) = T(\mu(M_\Gamma)) = T(M_{T(\mu(M_\Gamma))}) = T(f(M_\Gamma)) = T(M_{T(\mu_f(M_\Gamma))}) = T(\mu_f(M_\Gamma)) = T(\{v \in M_\Gamma : \forall W \in \mathbf{D}, \text{ if } v \in W \subseteq M_\Gamma, \text{ then } v \in f(W)\}) = T(\{v \in M_\Gamma : \forall \Delta \subseteq \mathcal{F}, \text{ if } v \in M_\Delta \subseteq M_\Gamma, \text{ then } v \in f(M_\Delta)\}) = T(\{v \in M_\Gamma : \forall \Delta \subseteq \mathcal{F}, \text{ if } v \in M_\Delta \subseteq M_\Gamma, \text{ then } v \in M_{\vdash(\Delta)}\})$.

Direction: “ \leftarrow ”.

Suppose \vdash satisfies (~ 5).

Let f be the function from \mathbf{D} to \mathbf{D} such that $\forall \Gamma \subseteq \mathcal{F}$, we have $f(M_\Gamma) = M_{\vdash(\Gamma)}$.

Note that f is well-defined. Indeed, if $\Gamma, \Delta \subseteq \mathcal{F}$ and $M_\Gamma = M_\Delta$, then, by (~ 5), $\vdash(\Gamma) = \vdash(\Delta)$.

In addition, by (~ 5), we clearly have $\forall \Gamma \subseteq \mathcal{F}, \vdash(\Gamma) = T(\mu_f(M_\Gamma))$.

And finally, by Lemma 30, μ_f is a coherent choice function. \blacksquare

3.3 The discriminative and definability preserving case

In this section, we will characterize certain families of DP preferential-discriminative consequence relations. To do so, we will develop new techniques (especially Lemmas 40 and 41 below). We need basic notations and an inductive construction:

Notation 34 \mathbb{N} denotes the natural numbers including 0: $\{0, 1, 2, \dots\}$.

\mathbb{N}^+ denotes the strictly positive natural numbers: $\{1, 2, \dots\}$.

\mathbb{Z} denotes the integers.

Let $i, j \in \mathbb{Z}$. Then, $[i, j]$ denotes the set of all $k \in \mathbb{Z}$ such that $i \leq k \leq j$.

Let \mathcal{L} be a language, \vee a binary connective of \mathcal{L} , \mathcal{F} the set of all wffs of \mathcal{L} , and $\beta_1, \beta_2, \dots, \beta_r \in \mathcal{F}$.

Whenever we write $\beta_1 \vee \beta_2 \vee \dots \vee \beta_r$, we mean $(\dots((\beta_1 \vee \beta_2) \vee \beta_3) \vee \dots \vee \beta_{r-1}) \vee \beta_r$.

Definition 35 Let \mathcal{L} be a language, \neg a unary connective of \mathcal{L} , \mathcal{F} the set of all wffs of \mathcal{L} , $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ a semantic structure, \vdash a relation on $\mathcal{P}(\mathcal{F}) \times \mathcal{F}$, and $\Gamma \subseteq \mathcal{F}$. Then,

$$H_1(\Gamma) := \{\neg\beta \in \mathcal{F} : \beta \in \vdash(\Gamma, \vdash(\Gamma)) \setminus \vdash(\Gamma) \text{ and } \neg\beta \notin \vdash(\Gamma, \vdash(\Gamma))\}.$$

Let $i \in \mathbb{N}$ with $i \geq 2$. Then,

$$H_i(\Gamma) := \{\neg\beta \in \mathcal{F} : \left\{ \begin{array}{l} \beta \in \vdash(\Gamma, \vdash(\Gamma), H_1(\Gamma), \dots, H_{i-1}(\Gamma)) \setminus \vdash(\Gamma) \text{ and} \\ \neg\beta \notin \vdash(\Gamma, \vdash(\Gamma), H_1(\Gamma), \dots, H_{i-1}(\Gamma)) \end{array} \right\}.$$

$$H(\Gamma) := \bigcup_{i \in \mathbb{N}^+} H_i(\Gamma).$$

Definition 36 Suppose \mathcal{L} is a language, \neg a unary connective of \mathcal{L} , \vee a binary connective of \mathcal{L} , \mathcal{F} the set of all wffs of \mathcal{L} , $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ a semantic structure, and \sim a relation on $\mathcal{P}(\mathcal{F}) \times \mathcal{F}$. Then, consider the following conditions: $\forall \Gamma, \Delta \subseteq \mathcal{F}, \forall \alpha, \beta \in \mathcal{F}$,

- (\sim 6) if $\beta \in \vdash(\Gamma, \sim(\Gamma)) \setminus \sim(\Gamma)$ and $\neg\alpha \in \vdash(\Gamma, \sim(\Gamma), \neg\beta)$, then $\alpha \notin \sim(\Gamma)$;
- (\sim 7) if $\alpha \in \vdash(\Gamma, \sim(\Gamma)) \setminus \sim(\Gamma)$ and $\beta \in \vdash(\Gamma, \sim(\Gamma), \neg\alpha) \setminus \sim(\Gamma)$, then $\alpha \vee \beta \notin \sim(\Gamma)$;
- (\sim 8) if $\alpha \in \sim(\Gamma)$, then $\neg\alpha \notin \vdash(\Gamma, \sim(\Gamma))$;
- (\sim 9) if $\Delta \subseteq \vdash(\Gamma)$, then $\sim(\Gamma) \cup H(\Gamma) \subseteq \vdash(\Delta, \sim(\Delta), H(\Delta), \Gamma)$;
- (\sim 10) if $\Gamma \subseteq \vdash(\Delta) \subseteq \vdash(\Gamma, \sim(\Gamma), H(\Gamma))$, then $\sim(\Gamma) \cup H(\Gamma) \subseteq \vdash(\Delta, \sim(\Delta), H(\Delta))$;
- (\sim 11) if Γ is consistent, then $\sim(\Gamma)$ is consistent, $\Gamma \subseteq \sim(\Gamma)$, and $\vdash(\sim(\Gamma)) = \sim(\Gamma)$.

Note that those conditions are purely syntactic when there is a proof system available for \vdash .

Proposition 37 Suppose \mathcal{L} is a language, \neg a unary connective of \mathcal{L} , \vee and \wedge binary connectives of \mathcal{L} , \mathcal{F} the set of all wffs of \mathcal{L} , $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ a semantic structure satisfying (A1) and (A3), and \sim a relation on $\mathcal{P}(\mathcal{F}) \times \mathcal{F}$. Then,

- (0) \sim is a CP DP preferential-discriminative consequence relation iff (\sim 0), (\sim 6), (\sim 7), (\sim 8), (\sim 9), and (\sim 11) hold;
- (1) \sim is a CP DP LM preferential-discriminative consequence relation iff (\sim 0), (\sim 6), (\sim 7), (\sim 8), (\sim 9), (\sim 10), and (\sim 11) hold.

Suppose $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ satisfies (A2) too. Then,

- (2) \sim is a DP preferential-discriminative consequence relation iff (\sim 0), (\sim 6), (\sim 7), (\sim 8), and (\sim 9) hold;
- (3) \sim is a DP LM preferential-discriminative consequence relation iff (\sim 0), (\sim 6), (\sim 7), (\sim 8), (\sim 9), and (\sim 10) hold.

The proof of Proposition 37 has been relegated at the end of Section 3.3. We need first Notation 34, Definition 38 and Lemmas 39, 40, and 41 below. Here are some purely technical tools:

Definition 38 Suppose \mathcal{L} is a language, \neg a unary connective of \mathcal{L} , \vee a binary connective of \mathcal{L} , \mathcal{F} the set of all wffs of \mathcal{L} , $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ a semantic structure satisfying (A1), \sim a relation on $\mathcal{P}(\mathcal{F}) \times \mathcal{F}$, and $\Gamma \subseteq \mathcal{F}$. Then,

$$M_{\Gamma}^1 := \{v \in M_{\Gamma, \sim(\Gamma)} : \exists \beta \in T(M_{\Gamma, \sim(\Gamma)}) \setminus \sim(\Gamma), v \notin M_{\neg\beta}\}.$$

Let $i \in \mathbf{N}$ with $i \geq 2$. Then,

$$M_{\Gamma}^i := \{v \in M_{\Gamma, \sim(\Gamma)} \setminus M_{\Gamma}^1 \cup \dots \cup M_{\Gamma}^{i-1} : \exists \beta \in T(M_{\Gamma, \sim(\Gamma)} \setminus M_{\Gamma}^1 \cup \dots \cup M_{\Gamma}^{i-1}) \setminus \sim(\Gamma), v \notin M_{\neg\beta}\}.$$

$$M'_{\Gamma} := \bigcup_{i \in \mathbf{N}^+} M_{\Gamma}^i$$

$$n(\Gamma) := |\{i \in \mathbf{N}^+ : M_{\Gamma}^i \neq \emptyset\}|$$

Suppose $M_\Gamma^1 \neq \emptyset$. Then, we denote by β_Γ^1 an element of \mathcal{F} , chosen arbitrarily, such that $\exists r \in \mathbb{N}^+, \exists v_1, v_2, \dots, v_r \in \mathcal{V}$, and $\exists \beta_1, \beta_2, \dots, \beta_r \in \mathcal{F}$ with $M_\Gamma^1 = \{v_1, v_2, \dots, v_r\}$,

$$\beta_\Gamma^1 = \beta_1 \vee \beta_2 \vee \dots \vee \beta_r,$$

and $\forall j \in [1, r], \beta_j \notin \vdash(\Gamma)$, $M_{\Gamma, \vdash(\Gamma)} \subseteq M_{\beta_j}$, and $v_j \notin M_{\neg\beta_j}$.
As $M_\Gamma^1 \neq \emptyset$ and M_Γ^1 is finite (thanks to (A1)), such an element exists.

Suppose $i \in \mathbb{N}, i \geq 2$, and $M_\Gamma^i \neq \emptyset$.

Then, we denote by β_Γ^i an element of \mathcal{F} , chosen arbitrarily, such that $\exists r \in \mathbb{N}^+, \exists v_1, v_2, \dots, v_r \in \mathcal{V}$, and $\exists \beta_1, \beta_2, \dots, \beta_r \in \mathcal{F}$ with $M_\Gamma^i = \{v_1, v_2, \dots, v_r\}$,

$$\beta_\Gamma^i = \beta_1 \vee \beta_2 \vee \dots \vee \beta_r,$$

and $\forall j \in [1, r], \beta_j \notin \vdash(\Gamma)$, $M_{\Gamma, \vdash(\Gamma)} \setminus M_\Gamma^1 \cup \dots \cup M_\Gamma^{i-1} \subseteq M_{\beta_j}$, and $v_j \notin M_{\neg\beta_j}$.
As $M_\Gamma^i \neq \emptyset$ and M_Γ^i is finite, such an element exists.

Suppose $M'_\Gamma \neq \emptyset$. Then,

$$\beta_\Gamma := \beta_\Gamma^1 \vee \beta_\Gamma^2 \vee \dots \vee \beta_\Gamma^{n(\Gamma)}$$

As $M'_\Gamma \neq \emptyset$, $n(\Gamma) \geq 1$. In addition, we will show in Lemma 39 below that $n(\Gamma)$ is finite and $\forall i \in \mathbb{N}^+$ with $i \leq n(\Gamma)$, $M_\Gamma^i \neq \emptyset$. Thus, β_Γ is well-defined.

$$F(\Gamma) := \begin{cases} \{\neg\beta_\Gamma\} & \text{if } M'_\Gamma \neq \emptyset \\ \emptyset & \text{otherwise} \end{cases}$$

$$G(\Gamma) := \{\alpha \in \mathcal{F} : \alpha \notin \vdash(\Gamma), \neg\alpha \notin \vdash(\Gamma), \text{ and } T_d(M_{\Gamma, \vdash(\Gamma)}, \alpha) \subseteq \vdash(\Gamma)\}$$

Here are some quick results about the purely technical tools defined just above:

Lemma 39 Suppose \mathcal{L} is a language, \neg a unary connective of \mathcal{L} , \vee a binary connective of \mathcal{L} , \mathcal{F} the set of all wffs of \mathcal{L} , $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ a semantic structure satisfying (A1), \vdash a relation on $\mathcal{P}(\mathcal{F}) \times \mathcal{F}$, $\Gamma \subseteq \mathcal{F}$, and $i, j \in \mathbb{N}^+$. Then,

- (0) if $i \neq j$, then $M_\Gamma^i \cap M_\Gamma^j = \emptyset$;
- (1) if $M_\Gamma^i = \emptyset$, then $M_\Gamma^{i+1} = \emptyset$;
- (2) $T_d(M_{\Gamma, \vdash(\Gamma)}) \subseteq \vdash(\Gamma)$ iff $M_\Gamma^1 = \emptyset$;
- (3) if $i \geq 2$, then $T_d(M_{\Gamma, \vdash(\Gamma)} \setminus M_\Gamma^1 \cup \dots \cup M_\Gamma^{i-1}) \subseteq \vdash(\Gamma)$ iff $M_\Gamma^i = \emptyset$;
- (4) $n(\Gamma)$ is finite;
- (5) if $i \leq n(\Gamma)$, then $M_\Gamma^i \neq \emptyset$;
- (6) if $i > n(\Gamma)$, then $M_\Gamma^i = \emptyset$;
- (7) if $M'_\Gamma \neq \emptyset$, then $M'_\Gamma = M_\Gamma^1 \cup \dots \cup M_\Gamma^{n(\Gamma)}$;
- (8) $T_d(M_{\Gamma, \vdash(\Gamma)} \setminus M'_\Gamma) \subseteq \vdash(\Gamma)$.

Proof *Proofs of (0), (1), (2), and (3).* Trivial.

Proof of (4). Obvious by (0) and (A1).

Proof of (5). Suppose $\exists i \in \mathbb{N}^+$, $M_\Gamma^i = \emptyset$ and $i \leq n(\Gamma)$.

Then, by (1), $\forall j \in \mathbb{N}^+$, $j \geq i$, $M_\Gamma^j = \emptyset$.

Thus, $|\{j \in \mathbb{N}^+ : M_\Gamma^j \neq \emptyset\}| \leq i - 1 < n(\Gamma)$, which is impossible.

Proof of (6). Suppose $\exists i \in \mathbb{N}^+$, $M_\Gamma^i \neq \emptyset$ and $i > n(\Gamma)$.

Then, by (1), $\forall j \in \mathbb{N}^+$, $j \leq i$, $M_\Gamma^j \neq \emptyset$.

Thus, $|\{j \in \mathbb{N}^+ : M_\Gamma^j \neq \emptyset\}| \geq i > n(\Gamma)$, which is impossible.

Proof of (7). Obvious by (6).

Proof of (8). Case 1: $M'_\Gamma = \emptyset$.

Then, $T_d(M_{\Gamma, \sim(\Gamma)} \setminus M'_\Gamma) = T_d(M_{\Gamma, \sim(\Gamma)})$. In addition, $M_\Gamma^1 = \emptyset$. Thus, by (2), we are done.

Case 2: $M'_\Gamma \neq \emptyset$.

Then, by (7), $T_d(M_{\Gamma, \sim(\Gamma)} \setminus M'_\Gamma) = T_d(M_{\Gamma, \sim(\Gamma)} \setminus M_\Gamma^1 \cup \dots \cup M_\Gamma^{n(\Gamma)})$.

In addition, $n(\Gamma) + 1 \geq 2$ and, by (6), $M_\Gamma^{n(\Gamma)+1} = \emptyset$. Thus, by (3), we are done. ■

We turn to an important lemma. Its main goal is to show that the conditions (\sim 6), (\sim 7), and (\sim 8) are sufficient to establish the following important equality: $\sim(\Gamma) = T_d(M_{\Gamma, \sim(\Gamma), H(\Gamma)})$, which provides a semantic definition of \sim (in the discriminative manner).

Lemma 40 Suppose \mathcal{L} is a language, \neg a unary connective of \mathcal{L} , \vee and \wedge binary connectives of \mathcal{L} , \mathcal{F} the set of all wffs of \mathcal{L} , $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ a semantic structure satisfying (A1) and (A3), \sim a relation on $\mathcal{P}(\mathcal{F}) \times \mathcal{F}$ satisfying (\sim 6), (\sim 7), and (\sim 8), and $\Gamma \subseteq \mathcal{F}$. Then,

- (0) if $M'_\Gamma \neq \emptyset$, then $\beta_\Gamma \notin \sim(\Gamma)$;
- (1) if $M'_\Gamma \neq \emptyset$, then $M_{\Gamma, \sim(\Gamma)} \subseteq M_{\beta_\Gamma}$;
- (2) if $M'_\Gamma \neq \emptyset$, then $M'_\Gamma \cap M_{\neg\beta_\Gamma} = \emptyset$;
- (3) if $M'_\Gamma \neq \emptyset$, then $M_{\Gamma, \sim(\Gamma)} \setminus M'_\Gamma \subseteq M_{\neg\beta_\Gamma}$;
- (4) $M_{\Gamma, \sim(\Gamma)} \setminus M'_\Gamma = M_{\Gamma, \sim(\Gamma), F(\Gamma)}$;
- (5) $\sim(\Gamma) = T_d(M_{\Gamma, \sim(\Gamma), F(\Gamma)})$;
- (6) $M_{\Gamma, \sim(\Gamma), H(\Gamma)} = M_{\Gamma, \sim(\Gamma), F(\Gamma)}$;
- (7) $\sim(\Gamma) = T_d(M_{\Gamma, \sim(\Gamma), H(\Gamma)})$.

Proof *Proof of (0), (1), and (2).* Suppose $M'_\Gamma \neq \emptyset$.

Then, it suffices to show by induction: $\forall i \in [1, n(\Gamma)]$,

$$p_3(i) \quad (M_\Gamma^1 \cup \dots \cup M_\Gamma^i) \cap M_{\neg(\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^i)} = \emptyset;$$

$$p_2(i) \quad M_{\Gamma, \sim(\Gamma)} \subseteq M_{\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^i};$$

$$p_1(i) \quad \beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^i \notin \sim(\Gamma).$$

As $M_\Gamma^1 \neq \emptyset$, $\exists r \in \mathbb{N}^+$, $\exists v_1, v_2, \dots, v_r \in \mathcal{V}$, and $\exists \beta_1, \beta_2, \dots, \beta_r \in \mathcal{F}$, $M_\Gamma^1 = \{v_1, \dots, v_r\}$,

$\beta_\Gamma^1 = \beta_1 \vee \dots \vee \beta_r$, and $\forall j \in [1, r]$, $\beta_j \notin \vdash(\Gamma)$, $M_{\Gamma, \vdash(\Gamma)} \subseteq M_{\beta_j}$, and $v_j \notin M_{\neg\beta_j}$.

Then, it can be shown that:

(0.0) $p_3(1)$ holds;

(0.1) $p_2(1)$ holds;

(0.2) $p_1(1)$ holds.

Now, let $i \in [1, n(\Gamma) - 1]$ and suppose $p_1(i)$, $p_2(i)$, and $p_3(i)$ hold.

As $M_\Gamma^{i+1} \neq \emptyset$, $\exists r \in \mathbb{N}^+$, $\exists v_1, v_2, \dots, v_r \in \mathcal{V}$, and $\exists \beta_1, \beta_2, \dots, \beta_r \in \mathcal{F}$,

$M_\Gamma^{i+1} = \{v_1, \dots, v_r\}$, $\beta_\Gamma^{i+1} = \beta_1 \vee \dots \vee \beta_r$, and

$\forall j \in [1, r]$, $\beta_j \notin \vdash(\Gamma)$, $M_{\Gamma, \vdash(\Gamma)} \setminus M_\Gamma^1 \cup \dots \cup M_\Gamma^i \subseteq M_{\beta_j}$, and $v_j \notin M_{\neg\beta_j}$.

Then, it can be shown that:

(0.3) $p_3(i+1)$ holds;

(0.4) $p_2(i+1)$ holds.

Before turning to $p_1(i+1)$, we need the following:

(0.5) $\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^i \vee \beta_1 \vee \beta_2 \vee \dots \vee \beta_r \notin \vdash(\Gamma)$;

(0.6) $p_1(i+1)$ holds.

Proof of (0.0). If $v_j \in M_\Gamma^1$, then $v_j \notin M_{\neg\beta_j}$. But, by (A3), $M_{\neg\beta_\Gamma^1} \subseteq M_{\neg\beta_j}$.

Proof of (0.1). We have $M_{\Gamma, \vdash(\Gamma)} \subseteq M_{\beta_1}$ which is, by (A3), a subset of $M_{\beta_\Gamma^1}$.

Proof of (0.2). It suffices to show by induction: $\forall j \in [1, r]$,

$q(j) \quad \beta_1 \vee \dots \vee \beta_j \notin \vdash(\Gamma)$.

Obviously, $q(1)$ holds.

Let $j \in [1, r-1]$. Suppose $q(j)$. We show $q(j+1)$.

By (A3), we have $M_{\Gamma, \vdash(\Gamma)} \subseteq M_{\beta_1 \vee \dots \vee \beta_j}$.

On the other hand, $M_{\Gamma, \vdash(\Gamma), \neg(\beta_1 \vee \dots \vee \beta_j)} \subseteq M_{\Gamma, \vdash(\Gamma)} \subseteq M_{\beta_{j+1}}$.

Thus, by $q(j)$ and ($\vdash 7$) (where α is $\beta_1 \vee \dots \vee \beta_j$ and β is β_{j+1}), we get $\beta_1 \vee \dots \vee \beta_{j+1} \notin \vdash(\Gamma)$.

Proof of (0.3). Let $v \in M_\Gamma^1 \cup \dots \cup M_\Gamma^{i+1}$. We show $v \notin M_{\neg(\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^{i+1})}$.

Case 1: $v \in M_\Gamma^1 \cup \dots \cup M_\Gamma^i$.

Then, by $p_3(i)$, we have $v \notin M_{\neg(\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^i)}$. But, by (A3), $M_{\neg(\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^{i+1})} \subseteq M_{\neg(\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^i)}$.

Case 2: $v \in M_\Gamma^{i+1}$.

Then, $\exists j \in [1, r]$, $v = v_j$. Thus, $v \notin M_{\neg\beta_j}$. But, by (A3), $M_{\neg(\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^{i+1})} \subseteq M_{\neg\beta_\Gamma^{i+1}} \subseteq M_{\neg\beta_j}$.

Proof of (0.4). By $p_2(i)$, $M_{\Gamma, \vdash(\Gamma)} \subseteq M_{\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^i}$ which is, by (A3), a subset of $M_{\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^{i+1}}$.

Proof of (0.5). It suffices to show by induction $\forall j \in [1, r]$:

$q(j) \quad \beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^i \vee \beta_1 \vee \dots \vee \beta_j \notin \vdash(\Gamma)$.

We will show:

(0.5.0) $M_{\Gamma, \vdash(\Gamma), \neg(\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^i)} \subseteq M_{\beta_1}$.

Then, by $p_1(i)$, $p_2(i)$, (0.5.0), and ($\vdash 7$) (where α is $\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^i$ and β is β_1), $q(1)$ holds.

Now, let $j \in [1, r-1]$ and suppose $q(j)$.

Then, we will show:

(0.5.1) $M_{\Gamma, \vdash(\Gamma), \neg(\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^i \vee \beta_1 \vee \dots \vee \beta_j)} \subseteq M_{\beta_{j+1}}$.

In addition, by $p_2(i)$ and (A3), we get:

(0.5.2) $M_{\Gamma, \vdash(\Gamma)} \subseteq M_{\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^i \vee \beta_1 \vee \dots \vee \beta_j}$.

By, (0.5.1), (0.5.2), $q(j)$, and (~ 7) (where α is $\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^i \vee \beta_1 \vee \dots \vee \beta_j$ and β is β_{j+1}), we get that $q(j+1)$ holds.

Proof of (0.5.0). Let $v \in M_{\Gamma, \sim(\Gamma), \neg(\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^i)}$. Then, $v \in M_{\neg(\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^i)}$. Thus, by $p_3(i)$, $v \notin M_\Gamma^1 \cup \dots \cup M_\Gamma^i$. Therefore, $v \in M_{\Gamma, \sim(\Gamma)} \setminus M_\Gamma^1 \cup \dots \cup M_\Gamma^i \subseteq M_{\beta_1}$.

Proof of (0.5.1). Let $v \in M_{\Gamma, \sim(\Gamma), \neg(\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^i \vee \beta_1 \vee \dots \vee \beta_j)}$. Then, by (A3), $v \in M_{\neg(\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^i)}$. Therefore, by $p_3(i)$, $v \notin M_\Gamma^1 \cup \dots \cup M_\Gamma^i$. Therefore, $v \in M_{\Gamma, \sim(\Gamma)} \setminus M_\Gamma^1 \cup \dots \cup M_\Gamma^i \subseteq M_{\beta_{j+1}}$.

Proof of (0.6). By $p_2(i)$ and (A3), we get $M_{\Gamma, \sim(\Gamma)} \subseteq M_{\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^i} \subseteq M_{\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^i \vee \beta_1 \vee \dots \vee \beta_r}$. In addition, by (A3), we get $M_{\neg(\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^i \vee \beta_1 \vee \dots \vee \beta_r)} = M_{\neg(\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^{i+1})}$. Therefore, by (0.5) and (~ 6) (where α is $\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^{i+1}$ and β is $\beta_\Gamma^1 \vee \dots \vee \beta_\Gamma^i \vee \beta_1 \vee \dots \vee \beta_r$), we get that $p_1(i+1)$ holds.

Proof of (3). Suppose $M'_\Gamma \neq \emptyset$, $v \in M_{\Gamma, \sim(\Gamma)} \setminus M'_\Gamma$, and $v \notin M_{\neg\beta_\Gamma}$. Then, by (0), (1), and definition of M_Γ^i , we get $v \in M_\Gamma^{n(\Gamma)+1}$, which is impossible by Lemma 39 (6).

Proof of (4). Case 1: $M'_\Gamma \neq \emptyset$.
By (3), we get one direction: $M_{\Gamma, \sim(\Gamma)} \setminus M'_\Gamma \subseteq M_{\Gamma, \sim(\Gamma), \neg\beta_\Gamma}$.
By (2), we get the other direction: $M_{\Gamma, \sim(\Gamma), \neg\beta_\Gamma} \subseteq M_{\Gamma, \sim(\Gamma)} \setminus M'_\Gamma$.
Case 2: $M'_\Gamma = \emptyset$.
Then, obviously, $M_{\Gamma, \sim(\Gamma)} \setminus M'_\Gamma = M_{\Gamma, \sim(\Gamma)} = M_{\Gamma, \sim(\Gamma), F(\Gamma)}$.

Proof of (5). Direction: “ \subseteq ”.
Case 1: $M'_\Gamma \neq \emptyset$.
Suppose the contrary of what we want to show, i.e. suppose $\exists \alpha \in \sim(\Gamma)$, $\alpha \notin T_d(M_{\Gamma, \sim(\Gamma), \neg\beta_\Gamma})$. Then, $M_{\Gamma, \sim(\Gamma), \neg\beta_\Gamma} \subseteq M_{\sim(\Gamma)} \subseteq M_\alpha$. Thus, $M_{\Gamma, \sim(\Gamma), \neg\beta_\Gamma} \subseteq M_{\neg\alpha}$. Consequently, by (0), (1), and (~ 6) , we get $\alpha \notin \sim(\Gamma)$, which is impossible.
Case 2: $M'_\Gamma = \emptyset$.
Let $\alpha \in \sim(\Gamma)$. Then, $M_{\Gamma, \sim(\Gamma)} \subseteq M_{\sim(\Gamma)} \subseteq M_\alpha$. In addition, by (~ 8) , $M_{\Gamma, \sim(\Gamma)} \not\subseteq M_{\neg\alpha}$. Consequently, $\alpha \in T_d(M_{\Gamma, \sim(\Gamma)}) = T_d(M_{\Gamma, \sim(\Gamma), F(\Gamma)})$.
Direction: “ \supseteq ”. Obvious by (4) and Lemma 39 (8).

Proof of (6). Direction: “ \subseteq ”.
Case 1: $M'_\Gamma = \emptyset$.
Case 1.1: $H_1(\Gamma) \neq \emptyset$.
Then, $\exists \alpha \in \mathcal{F}$, $\alpha \notin \sim(\Gamma)$, $M_{\Gamma, \sim(\Gamma)} \subseteq M_\alpha$, and $M_{\Gamma, \sim(\Gamma)} \not\subseteq M_{\neg\alpha}$. Thus, $\alpha \in T_d(M_{\Gamma, \sim(\Gamma)})$. Therefore, by (5), $\alpha \in \sim(\Gamma)$, which is impossible.
Case 1.2: $H_1(\Gamma) = \emptyset$.
Clearly, $\forall i \in \mathbb{N}^+$, if $H_i(\Gamma) = \emptyset$, then $H_{i+1}(\Gamma) = \emptyset$. Therefore, $H(\Gamma) = \emptyset = F(\Gamma)$.
Case 2: $M'_\Gamma \neq \emptyset$.
As, $M'_\Gamma \subseteq M_{\Gamma, \sim(\Gamma)}$, we get, by (2), $M_{\Gamma, \sim(\Gamma)} \not\subseteq M_{\neg\beta_\Gamma}$.
Thus, by (0) and (1), we get $\neg\beta_\Gamma \in H_1(\Gamma) \subseteq H(\Gamma)$. Therefore, $M_{H(\Gamma)} \subseteq M_{F(\Gamma)}$.
Direction: “ \supseteq ”.
Case 1: $M'_\Gamma = \emptyset$.
Verbatim the proof of Case 1 of direction “ \subseteq ”.

Case 2: $M'_\Gamma \neq \emptyset$.

Then, the following holds:

$$(6.0) \quad \forall i \in \mathbb{N}^+, M_{\Gamma, \sim(\Gamma), \neg\beta_\Gamma} \subseteq M_{\Gamma, \sim(\Gamma), H_1(\Gamma), \dots, H_i(\Gamma)}.$$

Now, suppose the contrary of what we want to show, i.e. suppose

$\exists v \in M_{\Gamma, \sim(\Gamma), \neg\beta_\Gamma}, v \notin M_{\Gamma, \sim(\Gamma), H(\Gamma)}$. Then, $v \notin M_{H(\Gamma)}$. But, clearly, $M_{H(\Gamma)} = \bigcap_{i \in \mathbb{N}^+} M_{H_i(\Gamma)}$. Therefore, $\exists i \in \mathbb{N}^+, v \notin M_{H_i(\Gamma)}$, which is impossible by (6.0).

Proof of (6.0). We show by induction: $\forall i \in \mathbb{N}^+$,

$$p(i) \quad M_{\Gamma, \sim(\Gamma), \neg\beta_\Gamma} \subseteq M_{\Gamma, \sim(\Gamma), H_1(\Gamma), \dots, H_i(\Gamma)}.$$

We will show

$$(6.0.0) \quad p(1) \text{ holds.}$$

Let $i \in \mathbb{N}^+$, suppose $p(i)$ holds, and suppose $p(i+1)$ does not hold.

Then, $\exists v \in M_{\Gamma, \sim(\Gamma), \neg\beta_\Gamma}, v \notin M_{\Gamma, \sim(\Gamma), H_1(\Gamma), \dots, H_{i+1}(\Gamma)}$.

Thus, $\exists j \in [1, i+1], v \notin M_{H_j(\Gamma)}$.

Case 1: $j = 1$.

Then, $\exists \beta \in \mathcal{F}, M_{\Gamma, \sim(\Gamma)} \subseteq M_\beta, \beta \notin \sim(\Gamma)$, and $v \notin M_{\neg\beta}$.

Thus $v \in M_\Gamma^1 \cap M_{\neg\beta_\Gamma}$, which is impossible by (2).

Case 2: $j \geq 2$.

Then, $\exists \beta \in \mathcal{F}, M_{\Gamma, \sim(\Gamma), H_1(\Gamma), \dots, H_{j-1}(\Gamma)} \subseteq M_\beta, \beta \notin \sim(\Gamma)$, and $v \notin M_{\neg\beta}$.

But, by Lemma 39 (7), by (4), and $p(i)$, we get

$$M_{\Gamma, \sim(\Gamma)} \setminus M_\Gamma^1 \cup \dots \cup M_\Gamma^{n(\Gamma)} = M_{\Gamma, \sim(\Gamma)} \setminus M'_\Gamma = M_{\Gamma, \sim(\Gamma), \neg\beta_\Gamma} \subseteq M_{\Gamma, \sim(\Gamma), H_1(\Gamma), \dots, H_i(\Gamma)} \subseteq M_{\Gamma, \sim(\Gamma), H_1(\Gamma), \dots, H_{j-1}(\Gamma)} \subseteq M_\beta.$$

Therefore, $v \in M_\Gamma^{n(\Gamma)+1}$, which is impossible by Lemma 39 (6).

Proof of (6.0.0). Suppose the contrary of what we want to show, i.e.

suppose $\exists v \in M_{\Gamma, \sim(\Gamma), \neg\beta_\Gamma}, v \notin M_{\Gamma, \sim(\Gamma), H_1(\Gamma)}$.

Then, $v \notin M_{H_1(\Gamma)}$. Thus, $\exists \beta \in \mathcal{F}, M_{\Gamma, \sim(\Gamma)} \subseteq M_\beta, \beta \notin \sim(\Gamma)$, and $v \notin M_{\neg\beta}$.

Thus $v \in M_\Gamma^1$. Therefore, $v \in M'_\Gamma \cap M_{\neg\beta_\Gamma}$, which is impossible by (2).

Proof of (7). Obvious by (5) and (6). ■

We turn to a second important lemma. Its main purpose is to show that any DP choice function μ representing (in the discriminative manner) a relation \sim satisfies the following remarkable equality: $\mu(M_\Gamma) = M_{\Gamma, \sim(\Gamma), H(\Gamma)}$, which enables us to define μ from \sim .

Lemma 41 Suppose \mathcal{L} is a language, \neg a unary connective of \mathcal{L} , \vee and \wedge binary connectives of \mathcal{L} , \mathcal{F} the set of all wffs of \mathcal{L} , $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ a semantic structure satisfying (A1) and (A3), $\mathbf{V} \subseteq \mathcal{P}(\mathcal{V})$, μ a DP choice function from \mathbf{D} to \mathbf{V} , \sim the relation on $\mathcal{P}(\mathcal{F}) \times \mathcal{F}$ such that $\forall \Gamma \subseteq \mathcal{F}, \sim(\Gamma) = T_d(\mu(M_\Gamma))$, and $\Gamma \subseteq \mathcal{F}$. Then:

$$(0) \quad \mu(M_\Gamma) \subseteq M_{\Gamma, \sim(\Gamma)};$$

$$(1) \quad \sim \text{ satisfies } (\sim 6);$$

$$(2) \quad \sim \text{ satisfies } (\sim 7);$$

$$(3) \quad \sim \text{ satisfies } (\sim 8);$$

$$(4) \quad M'_\Gamma \cap \mu(M_\Gamma) = \emptyset;$$

- (5) $M_{\Gamma, \vdash(\Gamma), T_c(\mu(M_\Gamma))} = \mu(M_\Gamma)$;
(6) if $M'_\Gamma \neq \emptyset$, then $M_{\Gamma, \vdash(\Gamma), H(\Gamma)} = \mu(M_\Gamma)$.

If $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ satisfies (A2) too, then:

- (7) if $M'_\Gamma = \emptyset$, then $M_{G(\Gamma)} = M_{T_c(\mu(M_\Gamma))}$;
(8) if $M'_\Gamma = \emptyset$, then $M_{\Gamma, \vdash(\Gamma)} \subseteq M_{G(\Gamma)}$;
(9) $M_{\Gamma, \vdash(\Gamma), H(\Gamma)} = \mu(M_\Gamma)$.

If μ is coherency preserving, then again:

- (10) $M_{\Gamma, \vdash(\Gamma), H(\Gamma)} = \mu(M_\Gamma)$.

Proof *Proof of (0).* We show $\mu(M_\Gamma) \subseteq M_{\vdash(\Gamma)}$. Let $v \in \mu(M_\Gamma)$ and $\alpha \in \vdash(\Gamma)$. Then, $\alpha \in T_d(\mu(M_\Gamma))$. Thus, $\mu(M_\Gamma) \subseteq M_\alpha$. Thus, $v \in M_\alpha$ and we are done. In addition, obviously, $\mu(M_\Gamma) \subseteq M_\Gamma$. Therefore, $\mu(M_\Gamma) \subseteq M_\Gamma \cap M_{\vdash(\Gamma)} = M_{\Gamma, \vdash(\Gamma)}$.

Proof of (1). Let $\alpha, \beta \in \mathcal{F}$ and suppose $\beta \in \vdash(\Gamma, \vdash(\Gamma)) \setminus \vdash(\Gamma)$ and $\neg\alpha \in \vdash(\Gamma, \vdash(\Gamma), \neg\beta)$. Then, by (0), $\mu(M_\Gamma) \subseteq M_{\Gamma, \vdash(\Gamma)} \subseteq M_\beta$. But, $\beta \notin \vdash(\Gamma) = T_d(\mu(M_\Gamma))$. Thus, $\mu(M_\Gamma) \subseteq M_{\neg\beta}$. Consequently, $\mu(M_\Gamma) \subseteq M_{\Gamma, \vdash(\Gamma), \neg\beta} \subseteq M_{\neg\alpha}$. Therefore, $\alpha \notin T_d(\mu(M_\Gamma)) = \vdash(\Gamma)$.

Proof of (2). Let $\alpha, \beta \in \mathcal{F}$ and suppose $\alpha \in \vdash(\Gamma, \vdash(\Gamma)) \setminus \vdash(\Gamma)$ and $\beta \in \vdash(\Gamma, \vdash(\Gamma), \neg\alpha) \setminus \vdash(\Gamma)$. Then, by (0), $\mu(M_\Gamma) \subseteq M_{\Gamma, \vdash(\Gamma)} \subseteq M_\alpha$. But, $\alpha \notin T_d(\mu(M_\Gamma))$. Thus, $\mu(M_\Gamma) \subseteq M_{\neg\alpha}$. Thus, $\mu(M_\Gamma) \subseteq M_{\Gamma, \vdash(\Gamma), \neg\alpha} \subseteq M_\beta$. But, $\beta \notin T_d(\mu(M_\Gamma))$. Therefore $\mu(M_\Gamma) \subseteq M_{\neg\beta}$. Thus, by (A3), $\mu(M_\Gamma) \subseteq M_{\neg\alpha} \cap M_{\neg\beta} = M_{\neg(\alpha \vee \beta)}$. Consequently, $\alpha \vee \beta \notin T_d(\mu(M_\Gamma)) = \vdash(\Gamma)$.

Proof of (3). Let $\alpha \in \vdash(\Gamma)$. Then, $\alpha \in T_d(\mu(M_\Gamma))$. Thus, $\mu(M_\Gamma) \not\subseteq M_{\neg\alpha}$. Thus, by (0), $M_{\Gamma, \vdash(\Gamma)} \not\subseteq M_{\neg\alpha}$.

Proof of (4). Case 1: $M'_\Gamma = \emptyset$. Obvious.

Case 2: $M'_\Gamma \neq \emptyset$.

It is sufficient to show by induction: $\forall i \in [1, n(\Gamma)]$,

$$p(i) \quad (M_\Gamma^1 \cup \dots \cup M_\Gamma^i) \cap \mu(M_\Gamma) = \emptyset.$$

We will show:

$$(4.0) \quad p(1) \text{ holds.}$$

Let $i \in [1, n(\Gamma) - 1]$. Suppose $p(i)$. We show $p(i + 1)$.

$$\text{Case 1: } M_\Gamma^{i+1} \cap \mu(M_\Gamma) = \emptyset.$$

Then, by $p(i)$, we obviously get $p(i + 1)$.

$$\text{Case 2: } \exists v \in M_\Gamma^{i+1} \cap \mu(M_\Gamma).$$

Then, $\exists \beta \in \mathcal{F}$, $\beta \notin \vdash(\Gamma)$, $M_{\Gamma, \vdash(\Gamma)} \setminus M_\Gamma^1 \cup \dots \cup M_\Gamma^i \subseteq M_\beta$, and $v \notin M_{\neg\beta}$.

Therefore, by (0) and $p(i)$, $\mu(M_\Gamma) \subseteq M_{\Gamma, \vdash(\Gamma)} \setminus M_\Gamma^1 \cup \dots \cup M_\Gamma^i \subseteq M_\beta$. But, $\mu(M_\Gamma) \not\subseteq M_{\neg\beta}$.

Consequently, $\beta \in T_d(\mu(M_\Gamma)) = \vdash(\Gamma)$, which is impossible.

Proof of (4.0). Suppose the contrary of $p(1)$, i.e. suppose $\exists v \in M_\Gamma^1 \cap \mu(M_\Gamma)$.

Then, $\exists \beta \in \mathcal{F}$, $\beta \notin \vdash(\Gamma)$, $M_{\Gamma, \vdash(\Gamma)} \subseteq M_\beta$ and $v \notin M_{\neg\beta}$.

Therefore, by (0), $\mu(M_\Gamma) \subseteq M_\beta$. On the other hand, $\mu(M_\Gamma) \not\subseteq M_{\neg\beta}$.

Therefore, $\beta \in T_d(\mu(M_\Gamma)) = \vdash(\Gamma)$, which is impossible.

Proof of (5). As $\mu(M_\Gamma) \in \mathbf{D}$, $\exists \Gamma' \subseteq \mathcal{F}$, $M_{\Gamma'} = \mu(M_\Gamma)$.

Therefore, $M_{T(\mu(M_\Gamma))} = M_{T(M_{\Gamma'})} = M_{\Gamma'} = \mu(M_\Gamma)$.

Thus, $M_{\Gamma, \sim(\Gamma), T_c(\mu(M_\Gamma))} = M_{\Gamma, T_d(\mu(M_\Gamma)), T_c(\mu(M_\Gamma))} = M_{\Gamma, T(\mu(M_\Gamma))}$. But, $\Gamma \subseteq T(\mu(M_\Gamma))$.

Therefore, $M_{\Gamma, T(\mu(M_\Gamma))} = M_{T(\mu(M_\Gamma))} = \mu(M_\Gamma)$.

Proof of (6). Suppose $M'_\Gamma \neq \emptyset$. Direction: “ \subseteq ”.

Case 1: $\exists v \in M_{\Gamma, \sim(\Gamma)} \setminus M_\Gamma^1 \cup \dots \cup M_\Gamma^{n(\Gamma)}$, $v \notin M_{T_c(\mu(M_\Gamma))}$.

Then, $\exists \alpha \in T_c(\mu(M_\Gamma))$, $v \notin M_\alpha$.

By Lemma 40 (3), Lemma 39 (7), and (A3), $M_{\Gamma, \sim(\Gamma)} \setminus M_\Gamma^1 \cup \dots \cup M_\Gamma^{n(\Gamma)} \subseteq M_{\neg\beta_\Gamma} \subseteq M_{\neg(\beta_\Gamma \wedge \alpha)}$.

By (0) and Lemma 40 (1), $\mu(M_\Gamma) \subseteq M_{\beta_\Gamma} \cap M_\alpha = M_{\neg\neg(\beta_\Gamma \wedge \alpha)}$.

Therefore, $\neg(\beta_\Gamma \wedge \alpha) \notin T_d(\mu(M_\Gamma)) = \sim(\Gamma)$.

In addition, $v \notin M_\alpha \supseteq M_{\neg\neg(\beta_\Gamma \wedge \alpha)}$.

Consequently, $v \in M_\Gamma^{n(\Gamma)+1}$ (take $\neg(\beta_\Gamma \wedge \alpha)$ for the β of the definition of M_Γ^i).

Therefore, by Lemma 39 (6), we get a contradiction.

Case 2: $M_{\Gamma, \sim(\Gamma)} \setminus M_\Gamma^1 \cup \dots \cup M_\Gamma^{n(\Gamma)} \subseteq M_{T_c(\mu(M_\Gamma))}$.

Then, by Lemma 40 (6), Lemma 40 (4), Lemma 39 (7), and by (5), we get

$M_{\Gamma, \sim(\Gamma), H(\Gamma)} = M_{\Gamma, \sim(\Gamma)} \setminus M_\Gamma^1 \cup \dots \cup M_\Gamma^{n(\Gamma)} \subseteq M_{\Gamma, \sim(\Gamma), T_c(\mu(M_\Gamma))} = \mu(M_\Gamma)$.

Direction: “ \supseteq ”.

By (0), (4), Lemma 40 (4), and Lemma 40 (6), we get

$\mu(M_\Gamma) \subseteq M_{\Gamma, \sim(\Gamma)} \setminus M'_\Gamma = M_{\Gamma, \sim(\Gamma), F(\Gamma)} = M_{\Gamma, \sim(\Gamma), H(\Gamma)}$.

Proof of (7). Suppose $M'_\Gamma = \emptyset$. Direction: “ \supseteq ”.

Suppose the contrary of what we want to show, i.e. suppose $\exists v \in M_{T_c(\mu(M_\Gamma))}$, $v \notin M_{G(\Gamma)}$.

Then, $\exists \alpha \in G(\Gamma)$, $v \notin M_\alpha$.

Case 1: $\alpha \in T(M_{\Gamma, \sim(\Gamma)})$.

As $\alpha \in G(\Gamma)$, $\alpha \notin \sim(\Gamma)$. Thus, by Lemma 40 (5), $\alpha \notin T_d(M_{\Gamma, \sim(\Gamma)})$.

Therefore, $\alpha \in T_c(M_{\Gamma, \sim(\Gamma)})$. Consequently, by (0), $\alpha \in T_c(\mu(M_\Gamma))$.

Thus, $v \in M_\alpha$, which is impossible.

Case 2: $\neg\alpha \in T(M_{\Gamma, \sim(\Gamma)})$.

As $\alpha \in G(\Gamma)$, $\neg\alpha \notin \sim(\Gamma)$. Thus, by Lemma 40 (5), $\neg\alpha \notin T_d(M_{\Gamma, \sim(\Gamma)})$.

Therefore, $\neg\alpha \in T_c(M_{\Gamma, \sim(\Gamma)})$. Consequently, by (A3), $\alpha \in T_c(M_{\Gamma, \sim(\Gamma)})$.

Therefore, by (0), $\alpha \in T_c(\mu(M_\Gamma))$. Thus, $v \in M_\alpha$, which is impossible.

Case 3: $\alpha \notin T(M_{\Gamma, \sim(\Gamma)})$ and $\neg\alpha \notin T(M_{\Gamma, \sim(\Gamma)})$.

Then, by (A2), $M_{\Gamma, \sim(\Gamma), \alpha} \not\subseteq M_{\neg\alpha}$. Therefore, $\alpha \in T_d(M_{\Gamma, \sim(\Gamma), \alpha})$.

But, $\alpha \in G(\Gamma)$. Thus, $T_d(M_{\Gamma, \sim(\Gamma), \alpha}) \subseteq \sim(\Gamma)$. Thus, $\alpha \in \sim(\Gamma)$. Thus, $\alpha \notin G(\Gamma)$, impossible.

Direction: “ \subseteq ”.

Suppose the contrary of what we want to show, i.e. suppose $\exists v \in M_{G(\Gamma)}$, $v \notin M_{T_c(\mu(M_\Gamma))}$.

Then, we will show:

(7.0) $\exists \alpha \in T_c(\mu(M_\Gamma))$, $|M_{\Gamma, \sim(\Gamma), \alpha}| < |\mu(M_\Gamma)|$

But, $\mu(M_\Gamma) \subseteq M_\alpha$ and, by (0), $\mu(M_\Gamma) \subseteq M_{\Gamma, \sim(\Gamma)}$. Therefore, $\mu(M_\Gamma) \subseteq M_{\Gamma, \sim(\Gamma), \alpha}$.

Thus, $|\mu(M_\Gamma)| \leq |M_{\Gamma, \sim(\Gamma), \alpha}|$, which is impossible by (7.0).

Proof of (7.0). We have $\exists \delta \in T_c(\mu(M_\Gamma))$, $v \notin M_\delta$.

By (A1), $|M_{\Gamma, \sim(\Gamma), \delta}|$ is finite. To show (7.0), it suffices to show by induction (in the decreasing direction): $\forall i \in \mathbf{Z}$ with $i \leq |M_{\Gamma, \sim(\Gamma), \delta}|$,

$p(i) \exists \alpha \in T_c(\mu(M_\Gamma)), v \notin M_\alpha$ and $|M_{\Gamma, \sim(\Gamma), \alpha}| - |\mu(M_\Gamma)| \leq i$.

Obviously, $p(|M_{\Gamma, \sim(\Gamma), \delta}|)$ holds (take δ).

Let $i \in \mathbb{Z}$ with $i \leq |M_{\Gamma, \sim(\Gamma), \delta}|$ and suppose $p(i)$ holds. We show $p(i-1)$.

We have $\exists \alpha \in T_c(\mu(M_\Gamma)), v \notin M_\alpha$ and $|M_{\Gamma, \sim(\Gamma), \alpha}| - |\mu(M_\Gamma)| \leq i$.

Case 1: $T_d(M_{\Gamma, \sim(\Gamma), \alpha}) \subseteq \sim(\Gamma)$.

As $\alpha \in T_c(\mu(M_\Gamma))$ and (A3) holds, we get $\neg\alpha \in T_c(\mu(M_\Gamma))$.

But, $T_c(\mu(M_\Gamma)) \cap T_d(\mu(M_\Gamma)) = \emptyset$. Thus, neither α nor $\neg\alpha$ belongs to $T_d(\mu(M_\Gamma)) = \sim(\Gamma)$.

Consequently, $\alpha \in G(\Gamma)$. Thus, $v \in M_\alpha$, which is impossible.

Case 2: $\exists \beta \in T_d(M_{\Gamma, \sim(\Gamma), \alpha}), \beta \notin \sim(\Gamma)$.

By (0), $\mu(M_\Gamma) \subseteq M_{\Gamma, \sim(\Gamma)}$. On the other hand, $\mu(M_\Gamma) \subseteq M_\alpha$. Thus, $\mu(M_\Gamma) \subseteq M_{\Gamma, \sim(\Gamma), \alpha} \subseteq M_\beta$.

But, $\beta \notin \sim(\Gamma) = T_d(\mu(M_\Gamma))$. Therefore, $\mu(M_\Gamma) \subseteq M_{-\beta}$.

Consequently, $\mu(M_\Gamma) \subseteq M_\alpha \cap M_{-\beta} = M_{\alpha \wedge \neg\beta}$ and $\mu(M_\Gamma) \subseteq M_{-\alpha} \subseteq M_{\neg(\alpha \wedge \neg\beta)}$.

Therefore, $\alpha \wedge \neg\beta \in T_c(\mu(M_\Gamma))$.

Moreover, $v \notin M_\alpha \supseteq M_{\alpha \wedge \neg\beta}$.

In addition, $M_{\Gamma, \sim(\Gamma), \alpha \wedge \neg\beta} \subseteq M_{\Gamma, \sim(\Gamma), \alpha}$, whilst $M_{\Gamma, \sim(\Gamma), \alpha} \not\subseteq M_{-\beta} \supseteq M_{\Gamma, \sim(\Gamma), \alpha \wedge \neg\beta}$.

Thus $|M_{\Gamma, \sim(\Gamma), \alpha \wedge \neg\beta}| \leq |M_{\Gamma, \sim(\Gamma), \alpha}| - 1$. Thus, $|M_{\Gamma, \sim(\Gamma), \alpha \wedge \neg\beta}| - |\mu(M_\Gamma)| \leq i - 1$.

Therefore, $p(i-1)$ holds (take $\alpha \wedge \neg\beta$).

Proof of (8). Suppose $M'_\Gamma = \emptyset$.

Now, suppose the contrary of what we want to show, i.e. $\exists v \in M_{\Gamma, \sim(\Gamma)}, v \notin M_{G(\Gamma)}$.

Then, $\exists \alpha \in G(\Gamma), v \notin M_\alpha$.

Case 1: $\alpha \in T(M_{\Gamma, \sim(\Gamma)})$.

As, $\alpha \in G(\Gamma), \alpha \notin \sim(\Gamma)$. Therefore, by Lemma 40 (5), $\alpha \notin T_d(M_{\Gamma, \sim(\Gamma)})$.

Thus, $\alpha \in T_c(M_{\Gamma, \sim(\Gamma)})$. Therefore, $M_{\Gamma, \sim(\Gamma)} \subseteq M_\alpha$. Consequently, $v \in M_\alpha$, which is impossible.

Case 2: $\neg\alpha \in T(M_{\Gamma, \sim(\Gamma)})$.

As, $\alpha \in G(\Gamma), \neg\alpha \notin \sim(\Gamma)$. Therefore, by Lemma 40 (5), $\neg\alpha \notin T_d(M_{\Gamma, \sim(\Gamma)})$.

Thus, $\neg\alpha \in T_c(M_{\Gamma, \sim(\Gamma)})$. Therefore, by (A3), $M_{\Gamma, \sim(\Gamma)} \subseteq M_{\neg\neg\alpha} = M_\alpha$.

Consequently, $v \in M_\alpha$, which is impossible.

Case 3: $\alpha \notin T(M_{\Gamma, \sim(\Gamma)})$ and $\neg\alpha \notin T(M_{\Gamma, \sim(\Gamma)})$.

Then, by (A2), $M_{\Gamma, \sim(\Gamma), \alpha} \not\subseteq M_{-\alpha}$. Thus, $\alpha \in T_d(M_{\Gamma, \sim(\Gamma), \alpha})$. But, $\alpha \in G(\Gamma)$. Thus, $\alpha \notin \sim(\Gamma)$.

Therefore, $T_d(M_{\Gamma, \sim(\Gamma), \alpha}) \not\subseteq \sim(\Gamma)$. Consequently, $\alpha \notin G(\Gamma)$, which is impossible.

Proof of (9). Case 1: $M'_\Gamma = \emptyset$.

By Lemma 40 (6), $M_{\Gamma, \sim(\Gamma), H(\Gamma)} = M_{\Gamma, \sim(\Gamma), F(\Gamma)} = M_{\Gamma, \sim(\Gamma)}$.

But, by (8), (7), and (5), $M_{\Gamma, \sim(\Gamma)} = M_{\Gamma, \sim(\Gamma), G(\Gamma)} = M_{\Gamma, \sim(\Gamma), T_c(\mu(M_\Gamma))} = \mu(M_\Gamma)$.

Case 2: $M'_\Gamma \neq \emptyset$. Obvious by (6).

Proof of (10).

Case 1: $M'_\Gamma = \emptyset$.

Case 1.1: $\exists v \in M_{\Gamma, \sim(\Gamma)}, v \notin M_{T_c(\mu(M_\Gamma))}$.

Case 1.1.1: Γ is not consistent.

Then, $\exists \alpha \in T_c(\mu(M_\Gamma)), v \notin M_\alpha$ and, as Γ is not consistent, $\exists \beta \in \mathcal{F}, M_\Gamma \subseteq M_\beta$ and $M_\Gamma \subseteq M_{-\beta}$.

We have $M_{\Gamma, \sim(\Gamma)} \subseteq M_\Gamma \subseteq M_\beta \subseteq M_{\beta \vee \neg\alpha}$.

Moreover, $\mu(M_\Gamma) \subseteq M_\Gamma \subseteq M_{-\beta}$. Thus, $\mu(M_\Gamma) \subseteq M_{-\beta} \cap M_\alpha = M_{\neg(\beta \vee \neg\alpha)}$.

Therefore, $\beta \vee \neg\alpha \notin T_d(\mu(M_\Gamma)) = \sim(\Gamma)$.

In addition, $v \notin M_\alpha \supseteq M_{\neg(\beta \vee \neg\alpha)}$.

Consequently, $v \in M_\Gamma^1$ (take $\beta \vee \neg\alpha$ for the β of the definition of M_Γ^1).

Thus, $v \in M'_\Gamma$, which is impossible.

Case 1.1.2: Γ is consistent.

Thus, $M_\Gamma \in \mathbf{C}$. Therefore, as μ is coherency preserving, $\mu(M_\Gamma) \in \mathbf{C}$. Thus, $T_c(\mu(M_\Gamma)) = \emptyset$.

Therefore, $M_{T_c(\mu(M_\Gamma))} = \mathcal{V}$. Thus, $v \in M_{T_c(\mu(M_\Gamma))}$, which is impossible.

Case 1.2: $M_{\Gamma, \sim(\Gamma)} \subseteq M_{T_c(\mu(M_\Gamma))}$.

Then, by Lemma 40 (6), $M_{\Gamma, \sim(\Gamma), H(\Gamma)} = M_{\Gamma, \sim(\Gamma), F(\Gamma)} = M_{\Gamma, \sim(\Gamma)} = M_{\Gamma, \sim(\Gamma), T_c(\mu(M_\Gamma))}$.

Therefore, by (5), $M_{\Gamma, \sim(\Gamma), H(\Gamma)} = \mu(M_\Gamma)$.

Case 2: $M'_\Gamma \neq \emptyset$. Obvious by (6). ■

Now comes the proof of **Proposition 37** (which is stated at the beginning of Section 3.3).

Proof *Proof of (0)*. Direction: “ \rightarrow ”.

There exists a CP DP coherent choice function μ from \mathbf{D} to $\mathcal{P}(\mathcal{V})$ such that

$\forall \Gamma \subseteq \mathcal{F}$, $\sim(\Gamma) = T_d(\mu(M_\Gamma))$.

We will show:

(0.0) \sim satisfies (~ 0).

By Lemma 41 (1), (2), and (3), \sim satisfies (~ 6), (~ 7), and (~ 8).

By Lemma 41 (10) and Coherence of μ , \sim satisfies (~ 9).

We will show:

(0.1) \sim satisfies (~ 11).

Direction: “ \leftarrow ”.

Suppose \sim satisfies (~ 0), (~ 6), (~ 7), (~ 8), (~ 9), and (~ 11).

Then, let μ be the function from \mathbf{D} to $\mathcal{P}(\mathcal{V})$ such that $\forall \Gamma \subseteq \mathcal{F}$, $\mu(M_\Gamma) = M_{\Gamma, \sim(\Gamma), H(\Gamma)}$.

We will show:

(0.2) μ is well-defined.

Clearly, μ is a DP choice function.

In addition, as \sim satisfies (~ 9), μ is coherent.

We will show:

(0.3) μ is CP.

And finally, by Lemma 40 (7), $\forall \Gamma \subseteq \mathcal{F}$, $\sim(\Gamma) = T_d(\mu(M_\Gamma))$.

Proof of (0.0). Let $\Gamma, \Delta \subseteq \mathcal{F}$ and suppose $\vdash(\Gamma) = \vdash(\Delta)$. Then, $M_\Gamma = M_\Delta$.

Therefore, $\sim(\Gamma) = T_d(\mu(M_\Gamma)) = T_d(\mu(M_\Delta)) = \sim(\Delta)$.

Proof of (0.1). Let $\Gamma \subseteq \mathcal{F}$ and suppose Γ is consistent.

Then, $M_\Gamma \in \mathbf{D} \cap \mathbf{C}$. Thus, as μ is CP, $\mu(M_\Gamma) \in \mathbf{C}$. Therefore, $T_d(\mu(M_\Gamma)) = T(\mu(M_\Gamma))$.

Consequently, $\Gamma \subseteq T(M_\Gamma) \subseteq T(\mu(M_\Gamma)) = T_d(\mu(M_\Gamma)) = \sim(\Gamma)$.

In addition, $M_{\sim(\Gamma)} = M_{T_d(\mu(M_\Gamma))} = M_{T(\mu(M_\Gamma))}$. But, $\mu(M_\Gamma) \in \mathbf{C}$. Thus, $M_{T(\mu(M_\Gamma))} \in \mathbf{C}$.

Consequently, $\sim(\Gamma)$ is consistent.

And finally, $\sim(\Gamma) = T_d(\mu(M_\Gamma)) = T(\mu(M_\Gamma)) = T(M_{T(\mu(M_\Gamma))}) = T(M_{\sim(\Gamma)}) = \vdash(\sim(\Gamma))$.

Proof of (0.2). Let $\Gamma, \Delta \subseteq \mathcal{F}$ and suppose $M_\Gamma = M_\Delta$.

Then, $\vdash(\Gamma) = \vdash(\Delta)$. Thus, by (~ 0), $\sim(\Gamma) = \sim(\Delta)$.

Consequently, $H(\Gamma) = H(\Delta)$. Therefore, $M_{\Gamma, \sim(\Gamma), H(\Gamma)} = M_{\Delta, \sim(\Delta), H(\Delta)}$.

Proof of (0.3). Suppose $V \in \mathbf{D} \cap \mathbf{C}$. Then, $\exists \Gamma \subseteq \mathcal{F}$, $V = M_\Gamma$.

Case 1: $H_1(\Gamma) \neq \emptyset$.

Thus, $\exists \beta \in \mathcal{F}$, $\beta \notin \sim(\Gamma)$ and $M_{\Gamma, \sim(\Gamma)} \subseteq M_\beta$.

By (\sim 11), $\Gamma \subseteq \sim(\Gamma)$ and $\vdash(\sim(\Gamma)) = \sim(\Gamma)$. Thus, $M_{\Gamma, \sim(\Gamma)} = M_{\sim(\Gamma)}$. Thus, $M_{\sim(\Gamma)} \subseteq M_{\beta}$. Therefore, $\beta \in T(M_{\sim(\Gamma)}) = \vdash(\sim(\Gamma)) = \sim(\Gamma)$, which is impossible.

Case 2: $H_1(\Gamma) = \emptyset$.

Then, $H(\Gamma) = \emptyset$. Thus, $\mu(V) = \mu(M_{\Gamma}) = M_{\Gamma, \sim(\Gamma), H(\Gamma)} = M_{\sim(\Gamma)}$.

But, by (\sim 11), $\sim(\Gamma)$ is consistent. Therefore, $M_{\sim(\Gamma)} \in \mathbf{C}$.

proof of (1). Direction: “ \rightarrow ”.

Verbatim the proof of (0), except that in addition μ is LM.

Then, by Lemma 41 (10) and LM, \sim satisfies (\sim 10).

Direction: “ \leftarrow ”.

Verbatim the proof of (0), except that in addition \sim satisfies (\sim 10).

Then, by definition of μ and (\sim 10), μ is LM.

Proof of (2). Direction: “ \rightarrow ”.

Verbatim the proof of (0), except that μ is no longer CP, whilst (A2) now holds.

Note that, in (0), CP was used only to show (\sim 11) and (\sim 9).

But, (\sim 11) is no longer required to hold.

In addition, by Lemma 41 (9) and Coherence of μ , (\sim 9) holds.

Direction: “ \leftarrow ”.

Verbatim the proof of (0), except that (\sim 11) does no longer hold, whilst (A2) now holds.

However, in (0), (\sim 11) was used only to show that μ is CP, which is no longer required.

Note that we do not need to use (A2) in this direction.

Proof of (3). Direction “ \rightarrow ”.

Verbatim the proof of (0), except that μ is no longer CP, whilst μ is now LM and (A2) now holds.

Note that, in (0), CP was used only to show (\sim 11) and (\sim 9).

But, (\sim 11) is no longer required.

In addition, by Lemma 41 (9) and Coherence of μ , (\sim 9) holds.

Similarly, by Lemma 41 (9) and Local Monotonicity of μ , (\sim 10) holds.

Direction: “ \leftarrow ”.

Verbatim the proof of (0), except that (\sim 11) does no longer hold, whilst (\sim 10) and (A2) now holds.

Note that, in (0), (\sim 11) was used only to show that μ is CP, which is no longer required.

Now, by definition of μ and by (\sim 10), μ is LM.

Note that we do not need to use (A2) in this direction. ■

3.4 The discriminative and not necessarily definability preserving case

Unlike in Section 3.3, the conditions of this section will not be purely syntactic. The translation of properties like Coherence in syntactic terms is blocked because we do no longer have the following useful equality: $\mu(M_{\Gamma}) = M_{\Gamma, \sim(\Gamma), H(\Gamma)}$, which holds when the choice functions under consideration are definability preserving (but this is not the case here). Thanks to Lemmas 30 and 31 (stated in Section 3.2), we will provide a solution with semi-syntactic conditions.

Notation 42 Let \mathcal{L} be a language, \neg a unary connective of \mathcal{L} , \mathcal{F} the set of all wffs of \mathcal{L} , $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ a semantic structure, and \sim a relation on $\mathcal{P}(\mathcal{F}) \times \mathcal{F}$.

Then, consider the following condition: $\forall \Gamma \subseteq \mathcal{F}$,

(\sim 12) $\vdash(\Gamma, \sim(\Gamma), H(\Gamma)) = T(\{v \in M_{\Gamma} : \forall \Delta \subseteq \mathcal{F}, \text{ if } v \in M_{\Delta} \subseteq M_{\Gamma}, \text{ then } v \in M_{\sim(\Delta), H(\Delta)}\})$.

Proposition 43 Let \mathcal{L} be a language, \neg a unary connective of \mathcal{L} , \vee and \wedge binary connectives of \mathcal{L} , \mathcal{F} the set of all wffs of \mathcal{L} , $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ a semantic structure satisfying (A1) and (A3), and \sim a relation on $\mathcal{P}(\mathcal{F}) \times \mathcal{F}$. Then,

(0) \sim is a CP preferential-discriminative consequence relation iff (~ 0) , (~ 6) , (~ 7) , (~ 8) , (~ 11) and (~ 12) hold.

Suppose $\langle \mathcal{F}, \mathcal{V}, \models \rangle$ satisfies (A2) too. Then,

(1) \sim is a preferential-discriminative consequence relation iff (~ 0) , (~ 6) , (~ 7) , (~ 8) , and (~ 12) hold.

Proof *Proof of (1). Direction: “ \rightarrow ”.*

There exists a coherent choice function μ from \mathbf{D} to $\mathcal{P}(\mathcal{V})$ such that $\forall \Gamma \subseteq \mathcal{F}$, $\sim(\Gamma) = T_d(\mu(M_\Gamma))$. Then, \sim satisfies obviously (~ 0) .

Let f be the function from \mathbf{D} to \mathbf{D} such that $\forall V \in \mathbf{D}$, $f(V) = M_{T(\mu(V))}$.

Then, by Lemma 31, $\forall V \in \mathbf{D}$, $f(V) = M_{T(\mu_f(V))}$.

Moreover, $\forall \Gamma \subseteq \mathcal{F}$, $f(M_\Gamma) = M_{T(\mu(M_\Gamma))} \subseteq M_{T(M_\Gamma)} = M_\Gamma$.

Therefore, f is a choice function.

Obviously, f is DP.

In addition, $\forall \Gamma \subseteq \mathcal{F}$, $\sim(\Gamma) = T_d(\mu(M_\Gamma)) = T_d(M_{T(\mu(M_\Gamma))}) = T_d(f(M_\Gamma))$.

Consequently, by Lemma 41 (1), (2), and (3), \sim satisfies (~ 6) , (~ 7) , and (~ 8) .

In addition, by Lemma 41 (9), $\forall \Gamma \subseteq \mathcal{F}$, $f(M_\Gamma) = M_{\Gamma, \sim(\Gamma), H(\Gamma)}$.

We show that \sim satisfies (~ 12) . Let $\Gamma \subseteq \mathcal{F}$.

Then, $\vdash(\Gamma, \sim(\Gamma), H(\Gamma)) = T(M_{\Gamma, \sim(\Gamma), H(\Gamma)}) = T(f(M_\Gamma)) = T(M_{T(\mu_f(M_\Gamma))}) = T(\mu_f(M_\Gamma)) = T(\{v \in M_\Gamma : \forall W \in \mathbf{D}, \text{if } v \in W \subseteq M_\Gamma, \text{ then } v \in f(W)\}) = T(\{v \in M_\Gamma : \forall \Delta \subseteq \mathcal{F}, \text{if } v \in M_\Delta \subseteq M_\Gamma, \text{ then } v \in f(M_\Delta)\}) = T(\{v \in M_\Gamma : \forall \Delta \subseteq \mathcal{F}, \text{if } v \in M_\Delta \subseteq M_\Gamma, \text{ then } v \in M_{\Delta, \sim(\Delta), H(\Delta)}\}) = T(\{v \in M_\Gamma : \forall \Delta \subseteq \mathcal{F}, \text{if } v \in M_\Delta \subseteq M_\Gamma, \text{ then } v \in M_{\sim(\Delta), H(\Delta)}\})$.

Direction: “ \leftarrow ”.

Suppose (~ 0) , (~ 6) , (~ 7) , (~ 8) , and (~ 12) hold.

Let f be the function from \mathbf{D} to \mathbf{D} such that $\forall \Gamma \subseteq \mathcal{F}$, $f(M_\Gamma) = M_{\Gamma, \sim(\Gamma), H(\Gamma)}$.

By (~ 0) , f is well-defined.

By Lemma 40 (7), $\forall \Gamma \subseteq \mathcal{F}$, $\sim(\Gamma) = T_d(M_{\Gamma, \sim(\Gamma), H(\Gamma)})$.

Therefore, $\forall \Gamma \subseteq \mathcal{F}$, $\sim(\Gamma) = T_d(f(M_\Gamma))$.

By (~ 12) , $\forall \Gamma \subseteq \mathcal{F}$, $f(M_\Gamma) = M_{T(\mu_f(M_\Gamma))}$.

Therefore, $\forall \Gamma \subseteq \mathcal{F}$, $\sim(\Gamma) = T_d(f(M_\Gamma)) = T_d(M_{T(\mu_f(M_\Gamma))}) = T_d(\mu_f(M_\Gamma))$.

But, by Lemma 30, μ_f is a coherent choice function.

Proof of (0). Direction: “ \rightarrow ”.

Verbatim the proof of (1), except that (A2) does no longer hold, whilst μ is now CP.

Note that (A2) was used only to apply Lemma 41 (9) to get $\forall \Gamma \subseteq \mathcal{F}$, $f(M_\Gamma) = M_{\Gamma, \sim(\Gamma), H(\Gamma)}$.

But, we will get this equality by another mean.

Indeed, if $V \in \mathbf{D} \cap \mathbf{C}$, then, as μ is CP, $\mu(V) \in \mathbf{C}$, thus $M_{T(\mu(V))} \in \mathbf{C}$, thus $f(V) \in \mathbf{C}$.

Therefore f is CP.

Consequently, by Lemma 41 (10), we get $\forall \Gamma \subseteq \mathcal{F}$, $f(M_\Gamma) = M_{\Gamma, \sim(\Gamma), H(\Gamma)}$.

In addition, by verbatim the proof of (0.1) of Proposition 37, \sim satisfies (~ 11) .

Direction: “ \leftarrow ”.

Verbatim the proof of (1), except that (A2) does no longer hold, whilst \sim now satisfies (~ 11) .

But, in this direction, (A2) was not used in (0).

It remains to show that μ_f is CP.

By verbatim the proof of (0.3) of Proposition 37, we get that f is CP.

Let $V \in \mathbf{D} \cap \mathbf{C}$. Then, $f(V) \in \mathbf{C}$. Thus, $M_{T(\mu_f(V))} \in \mathbf{C}$. Thus, $\mu_f(V) \in \mathbf{C}$ and we are done. ■

4 Conclusion

We provided, in a general framework, characterizations for families of preferential(-discriminative) consequence relations. Note that we have been inspired by the work of Schlechta in the non-discriminative case, whilst we developed new techniques and ideas in the discriminative case. In many cases, our conditions are purely syntactic. In fact, when the choice functions under consideration are not necessarily definability preserving, we provided solutions with semi-syntactic conditions. We managed to do so thanks to Lemmas 30 and 31. An interesting thing is that we used them both in the plain and the discriminative versions. This suggests that they can be used in yet other versions. In addition, we are quite confident that Lemmas 40 and 41 can be used to characterize other families of consequence relations defined in the discriminative manner by DP choice functions (not necessarily coherent, unlike all the families investigated here).

5 Acknowledgements

I owe very much to Karl Schlechta for his hints, advice, constructive criticism, and more. I acknowledge also Arnon Avron for discussions and the referees for their help about the presentation of the paper.

References

- [AA94] O. Arieli and A. Avron. Logical bilattices and inconsistent data. In *Proc. 9th IEEE Annual Symp. on Logic in Computer Science*, pages 468–476. IEEE Press, 1994.
- [AA96] O. Arieli and A. Avron. Reasoning with logical bilattices. *Journal of Logic, Language and Information*, 5(1):25–63, 1996.
- [AA98] O. Arieli and A. Avron. The value of the four values. *Artificial Intelligence*, 102:97–141, 1998.
- [AA00] O. Arieli and A. Avron. Bilattices and paraconsistency. In D. Batens, C. Mortensen, G. Priest, and J. Van-Benedem, editors, *Frontiers in Paraconsistent Logic*, pages 11–28. Research Studies Press, 2000.
- [Aiz85] M. A. Aizerman. New problems in the general choice theory: Review of a research trend. *Social Choice and Welfare*, 2:235–282, 1985.
- [AL01a] A. Avron and I. Lev. A formula-preferential base for paraconsistent and plausible reasoning systems. In “*Inconsistency in Data and Knowledge*” workshop (KRR-4) at the 17th International Joint Conference on Artificial Intelligence (IJCAI’01), pages 60–70, 2001.

- [AL01b] A. Avron and I. Lev. Formula-preferential systems for paraconsistent non-monotonic reasoning. In *International Conference on Artificial Intelligence (ICAI'01)*, pages 816–820, 2001.
- [AM81] M. A. Aizerman and A.V. Malishevski. General theory of best variants choice: Some aspects. *IEEE Transactions on Automatic Control*, 26:1030–1040, 1981.
- [Arr59] K. J. Arrow. Rational choice functions and orderings. *Economica*, 26:121–127, 1959.
- [Avr91] A. Avron. Natural 3-valued logics : characterization and proof theory. *The Journal of Symbolic Logic*, 56(1):276–294, 1991.
- [Bat98] D. Batens. Inconsistency-adaptive logics. In Ewa Orłowska, editor, *Logic at Work. Essays dedicated to the memory of Elena Rasiowa*. Springer-Verlag, 1998.
- [Bel77a] N. D. Belnap. How computers should think. In G. Ryle, editor, *Contemporary Aspects of Philosophy*, pages 30–56. Oriel Press, 1977.
- [Bel77b] N. D. Belnap. A useful four-valued logic. In J.M. Dunn and G. Epstein, editors, *Modern Uses of Multiple-Valued Logic*, pages 7–37. Oriel Press, 1977.
- [Che54] H. Chernoff. Rational selection of decision functions. *Econometrica*, 26:121–127, 1954.
- [CLM99] W. A. Carnielli and M. Lima-Marques. Society semantics for multiple-valued logics. In W. A. Carnielli and I. M. L. D’Ottaviano, editors, *Advances in Contemporary Logic and Computer Science*, volume 235 of Contemporary Mathematics Series, pages 33–52. American Mathematical Society, 1999.
- [CMdA00] W. A. Carnielli, J. Marcos, and S. de Amo. Formal inconsistency and evolutionary databases. *Logic and Logical Philosophy*, 8:115–152, 2000.
- [dACM02] S. de Amo, W. A. Carnielli, and J. Marcos. A logical framework for integrating inconsistent information in multiple databases. In Thomas Eiter and Berlin Klaus-Dieter Schewe, Springer-Verlag, editors, *Second International Symposium on Foundations of Information and Knowledge Systems*, pages 67–84, 2002.
- [DdC70] I.M.L. D’Ottaviano and N.C.A. da Costa. Sur un problème de Jaśkowski, Sciences. In *Comptes Rendus de l’Académie des Sciences de Paris*, volume 270, pages 1349–1353, 1970.
- [Eps90] R.L. Epstein. *The Semantic foundation of logic. Vol. I: propositional logics*. Kluwer Academic Publisher, 1990.
- [Gab85] D. M. Gabbay. Theoretical foundations for non-monotonic reasoning in expert systems. In eds. K. R. Apt, editor, *Logics and Models of Concurrent Systems*, pages 439–457. Springer-Verlag, 1985.
- [Han69] B. Hansson. An analysis of some deontic logics. *Nôus*, 3:373–398, 1969. Reprinted in R. Hilpinen ed., *Deontic Logic: Introductory and Systematic Readings*. Reidel, Dordrecht 1971, pages 121-147.
- [Imi87] T. Imielinski. Results on translating defaults to circumscription. *Artificial Intelligence*, 32:131–146, 1987.

- [KL92] M. Kifer and E. L. Lozinskii. A logic for reasoning with inconsistency. *Journal of Automated Reasoning*, 9(2):179–215, 1992.
- [KLM90] S. Kraus, D. Lehmann, and M. Magidor. Nonmonotonic reasoning, preferential models, and cumulative logics. *Artificial Intelligence*, 44(1-2):167–207, 1990.
- [KM02] S. Konieczny and P. Marquis. Three-valued Logics for Inconsistency Handling. In *Proceedings of the 8th European Conference on Logics in Artificial Intelligence (JELIA'02)*, pages 332–344. Springer-Verlag, 2002.
- [Leh01] D. Lehmann. Nonmonotonic Logics and Semantics. *Journal of Logic and Computation*, 11(2):229–256, 2001.
- [Leh02] D. Lehmann. Connectives in Quantum and other Cumulative Logics. Technical report, ASL European Summer Meeting LC02, Muenster, Germany. Leibniz Center for Research in Computer Science TR-2002-28, 2002.
- [LM92] D. Lehmann and M. Magidor. What does a conditional knowledge base entail? *Artificial Intelligence*, 55(1):1–60, 1992.
- [Mak89] D. Makinson. General theory of cumulative inference. In M. Reinfran, J. de Cleer, M. L. Ginsberg, and E. Sandewall, editors, *2nd International workshop on Non-monotonic reasoning*, pages 1–18. Springer-Verlag, 1989.
- [Mak94] D. Makinson. General patterns in nonmonotonic reasoning. *Handbook of logic in artificial intelligence and logic programming: nonmonotonic reasoning and uncertain reasoning*, 3:35–110, 1994.
- [Mak05] D. C. Makinson. *Bridges from Classical to Nonmonotonic Logic*. London: Kings College Publications, 2005.
- [Mou85] H. Moulin. Choices functions over a finite set: A summary. *Social Choice and Welfare*, 2:147–160, 1985.
- [Pri91] G. Priest. Minimally inconsistent LP. *Studia Logica*, 50:321–331, 1991.
- [Sch92] K. Schlechta. Some results on classical preferential models. *Journal of Logic and Computation*, 2(6):675–686, 1992.
- [Sch96] K. Schlechta. Some completeness results for stoppered and ranked classical preferential models. *Journal of Logic and Computation*, 6(4):599–622, 1996.
- [Sch00] K. Schlechta. New techniques and completeness results for preferential structures. *The Journal of Symbolic Logic*, 65(2):719–746, 2000.
- [Sch04] K. Schlechta. *Coherent systems*. Elsevier, 2004.
- [Sen70] A. K. Sen. *Collective Choice and Social Welfare*. Holden-Day, San Francisco, CA, 1970.
- [Sho87] Y. Shoham. A semantical approach to nonmonotonic logics. In *Tenth International Joint Conference on Artificial Intelligence (IJCAI)*, pages 388–392, 1987.
- [Sho88] Y. Shoham. *Reasoning About Change*. MIT Press, Cambridge USA, 1988.