

Dynamic Analysis of a One-Parameter Chaotic System in Complex Field

XIU ZHAO^{®1}, JIAN LIU^{®1}, HONGJUN LIU^{®1}, AND FANGFANG ZHANG^{®2}

¹School of Mathematical Sciences, University of Jinan, Jinan 250022, China
 ²Department of Electrical Engineering and Automation, Qilu University of Technology (Shandong Academy of Sciences), Jinan 250353, China
 Corresponding author: Jian Liu (ss_liuj@ujn.edu.cn)

This work was supported by the National Nature Science Foundation of China under Grant 61773010 and Grant 61703180.

ABSTRACT Chaotic dynamics analysis of complex-variable chaotic systems (CVCSs) is an important problem in real secure communication and encryption. In this paper, a simple one-parameter chaotic system in complex field is proposed, whose nonlinear terms are the same as Lorenz system but the linear terms are much simpler. The proposed system has circular equilibria and therefore multi-stability can be measured by phase portraits, bifurcation diagrams and Lyapunov exponent spectrum. Its basin of attraction is filled with initial points leading to chaotic behaviors. The coexistence of infinitely many attractors is found in the proposed system, which is not reported in the existing complex-variable Lorenz system. Finally, two complexity indexes are used to measure dynamic characteristic with respect to parameter.

INDEX TERMS Coexisting attractors, extreme multistability, complex-variable chaotic system, complexity.

I. INTRODUCTION

Up to now, chaos is a very important research topic and received much attention in complicated nonlinear dynamics and chaotic cryptography [1]–[4]. As a model for convection between parallel plates, the Lorenz system with real variables is always a paradigm of chaos. In 1982, the Lorenz system in the complex domain was unexpectedly discovered from laser physics and the geophysical flows [5], [6], and widely attracted attentions of researchers in multiple fields. Since then, complex-variable chaotic systems (CVCSs) have been extensively investigated in various important fields, such as detuned lasers [7], [8], fluids [9], electromagnetic fields [10] and image encryption [11]–[13].

Due to the existence of complex variables, the dynamic behaviors of complex system are more complex and diverse [14], [15]. Real variable is a special case of complex variable, so the conclusions of complex chaotic system are also applicable to real chaotic system. In recent years, the stabilization and synchronization [14]–[19] of CVCSs stimulated a great deal of interest among scientist for the wide scope of applications. Especially in the chaotic secure communication field, the complex variable can increase the content and security of the transmitted information [12], [20].

The coexistence of multi-attractors for the same group of parameters can be observed in a number of nonlinear circuits or systems [1], [21], [22], this phenomenon is called multistability. System displays different attractors depending on the different initial values. The coexistence of attractors exhibits a complex diversity of attractors of a nonlinear dynamical system. At present, the coexistence of attractors becomes one of the hot spots in international nonlinear dynamics research [23], [24]. Some interesting properties of multiple coexisting attractors are found not only in the novel 4D-hyperjerk system [25] and 5-D memristive oscillator, but also in the simple three-dimensional system, such as Sprott B system [26] and Lü system [27]. Recently, it is found that the coexistence of infinitely many attractors may be coined when a periodic function is introduced into the offsetboostable variable [23]. The phenomenon that system has a great many coexisting attractors is defined as extreme multistability. Bao et al. investigated the coexistence of infinitely many attractors through analyzing the stability distribution of line equilibrium point in a memristive circuit [28]. Similarly, extreme multistability is also observed in an extremely simple chaotic system [29]. It is very exciting that a memristive hyperchaotic system without any equilibrium points can still generate coexisting infinitely many attractors [30].

The associate editor coordinating the review of this manuscript and approving it for publication was Hassen Ouakad.

In the literature mentioned above, the phenomenon of coexisting attractors is widely considered in real-variable systems. However, the coexistence of attractors, particularly that of infinitely many chaotic attractors, is rarely reported in CVCSs so far. CVCSs have more important significance in practical application bacause they possess a diversity of coexisting attractors and exhibit very complicated basins of attraction. The main goal of present work is to investigate coexistence of attractors in one-parameter CVCSs.

The rest of this paper is structured as follows. In Sec. II, we introduce the mathematical model of one-parameter CVCSs and discuss the stability of circular equilibria, dissipation and symmetry. In Sec. III, the dynamical characteristics are investigated and the coexistence of infinitely many chaotic and periodic attractors depending on initial values are therefore discovered. In Sec. IV, spectral entropy (SE) and C_0 complexity algorithms are applied to analyze the complexity of the CVCSs. The last section concludes the paper.

II. THE PROPOSED COMPLEX-VARIABLE CHAOTIC SYSTEMS

The Sprott B system is algebraically the simplest one [31]. Assigning $x \rightarrow -y, y \rightarrow -x, z \rightarrow -z$, transform the Sprott B system into

$$\begin{cases} \dot{x} = -x + y, \\ \dot{y} = -xz, \\ \dot{z} = xy - a, \end{cases}$$
(1)

where (x, y, z) is the real state vector, and *a* is a positive constant parameter. The attractors of system (1) are similar to the attractors of Lorenz and Chen systems [32]. And its simplest 4-D extension has coexisting hidden attractors [33]. In this paper, we propose the complex extension of the system:

$$\begin{cases} \dot{z}_1 = -z_1 + z_2, \\ \dot{z}_2 = -z_1 z_3, \\ \dot{z}_3 = \frac{1}{2} (\bar{z}_1 z_2 + z_1 \bar{z}_2) - a, \end{cases}$$
(2)

where $z_1 = u_1 + ju_2$ and $z_2 = u_3 + ju_4$ are complex state variables, $z_3 = u_5$ is real state variable, and *a* is positive constant parameter. It is clear that system (2) has the same nonlinear parts as the complex-variable Lorenz and Chen systems, but has much simpler linear parts. Assume that a = 4.8 and the initial values $z_0 = (0.5 + 0.5j,$ 1 + j, 1). Then system (2) has Lyapunov exponents (0.0526, 0, -0.9533, -47.8200, -116.2680), and the corresponding Kaplan-Yorke dimension $D_{KY} = 2.048$. Due to its fractional dimension and positive Lyapunov exponent, system (2) is chaotic and the orbit is a bond chaotic attractor as in Fig. 1. It is apparent that bond-orbital attractor [34] is quite different from butterfly attractor in [32].



FIGURE 1. 3-D space of the bond chaotic attractor for system (2) when a = 4.8.

System (2) can be rewritten as a real-variable system of the form

$$\begin{cases}
\dot{u}_1 = -u_1 + u_3, \\
\dot{u}_2 = -u_2 + u_4, \\
\dot{u}_3 = -u_1 u_5, \\
\dot{u}_4 = -u_2 u_5, \\
\dot{u}_5 = u_1 u_3 + u_2 u_4 - a.
\end{cases}$$
(3)

System (3) has rotational symmetry with respect to the u_5 -axis, due to its invariance under the coordinates transform from $(u_1, u_2, u_3, u_4, u_5)$ to $(-u_1, -u_2, -u_3, -u_4, u_5)$. In particular, the symmetry about the u_5 -axis for any choice of parameter *a* is accurate. The corresponding Jacobian matrix of system (3) is

$$J = \begin{bmatrix} -1 & 0 & 1 & 0 & 0 \\ 0 & -1 & 0 & 1 & 0 \\ -u_5 & 0 & 0 & 0 & -u_1 \\ 0 & -u_5 & 0 & 0 & -u_2 \\ u_3 & u_4 & u_1 & u_2 & 0 \end{bmatrix}.$$

If $u_5 \neq 0$ and $u_1^2 + u_2^2 + u_1 u_3 + u_2 u_4 \neq 0$, system (3) has the full rank because the determinant of the Jacobian matrix is $-u_5(u_1^2 + u_2^2 + u_1 u_3 + u_2 u_4)$, which means that system (3) is a real five-dimensional system.

The rate of hypervolume contraction is given by the Lie derivative

$$\nabla V = \sum_{l=1}^{5} \frac{\partial \dot{u}_l}{\partial u_l} = -2.$$

Hence, system (3) is dissipative. This means each volume element V_0 containing the system trajectories shrinks to zero when $t \to \infty$ at an exponential rate -2. From the algebraic equations: $-u_1 + u_3 = 0$, $-u_2 + u_4 = 0$, $-u_1u_5 = 0$, $-u_2u_5 = 0$, $u_1u_3 + u_2u_4 - a = 0$, one can derive that system (3) has infinite equilibrium points. Let $u_1 = u_3 = \sqrt{a} \cos \theta$ and $u_2 = u_4 = \sqrt{a} \sin \theta$, where $\theta \in [0, 2\pi]$, the non-isolated (nontrivial) fixed points can be described as

$$E = (\sqrt{a}\cos\theta, \sqrt{a}\sin\theta, \sqrt{a}\cos\theta, \sqrt{a}\sin\theta, 0).$$



FIGURE 2. Bifurcation diagram and Lyapunov exponent spectrum versus parameter *a* of system (3).

The characteristic equation of the corresponding Jacobian matrix is obtained as $\mu(\mu + 1)(\mu^3 + \mu^2 + a\mu + 2a) = 0$. According to Routh-Hurwitz criterion, system (3) has one eigenvalue 0, two negative real eigenvalues and a pair of conjugate complex eigenvalues with positive real part provided a > 0. For example, when a = 1, eigenvalues are $\mu_1 = -1.3532, \mu_2 = -1, \mu_3 = 0, \mu_4 = 0.1766 + 1.2028j, \mu_5 = 0.1766 - 1.2028j$, which means that circular equilibria are non-hyperbolic and saddle-foci with index 2 (unstable).

III. CHAOTIC ATTRACTORS

To verify existence of the chaotic attractors, we use two effective indexes, Lyapunov exponents and bifurcation diagram, to characterize statistical properties. Lyapunov exponents are the average exponential divergence rate of two closed orbits, which are used to quantitatively express the stability of the orbits and its sensitivity to the initial conditions. For chaotic attractors, it possess the positive Lyapunov exponents. According to the definition, Lyapunov exponents are adopted as

$$\lambda_l = \lim_{t \to \infty} \frac{1}{t} \ln \left| \frac{\delta u_l(u_0, t)}{\delta u_l(u_0, 0)} \right|.$$
(4)

For the initial condition $U_0 = (0.5, 0.5, 1, 1, 1)$, the bifurcation diagram and Lyapunov exponent spectrum versus parameter *a* are plotted in Fig. 3. It can be seen from Fig. 2(a) that the bifurcation of system (3) with respect to *a* is reverse period-doubling bifurcation. Fig. 2(b) shows that system (3)



FIGURE 3. 3D views of chaotic attractors of system (3) when a = 0.12.



FIGURE 4. 3D views of the two-butterfly chaotic attractor of system (3) when a = 3.61.

has periodic behaviors when $a \in (0.24, 0.32), (2.4, 2.53)$ and (5.02, 10). The chaotic attractors are clear for $a \in (0, 0.23)$, (0.33, 2.39) and (2.54, 5.01), respectively. Note that there exist always two zero Lyapunov exponents in Fig. 2(b).

The Lyapunov exponents of system (3) are (0.0197, 0, -0.0113, -0.987, -1.0127) when a = 0.12. The chaotic attractor and the infinite equilibria when a = 0.12 are shown in Fig 3, where red circle represents the equilibrium points, gray plots are projections of the attractor onto the different planes. Interestingly, a two-butterfly chaotic attractor with Lyapunov exponents (0.2866, 0, 0, -0.9983, -1.2876) is found when a = 3.61, as shown in Fig. 4.

IV. COEXISTING ATTRACTORS

A. COEXISTENCE OF TWO ATTRACTORS

The coexisting bifurcation diagram and coexisting Lyapunov exponent spectrum are used to examine the coexisting attractors of the system. The coexisting bifurcation diagram about state variable u_3 and coexisting Lyapunov exponent spectrum versus $a \in (0, 8)$ are plotted in Fig. 5, respectively. As shown in Fig. 5(a), the initial values for the blue orbit is $U_0 = (0.5, 0.5, 1, 1, 1)$, while those for the blue one is $U_1 = (0.5, 0, 0.5, 0.1, 0.1)$. It is obvious that $|\max(u_3)|$ starting from U_1 is larger than that from U_0 at $a \in (0, 0.23), (0.33, 2.39)$ and (2.54, 5.01), which means that there are two different amplitudes attractors. In Fig. 5(b),



(a) Coexisting bifurcation diagram



(b) Coexisting Lyapunov exponent spectrum

FIGURE 5. Coexisting bifurcation diagram and coexisting Lyapunov exponent spectrum of system (3) for the initial conditions $U_0 = (0.5, 0.5, 1, 1, 1)$ and $U_1 = (0.5, 0, 0.5, 0.1, 0.1)$.

the blue and green trajectories express the first two Lyapunov exponents of the system with initial value U_0 , while the red and pink trajectories represent the case with U_1 .

It is very interesting that coexisting butterfly attractors with asymmetric initial conditions are shown in Fig. 6, where green attractors start form red initial conditions U_0 and pink attractors start form blue U_1 . Fig. 6 illustrates the coexistence of periodic attractors when a = 2.4, coexistence of periodic attractor and chaotic attractor when a = 2.5, coexistence of two-butterfly chaotic attractors when a = 3.61 and coexistence of chaotic attractors when a = 4.56.

B. COEXISTENCE OF SIX CHAOTIC ATTRACTORS

The basin of attraction is a useful method to examine the coexisting attractors of the system, which changes with the initial values [28], [33]. When parameter of the system is set as a = 4 and the initial value is given as $(0.5, 1, u_3, 1, u_5)$, the basin of attraction with respect to initial values is shown in Fig. 7(a). The chaotic sea is filled with points leading to chaotic attractors when $u_3 \in [-10, 10]$ and $u_5 \in [-10, 10]$, where different colors represent different chaotic attractors in Fig. 7(a), respectively.

Fig. 7(b) depicts six typical chaotic attractors which are observed in the basin of attraction. And the initial values of the attractors are selected from different color



(a) Coexistence of periodic attractors (a = 2.4)



(b) Coexistence of chaotic attractor and periodic attractor (a = 2.5)



(c) Coexistence of two-butterfly attractors (a = 3.61)



FIGURE 6. Coexistence of attractors in the (u_2, u_5) plane for the initial conditions $U_0 = (0.5, 0.5, 1, 1, 1)$ and $U_1 = (0.5, 0, 0.5, 0.1, 0.1)$.

points in the basin of attraction. In Fig. 7(b), the red attractor corresponds to the initial value (0.5, 1, 1, 1, -6) with LEs (0.2066, 0, 0, -1.0036, -1.2121), the blue attractor corresponds to the initial value (0.5, 1, 4, 1, -6) with



FIGURE 7. The basin of attraction and the coexisting chaotic attractors with initial condition $(0.5, 1, u_3, 1, u_5)$ when a = 4.

LEs (0.2371, 0, 0, -0.9983, -1.2396), the green attractor corresponds to the initial value (0.5, 1, -2.8, 1, 6) with LEs (0.2269, 0, 0, -0.9977, -1.2330), the yellow attractor corresponds to the initial value (0.5, 1, 6, 1, 6) with LEs (0.2053, 0, 0, -1.0034, -1.2048), the cyan attractor corresponds to the initial value (0.5, 1, 8, 1, -3) with LEs (0.2497, 0, 0, -0.9978, -1.2482) and the pink attractor corresponds to the initial value (0.5, 1, 6.8, 1, -10) with LEs (0.2061, 0, 0, -1.0012, -1.2001), respectively.

C. THE COEXISTENCE OF INFINITELY MANY ATTRACTORS

Extreme multi-stability means that the coexistence of an infinite number of attractors exists in a nonlinear system. And it can be obtained in a nonlinear dynamical circuit or system [28], [33], [35]. To further confirm the infinite number of coexisting attractors, the coexisting bifurcation diagrams with the change of initial conditions and phase diagram of attractors with different amplitudes are drawn.

Here, the parameter a = 5 is set and the initial conditions versus *b* are considered. When *b* is gradually increased from -2 to 2, coexisting bifurcation diagrams are shown in Fig. 8, where the green, cyan, pink, red and blue branches are yielded from initial conditions (*b*, 0.5, 0.5, 1, 1), (0.5, *b*, 0.5, 1, 1), (0.5, 0.5, *b*, 1, 1), (0.5, 0.5, 0.5, 1, 1), (0.5, 0.5, 0.5, 1, b), respectively. The coexistence of attractors with different amplitudes is found, which means that coexistence phenomenon of infinitely many attractors in the





(b) Coexisting bifurcation diagram versus b

FIGURE 8. Coexisting bifurcation diagram versus initial conditions when a = 5.

proposed system. We only pick initial values selected randomly in Fig. 8, and the coexistence of infinitely many attractors as plotted in Fig. 9.

In Fig. 9, the coexistence of infinitely many chaotic attractors at a = 5 and the the coexistence of infinitely many periodic attractors at a = 7 with different initial conditions are discovered, where the black attractors match the initial value (0.5, 0.5, 0.5, 2, 1), the yellow attractors match (0.5, 0.5, 0.5, 1, 1.5), the blue attractors match (0.5, 0.5, 0.5, 1, 1.5), the blue attractors match (0.5, 0.5, 0.5, 1, 1.5), the purple attractors match (0.5, 0.5, 0.5, 1, -1.6), the pink attractors match (0.5, 0.5, -1.4, 1, 1), the green attractors match (0.5, 0.5, 0.5, 1, 1) and the red attractors match (0.5, -1.5, 0.5, 1, 1). In fact, for more different initial values, more attractors could be found. Fig. 9 clearly disclose the coexistence of infinite many chaotic and periodic attractors, which indicates that there indeed exists extreme multistability in system.

It is worth mentioning that the proposed system (2) has the coexistence of infinitely many attractors because of its infinite equilibrium points. This property of extreme multistability is not reported in the Lorenz CVCSs in [15], [16], [19] and the real-variable systems in [28], [31]–[33], [35].

V. SPECTRAL ENTROPY (SE) AND C₀ COMPLEXITY

Complexity is an important reference to measure dynamic characteristics of a chaotic system. Up to now, there are several main algorithms for measuring the complexity of



(a) The coexistence of infinitely many chaotic attractors



(b) The coexistence of infinitely many periodic attractors





FIGURE 10. SE and C_0 complexity versus parameter *a* for the initial condition $U_0 = (0.5, 0.5, 1, 1, 1)$.

chaotic sequences, such as intensive statistical method [36], multiscale entropy (MSE) [37], spectral entropy (SE) [38], and C_0 complexity [39]. In particular, C_0 and SE complexity

algorithms often have been used to analyze information security of the system in chaotic secure communication, because of their less parameters, faster calculation speed, and higher accuracy [20]. The SE value is obtained through the energy distribution in the Fourier transform domain and Shannon entropy. The main idea of C_0 complexity is to decompose the sequence into regular and irregular components. And its measure value is the proportion of irregular components in the sequence. Here, the SE and C_0 complexity of the complex system versus parameter a at the initial condition $U_0 = (0.5, 0.5, 1, 1, 1)$ are shown in Fig. 10, they have the similar changing trend, although the values are different. Compared with the Lyapunov exponent spectrum and bifurcation diagram in Fig. 10, SE and C₀ complexity are consistent, which means that complexity can also reflect the dynamic characteristics of a chaotic complex system.

The value of SE complexity for proposed CVCSs has obvious change for the period, whereas it has relatively small change for the chaos like the real-variable system [21]. The value of C_0 complexity has obvious change for both period and chaos. It is noted that the SE complexity of two-butterfly attractor at a = 3.61 is biggest in Fig. 10(a), and the C_0 complexity of chaotic attractor with a = 4.56 is the biggest in Fig. 10(b).

VI. CONCLUSION

This paper studied dynamical properties of complex-variable chaotic systems with circular equilibria. Numerical simulations displayed that the proposed system had a number of rich and interesting dynamic phenomenon. Furthermore, chaotic sea about chaotic attractors was revealed in the basin of attraction. In particular, the coexistence of infinitely many chaotic and periodic attractors with different values of control parameter were discovered in coexisting bifurcation diagram versus initial conditions. The results of spectral entropy (SE) and C_0 complexity were matched well with Lyapunov exponent spectrum and bifurcation model. Only some preliminary and empirical experimental results were reported, much more underlying dynamical properties and their applications deserve further exploration.

ACKNOWLEDGMENT

The author would like to thank the anonymous referees for their valuable suggestions and questions.

REFERENCES

- N. Wang, C. Li, H. Bao, M. Chen, and B. Bao, "Generating multiscroll Chua's attractors via simplified piecewise-linear Chua's diode," *IEEE Trans. Circuits Syst. I, Reg. Papers*, vol. 66, no. 12, pp. 4767–4779, Dec. 2019.
- [2] C. Li, Y. Zhang, and E. Y. Xie, "When an attacker meets a cipherimage in 2018: A year in review," J. Inf. Secur. Appl., vol. 48, Oct. 2019, Art. no. 102361.
- [3] Z. Hua, B. Xu, F. Jin, and H. Huang, "Image encryption using josephus problem and filtering diffusion," *IEEE Access*, vol. 7, pp. 8660–8674, 2019.
- [4] C. Li, B. Feng, S. Li, J. Kurths, and G. Chen, "Dynamic analysis of digital chaotic maps via state-mapping networks," *IEEE Trans. Circuits Syst. I, Reg. Papers*, vol. 66, no. 6, pp. 2322–2335, Jun. 2019.

- IEEE Access^{*}
- [5] A. Fowler, J. Gibbon, and M. Mcguinness, "The complex Lorenz equations," *Phys. D, Nonlinear Phenomena*, vol. 4, no. 2, pp. 139–163, Jan. 1982.
- [6] J. Gibbon and M. Mcguinness, "The real and complex Lorenz equations in rotating fluids and lasers," *Phys. D, Nonlinear Phenomena*, vol. 5, no. 1, pp. 108–122, May 1982.
- [7] H. Zeghlache and P. Mandel, "Influence of detuning on the properties of laser equations," J. Opt. Soc. Amer. B, vol. 2, no. 1, p. 18, Jan. 1985.
- [8] C.-Z. Ning and H. Haken, "Detuned lasers and the complex Lorenz equations: Subcritical and supercritical Hopf bifurcations," *Phys. Rev. A, Gen. Phys.*, vol. 41, no. 7, pp. 3826–3837, Jul. 2002.
- [9] G. Van Tartwijk and G. Agrawal, "Absolute instabilities in lasers with host-induced nonlinearities and dispersion," *IEEE J. Quantum Electron.*, vol. 34, no. 10, pp. 1854–1860, Oct. 1998.
- [10] G. M. Mahmoud and T. Bountis, "The dynamics of systems of complex nonlinear oscillators: A review," *Int. J. Bifurcation Chaos*, vol. 14, no. 11, pp. 3821–3846, Nov. 2004.
- [11] C. Li, D. Lin, B. Feng, J. Lü, and F. Hao, "Cryptanalysis of a chaotic image encryption algorithm based on information entropy," *IEEE Access*, vol. 6, pp. 75834–75842, 2018.
- [12] C. Li, D. Lin, J. Lü, and F. Hao, "Cryptanalyzing an image encryption algorithm based on autoblocking and electrocardiography," *IEEE MultimediaMag.*, vol. 25, no. 4, pp. 46–56, Oct./Dec. 2018.
- [13] H. Liu, Y. Zhang, A. Kadir, and Y. Xu, "Image encryption using complex hyper chaotic system by injecting impulse into parameters," *Appl. Math. Comput.*, vol. 360, pp. 83–93, Nov. 2019.
- [14] F.-F. Zhang, S.-T. Liu, and W.-Y. Yu, "Modified projective synchronization with complex scaling factors of uncertain real chaos and complex chaos," *Chin. Phys. B*, vol. 22, no. 12, Dec. 2013, Art. no. 120505.
- [15] J. Liu, S. Liu, and C. Yuan, "Adaptive complex modified projective synchronization of complex chaotic (hyperchaotic) systems with uncertain complex parameters," *Nonlinear Dyn.*, vol. 79, no. 2, pp. 1035–1047, Jan. 2015.
- [16] J. Liu, S. Liu, and J. C. Sprott, "Adaptive complex modified hybrid function projective synchronization of different dimensional complex chaos with uncertain complex parameters," *Nonlinear Dyn.*, vol. 83, nos. 1–2, pp. 1109–1121, Jan. 2016.
- [17] F. Nian, X. Wang, Y. Niu, and D. Lin, "Module-phase synchronization in complex dynamic system," *Appl. Math. Comput.*, vol. 217, no. 6, pp. 2481–2489, Nov. 2010.
- [18] X. Wang and Y. He, "Projective synchronization of fractional order chaotic system based on linear separation," *Phys. Lett. A*, vol. 372, no. 4, pp. 435–441, Jan. 2008.
- [19] J. Liu and S. Liu, "Complex modified function projective synchronization of complex chaotic systems with known and unknown complex parameters," *Appl. Math. Model.*, vol. 48, pp. 440–450, Aug. 2017.
- [20] F. Yang, J. Mou, J. Liu, C. Ma, and H. Yan, "Characteristic analysis of the fractional-order hyperchaotic complex system and its image encryption application," *Signal Process.*, vol. 169, Apr. 2020, Art. no. 107373.
- [21] N. Wang, G. Zhang, and H. Bao, "Bursting oscillations and coexisting attractors in a simple memristor-capacitor-based chaotic circuit," *Nonlinear Dyn.*, vol. 97, no. 2, pp. 1477–1494, 2019.
- [22] T. Lu, C. Li, S. Jafari, and F. Min, "Controlling coexisting attractors of conditional symmetry," *Int. J. Bifurcation Chaos*, vol. 29, no. 14, 2019, Art. no. 1950207.
- [23] C. Li, T. Lu, G. Chen, and H. Xing, "Doubling the coexisting attractors," *Chaos*, vol. 29, no. 5, May 2019, Art. no. 051102.
- [24] A. Bayani, K. Rajagopal, A. J. M. Khalaf, S. Jafari, G. Leutcho, and J. Kengne, "Dynamical analysis of a new multistable chaotic system with hidden attractor: Antimonotonicity, coexisting multiple attractors, and offset boosting," *Phys. Lett. A*, vol. 383, no. 13, pp. 1450–1456, Apr. 2019.
- [25] G. Leutcho, J. Kengne, and L. K. Kengne, "Dynamical analysis of a novel autonomous 4-D hyperjerk circuit with hyperbolic sine nonlinearity: Chaos, antimonotonicity and a plethora of coexisting attractors," *Chaos, Solitons Fractals*, vol. 107, pp. 67–87, Feb. 2018.
- [26] Q. Lai and S. Chen, "Generating multiple chaotic attractors from Sprott B system," *Int. J. Bifurcation Chaos*, vol. 26, no. 11, Oct. 2016, Art. no. 1650177.
- [27] Q. Lai, B. Norouzi, and F. Liu, "Dynamic analysis, circuit realization, control design and image encryption application of an extended Lü system with coexisting attractors," *Chaos, Solitons Fractals*, vol. 114, pp. 230–245, Sep. 2018.

- [28] B. Bao, T. Jiang, G. Wang, P. Jin, H. Bao, and M. Chen, "Two-memristorbased Chua's hyperchaotic circuit with plane equilibrium and its extreme multistability," *Nonlinear Dyn.*, vol. 89, no. 2, pp. 1157–1171, 2017.
- [29] Q. Lai, P. D. K. Kuate, F. Liu, and H. H.-C. Iu, "An extremely simple chaotic system with infinitely many coexisting attractors," *IEEE Trans. Circuits Syst. II, Exp. Briefs*, to be published.
- [30] B. Bao, H. Bao, N. Wang, M. Chen, and Q. Xu, "Hidden extreme multistability in memristive hyperchaotic system," *Chaos, Solitons Fractals*, vol. 94, pp. 102–111, Jan. 2017.
- [31] J. C. Sprott, "Some simple chaotic flows," *Phys. Rev. E, Stat. Phys. Plasmas Fluids Relat. Interdiscip. Top.*, vol. 50, no. 2, pp. R647–R650, Jul. 2002.
- [32] X. Wang and G. Chen, "A gallery of lorenz-like and Chen-like attractors," *Int. J. Bifurcation Chaos*, vol. 23, no. 04, Apr. 2013, Art. no. 1330011.
- [33] C. Li and J. C. Sprott, "Coexisting hidden attractors in a 4-D simplified lorenz system," *Int. J. Bifurcation Chaos*, vol. 24, no. 3, Mar. 2014, Art. no. 1450034.
- [34] X. Zhang, C. Wang, W. Yao, and H. Lin, "Chaotic system with bondorbital attractors," *Nonlinear Dyn.*, vol. 97, no. 4, pp. 2159–2174, Sep. 2019.
- [35] C. Li, J. C. Sprott, Y. Liu, Z. Gu, and J. Zhang, "Offset boosting for breeding conditional symmetry," *Int. J. Bifurcation Chaos*, vol. 28, no. 14, Dec. 2018, Art. no. 1850163.
- [36] H. Larrondo, C. González, M. Martín, A. Plastino, and O. Rosso, "Intensive statistical complexity measure of pseudorandom number generators," *Phys. A, Stat. Mech. Appl.*, vol. 356, no. 1, pp. 133–138, Oct. 2005.
- [37] M. Borowiec, A. Rysak, D. Betts, C. Bowen, H. Kim, and G. Litak, "Complex response of a bistable laminated plate: Multiscale entropy analysis," *Eur. Phys. J. Plus*, vol. 129, no. 10, p. 211, 2014.
- [38] K. Sun, S. He, H. Yi, and L. Yin, "Complexity analysis of chaotic pseudorandom sequences based on spectral entropy algorithm," *Acta Phys. Sinica*, vol. 2, no. 1, 2013, p. 010501.
- [39] K. Sun, S. He, C. Zhu, and Y. He, "Analysis of chaotic complexity characteristics based on C₀ algorithm," *Acta Electron. Sinica*, vol. 41, no. 9, pp. 1765–1771, 2013.



XIU ZHAO received the B.S. degree in applied mathematics from the University of Jinan, Jinan, China, in 2018, where she is currently pursuing the M.S. degree in analysis and control of complex systems. Her research interests mainly include nonlinear dynamics, synchronization, and control of complex systems.



JIAN LIU received the B.S. degree in mathematics from Qufu Normal University, and the M.S. and Ph.D. degrees in control science and control engineering from Shandong University, China. She is currently a Full Professor with the School of Mathematical Sciences, University of Jinan, Shandong, China. Her research interest mainly focus on nonlinear dynamics, synchronization, and control of complex systems.



HONGJUN LIU received the B.S. degree in computer application technology from Shandong University, Shandong, China, in 1995, the M.S. degree in computer application technology from the Ocean University of China, Shandong, in 2005, and the Ph.D. degree from the Dalian University of Technology, Dalian, China, in 2014. He is currently an Associate Professor with the School of Mathematical Sciences, University of Jinan, Shandong. He did a lot of research in the design

and cryptanalysis of chaos-based symmetric cryptography: block ciphers, hash functions, and stream ciphers.



FANGFANG ZHANG received the B.S. degree from Northeast Petroleum University, the M.S. degree from the Beijing University of Technology and the Ph.D. degree from Shandong University, China. She is currently an Associate Professor with the Department of Electrical Engineering and Automation, Qilu University of Technology (Shandong Academy of Sciences), Shandong, China. Her interests include nonlinear dynamics, the complex chaotic systems, and chaotic secure communication.

...