Invariance Principles for Switched Systems with Restrictions

J. L. Mancilla-Aguilar * and R.A. García †

November 6, 2018

Abstract

In this paper we consider switched nonlinear systems under average dwell time switching signals, with an otherwise arbitrary compact index set and with additional constraints in the switchings. We present invariance principles for these systems and derive by using observability-like notions some convergence and asymptotic stability criteria. These results enable us to analyze the stability of solutions of switched systems with both state-dependent constrained switching and switching whose logic has memory, i.e., the active subsystem only can switch to a prescribed subset of subsystems.

1 Introduction

A switched system is a family of continuous-time dynamical subsystems and a rule, usually time or state-dependent, that orchestrates the switching between them. At first glance switched systems may look simple; nevertheless their behavior may be very complicated, being a classical example of this fact, divergent trajectories obtained by switching among asymptotically stable subsystems (see [11]). Consequently, the stability analysis of such systems turned out to be an important and challenging problem which has received considerable attention in the recent literature (see [4], [11], [12] and references therein). Although the stability of switched systems under arbitrary switching laws can be assured by the existence of a common Lyapunov function

^{*}Department of Physics & Mathematics, Instituto Tecnológico de Buenos Aires jmancill@itba.edu.ar.

[†]Department of Physics & Mathematics, Instituto Tecnológico de Buenos Aires ragarcia@itba.edu.ar.

(CLF) for all the switching modes ([11], [15]), this type of stability condition is deemed to be too conservative when a particular type of switching logic is considered. In fact, switched systems that do not share a CLF may be stable under restricted switching signals. Restrictions on the set of admissible switching signals of a certain switched system arise naturally from physical constraints of the system, from design strategies (e.g. discontinuous control feedback laws), or from the knowledge about possible switching logic of the switched system, e.g., partitions of the state space and their induced switching rules. Multiple Lyapunov functions (MLF) have been shown to be very useful tools for the stability analysis of switched systems with constrained switchings. In this context each switching mode may have its own Lyapunov function (see [4] and references therein). However, some additional conditions are necessary to assure that the value of each Lyapunov function on its corresponding mode will decrease. Sufficient conditions for asymptotic stability of switched systems with MLF can be found in [4], [11] and references therein. When the derivative of a candidate Lyapunov function with respect to (w.r.t) each mode is only *non-positive*, the convergence of the solutions of the switching system to an equilibrium point, and consequently the asymptotic stability, can be derived from one of the various extensions to switched systems of LaSalle's invariance principle for differential equations (see [8, 9]). Hespanha in [6] introduced an invariance principle for switched linear systems under *persistently dwell-time* switching signals and in [7] Hespanha et al. extended some of those results to a family of nonlinear systems. Bacciotti and Mazzi presented in [1] an invariance principle for switched systems with dwell-time signals. An invariance principle for switched nonlinear systems with average dwell-time signals that satisfy state-dependent constraints was derived by Mancilla-Aguilar and García in [16] from the sequential compactness of particular classes of trajectories of switched systems. Based on invariance results for hybrid systems ([17]), Goebel *et al.* in [5] obtained recently invariance results for switched systems under various types of switching signals. Lee and Jiang in [10] gave a generalized version of Krasovskii-LaSalle Theorem for time-varying switched systems. Under certain ergodicity conditions on the switching signal, some stability results were also obtained in [3, 18, 19].

Most of the invariance results for switched systems already published only consider restrictions originated by the timing of the switchings or by the state dependence of it. Nevertheless there is also an important restriction to take into account: the fact that not all the subsystems may be accessible from a particular one, i.e. the case in which the switching logic has memory. This restriction is clearly exhibited, for example, in switched systems which are the continuous portion of a hybrid automaton (see [4], [14]). In this regard, the invariance principles developed for hybrid systems in [14] and in [17] could be useful in the analysis of switched systems with this class of restriction in the switchings.

In this paper we present invariance results that hold for trajectories of switched systems with a non necessarily finite number of subsystems and whose switching signals verify an average dwell time condition and belong to a family for which a certain property \mathbf{P} holds. As various of the restricted switching classes mentioned above satisfy \mathbf{P} , these results enable us to obtain in an unified way invariance theorems for all of them. Based on these invariance results, we derive new convergence and stability criteria that recover, generalize and strengthen some results previously obtained. In particular:

- Theorem 3.2 extends LaSalle's invariance principle to switched systems with different restrictions on the switching signals by involving both backward and forward invariance as in, for example, [9, Theorem 6.4].
- Theorems 1 and 2 in [1], Corollary 5.6 of [5] and Proposition 4.1 of [16] follow from Theorem 3.3.
- The first conclusion of Corollary 4.10 in [5] is a particular case of Theorem 4.1.
- Theorem 4.7 is an improvement of Theorem 15 in [18]. In fact the hypotheses of Theorem 4.7 are weaker since the existence of a Common Joint Lyapunov Function is not assumed.
- Corollary 4.1, whose hypotheses are weaker than those of Theorem 3 in [3], improves it.

The paper presents two groups of statements. First we present statements about invariance of sets to which bounded trajectories of the switched systems converge (Theorems 3.2, 3.3 and 3.4). These statements involve either continuous functions which are nonincreasing along forward complete trajectories of the switched system or appropriately fast vanishing "outputs". Finally we present results about convergence and asymptotic stability (Theorems 4.1 to 4.7) that rely on observability-like conditions on the functions which bound the derivatives of nonincreasing functions as those mentioned above.

The article unfolds as follows. Section 2. contains the basic definitions. In Section 3. we present invariance principles for switched systems with constrained switching. Convergence and stability results are given in Section 4. Finally Section 5. contains some conclusions.

2 Basic definitions

In this work we consider switched systems described by

$$\dot{x} = f(x,\sigma) \tag{1}$$

where x takes values in \mathbb{R}^n , $\sigma : \mathbb{R} \to \Gamma$, with Γ a compact metric space, is a *switching signal*, *i.e.*, σ is piecewise constant (it has at most a finite number of jumps in each compact interval) and is continuous from the right and $f : \operatorname{dom}(f) \to \mathbb{R}^n$, with $\operatorname{dom}(f)$ a closed subset of $\mathbb{R}^n \times \Gamma$, is continuous.

For each $\gamma \in \Gamma$, let $\chi_{\gamma} = \{\xi \in \mathbb{R}^n : (\xi, \gamma) \in \text{dom}(f)\}$ and $f_{\gamma} : \chi_{\gamma} \to \mathbb{R}^n$ be defined by $f_{\gamma}(\xi) = f(\xi, \gamma)$; then χ_{γ} is closed and f_{γ} is continuous for any $\gamma \in \Gamma$. We note that when Γ is finite, these last two conditions imply that dom(f) is closed and that f is continuous on dom(f). In the sequel we denote with \mathcal{S} the set of all the switching signals.

Given $\sigma \in \mathcal{S}$, a solution of (1) corresponding to σ is a locally absolutely continuous function $x : I_x \to \mathbb{R}^n$, with $I_x \subset \mathbb{R}$ a nonempty interval, such that $(x(t), \sigma(t)) \in \text{dom}(f)$ for all $t \in I_x$ and $\dot{x}(t) = f(x(t), \sigma(t))$ for almost all $t \in I_x$. The solution x is complete if $I_x = \mathbb{R}$ and forward complete if $\mathbb{R}_{\geq 0} \subset I_x$. A pair (x, σ) is a trajectory of (1) if $\sigma \in \mathcal{S}$ and x is a solution of (1) corresponding to σ . The trajectory is complete or forward complete if x is complete or forward complete, respectively. Given a subset \mathcal{O} of \mathbb{R}^n , we say that the trajectory (x, σ) is precompact relative to \mathcal{O} if there exists a compact set $B \subset \mathcal{O}$ such that $x(t) \in B$ for all $t \in I_x$. When $\mathcal{O} = \mathbb{R}^n$ we simply say that (x, σ) is precompact.

Remark 2.1 Note that we do not suppose that $\operatorname{dom}(f) = \mathbb{R}^n \times \Gamma$. In this way we can take into account, in the analysis of the asymptotic behavior of a given trajectory (x, σ) of (1), some kind of state-dependent constraints which the trajectory under study must satisfy. In fact, in some situations we are not interested in the behavior of an arbitrary forward complete trajectory (x, σ) of a switched system (1) (with $\operatorname{dom}(f) = \mathbb{R}^n \times \Gamma$) but only of one of those that verify the constraint

$$x(t) \in \chi_{\sigma(t)} \quad for \ all \ t \in I_x, \tag{2}$$

where $\{\chi_{\gamma} : \gamma \in \Gamma\}$ is a collection of subsets of \mathbb{R}^n . If we consider the map \tilde{f} , which is the restriction of f to the set dom $(\tilde{f}) = \{(\xi, \gamma) : \xi \in \chi_{\gamma}\}$, and if dom (\tilde{f}) is closed in $\mathbb{R}^n \times \Gamma$, then the set of trajectories (x, σ) of (1) which verify (2) coincides with the set of trajectories of

$$\dot{x} = f(x, \sigma). \tag{3}$$

It must be pointed out that in this way we can consider the system as if its switching is state-independent, and focus on the restrictions imposed to it by the timing of the discontinuities of the switching signal and/or by the accessibility to certain subsystems from another ones.

In this paper we consider forward complete solutions of (1) corresponding to switching signals σ which belong to particular subclasses of S. Let $\Lambda(\sigma)$ be the set of times where σ has a jump (switching time). Following [6] we say that $\sigma \in S$ has a dwell-time $\tau_D > 0$ if $|t - t'| \ge \tau_D$ for any pair $t, t' \in \Lambda(\sigma)$ such that $t \neq t'$.

A switching signal σ has an average dwell-time $\tau_D > 0$ and a chatter bound $N_0 \in \mathbb{N}$ if the number of switching times of σ in any open finite interval $(\tau_1, \tau_2) \subset \mathbb{R}$ is bounded by $N_0 + (\tau_2 - \tau_1)/\tau_D$, i.e. $\operatorname{card}(\Lambda(\sigma) \cap (\tau_1, \tau_2)) \leq N_0 + (\tau_2 - \tau_1)/\tau_D$.

We denote by $S_a[\tau_D, N_0]$ the set of all the switching signals which have an average dwell-time $\tau_D > 0$ and a chatter bound $N_0 \in \mathbb{N}$ and by $\mathcal{T}_a[\tau_D, N_0]$ the set of all the complete trajectories (x, σ) of (1) with $\sigma \in S_a[\tau_D, N_0]$ and let $S_a = \bigcup_{\tau_D > 0, N_0 > 0} S_a[\tau_D, N_0]$ and $\mathcal{T}_a = \bigcup_{\tau_D > 0, N_0 > 0} \mathcal{T}_a[\tau_D, N_0]$. We note that the set of switching signals σ which have a dwell-time $\tau_D > 0$ coincides with $S_a[\tau_D, 1] := S_d[\tau_D]$. We denote by $\mathcal{T}_d[\tau_D]$ the set of all the complete trajectories (x, σ) of (1) with $\sigma \in S_d[\tau_D]$ and let $S_d = \bigcup_{\tau_D > 0} S_d[\tau_D]$ and $\mathcal{T}_d = \bigcup_{\tau_D > 0} \mathcal{T}_d[\tau_D]$.

For Γ a finite set and T > 0, we denote by $\mathcal{S}_e[T]$ the family of all the switching signals σ which verify the following "ergodicity" condition: for every $t_0 \geq 0$ and every $\gamma \in \Gamma$, $\sigma^{-1}(\gamma) \cap [t_0, t_0 + T] \neq \emptyset$.

 $\mathcal{T}_e[T]$ will denote the set of complete trajectories (x, σ) with $\sigma \in \mathcal{S}_e[T]$ and $\mathcal{S}_e = \bigcup_{T>0} \mathcal{S}_e[T]$ and $\mathcal{T}_e = \bigcup_{T>0} \mathcal{T}_e[T]$.

The families of switching signals already introduced have no restrictions on the accessibility from any subsystem to another. The family of switching signals —and their corresponding trajectories— that we introduce next, takes into account the case in which the switching logic has memory, i.e. when a subsystem corresponding to an index $\gamma \in \Gamma$ can only switch to subsystems corresponding to modes γ' that belong to a certain subset $\Gamma_{\gamma} \subset \Gamma$.

Given a set-valued map $H : \Gamma \rightsquigarrow \Gamma$, \mathcal{S}^H is the set of all the switching signals σ which verify the condition $\sigma(t) \in H(\sigma(t^-))$ for every time $t \in \Lambda(\sigma)$. Here $\sigma(t^-) = \lim_{s \to t^-} \sigma(s)$. \mathcal{T}^H denotes the set of all the complete trajectories (x, σ) with $\sigma \in \mathcal{S}^H$. This class of switching signals enable us, for example, to model the restrictions imposed by the discrete process of a hybrid system whose continuous portion is as in (1) (see [4]).

3 Invariance results for trajectories which satisfy a dwell-time condition

In this section we present some invariance results that enable us to characterize the asymptotic behavior of a precompact forward complete trajectory (x, σ) of (1) with σ belonging to a certain subclass of S_a . The consideration of such subclass allows us to obtain in an unified way invariance results for systems whose switching signals undergo different restrictions.

We recall that a point $\xi \in \mathbb{R}^n$ belongs to $\Omega(x)$, the ω -limit set of $x : I_x \to \mathbb{R}^n$, with $\mathbb{R}_{\geq 0} \subset I_x$, if there exists a strictly increasing sequence of times $\{s_k\} \subset I_x$ with $\lim_{k\to\infty} s_k = +\infty$ and $\lim_{k\to\infty} x(s_k) = \xi$. The ω -limit set $\Omega(x)$ is always closed and, when x evolves in a compact set of \mathbb{R}^n , it is nonempty, compact, connected if x is continuous, and $x \to \Omega(x)$ (for a set $M \subset \mathbb{R}^n, x \to M$ if $\lim_{t\to+\infty} d(x(t), M) = 0$, being $d(\xi, M) = \inf_{\nu \in M} |\nu - \xi|$).

As was done in [16], we will associate to each forward complete trajectory (x, σ) of (1) with $\sigma \in S_a$, the nonempty set $\Omega^{\sharp}(x, \sigma) \subset \mathbb{R}^n \times \Gamma$ that we introduce in the following

Definition 3.1 Given a forward complete trajectory (x, σ) of (1) with $\sigma \in S_a$, a point $(\xi, \gamma) \in \mathbb{R}^n \times \Gamma$ belongs to $\Omega^{\sharp}(x, \sigma)$ if there exists a strictly increasing and unbounded sequence $\{s_k\} \subset \mathbb{R}_{\geq 0}$ such that

- 1. $\lim_{k \to \infty} \tau_{\sigma}^{1}(s_k) s_k = r, \ 0 < r \le \infty,$
- 2. $\lim_{k\to\infty} x(s_k) = \xi$ and $\lim_{k\to\infty} \sigma(s_k) = \gamma$.

Here, for any $t \in \mathbb{R}$, $\tau_{\sigma}^{1}(t) = \inf\{s \in \Lambda(\sigma) : t < s\}$ if $\{s \in \Lambda(\sigma) : t < s\} \neq \emptyset$ and $\tau_{\sigma}^{1}(t) = +\infty$ in other case (i.e. $\tau_{\sigma}^{1}(t)$ is the first switching time greater than t).

Let $\pi_1 : \mathbb{R}^n \times \Gamma \to \mathbb{R}^n$ be the projection onto the first component. Then the following relation between $\Omega(x)$ and $\Omega^{\sharp}(x, \sigma)$ holds.

Lemma 3.1 Let (x, σ) be a forward complete trajectory of (1) with $\sigma \in S_a$ that is precompact relative to $\mathcal{O} \subset \mathbb{R}^n$. Then $\Omega^{\sharp}(x, \sigma) \subset \operatorname{dom}(f) \cap (\mathcal{O} \times \Gamma)$ and $\Omega(x) = \pi_1(\Omega^{\sharp}(x, \sigma))$.

Proof. That $\Omega^{\sharp}(x,\sigma) \subset \operatorname{dom}(f) \cap (\mathcal{O} \times \Gamma)$ follows from the fact that for all $t \in I_x(x(t),\sigma(t))$ belongs to a compact subset of $\operatorname{dom}(f) \cap (\mathcal{O} \times \Gamma)$ and from the definition of $\Omega^{\sharp}(x,\sigma)$. The proof of the other assertion follows *mutatis mutandis* from the proof of Lemma 4.1 in [16].

In order to see that the set $\Omega^{\sharp}(x,\sigma)$ enjoys certain kind of invariance property, let us introduce the following **Definition 3.2** Given a family \mathcal{T}^* of complete trajectories of (1), we say that a nonempty subset $M \subset \mathbb{R}^n \times \Gamma$ is weakly-invariant w.r.t \mathcal{T}^* if for each $(\xi, \gamma) \in M$ there is a trajectory $(x, \sigma) \in \mathcal{T}^*$ such that $x(0) = \xi$, $\sigma(0) = \gamma$ and $(x(t), \sigma(t)) \in M$ for all $t \in \mathbb{R}$.

This notion of weak invariance differs from the one introduced in [16], in that the last one involves only forward invariance while the introduced here also involves backward invariance.

Let us introduce now the following class of switching signals.

Definition 3.3 We say that a family of switching signals S^* has the property **P** if

- 1. $\mathcal{S}^* \subset \mathcal{S}_a[\tau_D, N_0]$ for some $\tau_D > 0$ and some $N_0 \in \mathbb{N}$;
- 2. for any s > 0 and any $\sigma \in \mathcal{S}^*$, $\sigma(\cdot + s) \in \mathcal{S}^*$;
- 3. for every sequence $\{\sigma_k\} \subset S^*$, there exist $\sigma^* \in S^*$ and a subsequence $\{\sigma_{k_l}\}$ such that $\lim_{l\to\infty} \sigma_{k_l}(t) = \sigma^*(t)$ for almost all $t \in \mathbb{R}$.

Lemma 3.2 The following classes of switching signals have the property P:

- 1. $S_a[\tau_D, N_0]$ for every $\tau_D > 0$ and every $N_0 \in \mathbb{N}$;
- 2. $\mathcal{S}_d[\tau_D] \cap \mathcal{S}^H$ for all $\tau_D > 0$ and every $H : \Gamma \rightsquigarrow \Gamma$ such that the set $\operatorname{Graph}(H) = \{(\gamma, \gamma') \in \Gamma \times \Gamma : \gamma' \in H(\gamma)\}$ is closed;

3. $S_d[\tau_D] \cap S_e[T]$ for every $\tau_D > 0$ and every T > 0.

Proof. See Appendix.

The next result will be instrumental in what follows.

Theorem 3.1 Let S^* be a family of switching signals which verifies property \mathbf{P} and let \mathcal{T}^* be the set of all the complete trajectories $(\overline{x}, \overline{\sigma})$ of (1) with $\overline{\sigma} \in S^*$. Then, if (x, σ) is a precompact forward complete trajectory of (1) such that $\sigma \in S^*$, $\Omega^{\sharp}(x, \sigma)$ is weakly-invariant w.r.t \mathcal{T}^* .

Proof. See Appendix.

Remark 3.1 Since the weak invariance of $\Omega^{\sharp}(x, \sigma)$ is a cornerstone of the results that we present below (Theorems 3.2 to 3.4), Theorem 3.1 enables us to obtain in a unified way invariance results not only for the different switching signals explicitly mentioned in Lemma 3.2 but also for any other that verifies property **P**.

Remark 3.2 At first glance, it would seem more natural to consider for a given precompact forward complete trajectory (x, σ) of (1) its ω -limit set $\Omega(x, \sigma)$ instead of $\Omega^{\sharp}(x, \sigma) \subset \Omega(x, \sigma)$. Nevertheless, there exist forward complete trajectories (x, σ) of (1) with $\sigma \in S_a$ such that $\Omega(x, \sigma)$ is not weaklyinvariant for any family of trajectories of that switched system.

Next, we present two invariance results that involve the existence of a function V which is nonincreasing along a trajectory of (1). In order to do so, we introduce the following class of functions.

Definition 3.4 We say that a function $V : \operatorname{dom}(V) \to \mathbb{R}$ belongs to class \mathcal{V} , if it verifies

- 1. dom $(V) \subset \mathbb{R}^n \times \Gamma$.
- 2. For every $\gamma \in \Gamma$, $\mathcal{D}_{\gamma} := \{\xi \in \mathbb{R}^n : (\xi, \gamma) \in \operatorname{dom}(V)\}$ is an open set.
- 3. Let $\mathcal{O} := \pi_1(\operatorname{dom}(V))$. Then $\mathcal{O}_{\gamma} := \mathcal{O} \cap \chi_{\gamma} \subset \mathcal{D}_{\gamma} \ \forall \gamma \in \Gamma$
- 4. For all $\gamma \in \Gamma$, $V_{\gamma}(\cdot) := V(\cdot, \gamma)$ is differentiable on \mathcal{O}_{γ} .

We note that $\operatorname{dom}(f) \cap (\mathcal{O} \times \Gamma) = \bigcup_{\gamma \in \Gamma} (\mathcal{O}_{\gamma} \times \{\gamma\}) \subset \operatorname{dom}(f) \cap \operatorname{dom}(V).$

We also note that when Γ is finite, the restriction of any function $V \in \mathcal{V}$ to dom $(f) \cap (\mathcal{O} \times \Gamma)$ is continuous.

In what follows, for a function $V \in \mathcal{V}$, let $Z_V = \{(\xi, \gamma) \in \text{dom}(f) \cap (\mathcal{O} \times \Gamma) : \nabla V_{\gamma}(\xi) f_{\gamma}(\xi) = 0\}.$

Assumption 1 The forward complete trajectory (x, σ) of (1) verifies the following: there exists a function $V \in \mathcal{V}$ whose restriction to dom $(f) \cap (\mathcal{O} \times \Gamma)$ is continuous, (x, σ) is precompact relative to \mathcal{O} and $v(t) = V(x(t), \sigma(t))$ is nonincreasing on $[0, +\infty)$.

Theorem 3.2 Let S^* be a family of switching signals which has property **P** and let \mathcal{T}^* be the set of all the complete trajectories (x, σ) of (1) with $\sigma \in S^*$. Suppose that (x, σ) , with $\sigma \in S^*$, is a forward complete trajectory of (1) for which Assumption 1 holds. Then there exists $c \in \mathbb{R}$ such that $x \to \pi_1(M(c))$, where M(c) is the maximal weakly-invariant set w.r.t. \mathcal{T}^* contained in $V^{-1}(c) \cap Z_V$.

Proof. Since $\Omega^{\sharp}(x,\sigma)$ is weakly-invariant w.r.t. \mathcal{T}^* and, from Lemma 3.1, $x \to \pi_1(\Omega^{\sharp}(x,\sigma))$, we only have to prove that $\Omega^{\sharp}(x,\sigma) \subset V^{-1}(c) \cap Z_V$ for some $c \in \mathbb{R}$.

As (x, σ) is precompact relative to \mathcal{O} , there exists a compact set $B \subset \mathcal{O}$ such that $x(t) \in B$ for all $t \in I_x$. Therefore $(x(t), \sigma(t))$ belongs to the compact set dom $(f) \cap (B \times \Gamma)$ for all $t \in I_x$. Thus v(t) is bounded, since V is continuous on dom $(f) \cap (B \times \Gamma)$, and nonincreasing by hypothesis; in consequence there exists $\lim_{t\to+\infty} v(t) = c$.

Let $(\xi, \gamma) \in \Omega^{\sharp}(x, \sigma)$. Then there exists a strictly increasing and unbounded sequence $\{s_k\}$ which verifies 1. and 2. of Definition 3.1. Since $(x(s_k), \sigma(s_k)) \to (\xi, \gamma)$ as $k \to \infty$, $c = \lim_{k \to \infty} v(s_k) = \lim_{k \to \infty} V(x(s_k), \sigma(s_k))$ $= V(\xi, \gamma)$ and $(\xi, \gamma) \in V^{-1}(c)$. Let us show that (ξ, γ) also belongs to Z_V .

As $\Omega^{\sharp}(x,\sigma)$ is weakly-invariant w.r.t. \mathcal{T}^* there exists $(x^*,\sigma^*) \in \mathcal{T}^*$ such that $(x^*(0),\sigma^*(0)) = (\xi,\gamma)$ and $(x^*(t),\sigma^*(t)) \in \Omega^{\sharp}(x,\sigma)$ for all $t \in \mathbb{R}$. Then, taking into account that $\Omega^{\sharp}(x,\sigma) \subset V^{-1}(c)$, $V(x^*(t),\sigma^*(t)) = c$ for all $t \in \mathbb{R}$. In particular, since $\sigma^*(t) = \gamma$ on $[0,\tau)$ for τ small enough, then $V(x^*(t),\gamma) = c$ on $[0,\tau)$. Therefore $\nabla V_{\gamma}(\xi) f_{\gamma}(\xi) = 0$, and $(\xi,\gamma) \in Z_V$.

Remark 3.3 We note that Theorem 3.2 is an extension to switched systems of the well known LaSalle's invariance principle for differential equations (see, for example, [9, Theorem 6.4]).

In the sequel, for any $\sigma \in S$ and any $\gamma \in \Gamma$, let $\mathcal{I}_{\sigma,\gamma} = \sigma^{-1}(\gamma) \cap [0, +\infty)$.

When Γ is a finite set, we can relax the nonincreasing condition in Assumption 1 as follows.

Assumption 2 The forward complete trajectory (x, σ) of (1) verifies the following: there exists a function $V \in \mathcal{V}$ such that (x, σ) is precompact relative to \mathcal{O} and $v(t) = V(x(t), \sigma(t))$ is nonincreasing on $\mathcal{I}_{\sigma,\gamma}$, for all $\gamma \in \Gamma$.

Remark 3.4 Assumptions of this kind are standard when the stability analysis of the zero solution of a switched system is performed by means of multiple Lyapunov functions (see [4], [11]).

In what follows, when Γ is a finite set, we identify it with the set $\{1, \ldots, N\} \subset \mathbb{N}$, where $N = \operatorname{card}(\Gamma)$.

Theorem 3.3 Suppose that Γ is finite and let S^* and \mathcal{T}^* be as in Theorem 3.2. Suppose that (x, σ) , with $\sigma \in S^*$, is a forward complete trajectory of (1) for which Assumption 2 holds. Then there exists $\vec{c} = (c_1, \ldots, c_N) \in \mathbb{R}^N$ such that $x \to \pi_1(M(\vec{c}))$, where $M(\vec{c})$ is the maximal weakly-invariant set w.r.t. \mathcal{T}^* contained in $\bigcup_{\gamma \in \Gamma} \{(\xi, \gamma) \in \operatorname{dom}(f) \cap (\mathcal{O} \times \Gamma) : V_{\gamma}(\xi) = c_{\gamma}\} \cap Z_V.$

Proof. For $\gamma \in \Gamma$ we define c_{γ} as follows:

- 1. $c_{\gamma} = \lim_{t \to +\infty, t \in \mathcal{I}_{\sigma,\gamma}} v(t)$ if $\mathcal{I}_{\sigma,\gamma}$ is unbounded. (This limit exists since v is non-increasing and bounded on $\mathcal{I}_{\sigma,\gamma}$).
- 2. $c_{\gamma} = a$ for every γ such that $\mathcal{I}_{\sigma,\gamma}$ is bounded. Here $a \in \mathbb{R}$ is arbitrary.

Reasoning as in the proof of Theorem 3.2, in order to prove the thesis it suffices to show that $\Omega^{\sharp}(x,\sigma) \subset \bigcup_{\gamma \in \Gamma} \{(\xi,\gamma) \in \operatorname{dom}(f) \cap (\mathcal{O} \times \Gamma) : V_{\gamma}(\xi) = c_{\gamma} \} \cap Z_{V}.$

Let $(\xi, \gamma) \in \Omega^{\sharp}(x, \sigma)$. Then there exists a strictly increasing and unbounded sequence $\{s_k\}$ which verifies 1. and 2. of Definition 3.1. Since $\sigma(s_k) \to \gamma$ and Γ is a finite set, $\sigma(s_k) = \gamma$ for k large enough and for those $k, s_k \in \mathcal{I}_{\sigma,\gamma}$. It follows that $V(x(s_k), \gamma) \to c_{\gamma}$ as $k \to \infty$ and in consequence, $V(\xi, \gamma) = c_{\gamma}$ and $(\xi, \gamma) \in \bigcup_{\gamma' \in \Gamma} \{(\xi', \gamma') \in \operatorname{dom}(f) \cap (\mathcal{O} \times \Gamma) : V(\xi', \gamma') = c_{\gamma'} \}$. That $(\xi, \gamma) \in Z_V$ can be proved in the same way as in the proof of Theorem 3.2.

Remark 3.5 Some invariance results for switched systems reported in the literature can be derived from Theorem 3.3. In particular [1, Theorems 1 and 2], [16, Proposition 4.1] and [5, Corollary 5.6].

The following invariance result involves weakly meagre functions. We recall that a function $y : \mathbb{R}_{\geq 0} \to \mathbb{R}$ is weakly meagre if $\lim_{k\to\infty} (\inf\{|y(t)| : t \in I_k\}) = 0$ for every family $\{I_k : k \in \mathbb{N}\}$ of nonempty and pairwise disjoint intervals in $\mathbb{R}_{\geq 0}$ with $\inf\{\mu(I_k) : k \in \mathbb{N}\} > 0$, where μ stands for the Lebesgue measure (see [13]). We note that, for example, any function $y \in L^p([0,\infty))$ with p > 0 is weakly meagre. More generally, if there exist positive numbers τ and p such that $\int_t^{t+\tau} |y(s)|^p ds$ converges to 0 as $t \to +\infty$, then y is weakly meagre.

Theorem 3.4 Let S^* and T^* be as in Theorem 3.2. Suppose that (x, σ) , with $\sigma \in S^*$, is a forward complete trajectory of (1). Suppose in addition that there exists a continuous function $h : \operatorname{dom}(h) \to \mathbb{R}$, with $\operatorname{dom}(h) \subset \mathbb{R}^n \times \Gamma$, such that $(x(t), \sigma(t))$ evolves in a compact subset K of $\operatorname{dom}(h)$ for all $t \ge 0$ and that $y(\cdot) = h(x(\cdot), \sigma(\cdot))$ is weakly meagre. Then $x \to \pi_1(M^*)$, where M^* is the maximal weakly invariant set w.r.t. T^* contained in $h^{-1}(0) \cap \operatorname{dom}(f)$.

Proof. Since $(x(t), \sigma(t))$ evolves in the compact subset K of dom(h) for all $t \ge 0$, we have that (x, σ) is precompact and that $\Omega^{\sharp}(x, \sigma) \subset \operatorname{dom}(h) \cap$ dom(f). By similar considerations as those in the proofs of the previous invariance results, it suffices to show that $\Omega^{\sharp}(x, \sigma) \subset h^{-1}(0)$.

Let $(\xi^*, \gamma^*) \in \Omega^{\sharp}(x, \sigma)$. Then there exists a strictly increasing and unbounded sequence $\{s_k\}$ which verifies 1. and 2. of Definition 3.1 with (ξ^*, γ^*) instead of (ξ, γ) . We can assume that $\tau^1_{\sigma}(s_k) - s_k \geq 3r/4$ for all k.

We will construct by recursion a sequence of times $\{s_m^*\}$ and a subsequence $\{s_{k_m}\}$ of $\{s_k\}$ such that $s_{k_m} \leq s_m^* \leq s_{k_m} + 2^{-m}r$ and $|y(s_m^*)| \leq 1/m$ for all m.

For any $m \in \mathbb{N}$, let $r_m = 2^{-m}r$ and $I_k^m = [s_k, s_k + r_m]$ for all $k \in \mathbb{N}$. Since *h* is weakly meagre, $\lim_{k\to\infty} (\inf\{|y(t)| : t \in I_k^1\}) = 0$. Thus, there exist $k^* \in \mathbb{N}$ an a time $t \in I_{k^*}^1$ such that $|y(t)| \leq 1$. Let $s_1^* = t$ and $k_1 = k^*$.

Suppose that we have already defined $\{s_m^*\}_{m=1}^l$ and $\{s_{k_m}\}_{m=1}^l$. As h is weakly meagre then $\lim_{k\to\infty} (\inf\{|y(t)|:t\in I_k^{l+1}\}) = 0$, and hence there exist $k' \in \mathbb{N}$, with $k' > k_l$, and $t' \in I_{k'}^{l+1}$ such that $|y(t')| \leq 1/(l+1)$. We define $s_{l+1}^* = t'$ and $k_{l+1} = k'$.

Consider now the sequence $\{(x(s_m^*), \sigma(s_m^*))\}$; since $s_m^* \in [s_{k_m}, s_{k_m} + r_m]$, $r_m \leq r/2$ and $\tau_{\sigma}^1(s_{k_m}) - s_{k_m} \geq 3r/4$, then for all $m, \sigma(s_m^*) = \sigma(s_{k_m})$ and hence $\sigma(s_m^*) \to \gamma^*$.

Given that for every $t \ge 0$, $(x(t), \sigma(t))$ belongs to the compact set dom $(f) \cap K$ and since f is continuous on that set, there exists $M \ge 0$ such that $|\dot{x}(t)| \le M$ a.e. on I_x and in consequence, $|x(t) - x(s)| \le M|t - s|$ for all $t, s \in I_x$. Therefore $|x(s_m^*) - x(s_{k_m})| \le rM2^{-m}$ for all m. Taking into account that $x(s_{k_m}) \to \xi^*$, we have that $x(s_m^*) \to \xi^*$.

Finally, since by construction $h(x(s_m^*), \sigma(s_m^*)) = y(s_m^*) \to 0$ and h is continuous, then $h(\xi^*, \gamma^*) = 0$.

4 Convergence and stability results

In this section we derive, from the invariance principles presented in §3, some convergence and stability results for switched systems with constrained switchings.

4.1 Convergence results

Let us first introduce some observability-like definitions.

Given a subset $\mathcal{X} \subset \mathbb{R}^n$, a continuous map $g : \mathcal{X} \to \mathbb{R}^n$ and a function $h : \mathcal{X} \to \mathbb{R}$, we say that for a given τ ($\tau > 0$ or $\tau = \infty$) a point $\xi \in \mathcal{X}$ belongs to the set $\mathcal{X}^f(g, h, \tau)$ (resp. $\mathcal{X}^b(g, h, \tau)$) if there exists a solution $\varphi : [0, \tau] \to \mathcal{X}$ (resp. $\varphi : [-\tau, 0] \to \mathcal{X}$) of $\dot{x} = g(x)$ such that $\varphi(0) = \xi$ and $h(\varphi(t)) = 0$ for all $t \in [0, \tau]$ (resp. $t \in [-\tau, 0]$).

Let also the sets $\mathcal{X}^{f}(g,h) = \bigcup_{\tau>0} \mathcal{X}^{f}(g,h,\tau), \mathcal{X}^{b}(g,h) = \bigcup_{\tau>0} \mathcal{X}^{b}(g,h,\tau)$ and $\mathcal{X}(g,h) = \mathcal{X}^{f}(g,h) \cup \mathcal{X}^{b}(g,h).$

Remark 4.1

1. The set $\mathcal{X}^{f}(g,h,\infty)$ ($\mathcal{X}^{b}(g,h,\infty)$) coincides with the maximal weakly forward(backward) invariant set w.r.t. g contained in the set { $\xi \in \mathcal{X}$: $h(\xi) = 0$ }.

We recall that a subset $K \subset \mathbb{R}^n$ is weakly forward(backward) invariant w.r.t g if for each $\xi \in K$ there exists a solution $\varphi : [0, \infty) \to \mathbb{R}^n$ $(\varphi : (-\infty, 0] \to \mathbb{R}^n)$ of $\dot{x} = g(x)$ such that $\varphi(0) = \xi$ and $\varphi(t) \in K$ for all $t \ge 0$ ($t \le 0$).

- 2. If we consider the system with outputs $\dot{x} = g(x)$, y = h(x) and state space \mathcal{X} , with $0 \in \mathcal{X}$, g(0) = 0 and h(0) = 0, then the set $\mathcal{X}^{f}(g,h)$ coincides with the set of states ξ that cannot be instantaneously distinguished from the zero state through the output y. In the particular case in which g is a linear function, i.e., $g(\xi) = A\xi$ and $h(\xi) = \xi^{T}C^{T}C\xi$, and C is a matrix, then $\mathcal{X}(g,h) \subset \mathcal{X} \cap \mathcal{U}$, being \mathcal{U} the unobservable subspace of (C, A).
- 3. When g and h are smooth functions we have that

$$\mathcal{X}(g,h) \subset \{\xi \in \mathcal{X} : L^k_a h(\xi) = 0 \; \forall k \in \mathbb{N}_0\},\$$

with $L_a^k h$ the k-th. Lie derivative of h along g.

Let us introduce the following assumptions, in order to obtain some convergence criteria based on the invariance results given in §3 and on the observability-like notions already introduced.

Assumption 3 For the forward complete trajectory (x, σ) of (1) there exist a function $V \in \mathcal{V}$ and a family of functions $\{W_{\gamma} : \mathcal{O}_{\gamma} \to \mathbb{R}, \gamma \in \Gamma\}$ such that (x, σ) and V satisfy Assumption 1 and in addition

$$-\nabla V_{\gamma}(\xi) f_{\gamma}(\xi) \ge W_{\gamma}(\xi) \ge 0 \quad \forall \xi \in \mathcal{O}_{\gamma}, \quad \forall \gamma \in \Gamma.$$
(4)

Assumption 4 For the forward complete trajectory (x, σ) of (1) there exist a function $V \in \mathcal{V}$ which is bounded on compact subsets of $\operatorname{dom}(f) \cap (\mathcal{O} \times \Gamma)$ and a family of functions $\{W_{\gamma} : \mathcal{O}_{\gamma} \to \mathbb{R}, \gamma \in \Gamma\}$ such that (x, σ) is precompact relative to $\mathcal{O}, v(t) = V(x(t), \sigma(t))$ is nonincreasing on $[0, +\infty)$, (4) holds and in addition $W(\xi, \gamma) := W_{\gamma}(\xi)$ is continuous on $\operatorname{dom}(W) = \operatorname{dom}(f) \cap (\mathcal{O} \times \Gamma)$.

Assumption 5 For the forward complete trajectory (x, σ) of (1) there exist a function $V \in \mathcal{V}$ and a family of functions $\{W_{\gamma} : \mathcal{O}_{\gamma} \to \mathbb{R}, \gamma \in \Gamma\}$ such that (x, σ) and V satisfy Assumption 2 and in addition (4) holds.

Theorem 4.1 Let (x, σ) be a forward complete trajectory of (1) with $\sigma \in S_a$. Then the following holds: 1. if (x, σ) verifies Assumption 3, then there exists $c \in \mathbb{R}$ such that

$$x \to \bigcup_{\gamma,\gamma' \in \Gamma} \left(\mathcal{O}^f_{\gamma}(f_{\gamma}, W_{\gamma}) \cap \mathcal{O}^b_{\gamma'}(f_{\gamma'}, W_{\gamma'}) \cap V_{\gamma}^{-1}(c) \cap V_{\gamma'}^{-1}(c) \right);$$

2. if (x, σ) verifies Assumption 4, then

$$x \to \bigcup_{\gamma,\gamma' \in \Gamma} \left(\mathcal{O}^f_{\gamma}(f_{\gamma}, W_{\gamma}) \cap \mathcal{O}^b_{\gamma'}(f_{\gamma'}, W_{\gamma'}) \right);$$

3. if Γ is finite and (x, σ) verifies Assumption 5, then there exists $\vec{c} \in \mathbb{R}^N$ such that

$$x \to \bigcup_{\gamma,\gamma' \in \Gamma} \left(\mathcal{O}^f_{\gamma}(f_{\gamma}, W_{\gamma}) \cap \mathcal{O}^b_{\gamma'}(f_{\gamma'}, W_{\gamma'}) \cap V_{\gamma}^{-1}(c_{\gamma}) \cap V_{\gamma'}^{-1}(c_{\gamma'}) \right);$$

Proof. Since $\sigma \in S_a$, there exist $\tau_D > 0$ and $N_0 \in \mathbb{N}$ such that $\sigma \in S_a[\tau_D, N_0]$.

In order to prove 1, let (x, σ) verify Assumption 3. As $S_a[\tau_D, N_0]$ has property **P** and (x, σ) verifies the hypotheses of Theorem 3.2, there exists a real number c such that $x \to \pi_1(M(c))$, where M(c) is the maximal weaklyinvariant set w.r.t. $\mathcal{T}_a[\tau_D, N_0]$ contained in $V^{-1}(c) \cap Z_V$.

Let $\xi \in \pi_1(M(c))$ and $\gamma \in \Gamma$ such that $(\xi, \gamma) \in M(c)$. From the weak invariance of M(c) w.r.t. $\mathcal{T}_a[\tau_D, N_0]$, there exists a trajectory $(x^*, \sigma^*) \in$ $\mathcal{T}_a[\tau_D, N_0]$ such that $(x^*(0), \sigma^*(0)) = (\xi, \gamma)$ and such that for every $t \in$ \mathbb{R} , $\nabla V_{\sigma^*(t)}(x^*(t))f_{\sigma^*(t)}(x^*(t)) = 0$ and $V_{\sigma^*(t)}(x^*(t)) = c$. Since for every $\gamma \in \Gamma$, $V_{\gamma}(\cdot)$ is continuous on \mathcal{O}_{γ} , we also have that $V_{\sigma^*(t^{-})}(x^*(t)) = c$ and $\nabla V_{\sigma^*(t^{-})}(x^*(t))f_{\sigma^*(t^{-})}(x^*(t)) = 0$ for all $t \in \mathbb{R}$. In view of (4), $W_{\sigma^*(t)}(x^*(t)) =$ $W_{\sigma^*(t^{-})}(x^*(t)) = 0$ for all $t \in \mathbb{R}$. Let us consider two cases.

Case 1. $0 \notin \Lambda(\sigma^*)$. Then, there exist $\tau_1 < 0 < \tau_2$ such that $\sigma^*(t) = \gamma$ for every $t \in [\tau_1, \tau_2]$. Hence, $\xi = x^*(0) \in \mathcal{O}^f_{\gamma}(f_{\gamma}, W_{\gamma}, \tau_2) \cap \mathcal{O}^b_{\gamma}(f_{\gamma}, W_{\gamma}, -\tau_1) \cap V_{\gamma}^{-1}(c)$.

Case 2. $0 \in \Lambda(\sigma^*)$. Let $\gamma' = \sigma^*(0^-)$; then, there exist $\tau_1 < 0 < \tau_2$ such that $\sigma^*(t) = \gamma'$ for all $t \in [\tau_1, 0)$ and $\sigma^*(t) = \gamma$ for all $t \in [0, \tau_2]$. In consequence, $\xi = x^*(0) \in \mathcal{O}^f_{\gamma}(f_{\gamma}, W_{\gamma}, \tau_2) \cap \mathcal{O}^b_{\gamma'}(f_{\gamma'}, W_{\gamma'}, -\tau_1) \cap V_{\gamma}^{-1}(c) \cap V_{\gamma'}^{-1}(c)$.

The proof of 3. is similar to that of 1. and we omit it.

We now demonstrate 2. Suppose that (x, σ) verifies Assumption 4; since (x, σ) is precompact relative to \mathcal{O} , there is a compact set $B \subset \mathcal{O}$ such that $x(t) \in B$ for all $t \geq 0$ and therefore $(x(t), \sigma(t))$ belongs to the compact set $\operatorname{dom}(f) \cap (B \times \Gamma) \subset \operatorname{dom}(W)$ for all $t \geq 0$. That $y(t) = W(x(t), \sigma(t))$ is

weakly meagre follows from the fact that $\int_0^\infty y(t)dt$ is finite. Let us prove this last fact.

As $v(t) = V(x(t), \sigma(t))$ in nonincreasing and differentiable on each finite interval [a, b) in which σ is constant, (say $\sigma(t) = \gamma$ for all $t \in [a, b)$), from [20, Theorem 7.21] we have that

$$v(a) - v(b^-) \ge \int_a^b (-\dot{v})(s) ds = -\int_a^b \nabla V_\gamma(x(s)) f_\gamma(x(s)) ds.$$

Taking into account (4) and that v is nonincreasing on $[0, \infty)$, we then have that

$$0 \le \int_0^t y(s) ds \le V(x(0), \sigma(0)) - V(x(t), \sigma(t)) \quad \forall t \ge 0.$$

As $(x(t), \sigma(t))$ evolves in the compact set $\operatorname{dom}(f) \cap (B \times \Gamma)$ for all $t \ge 0$ and V is bounded on compact subsets of $\operatorname{dom}(f) \cap (\mathcal{O} \times \Gamma)$ then, for some $M \ge 0$, $|V(x(t), \sigma(t))| \le M$ for all $t \ge 0$. Thus $\int_0^t y(s) ds \le 2M$ for all $t \ge 0$ and the integrability of y follows.

Since (x, σ) verifies the hypotheses of Theorem 3.4 with W instead of h, $x \to \pi_1(M^*)$ where M^* is the maximal weakly-invariant set w.r.t. $\mathcal{T}_a[\tau_D, N_0]$ contained in dom $(f) \cap W^{-1}(0)$.

In order to prove that $\pi_1(M^*) \subset \bigcup_{\gamma,\gamma'\in\Gamma} (\mathcal{O}^f_{\gamma}(f_{\gamma},W_{\gamma})\cap \mathcal{O}^b_{\gamma'}(f_{\gamma'},W_{\gamma'}))$, we can proceed as in the proof of 1., but using now the fact that for all $(\xi,\gamma)\in M^*$ there exists $(x^*,\sigma^*)\in\mathcal{T}_a[\tau_D,N_0]$ such that $0=W_{\sigma^*(t)}(x^*(t))=W_{\sigma^*(t^-)}(x^*(t))$ for all $t\in\mathbb{R}$ (being the last equality due to the continuity of W on its domain).

Remark 4.2 Theorem 4.1 gives a more accurate result than Theorem 8 in [6] in the case when the switching signal $\sigma \in S_a$ (instead of $\sigma \in S_{p-dwell}$ as is considered there). In fact, it can be shown that the hypotheses of that theorem imply that the forward complete trajectory (x, σ) of the linear switched system $\dot{x} = A_{\sigma(t)}x$ is precompact and verifies Assumption 4 with $V(\xi, \gamma) = \xi^T P_{\gamma}\xi$ and $W(\xi, \gamma) = \xi^T C_{\gamma}^T C_{\gamma}\xi$. So, by applying Theorem 4.1, and taking into account 2. of Remark 4.1, it results that $x \to \bigcup_{\gamma \in \Gamma} \mathcal{U}_{\gamma}$, where \mathcal{U}_{γ} is the unobservable subspace of the pair (C_{γ}, A_{γ}) . On the other hand, Theorem 8 in [6] asserts that $x \to \mathcal{M}$, where \mathcal{M} is the smallest subspace which contains $\bigcup_{\gamma \in \Gamma} \mathcal{U}_{\gamma}$ and is A_{γ} -invariant for all $\gamma \in \Gamma$.

Remark 4.3 If in addition to the hypotheses of Theorem 4.1, we have that for some $x_e \in \bigcup_{\gamma \in \Gamma} \mathcal{O}_{\gamma}$, either for all $\gamma \in \Gamma$, $\mathcal{O}_{\gamma}^f(f_{\gamma}, W_{\gamma}) \subset \{x_e\}$ or for all $\gamma \in \Gamma$, $\mathcal{O}_{\gamma}^b(f_{\gamma}, W_{\gamma}) \subset \{x_e\}$, then $x \to x_e$. The first conclusion of Corollary 4.10 in [5] is a particular case of this result. We note that, according to the particular geometry of each χ_{γ} , it could happen that $\mathcal{O}_{\gamma}^{b}(f_{\gamma}, W_{\gamma}) \neq \mathcal{O}_{\gamma}^{f}(f_{\gamma}, W_{\gamma})$ and even that one of those sets be void and the other one not.

In what follows let for each $\gamma \in \Gamma$, $E_{\gamma} = \{\xi \in \chi_{\gamma} : f_{\gamma}(\xi) = 0\}$ the set of equilibrium points of f_{γ} .

The following convergence result involves an "ergodicity" condition on the switching signals considered.

Theorem 4.2 Suppose that Γ is a finite set. Let (x, σ) , with $\sigma \in S_e \cap S_d$, be a forward complete trajectory of (1). Then the following holds:

- 1. if (x, σ) verifies Assumption 3 and if for every $\gamma \in \Gamma$, either $\mathcal{O}^b_{\gamma}(f_{\gamma}, W_{\gamma})$ $= E_{\gamma} \cap \mathcal{O}_{\gamma} \text{ or } \mathcal{O}^f_{\gamma}(f_{\gamma}, W_{\gamma}) = E_{\gamma} \cap \mathcal{O}_{\gamma}, \text{ then there exists } c \in \mathbb{R} \text{ such that}$ $x \to \bigcap_{\gamma \in \Gamma} (E_{\gamma} \cap V_{\gamma}^{-1}(c)).$ If, in addition, for each $c \in \mathbb{R}, \ \bigcap_{\gamma \in \Gamma} (E_{\gamma} \cap V_{\gamma}^{-1}(c))$ is a discrete set, then $x \to x_e$ for some $x_e \in \bigcap_{\gamma \in \Gamma} (E_{\gamma} \cap \mathcal{O}_{\gamma}).$
- 2. If (x, σ) verifies Assumption 5 and if for every $\gamma \in \Gamma$, either $\mathcal{O}^b_{\gamma}(f_{\gamma}, W_{\gamma})$ $= E_{\gamma} \cap \mathcal{O}_{\gamma} \text{ or } \mathcal{O}^f_{\gamma}(f_{\gamma}, W_{\gamma}) = E_{\gamma} \cap \mathcal{O}_{\gamma}, \text{ then } x \to \cap_{\gamma \in \Gamma}(E_{\gamma} \cap \mathcal{O}_{\gamma}).$ If, in addition, $\cap_{\gamma \in \Gamma}(E_{\gamma} \cap \mathcal{O}_{\gamma})$ is a discrete set, then $x \to x_e$ for some $x_e \in \cap_{\gamma \in \Gamma}(E_{\gamma} \cap \mathcal{O}_{\gamma})$

Proof. As $\sigma \in S_e \cap S_d$, then there exist T > 0 and $\tau_D > 0$ such that $\sigma \in S_e[T] \cap S_d[\tau_D]$.

Let us prove 1. first. Since $S_e[T] \cap S_d[\tau_D]$ has property **P** and (x, σ) verifies the hypotheses of Theorem 3.2, there exists $c \in \mathbb{R}$ such that $x \to \pi_1(M(c))$, where M(c) is the maximal weakly-invariant set w.r.t. $\mathcal{T}_e[T] \cap \mathcal{T}_d[\tau_D]$ contained in $V^{-1}(c) \cap Z_V$. So, it suffices to show that $\pi_1(M(c)) \subset \bigcap_{\gamma \in \Gamma} (E_\gamma \cap V_\gamma^{-1}(c))$.

Let $(\xi, \gamma) \in M(c)$; then there exists a trajectory $(x^*, \sigma^*) \in \mathcal{T}_e[T] \cap \mathcal{T}_d[\tau_D]$ such that $(x^*(0), \sigma^*(0)) = (\xi, \gamma)$ and $(x^*(t), \sigma^*(t)) \in M(c)$ for all $t \in \mathbb{R}$. Then, reasoning as in the proof of Theorem 4.1, $V_{\sigma^*(t^-)}(x^*(t)) \equiv V_{\sigma^*(t)}(x^*(t)) \equiv c$ and, from (4), $W_{\sigma^*(t^-)}(x^*(t)) \equiv W_{\sigma^*(t)}(x^*(t)) \equiv 0$.

Taking into account that x^* is continuous and that either $\mathcal{O}^b_{\gamma}(f_{\gamma}, W_{\gamma}) = E_{\gamma} \cap \mathcal{O}_{\gamma}$ or $\mathcal{O}^f_{\gamma}(f_{\gamma}, W_{\gamma}) = E_{\gamma} \cap \mathcal{O}_{\gamma}$, it follows that $x^*(t) \equiv \xi$ and that $\xi \in E_{\sigma^*(t)} \cap \mathcal{O}_{\sigma^*(t)} \cap V^{-1}_{\sigma^*(t)}(c) = E_{\sigma^*(t)} \cap V^{-1}_{\sigma^*(t)}(c)$ for all $t \in \mathbb{R}$.

As $\sigma^* \in \mathcal{S}_e[T]$, then $\sigma^*(\mathbb{R}) = \Gamma$ and, in consequence, $\xi \in E_\gamma \cap V_\gamma^{-1}(c)$ for all $\gamma \in \Gamma$.

In the case that for every $c \in \mathbb{R}$, $\bigcap_{\gamma \in \Gamma} (E_{\gamma} \cap V_{\gamma}^{-1}(c))$ is a discrete set, that $x \to x_e$ with $x_e \in \bigcap_{\gamma \in \Gamma} (E_{\gamma} \cap \mathcal{O}_{\gamma})$, follows from the facts that $x \to \Omega(x)$, that $\Omega(x)$ is a nonempty connected set and that $\Omega(x) \subset \bigcap_{\gamma \in \Gamma} (E_{\gamma} \cap V_{\gamma}^{-1}(c))$ for some $c \in \mathbb{R}$. The proof of 2. is similar to that of 1., so we omit it.

In the sequel we give sufficient conditions for the convergence to a given equilibrium point x_e of (1), i.e. a point x_e that verifies $f_{\gamma}(x_e) = 0$ for all $\gamma \in \Gamma$ such that $x_e \in \chi_{\gamma}$. We assume, without loss of generality, that x_e is the origin.

Assumption 6 0 is an equilibrium point of (1).

Assumption 7 For every $\gamma \in \Gamma$ such that $0 \in \chi_{\gamma}$, the initial value problem $\dot{x} = f_{\gamma}(x), x(0) = 0$ has a unique solution.

Theorem 4.3 Suppose that assumptions 6 and 7 hold and let (x, σ) be a forward complete trajectory of (1) with $\sigma \in S_a$.

1. If Assumption 4 is verified, $0 \in \mathcal{O}$ and the following holds

(a)
$$\mathcal{O}^f_{\gamma}(f_{\gamma}, W_{\gamma}, \infty) \cap \mathcal{O}^b_{\gamma}(f_{\gamma}, W_{\gamma}, \infty) \subset \{0\} \text{ for every } \gamma \in \Gamma,$$

$$(b) \ \mathcal{O}^b_{\gamma}(f_{\gamma}, W_{\gamma}) \cap \mathcal{O}^f_{\gamma'}(f_{\gamma'}, W_{\gamma'}) \subset \{0\}, \quad \forall \gamma \neq \gamma',$$

then $x \to 0$.

If Γ is finite, the same holds if we suppose that (x, σ) verifies Assumption 5 instead of Assumption 4.

- 2. If Assumption 3 is verified, $0 \in \mathcal{O}$, 1.(i) holds and
 - (a) $\mathcal{O}^b_{\gamma}(f_{\gamma}, W_{\gamma}) \cap \mathcal{O}^f_{\gamma'}(f_{\gamma'}, W_{\gamma'}) \cap V_{\gamma}^{-1}(c) \cap V_{\gamma'}^{-1}(c) \subset \{0\}, \quad \forall \gamma \neq \gamma' \in \Gamma, \quad \forall c \in \mathbb{R},$

then $x \to 0$.

Proof. Since $\sigma \in S_a$, there exist $\tau_D > 0$ and $N_0 \in \mathbb{N}$ such that $\sigma \in S_a[\tau_D, N_0]$.

We first prove 2. By using the same arguments as in the proof of the first part of Theorem 4.1, it follows that there exists $c \in \mathbb{R}$ such that $x \to \pi_1(M(c))$, with M(c) the maximal weakly-invariant set w.r.t. $\mathcal{T}_a[\tau_D, N_0]$ contained in $V^{-1}(c) \cap Z_V$. So, it suffices to show that $M(c) \subset \{0\} \times \Gamma$.

Let $(\xi, \gamma) \in M(c)$; then there exists a trajectory $(x^*, \sigma^*) \in \mathcal{T}_a[\tau_D, N_0]$ such that $(x^*(0), \sigma^*(0)) = (\xi, \gamma)$ and $(x^*(t), \sigma^*(t)) \in M(c)$ for all $t \in \mathbb{R}$. Once again, as in Theorem 4.1 we have that $V_{\sigma^*(t^-)}(x^*(t)) \equiv V_{\sigma^*(t)}(x^*(t)) \equiv c$ and $W_{\sigma^*(t^-)}(x^*(t)) \equiv W_{\sigma^*(t)}(x^*(t)) \equiv 0$. We will consider two cases.

Case 1. σ^* has no switching times, i.e. $\sigma^*(t) = \gamma$ for all $t \in \mathbb{R}$. Then for every $t \in \mathbb{R}, x^*(t) \in \mathcal{O}_{\gamma}$ and $W_{\gamma}(\varphi(t)) = 0$ and, since $x^*(t) \in \mathcal{O}_{\gamma}^f(f_{\gamma}, W_{\gamma}, \infty) \cap \mathcal{O}_{\gamma}^b(f_{\gamma}, W_{\gamma}, \infty) \subset \{0\}, x^*(t) = 0$ for all t.

Case 2. σ^* has a switching time t^* . Then, there exists $\tau > 0$ such that, if $\varphi(t) = x^*(t+t^*), \ \hat{\gamma} = \sigma^*(t^{*-}) \text{ and } \gamma' = \sigma^*(t^*), \ \hat{\gamma} \neq \gamma' \text{ and}$

1. $\varphi : [-\tau, 0] \to \mathcal{O}_{\hat{\gamma}}$ is solution of $\dot{z} = f_{\hat{\gamma}}(z)$ on $[-\tau, 0]$ and $\varphi : [0, \tau] \to \mathcal{O}_{\gamma'}$ is solution of $\dot{z} = f_{\gamma'}(z)$ on $[0, \tau]$;

2.
$$V_{\hat{\gamma}}(\varphi(t)) = c \text{ on } [-\tau, 0] \text{ and } V_{\gamma'}(\varphi(t)) = c \text{ on } [0, \tau];$$

3. $W_{\hat{\gamma}}(\varphi(t)) = 0$ on $[-\tau, 0]$ and $W_{\gamma'}(\varphi(t)) = 0$ on $[0, \tau]$.

Thus $x^*(t^*) = \varphi(0) \in \mathcal{O}^b_{\hat{\gamma}}(f_{\hat{\gamma}}, W_{\hat{\gamma}}) \cap \mathcal{O}^f_{\gamma'}(f_{\gamma'}, W_{\gamma'}) \cap V^{-1}_{\hat{\gamma}}(c) \cap V^{-1}_{\gamma'}(c) \subset \{0\}.$

That $x^*(0) = 0$ follows from the fact that, due to assumptions 6 and 7, any initial value problem $\dot{z} = f_{\hat{\gamma}}(z), z(0) = 0$ has the unique solution $z(t) \equiv 0$ when $0 \in \mathcal{O}_{\hat{\gamma}}$.

In order to prove 1. we note that in the case in which Γ is finite and (x, σ) verifies Assumption 5 then, due to Theorem 3.3 (with $S^* = S_a[\tau_D, N_0]$), it suffices to show that for any $\vec{c} \in \mathbb{R}^N$, the set $M(\vec{c})$ of that theorem is a subset of $\{0\} \times \Gamma$. Since the proof of this fact is similar to that of $M(c) \subset \{0\} \times \Gamma$ given above, we omit it.

Suppose now that (x, σ) verifies Assumption 4. It follows from the proof of Theorem 4.1 that $y(t) = W(x(t), \sigma(t))$ is weakly meagre. Since W is continuous on dom $(f) \cap (\mathcal{O} \times \Gamma)$, (x, σ) verifies the hypotheses of Theorem 3.4 and in consequence $x \to \pi_1(M^*)$, being M^* the maximal weakly-invariant set w.r.t. $\mathcal{T}_a[\tau_D, N_0]$ contained in dom $(f) \cap W^{-1}(0)$. The proof of $M^* \subset \{0\} \times \Gamma$ in similar to that of $M(c) \subset \{0\} \times \Gamma$, so we only delineate it. Let $(\xi, \gamma) \in M^*$; then there exists $(x^*, \sigma^*) \in \mathcal{T}_a[\tau_D, N_0]$ such that $(x^*(0), \sigma^*(0)) = (\xi, \gamma)$ and, as in Theorem 4.1, $W_{\sigma^*(t^-)}(x^*(t)) = W_{\sigma^*(t)}(x^*(t)) = 0$ for all $t \in \mathbb{R}$. If σ^* has no switching times, then $x^*(0) = 0$ due to 1.(i).

If σ^* has a switching time t^* we considerer τ , φ , $\hat{\gamma}$ and γ' as in the proof of Case 2. above. Then φ verifies 1. of that case, $W_{\hat{\gamma}}(\varphi(t)) = 0$ on $[-\tau, 0]$ and $W_{\gamma'}(\varphi(t)) = 0$ on $[0, \tau]$. Thus, from 1.(ii) we deduce that $x^*(t^*) = \varphi(0) = 0$ and a posteriori that $x^*(0) = 0$.

When Γ is finite and σ belongs to $\mathcal{S}_d \cap \mathcal{S}^H$, hypothesis 2. of Theorem 4.3 can be weakened as follows.

Given a set-valued map $H : \Gamma \rightsquigarrow \Gamma$, a finite sequence $\{\gamma_i\}_{i=1}^m \subset \Gamma, m \geq 3$, is a simple cycle of H if $\gamma_1 = \gamma_m, \gamma_{i+1} \in H(\gamma_i)$ for all $i = 1, \ldots, m-1$ and if $\gamma_i = \gamma_j$ and i < j then i = 1 and j = m.

Theorem 4.4 Suppose that Γ is finite, that $H : \Gamma \rightsquigarrow \Gamma$ and that (x, σ) is a forward complete trajectory of (1) with $\sigma \in S_d \cap S^H$. Suppose in addition that assumptions 6 and 7 hold.

- 1. If Assumption 5 holds, $0 \in \mathcal{O}$ and
 - (a) $\mathcal{O}^f_{\gamma}(f_{\gamma}, W_{\gamma}, \infty) \subset \{0\}$ for every $\gamma \in \Gamma$ or $\mathcal{O}^b_{\gamma}(f_{\gamma}, W_{\gamma}, \infty) \subset \{0\}$ for every $\gamma \in \Gamma$,

(b) for each simple cycle $\{\gamma_i\}_{i=1}^m$ of H there exists $j \in \{1, \ldots, m-1\}$ such that

$$\mathcal{O}^{b}_{\gamma_{j}}(f_{\gamma_{j}}, W_{\gamma_{j}}) \cap \mathcal{O}^{f}_{\gamma_{j+1}}(f_{\gamma_{j+1}}, W_{\gamma_{j+1}}) \subset \{0\},$$
(5)

then $x \to 0$.

- 2. The same conclusion as in 1. holds if we replace Assumption 5 by Assumption 3 and condition 1. (ii) by the weaker one:
 - (a) for every $c \in \mathbb{R}$ and for each simple cycle $\{\gamma_i\}_{i=1}^m$ of H there exists $j \in \{1, \dots, m-1\}$ such that

$$\mathcal{O}_{\gamma_j}^b(f_{\gamma_j}, W_{\gamma_j}) \cap \mathcal{O}_{\gamma_{j+1}}^f(f_{\gamma_{j+1}}, W_{\gamma_{j+1}}) \cap V_{\gamma_j}^{-1}(c) \cap V_{\gamma_{j+1}}^{-1}(c) \subset \{0\}.$$
(6)

Proof. As $\sigma \in \mathcal{S}_d \cap \mathcal{S}^H$, there exists $\tau_D > 0$ such that $\sigma \in \mathcal{S}_d[\tau_D]$.

Suppose that (x, σ) verifies Assumption 5. Since $\mathcal{S}_d[\tau_D] \cap \mathcal{S}^H$ has property **P** and (x, σ) verifies the hypotheses of Theorem 3.3, there exists $\vec{c} \in \mathbb{R}^N$ such that $x \to \pi_1(M(\vec{c}))$, with $M(\vec{c})$ as in that theorem (with $\mathcal{T}^* = \mathcal{T}_d[\tau_D] \cap \mathcal{T}^H$). So, it suffices to show that $M(\vec{c}) \subset \{0\} \times \Gamma$.

Let $(\xi, \gamma) \in M(\vec{c})$; then there exists a trajectory $(x^*, \sigma^*) \in \mathcal{T}_d[\tau_D] \cap \mathcal{T}^H$ such that $(x^*(0), \sigma^*(0)) = (\xi, \gamma)$ and $(x^*(t), \sigma^*(t)) \in M(\vec{c})$ for all $t \in \mathbb{R}$. So, reasoning as in Theorem 4.1, $V_{\sigma^*(t^-)}(x^*(t)) = c_{\sigma^*(t^-)}, V_{\sigma^*(t)}(x^*(t)) = c_{\sigma^*(t)}$ and $W_{\sigma^*(t^-)}(x^*(t)) = W_{\sigma^*(t)}(x^*(t)) = 0$ for all $t \in \mathbb{R}$. We distinguish two cases:

Case 1. σ^* has a finite number of switching times, $t_0 < t_1 < \cdots < t_l$. Suppose first that for every $\gamma \in \Gamma$, $\mathcal{O}^f_{\gamma}(f_{\gamma}, W_{\gamma}, \infty) \subset \{0\}$ and let $\varphi(t) = x^*(t+t_l)$ for $t \geq 0$ and $\gamma_l = \sigma^*(t_l)$. Then φ is a solution of $\dot{z} = f_{\gamma_l}(z)$, $\varphi(t) \in \mathcal{O}_{\gamma_l}$ and $W_{\gamma_l}(\varphi(t)) = 0$ for all $t \geq 0$. In consequence $x^*(t_l) = 0$, since $x^*(t_l) = \varphi(0) \in \mathcal{O}^f_{\gamma_l}(f_{\gamma}, W_{\gamma}, \infty) \subset \{0\}$.

That $\xi = x^*(0) = 0$, follows from the fact that for every $\gamma \in \Gamma$ such that $0 \in \mathcal{O}_{\gamma}$, the unique solution of the initial value problem $\dot{z} = f_{\gamma}(z), \ z(0) = 0$ is $z(t) \equiv 0$.

In the case when for every $\gamma \in \Gamma$, $\mathcal{O}^b_{\gamma}(f_{\gamma}, W_{\gamma}, \infty) \subset \{0\}$, we proceed in a similar way, but considering instead $\varphi(t) = x^*(t+t_0)$ for $t \leq 0$.

Case 2. σ^* has an infinite number of switching times. Since $\sigma^* \in \mathcal{S}^H$, there exists a finite sequence of consecutive switching times $\{t_k\}_{k=1}^m$, such that the sequence $\{\gamma_k\}_{k=1}^m$, with $\gamma_k = \sigma^*(t_k)$, is a simple cycle of H. By hypothesis there exists an index $j \in \{1, \ldots, m-1\}$ for which (5) holds. For such j we consider the function $\varphi : [-\tau_D, \tau_D] \to \mathbb{R}^n$ defined by $\varphi(t) = x^*(t+t_{j+1})$. Since $(x^*, \sigma^*) \in \mathcal{T}_d[\tau_D]$ we have, for any $0 < \tau < \tau_D$, that

1. $\varphi : [-\tau, 0] \to \mathcal{O}_{\gamma_j}$ is solution of $\dot{z} = f_{\gamma_j}(z)$ on $[-\tau, 0]$ and $\varphi : [0, \tau] \to \mathcal{O}_{\gamma_{j+1}}$ is solution of $\dot{z} = f_{\gamma_{j+1}}(z)$ on $[0, \tau]$;

2.
$$W_{\gamma_j}(\varphi(t)) = 0$$
 on $[-\tau, 0]$ and $W_{\gamma_{j+1}}(\varphi(t)) = 0$ on $[0, \tau]$.

Therefore

$$\varphi(0) \in \mathcal{O}^b_{\gamma_j}(f_{\gamma_j}, W_{\gamma_j}) \cap \mathcal{O}^f_{\gamma_{j+1}}(f_{\gamma_{j+1}}, W_{\gamma_{j+1}})$$

and, by (5), $x^*(t_{j+1}) = \varphi(0) = 0$. By using arguments similar to those of the proof of case 1, we conclude that $\xi = x^*(0) = 0$.

Since the proof of 2. is similar to that of 1., we omit it.

Remark 4.4 It can be seen that Theorem 4.4 and Theorem 4.3 (supposing in Part 1. that Γ is finite and that Assumption 5 holds) remain valid if, instead of Assumption 7, we suppose that the function V in assumptions 3 and 5 verifies the following: for each $\gamma \in \Gamma$ such that $0 \in \chi_{\gamma}, V_{\gamma}^{-1}(0) \cap \chi_{\gamma} = \{0\}$. This condition is fulfilled when, for example, $V_{\gamma}(\cdot)$ is positive definite on χ_{γ} for every γ such that $0 \in \chi_{\gamma}$.

4.2 Stability criteria

Combining the convergence results already presented with well known sufficient Lyapunov conditions for the local (global) stability of a family \mathcal{T} of forward complete trajectories of (1), we can derive some new local (global) asymptotic stability criteria.

We recall that a family \mathcal{T} of forward complete trajectories of (1) is

1. locally uniformly stable (LUS) if there exist a positive number r > 0and a function $\alpha : [0, r] \to \mathbb{R}$ of class \mathcal{K}^1 such that for all $(x, \sigma) \in \mathcal{T}$

$$|x(t_0)| \le r \Rightarrow |x(t)| \le \alpha(|x(t_0)|) \quad \forall t \ge t_0, \forall t_0 \ge 0;$$

2. globally uniformly stable (GUS) if there exists a function $\alpha : [0, \infty) \to \mathbb{R}$ of class \mathcal{K}_{∞} such that for all $(x, \sigma) \in \mathcal{T}$

$$|x(t)| \le \alpha(|x(t_0)|) \quad \forall t \ge t_0, \forall t_0 \ge 0;$$

3. locally asyptotically stable (LAS) if it is LUS and there exists $\eta > 0$ such that for all $(x, \sigma) \in \mathcal{T}$ with $|x(0)| < \eta, x \to 0$;

¹ As usual, by a \mathcal{K} -function we mean a function $\alpha : [0, r] \to \mathbb{R}_{\geq 0}$ that is strictly increasing and continuous, and satisfies $\alpha(0) = 0$. A \mathcal{K}_{∞} -function is one of class \mathcal{K} for which $r = +\infty$ and that is in addition unbounded.

4. globally asyptotically stable (GAS) is it is GUS and for all $(x, \sigma) \in \mathcal{T}, x \to 0$.

The different stability results that we present next, require the introduction of the following pair of functions.

Definition 4.1 We say that a pair (V, W) is a weak Lyapunov pair for the family \mathcal{T} of forward complete trajectories of (1) if

1. $V \in \mathcal{V}, 0 \in \mathcal{O}$ and there exist functions α_1 and α_2 of class \mathcal{K} such that

$$\alpha_1(|\xi|) \le V(\xi, \gamma) \le \alpha_2(|\xi|) \quad \forall \xi \in \mathcal{O}_\gamma, \forall \gamma \in \Gamma.$$
(7)

- 2. $W : \operatorname{dom}(f) \cap (\mathcal{O} \times \Gamma) \to \mathbb{R}$ is such that (4) holds with $W_{\gamma}(\cdot) = W(\cdot, \gamma)$.
- 3. For every $(x, \sigma) \in \mathcal{T}$, the following is verified: $x(t) \in \mathcal{O} \quad \forall t \in [a, b] \subset [0, +\infty) \Rightarrow v(t) = V(x(t), \sigma(t))$ is nonincreasing on [a, b].

We say that a pair (V, W) is a F-weak Lyapunov pair for the family \mathcal{T} of forward complete trajectories of (1) if V and W satisfy 1. and 2. and the following condition, which is weaker than 3.

4. For every $(x, \sigma) \in \mathcal{T}$, the following holds: $x(t) \in \mathcal{O}$ for all $t \in [a, b] \subset [0, +\infty) \Rightarrow$ for every $\gamma \in \Gamma$ $v(t) = V(x(t), \gamma)$ is nonincreasing on $[a, b] \cap \sigma^{-1}(\gamma)$.

By using standard techniques (like those in [2] or in [11]) it is not hard to prove that the existence of a weak Lyapunov pair (or a F-weak Lyapunov pair when Γ is finite) for a family of trajectories \mathcal{T} of (1), implies that \mathcal{T} is LUS and that it is GUS if, in addition, $\mathcal{O} = \mathbb{R}^n$ and V is radially unbounded, i.e. there exist functions α_1 and α_2 of class \mathcal{K}_{∞} such that (7) holds.

Theorem 4.5 Suppose that Assumption 6 holds and let \mathcal{T} be a family of forward complete trajectories of (1) such that for every $(x, \sigma) \in \mathcal{T}, \sigma \in \mathcal{S}_a$. Then \mathcal{T} is LAS if one of the following conditions holds:

- 1. there exists a weak Lyapunov pair (V, W) for \mathcal{T} such that the restriction of V to dom $(f) \cap (\mathcal{O} \times \Gamma)$ is continuous and 1.(i) and 2.(ii) of Theorem 4.3 hold.
- 2. Assumption 7 holds and there exists a weak Lyapunov pair (V, W) for \mathcal{T} such that W is continuous and 1.(i) and 1.(ii) of Theorem 4.3 hold.

- 3. Γ is finite and there exists a F-weak Lyapunov pair (V, W) for \mathcal{T} such that 1.(i) and 1.(ii) of Theorem 4.3 hold.
- If, in addition, $\mathcal{O} = \mathbb{R}^n$ and V is radially unbounded, then \mathcal{T} is GAS.

Theorem 4.6 Suppose that Γ is finite and that Assumption 6 holds. Let \mathcal{T} be a family of forward complete trajectories of (1) such that for every $(x,\sigma) \in \mathcal{T}, \sigma \in \mathcal{S}_d \cap \mathcal{S}^H$, with $H : \Gamma \rightsquigarrow \Gamma$. Then \mathcal{T} is LAS if one of the following holds.

- 1. There exists a weak Lyapunov pair (V, W) such that 1.(i) and 2.(iii) of Theorem 4.4 hold.
- 2. There exists a F-weak Lyapunov pair (V,W) such that 1.(i) and 1.(ii) of Theorem 4.4 hold.
- If, in addition, $\mathcal{O} = \mathbb{R}^n$ and V is radially unbounded, then \mathcal{T} is GAS.

Theorem 4.7 Suppose that Γ is finite and let \mathcal{T} be a family of forward complete trajectories of (1) such that for every $(x, \sigma) \in \mathcal{T}, \sigma \in \mathcal{S}_e \cap \mathcal{S}_d$. Suppose that there exists a F-weak Lyapunov pair (V, W) for \mathcal{T} such that for all $\gamma \in \Gamma$, either $\mathcal{O}^b_{\gamma}(f_{\gamma}, W_{\gamma}) = E_{\gamma} \cap \mathcal{O}_{\gamma}$ or $\mathcal{O}^f_{\gamma}(f_{\gamma}, W_{\gamma}) = E_{\gamma} \cap \mathcal{O}_{\gamma}$ and that $\bigcap_{\gamma \in \Gamma} (E_{\gamma} \cap \mathcal{O}_{\gamma}) = \{0\}$. Then \mathcal{T} is LAS.

If, in addition, $\mathcal{O} = \mathbb{R}^n$ and V is radially unbounded, then \mathcal{T} is GAS.

Proof of Theorems 4.5, 4.6 and 4.7. Since the hypotheses of the three theorems imply that \mathcal{T} is locally uniformly stable (LUS), we only need to prove that there exists $\eta > 0$ such that for every $(x, \sigma) \in \mathcal{T}$, $|x(0)| < \eta$ implies that $x \to 0$.

Since \mathcal{T} is LUS, there exist $\eta > 0$ and $\rho > 0$ such that, for every $(x, \sigma) \in \mathcal{T}$ with $|x(0)| < \eta$, $x(t) \in B = \{\xi \in \mathbb{R}^n : |\xi| \le \rho\} \subset \mathcal{O}$ for all $t \ge 0$. Therefore $(x, \sigma) \in \mathcal{T}$ is precompact relative to \mathcal{O} whenever $|x(0)| < \eta$. Then, due to 1. of Remark 4.4, to Theorem 4.3 in the case of Theorem 4.5 and to Theorem 4.4 in the case of Theorem 4.6, we have that $x \to 0$ for any $(x, \sigma) \in \mathcal{T}$ such that $|x(0)| < \eta$.

In the case of Theorem 4.7, due to Theorem 4.2 we have that for every $(x, \sigma) \in \mathcal{T}$ such that $|x(0)| < \eta, x \to \bigcap_{\gamma \in \Gamma} (E_{\gamma} \cap \mathcal{O}_{\gamma}) = \{0\}$. In consequence the local asymptotic stability of \mathcal{T} follows.

When $\mathcal{O} = \mathbb{R}^n$ and V is radially unbounded, we have that \mathcal{T} is GUS. That $x \to 0$ for every $(x, \sigma) \in \mathcal{T}$ follows by using the fact that any trajectory of \mathcal{T} is precompact and the same arguments as above. **Remark 4.5** Theorem 4.7 strengthens Theorem 15 in [18] (which is the extension of the main result of [3] to nonlinear switched systems). In fact, the hypotheses of Theorem 4.7 are weaker than those of that theorem since, on one hand, even when restricted to the case $V(\xi, \gamma) = V(\xi)$ and $\mathcal{O} = \mathbb{R}^n$ (as that theorem considers) the condition $\bigcap_{\gamma \in \Gamma} E_{\gamma} = \{0\}$ is weaker than the hypothesis that V is a common joint Lyapunov function as is assumed in that work and, on the other hand, the condition $\mathcal{O}^f_{\gamma}(f_{\gamma}, W_{\gamma}) = E_{\gamma}$ is weaker than the condition $M \cap Z_{\gamma} = E_{\gamma}$ (with $Z_{\gamma} = \{\xi : W_{\gamma}(\xi) = 0\}$) considered in [18], since it always holds that $\mathcal{O}_{\gamma}(f_{\gamma}, W_{\gamma}) \subset M \cap Z_{\gamma}$ and sometimes the inclusion is strict.

From Theorem 4.7 and Remark 4.1.2. we can easily derive the following result, that contains as a particular case Theorem 1 of [3].

Corollary 4.1 Assume that Γ is finite and that $f_{\gamma}(\xi) = A_{\gamma}\xi$ with $A_{\gamma} \in \mathbb{R}^{n \times n}$ for all $\xi \in \mathbb{R}^n$. Let \mathcal{T} be a family of forward complete trajectories of (1) such that for all $(x, \sigma) \in \mathcal{T}, \sigma \in S_d \cap S_e$. Suppose that there exists a family of positive definite matrixes $\{P_{\gamma}, \gamma \in \Gamma\} \subset \mathbb{R}^{n \times n}$ and a family of matrixes $\{C_{\gamma}, \gamma \in \Gamma\}$ such that

- 1. $P_{\gamma}A_{\gamma} + A_{\gamma}^T P_{\gamma} \leq -C_{\gamma}^T C_{\gamma}$ for all $\gamma \in \Gamma$;
- 2. $v(t) = x^T(t)P_{\sigma(t)}x(t)$ is nonincreasing on $[0,\infty)$ for all $(x,\sigma) \in \mathcal{T}$;
- 3. for every $\gamma \in \Gamma$, \mathcal{U}_{γ} , the unobservable subspace of the pair (C_{γ}, A_{γ}) , coincides with ker (A_{γ}) ;
- $4. \ \cap_{\gamma \in \Gamma} \ker(A_{\gamma}) = \{0\}.$

The, \mathcal{T} is GAS.

5 Conclusions

In this paper we have obtained some invariance results for switched systems which satisfy a dwell-time condition. These results enable us to study, in an unified way, properties of bounded trajectories of switched systems whose switchings are subjected not only to state-dependent constraints, but also to restrictions on the accessibility from each subsystem to other ones.

We also derived from these results some convergence and stability criteria. These criteria involve observability-like conditions on functions which bound the derivatives of some continuous functions that are nonincreasing along complete trajectories of the switched systems.

ACKNOWLEDGMENTS

This work was partially supported by FONCyT Project 31255.

References

- A. Bacciotti and L. Mazzi. An invariance principle for nonlinear switched systems. Systems Control Lett., 54:1109–1119, 2005.
- [2] M. Branicky. Multiple Lyapunov functions and other analysis tools for switched and hybrid systems. *IEEE Trans. Automat. Control*, 43:475–482, 1998.
- [3] D. Cheng, J. Wang and X. Hu. An extension of LaSalle's invariance principle and its applications to multi-agent consensus. *IEEE Trans. Automat. Control*, 53:1765–1770, 2008.
- [4] R.A. DeCarlo, M.S. Branicky, S. Pettersson and B. Lennartson. Perspectives and results on the stability and stabilizability of hybrid systems. *Proc. IEEE*, 88:1069–1082, 2000.
- [5] R. Goebel, R.G. Sanfelice and A. Teel. Invariance principles for switching systems via hybrid systems techniques. *Systems Control Lett.*, 57:980–986, 2008.
- [6] J.P Hespanha. Uniform stability properties of switched linear systems: extensions of LaSalle's invariance principle. *IEEE Trans. Automat. Control*, 49:470–482, 2004.
- [7] J.P Hespanha, D. Liberzon, D. Angeli and E. Sontag. Nonlinear observability notions and stability of switched systems. *IEEE Trans. Automat. Control*, 50:154–168, 2005.
- [8] J.P. LaSalle. Some extensions of Lyapunov's second method. IRE Trans. Circuit Theory, 7:520–527, 1960.
- [9] J.P. LASALLE, The Stability of Dynamical Systems, CBMS-NSF Regional Conference Series in Applied Mathematics, SIAM, Philadelphia, 1976.
- [10] T.C. Lee and Z.P. Jiang. Uniform asymptotic stability of nonlinear switched systems with an application to mobile robots. *IEEE Trans. Au*tomat. Control, 53:1235–1252, 2008.
- [11] D. LIBERZON, Switching in Systems and Control, Birkhäuser, Boston, 2003.

- [12] H. Lin and P.J. Antsaklis. Stability and stabilizability of switched linear systems: A survey of recent results. *IEEE Trans. Automat. Control*, 54: 308–322, 2009.
- [13] H. Logemann and E.P. Ryan. Asymptotic behaviour of nonlinear systems. Amer. Math. Monthly, 111:864–889, 2004.
- [14] J. Lygeros, K. H. Johansson, S. M. Simić, J. Zhang and S. Sastry. Dynamical properties of hybrid automata. *IEEE Trans. Automat. Control*, 48:2–17, 2003.
- [15] J.L. Mancilla Aguilar and R.A.García. A converse Lyapunov theorem for nonlinear switched systems. Systems Control Lett., 41:67–71, 2000.
- [16] J.L. Mancilla Aguilar and R.A.García. An extension of LaSalle's invariance principle for switched systems. *Systems Control Lett.*, 55:376–384, 2006.
- [17] R. Sanfelice, R. Goebel and A. Teel. Invariance principles for hybrid systems with connections to detectability and asymptotic stability. *IEEE Trans. Automat. Control*, 52:2282–2297, 2007.
- [18] J. Wang and D. Cheng. Stability of switched nonlinear systems via extensions of LaSalle's invariance principle. *Sci. China, Ser. F: Information Sciences*, 52:84–90, 2009.
- [19] J. Wang, D. Cheng and X. Hu. An extension of LaSalle's invariance principle for a class of switched linear systems. *Systems Control Lett.*, 58: 754–758, 2009.
- [20] R WHEEDEN AND A ZYGMUND, Measure and Integral: an Introduction to Real Analysis, Pure and Applied Mathematics, Vol. 43. Marcel Dekker, Inc., New York-Basel, 1977.

A Proof of Lemma 3.2

The following lemma is used in the proofs of Lemma 3.2 and Theorem 3.1.

Lemma A.1 Let $\{\sigma_k\}$ be a sequence of switching signals in $S_a[\tau_D, N_0]$ with $\tau_D > 0$ and $N_0 \in \mathbb{N}$. Then there exist a subsequence $\{\sigma_{k_l}\}$ and a switching signal $\sigma^* \in S_a[\tau_D, N_0]$ such that

1. $\lim_{l\to\infty} \sigma_{k_l}(t) = \sigma^*(t)$ for almost all $t \in \mathbb{R}$;

2. for each $t \in \mathbb{R}$ there exists a sequence of times $\{r_l(t)\}$ such that

$$\lim_{l\to\infty}r_l(t)=t,\ \lim_{l\to\infty}\sigma_{k_l}(r_l(t))=\sigma^*(t)\ and\ \lim_{l\to\infty}\tau^1_{\sigma_{k_l}}(r_l(t))-r_l>0.$$

Proof. Let $m \in \mathbb{N}$. By applying Lemma A.1 in [16] to the sequence $\{\sigma_k(\cdot - m)\}$, the thesis holds with $[-m, +\infty)$ instead of \mathbb{R} . In addition, for any $t \geq -m$ the sequence $\{r_l(t)\}$ can be chosen to satisfy the condition $r_l(t) > -m$ for all l.

Due to this fact, for each $m \in \mathbb{N}$ we can construct recursively a sequence of positive integers $\{k_i^m\}_{j \in \mathbb{N}}$ such that:

- a) $\{k_j^{m+1}\}_{j\in\mathbb{N}}$ is a subsequence of $\{k_j^m\}_{j\in\mathbb{N}}$ for all $m\in\mathbb{N}$;
- b) For each $m \in \mathbb{N}$ there exists $\sigma_m^* \in \mathcal{S}_a[\tau_D, N_0]$ such that 1. $\lim_{j \to \infty} \sigma_{k_j^m}(t) = \sigma_m^*(t)$ for almost all $t \ge -m$; 2. for each $t \in [-m, \infty)$ there exists a sequence of times $\{r_j^m(t)\}_{j \in \mathbb{N}}$, with $r_j^m(t) > -m$ for all $j \in \mathbb{N}$, such that $\lim_{j \to \infty} r_j^m(t) = t$, $\lim_{j \to \infty} \sigma_k^m(t) = \sigma_k^*(t)$ and $\lim_{j \to \infty} \sigma_k^m(t) = r_k^m(t)$.

$$\lim_{j \to \infty} r_j^m(t) = t, \quad \lim_{j \to \infty} \sigma_{k_j^m}(r_j^m(t)) = \sigma_m^*(t) \text{ and } \lim_{l \to \infty} \tau_{\sigma_{k_j^m}}^1(r_j^m(t)) - r_j^m(t) > 0$$

In addition, if $m \ge 2$ and $t \ge -m - 1$, $\{r_j^m(t)\}_{j \in \mathbb{N}}$ is a subsequence of $\{r_j^{m-1}(t)\}_{j \in \mathbb{N}}$.

Let us define $\{\sigma_{k_l}\}$ by $\sigma_{k_l} = \sigma_{k_l^l}$ for all $l \in \mathbb{N}$, σ^* by $\sigma^*(t) = \sigma_l^*(t)$ if $t \in [-l, -l+1)$ with $l \ge 2$, and by $\sigma^*(t) = \sigma_1(t)$ if $t \in [-1, \infty)$ and $\{r_l(t)\}$ by $r_l(t) = r_l^l(t)$ for all $t \in \mathbb{R}$ and all $l \in \mathbb{N}$. Taking into account that for all $t \ge -m$, $\{k_l^l\}_{l \ge m}$ and $\{r_l^l(t)\}_{l \ge m}$ are subsequences of $\{k_l^m\}_{l \in \mathbb{N}}$ and of $\{r_l^m(t)\}_{l \in \mathbb{N}}$ respectively, the theorem follows.

Proof of Lemma 3.2. Since it is clear that the three families of switching signals verify 1. and 2. of definition 3.3, we only have to prove that they also verify property 3. That this fact is true for $S_a[\tau_D, N_0]$ follows from Lemma A.1, and next we proceed to show that the other families of switching signals also have this property.

Suppose that $\sigma_k \in \mathcal{S}_d[\tau_D] \cap \mathcal{S}^H$ for all k. Then, from Lemma A.1 and the fact that $\mathcal{S}_d[\tau_D] = \mathcal{S}_a[\tau_D, 1]$, there exist a subsequence $\{\sigma_{k_l}\}$ and $\sigma^* \in \mathcal{S}_d[\tau_D]$ such that $\sigma_{k_l} \to \sigma^*$ a.e. on \mathbb{R} . Let $t \in \mathbb{R}$ be a switching time of σ^* ; since $\sigma_{k_l}(\tau) \to \sigma^*(\tau)$ for almost all τ , there exist s < t < s' such that $\sigma_{k_l}(s) \to \sigma^*(s)$, $\sigma_{k_l}(s') \to \sigma^*(s')$ and $s' - s < \tau_D$. We note that $\sigma^*(s') = \sigma^*(t) \neq \sigma^*(t^-) = \sigma^*(s)$.

Then, for l large enough, $\sigma_{k_l}(s) \neq \sigma_{k_l}(s')$, and in consequence, σ_{k_l} has a unique switching time in the interval (s, s']. Thus, $(\sigma_{k_l}(s), \sigma_{k_l}(s')) \in$ Graph(H) for l large enough and, in consequence, since Graph(H) is closed,

$$(\sigma^*(t^-), \sigma^*(t)) = (\sigma^*(s), \sigma^*(s')) \in \operatorname{Graph}(H).$$

Therefore $\sigma^* \in \mathcal{S}_d[\tau_D] \cap \mathcal{S}^H$.

Suppose now that $\{\sigma_k\} \subset S_d[\tau_D] \cap S_e[T]$. By the same arguments of the previous case, there exist a subsequence $\{\sigma_{k_l}\}$ and $\sigma^* \in S_d[\tau_D]$ such that $\sigma_{k_l} \to \sigma^*$ a.e. on \mathbb{R} .

Let $t_0 \ge 0, \ \gamma \in \Gamma$ and $\epsilon > 0$ be fixed, and suppose that

$$\sigma^{*-1}(\gamma) \cap [t_0, t_0 + T + \epsilon] = \emptyset.$$
(8)

Let $\epsilon' > 0$ such that $\epsilon' < \min\{\epsilon/2, \tau_D/2\}$ and

$$\lim_{k \to \infty} \sigma_k(t_0 + \epsilon') = \sigma^*(t_0 + \epsilon') = \gamma_0 \neq \gamma,$$
$$\lim_{k \to \infty} \sigma_k(t_0 + T + \epsilon - \epsilon') = \sigma^*(t_0 + T + \epsilon - \epsilon') = \gamma_1 \neq \gamma.$$

Then, since Γ is a finite set, there exists $K_0 \in \mathbb{N}$ such that

$$\sigma_k(t_0 + \epsilon') = \gamma_0 \text{ and } \sigma_k(t_0 + T + \epsilon - \epsilon') = \gamma_1 \quad \forall k \ge K_0.$$
(9)

Let $I = [t_0 + \epsilon', t_0 + T + \epsilon - \epsilon']$; as the length of I is greater than T and (9) holds, then for each $k \geq K_0$ there exists $t_k \in I$ such that $\sigma_k(t_k) = \gamma$ and $\sigma_k(t_k^-) \neq \gamma$. From the compactness of I, there exists a subsequence $\{t_{k_l}\} \subset \{t_k\}$ that converges to, say, $t_{\gamma} \in I$.

Let $\epsilon'' \in (0, \epsilon'/2)$ and $L_0 \in \mathbb{N}$ such that $t_{k_l} \in (t_\gamma - \epsilon'', t_\gamma + \epsilon'')$ for every $l \geq L_0$. Since for all $l \geq L_0$, $\sigma_{k_l}(s) = \gamma \forall s \in [t_{k_l}, t_{k_l} + \tau_D)$, then for those l's $\sigma_{k_l}(s) = \gamma \forall s \in [t_\gamma + \epsilon'', t_\gamma + \epsilon') \subset [t_0, t_0 + T + \epsilon]$. Hence, there exists $t \in [t_\gamma + \epsilon'', t_\gamma + \epsilon')$ such that $\gamma = \lim_{l \to \infty} \sigma_{k_l}(t) = \lim_{k \to \infty} \sigma_k(t) = \sigma^*(t)$, which contradicts (8). In consequence for every $\epsilon > 0$, every $t_0 \geq 0$ and every $\gamma \in \Gamma$, $\sigma^{*-1}(\gamma) \cap [t_0, t_0 + T + \epsilon] \neq \emptyset$.

Suppose now that for certain $t_0 \geq 0$ and $\gamma \in \Gamma$, $\sigma^{*-1}(\gamma) \cap [t_0, t_0 + T] = \emptyset$; then $\sigma^*(t_0 + T) \neq \gamma$ and since σ^* in right-continuous, there exists $\epsilon > 0$ such that $\sigma^*(t_0 + T + s) \neq \gamma \forall s \in [0, \epsilon]$ and then $\sigma^{*-1}(\gamma) \cap [t_0, t_0 + T + \epsilon] = \emptyset$ which is a contradiction. In consequence $\sigma^{*-1}(\gamma) \cap [t_0, t_0 + T] \neq \emptyset$ for every $t_0 \geq 0$ and every $\gamma \in \Gamma$, and the lemma follows.

B Proof of Theorem 3.1

The next result, that shows that under suitable hypotheses certain families of trajectories of system (1) enjoy a certain kind of sequential compactness, is used in the proof of Theorem 3.1.

We say that a sequence $\{(x_k, \sigma_k)\}$ of trajectories of (1) is uniformly precompact if there exists a compact set $B \subset \mathbb{R}^n$ such that $x_k(t) \in B$ for all $t \in I_{x_k}$.

Consider next the family \mathcal{I} composed by the intervals I = [a, b) or $I = (-\infty, b)$ with b finite or $b = +\infty$.

Proposition B.1 Let $\{(x_k, \sigma_k)\}$ be a uniformly precompact sequence of trajectories of (1) such that for a certain interval $I \in \mathcal{I}$ and certain $\tau_D > 0$ and $N_0 \in \mathbb{N}$, $\sigma_k \in \mathcal{S}_a[\tau_D, N_0]$ and $I_{x_k} = I$ for all k. Then, there exist a subsequence $\{(x_{k_l}, \sigma_{k_l})\}$ and a trajectory (x^*, σ^*) of (1) with $I_{x^*} = I$ and $\sigma^* \in \mathcal{S}_a[\tau_D, N_0]$ such that

- 1. $x_{k_l} \rightarrow x^*$ uniformly on compact subsets of I;
- 2. $\sigma_{k_l} \to \sigma^*$ a.e. on \mathbb{R} .

Proposition B.1 makes use of the assertions of the following lemma.

Lemma B.1 Let $\{(x_k, \sigma_k)\}$ be a sequence such that for all $k \in \mathbb{N}, x_k : [a, b] \to \mathbb{R}^n$ is absolutely continuous, $\sigma_k : [a, b] \to \Gamma$ and $\dot{x}(t) = f(x_k(t), \sigma_k(t))$ a.e. on [a, b]. Suppose that there exist a compact set $B \subset \mathbb{R}^n$ such that $x_k([a, b]) \subset B$ for all k, and a function $\sigma : [a, b] \to \Gamma$ such that $\lim_{k\to\infty} \sigma_k(t) = \sigma(t)$ for almost all $t \in [a, b]$. Then there exist a subsequence $\{x_{k_j}\}$ of $\{x_k\}$ and an absolutely continuous function $x : [a, b] \to \mathbb{R}^n$ such that $x_{k_j} \to x$ uniformly on [a, b] and $\dot{x}(t) = f(x(t), \sigma(t))$ a.e. on [a, b].

Proof. Since for all $k \in \mathbb{N}$, $(x_k(t), \sigma_k(t)) \in \text{dom}(f) \cap (B \times \Gamma), \dot{x}_k(t) = f(x_k(t), \sigma_k(t))$ a.e. [a, b] and f is continuous on the compact set $\text{dom}(f) \cap (B \times \Gamma)$, there exists M > 0 such that for all $k \in \mathbb{N}, |\dot{x}_k(t)| \leq M$ for almost all $t \in [a, b]$. In consequence $\{x_k\}$ is uniformly Lipschitz, since $|x_k(t) - x_k(s)| \leq M |t - s|$ for all $t, s \in [a, b]$ and all k, and therefore equicontinuous.

Then, due to Arzelà-Ascoli Theorem, there exist a subsequence $\{x_{k_j}\}$ and a continuous function $x : [a, b] \to \mathbb{R}^n$ such that $x_{k_j} \to x$ uniformly on [a, b].

We note that $(x(t), \sigma(t)) \in \text{dom}(f)$ a.e. on [a, b] since for every $t \in [a, b]$ such that $\lim_{k\to\infty} \sigma_k(t) = \sigma(t)$ and that $(x_k(t), \sigma_k(t)) \in \text{dom}(f)$ for all $k \in \mathbb{N}$, $(x(t), \sigma(t)) \in \text{dom}(f)$.

Given that $\lim_{j\to\infty} \dot{x}_{k_j}(t) = \lim_{j\to\infty} f(x_{k_j}(t), \sigma_{k_j}(t)) = f(x(t), \sigma(t))$ for almost all $t \in [a, b]$ and that $\{\dot{x}_{k_j}\}$ is majorized by the constant function $\rho(t) = M$ on [a, b], by applying Lebesgue's Dominated Convergence Theorem we have that

$$x(t) = \lim_{j \to \infty} \left(x_{k_j}(a) + \int_a^t \dot{x}_{k_j}(s) ds \right)$$
$$= x(a) + \int_a^t f(x(s), \sigma(s)) ds,$$

and the lemma follows.

Proof of Proposition B.1 Due to Lemma A.1 we can suppose without loss of generality that there exists a switching signal $\sigma^* \in \mathcal{S}_a[\tau_D, N_0]$ such that $\lim_{k\to\infty} \sigma_k(t) = \sigma^*(t)$ for almost all $t \in \mathbb{R}$. Consider a sequence of intervals $\{[a_k, b_k]\}$ such that for all $k \in \mathbb{N}, [a_k, b_k] \subset [a_{k+1}, b_{k+1}] \subset I$ and $\bigcup_{k \in \mathbb{N}} [a_k, b_k] = I$ and let a compact set B such that $x_k(I) \subset B$ for all k. Since $x_k([a_1, b_1]) \subset B$, it follows from Lemma B.1 that there exist a subsequence $\{(x_k^1, \sigma_k^1)\}$ and an absolutely continuous function $x^1 : [a_1, b_1] \to \mathbb{R}^n$ such that $x_k^1 \to x^1$ uniformly on $[a_1, b_1]$ and $\dot{x}^1(t) = f(x^1(t), \sigma^*(t))$ a.e. $[a_1, b_1]$. Applying Lemma B.1 to the sequence $\{(x_k^1, \sigma_k^1)\}$ on the interval $[a_2, b_2]$ we can assure the existence of a subsequence $\{(x_k^2, \sigma_k^2)\}$ of $\{(x_k^1, \sigma_k^1)\}$ and an absolutely continuous function $x^2 : [a_2, b_2] \to \mathbb{R}^n$ such that $x_k^2 \to x^2$ uniformly on $[a_2, b_2]$ and $\dot{x}^2(t) = f(x^2(t), \sigma^*(t))$ a.e. $[a_2, b_2]$. Proceeding in this way, it follows that for each $l \in \mathbb{N}$ we can construct recursively a subsequence $\{(x_k^l, \sigma_k^l)\}$ of $\{(x_k, \sigma_k)\}$ such that:

- 1. $\{x_k^{l+1}\}$ is a subsequence of $\{x_k^l\}$ for all $l \in \mathbb{N}$;
- 2. for each $l \in \mathbb{N}$ there exists an absolutely continuous function x^{l} : $[a_{l}, b_{l}] \to \mathbb{R}^{n}$ such that $x_{k}^{l} \to x^{l}$ uniformly on $[a_{l}, b_{l}]$ and $\dot{x}^{l}(t) = f(x^{l}(t), \sigma^{*}(t))$ a.e. $[a_{l}, b_{l}]$.

Let now the subsequence $\{(x_l^*, \sigma_l^*)\}$ of $\{(x_k, \sigma_k)\}$ defined for each $l \in \mathbb{N}$ as $x_l^*(t) = x_l^l(t)$ for all $t \in I$ and $\sigma_l^*(t) = \sigma_l^l(t)$ for all $t \in \mathbb{R}$ and let $x^* : I \to \mathbb{R}^n$ be the function given by $x^*(t) = x^l(t)$ for all $t \in [a_l, b_l]$ and all $l \in \mathbb{N}$. Clearly, x^* is locally absolutely continuous on I and we show next that $x_l^* \to x^*$ uniformly on compact subsets of I.

Let $K \subset I$ be compact; then there exists $j \in \mathbb{N}$ such that $K \subset [a_j, b_j]$. Since for all $l \geq j$, $x_l^* \in \{x_k^j\}$, it follows that $x_l^* \to x^j$ uniformly on $[a_j, b_j]$. The uniform convergence of $\{x_l^*\}$ to x^* follows from the fact that $x^*(t) = x^j(t)$ for all $t \in [a_j, b_j]$.

Let now for all $l \in \mathbb{N}$, $k_l \in \mathbb{N}$ such that $(x_{k_l}, \sigma_{k_l}) = (x_l^*, \sigma_l^*)$. Then the subsequence $\{(x_{k_l}, \sigma_{k_l})\}$ verifies 1. and 2. of the thesis.

The proof finishes by showing that (x^*, σ^*) is a trajectory of (1). By construction we have that $\dot{x}^*(t) = f(x^*(t), \sigma^*(t))$ for almost all $t \in I$. It remains to prove that $(x^*(t), \sigma^*(t))$ belongs to dom(f) for all $t \in I$.

Let $t \in I$; then there exists a sequence of times $\{t_k\}$, such that: i) for all k, $t_k \in I$, $t_k > t$, $(x^*(t_k), \sigma^*(t_k)) \in \text{dom}(f)$ and $\sigma^*(t_k) = \sigma^*(t)$, ii) $\lim_{k \to \infty} t_k = t$. Hence, $(x^*(t_k), \sigma^*(t_k)) \to (x^*(t), \sigma^*(t))$ and $(x^*(t), \sigma^*(t)) \in \text{dom}(f)$ since dom(f) is closed.

From Proposition B.1 and by arguments similar to those used in its proof, we can derive the following

Proposition B.2 Let $\tau_D > 0$ and $N_0 \in \mathbb{N}$ and let $\{(x_k, \sigma_k)\}$ be an uniformly precompact sequence of trajectories of (1) such that $\sigma_k \in \mathcal{S}_a[\tau_D, N_0]$ and $[-k, \infty) \subset I_{x_k}$ for all k. Then, there exist a subsequence $\{(x_{k_l}, \sigma_{k_l})\}$ and a trajectory $(x^*, \sigma^*) \in \mathcal{T}_a[\tau_D, N_0]$ such that

- 1. $x_{k_l} \to x^*$ uniformly on compact subsets of \mathbb{R} ;
- 2. $\sigma_{k_l} \to \sigma^*$ a.e. on \mathbb{R} .

Proof of Theorem 3.1. Without loss of generality we can suppose that $I_x = [0, +\infty)$. Let $(\xi, \gamma) \in \Omega^{\sharp}(x, \sigma)$; then there exists a strictly increasing and unbounded sequence $\{s_k\}$ which verifies 1. and 2. of Definition 3.1. Let $\sigma_k(\cdot) = \sigma(\cdot + s_k)$ and $x_k(\cdot) = x(\cdot + s_k)$ and note that $\sigma_k \in S^*$ for all k since S^* has property **P**. As the sequence $\{(x_k, \sigma_k)\}$ is uniformly precompact, $I_{x_k} = [-k, \infty)$ and, due to 1. of Definition 3.3, $S^* \subset S_a[\tau_D, N_0]$ for some $\tau_D > 0$ and some $N_0 \in \mathbb{N}$, then from Proposition B.2 and taking into account 3. of Definition 3.3, there exist a subsequence (x_{k_l}, σ_{k_l}) and a trajectory $(x^*, \sigma^*) \in \mathcal{T}^*$ such that $\{x_{k_l}\}$ converges to x^* uniformly on compact subsets of \mathbb{R} and $\{\sigma_{k_l}\}$ converges to σ^* a.e. on \mathbb{R} .

Due to Lemma A.1, we can assume without loss of generality that $\{\sigma_{k_l}\}$ verifies condition 2 of that lemma.

The proof ends provided that we show that $(x^*(0), \sigma^*(0)) = (\xi, \gamma)$ and that $(x^*(t), \sigma^*(t)) \in \Omega^{\sharp}(x, \sigma)$ for all $t \in \mathbb{R}$.

Let us prove first that $(x^*(0), \sigma^*(0)) = (\xi, \gamma)$. From the fact that $x_k(0) = x(s_k)$, from 2. of Definition 3.1 and due to the convergence of $\{x_{k_l}\}$ to x^* , it holds that $x^*(0) = \xi$.

According to 2. of Lemma A.1, there exists a sequence $\{r_l\} \subset \mathbb{R}$ such that $\lim_{l\to\infty} r_l = 0$, $\lim_{l\to\infty} \tau^1_{\sigma_{k_l}}(r_l) - r_l > 0$ and $\lim_{l\to\infty} \sigma_{k_l}(r_l) = \sigma^*(0)$. From item 1. of Definition 3.1 and the fact that $\tau^1_{\sigma_k}(0) = \tau^1_{\sigma}(s_k) - s_k$ for all $k \in \mathbb{N}$, it follows that $\lim_{l\to\infty} \tau^1_{\sigma_{k_l}}(0) > 0$. Then $r_l < \tau^1_{\sigma_{k_l}}(0)$ for l large enough and therefore, $\sigma^*(0) = \lim_{l\to\infty} \sigma_{k_l}(r_l) = \lim_{l\to\infty} \sigma_{k_l}(0) = \lim_{l\to\infty} \sigma(s_{k_l}) = \gamma$.

In order to prove that $(x^*(t), \sigma^*(t)) \in \Omega^{\sharp}(x, \sigma)$ for all $t \in \mathbb{R}, t \neq 0$, let one such time t and let $\{r_l(t)\}$ be a sequence as in 2. of Lemma A.1. In order to simplify the notation, we will write r_l instead of $r_l(t)$.

Consider the unbounded sequence $\{s'_l\}$, defined by $s'_l = r_l + s_{k_l}$, which we assume, without loss of generality, strictly increasing. Since $\tau_{\sigma}^1(s'_l) = \tau_{\sigma_{k_l}}^1(r_l) + s_{k_l}$, we have that $\lim_{l\to\infty} \tau_{\sigma}^1(s'_l) - s'_l > 0$. So $\{s'_l\}$ satisfies condition 1. of Definition 3.1.

That $\lim_{l\to\infty} \sigma(s'_l) = \lim_{l\to\infty} \sigma_{k_l}(r_l) = \sigma^*(t)$ follows from 2. of Lemma A.1. From the uniform convergence of $\{x_{k_l}\}$ to x^* on compact sets and the continuity of x^* we have that $\lim_{l\to\infty} x(s'_l) = \lim_{l\to\infty} x_{k_l}(r_l) = x^*(t)$. Hence $(x(s'_l), \sigma(s'_l)) \to (x^*(t), \sigma^*(t))$ as $l \to \infty$ and in consequence $(x^*(t), \sigma^*(t)) \in \Omega^{\sharp}(x, \sigma)$.