Uniqueness of a constrained variational problem and large deviations of buffer size

Adam Shwartz, and Alan Weiss *[†]

November 20, 2018

Abstract

We show global uniqueness of the solution to a class of constrained variational problems, using scaling properties. This is used to establish the essential uniqueness of solutions of a large deviations problem in multiple dimensions. The result is motivated by models of buffers, and in particular the probability of, and typical path to overflow in the limit of small buffers, which we analize.

keywords Uniqueness of Variational problems, Large Deviations, Buffer overflow, AMS model.

1 Introduction

We investigate uniqueness of solutions to variational problems that arise in sample-path large deviations. Our motivation comes from models of buffers in telecommunication systems. The original model was developed by Anick, Mitra, and Sondhi [1]. Weiss [11] cast this model in the framework of samplepath large deviations and showed for the probability of buffer overflow

$$\lim_{n \to \infty} \frac{1}{n} \log \mathbb{P}(b_n(t) \ge B) = \inf_{(\vec{r}, T) \in G(B)} \int_0^T \ell(\vec{r}(t), \vec{r}'(t)) \, dt, \tag{1.1}$$

^{*}Adam Shwartz is the Julius M. and Bernice Naiman Chair in Engineering, at the Faculty of Electrical Engineering, Technion—Israel Institute of Technology, Haifa 32000, Israel. email adam@ee.technion.ac.il. His work was supported in part by the fund for promotion of research at the Technion and by the Dent charitable trust-non military research fund.

[†]Alan Weiss is with Mathworks, Inc., Natick, MA 01760-2098. email aweiss@mathworks.com

where $\ell(\vec{r}(t), \vec{r}'(t))$ is a positive "cost" function, and G(B) is the set of paths \vec{r} (d-dimensional functions of t) and terminal times T satisfying a buffer overflow property (see § 7). Botvich and Duffield [2] (and independently Courcoubetis and Weber [3] and Simonian and Guibert [7]) analyze the probability of buffer overflow for more general traffic models. For a superposition of n traffic sources, each bringing independent traffic A(t) to the buffer in time (0, t), and with the buffer draining at rate nc for some constant c, they show that the steady-state probability that the scaled buffer content $b_n(t) = b(t)/n$ exceeds a level B is approximately

$$\lim_{n \to \infty} \frac{1}{n} \log \mathbb{P}\left(b_n(t) \ge B\right) = -\inf_{t \ge 0} \sup_{\theta} \Big\{ \theta(b+ct) - \log E\left(\exp(\theta A(t))\right) \Big\}.$$
(1.2)

For more recent results see e.g. Mandjes and Mannersalo [6] and references therein. Mandjes and Ridder [5] showed how to use the solution of (1.2) to find a minimal sample path $\vec{r}(t)$ in (1.1); that is, a solution to the variational problem. If this solution is unique, then this result is quite useful, as shown by Freidlin and Wentzell [4]: if $\vec{z}_n(t)$ represents the scaled state of the system at time t, $b_n(t)$ the scaled buffer size, and $\vec{r}(t)$ is the unique solution to the variational problem (1.1), then for each fixed $s \leq T$,

$$\lim_{n \to \infty} \mathbb{P}_{ss} \left(\sup_{s \le t \le T} \left| \vec{z}_n(t) - \vec{r}(t) \right| < \varepsilon \ \middle| \ b_n(T) \ge B \right) = 1.$$
 (1.3)

Here \mathbb{P}_{ss} is steady-state probability and $\mathbb{P}(A|B)$ is the probability of A conditioned on B. Thus, conditioned on the buffer overflowing, we can determine beforehand most likely how it did so. So the sample-path approach yields interesting information when the minimizing $\vec{r}(t)$ is unique.

The general problem of uniqueness for variational problems is difficult. In general, it is difficult to prove local uniqueness for variational problems, and global uniqueness is even more difficult. In addition, our problem is an optimization problem under a constraint (buffer exceeds B). Constraints can often be handled using Lagrange multipliers, so that standard uniqueness methods can be applied. However, this requires a proof that the multiplier principle holds in this case. Note that due to our application, the minimum must be shown to be unique in the class of absolutely continuous paths, not simply C^1 or piecewise C^1 paths. Finally, in terms of the application, there seem to be no direct probabilistic methods for establishing uniqueness.

In this note we consider a constrained variational problem where the constraint is given by a single functional. We assume a certain scaling property of the constraining functional together with homogeneity, convexity and superlinear growth of the Lagrangian ℓ to show global uniqueness of the minimizing solution. For our motivating model this means the following. In previous studies of this model ([8], Chapter 13) we showed, through laborious calculations, the applicability of the multiplier principle as well as uniqueness, for a simple one-dimensional case. We show here that the assumptions of our new uniqueness result hold for a constant coefficient process. Then we approximate our model with a constant coefficient one and show that, in the limit as the buffer size $B \to 0$, the variational problem has an *essentially* unique solution, in the following sense. There is a unique path $\vec{r}^*(t)$, $0 \le t \le T$, for the constant coefficient process, such that any minimizing path $\vec{r}_B(t)$ of the real process is close to a scaled version of $\vec{r}^*(t)$:

$$\lim_{B \to 0} \frac{1}{\sqrt{B}} \left(\vec{r}_B(t\sqrt{B}) - \vec{r}_B(0) \right) + \vec{r}^*(0) = \vec{r}^*(t), \ 0 \le t < T,$$
(1.4)

uniformly in t, and $\vec{r}_B(0) \to \vec{r}^*(0)$. In other words, for each $\varepsilon > 0$ there is a $\delta > 0$ such that for all $B < \delta$,

$$\sup_{0 \le t \le T-\delta} \left| \frac{1}{\sqrt{B}} \left(\vec{r}_B \left(t \sqrt{B} \right) - \vec{r}_B(0) \right) - \left(\vec{r}^*(t) - \vec{r}^*(0) \right) \right| < \varepsilon.$$
(1.5)

The conditions under which this result holds are mild.

Note that this result together with a local uniqueness result for the original model—for small enough B— imply that there is a unique solution for all B small enough, for each given starting point. We do not give general conditions for the uniqueness of the full buffer model, but some discussions appear in § 8. Our result does not imply full sample-path uniqueness for the stochastic model with non-constant rates, although we are certain that uniqueness does hold. In paticular, we have no effective bounds on the error (the relationship between ε and δ in (1.5)).

In § 2 we introduce the general variational problem, and in § 3 we state the abstract assumptions under which uniqueness holds. Sections 4 and 5 prove existence and uniqueness resp. In § 6 we derive a property of the shape of optimal paths for buffer problems. In § 7 we formulate the stochastic buffer problem and obtain some properties, in § 8 we prove weak uniqueness and in § 9 we show a strong uniqueness result and discuss the desired full uniqueness result.

2 The variational problem

We assume throughout that the functional $I_{[0,T]}(\vec{r})$ on absolutely continuous paths $\vec{r}(t)$ (*d*-dimensional functions of a real variable) has a representation

$$I_{[0,T]}(\vec{r}) \stackrel{\triangle}{=} \int_0^T \ell\left(\vec{r}(t), \frac{d}{dt}\vec{r}(t)\right) dt, \qquad (2.1)$$

where $\ell(\vec{x}, \vec{y})$ is a positive function, convex in \vec{y} . The constant coefficient problem has $\ell(\vec{x}, \vec{y}) = \ell(\vec{y})$ —independent of \vec{x} . We prove our uniqueness result for a general class of constraints. So, let \mathcal{B} be a given positive functional on absolutely continuous paths. Our optimization problem is

Problem A. Given \vec{x} and B > 0, find a time T and an absolutely continuous path $\vec{r}(t)$ that minimize $I_{[0,T]}(\vec{r})$ subject to $\mathcal{B}(\vec{r},T) \geq B$.

Denote this minimal cost by \mathcal{I} . This is a "free time" problem; hence straightofrward convexity arguments cannot be used to infer uniqueness.

3 Assumptions

Throughout, we use the sup norm on the space of (measurable) functions. The following assumptions apply to the constant coefficient problem. We shall use all save the last assumption in order to prove existence of solutions to Problem A, and a different subset to prove uniqueness of solutions.

Assumption 1

- $[A] \ \ell(\vec{y})$ is positive and strictly convex in \vec{y} .
- $[B] \lim_{|\vec{y}| \to \infty} \ell(\vec{y}) / |\vec{y}| = \infty.$
- $|C| \mathcal{B}(\vec{r},t)$ is positive, continuous in (\vec{r},t) , and $\mathcal{B}(\vec{r},0) = 0$.
- [D] For some \vec{s} , $\ell(\vec{s}) < \infty$ and $\lim_{t\to\infty} \mathcal{B}(\vec{r},t) = \infty$ for $\vec{r}(t) = \vec{x} + t\vec{s}$.
- [E] Given \vec{x} and B there is a $T_1 < \infty$ such that any path $\vec{r}(t)$ on $0 \le t \le T$ with $T > T_1$, $\vec{r}(0) = \vec{x}$ and $\mathcal{B}(\vec{r}, T) \ge B$ has $I_{[0,T]}(\vec{r}) > \mathcal{I} + 1$.
- [F] For all $\vec{r}(t)$ and $\vec{z}(t)$ on $0 \le t \le T$ with $\vec{r}(0) = \vec{z}(0) = \vec{x}$, \mathcal{B} satisfies
 - 1. For any $\gamma \in (0,1)$ and $0 \le t \le T$ we have

$$\mathcal{B}(\gamma \vec{r} + (1 - \gamma)\vec{z}, t) \ge \gamma \mathcal{B}(\vec{r}, t) + (1 - \gamma)\mathcal{B}(\vec{z}, t).$$
(3.1)

2. There exists a positive increasing function $f(\alpha)$, defined for $\alpha > 0$, that satisfies $f(\alpha)f(1/\alpha) \equiv 1$, and is either the identity ($f(\alpha) = \alpha$), or is strictly convex for $\alpha \geq 1$ such that for every $\vec{r}(t)$ on $0 \leq t \leq T$, the scaled path $\vec{y}(t) \stackrel{\triangle}{=} \alpha \vec{r}(t/\alpha)$ satisfies

$$\mathcal{B}(\vec{y}, t) = f(\alpha)\mathcal{B}(\vec{r}, \alpha t) \quad 0 \le t \le \alpha T \tag{3.2}$$

Note that Assumption 1 [A] implies for any $\gamma \in (0, 1)$ we have

$$I_{[0,T]}(\gamma \vec{r} + (1-\gamma)\vec{z}) \le \gamma I_{[0,T]}(\vec{r}) + (1-\gamma)I_{[0,T]}(\vec{z}),$$
(3.3)

where the inequality is strict unless $\vec{r}(t) \equiv \vec{z}(t)$. Furthermore, the representation of I as an integral implies that, for \vec{y} defined in terms of \vec{r} as in Assumption 1 [F] 2, $I_{[0,\alpha T]}(\vec{y}) = \alpha I_{[0,T]}(\vec{r})$.

The prototypical $f(\alpha)$ satisfying [F] 2 is $f(\alpha) = \alpha^{\beta}$ for $\beta > 1$. Our motivating example has $\beta = 2$; see (6.1). The conditions do not imply that $f(\alpha)$ is convex for $\alpha < 1$. They imply that f(1) = 1 and $f(\alpha) \to 0$ as $\alpha \to 0$.

4 Existence

The real import of this paper is uniqueness; for completeness, we prove that for the constant coefficient problem, there exist solutions to Problem A under our assumptions.

Theorem 1 Under Assumption 1 parts [A]-[E], there exists a solution to Problem A.

Proof: By Assumption 1 [D] the linear function $\vec{r}(t) = \vec{x} + t\vec{s}$ has $I_{[0,T]}(\vec{r}) < \infty$ and makes $\mathcal{B}(\vec{r},T) \ge B$ for T large enough. Therefore we are not minimizing over an empty set. By Assumption 1 [E] we may restrict to minimizations over sets of bounded time.

Take a minimizing sequence \vec{r}_i on $[0, T_i]$ with $\mathcal{B}(\vec{r}_i, T_i) \geq B$ and $\vec{r}_i(0) = \vec{x}$. To prove existence of an optimal \vec{r} we need to show that a converging subsequence of approximate minimizers has a minimal limit. By [8, Lemma 5.18], under Assumption 1 [A] and [B], a set of paths $\vec{r}_i(t)$ over a bounded interval [0, T], having $\vec{r}_i(0)$ in a compact set, and with bounded *I*-functions, is uniformly absolutely continuous, hence pre-compact. Therefore, since any minimizing sequence has bounded *T*, there exists a convergent subsequence, and the limiting path $\vec{r}(t)$ is absolutely continuous. Again under Assumption 1 [A] and [B], [8, Lemma 5.42] shows that the functional $I_{[0,T]}$ is lower semicontinuous in \vec{r} ; therefore, the limiting path satisfies $I_{[0,T]}(\vec{r}) \leq \lim_{i} I_{[0,T]}(\vec{r}_i)$. The continuity of \mathcal{B} in (\vec{r}, T) shows that $\mathcal{B}(\vec{r}, T) \geq B$.

5 Uniqueness

In this section we establish that the solution to the constant coefficient problem is unique. Note that this is only interesting because Problem A has the constraint $\mathcal{B}(\vec{r},T) \geq B$ and T is not determined. If we were simply trying to minimize $I_{[0,T]}(\vec{r})$ with fixed initial and final points, it is trivial to show that the unique solution is a straight line path (e.g., [8, Lemma 5.15]).

Theorem 2 Fix \vec{x} and B > 0. Suppose Assumption 1 [A], [B], [C], and [F] hold. If $\vec{r}(t) \ 0 \le t \le T$ and $\vec{y}(t) \ 0 \le t \le T_1$ solve Problem A, then $T = T_1$ and $\vec{r}(t) = \vec{y}(t), \ 0 \le t \le T$.

Note: in the proof, we need to rescale time in part because the optimal paths may live on different time intervals.

Proof: Without loss of generality assume $T_1 \ge T$. Define $\alpha = T_1/T \ge 1$. By assumption,

$$I_{[0,T]}(\vec{r}) = \mathcal{I} \qquad \qquad \mathcal{B}(\vec{r},T) \ge B$$
$$I_{[0,\alpha T]}(\vec{y}) = \mathcal{I} \qquad \qquad \mathcal{B}(\vec{y},\alpha T) \ge B.$$

Take $\vec{u}(t) \stackrel{\triangle}{=} \alpha \vec{r}(t/\alpha)$. Then $I_{[0,\alpha T]}(\vec{u}) = \alpha \mathcal{I}$ and $\mathcal{B}(\vec{u},\alpha T) \ge f(\alpha)B$. Now for any $0 < \gamma < 1$ define $\vec{v} = \gamma \vec{y}(t) + (1-\gamma)\vec{u}(t)$. Then we have

$$I_{[0,\alpha T]}(\vec{v}) \le (\gamma + (1-\gamma)\alpha)\mathcal{I}, \quad \mathcal{B}(\vec{v},\alpha T) \ge (\gamma + (1-\gamma)f(\alpha))B.$$
(5.1)

Now scale \vec{v} to get $\vec{w}(t) = \delta \vec{v}(t/\delta)$. Then we have

$$I_{[0,\delta\alpha T]}(\vec{w}) \le \delta(\gamma + (1-\gamma)\alpha)\mathcal{I}, \quad \mathcal{B}(\vec{w},\delta\alpha T) \ge f(\delta)(\gamma + (1-\gamma)f(\alpha))B \quad (5.2)$$

where the first inequality is strict unless $\vec{u} = \vec{y}$. Note that since f(1) = 1, so $\gamma + (1 - \gamma)f(\alpha) = \gamma f(1) + (1 - \gamma)f(\alpha)$, we have by (5.2) that

$$\mathcal{B}(\vec{w}, \delta\alpha T) > f(\delta)f(\gamma + (1 - \gamma)\alpha)B \tag{5.3}$$

by the strict convexity of f, as long as $\alpha > 1$ and f is not the identity. If there is strict inequality in (5.3), we may find a $\delta < 1/(\gamma + (1 - \gamma)\alpha)$ so that $\mathcal{B}(\vec{w}, \delta \alpha T) > B$. But then by (5.2), $I_{[0,\delta \alpha T]}(\vec{w}) < \mathcal{I}$, contradicting the assumed minimality of \mathcal{I} . If f is the identity, so that inequality (5.3) is not strict, then we choose $\delta = 1/(\gamma + (1 - \gamma)\alpha)$, and get a contradiction to the assumed minimality of \mathcal{I} from the strict inequality in (5.2).

6 Integral constraint: concave buffer

This section illustrates our assumptions on \mathcal{B} , and derives a result of independent interest for the study of properties of buffer models. This result is not needed for the main argument. We consider the constant coefficient problem, and a functional \mathcal{B} that is a the traditional buffer model, and show that optimizing paths have to be concave in a preferred direction. For a given vector \vec{a} with positive components, we take

$$\frac{d}{dt}\mathcal{B}(\vec{r},t) = \begin{cases} \langle \vec{r}(t), \vec{a} \rangle & \text{if } \langle \vec{r}(t), \vec{a} \rangle > 0 \text{ or } \mathcal{B}(\vec{r},t) > 0 \\ 0 & \text{otherwise,} \end{cases}$$
(6.1)

$$\mathcal{B}(\vec{r},0) = 0. \tag{6.2}$$

Clearly \mathcal{B} satisfies Assumption 1 [C]–[D]¹; we discuss [E] later. We now show [F].

Lemma 3 For any integrable function g(t), we have

$$u_{\alpha}(t) \stackrel{\triangle}{=} \int_{\alpha s}^{\alpha t} \alpha g(w/\alpha) \, dw = \alpha^2 u_1(t). \tag{6.3}$$

Lemma 4 $\mathcal{B}(\vec{r},t)$ defined by (6.1), (6.2) satisfies

$$\mathcal{B}(\vec{r},t) = \sup_{0 \le s \le t} \int_{s}^{t} \langle \vec{r}(w), \vec{a} \rangle \, dw.$$
(6.4)

Corollary 5 Given $\vec{r}(t)$ set $\vec{y}(t) \stackrel{\triangle}{=} \alpha \vec{r}(t/\alpha)$. Then $\mathcal{B}(\vec{y},T) = \alpha^2 \mathcal{B}(\vec{r},\alpha T)$.

These results are immediate; for a proof of Lemma 4 see [8, §13.2], [9].

We begin with an intuitive result: optimal paths never have $\langle \vec{r}(t), \vec{a} \rangle < 0$. If they did, then the buffer could be increased and the cost *I* decreased by eliminating the point. The proof is exactly this argument, but we must account for the shifting of time and position (shift segments of paths to make them continuous) that come from eliminating intervals of time.

Lemma 6 Any minimal $\vec{r}(t)$, $0 \leq t \leq T$ with $\langle \vec{r}(0), \vec{a} \rangle \geq 0$ satisfies $\langle \vec{r}(t), \vec{a} \rangle \geq 0$, $t \in [0, T]$.

¹[D] holds provided $\ell(\vec{s}) < \infty$ for some \vec{s} with $\langle \vec{s}, \vec{a} \rangle > 0$.

Proof: Eliminate any times with $\langle \vec{r}(t), \vec{a} \rangle < 0$, then reconstitute \vec{r} . Mathematically, let

$$s(t) = \int_0^t \mathbf{1}_{\langle \vec{r}(u), \vec{a} \rangle \ge 0} \, du; \tag{6.5}$$

Let $\tau(q) = \inf\{t : s(t) = q\}$ be the first time by which $\langle \vec{r}(u), \vec{a} \rangle \ge 0$ for a total time of q. Then $\tau(s(p)) = s(\tau(p)) = p$, that is, τ and s are inverses. Then define

$$\vec{r}_1(t) = \vec{r}(0) + \int_0^{\tau(t)} \frac{d}{du} \vec{r}(s(u)) \, du.$$
(6.6)

This is well defined since \vec{r} is absolutely continuous. The path $\vec{r}_1(t)$ has the same increments as the path $\vec{r}(t)$ once the times $\langle \vec{r}(t), \vec{a} \rangle < 0$ are eliminated, and it is continuous. Clearly $I_{[0,T]}(\vec{r}) \geq I_{[0,s(T)]}(\vec{r}_1)$. But if $\mathcal{B}(\vec{r},T) = B > 0$ then $\mathcal{B}(\vec{r}_1,s(T)) \geq B$, since the two functions are equal at corresponding (shifted) times whenever $\langle \vec{r}(t), \vec{a} \rangle \geq 0$; see (6.1) or (6.4). In fact, if $\langle \vec{r}(t), \vec{a} \rangle < 0$ anywhere, then $\mathcal{B}(\vec{r}_1, s(T)) > B$. But in this case, by the continuity of \mathcal{B} (assumption 1 [C]) and its definition, there is some $T_1 < s(T)$ so that $\mathcal{B}(\vec{r}_1, T_1) = B$ and, by definition (2.1), since ℓ is positive, $I_{[0,T]}(\vec{r}) > I_{[0,T_1]}(\vec{r}_1)$, so that \vec{r} is not optimal .

Lemma 7 Given B, T and \vec{x} with $\langle \vec{x}, \vec{a} \rangle \ge 0$, any $\vec{r}(t)$ with $\vec{r}(0) = \vec{x}$ that has $\mathcal{B}(\vec{r}, T) = B$ and has minimal cost $I_{[0,T]}(\vec{r})$, has the property that the function $\langle \vec{r}(t), \vec{a} \rangle$ is concave, $0 \le t \le T$.

Proof: Suppose that the function $\langle \vec{r}(t), \vec{a} \rangle$ lies strictly below its concave envelope for some L < t < M. Let v(t) denote the concave envelope of $\langle \vec{r}(t), \vec{a} \rangle$; hence v(t) is linear on [L, M]. By [8, Lemma 5.15], replacing \vec{r} with a linear function over the interval [L, M] strictly decreases I, and since $v(t) > \langle \vec{r}(t), \vec{a} \rangle$ for L < t < M, by (6.1), $\mathcal{B}(\vec{r}, t)$ increases for t > L.

7 Stochastic buffer model

This section gives an overview of the connection between a class of stochastic traffic and buffer models and the variational Problem A. Precise definitions and assumptions will be given later. There are two components in the models: a Markov chain $\vec{x}(t)$ (called the state of the system) and the buffer itself. The Markov chain operates in continuous time and its state lies in the *K*-dimensional lattice with positive integer components. There are J possible transitions, with transition rates $\lambda_i(\vec{x})$ in direction \vec{e}_i , $1 \leq i \leq J$.

This means that the state jumps from \vec{x} to $\vec{x} + \vec{e_i}$ with Poisson rate $\lambda_i(\vec{x})$. The interpretation of the state is that there are x_i sources of traffic in state *i*. There is a single buffer with continuous contents B(t). The buffer contents are filled by each source in state *i* at rate $a_i > 0$, and the total buffer drain rate is *C*. Mathematically, the buffer satisfies the following equation (cf. (6.1)–(6.4)):

$$\frac{d}{dt}B(t) = \begin{cases} \langle \vec{x}(t), \vec{a} \rangle - C & \text{if } \langle \vec{x}(t), \vec{a} \rangle > C \text{ or } B(t) > 0\\ 0 & \text{otherwise.} \end{cases}$$
(7.1)

If B(0) = 0 then it is easy to see that

$$B(t) = \sup_{0 \le s \le t} \int_{u=s}^{t} \langle a, \vec{x}(u) \rangle \, du.$$
(7.2)

We define $\mathcal{B}(\vec{x}, t)$ as this B(t) for a given (state) sample path $\vec{x}(t)$.

As an example, suppose that there are K = 2 types of traffic sources, phone and data. Type 1 sources, phone, generate traffic at rate 1, arrive at Poisson rate λ , and depart at rate μ each. Type 2 sources, data, generate traffic with rate 5, arrive at rate θ , and depart at rate ψ each. Furthermore, a phone source can turn into a data source (a person talks for a while, then starts sending a fax); this occurs at rate γ for each source in state 1, and, of course, a data source can become a phone source with rate δ . We suppose the buffer drain rate is 100. This leads to a model with the following jump rates and transitions:

$$\vec{e}_1 = (1,0) \qquad \lambda_1 = \lambda \qquad \vec{e}_2 = (-1,0) \qquad \lambda_2 = \mu x_1 \\ \vec{e}_3 = (0,1) \qquad \lambda_3 = \theta \qquad \vec{e}_4 = (0,-1) \qquad \lambda_4 = \psi x_2 \\ \vec{e}_5 = (-1,1) \qquad \lambda_5 = \gamma x_1 \qquad \vec{e}_6 = (1,-1) \qquad \lambda_6 = \delta x_2 \\ \frac{d}{dt} B(t) = \begin{cases} x_1(t) + 5x_2(t) - 100 & \text{if } x_1(t) + 5x_2(t) > 100 \text{ or } B(t) > 0 \\ 0 & \text{otherwise.} \end{cases}$$

This is an open model, since sources arrive and depart. For $\lambda_i = 0, 1 \leq i \leq 4$, this is a closed model: transitions are between types of sources, and the total number of sources is constant. We could also allow for correlated arrivals and departures. For example, we could have a person arrive and start to talk on the phone and use his email at the same time; this would be an arrival with $e_7 = (1, 1)$. Similarly, this type of person could depart ($e_8 = (-1, -1)$), stop one of his activities (e_2 or e_4), or a couple could start their day together with $e_9 = (2, 2)$ etc.

We suppose that the process $\vec{x}(t)$ may be scaled to $\vec{z}_n(t)$ with jumps of \vec{z}_n having size \vec{e}_i/n with rate $n\lambda_i(\vec{z}_n(t))$. The scaled buffer $b_n(t)$ is defined similarly to (7.1):

$$\frac{d}{dt}b_n(t) = \begin{cases} \langle z_n(t), \vec{a} \rangle - C & \text{if } \langle z_n(t), \vec{a} \rangle > C \text{ or } b_n(t) > 0\\ 0 & \text{otherwise.} \end{cases}$$
(7.3)

In some models C is also scaled by n so as to have a finite limit as $n \to \infty$. In general we suppose that the jump rates $\lambda_i(\vec{x})$ keep each component of $\vec{z}_n(t)$ positive; that is, we do not consider "sources" that might drain the buffer. Clearly $b_n(t) = \mathcal{B}(\vec{z}_n, t)$ under appropriate scaling.

We showed in [10] that, under a wide variety of assumptions on the $\lambda_i(\vec{x})$, the process $\vec{z}_n(t)$ satisfies a large deviations principle. Furthermore, we showed in [9, 8, Chapter 13], that in a buffer model very similar to the one just described, statistical properties of the buffer size process may also be approximated by solutions to variational problems. (There have been many other analyses of these problems; typically, they consider steady state statistics. See, e.g. Mandjes and Ridder [5] or, for a non-sample-path approach Botvich and Duffield [2].) In particular, it was shown that the steady-state probability that the buffer exceeds a level B is given by $\exp(-nI+o(n))$, where I is the solution to a variational problem we describe below.

7.1 The variational problem for Buffer overflow

Given the rates and jump directions, define the local cost function by

$$\ell(\vec{x}, \vec{y}) \stackrel{\triangle}{=} \sup_{\vec{\theta} \in \mathbb{R}^K} \left(\langle \vec{\theta}, \vec{y} \rangle - \sum_i \lambda_i(\vec{x}) \left(\exp\langle \vec{\theta}, \vec{e}_i \rangle - 1 \right) \right).$$
(7.4)

Then define the action functional, or cost of an absolutely continuous function $\vec{r}(t) \in \mathbb{R}^{K}$

$$I_{[0,T]}(\vec{r}) \stackrel{\triangle}{=} \int_0^T \ell\left(\vec{r}(t), \frac{d}{dt}\vec{r}(t)\right) dt.$$
(7.5)

The variational problem associated with making a buffer exceed level B is given as follows. We assume that the process is typically near its steady state value, denoted by \vec{q} , and that at that point the capacity C suffices to serve the traffic, namely $\langle \vec{q}, \vec{a} \rangle < C$. Let \mathcal{I} be defined by

$$\mathcal{I} = \inf \left\{ I_{[0,T]}(\vec{r}) : \vec{r}(0) = \vec{q}, \ T < \infty, \ \mathcal{B}(\vec{r},T) \ge B \right\}.$$
 (7.6)

Usually the optimizing T is infinite, since the time to leave \vec{q} is infinite, so we perform a standard shifting of time so that the upcrossing of the path \vec{r} to the hyperplane $\langle \vec{r}, \vec{a} \rangle = C$ takes place at time 0. The time T after time 0 required to fill the buffer is then typically finite. If there is a unique solution (T, \vec{r}) with $I_{[-\infty,T]}(\vec{r}) = \mathcal{I}$ and $\mathcal{B}(\vec{r}, T) \geq B$ then equation (1.3) holds; in other words, given a buffer overflow, we know how it was almost certain to have happened. For a proof of this assertion see [8, Chapter 13].

The problem of finding the minimal cost path \vec{r} starting from \vec{q} and making $\mathcal{B}(\vec{r},T) \geq B$ naturally splits into finding the cheapest path from \vec{q} to sone \vec{x} with $\langle \vec{x}, \vec{a} \rangle = C$, followed by the cheapest path starting from \vec{x} making $\mathcal{B}(\vec{r},T) \geq B$, and then minimizing the total cost I over all such \vec{x} . This is clearly valid because the buffer cannot begin to increase until $\langle \vec{r}, \vec{a} \rangle > C$. It is also clear that the cost $I_{[0,T]}(\vec{r})$ of making a buffer Btends to zero as $B \to 0$ (since we can take a small period of time to make the small buffer, and ℓ is finite for some directions). Therefore, if there is a unique upcrossing point \vec{x}^* , then for small B, the optimal point x (in the full problem of making a small buffer) will be close to \vec{x}^* , the optimal upcrossing point.

7.2 Typical behavior

To describe the typical behavior of the state and buffer size processes, define the process $\vec{z}_{\infty}(t)$

$$\vec{v}(\vec{x}) \stackrel{\triangle}{=} \sum_{i} \lambda_i(\vec{x}) \vec{e}_i, \quad \frac{d}{dt} \vec{z}_{\infty}(t) = \vec{v}(\vec{z}_{\infty}(t)).$$
(7.7)

Kurtz's theorem [8, Chapter 5.1] states that, over finite time intervals, the process $\vec{z}_n(t)$ is extremely likely to remain close to the path $\vec{z}_{\infty}(t)$ that starts at the same point $\vec{z}_{\infty}(0) = \vec{z}_n(0)$. The intuition is that \vec{v} is the average drift of the process $\vec{z}_n(t)$, in the sense that

$$\lim_{\delta \downarrow 0} \lim_{n \to \infty} \frac{\vec{z}_n(t+\delta) - \vec{z}_n(t)}{\delta} - \vec{v}(\vec{z}_n(t)) = 0 \text{ with probability one.}$$
(7.8)

From this one can infer [9, Cor. 8] that the buffer size process b_n is also extremely likely to remain close to a limiting path, which can be calculated.

We will show next that given an initial point \vec{x} with $\langle \vec{x}, \vec{a} \rangle = C$, and given a small B, there is an *essentially unique* path \vec{r} with $\vec{r}(0) = \vec{x}$ and $\mathcal{B}(\vec{r},T) \geq B$). This clearly does not solve the full uniqueness problem, because although we know that $\vec{x} \approx \vec{x}^*$, for each B we have no idea whether there is a unique associated point \vec{x} .

8 Weak uniqueness

We now state the assumptions under which the probabilistic problem of the buffer being large is described by the deterministic variational Problem A.

Assumption 2

- A. The processes $\vec{z}_n(t)$ and $b_n(t)$ satisfy a sample-path large deviations principle; the rate function for \vec{z}_n is $I_{[0,T]}$.
- B. There is a unique globally attracting point \vec{q} of $\vec{z}_{\infty}(t)$.
- C. $\langle \vec{q}, \vec{a} \rangle < C$.
- D. The Freidlin-Wentzell theory [8, Chapter 6] applies to both $\vec{z}_n(t)$ and $b_n(t)$.
- E. There is a unique endpoint \vec{x}^* of the upcrossing problem from \vec{q} to $\{\vec{x}: \langle \vec{x}, \vec{a} \rangle = C\}.$
- F. $\langle \vec{x}^*, \vec{v}(\vec{x}^*) \rangle < 0.$

Since \mathcal{B} is a continuous functional, the assumption that the buffer process satisfies a large deviations principle is a consequence of the fact that \vec{z}_n satisfies the principle, and the rate function for b_n can be derived from $I_{[0,T]}$ [8, §2.3]: see e.g. [9, Cor. 8].

Theorem 8 Let \mathbb{P}_{ss} denote steady state probability. Under Assumption 2,

$$\lim_{n \to \infty} \frac{1}{n} \log \mathbb{P}_{ss}(\mathcal{B}(\vec{z}_n, t) \ge B) = -\inf_{\vec{r}, T} \left\{ I_{[0,T]}(\vec{r}) : \vec{r}(0) = \vec{q}, \ \mathcal{B}(\vec{r}, T) \ge B \right\}.$$

Furthermore, if \vec{r} is unique to within a time shift, then (shifting time so that the buffer overflow takes place at time t = 0) the estimate (1.3) holds. That is, $\vec{z}_n(t) \approx \vec{r}(t)$ over each fixed interval of time before the time of overflow, assuming that overflow occurs at time T.

The proof of Theorem 8 is standard from the Freidlin-Wentzell theory, and will not be given here; see [4] or [8, Chapter 6, 13]. However, we must explain when Assumption 2 can hold. Part A was proved in [10], under the following conditions. The process $\vec{z}_n(t)$ is assumed to live on the positive quadrant G (in fact, the domain G in [10] is quite general). The jump rates $\lambda_i(\vec{x})$ were assumed Lipschitz continuous, strictly positive in the interior of the domain G, and $\lambda_i(\vec{x}) \to 0$ as $x \to \vec{y} \in \partial G$ for every *i* such that a jump in direction $\vec{e_i}$ from \vec{y} would take $\vec{z_n}$ out of G. The associated $\vec{e_i}$ were assumed to positively span all of \mathbb{R}^K (that is, for any $\vec{x} \in \mathbb{R}^K$ there are $\beta_i \geq 0$ such that $\vec{x} = \sum_i \vec{e_i} \beta_i$). There were some further technical conditions that are quite minor and will not be described here. Part B of Assumption 2 is simply that the process $\vec{z_n}(t)$ tends to go to a small neighborhood of \vec{q} . Part C means that the buffer B(t) tends to decrease when the process is in its most likely state. Part D implies that steady-state quantities can be calculated from the transient sample-path large deviations principle [8, Chapters 6,13]. Part F means once the process $\vec{z_n}(t)$ reaches the most likely point where the buffer can begin to fill, the process tends to go in a direction that has the buffer decrease. This means that, most likely, even when the buffer begins to fill, it will almost immediately begin to decrease.

The assumptions can be largely eliminated if we change the question we ask. Suppose we want to know how the buffer fills, starting from some point \vec{x} with $\langle \vec{x}, \vec{a} \rangle = C$, and how likely it is to fill to a level B.

Theorem 9 Under Assumption 2 A, given \vec{x} with $\langle \vec{x}, \vec{a} \rangle = C$,

$$\lim_{n \to \infty} \frac{1}{n} \log \mathbb{P}(\mathcal{B}(\vec{z}_n, T) \ge B) = -\inf \{ I_{[0,T]}(\vec{r}) : \vec{r}(0) = \vec{x}, \ T, \ \mathcal{B}(\vec{r}, T) \ge B \}.$$

Furthermore, if \vec{r} (and $T < \infty$) are unique then for every $\varepsilon > 0$

$$\lim_{n \to \infty} \mathbb{P}\left(\sup_{0 \le t \le T} \left| \vec{z}_n(t) - \vec{r}(t) \right| < \varepsilon \mid \mathcal{B}(\vec{z}_n, T) \ge B\right) = 1.$$
(8.1)

The proof is standard: see [9].

It is because of Theorem 8 and (8.1) of Theorem 9 that we are motivated to study uniqueness for solutions to Problem A. This concludes our discussion of the connection between the probability and variational problems. We return to the study of the variational problem.

Suppose that there is a unique upcrossing point \vec{x}^* , the lowest cost point from \vec{q} to the hyperplane $\langle \vec{x}, \vec{a} \rangle = C$. Consider the local cost function $\ell(\vec{x}^*, \vec{y})$, which is defined for any (\vec{x}, \vec{y}) in (7.4). When viewed as a constant coefficient cost $\ell(\vec{y})$, by [8, Chapter 5.2] $\ell(\vec{x}^*, \vec{y})$ satisfies Assumption 1 [A], [B] and [E] and, for nontrivial Markov models, also [D]. When combined with the functional \mathcal{B} given in (7.1), it is easy to check that all of Assumption 1 [A]–[F] hold (see [8, Chapter 5.2] for properties of $\ell(\vec{y})$, Chapter 13 for properties of \mathcal{B}). Therefore, the solution to Problem A for constant coefficient cost function, with paths starting at \vec{x}^* making buffer at least B, exists and is unique. For the case B = 1 we call the resulting solution $\vec{r}(t)$, and the time *T*. Actually, the path may start at any \vec{x}_1 with $\langle \vec{x}, \vec{a} \rangle = C$, as long as we take the cost function given by $\ell(\vec{x}^*, \vec{y})$, not $\ell(\vec{x}_1, \vec{y})$, and then the solution will be $\vec{r}(t) + \vec{x}_1 - \vec{x}^*$, since $I_{[0,T]}$ is invariant under shifts, while \mathcal{B} is invariant under shifts along the hyperplane $\{\vec{x} : \langle \vec{x}, \vec{a} \rangle = 0\}$. Furthermore, our scaling property gives the optimal path for any $B \neq 1$ as a scaled version of \vec{r} , namely $\alpha \vec{r}(t/\alpha) + \vec{x}^* - \alpha \vec{x}^*$, for $0 \leq t \leq T\alpha$, where $\alpha = \sqrt{B}$.

We now switch back to cost function $\ell(\vec{x}, \vec{y})$ depending on \vec{x} . Many properties of this function are detailed in [8, Chapter 5]. If the positive cone spanned by the \vec{e}_i is the entire space \mathbb{R}^K , and if all $\lambda_i(\vec{x}) > 0$, then $0 \leq \ell(\vec{x}, \vec{y}) < \infty$, and $\ell(\vec{x}, \vec{y})$ is convex in \vec{y} . [8, Thm. 5.35] shows that for any $\varepsilon > 0$, c > 0, there exists a $\delta > 0$ such that if $J_{[0,T]}(\vec{r})$ is defined to be the same as $I_{[0,T]}(\vec{r})$ except using the local cost $\ell(\vec{x}^*, \frac{d}{dt}\vec{r}(t))$, and if $J_{[0,T]}(\vec{r}) \leq c$ and $|\lambda_i(\vec{r}(t)) - \lambda_i(\vec{x}^*)| < \delta$ then $|I_{[0,T]}(\vec{r}) - J_{[0,T]}(\vec{r})| < \varepsilon$. In other words, the functional $I_{[0,T]}$ is uniformly continuous in sets of bounded cost. Also, [8, Prop. 5.46] shows that for any $c, T < \infty$, the set of $\vec{r}(t)$ with $\vec{r}(0)$ in a compact set, and with $I_{[0,T]}(\vec{r}) \leq c$, is compact. Compactness is in the sup norm topology for the paths in $[0,T] \to \mathbb{R}^K$.

The basis of our argument is that when B is small, paths $\vec{r}_B(t)$, $0 \leq t \leq T_B$, can stay very near $\vec{r}_B(0)$ and have T_B small, and still make $\mathcal{B}(\vec{r}_B, T_B) = B$. So the local cost function $\ell(\vec{r}_B(t), \frac{d}{dt}\vec{r}_B(t))$ is very nearly equal to $\ell(\vec{r}_B(0), \frac{d}{dt}\vec{r}_B(t))$, a constant coefficient cost, for which we know uniqueness applies. This motivates the following cost function, which is based on considering a small neighborhood of $\vec{r}_B(0)$. Given B and $\vec{r}_B(0)$ with $\langle \vec{r}_B(0), \vec{a} \rangle = C$, define

$$\ell_B(\vec{x}, \vec{y}) \stackrel{\triangle}{=} \ell\left(\vec{r}_B(0) + \sqrt{B}(\vec{x} - \vec{r}_B(0)), \vec{y}\right). \tag{8.2}$$

For any such path $\vec{r}_B(t)$ that has $\mathcal{B}(\vec{r}(T_B)) = B$ we define a scaled path

$$\vec{s}_B(t) \stackrel{\triangle}{=} \vec{r}_B(0) + \frac{1}{\sqrt{B}} \left(\vec{r}_B(t\sqrt{B}) - \vec{r}_B(0) \right), \ 0 \le t \le T_B/\sqrt{B}.$$
(8.3)

A simple calculation shows that

$$\mathcal{B}\left(\vec{s}_B, T_B/\sqrt{B}\right) = 1. \tag{8.4}$$

Now we find

$$\int_{0}^{T_{B}/\sqrt{B}} \ell_{B}\left(\vec{s}_{B}(t), \frac{d}{dt}\vec{s}_{B}(t)\right) dt
= \int_{0}^{T_{B}/\sqrt{B}} \ell_{B}\left(\vec{r}_{B}(0) + \frac{1}{\sqrt{B}}(\vec{r}_{B}(t\sqrt{B}) - \vec{r}_{B}(0)), \frac{d}{dt}\vec{r}_{B}(t\sqrt{B})\right) dt
= \int_{0}^{T_{B}/\sqrt{B}} \ell\left(\vec{r}_{B}(0) + (\vec{r}_{B}(t\sqrt{B}) - \vec{r}_{B}(0)), \frac{d}{dt}\vec{r}_{B}(t\sqrt{B})\right) dt$$

$$= \frac{1}{\sqrt{B}} \int_{0}^{T_{B}} \ell\left(\vec{r}_{B}(u), \frac{d}{du}\vec{r}_{B}(u)\right) du.$$
(8.5)

It is easy to show that, under Assumption 2 E, as $B \to 0$, minimal solutions to Problem A starting from $\vec{r}_B(0)$ with $\langle \vec{r}_B(0), \vec{a} \rangle = C$ must have T_B/\sqrt{B} bounded, since there are paths $\vec{r}(t)$ that have T/\sqrt{B} bounded, and have cost of order \sqrt{B} , and any path that stays above $\langle \vec{r}(t), \vec{a} \rangle = C$ for time T has cost that is bounded below by a function linear in T. Therefore, if we consider solutions to Problem A with buffer size B small, then the scaled problem with ℓ_B cost function has solution \vec{s}_B in bounded time T_B/\sqrt{B} .

Theorem 10 Suppose all of Assumption 2 holds, and further that the upcrossing point \vec{x}^* is unique. For any set of paths $\vec{r}_B(t)$ satisfying $\mathcal{B}(\vec{r}(T_B)) = B$ and with minimal cost, starting from the point \vec{q} , the scaled paths $\vec{s}_B(t)$ satisfy the following.

$$\lim_{B \to 0} \frac{T_B}{\sqrt{B}} = T \tag{8.6}$$

$$\lim_{B \to 0} \vec{s}_B(t) = \vec{r}(t), \ 0 \le t \le T, \ uniformly \ in \ t.$$
(8.7)

This theorem gives a connection between solutions to Problem A for constant coefficient costs, and solutions with costs depending on \vec{x} , but for small buffer B. Proof of existence of solutions to Problem A for non constant coefficient costs is the same as the proof for Theorem 1.

Proof: By [8, Prop. 5.46], the paths $\vec{s}_B(t)$ lie in a compact set, since they have bounded cost and their initial points $\vec{s}_B(0)$ approach \vec{x}^* . Take a convergent subsequence of \vec{s}_B . By [8, Thm. 5.35], the costs (based on ℓ_B) of these paths converge to the cost of the constant coefficient path starting at \vec{x}^* and making buffer size 1 in time T. But Theorem 2 shows that this path is unique among paths that start at \vec{x}^* and make buffer size 1. Therefore any limiting sequence of the \vec{s}_B must converge to the same path, and do so uniformly, and their times must converge also. We note that the point \vec{x}^* will be unique if the Markov chain can be written as a superposition of sources, either an open model (with productform Poisson steady-state distribution) or a closed model (with multinomial distribution). In these cases, \vec{x}^* is the solution of Sanov's theorem (see [8, Chapter 2]). Explicitly, if the steady-state distribution of the chain is $\pi(i)$, $1 \leq i \leq K$, then the upcrossing point is given by the minimum entropy point (i.e., minimum Kullback-Leibler measure) on the hyperplane $\langle \vec{x}, \vec{a} \rangle = C$. If the level curves of this entropy are strictly convex, then the point \vec{x}^* is unique. But it is easy to see that these level curves are strictly convex for the two models under study, and the strict convexity also holds in the limit as we scale $n \to \infty$.

9 Strong uniqueness

The preceding result may be combined with traditional variational calculus to show strong (true) uniqueness for the solution to Problem A from a given initial point $\vec{r}(0)$ where $\langle \vec{r}(0), \vec{a} \rangle = C$. As mentioned before, this result still does not give full uniqueness for the steady-state variational problem, because that problem has $\vec{r}(0) = \vec{q}$ where \vec{q} is the unique attracting point for the fluid limit $\vec{z}_{\infty}(t)$, and which satisfies $\langle \vec{q}, \vec{a} \rangle < C$. Nevertheless, it is a step in that direction.

Theorem 11 If the positive cone of the $\{\vec{e}_i\}$ equals \mathbb{R}^K , and $\log \lambda_i(\vec{x})$ are Lipschitz and C^2 functions, then given \vec{x}^* with $\langle \vec{x}^*, \vec{a} \rangle = C$, there are B_0 and $\delta > 0$ so that for all $B \leq B_0$ and $|\vec{x} - \vec{x}^*| \leq \delta$ with $\langle \vec{x}, \vec{a} \rangle = C$, the solution to Problem A starting at $\vec{r}(0) = \vec{x}$ is unique.

References

- Anick, D. D. Mitra, and M.M. Sondhi, Stochastic theory of a datahandling system with multiple sources, Bell System Tech. J., 61, pp. 1871–1894, 1982.
- Botvich, D. and N. Duffield, Large deviations, the shape of the loss curve, and economies of scale in large multiplexers, Queueing Systems 20, pp. 293–320, 1995.
- [3] Courcoubetis, C. and R. Weber, Buffer overflow asymptotics for a buffer handling many traffic sources, J. Appl. Prob., 33, pp. 886–903, 1996.

- [4] Freidlin, M.I. and A.D. Wentzell. Random Perturbations of Dynamical Systems, Springer-Verlag, New York, 1998.
- [5] Mandjes, M. and A. Ridder, Optimal trajectory to overflow in a queue fed by a large number of sources, Queueing Systems, 31, pp. 137–170, 1999.
- [6] Mandjes, M. and P. Mannersalo, Queueing systems fed by many exponential on-off sources: an infinite intersection approach, Queueing Systems 54, pp. 5-20, 2006.
- [7] Simonian, A. and and J. Guibert., Large deviations approximation for fluid sources fed by a large number of on/off sources, IEEE J. Selected Areas Commun. 13, pp. 1017–1027, 1995.
- [8] Shwartz, A. and A. Weiss, *Large Deviations for Performance Analysis*, Chapman and Hall, 1995.
- [9] Shwartz, A. and A. Weiss, Multiple time scales in Markovian ATM models I. Formal calculations, Technion, 1999.
- [10] Shwartz, A. and A. Weiss, Large deviations with diminishing rates, Math. of Oper. Res. 30 pp. 281–310, 2005.
- [11] Weiss, A. A new technique for analyzing large traffic systems, Adv. Appl. Prob., 18, pp. 506–532, 1986.