CONVERGENCE PROPERTIES FOR DISCRETE-TIME NONLINEAR SYSTEMS

DUC N. TRAN, BJÖRN S. RÜFFER, AND CHRISTOPHER M. KELLETT

ABSTRACT. Three similar convergence notions are considered. Two of them are the long established notions of convergent dynamics and incremental stability. The other is the more recent notion of contraction analysis. All three convergence notions require that all solutions of a system converge to each other. In this paper, we investigate the differences between these convergence properties for discrete-time, timevarying nonlinear systems by comparing the properties in pairs and using examples. We also demonstrate a time-varying smooth Lyapunov function characterization for each of these convergence notions, and, with appropriate assumptions, we provide several sufficient conditions to establish relationships between these properties in terms of Lyapunov functions.

1. INTRODUCTION

Convergence properties, namely incremental stability [1–3], convergent dynamics [4–6], and contraction analysis [7], are established methods to characterize the asymptotic behavior of one solution with respect to any other solution of a (nonlinear) dynamical system. In particular, all convergence properties impose conditions that all solutions "forget" their initial conditions and converge to each other. This is a highly desirable property in solving problems in nonlinear output stabilization and regulation [3, 5], synchronization [2, 8–10], observer design [2, 7], steady-state and frequency response analysis of nonlinear systems [11–13], and others. These three notions of convergence are similar and largely related to each other. However, the three methods were all derived independently, were motivated distinctly, and employ different tool sets. As a consequence, their mutual relationships are, in general, not fully understood even for continuous-time nonlinear systems in the literature. Thus, the purpose of this study is to investigate explicit differences between these properties.

Incremental stability describes the asymptotic property of differences between any two solutions. Specifically, an augmented system is formed by two "copies" of the original system. Then, global asymptotic stability of a special closed set (called the diagonal) with respect to the augmented system can be demonstrated to be equivalent to incremental stability of the original system. Thus, a Lyapunov characterization for incremental stability of the original system can be derived from a classical Lyapunov characterization of global asymptotic stability. Historically, the origin of incremental stability goes back to works on forced oscillators by German mathematicians Trefftz [14] and Reissig [15–17], without the property being named. It was LaSalle who in [18] attributes the term "extreme stability" to Reissig's work [15]. Subsequently, the work of Yoshizawa [1, Chapter IV] made significant contributions to the study of extreme stability. In particular, for continuous-time, time-varying systems, a continuous Lyapunov function characterization of extreme stability was demonstrated [1, Theorem 21.1]. Similar results are found in [2, Theorem 1] and [19, Theorem 5] for time-invariant and time-varying systems, respectively, where in both works the property is termed incremental stability. Incremental stability for input-output operators is presented in [20], and in [21] conclusions about (asymptotic) stability of equilibria and periodic solutions are drawn from linear incremental gain properties. Incremental Input-to-State Stability [2] is an extension of incremental stability to systems with input using the Input-to-State Stability (ISS) framework [22].

Convergent dynamics, on the other hand, requires the existence of a unique and asymptotically stable reference solution that is bounded for all (backward and forward) time. However, a priori knowledge of this reference solution is not necessarily required. Every other solution, then, converges to this reference solution asymptotically. A convenient, sometimes easy to check, sufficient condition of convergent dynamics is the Demidovič condition. Essentially, using a quadratic Lyapunov function, exponential convergence of all solutions to a reference solution can be demonstrated (see, for instance, [6]). The existence and boundedness of the reference solution usually relies on Yakubovich's Lemma; i.e., that a compact and positively invariant set contains at least one bounded solution defined for all times. This result was demonstrated in [23, Lemma 2] based on ideas of Demidovič in [24,25]. Convergent dynamics (or convergent systems) was pioneered by Demidovič [4] (only available in Russian), see also a historical perspective of developments for convergent dynamics in [6]. Results on the convergent dynamics property for periodic systems are discussed in [26]. A converse Lyapunov theorem for globally convergent systems was introduced in [19, Theorem 7]. More details and further extensions of convergent dynamics to systems with input (input-to-state convergence) and applications to output regulation problems can be found in [5].

Lastly, contraction analysis [7] was inspired by fluid mechanics and differential geometry. Contraction analysis utilizes local analysis of the linearization along every trajectory to establish globally convergent behavior. This method, in essence, generalizes linear eigenvalue analysis to nonlinear systems. As a result, contraction analysis provides a characterization for exponential convergence of one solution with respect to any other solution. The contraction analysis approach, thus, independently extends the Demidovič sufficient condition to a necessary and sufficient condition. (Yet another approach, dating back at least to the 1950s, that allows linearizing nonlinear finite-dimensional systems into infinite-dimensional linear systems is via the Koopman operator [27]—the adjoint of the transfer or Frobenius-Perron operator.) Contraction analysis (or contraction metrics) was first introduced in [7]. Since then, developments based on differential geometry have been made in [28], [29] including converse theorems on contraction metrics for an equilibrium in [30]. A historical perspective of contraction analysis, including earlier closely related concepts, is presented in [31]. Various applications of contraction analysis can be found in [8,32].

In the continuous-time setting, several relationships have been established between these properties. Available comparisons include convergent dynamics versus incremental stability [19] and extending contraction analysis and incremental stability to the same differential geometric framework [28]. In [33] it is shown that exponential incremental stability is equivalent to exponential (asymptotic) stability of one and of all trajectories of a system, and how this relates to the linearization of the system, assuming the state space is closed and convex.

In this paper, we study and compare the three previously discussed convergence properties for time-varying nonlinear systems, albeit, in contrast to most of the existing literature, in discrete-time. In particular, we show that convergent dynamics and incremental stability are two distinct properties. Similarly, convergent dynamics and contraction analysis are two distinct properties. However, contraction analysis is a special case of incremental stability, namely, exponential incremental stability. Furthermore, we establish various sufficient conditions to demonstrate relationships between the three convergence properties. This is done by either tightening the regularity requirements on the dynamics or by adding an assumption on the state space. We also provide time-dependent smooth Lyapunov function characterizations for each of the properties. Furthermore, we present discrete-time analogues of the Demidovič condition for incremental stability (similar to the continuous-time result [19, Equation 15]), convergent dynamics (similar to the continuous-time result [6, Theorem 1]), and contraction analysis.

To derive the three Lyapunov function characterizations, we rely heavily on a converse Lyapunov result for time-varying systems with an asymptotically stable set [34, Theorem 1] and a stronger version of this theorem for time-varying systems with an exponentially stable set [34, Theorem 2]. Specifically, we use a result [2, Lemma 2.3] to convert global asymptotic (or exponential) incremental stability to global asymptotic (or exponential) stability of a closed set with respect to an augmented system. For convergent dynamics, we first apply a nonlinear change of coordinates with respect to the reference solution, then we use a standard Lyapunov result [34, Theorem 1] (similar to [35, Theorem 23] for the continuous-time case) to obtain the desired characterization. Lastly, we demonstrate that contraction analysis, in fact, is equivalent to exponential incremental stability, and hence, is equivalent to the existence of an exponential incremental Lyapunov function.

To demonstrate the differences between the three considered convergence notions, we construct examples of systems that individually satisfy one of the three properties but not the others. Specifically, we exploit the fact that global incremental stability (any two trajectories tend to each other asymptotically) is a stronger property than global asymptotic convergence to a single trajectory. From there, we are able to construct a system (Example 1) showing that convergent dynamics does not imply incremental stability. On the other hand, we exploit another fact of convergent systems: that there must exist at least one bounded solution. This is not necessarily true for asymptotically incrementally stable systems. Thus, we show that incremental stability does not imply convergent dynamics (Example 2).

In contrast to the other two properties, contraction analysis explicitly requires continuously differentiable dynamics. Given a system with differentiable right hand side, we are able to show that contraction analysis is equivalent to exponential incremental stability (Theorem 15). As a consequence, contraction analysis does not imply convergent dynamics. Finally, we employ the fact that the convergence rate in contraction analysis is always exponential. This is not necessarily true for convergent systems (Example 3). Therefore, contraction analysis and convergent dynamics are distinct convergence notions.

The paper is organized as follows: the necessary technical assumptions, notational conventions, and definitions of convergence properties are provided in Section 2. Then, Lyapunov function characterizations for these notions are presented in Section 3. In Section 4, comparisons in pairs for the three properties are presented together with various examples and sufficient conditions. Conclusions and future research indications are provided in Section 5. Several proofs are collected in the appendix.

2. PRELIMINARIES

Let \mathbb{Z} denote the set of integers and $\mathbb{Z}_{\geq k_0}$ the set of integers greater or equal to k_0 . We consider discrete-time nonlinear time-varying systems described by the difference equation

$$x(k+1) = f(k, x(k)), \quad x(k) \in \mathbb{R}^n, k \in \mathbb{Z},$$
(1)

where $f: \mathbb{Z} \times \mathbb{R}^n \to \mathbb{R}^n$ is continuous and $x(k_0) = \xi \in \mathbb{R}^n$ for $k_0 \in \mathbb{Z}$. For any $k_0 \in \mathbb{Z}$ and $\xi \in \mathbb{R}^n$, the solution of system (1) is a function $\phi: \mathbb{Z}_{\geq k_0} \to \mathbb{R}^n$, parameterized by initial state and time, and satisfying $\phi(k_0; k_0, \xi) = \xi$ and equation (1); i.e., $\phi(k+1; k_0, \xi) = f(k, \phi(k; k_0, \xi))$ for all $k, k_0 \in \mathbb{Z}$ such that $k \geq k_0$. We use standard comparison function classes¹ $\mathcal{K}, \mathcal{L}, \mathcal{K}_\infty$, and \mathcal{KL} (see [36]). For a vector $y \in \mathbb{R}^n$, a matrix $P \in \mathbb{R}^{n \times n}$, and a positive definite matrix Q we denote the Euclidean norm by |y|, the induced (matrix) norm by $||P|| = \max_{x \in \mathbb{R}^n, x \neq 0} \frac{|Px|}{|x|}$, and the induced norm of y with respect to matrix Q by $||y||_Q = \sqrt{y^T Q y}$. Given two symmetric matrices A and B in $\mathbb{R}^{n \times n}$ we write $A \preceq B$ if for all $x \in \mathbb{R}^n, x^T A x \leq x^T B x$. We say that a set $\mathbb{X} \subseteq \mathbb{R}^n$ is positively invariant under (1) if for all $\xi \in \mathbb{X}$ and $k, k_0 \in \mathbb{Z}$ such that $k \geq k_0$, we have $\phi(k; k_0, \xi) \in \mathbb{X}$.

In the following subsections, we recall the definitions of incremental stability, convergent dynamics, and contraction analysis.

2.1. Incremental Stability. The following definition of asymptotic incremental stability is a discrete-time analogue to that in [2, Definition 2.1].

¹Recall that $\alpha : \mathbb{R}_{\geq 0} \to \mathbb{R}_{\geq 0}$ is of class- \mathcal{K} if it is continuous, zero at zero, and strictly increasing. If $\alpha \in \mathcal{K}$ is unbounded, it is of class- \mathcal{K}_{∞} . A function $\sigma : \mathbb{R}_{\geq 0} \to \mathbb{R}_{\geq 0}$ is of class- \mathcal{L} if it is continuous, strictly decreasing, and $\lim_{t\to\infty} \sigma(t) = 0$. A function $\beta : \mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0} \to \mathbb{R}_{\geq 0}$ is of class- $\mathcal{K}\mathcal{L}$ if it is class- \mathcal{K} in its first argument and class- \mathcal{L} in its second argument. By convention, $\beta \in \mathcal{K}\mathcal{L}$ satisfies $\beta(0, t) = 0$ for all $t \in \mathbb{R}_{>0}$.

Definition 1. System (1) is uniformly asymptotically incrementally stable in a positively invariant set $\mathbb{X} \subseteq \mathbb{R}^n$ if there exists $\beta \in \mathcal{KL}$ such that

$$|\phi(k;k_0,\xi_1) - \phi(k;k_0,\xi_2)| \le \beta(|\xi_1 - \xi_2|, k - k_0), \tag{2}$$

holds for all $\xi_1, \xi_2 \in \mathbb{X}$ and $k, k_0 \in \mathbb{Z}$ such that $k \geq k_0$. In case $\mathbb{X} = \mathbb{R}^n$, we say that system (1) is uniformly globally asymptotically incrementally stable.

As is standard, *uniform* here refers to the fact that the bound in (2) depends only on the elapsed time $k-k_0$. A strictly stronger property requires an exponential rate of convergence.

Definition 2. System (1) is uniformly exponentially incrementally stable in a positively invariant set $\mathbb{X} \subseteq \mathbb{R}^n$ if there exist $\kappa \geq 1$ and $\lambda > 1$ such that

$$|\phi(k;k_0,\xi_1) - \phi(k;k_0,\xi_2)| \le \kappa |\xi_1 - \xi_2| \lambda^{-(k-k_0)}$$
(3)

holds for all $\xi_1, \xi_2 \in \mathbb{X}$ and $k, k_0 \in \mathbb{Z}$ such that $k \ge k_0$. In case $\mathbb{X} = \mathbb{R}^n$, we say that system (1) is uniformly globally exponentially incrementally stable.

2.2. Convergent Dynamics. The following definition of convergent dynamics for discrete-time systems is recalled from [11, Definition 1]. This definition is a discrete-time analogue to the continuous-time definition in [6, Definition 1].

Definition 3. System (1) is uniformly convergent in a positively invariant set $\mathbb{X} \subseteq \mathbb{R}^n$ if

- (1) there exists a unique bounded solution $\bar{x}(k)$ of system (1) defined in \mathbb{X} for all $k \in \mathbb{Z}$;
- (2) there exists a function $\beta \in \mathcal{KL}$ such that, for all $\xi \in \mathbb{X}$, $k, k_0 \in \mathbb{Z}$ with $k \geq k_0$, we have

$$|\phi(k;k_0,\xi) - \bar{x}(k)| \le \beta(|\xi - \bar{x}(k_0)|, k - k_0).$$
(4)

In case $\mathbb{X} = \mathbb{R}^n$, we say that system (1) is uniformly globally convergent. As in the previous definition, we say system (1) is uniformly (globally) exponentially convergent if the \mathcal{KL} -function β can be chosen to be $\beta(s, k - k_0) = \kappa s \lambda^{-(k-k_0)}$ for some $\kappa \geq 1$ and $\lambda > 1$.

Remark 4. In continuous-time, the definition of convergent dynamics [6, Definition 1] requires an additional condition that system (1) is forward complete. In discrete-time, such a condition is not required because, by definition, the range of f is \mathbb{R}^n .

Remark 5. In fact, condition 2) and the boundedness of $\bar{x}(k)$ in Definition 3 also imply that $\bar{x}(k)$ is unique. Indeed, let $\hat{x}(k)$ be another solution defined and bounded for all $k \in \mathbb{Z}$, then $|\hat{x}(k_0) - \bar{x}(k_0)|$ is bounded for all $k_0 \in \mathbb{Z}$. Since (4) holds for all solutions, taking the limit for $k_0 \to -\infty$ in (4), it follows that $|\hat{x}(k) - \bar{x}(k)| \leq 0$ for all $k \in \mathbb{Z}$. Hence, $\hat{x} \equiv \bar{x}$.

2.3. Contraction Analysis. Unlike the previous two properties, contraction analysis explicitly requires the dynamics of the considered system to be continuously differentiable. As a consequence, in this subsection, we make a standing assumption that the mapping f (or the right hand side) of system (1) is continuously differentiable in x on \mathbb{R}^n . We are now ready to state the formal definition (following [7]) of contraction analysis.

Definition 6. Suppose the right hand side of system (1) is continuously differentiable in x on a positively invariant set $\mathbb{X} \subseteq \mathbb{R}^n$. Then system (1) is uniformly contracting in \mathbb{X} if there exist a nonsingular-matrix-valued function $\Theta: \mathbb{Z} \times \mathbb{X} \to \mathbb{R}^{n \times n}$ and constants $\mu, \eta, \rho > 0$ such that, for all $x \in \mathbb{X}, k \in \mathbb{Z}$, we have

$$\eta I \preceq \Theta(k, x)^T \Theta(k, x) \preceq \rho I, \tag{5}$$

$$F(k,x)^T F(k,x) - I \preceq -\mu I, \tag{6}$$

where the matrix F(k, x) is given by

$$F(k,x) = \Theta(k+1,x)\frac{\partial f}{\partial x}(k,x)\Theta(k,x)^{-1}.$$
(7)

In case $\mathbb{X} = \mathbb{R}^n$, we say that system (1) is uniformly globally contracting.

Remark 7. While the previous definitions for convergence properties are all trajectory-based, Definition 6 is presented in a manner consistent with the literature in terms of the existence of the matrix-valued function Θ . As we will demonstrate in Theorem 15 in the sequel, contraction analysis is equivalent to uniform exponential incremental stability. Hence, an alternate definition of contraction analysis involves a simple nomenclature change in Definition 2.

Remark 8. Note that [7, Definition 3] includes (6) but not explicitly both bounds in (5). The lower bound in (5) is required to guarantee contraction as the following example illustrates. The upper bound guarantees uniformity of the contraction property with respect to initial time, which is generally desirable. Consider the system

$$x(k+1) = (k^2 + 1)x(k), \quad x(k) \in \mathbb{R}^n, k \in \mathbb{Z}.$$
(8)

Choose $\Theta(k,x) \coloneqq \frac{1}{k^2+1}$, then $F(k,x) = \frac{1}{k^2+2k+2}$ and, thus, $F(k,x)^T F(k,x) = \frac{1}{(k^2+2k+2)^2} < \frac{1}{4}$; i.e., condition (6) is satisfied. However, the origin is an equilibrium and is unstable by inspection. Thus, this system is not uniformly globally contracting. Note that for this particular choice of $\Theta(k,x)$, the lower bound of condition (5) is not satisfied.

Given (1), similar to [28], we define the variational system

$$\begin{cases} x(k+1) = f(k, x(k)), \\ x_{\delta}(k+1) = \frac{\partial f}{\partial x}(k, x(k))x_{\delta}(k). \end{cases}$$

As in [7], we refer to (9b) as the displacement dynamics where x_{δ} is called a displacement of x and is sometimes denoted by δx in the literature.

3. LYAPUNOV FUNCTION CHARACTERIZATIONS

In this section, we present time-varying Lyapunov function characterizations for the three considered convergence properties.

 $\mathbf{6}$

3.1. Incremental Stability. A Lyapunov function characterization of incremental stability for continuous-time systems was first presented in [1, Theorem 21.1 & Theorem 21.2]. Subsequently, similar results for time-invariant and time-varying systems were given in [2, Theorem 1] and [19, Theorem 5], respectively. The following theorem is a discrete-time analogue to [19, Theorem 5]. By a smooth function we mean one that is infinitely often differentiable.

Theorem 9. System (1) is uniformly globally asymptotically incrementally stable if and only if there exist a smooth function $V: \mathbb{Z} \times \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}_{\geq 0}$, functions $\alpha_1, \alpha_2 \in \mathcal{K}_{\infty}$, and $\alpha_3: \mathbb{R}_{\geq 0} \to \mathbb{R}_{\geq 0}$ positive definite, such that

$$\alpha_1(|x_1 - x_2|) \le V(k, x_1, x_2) \le \alpha_2(|x_1 - x_2|), \tag{10}$$

$$V(k+1, f(k, x_1), f(k, x_2)) - V(k, x_1, x_2) \le -\alpha_3(|x_1 - x_2|)$$
(11)

hold for all $x_1, x_2 \in \mathbb{R}^n$ and $k \in \mathbb{Z}$.

We remark that, in accordance with [34, Lemma 2.8], the function α_3 provided by the above result may be assumed to be of class \mathcal{K}_{∞} without loss of generality.

The proof of Theorem 9 is contained in the appendix. A function V satisfying (10)–(11) is called an incremental stability Lyapunov function.

3.2. Convergent Dynamics. The following theorem is a time-varying Lyapunov function characterization of discrete-time globally convergent systems. This result is analogous to a continuous-time result presented in [19, Theorem 7].

Theorem 10. Assume that system (1) is uniformly globally convergent. Then, there exist a smooth function $V \colon \mathbb{Z} \times \mathbb{R}^n \to \mathbb{R}_{\geq 0}$, a constant $c \geq 0$, and functions $\alpha_1, \alpha_2, \alpha_3 \in \mathcal{K}_{\infty}$ such that

$$\alpha_1(|x - \bar{x}(k)|) \le V(k, x) \le \alpha_2(|x - \bar{x}(k)|), \tag{12}$$

$$V(k+1, f(k, x)) - V(k, x) \le -\alpha_3(|x - \bar{x}(k)|),$$
(13)

$$V(k,0) \le c < +\infty \tag{14}$$

hold for all $x \in \mathbb{R}^n$ and $k \in \mathbb{Z}$. Conversely, if a smooth function $V : \mathbb{Z} \times \mathbb{R}^n \to \mathbb{R}_{\geq 0}$, a constant $c \geq 0$, functions $\alpha_1, \alpha_2, \in \mathcal{K}_{\infty}$, and $\alpha_3 : \mathbb{R}_{\geq 0} \to \mathbb{R}_{\geq 0}$ positive definite are given such that for some trajectory $\bar{x} : \mathbb{Z} \to \mathbb{R}^n$ estimates (12)–(14) hold, then system (1) is uniformly globally convergent and the solution $\bar{x}(k)$ is the unique bounded solution as in Definition 3. The same statement holds for uniform global exponential convergence if the \mathcal{K} functions α_1, α_2 , and α_3 are replaced by quadratic functions.

The proof of Theorem 10 is contained in the appendix. A function V satisfying (12)–(14) is called a convergent dynamics Lyapunov function.

3.3. Contraction Analysis. The following theorem is a Lyapunov function characterization for globally contracting systems. Note that this result is a special case of the Lyapunov function characterization for uniform global asymptotic incremental stability. **Theorem 11.** Assume the right hand side f of (1) is continuously differentiable in x. System (1) is globally contracting if and only if there exist a smooth function $V : \mathbb{Z} \times \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}_{\geq 0}$, and constants $c_1, c_2, c_3 \in \mathbb{R}_{>0}$ such that

$$|c_1|x_1 - x_2|^2 \le V(k, x_1, x_2) \le c_2|x_1 - x_2|^2,$$
 (15)

$$V(k+1, f(k, x_1), f(k, x_2)) - V(k, x_1, x_2) \le -c_3 |x_1 - x_2|^2$$
(16)

hold for all $x_1, x_2 \in \mathbb{R}^n$ and $k \in \mathbb{Z}$.

/

The proof of Theorem 11 is contained in the appendix. A function V satisfying (15)–(16) is called a contraction analysis Lyapunov function.

4. Comparisons

In this section, we compare the three previously discussed properties in pairs. While any asymptotically stable linear system satisfies all three properties, there are distinct differences when nonlinear systems are considered. We provide several examples of systems that highlight the essential differences among the three convergence notions. Several sufficient conditions are also proposed to establish mutual relationships. Finally, we provide a discrete-time Demidovič condition that is sufficient for all three properties.

4.1. Incremental Stability vs. Convergent Dynamics. The following example, which is adapted from [19, Example 3], provides a system that is uniformly globally convergent but not uniformly globally asymptotically incrementally stable.

Example 1. Consider the system

$$x(k+1) = f(k, x(k)), \qquad x(k) \in \mathbb{R}^2,$$
 (17)

with

$$f(k,x) \coloneqq \frac{1}{2} \begin{bmatrix} \cos(|z(k)|) & -\sin(|z(k)|) \\ \sin(|z(k)|) & \cos(|z(k)|) \end{bmatrix} z(k) + \begin{bmatrix} \cos(k+1) \\ \sin(k+1) \end{bmatrix},$$

where $\bar{x}(k) \coloneqq (\cos(k), \sin(k))^T$ and $z(k) \coloneqq x(k) - \bar{x}(k)$. Clearly, $\bar{x}(k)$ is a bounded solution of (17). Now, consider the time-varying, quadratic Lyapunov function $V(k, x) = |z(k)|^2$. We compute

$$V(k+1, f(x)) - V(k, x) = |f(k, x) - \bar{x}(k+1)|^2 - |z(k)|^2$$

= $\frac{1}{4}|z(k)|^2 - |z(k)|^2 = -\frac{3}{4}|z(k)|^2 \le 0.$

Hence, appealing to Theorem 10, system (17) is uniformly globally convergent. Rewriting z(k) in polar coordinates $(r(k), \theta(k))$ yields

$$r(k+1) = \frac{r(k)}{2},$$
 $\theta(k+1) = \theta(k) + r(k),$

whose explicit solution for an initial condition $(r_0, \theta_0) \in \mathbb{R} \times \mathbb{R}$ is

$$r(k) = \frac{r_0}{2^{k-k_0}}, \qquad \qquad \theta(k) = \theta_0 + r_0 \sum_{\kappa=0}^{k-k_0-1} \frac{1}{2^{\kappa}}.$$



FIGURE 1. The two trajectories of the system defined in Example 1 start on the positive real half line with an initial separation of π at time k = 0 and the initial distances to the origin are M + 1 and $M + 1 + \pi$. At time k = 1, the polar arguments of the trajectories are shifted by 180° so that the separation distance is $M + \frac{\pi}{2}$.

At time $k_0 = 0$ consider two initial conditions in Cartesian coordinates $\xi_1 = (M + 1, 0)$ and $\xi_2 = (M + \pi + 1, 0)$ for some $M \in \mathbb{R}_{>0}$ as shown in Figure 1. The polar coordinates of z(0) corresponding to ξ_1 and ξ_2 are $(r_1^0, \theta_1^0) = (M, 0)$ and $(r_2^0, \theta_2^0) = (M + \pi, 0)$. The initial separation is $|\xi_1 - \xi_2| = \pi$. At the time instance k = 1, the polar coordinates of z(1) are $(r_1(1), \theta_1(1)) = (\frac{M}{2}, M)$ and $(r_2(1), \theta_2(1)) = (\frac{M+\pi}{2}, M+\pi)$, hence, the angle difference is π . Consequently, the distance between two states is $|\phi(1; 0, \xi_1) - \phi(1; 0, \xi_2)| = r_1(1) + r_2(1) = \frac{M}{2} + \frac{M+\pi}{2} = M + \frac{\pi}{2}$. Since $M \in \mathbb{R}_{>0}$ is arbitrary, for any function $\beta \in \mathcal{KL}$, we can always choose a sufficiently large M such that (2) is violated. Thus, system (17) is not incrementally stable.

The above example exploits the fact that global asymptotic convergence of any two trajectories to each other (incremental stability) is a stronger property than global asymptotic convergence of all trajectories to a single trajectory (convergent dynamics). This is a direct consequence of the triangle inequality.

The following theorem provides a connection from convergent dynamics to incremental stability by restricting the state space of system (1) to be a compact and positively invariant set.

Theorem 12. Suppose system (1) is uniformly convergent on a compact and positively invariant set $\mathbb{X} \subset \mathbb{R}^n$. Further assume that the right hand side f of system (1) is locally Lipschitz continuous in x on \mathbb{X} . Then, system (1) is uniformly asymptotically incrementally stable on \mathbb{X} . Moreover, if the attraction rate in the uniform convergence property is exponential, then the system is uniformly exponentially incrementally stable.

The proof of Theorem 12 is contained in the appendix. Conversely, incremental stability does not imply convergent dynamics as shown by the following example adapted from [19, Example 4].

Example 2. The system

$$x(k+1) = -\frac{k}{2} - 1 + \frac{x(k)}{2}, \text{ with } x(k_0) = \xi \in \mathbb{R}$$
(18)

has the explicit solution

$$\phi(k;k_0,\xi) = \frac{k_0}{2^{k-k_0}} - k + \frac{\xi}{2^{k-k_0}}, \quad \forall k \ge k_0.$$
(19)

For any two initial conditions $\xi_1, \xi_2 \in \mathbb{R}$,

$$|\phi(k;k_0,\xi_1) - \phi(k;k_0,\xi_2)| = \frac{|\xi_1 - \xi_2|}{2^{k-k_0}} = \beta(|\xi_1 - \xi_2|, k - k_0),$$

where $\beta \in \mathcal{KL}$ is defined by $\beta(s,r) = \frac{s}{2^r}$ for all $s,r \in \mathbb{R}_{\geq 0}$. Hence, system (18) is uniformly globally asymptotically incrementally stable. However, for the specific initial condition $(\xi, k_0) = (0,0)$, it is straightforward that the solution (19) from this initial condition is unbounded. Therefore, system (18) is not uniformly globally convergent. \Box

The above example exploits the fact that convergent dynamics requires the existence of at least one bounded solution. Whereas solutions of an incrementally stable system can be unbounded.

The following theorem provides a connection from incremental stability to convergent dynamics with an assumption that the state space of system (1) is compact and positively invariant.

Theorem 13. Suppose system (1) is uniformly globally asymptotically incrementally stable and there exists a compact set $\mathbb{X} \subset \mathbb{R}^n$ that is positively invariant under (1). Then system (1) is uniformly globally convergent. Moreover, if the incremental attraction rate is exponential, then it is also exponential for the convergence property.

The proof of Theorem 13 is contained in the appendix.

The following theorem provides a sufficient condition for system (1) to satisfy all three convergence properties. It is the discrete-time analogue of the Demidovič result [6, Theorem 1].

Theorem 14 (Discrete-time Demidovič condition). Assume the right hand side f of system (1) is continuously differentiable in x. Suppose there exists a positive definite matrix P such that the matrix

$$J(k,x) \coloneqq \frac{\partial f}{\partial x}(k,x)^T P \frac{\partial f}{\partial x}(k,x) - \rho P$$
(20)

is negative semidefinite uniformly in $(k, x) \in \mathbb{Z} \times \mathbb{R}^n$ for some $\rho \in (0, 1)$. Then system (1) is uniformly globally exponentially incrementally stable and globally contracting. Furthermore, if there exists $c \geq 0$ such that

$$\sup_{k\in\mathbb{Z}} |f(k,0)| = c < \infty,$$
(21)

then system (1) is uniformly globally exponentially convergent.

The proof of Theorem 14 is contained in the appendix.

4.2. Contraction Analysis vs. Incremental Stability. Assuming the right hand side of system (1) is continuously differentiable, we demonstrate that contraction analysis is a strictly stronger property than asymptotic incremental stability. In fact, contraction analysis is equivalent to exponential incremental stability.

Theorem 15. Suppose the right hand side of system (1) is continuously differentiable in x on \mathbb{R}^n . System (1) is uniformly globally exponentially incrementally stable if and only if it is uniformly globally contracting.

The proof of Theorem 15 is contained in the appendix.

4.3. Convergent Dynamics vs. Contraction Analysis. System (17) in Example 1 is globally convergent, however, it is not globally contracting because it is not differentiable at x = 0. Hence, a uniformly globally convergent system is not necessarily globally contracting.

However, even in the case where the system dynamics are continuously differentiable, global convergent dynamics is still different from contraction analysis. This is due to the fact that contraction analysis requires an exponential convergence rate whereas convergent dynamics only requires asymptotic convergence. The following example, adapted from [37, Example 1], provides a (time-invariant) system that is asymptotically convergent but not exponentially convergent.

Example 3. The system

$$x(k+1) = f(x(k)) \coloneqq \frac{x(k)}{\sqrt{x(k)^2 + 1}},$$
(22)

with $x(k_0) = \xi \in \mathbb{R}$ has the solution

$$\phi(k;k_0,\xi) = \frac{\xi}{\sqrt{(k-k_0)\xi^2 + 1}}, \quad \forall k \in \mathbb{Z}, k \ge k_0.$$
(23)

The zero solution $\bar{x}(k) = 0$ is (uniformly) globally asymptotically stable. Indeed, for the Lyapunov function $V(x) = x^2$, we have, for all $x \neq 0$,

$$V(f(x)) - V(x) = -\frac{x^4}{x^2 + 1} < 0.$$
(24)

Hence, system (22) is (uniformly) globally convergent. By inspection of (23), we see that the convergence to the zero solution is not exponential, hence, by Theorem 15, system (22) is not globally contracting. \Box

Example 4. Returning to Example 2, system (18) is not convergent since the solution passing through $(k_0,\xi) = (0,0)$ is unbounded. However, system (18) is globally contracting. Indeed, take $\Theta(k,x) = 1$ for all $k \in \mathbb{Z}$ and $x \in \mathbb{R}, \mu = \frac{3}{4}$, and note that $\frac{\partial f}{\partial x}(k,x) = \frac{1}{2}$ for all $k \in \mathbb{Z}$ and $x \in \mathbb{R}^n$. With F(k,x) given by (7) we see that

$$F(k,x)^T F(k,x) - 1 = -\mu.$$

Hence, system (18) is globally contracting.

Using the results of Theorem 13 and Theorem 15, we provide sufficient conditions under which global contraction implies convergent dynamics.



FIGURE 2. Relationships between different convergence properties assuming continuously differential dynamics f. Note the abbreviations: IS for (global asymptotic) *Incremental Stability*, EIS for (global) *Exponential Incremental Stability*, CD for (uniform global) *Convergent Dynamics*, ECD for (uniform global) *Exponential Convergent Dynamics*, and CA for (global) *Contraction Analysis*. The blue, dashed implications do not hold in general, but for the case of a compact (and invariant) state space X.

Theorem 16. Suppose the mapping f of system (1) is continuously differentiable in x on a compact and positively invariant set $\mathbb{X} \subset \mathbb{R}^n$. If system (1) is uniformly contracting in \mathbb{X} , then system (1) is uniformly exponentially convergent in \mathbb{X} .

The proof of Theorem 16 is contained in the appendix.

5. Conclusions

This paper contributes discrete-time, time-varying, smooth Lyapunov function characterizations for incremental stability, convergent dynamics, and contraction analysis. The paper also contributes examples of systems that highlight the essential differences as well as similarities among the three considered notions of stability. Moreover, with appropriate assumptions, we present several conditions that provide connections between each of the three considered convergence properties. Overall, assuming the right hand side of the considered system is continuously differentiable, the relationships between the three convergence properties can be summarized as in Figure 2.

References

- T. Yoshizawa, Stability Theory by Liapunov's Second Method. Mathematical Society of Japan, 1966.
- [2] D. Angeli, "A Lyapunov approach to incremental stability properties," *IEEE Trans. Autom. Control*, vol. 47, no. 3, pp. 410–421, March 2002.
- [3] M. Zamani and P. Tabuada, "Backstepping design for incremental stability," *IEEE Trans. Autom. Control*, vol. 56, no. 9, pp. 2184–2189, Sept 2011.
- [4] B. P. Demidovič, Lektsii po matematicheskoi teorii ustoichivosti. (Russian) [Lectures on the mathematical theory of stability]. Moscow: Nauka, 1967.

- [5] A. Pavlov, N. van de Wouw, and H. Nijmeijer, Uniform Output Regulation of Nonlinear Systems: A Convergent Dynamics Approach. Boston: Birkhäuser, 2006.
- [6] A. Pavlov, A. Pogromsky, N. van de Wouw, and H. Nijmeijer, "Convergent dynamics, a tribute to Boris Pavlovich Demidovich," *Systems Control Lett.*, vol. 52, no. 34, pp. 257 – 261, 2004.
- [7] W. Lohmiller and J.-J. E. Slotine, "On contraction analysis for non-linear systems," Automatica J. IFAC, vol. 34, no. 6, pp. 683–696, 1998.
- [8] Q.-C. Pham and J.-J. E. Slotine, "Stable concurrent synchronization in dynamic system networks," *Neural Netw.*, vol. 20, no. 1, pp. 62 – 77, 2007.
- [9] C. Cai and G. Chen, "Synchronization of complex dynamical networks by the incremental ISS approach," *Phys. A*, vol. 371, no. 2, pp. 754 – 766, 2006.
- [10] G. B. Stan and R. Sepulchre, "Analysis of interconnected oscillators by dissipativity theory," *IEEE Trans. Autom. Control*, vol. 52, no. 2, pp. 256–270, Feb 2007.
- [11] A. Pavlov and N. van de Wouw, "Steady-state analysis and regulation of discretetime nonlinear systems," *IEEE Trans. Autom. Control*, vol. 57, no. 7, pp. 1793–1798, July 2012.
- [12] A. Pavlov, N. van de Wouw, and H. Nijmeijer, "Frequency response functions for nonlinear convergent systems," *IEEE Trans. Autom. Control*, vol. 52, no. 6, pp. 1159–1165, June 2007.
- [13] A. Pavlov and N. van de Wouw, "Convergent discrete-time nonlinear systems: The case of PWA systems," in *Proc. IEEE American Contr. Conf.*, June 2008, pp. 3452– 3457.
- [14] E. Trefftz, "Zu den Grundlagen der Schwingungstheorie," Math. Ann., vol. 95, no. 1, pp. 307–312, 1926.
- [15] R. Reissig, "Zur Theorie der erzwungenen Schwingungen," Math. Nachr., vol. 13, pp. 309–312, 1955.
- [16] —, "Über eine nichtlineare Differentialgleichung 2. Ordnung," Math. Nachr., vol. 13, pp. 313–318, 1955.
- [17] —, "Erzwungene Schwingungen mit zäher und trockener Reibung (Ergänzung)," Math. Nachr., vol. 12, pp. 249–252, 1954.
- [18] J. P. LaSalle, "A study of synchronous asymptotic stability," Ann. of Math. (2), vol. 65, no. 3, pp. 571–581, 1957.
- [19] B. S. Rüffer, N. van de Wouw, and M. Mueller, "Convergent systems vs. incremental stability," Systems Control Lett., vol. 62, no. 3, pp. 277–285, 2013.
- [20] V. Fromion, S. Monaco, and D. Normand-Cyrot, "Asymptotic properties of incrementally stable systems," *IEEE Trans. Autom. Control*, vol. 41, no. 5, pp. 721–723, 1996.
- [21] V. Fromion and G. Scorletti, "The behavior of incrementally stable discrete time systems," Systems Control Lett., vol. 46, no. 4, pp. 289–301, 2002.
- [22] E. D. Sontag, "Smooth stabilization implies coprime factorization," *IEEE Trans. Autom. Control*, vol. 34, no. 4, pp. 435–443, April 1989.
- [23] V. A. Yakubovich, "The matrix-inequality method in the theory of the stability of nonlinear control systems. I: The absolute stability of forced vibrations." Autom. Rem. Control, vol. 25, pp. 905–917, 1965.
- [24] B. P. Demidovič, "On the dissipativity of a certain non-linear system of differential equations. I," Vestnik Moskov. Univ. Ser. I Mat. Meh., vol. 1961, no. 6, pp. 19–27, 1961.
- [25] —, "On the dissipative character of a certain non-linear system of differential equations. II," Vestnik Moskov. Univ. Ser. I Mat. Meh., vol. 1962, no. 1, pp. 3–8, 1962.
- [26] V. A. Pliss, Nonlocal problems of the theory of oscillations. Academic Press, 1966, translated from Russian.
- [27] Y. Lan and I. Mezič, "Linearization in the large of nonlinear systems and Koopman operator spectrum," *Phys. D*, vol. 242, no. 1, pp. 42–53, 2013.
- [28] F. Forni and R. Sepulchre, "A differential Lyapunov framework for contraction analysis," *IEEE Trans. Autom. Control*, vol. 59, no. 3, pp. 614–628, March 2014.
- [29] J. W. Simpson-Porco and F. Bullo, "Contraction theory on Riemannian manifolds," Systems Control Lett., vol. 65, pp. 74 – 80, 2014.

- [30] P. Giesl, "Converse theorems on contraction metrics for an equilibrium," J. Math. Anal. Appl., vol. 424, no. 2, pp. 1380–1403, 2015.
- [31] J. Jouffroy, "Some ancestors of contraction analysis," in Proc. Joint 44th IEEE Conf. Decis. Control and Europ. Contr. Conf., Dec 2005, pp. 5450–5455.
- [32] E. M. Aylward, P. A. Parrilo, and J.-J. E. Slotine, "Stability and robustness analysis of nonlinear systems via contraction metrics and SOS programming," *Automatica J. IFAC*, vol. 44, no. 8, pp. 2163 – 2170, 2008.
- [33] V. Fromion and G. Scorletti, "Connecting nonlinear incremental Lyapunov stability with the linearizations Lyapunov stability," in *Proc. Joint 44th IEEE Conf. Decis. Control and Europ. Contr. Conf.*, Dec 2005, pp. 4736–4741.
- [34] Z.-P. Jiang and Y. Wang, "A converse Lyapunov theorem for discrete-time systems with disturbances," Systems Control Lett., vol. 45, no. 1, pp. 49–58, 2002.
- [35] J. L. Massera, "Contributions to stability theory," Ann. of Math.(2), vol. 64, pp. 182–206, 1956, (Erratum: Ann. of Math.(2), 68:202, 1958.).
- [36] C. M. Kellett, "A compendium of comparison function results," Mathematics of Controls, Signals and Systems, vol. 26, no. 3, pp. 339–374, 2014.
- [37] D. N. Tran, C. M. Kellett, and P. M. Dower, "Qualitative equivalences of ISS and l_p-gain stability properties for discrete-time nonlinear systems," *Automatica J. IFAC*, vol. 77, pp. 360–369, 2017.
- [38] E. D. Sontag, "Comments on Integral Variants of ISS," Systems Control Lett., vol. 34, no. 1–2, pp. 93–100, 1998.
- [39] W. J. Rugh, Linear System Theory, 2nd ed. Pearson, 1995.

APPENDIX

As several proofs in this appendix rely on the same modification of a converse Lyapunov result originally due to Jiang and Wang [34], we state the modification here explicitly for the convenience of the reader.

Definition 17 (cf. [34]). Let \mathcal{A} be a closed, not necessarily compact, positively invariant set of (1). System (1) is uniformly global asymptotically stable (UGAS) with respect to \mathcal{A} if there exists a class- \mathcal{KL} function β such that

$$|\phi(k;k_0,\xi_0)|_{\mathcal{A}} \le \beta(|\xi_0|_{\mathcal{A}},k-k_0) \tag{25}$$

for all $k \geq k_0$ and all $\xi_0 \in \mathbb{R}^n$, where

$$|\xi|_{\mathcal{A}} \coloneqq \inf_{\eta \in \mathbb{R}^n} |\eta - \xi|$$

denotes the distance of ξ to \mathcal{A} .

Similarly, the notion of uniform global *exponential* stability (UGES) is defined by replacing estimate (25) with the exponential bound

$$|\phi(k;k_0,\xi_0)|_{\mathcal{A}} \le \kappa |\xi_0|_{\mathcal{A}} \lambda^{-(k-k_0)},\tag{26}$$

for some constants $\kappa \geq 1$ and $\lambda > 1$.

Theorem 18 (Jiang & Wang 2002). Assume system (1) is UGAS with respect to the closed set \mathcal{A} . Then there exist a smooth function $V : \mathbb{Z} \times \mathbb{R}^n \to \mathbb{R}_{\geq 0}$ and functions $\alpha_1, \alpha_2, \alpha_3 \in \mathcal{K}_{\infty}$, such that for all $k \in \mathbb{Z}$ and all $x \in \mathbb{R}^n$,

$$\alpha_1(|x|_{\mathcal{A}}) \le V(k, x) \le \alpha_2(|x|_{\mathcal{A}}) \tag{27}$$

and

$$V(k+1, f(k, x)) - V(k, x) \le -\alpha_3(|x|_{\mathcal{A}}).$$
 (28)

Conversely, if there exist a continuous function $V : \mathbb{Z} \times \mathbb{R}^n \to \mathbb{R}_{\geq 0}$, functions $\alpha_1, \alpha_2 \in \mathcal{K}_{\infty}$, and a positive definite function $\alpha_3 : \mathbb{R}_{\geq 0} \to \mathbb{R}_{\geq 0}$, such that (27)–(28) hold, then system (1) is UGAS with respect to the closed set \mathcal{A} .

The original result in [34, Theorem 1] asserts the existence of a function V defined on $\mathbb{Z}_{\geq 0}$ satisfying (27) and, for a positive definite function α_3 , (28). In contrast, the formulation above yields a function V defined on all of \mathbb{Z} . That the function α_3 can indeed be chosen to be of class- \mathcal{K}_{∞} instead of merely positive definite is already proven in [34, Lemma 2.8].

It is thus no surprise that the proof of the above result follows the original very closely.

Proof. Start by assuming that system (1) is UGAS with respect to the closed set \mathcal{A} . Applying Sontag's Lemma on \mathcal{KL} -estimates [38, Proposition 7], given $\beta \in \mathcal{KL}$, there exist $\rho_1, \rho^{-1} \in \mathcal{K}_{\infty}$ such that

$$\beta(|\xi_0|_{\mathcal{A}}, k - k_0) \le \rho^{-1}(\rho_1(|\xi_0|_{\mathcal{A}})e^{-(k-k_0)})$$
(29)

for all $\xi_0 \in \mathbb{R}^n$ and $k, k_0 \in \mathbb{Z}$ such that $k \ge k_0$. Hence,

$$\rho(|\phi(k;k_0,\xi_0)|_{\mathcal{A}}) \le \rho_1(|\xi_0|_{\mathcal{A}})e^{-(k-k_0)}.$$
(30)

Define $V: \mathbb{Z} \times \mathbb{R}^n \to \mathbb{R}_{\geq 0}$ by

$$V(k_0,\xi_0) \coloneqq \sum_{k=k_0}^{\infty} \rho(|\phi(k;k_0,\xi_0)|_{\mathcal{A}}).$$
 (31)

It is true that for all $k_0 \in \mathbb{Z}$ and $\xi_0 \in \mathbb{R}^n$

$$\rho(|\xi_0|_{\mathcal{A}}) = \rho(|\phi(k_0; k_0, \xi_0)|_{\mathcal{A}}) \le V(k_0, \xi_0)$$
$$\le \sum_{k=k_0}^{\infty} \rho_1(|\xi_0|_{\mathcal{A}})e^{-(k-k_0)} \le \frac{e}{e-1}\rho_1(|\xi_0|)$$
(32)

where the second inequality follows from (30) and (31). This means that the series defined in (31) is convergent. Moreover, for each $k_0 \in \mathbb{Z}$, ρ is a continuous function in ξ_0 , thus, V is continuous in ξ_0 .

Define $\alpha_1 := \rho$ and $\alpha_2(s) := \frac{e}{e-1}\rho_1(s)$ for all $s \in \mathbb{R}_{\geq 0}$. Observe that α_1 and α_2 are both of class- \mathcal{K}_{∞} . We immediately see that V satisfies condition (27).

It is only left to prove that V satisfies condition (28). Indeed, for arbitrary $k_0 \in \mathbb{Z}$ and $\xi_0 \in \mathbb{R}^n$ we have

$$V(k_0 + 1, f(k_0, \xi_0)) = \sum_{k=k_0+1}^{\infty} \rho(|\phi(k; k_0 + 1, f(k_0, \xi_0))|_{\mathcal{A}})$$

=
$$\sum_{k=k_0+1}^{\infty} \rho(|\phi(k; k_0 + 1, f(k_0, \xi_0))|_{\mathcal{A}})$$

+
$$\rho(|\phi(k_0; k_0, \xi_0)|_{\mathcal{A}}) - \rho(|\phi(k_0; k_0, \xi_0)|_{\mathcal{A}})$$

=
$$\sum_{k=k_0}^{\infty} \rho(|\phi(k; k_0, \xi_0)|) - \rho(|\phi(k_0; k_0, \xi_0)|_{\mathcal{A}})$$

=
$$V(k_0, \xi_0) - \rho(|\xi_0|_{\mathcal{A}}).$$

We conclude that the continuous function V satisfies (28) with $\alpha_3 \coloneqq \rho$. Lastly, an application of [34, Lemma 2.7] allows us to replace the continuous function V by a smooth one (which also modifies the α_i , i = 1, 2, 3).

The converse direction is standard and the same as in [34, Section 4.3.3] and thus omitted. $\hfill\square$

Corollary 19. If system (1) is UGES with respect to the closed set \mathcal{A} , then there exist a smooth function $V: \mathbb{Z} \times \mathbb{R}^n \to \mathbb{R}_{\geq 0}$ and constants $c_1, c_2, c_3 > 0$, such that for all $k \in \mathbb{Z}$ and all $x \in \mathbb{R}^n$,

$$c_1 |x|^2_{\mathcal{A}} \le V(k, x) \le c_2 |x|^2_{\mathcal{A}}$$
 (33)

and

$$V(k+1, f(k, x)) - V(k, x) \le -c_3 |x|_{\mathcal{A}}^2.$$
(34)

Conversely, the existence of a continuous function $V: \mathbb{Z} \times \mathbb{R}^n \to \mathbb{R}_{\geq 0}$ and positive constants c_1, c_2, c_3 satisfying (33)–(34) is sufficient for system (1) to be UGES with respect to the closed set \mathcal{A} .

Proof. This result follows as in the previous proof by observing that in the additional step of using Sontag's Lemma on \mathcal{KL} -estimates is not necessary due to the exponential estimate (26), which is equivalent to

$$|\phi(k;k_0,\xi_0)|_{\mathcal{A}}^2 \le \kappa^2 |\xi_0|_{\mathcal{A}}^2 (\lambda^2)^{-(k-k_0)}.$$

With these two results at hand, we can now turn to proving the main results in this paper.

Proof of Theorem 9. Consider the augmented system

$$\begin{cases} x_1(k+1) = f(k, x_1(k)) \\ x_2(k+1) = f(k, x_2(k)) \end{cases}$$
(35)

as in [2]. The diagonal is the set $\Delta := \{ [x^T, x^T]^T \in \mathbb{R}^{2n} : x \in \mathbb{R}^n \}$. Let $z = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \in \mathbb{R}^{2n}, x_1, x_2 \in \mathbb{R}^n$, then the distance from z to the diagonal is given by

$$|z|_{\Delta} \coloneqq \inf_{w \in \Delta} |w - z| = \frac{1}{\sqrt{2}} |x_1 - x_2|,$$
 (36)

where the equality is shown in [2]. Denote $F\left(k, \begin{bmatrix} x_1\\ x_2 \end{bmatrix}\right) \coloneqq \begin{bmatrix} f(k,x_1(k))\\ f(k,x_2(k)) \end{bmatrix}$ and rewrite (35) as

$$z(k+1) = F(k, z(k)),$$
(37)

where $z = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \in \mathbb{R}^{2n}$ and $k \in \mathbb{Z}$.

Appealing to the relationship (36), we see that system (1) is uniformly globally asymptotically incrementally stable if and only if system (37) is uniformly globally asymptotically stable with respect to the diagonal set Δ .

By application of Theorem 18, system (37) admits a smooth, time-varying Lyapunov function $V: \mathbb{Z} \times \mathbb{R}^{2n} \to \mathbb{R}_{\geq 0}$ for which there exist functions $\alpha_1, \alpha_2, \alpha_3 \in \mathcal{K}_{\infty}$ such that

$$\alpha_1(|z|_{\Delta}) \le V(k, z) \le \alpha_2(|z|_{\Delta}),\tag{38}$$

$$V(k, F(k, z)) - V(k, z) \le -\alpha_3(|z|_{\Delta})$$
(39)

hold for all $z \in \mathbb{R}^{2n}$ and $k \in \mathbb{Z}$. Using (36), it is clear that (38)–(39) are correspondingly equivalent to (10)–(11).

Again using Theorem 18, we conclude that system (1) is uniformly globally asymptotically incrementally stable if and only if system (1) admits a Lyapunov function satisfying (10)–(11).

Proof of Theorem 10. Assume that system (1) is uniformly globally convergent as in Definition 3, so that there exists a solution $\bar{x}(k)$ bounded for all $k \in \mathbb{Z}$; i.e., $\sup_{k \in \mathbb{Z}} |\bar{x}(k)| < \infty$.

After a change of coordinates $z(k) = x(k) - \bar{x}(k)$ to the original system, we obtain a new system

$$z(k+1) = f(x(k)) - f(\bar{x}(k)) = f(z(k) + \bar{x}(k)) - f(\bar{x}(k)) \rightleftharpoons g(k, z(k)).$$

Applying Theorem 18 to this new system yields a smooth function $W: \mathbb{Z} \times \mathbb{R}^n \to \mathbb{R}_{>0}$ and functions $\alpha_1, \alpha_2, \alpha_3 \in \mathcal{K}_{\infty}$ such that

$$\alpha_1(|z|) \le W(k, z) \le \alpha_2(|z|), \text{ and}$$

$$\tag{40}$$

$$W(k+1, g(k, z)) - W(k, z) \le -\alpha_3(|z|), \tag{41}$$

hold for all $z \in \mathbb{R}^n$ and $k \in \mathbb{Z}$. Reverting to the original coordinates and defining $V(k,x) := W(k, x - \bar{x}(k))$, we obtain (12)–(13) from (40)–(41). Moreover,

$$0 \le V(k,0) \le \alpha_2(|\bar{x}(k)|) \le \alpha_2\left(\sup_{k \in \mathbb{Z}} |\bar{x}(k)|\right) \eqqcolon c < \infty$$

establishes (14). This completes the proof of the first statement of the theorem.

Conversely, if (12)–(14) hold, then with the same coordinate change and by applying the forward result in Theorem 18, we see that system (1) is uniformly globally asymptotically stable with respect to the trajectory $\bar{x}(k)$. Thus, it is only left to show that $\bar{x}(k)$ is bounded and unique. Indeed,

$$|\bar{x}(k)| \le \alpha_1^{-1}(V(k,0)) \le \alpha_1^{-1}(c) < \infty$$

shows that $\bar{x}(k)$ is bounded. Lastly, uniqueness of $\bar{x}(k)$ follows from Remark 5.

Replacing every invocation of Theorem 18 in this proof by an invocation of Corollary 19 proves the statement about uniform global exponential convergence.

Proof of Theorem 11. The proof proceeds by demonstrating the equivalence of an exponential incremental Lyapunov function and exponential incremental stability and then appealing to Theorem 15.

Similar to the proof of Theorem 9, consider the augmented system (35) (also, its shorthand notation (37)). Since (36) holds, system (1) is uniformly globally exponentially incrementally stable if and only if the diagonal set Δ is uniformly globally exponentially stable with respect to system (37).

Applying Corollary 19, the closed set Δ is uniformly globally exponentially stable with respect to system (37) if and only if system (37) admits a smooth, time-varying Lyapunov function $V : \mathbb{Z} \times \mathbb{R}^{2n} \to \mathbb{R}_{\geq 0}$ for which there exist constants $c_1, c_2, c_3 \in \mathbb{R}_{>0}$ such that

$$c_1 |z|_{\Delta}^2 \le V(k, z) \le c_2 |z|_{\Delta}^2,$$
(42)

$$V(k, F(k, z)) - V(k, z) \le -c_3 |z|_{\Delta}^2$$
 (43)

hold for all $z \in \mathbb{R}^{2n}$ and $k \in \mathbb{Z}$. Using (36), it is clear that (42)–(43) are correspondingly equivalent to (15)–(16). Therefore, we conclude that system (1) admits an (exponential incremental) Lyapunov function (15)–(16) if and only if system (1) is uniformly globally exponentially incrementally stable.

Applying Theorem 15 yields that system (1) is uniformly globally exponentially incrementally stable if and only if system (1) is globally contracting. Note that the proof of Theorem 15 does not employ any results prior to Theorem 11 or Theorem 11 itself. This completes the proof.

Proof of Theorem 12. The proof follows that of [19, Theorem 8]. Denote the diameter of \mathbb{X} as $d_{\mathbb{X}} := \max_{x,y \in \mathbb{X}} |x - y|$. Fix any $\xi_1, \xi_2 \in \mathbb{X}$. Applying the triangle inequality, we see that

$$\begin{aligned} |\phi(k;k_0,\xi_1) - \phi(k;k_0,\xi_2)| \\ &\leq |\phi(k;k_0,\xi_1) - \bar{x}(k)| + |\phi(k;k_0,\xi_1) - \bar{x}(k)| \\ &\leq 2\beta(d_{\mathbb{X}},k-k_0) \end{aligned}$$
(44)

for all $x \in \mathbb{X}$ and $k, k_0 \in \mathbb{Z}$ such that $k \ge k_0$. Let $L \in \mathbb{R}_{>0}$ be the maximum Lipschitz constant for f on \mathbb{X} ; i.e.,

$$|f(k_0,\xi_1) - f(k_0,\xi_2)| \le L|\xi_1 - \xi_2| \tag{45}$$

for all $\xi_1, \xi_2 \in \mathbb{X}$. It immediately follows from (45) that

$$|\phi(k;k_0,\xi_1) - \phi(k;k_0,\xi_2)| \le L^{k-k_0} |\xi_1 - \xi_2|.$$
(46)

Combining (44) and (45), we see that

$$|\phi(k;k_0,\xi_1) - \phi(k;k_0,\xi_2)| \le \min\left\{L^{k-k_0}|\xi_1 - \xi_2|, 2\beta(d_{\mathbb{X}},k-k_0)\right\}.$$

From there, we can obtain a function $\hat{\beta} \in \mathcal{KL}$ such that

$$|\phi(k;k_0,\xi_1) - \phi(k;k_0,\xi_2)| \le \hat{\beta}(|\xi_1 - \xi_2|, k - k_0)$$

for all $\xi_1, \xi_2 \in \mathbb{X}$ and $k, k_0 \in \mathbb{Z}$ such that $k \geq k_0$. Therefore, system (1) is uniformly asymptotically incrementally stable on \mathbb{X} .

With only minor rewording of the proof we establish the assertion about exponential incremental stability.

Proof of Theorem 13. Suppose system (1) is uniformly globally asymptotically incrementally stable, then by applying Theorem 9, there exists a function V satisfying (10)–(11).

To derive the existence of a bounded solution $\bar{x}(k)$, we use a discrete-time version from [13, Lemma 2] of Yakubovich's Lemma [23, Lemma 2] which states if there exists a compact and positively invariant set X of system (1), then there exists a solution $\bar{x}(k)$ of system (1) defined on Z and satisfying $\bar{x}(k) \in X$ for all $k \in \mathbb{Z}$. Thus, we see that

$$\alpha_1(|x - \bar{x}(k)|) \le V(k, x, \bar{x}(k)) \le \alpha_2(|x - \bar{x}(k)|)$$

and

$$V(k+1, f(k, x), f(k, \bar{x}(k))) - V(k, x, \bar{x}(k)) \le -\alpha_3(|x - \bar{x}(k)|)$$

for all $x \in \mathbb{R}^n$ and $k \in \mathbb{Z}$. Moreover, since $\bar{x}(k)$ is bounded, we have

$$V(k, 0, \bar{x}(k)) \le \alpha_2(|0 - \bar{x}(k)|) < \infty$$
 (47)

for all $k \in \mathbb{Z}$.

Hence, the function $W(k, x) := V(k, x, \bar{x}(k))$ satisfies conditions (12)–(14). Thus, by applying Theorem 10, we conclude that system (1) is uniformly globally convergent.

Proof of Theorem 14. Consider any two initial conditions $\xi_1, \xi_2 \in \mathbb{R}^n$. Define a function $\Phi: [0,1] \to \mathbb{R}$ by

$$\Phi(s) \coloneqq \left(f(k,\xi_1) - f(k,\xi_2)\right)^T Pf(k,s\xi_1 + (1-s)\xi_2).$$
(48)

Then, Φ satisfies

$$\Phi(1) - \Phi(0) = (f(k,\xi_1) - f(k,\xi_2))^T P(f(k,\xi_1) - f(k,\xi_2)).$$
(49)

Applying the mean value theorem, there exists an $\bar{s} \in [0, 1]$ such that

$$\Phi(1) - \Phi(0) = \frac{d}{ds} \Phi(\bar{s}) = \left(f(k,\xi_1) - f(k,\xi_2) \right)^T P \frac{\partial f}{\partial x} \left(k, \bar{s}\xi_1 + (1-\bar{s})\xi_2 \right) \left(\xi_1 - \xi_2 \right).$$
(50)

Denote $\check{\xi} \coloneqq \bar{s}\xi_1 + (1-\bar{s})\xi_2$. From (49), (50), we obtain

$$(f(k,\xi_1) - f(k,\xi_2))^T P(f(k,\xi_1) - f(k,\xi_2)) = (f(k,\xi_1) - f(k,\xi_2))^T P \frac{\partial f}{\partial x}(k,\check{\xi})(\xi_1 - \xi_2)$$
(51)

We make the following claim:

Claim 1. For all $\xi_1, \xi_2 \in \mathbb{R}^n$ and $k \in \mathbb{Z}$, the following holds

$$(f(k,\xi_1) - f(k,\xi_2))^T P(f(k,\xi_1) - f(k,\xi_2))$$

$$\leq (\xi_1 - \xi_2)^T \left(\frac{\partial f}{\partial x}(k,\check{\xi})^T P \frac{\partial f}{\partial x}(k,\check{\xi})\right) (\xi_1 - \xi_2).$$
(52)

Proof of Claim 1: To simplify the notations, we denote $a = (f(k,\xi_1) - f(k,\xi_2))$ and $b = \left(\frac{\partial f}{\partial x}(k,\xi)\right)(\xi_1 - \xi_2)$, we want to show $a^T P a \leq b^T P b$. From (51), we have $a^T P a = a^T P b$. Thus,

$$b^{T}Pb - a^{T}Pa = b^{T}Pb - a^{T}Pa + 2(a^{T}Pa - a^{T}Pb)$$

= $(b^{T}Pb - b^{T}Pa) + (a^{T}Pa - a^{T}Pb)$
= $(a - b)^{T}P(a - b) \ge 0$ (53)

completes the proof of Claim 1.

As $J(k, x) \leq 0$, we see that

$$(\xi_1 - \xi_2)^T \left(\frac{\partial f}{\partial x}(k, \check{\xi})^T P \frac{\partial f}{\partial x}(k, \check{\xi})\right) (\xi_1 - \xi_2) \le \rho(\xi_1 - \xi_2)^T P(\xi_1 - \xi_2)$$

which, with Claim 1, implies that

$$(f(k,\xi_1) - f(k,\xi_2))^T P(f(k,\xi_1) - f(k,\xi_2)) \le \rho(\xi_1 - \xi_2)^T P(\xi_1 - \xi_2).$$
 (54)

Define a Lyapunov function candidate $V(k, \xi_1, \xi_2) := (\xi_1 - \xi_2)^T P(\xi_1 - \xi_2)$. Since P is positive definite, V satisfies condition (10). Next, using (54),

$$V(k+1, f(k,\xi_1), f(k,\xi_2)) - V(k,\xi_1,\xi_2) \le -(1-\rho)V(k,\xi_1,\xi_2),$$

for all $\xi_1, \xi_2 \in \mathbb{R}^n$ and $k \in \mathbb{Z}$. This implies that V satisfies condition (11). Therefore, by virtue of Theorem 9, system (1) is uniformly globally asymptotically incrementally stable.

It is straightforward to see that the function V defined above also satisfies (15) and (16) of Theorem 11. Hence, system (1) is globally contracting.

The additional condition (21) together with (54) implies that system (1) is uniformly globally (exponentially) convergent by applying [11, Theorem 1].

Proof of Theorem 15. " \Longrightarrow " Assume system (1) is globally uniformly uniformly exponentially incrementally stable. Let $\begin{bmatrix} \phi(k;k_0,\xi) \\ \phi_{\delta}(k;k_0,\xi_{\delta},\xi) \end{bmatrix}$ denote the solution of system (9a)–(9b) for the initial condition $\begin{bmatrix} \xi \\ \xi_{\delta} \end{bmatrix} \in \mathbb{R}^{2n}$, whereby we note that the solutions to the dynamics (9b) (denoted by ϕ_{δ}) depend on the dynamics (9a), so that ϕ_{δ} depends on a third parameter ξ identifying a particular reference trajectory of (9a). Parametrize the straight line segment connecting ξ and $\xi + \xi_{\delta}$ at time k_0 by a function γ_{k_0} : $[0, 1] \to \mathbb{R}^n$ given by

$$\gamma_{k_0}(s) \coloneqq \xi + s\xi_\delta,\tag{55}$$

where $s \in [0, 1]$. At time $k \ge k_0$, we denote the parametrized curve initiated from segment (55) by

$$\gamma_k(s) \coloneqq \phi(k; k_0, \xi + s\xi_{\delta}) = \phi(k; k_0, \gamma_{k_0}(s)).$$

We make the following claim:

Claim 2. The trajectory $\begin{bmatrix} \phi(k;k_0,\xi) \\ \phi_{\delta}(k;k_0,\xi_{\delta},\xi) \end{bmatrix}$ of system (9a)-(9b) is identical to $\begin{bmatrix} \phi(k;k_0,\xi) \\ \frac{d}{ds}\phi(k;k_0,\gamma_{k_0}(s)) \end{bmatrix}$; i.e., $\phi_{\delta}(k;k_0,\xi_{\delta},\xi) = \frac{d}{ds}\phi(k;k_0,\gamma_{k_0}(s))$ for all $k,k_0 \in \mathbb{Z}$ such that $k \geq k_0$.

Proof. By the chain rule,

$$\frac{d}{ds}f(k,\phi(k;k_0,\gamma_{k_0}(s))) = \frac{\partial f}{\partial x}(k,x)\frac{d}{ds}\phi(k;k_0,\gamma_{k_0}(s))$$

which implies that $\frac{d}{ds}\phi(k;k_0,\gamma_{k_0}(s))$ satisfies the displacement dynamics (9b). Thus, it is only left to prove that the initial conditions of both trajectories are the same; i.e., $\frac{d}{ds}\phi(k_0;k_0,\gamma_{k_0}(s)) = \xi_{\delta}$. Note that $\frac{d}{ds}\phi(k_0;k_0,\gamma_{k_0}(s)) = \frac{d}{ds}\gamma_{k_0}(s)$. Hence, by taking the derivative on both sides of (55) with respect to s, we obtain the desired result.

For any sufficiently small $\epsilon > 0$, Definition 2 yields $k \ge 1$, $\lambda > 1$ so that $|\phi(k; k_0, \gamma_{k_0}(s + \epsilon)) - \phi(k; k_0, \gamma_{k_0}(s))| \le \kappa |\gamma_{k_0}(s + \epsilon) - \gamma_{k_0}(s)|\lambda^{-(k-k_0)}$. (56) Dividing both sides of (56) by ϵ then sending $\epsilon \to 0$, we obtain

$$\left|\phi_{\delta}(k;k_{0},\xi_{\delta},\xi)\right| = \left|\frac{d}{ds}\phi(k;k_{0},\gamma_{k_{0}}(s))\right| \le \kappa |\xi_{\delta}|\lambda^{-(k-k_{0})}.$$
(57)

Thus, from the global exponential incremental stability of the subsystem (9a), by considering (9a)-(9b), we have established the global exponential stability of the subsystem (9b). As this system is linear, its global exponential stability is the same as global exponential incremental stability.

In what follows, we will construct a nonsingular matrix $\Theta(k, x)$ and, hence construct F(k, x) such that (6) is satisfied for all $x \in \mathbb{R}^n$ and $k \in \mathbb{Z}$.

To this end observe that the transfer matrix $\Phi(k, k_0; \xi)$ of the linear displacement dynamics (9b), which satisfies $\Phi(k, k_0; \xi)\xi_{\delta} = \phi(k; k_0, \xi_{\delta}, \xi)$ for all $\xi_{\delta} \in \mathbb{R}^n$ and all $k \geq k_0$, satisfies the exponential bound

$$\|\Phi(k,k_0;\xi)\| \le \kappa \lambda^{-(k-k_0)} \tag{58}$$

(with $\kappa \geq 1$, $\lambda > 1$) for $k \geq k_0$ independently of the initial condition ξ of the reference trajectory generated by (9a). From here we may follow, mutatis mutandis, the proof of [39, Theorem 23.3] and define an $n \times n$ matrix

$$Q(k,\xi) \coloneqq \sum_{j=k}^{\infty} \left(\Phi(j,k;\xi) \right)^T \Phi(j,k;\xi)$$

and noting that, due to (58), $||Q(k,\xi)|| \leq \frac{\kappa^2 \lambda^2}{\lambda^2 - 1}$ independently of k and ξ , so $Q(k,\xi)$ is well defined. Following the remainder of said proof, one establishes, mutatis mutandis, that for all $x \in \mathbb{R}^n$ and $k \in \mathbb{Z}$,

$$\eta I \preceq Q(k, x) \preceq \rho I, \tag{59}$$

$$\frac{\partial f}{\partial x}(k,x)^T Q(k+1,x) \frac{\partial f}{\partial x}(k,x) - Q(k,x) \preceq -\nu I$$
(60)

where η, ρ , and ν are positive constants.

Applying the Cholesky factorization on the uniformly positive definite matrix Q(k, x), there exists a nonsingular matrix $\Theta(k, x)$ such that, $Q(k, x) = \Theta(k, x)^T \Theta(k, x)$. With this construction of $\Theta(k, x)$, condition (5) is automatically satisfied by appealing to (59). Now, multiplying each side of (60) by $\Theta(k, x)^{-T}$ from the left and $\Theta(k, x)^{-1}$ from the right, we obtain

$$\Theta(k,x)^{-T} \frac{\partial f}{\partial x}(k,x)^{T} \\ \times \Theta(k+1,x)^{T} \Theta(k+1,x) \frac{\partial f}{\partial x}(k,x) \Theta(k,x)^{-1} \\ - \Theta(k,x)^{-T} Q(k,x) \Theta(k,x)^{-1} \\ \preceq -\nu \Theta(k,x)^{-T} \Theta(k,x)^{-1}$$

which, with (7), implies

$$F(k,x)^T F(k,x) - I \preceq -\nu \left(\Theta(k,x)\Theta(k,x)^T\right)^{-1}.$$
(61)

Next, since $Q(k, x) \preceq \rho I$ by (59), for any $s \in \mathbb{R}^n$ we have

$$\left(\Theta(k,x)^{-1}s\right)^T Q(k,x) \left(\Theta(k,x)^{-1}s\right) \le \rho\left(\Theta(k,x)^{-1}s\right)^T \left(\Theta(k,x)^{-1}s\right)^T \left(\Theta$$

Straightforward manipulations result in

$$\frac{1}{\rho}s^T s \le s^T \left(\Theta(k, x)\Theta(k, x)^T\right)^{-1} s.$$

Thus, we arrive at

$$-\nu \left(\Theta(k,x)\Theta(k,x)^T\right)^{-1} \preceq -\frac{\nu}{\rho}I.$$
(62)

Combining (61) with (62), we see that

$$F(k,x)^T F(k,x) - I \preceq -\frac{\nu}{\rho}I.$$

Therefore, condition (6) is satisfied for this construction of F(k, x) with $\mu := \frac{\nu}{\rho}$, in other words, system (1) is uniformly globally contracting. " \leftarrow " Now, assume that (1) is uniformly globally contracting. Define ma-

" \Leftarrow " Now, assume that (1) is uniformly globally contracting. Define matrix $Q(k,x) \coloneqq \Theta(k,x)^T \Theta(k,x)$. By the definition of $\Theta(k,x)$, condition (5) implies Q(k,x) is uniformly positive definite; i.e., there exists $\eta > 0, \rho > 0$ so that

$$\eta I \preceq Q(k, x) \preceq \rho I. \tag{63}$$

Multiplying each side of (6) with $\Theta(k, x)^T$ from the right and $\Theta(k, x)$ from the left, then expanding by using (7), we have

$$\Theta(k,x)^T \Big(\Theta(k,x)^{-T} \frac{\partial f}{\partial x}(k,x)^T \Theta(k+1,x)^T \\ \times \Theta(k+1,x) \frac{\partial f}{\partial x}(k,x) \Theta(k,x)^{-1} \Big) \Theta(k,x) \\ - \Theta(k,x)^T \Theta(k,x) \preceq -\mu \Theta(k,x)^T \Theta(k,x).$$

After straightforward simplifications, we see that

$$\frac{\partial f}{\partial x}(k,x)^T Q(k+1,x) \frac{\partial f}{\partial x}(k,x) - Q(k,x) \preceq -\mu Q(k,x) \preceq -\frac{\eta}{\mu} I \qquad (64)$$

where the second matrix inequality follows directly from the lower bound of (63). Thus, from (63) and (64), applying [39, Theorem 23.3], we conclude that the linear time-varying subsystem (9b) is uniformly exponentially stable. Furthermore, from the first inequality of (64), we have

$$\frac{\partial f}{\partial x}(k,x)^T Q(k+1,x) \frac{\partial f}{\partial x}(k,x) \preceq \beta Q(k,x)$$
(65)

for some $\beta \in (0, 1)$.

Pick any $\xi_1, \xi_2 \in \mathbb{R}^n$. Consider a straight line segment connecting ξ_1, ξ_2 and parametrized by a function $\gamma_{k_0} \colon [0,1] \to \mathbb{R}^n$, at k_0 , given by

$$\gamma_{k_0}(s) \coloneqq s\xi_1 + (1-s)\xi_2,$$
(66)

where $s \in [0, 1]$. The length of this segment at $k = k_0$ is $l_0 = |\xi_1 - \xi_2|$. Then, for any $k > k_0$,

$$\gamma_k(s) \coloneqq \phi(k; k_0, \gamma_{k_0}(s)) \tag{67}$$

is a curve connecting $\phi(k; k_0, \xi_1)$ to $\phi(k; k_0, \xi_2)$ parametrized by $s \in [0, 1]$. The length of the curve defined in (67) at time k is given by

$$l(k) = \int_0^1 \sqrt{\frac{d}{ds} \gamma_k(s)^T Q(k, \gamma_k(s)) \frac{d}{ds} \gamma_k(s)} ds.$$

Applying the chain rule to $\gamma_{k+1}(s)$ for $s \in [0,1]$ yields

$$\frac{d}{ds}\gamma_{k+1}(s) = \frac{d}{ds}\phi(k+1;k_0,\gamma_{k_0}(s))$$
$$= \frac{d}{ds}f(k,\phi(k;k_0,\gamma_{k_0}(s)))$$
$$= \frac{\partial f}{\partial x}(k,x)\frac{d}{ds}\phi(k;k_0,\gamma_{k_0}(s)) = \frac{\partial f}{\partial x}(k,x)\frac{d}{ds}\gamma_k(s).$$

We see that

$$\begin{aligned} \frac{d}{ds}\gamma_{k+1}(s)^T Q(k+1,\gamma_{k+1}(s)) \frac{d}{ds}\gamma_{k+1}(s) \\ &= \left(\frac{\partial f}{\partial x}(k,x) \frac{d}{ds}\gamma_k(s)\right)^T Q(k+1,\gamma_{k+1}(s)) \\ &\times \left(\frac{\partial f}{\partial x}(k,x) \frac{d}{ds}\gamma_k(s)\right) = \frac{d}{ds}\gamma_k(s)^T \times \\ &\left[\frac{\partial f}{\partial x}(k,x)^T Q(k+1,\gamma_{k+1}(s)) \frac{\partial f}{\partial x}(k,x)\right] \frac{d}{ds}\gamma_k(s) \\ &\leq \frac{d}{ds}\gamma_k(s)^T \beta Q(k,\gamma_k(s)) \frac{d}{ds}\gamma_k(s), \end{aligned}$$

where the final inequality follows from (65). Therefore,

$$\begin{split} l(k+1) - l(k) \\ &= \int_0^1 \sqrt{\frac{d}{ds} \gamma_{k+1}(s)^T Q(k+1,\gamma_{k+1}(s)) \frac{d}{ds} \gamma_{k+1}(s)} ds \\ &- \int_0^1 \sqrt{\frac{d}{ds} \gamma_k(s)^T Q(k,\gamma_k(s)) \frac{d}{ds} \gamma_k(s)} ds \\ &\leq (\sqrt{\beta} - 1) l(k) \end{split}$$

which, in turn, implies $l(k+1) \leq \frac{l(k)}{\lambda}$, where $\lambda = \frac{1}{\sqrt{\beta}} > 1$ since $\beta \in (0, 1)$. Consequently, it is straightforward to see that $l(k) \leq l_0 \lambda^{-(k-k_0)}$ and, hence,

$$|\phi(k;k_0,\xi_1) - \phi(k;k_0,\xi_2)| \le l(k) \le l_0 \lambda^{-(k-k_0)} = |\xi_1 - \xi_2| \lambda^{-(k-k_0)}$$

for all $\xi_1, \xi_2 \in \mathbb{R}^n$ and $k, k_0 \in \mathbb{Z}$ such that $k \ge k_0$. We conclude that (1) is globally exponentially incrementally stable.

Proof of Theorem 16. The proof of Theorem 16 is a consequence of Theorem 13 and Theorem 15. In particular, by Theorem 15, we see that a globally contracting system is globally exponentially incrementally stable. Then, by Theorem 13, we conclude that a globally contracting system whose

state evolves in a compact and positively invariant set, is uniformly globally convergent.

DUC N. TRAN AND CHRISTOPHER M. KELLETT ARE WITH THE SCHOOL OF ELECTRICAL ENGINEERING AND COMPUTER SCIENCE, THE UNIVERSITY OF NEWCASTLE, CALLAGHAN, NSW 2308, AUSTRALIA.

E-mail address: DucNgocAnh.TranQuon.edu.au *E-mail address*: Chris.KellettQnewcastle.edu.au

 $\rm Björn~S.$ Rüffer is with the School of Mathematical and Physical Sciences, The University of Newcastle, Callaghan, NSW 2308, Australia.

E-mail address: Bjorn.Ruffer@newcastle.edu.au