

Correspondence

Corrections to “Empirical Estimation of Leaf Chlorophyll Density in Winter Wheat Canopies Using Sentinel-2 Spectral Resolution”

Massimo Vincini, *Member, IEEE*, Stefano Amaducci,
and Ermes Frazzi

In the above paper [1], clarification of Equation (2) is needed. The power model (2), i.e.,

$$y = a \cdot x^b + c$$

was used with leaf chlorophyll density as the independent variable x and the VI values as the dependent variable y in order to evaluate the sensitivity function (3) (see Fig. 6). Conversely, the same model was used with the VI values as the independent variable and leaf chlorophyll density as the dependent variable for all other regressions between VI values and chlorophyll density (see Figs. 2 and 4, and Tables IV–VI).

REFERENCES

- [1] M. Vincini, S. Amaducci, and E. Frazzi, “Empirical estimation of leaf chlorophyll density in winter wheat canopies using Sentinel-2 spectral resolution,” *IEEE Trans. Geosci. Remote Sens.*, vol. 52, no. 6, pp. 3220–3235, Jun. 2014.

Manuscript received November 27, 2013. Date of publication January 24, 2014; date of current version March 3, 2014.

M. Vincini and E. Frazzi are with Centro Ricerca Analisi geoSpaziale e Telerilevamento (CRAT), Università Cattolica del Sacro Cuore, 29122 Piacenza, Italy (e-mail: massimo.vincini@unicatt.it; ermes.frazzi@unicatt.it).

S. Amaducci is with Istituto di Agronomia Generale, Genetica e Coltivazioni erbacee, Università Cattolica del Sacro Cuore, 29122 Piacenza, Italy (e-mail: stefano.amaducci@unicatt.it).

Digital Object Identifier 10.1109/TGRS.2013.2294451