# Critical pairs for the Product Singleton Bound

Diego Mirandola<sup>\*</sup> and Gilles Zémor<sup>†</sup>

September 6, 2018

#### Abstract

We characterize Product-MDS pairs of linear codes, i.e. pairs of codes C, D whose product under coordinatewise multiplication has maximum possible minimum distance as a function of the code length and the dimensions dim C, dim D. We prove in particular, for C = D, that if the square of the code C has minimum distance at least 2, and (C, C) is a Product-MDS pair, then either C is a generalized Reed-Solomon code, or Cis a direct sum of self-dual codes. In passing we establish coding-theory analogues of classical theorems of additive combinatorics.

Keywords: Error-correcting codes, Schur-product codes, Product Singleton Bound.

### 1 Introduction

Let F be a finite field. Given vectors  $x = (x_1, \ldots, x_n)$ ,  $y = (y_1, \ldots, y_n)$  of  $F^n$ , let us denote by xy the componentwise product of x and y,

$$xy = (x_1y_1, \dots, x_ny_n)$$

Given two linear codes  $C, D \subseteq F^n$ , let us denote by CD the *F*-linear subspace of  $F^n$  generated by all products  $xy, x \in C, y \in D$ . This product, sometimes called the Schur product, has usually been denoted by C \* D, but we wish to lighten notation. Likewise we shall denote the Schur square (henceforth square) of a code C by  $C^2$ : context should prevent confusion with cartesian products.

Products of codes turn up in a variety of situations, such as algebraic error correction, secret sharing and multiparty computation, algebraic complexity theory, lattice constructions, and lately cryptanalysis. For an exhaustive discussion on the connection between code products and these different topics, the reader is referred to [4]. A number of efforts have gone into describing the code-theoretic structure of code products, see [6, Chapter 12] and [15] for an extensive review of the current state of the art. In particular, [15] collects several technical results which will be cited explicitly in Section 2 and used later in this paper. Among these, the following bound on the minimum distance of products, which was first proved in [13], appears.

THEOREM 1.1 (Product Singleton Bound [13]). Let  $C, D \subseteq F^n$  be linear codes. Then

$$d_{\min}(CD) \le \max\{1, n - (\dim C + \dim D) + 2\}.$$
(1)

<sup>\*</sup>CWI Amsterdam and Mathematical Institute, Leiden University, The Netherlands, and Institut de Mathématiques de Bordeaux, Université de Bordeaux, France. Email: diego@cwi.nl.

<sup>&</sup>lt;sup>†</sup>Institut de Mathématiques de Bordeaux, Université de Bordeaux, France. Email: zemor@math.u-bordeaux.fr.

This paper was presented at the 2015 Ninth International Workshop on Coding and Cryptography and at the 2015 Information Theory and Applications Workshop. The second author is thankful for financial support from IdEx Bordeaux - CPU (ANR-10-IDEX-03-02).

A slightly stronger version of Theorem 1.1 is actually proved in [13], as is a version involving the product of more than two codes, but the above statement is really what motivates our discussion. We shall call the upper bound (1) the *Product Singleton Bound*, that can be thought of as a generalization of the classical Singleton Bound. Indeed, the classical Singleton Bound for a single code C is recovered by taking the code D in Theorem 1.1 to be of dimension 1 and minimum distance n.

Our goal in this paper is to characterize pairs (C, D) of codes that achieve equality in (1). We make the remark that if  $d_{\min}(CD)$  is allowed to be equal to 1, then pairs achieving equality in (1) can be almost anything, since typical pairs of codes will have a product equal to the whole space  $F^n$ . For a study of this phenomenon see [4, 14]. So we shall disregard the situation when  $d_{\min}(CD) = 1$  and call (C, D) a *Product-MDS (PMDS)* pair if it achieves equality in (1) and  $d_{\min}(CD) \ge 2$ .

As mentioned above, a PMDS pair can consist of an ordinary MDS code and a code of dimension 1. It is a natural question to ask what other PMDS pairs exist. It turns out that there is a surprisingly complete answer to this question. We shall show in particular that if (C, D) is a PMDS pair such that dim  $C \ge 2$ , dim  $D \ge 2$ , and  $d_{\min}(CD) \ge 3$ , then C and D can only be Reed-Solomon codes. By this we mean Reed-Solomon code in the widest sense, i.e. generalized, possibly extended or doubly extended in the terminology of [11], or Cauchy codes as in [7]. PMDS pairs with  $d_{\min}(CD) = 2$  will also be described quite precisely. To be more specific, in the symmetric case C = D we shall prove:

THEOREM 1.2. If (C, C) is a PMDS pair, then C is either a Reed-Solomon code or a direct sum of self-dual codes.

Self-duality in the above statement should be understood to be relative to a non-degenerate bilinear form which is not necessarily the standard inner product.

To establish these results we shall import methods from additive combinatorics and establish coding-theoretic analogues of the classical theorems of Kneser [10] and Vosper [17]. For background on and proofs of Kneser and Vosper's Theorems we refer to [16]. Kneser's Theorem implies in particular that if A, B are subsets of an abelian group such that

$$|A + B| < |A| + |B| - 1$$

then A + B must be periodic, i.e. there exists a non-zero element g of the abelian group that stabilizes A + B so that we have A + B + g = A + B. Our coding-theoretic variant of Kneser's Theorem will imply that if C and D are two codes such that

$$\dim CD < \dim C + \dim D - 1,$$

then the code C is necessarily the direct sum of two non-zero codes, which is equivalent to the existence of a non-constant vector x of  $F^n$  such that xCD = CD.

Vosper's Theorem is a characterization of pairs of subsets A, B of the integers modulo a prime p with the property that |A + B| = |A| + |B| - 1. It states that, excluding some degenerate cases, A, B must be arithmetic progressions with the same difference. We make the remark that if a code C has a generator matrix with rows  $g, g\alpha, \ldots, g\alpha^{k-1}$ , i.e. has a basis of elements in "geometric" progression then, provided g is of weight n and  $\alpha$  has distinct coordinates, C must be a Reed-Solomon code. This is why a code-theoretic version of Vosper's Theorem forces the appearance of Reed-Solomon codes. There will be some twists to the analogy however that we shall discuss later in the paper.

The paper is organized as follows. Section 2 sets up notation and preliminary results and states our primary contribution, Theorem 2.12. Section 3 states and proves the codingtheory equivalent of Kneser's Theorem. Section 4 is dedicated to a coding-theory version of Vosper's Theorem. Section 5 shows how to recover a version of Randriambololona's Product Singleton Bound as a straightforward consequence of Kneser's Theorem and goes on to derive the proof of Theorem 2.12. Section 6 concludes with some comments.

### 2 Preliminaries and statement of the main Theorem

Throughout the paper F will denote a finite field. We shall need, in a couple of occasions, to deal with fields that may be infinite in which case we will use the notation K.

Given a vector  $x \in K^n$ , we denote by  $\sup x$  its support and by wt(x) its weight. The support of a subvector space of  $K^n$  is defined as the union of the supports of all its vectors, and we shall say that a vector space in  $K^n$  has *full support* if its support is  $\{1, \ldots, n\}$ .

In this paper all codes will be linear. We will call them simply "codes" when the ambient space is  $F^n$ , and use the terminology of vector spaces in the general setting of  $K^n$ .

### 2.1 MDS Codes

Given a code  $C \subseteq F^n$ , we denote by  $C^{\perp}$  its dual with respect to the standard inner product in  $F^n$  and by  $d_{\min}(C)$  its minimum distance.

The classical Singleton Bound states

$$\dim C + d_{\min}(C) \le n+1.$$

A code which attains this bound is said to be Maximum Distance Separable (MDS). We recall the following well-known properties and characterizations of MDS codes [11].

LEMMA 2.1. Given a code  $C \subseteq F^n$ , the following statements are equivalent:

- 1. C is MDS,
- 2.  $C^{\perp}$  is MDS,
- 3. any  $\dim C$  columns of a generator matrix of C are linearly independent,
- 4. for any coordinate set I of size  $n + 1 \dim C$ , there is a word of C whose support equals I.

The following property is somewhat less standard.

LEMMA 2.2. Let  $C \subseteq F^n$  be a code. It is MDS if and only if any systematic generator matrix of C has all its rows of weight  $n + 1 - \dim C$ .

PROOF. It is clear that if C is MDS the property must hold. The converse implication is an immediate consequence of the following claim: if  $C \subseteq F^n$  is any code and  $x \in C$  is a codeword of minimal weight, then there is a systematic generator matrix of C whose first row is x.

We now prove this claim. Renumbering the coordinates, we may assume that  $\operatorname{supp} x = \{1, 2 + n - \operatorname{wt}(x), \ldots, n\}$  and that

$$x = (1, 0, \dots, 0, *, \dots, *),$$

where the stars denote non-zero entries. Let  $\{x_1 = x, x_2, \ldots, x_k\}$  be an *F*-basis of *C* containing *x*, where  $k := \dim C$ , and let *G* be the generator matrix of *C* whose rows are the  $x_i$ 's. If *G* can be made systematic in the first  $1 + n - \operatorname{wt}(x)$  positions then we are done. Otherwise, we obtain a contradiction as follows. We have that  $\operatorname{wt}(x) > 1$  and the rank of the matrix G restricted to its first 1 + n - wt(x) columns is  $\langle k$ . There exists therefore a linear combination

$$\tilde{x} = \sum_{i=2}^{k} \alpha_i x_i,$$

with  $\alpha_2, \ldots, \alpha_k \in F$ , which has zeros in positions  $\{2, \ldots, 1 + n - \operatorname{wt}(x)\}$ , but with  $\tilde{x} \neq 0$ . Now a suitable combination of x and  $\tilde{x}$  yields a non-zero word of weight smaller than  $\operatorname{wt}(x)$ , contradicting the minimality of  $\operatorname{wt}(x)$ .

Among MDS codes, Reed-Solomon codes, in the widest possible sense, will be prominent. A Reed-Solomon code of length n and dimension k is a code of the form

$$\{(g_1f(\alpha_1),\ldots,g_nf(\alpha_n)): f\in F[X]_{\leq k}\},\$$

where  $F[X]_{<k}$  denotes the space of polynomials of degree less than  $k, g_1, \ldots, g_n$  are nonzero elements of F, and  $\alpha_1, \ldots, \alpha_n$  are pairwise disjoint and belong to  $F \cup \{\infty\}$ , with the convention that for any  $f \in F[X]_{<k}, f(\infty)$  equals the coefficient of  $X^{k-1}$  in f. We shall call  $(\alpha_1, \ldots, \alpha_n)$  an evaluation-point sequence for the Reed-Solomon code.

This code family includes the codes called generalized, extended, and doubly-extended Reed-Solomon codes. From the geometric point of view, they may be thought of as the projective version of Reed-Solomon codes. In [7] they are named "Cauchy codes" and have also been called "Cauchy Reed-Solomon codes". We shall simply refer to them as "Reed-Solomon codes".

REMARK 2.3. If C and D are two Reed-Solomon codes with a common evaluation-point sequence  $\alpha$ , then the product CD is also Reed-Solomon with evaluation-point sequence  $\alpha$  and we have dim  $CD = \min\{n, \dim C + \dim D - 1\}$ . Theorem 2.5 below implies that  $\min\{n, \dim C + \dim D - 1\}$  is the minimum possible dimension of the product of MDS codes.

#### 2.2 Code products

For an arbitrary field K, the space  $K^n$  is, with the coordinatewise product, a commutative unitary K-algebra. Its unit element is the all-one vector, denoted by **1**. The multiplicative group of its invertible elements is  $(K^n)^{\times} = (K^{\times})^n$ , meaning that  $x \in K^n$  is invertible if and only if all entries of x are non-zero. Given  $x \in (K^n)^{\times}$ , its inverse is denoted by  $x^{-1}$ . The following simple observation will be freely used later.

LEMMA 2.4. Let  $x \in (K^n)^{\times}$ . For any vector space  $V \subseteq K^n$ , we have dim  $V = \dim xV$ .

The two results below relate the dimension of the product of two codes with the MDS property. The first one is taken from  $[15, \S 3.5]$ .

THEOREM 2.5. Let  $C, D \subseteq F^n$  be full-support codes. If (at least) one of them is MDS, then

 $\dim CD \ge \min\{n, \dim C + \dim D - 1\}.$ 

We also observe the following.

LEMMA 2.6. Let  $C, D \subseteq F^n$  be MDS codes such that

 $\dim CD = \dim C + \dim D - 1.$ 

Then CD is MDS.

PROOF. By Lemma 2.1, it suffices to show that for any choice of  $I \subseteq \{1, \ldots, n\}$  with  $|I| = d^* := n + 1 - \dim CD$  there exist  $x \in C, y \in D$  with  $\operatorname{supp} xy = I$ . Without loss of generality, assume that  $I = \{1, \ldots, d^*\}$ . As C and D are both MDS, there exist  $x \in C$  and  $y \in D$  such that  $\operatorname{supp} x = I \cup \{d^* + 1, \ldots, d_C\}$  and  $\operatorname{supp} y = I \cup \{n - (d_D - d^*) + 1, \ldots, n\}$ , where  $d_C$  and  $d_D$  denote the minimum distance of C and D respectively. One checks that  $d_C = n - (d_D - d^*)$ , hence indeed  $\operatorname{supp} xy = \operatorname{supp} x \cap \operatorname{supp} y = I$ .

#### 2.3 Stabilizer algebras and decomposable codes

Lemma 2.7 below classifies all subalgebras of  $K^n$ . For all i = 1, ..., n, let  $e_i$  denote the *i*-th unit vector in  $K^n$ . We call a vector of the form  $\sum_{i \in I} e_i$  for some  $I \subseteq \{1, ..., n\}$  a projector. In particular, **1** is the projector with support  $\{1, ..., n\}$ . A family of projectors is *disjoint* if the projectors have pairwise disjoint supports.

LEMMA 2.7. Any K-subalgebra of  $K^n$  admits a K-basis of disjoint projectors.

PROOF. Let  $A \subseteq K^n$  be a K-subalgebra. We argue by induction on  $k := \dim A$ . If k = 1then A is generated by a vector x whose non-zero coordinates must be all equal, otherwise  $x^2$  is not a K-multiple of x. If k > 1, pick  $x \in A, x \neq 0$  of minimal support with one of its coordinates equal to 1, and let  $\{x_1 = x, \ldots, x_k\}$  be a K-basis of A containing x. Then x is a projector, otherwise  $x^2 - x \neq 0$  would have smaller support. For all  $i = 2, \ldots, k$ , if  $\sup x_i$ and  $\sup x$  intersect, say in position j, then we can choose  $\lambda_i \in K$  so that  $x_i + \lambda_i x$  has a zero in position j, hence  $x(x_i + \lambda_i x)$  has support strictly smaller than x. By minimality of  $\sup x_i + \lambda_i x$ , we obtain that A is a direct sum  $A = \langle x \rangle \oplus \langle x_2, \ldots, x_k \rangle$  and the conclusion follows by applying the induction hypothesis to the second summand.

REMARK 2.8. Lemma 2.7 implies in particular that the number of subalgebras of  $K^n$  is finite. This fact will be useful later.

Let  $V \subseteq K^n$  be a K-vector space. We define  $St(V) := \{x \in K^n : xV \subseteq V\}$ , the stabilizer of V in  $K^n$ . As V is linear, St(V) is a K-algebra, hence Lemma 2.7 applies. In particular, St(V) has a basis of vectors whose entries are all 0's and 1's. This leads to the following lemma, that characterizes the stabilizer of a base-field extension<sup>1</sup>. For a proof we refer to [15, Proposition 2.24].

LEMMA 2.9. Let K'/K be a field extension. Let  $V \subseteq K^n$  be a K-vector space. Then

$$\operatorname{St}(V \otimes K') = \operatorname{St}(V) \otimes K'.$$

Let  $C \subseteq F^n$  be a code. As in the vector-space case, we can define its stabilizer and apply Lemma 2.7, which yields an *F*-basis  $\{\pi_1, \ldots, \pi_h\}$  of  $\operatorname{St}(C)$  of disjoint projectors, with  $h := \operatorname{dim} \operatorname{St}(C)$ . When h = 1 we say that *C* has trivial stabilizer. We have the following lemma, whose proof is straightforward.

LEMMA 2.10. Any full-support code  $C \subseteq F^n$  decomposes as

$$C = \pi_1 C \oplus \dots \oplus \pi_h C$$

where  $\{\pi_1, \ldots, \pi_h\}$  is an *F*-basis of disjoint projectors of St(C). Moreover, each summand  $\pi_i C$ , viewed as a code in  $F^{|\operatorname{supp} \pi_i|}$ , has trivial stabilizer and full support.

<sup>&</sup>lt;sup>1</sup>If K'/K is a field extension, the base-field extension  $V \otimes K'$ , where the tensor product is taken over K, of V is the K'-span of V.

Facts on stabilizers, including Lemmas 2.9 and 2.10, can be found in [15, from §2.6 onwards]. Following [15], let us say that a code is *indecomposable* if it has trivial stabilizer.

Lemma 2.10 states in particular that a full-support code has non-trivial stabilizer if and only if it decomposes as a direct sum of codes, and the dimension of the stabilizer equals the number of indecomposable components. It follows that all MDS codes that are not equal to  $F^n$  have trivial stabilizer.

We conclude this section with two refinements of the classical Singleton Bound, involving the dimension of St(C) beside the usual parameters. They naturally reduce to the classical Singleton Bound when the code C is indecomposable, i.e. dim St(C) = 1.

LEMMA 2.11. Let  $C \subseteq F^n$  be a code.

1. If  $d_{\min}(C) > 1$  then

$$d_{\min}(C) \le n - \dim C + 1 - (\dim \operatorname{St}(C) - 1).$$

2. If C has full support then

$$d_{\min}(C) \le \frac{n - \dim C}{\dim \operatorname{St}(C)} + 1.$$

PROOF. We may assume that C has full support, as the first claim in the general case follows immediately from the first claim in the full-support case. Set  $k := \dim C$ ,  $d := d_{\min}(C)$  and  $h := \dim \operatorname{St}(C)$ . By Lemma 2.10 we have that C is a direct sum  $C = C_1 \oplus \cdots \oplus C_h$  of full-support codes. For all  $i = 1, \ldots, h$ , let  $n_i, k_i$  and  $d_i$  denote the support size, the dimension and the minimum distance of  $C_i$  respectively. We have  $\sum_{i=1}^{h} n_i = n, \sum_{i=1}^{h} k_i = k$  and

$$d = \min_{i} \{d_i\} \le \min_{i} \{n_i - k_i\} + 1$$

by the classical Singleton Bound. In the case d > 1 we have  $n_i - k_i \ge d_i - 1 \ge 1$  for all i = 1, ..., h, hence, for all j = 1, ..., h,

$$n_j - k_j = n - k - \sum_{i \neq j} (n_i - k_i) \le n - k - (h - 1).$$

Putting everything together we have

$$d = \min_{i} \{d_i\} \le \min_{i} \{n_i - k_i\} + 1 \le n - k + 1 - (h - 1),$$

which proves the first claim. To prove the second claim, note that

$$n-k = \sum_{i=1}^{h} (n_i - k_i) \ge h \min_i \{n_i - k_i\},$$

hence  $\min_i \{n_i - k_i\} \leq (n - k)/h$  and the conclusion follows.

#### 2.4 Main Theorem

Our main result takes the following form.

THEOREM 2.12. Let  $C, D \subseteq F^n$  be codes such that the pair (C, D) is Product MDS. Then one of the following situations occurs.

 $\triangle$ 

- (i) C and D are MDS and, if none of them has dimension 1, they are Reed-Solomon codes with a common evaluation-point sequence.
- (ii) There is a partition of the coordinate set into non-empty subsets

$$\{1,\ldots,n\}=I_1\cup\cdots\cup I_h$$

and there exist h pairs  $(C_1, D_1), \ldots, (C_h, D_h)$  of codes of  $F^n$ , such that supp  $C_i =$  supp  $D_i = I_i$ , for all  $i = 1, \ldots, h$ , and such that C and D decompose as:

$$C = C_1 \oplus \dots \oplus C_h,$$
$$D = D_1 \oplus \dots \oplus D_h.$$

Furthermore, for all i = 1, ..., h, when  $C_i$  and  $D_i$  are identified with codes of  $F^{|I_i|}$ through the natural projection on their support, we have that  $C_i = (g_i D_i)^{\perp}$  for some  $g_i \in (F^{|I_i|})^{\times}$ .

REMARK 2.13. The codes  $C_i$  and  $D_i$  are mutually orthogonal relative to the non-degenerate bilinear form  $(x, y) \mapsto (x | g_i y) = (g_i x | y)$ , where  $(\cdot | \cdot)$  denotes the standard inner product. Hence the wording of Theorem 1.2 in the case C = D.

### 3 Kneser's Theorem

Kneser's Addition Theorem below involves the stabilizer  $St(X) = \{g \in G : g + X = X\}$  of a subset X of an abelian group G. The (Minkowski) sum A + B of two subsets A, B of G is defined as the set of sums a + b when a and b range over A and B respectively.

THEOREM 3.1 (Kneser [10]). Let G be an abelian group. Let  $A, B \subseteq G$  be non-empty, finite subsets. Then

$$|A + B| \ge |A| + |B| - |\operatorname{St}(A + B)|.$$

Kneser's original Theorem was transposed to the extension field setting by Hou, Leung and Xiang in [8]. Let L/K be a field extension. For K-linear subspaces  $S, T \subseteq L$ , we may consider the product of subspaces ST defined as the K-linear span of the set of elements of the form  $st, s \in S, t \in T$ . Hou et al.'s Theorem is concerned with the structure of pairs of subspaces whose product has small dimension. Again, the stabilizer of a K-subspace  $X \subseteq L$ is involved and is defined in the expected way  $St(X) = \{z \in L : zX \subseteq X\}$ .

THEOREM 3.2 (Generalized Kneser Theorem [8]). Let L/K be a separable field extension. Let  $S, T \subseteq L$  be non-zero, finite-dimensional K-vector spaces. Then

$$\dim ST \ge \dim S + \dim T - \dim \operatorname{St}(ST).$$

Remarkably, Kneser's original Theorem for groups can be recovered easily from Hou et al.'s version.

We will now proceed to show that there is a variant of Kneser's Theorem for the algebra induced by coordinatewise multiplication.

THEOREM 3.3. Let  $S, T \subseteq K^n$  be non-zero K-vector spaces. Then

$$\dim ST \ge \dim S + \dim T - \dim \operatorname{St}(ST).$$

REMARK 3.4. The products ST in Theorems 3.2 and 3.3 are in different algebras. The statement of Theorem 3.2 is the only instance of the paper where the product ST does not refer to a coordinatewise product.

REMARK 3.5. Assuming that Theorem 3.3 holds in the case of full-support S and T, the general case can be derived as follows. Let  $S_0, T_0 \subseteq K^{n_0}$  be the projections of S, T respectively on supp ST, where  $n_0 := |\operatorname{supp} ST|$ . The spaces  $S_0$  and  $T_0$  both have full support, hence

$$\dim S_0 T_0 \ge \dim S_0 + \dim T_0 - \dim \operatorname{St}(S_0 T_0).$$

Clearly dim  $S_0T_0 = \dim ST$  and dim  $\operatorname{St}(ST) = \dim \operatorname{St}(S_0T_0) + n - n_0$ . It remains to prove that

$$\dim S_0 + \dim T_0 \ge \dim S + \dim T - (n - n_0).$$

Let  $S_1, T_1 \subseteq K^{n-n_0}$  be the projections of S, T respectively on the complement of supp ST. Observe that supp  $S_1$  and supp  $T_1$  cannot intersect, hence dim  $S_1$ +dim  $T_1 \leq n-n_0$ . Moreover dim  $S \leq \dim S_0$ +dim  $S_1$  and dim  $T \leq \dim T_0$ +dim  $T_1$ . Putting everything together we obtain the desired inequality.

From here on "Kneser's Theorem" will refer to Theorem 3.3 rather than to the original result. Our proof is strongly inspired by Hou et al.'s proof of Theorem 3.2 [8], itself drawing upon the e-transform technique of additive combinatorics (see e.g. [16]).

If V is a K-subspace of  $K^n$ , we use the notation  $V^{\times}$  to mean the subset of invertible elements of V.

LEMMA 3.6. Let  $S, T \subseteq K^n$  be non-zero K-vector spaces. Assume that T has a basis of invertible elements. Then, for all  $x \in S^{\times}$ , there exist a K-algebra  $H_x \subseteq K^n$  and a K-vector space  $V_x \subseteq K^n$  such that  $H_x V_x = V_x$ ,  $xT \subseteq V_x \subseteq ST$  and

$$\dim V_x + \dim H_x \ge \dim S + \dim T.$$

PROOF. Assume that the lemma is proved for  $x = \mathbf{1}$ . Then, if  $S^{\times}$  is non-empty, for any  $x \in S^{\times}$  we may apply the result for the case  $x = \mathbf{1}$  to  $x^{-1}S$  and T. So we only need to prove the Lemma for  $\mathbf{1} \in S$  and  $x = \mathbf{1}$ . Analogously, we may assume that  $\mathbf{1} \in T$ .

We argue by induction on  $k := \dim S$ . If k = 1,  $H := K\mathbf{1}$  and V := T do the job. So assume that k > 1 and the result holds for smaller dimension. For each  $e \in T^{\times}$ , define

$$S(e) := S \cap Te^{-1}, \quad T(e) := T + Se.$$

We have  $S(e)T \subseteq ST$ ,  $S(e)Se \subseteq TS$ , therefore  $S(e)T(e) \subseteq ST$ . Furthermore,

$$\dim T(e) = \dim T + \dim Se - \dim(T \cap Se)$$
$$= \dim T + \dim S - \dim(Te^{-1} \cap S)$$

by Lemma 2.4, hence

$$\dim S(e) + \dim T(e) = \dim S + \dim T.$$

We distinguish two cases.

Assume that S(e) = S for all  $e \in T^{\times}$ , i.e.  $S \subseteq Te^{-1}$  for all  $e \in T^{\times}$ . Then, since T has a basis of invertible elements, we have  $ST \subseteq T$ . The result then holds with H the subalgebra generated by S and V := T.

Assume that there exists  $e \in T^{\times}$  such that  $S(e) \subsetneq S$ . Then  $0 < \dim S(e) < k$  hence the induction hypothesis applied to S(e) and T(e) gives an algebra H and a vector space V such that HV = V,

$$T \subseteq T(e) \subseteq V \subseteq S(e)T(e) \subseteq ST$$

and

$$\dim V + \dim H \ge \dim S(e) + \dim T(e) = \dim S + \dim T.$$

PROOF OF THEOREM 3.3. By Remark 3.5 we may assume that both S and T have full support. The key to the proof is the following observation. Assume that T has a basis of invertible elements. Recall that, by Lemma 3.6, for all  $x \in S^{\times}$  there exist a K-algebra  $H_x \subseteq K^n$  and a K-vector space  $V_x \subseteq F^n$  such that

$$H_x V_x = V_x \tag{2}$$

$$xT \subseteq V_x \subseteq ST \tag{3}$$

$$\dim V_x + \dim H_x \ge \dim S + \dim T. \tag{4}$$

Set  $k := \dim S$  and assume furthermore that there exists a K-basis  $\{x_1, \ldots, x_k\}$  of S contained in  $S^{\times}$  such that

$$H_{x_1} = \dots = H_{x_k} =: H. \tag{5}$$

Then  $ST = V_{x_1} + \cdots + V_{x_k}$  by (3), and therefore HST = ST by (2), in other words  $H \subseteq St(ST)$ . From (4) it follows therefore that

$$\dim ST + \dim \operatorname{St}(ST) \ge \dim V_{x_1} + \dim H \ge \dim S + \dim T,$$

hence the conclusion.

We shall first prove the Theorem when K is an infinite field, by showing in that case that T always has a basis of invertible elements and that there always exists a basis  $\{x_1, \ldots, x_n\}$  of invertible elements of S satisfying (5).

Since T has full support, it should be clear enough that it has a basis of invertible elements for K infinite. In this case Lemma 3.6 applies. Now fix a K-basis  $\{s_1, \ldots, s_k\}$  of S and define, for all  $\alpha \in K$ ,  $y_\alpha := \sum_{i=1}^k \alpha^{i-1} s_i \in S$ . For any choice of non-zero, pairwise distinct  $\alpha_1, \ldots, \alpha_k \in K$ , the matrix transforming  $s_1, \ldots, s_k$  into  $y_{\alpha_1}, \ldots, y_{\alpha_k}$  is Vandermonde, and therefore  $y_{\alpha_1}, \ldots, y_{\alpha_k}$  is also a K-basis of S. We now observe that the set  $\{\alpha \in K : y_\alpha \in S^\times\}$ is infinite: indeed its complement in K is finite, as it is a finite union of zero-sets of nonzero polynomials. That these polynomials are non-zero is guaranteed by the full-support property of S. On the other hand, the number of subalgebras of  $K^n$  is finite by Remark 2.8, in particular the number of subalgebras  $H_x$  guaranteed by Lemma 3.6 is finite. It follows that there exist  $\alpha_1, \ldots, \alpha_k$  such that  $\{x_1 = y_{\alpha_1}, \ldots, x_k = y_{\alpha_k}\}$  is a K-basis of S whose elements are all invertible and such that  $H_{x_1} = \cdots = H_{x_k}$ . This concludes the proof in the case K infinite.

Assume now that K is finite, and consider an infinite field extension K' of K, for example the rational function field K' := K(t), where t is transcendental over K. The infinite basefield case applies to K'-vector spaces. Our purpose is to draw our conclusion from this. Define the base-field extensions  $S' := S \otimes K', T' := T \otimes K'$ , where tensor products are taken over K. By construction S' and T' are K'-vector spaces and we have just proved that

$$\dim_{K'} S'T' \ge \dim_{K'} S' + \dim_{K'} T' - \dim_{K'} \operatorname{St}(S'T').$$

It is clear that  $S'T' = ST \otimes K'$ ,  $\dim_{K'} S' = \dim S$ ,  $\dim_{K'} T' = \dim T$  and  $\dim_{K'} S'T' = \dim ST$ , where non-indexed dimensions are taken over K. Moreover  $\operatorname{St}(S'T') = \operatorname{St}(ST) \otimes K'$  by Lemma 2.9 and the conclusion follows.

Theorem 3.3 implies in particular that if C and D are two codes such that CD has trivial stabilizer, i.e. is indecomposable, then we must have

$$\dim CD \ge \dim C + \dim D - 1. \tag{6}$$

The next section studies pairs of codes C, D such that CD is indecomposable and achieves equality in (6).

### 4 Vosper's Theorem

We start by recalling Vosper's Addition Theorem.

THEOREM 4.1 (Vosper [17]). Let G be an abelian group of prime order p. Let  $A, B \subseteq G$  be subsets, with  $|A|, |B| \ge 2$  and  $|A + B| \le p - 2$ . If

$$|A + B| = |A| + |B| - 1$$

then A and B are arithmetic progressions with the same difference.

We point out that an extension-field version of Vosper's Theorem for finite fields was recently proved in [1].

Since the stabilizer of a subset of a group G must be a subgroup, when G is of prime order and has no proper subgroup, Kneser's Addition Theorem 3.1 implies that subsets A, B of G such that  $A + B \neq G$  must satisfy

$$|A + B| \ge |A| + |B| - 1.$$

This result is known as the Cauchy-Davenport Inequality, see [12, 16]. Vosper's Theorem is therefore concerned with characterizing pairs of sets achieving equality in the Cauchy-Davenport Inequality.

In the algebra setting, the inequality (6) may be thought of as a code-product version of the Cauchy-Davenport Inequality. But contrary to the group case, the algebra  $F^n$  always has proper subalgebras (for n > 1) so we cannot hope to ensure (6) purely by a condition on  $F^n$ . However, we have seen that (6) holds when (at least one of) the codes involved is MDS (Theorem 2.5). The following theorem may be seen as a version of Vosper's Theorem for MDS codes, and is the main result of this section.

THEOREM 4.2. Let  $C, D \subseteq F^n$  be MDS codes, with dim C, dim  $D \ge 2$  and dim  $CD \le n-2$ . If

$$\dim CD = \dim C + \dim D - 1$$

then C and D are Reed-Solomon codes with a common evaluation-point sequence.

REMARK 4.3. The hypotheses dim C, dim  $D \ge 2$  clearly cannot be removed. The value n-2 is also best possible in the hypothesis dim  $CD \le n-2$ , since by taking C to be an arbitrary MDS (non Reed-Solomon) code, and taking  $D = C^{\perp}$ , we will have a pair of MDS codes such that dim  $CD = \dim C + \dim D - 1 = n - 1$ .

We introduce the following notation for Vandermonde-type matrices. Given a positive integer k and  $\alpha = (\alpha_1, \ldots, \alpha_n) \in (F \cup \{\infty\})^n$  we denote by  $V_k(\alpha)$  the  $k \times n$  matrix whose *i*-th column is  $(1, \alpha_i, \ldots, \alpha_i^{k-1})^T$  if  $\alpha_i \neq \infty, (0, \ldots, 0, 1)^T$  otherwise. Note that the possible presence of this last column makes  $V_k(\alpha)$  a Vandermonde matrix in a generalized sense. We remark that if the entries of  $\alpha$  are pairwise distinct then  $V_k(\alpha)$  has full rank. With this notation, a Reed-Solomon code of length n and dimension k is a code of the form gC, where  $g \in (F^{\times})^n$  (i.e. g has no zero entries) and C is generated by  $V_k(\alpha)$  for some  $\alpha \in (F \cup \{\infty\})^n$ with pairwise-distinct entries. The vector  $\alpha$  is an evaluation-point sequence of C (see the end of Section 2.1).

LEMMA 4.4. Let  $C \subseteq F^n$  be a full-support code with dim  $C \ge 2$  and  $d_{\min}(C) > 1$ . Assume that there exists a 2-dimensional MDS code  $A \subseteq F^n$ , generated by  $V_2(\alpha)$  for some  $\alpha \in F^n$ with pairwise distinct entries, such that

 $\dim AC = \dim C + 1 \le n - 1.$ 

Then C is generated by  $gV_{\dim C}(\alpha)$  for some  $g \in C$ .

PROOF. Since  $\alpha$  has at most one zero coordinate,  $d_{\min}(C) > 1$  implies that  $\dim \alpha C = \dim C$ . We therefore have

$$\dim AC = \dim(C + \alpha C) = 2\dim C - \dim(C \cap \alpha C),$$

hence

$$\dim(C \cap \alpha C) = \dim C - 1.$$

Moreover,  $C' = C \cap \alpha C$  has support strictly larger than its dimension, otherwise it would have minimum distance 1 and this would imply the existence of a word of weight 1 in C. We prove the lemma by induction on  $k := \dim C$ .

In the case k = 2, pick  $g' \in C \cap \alpha C$ , which exists as  $\dim(C \cap \alpha C) = 1$ , and let  $g \in C$  be such that  $g' = g\alpha$ . Then g and  $g' = g\alpha$  are linearly independent, as  $|\operatorname{supp} g| \ge |\operatorname{supp} g'| \ge 2$ and  $\alpha$  has pairwise distinct entries. It follows that C is generated by g and  $g\alpha$ , i.e. by  $gV_2(\alpha)$ .

Now assume that k > 2. We have

$$k = \dim C' + 1 \le \dim AC' \le \dim \alpha C = k$$

where the right inequality follows from the inclusion  $AC' = C' + \alpha C' \subseteq \alpha C$ , and the left inequality follows from Theorem 2.5 (recall that A is MDS). Strictly speaking, Theorem 2.5 only applies to full-support codes and C' may have a support of cardinality n - 1 if  $\alpha$  has a zero coordinate. But if this happens we may puncture A and C' by deleting this coordinate to obtain full-support codes of the same dimension as A and C and still apply Theorem 2.5.

Since  $C' \subseteq C$  we have  $d_{\min}(C') \geq d_{\min}(C) > 1$ , and we have just shown  $\dim AC' = \dim C' + 1 \leq (n-1) - 1$ , since  $\dim C' = \dim C - 1$ . Therefore the induction hypothesis applies to C', possibly after puncturing one zero coordinate to make C' full support. Hence C' is generated by  $g'V_{k-1}(\alpha)$  for some  $g' \in C'$ . Let  $g \in C$  be such that  $g' = g\alpha$ . The matrix whose rows are the elements of the set  $\{g, g' = g\alpha, \ldots, g'\alpha^{k-2} = g\alpha^{k-1}\} \subseteq C$  is  $gV_k(\alpha)$ , which has rank k as  $|\operatorname{supp} g| \geq |\operatorname{supp} C'| \geq k$ . It follows that this set is linearly independent and  $gV_k(\alpha)$  generates C.

LEMMA 4.5. Let  $C, D \subseteq F^n$  be MDS codes satisfying

$$\dim CD = \dim C + \dim D - 1.$$

Assume that there exists an index set  $I \subseteq \{1, ..., n\}$  with  $|I| \ge \dim CD$  such that the punctured codes  $C_I, D_I \subseteq F^{|I|}$  obtained by projecting C and D on the coordinates indexed by I are Reed-Solomon codes with a common evaluation-point sequence. Then C and D are Reed-Solomon codes with a common evaluation-point sequence.

PROOF. Set  $k := \dim C$ ,  $\ell := \dim D$ . Since  $|I| \ge \dim CD$  we have  $|I| \ge k$  and  $|I| \ge \ell$ and since C and D are MDS we must have  $\dim C_I = \dim C = k$ ,  $\dim D_I = \dim D = \ell$ . Note that we may suppose  $k, \ell \ge 2$ , otherwise there is nothing to prove.

Reformulating the hypothesis, there exist  $g_I, g'_I \in F^{|I|}, \alpha_I \in (F \cup \{\infty\})^{|I|}$ , where  $\alpha_I$  has pairwise-distinct entries, such that  $C_I$  and  $D_I$  are generated by  $g_I V_k(\alpha_I)$  and  $g'_I V_\ell(\alpha_I)$  respectively. In other words there are unique generator matrices  $G_C$  and  $G_D$  of C and D whose I-indexed columns form  $g_I V_k(\alpha_I)$  and  $g'_I V_\ell(\alpha_I)$  respectively. It also follows that  $g_I g'_I V_{k+\ell-1}(\alpha_I)$  generates  $C_I D_I$  (as  $k+\ell-1 \leq |I|$ ), dim  $C_I D_I = k+\ell-1 = \dim CD$  and there is a unique generator matrix  $G_{CD}$  of CD whose I-indexed columns form  $g_I g'_I V_{k+\ell-1}(\alpha_I)$ .

Let  $x_0, \ldots, x_{k-1}$  and  $y_0, \ldots, y_{\ell-1}$  denote the rows of  $G_C$  and  $G_D$  respectively.

The key observation is the following: let u, v, s, t be integers,  $0 \le u, s \le k-1$   $0 \le v, t \le \ell - 1$ , such that

$$u + v = s + t$$
.

Since  $x_u y_v$  and  $x_s y_t$  coincide in the *I*-indexed coordinates, and dim  $C_I D_I = \dim CD$ , the vectors  $x_u y_v$  and  $x_s y_t$  must coincide in every coordinate of  $\{1, \ldots, n\}$ . In other words, if  $\pi = (\pi_0, \pi_1, \ldots, \pi_{k-1})^T$  and  $\rho = (\rho_0, \rho_1, \ldots, \rho_{\ell-1})^T$  are the *j*-th column of  $G_C$  and  $G_D$  respectively, for some  $j \notin I$ , then

$$\pi_u \rho_v = \pi_s \rho_t.$$

We now exploit this property in order to prove the lemma. Pick two columns  $\pi, \rho$  of  $G_C, G_D$  as above.

First assume that  $\pi_0 \neq 0$  and  $\rho_0 \neq 0$ . Without loss of generality we may assume  $\pi_0 = \rho_0 = 1$ . It follows from  $\pi_0\rho_1 = \pi_1\rho_0$  that  $\rho_1 = \pi_1 =: \beta \in F$ . For all  $i \leq k - 1$ , it holds that  $\pi_i = \pi_i\rho_0 = \pi_{i-1}\rho_1$ . Applying this formula recursively we obtain  $\pi_i = \beta^i$  for all  $i \leq k - 1$ , i.e.  $\pi$  corresponds to the evaluation point  $\beta \in F$ . The same argument applies to  $\rho$ , which corresponds to the evaluation point  $\beta \in F$  as well.

Now assume that  $\pi_0 = 0$ . If  $\rho_0 \neq 0$ , then  $\pi_1 \rho_0 = \pi_0 \rho_1 = 0$  implies  $\pi_1 = 0$ . Continuing in this way, we see that if  $\pi_i = 0$ , then  $\pi_{i+1}\rho_0 = \pi_i\rho_1 = 0$  implies  $\pi_{i+1} = 0$  and by induction we obtain  $\pi = 0$  which contradicts the full-support property of the MDS code C. Therefore  $\rho_0 = 0$ . Assume without loss of generality that  $k \leq \ell$ . If  $k = \ell = 2$ , then both  $\pi_1$  and  $\rho_1$ are non zero as C and D have full support, hence the columns  $\pi$  and  $\rho$  correspond to the evaluation point  $\infty$ . If k = 2 and  $\ell \geq 3$  then as  $\rho_i \pi_1 = \rho_{i+1} \pi_0 = 0$  for all  $i < \ell - 1$  and as  $\pi_1 \neq 0$  it follows that  $\rho_i = 0$  for all  $i < \ell - 1$  and again the full-support property of D implies that the column  $\rho$  corresponds to the evaluation point  $\infty$ . If k > 2, then the same procedure that we applied to  $\pi_0, \rho_0$  again yields  $\pi_1 = \rho_1 = 0$ . Iterating in this way, we obtain that both  $\pi$  and  $\rho$  correspond to the evaluation point  $\infty$ .

We have proved that up to multiplication by vectors g, g', the codes C and D have generator matrices of the form  $V_k(\alpha)$  and  $V_\ell(\alpha)$ . Since C and D are MDS, the evaluationsequence  $\alpha$  must have distinct entries and C and D are Reed-Solomon codes with the same evaluation-point sequence.

PROOF OF THEOREM 4.2. Set  $k := \dim C$ ,  $\ell := \dim D$ ,  $k^* := \dim CD = k + \ell - 1$ . Let  $C_0, D_0 \subseteq F^{n_0}$  be the punctured codes obtained by projecting C, D on the first  $n_0 := k^* + 2$  coordinates. As  $C_0, D_0$  and  $C_0D_0$  are MDS, we have dim  $C_0 = \dim C$ , dim  $D_0 = \dim D$ , dim  $C_0D_0 = \dim CD$  and

$$k^* = \dim C_0 D_0 = \dim C_0 + \dim D_0 - 1 = n_0 - 2.$$
(7)

Define the code  $A \subseteq F^{n_0}$  by

$$A := (C_0 D_0)^{\perp}.$$

By Lemma 2.6 the code  $C_0D_0$  is MDS, therefore A is MDS and furthermore has dimension 2 by (7). Now observe that for any  $a \in A$ ,  $x \in C_0$ ,  $y \in D_0$ , orthogonality of A and  $C_0D_0$ translates into

$$(a \,|\, xy) = 0$$

which is equivalent to

$$(ax \mid y) = 0.$$

We have therefore  $(AC_0)^{\perp} \supseteq D_0$ , from which we deduce

$$\dim AC_0 \le n_0 - \dim D_0 = \dim C_0 + 1 \le n_0 - 1$$

whence

$$\dim AC_0 = \dim C_0 + 1 \tag{8}$$

by Theorem 2.5. Similarly we also have

$$\dim AD_0 = \dim D_0 + 1. \tag{9}$$

Now A is an MDS code of dimension 2 and therefore has a generator matrix with at most two zero entries. By puncturing one coordinate if need be, we obtain a generator matrix with at most one zero entry. The two rows of this matrix are clearly of the form  $g, g\alpha$  for some  $g \in F^n$  and  $\alpha \in F^n$  with pairwise distinct coordinates. Finally, consider that dim  $C_0 = n_0 - 1 - \dim D_0 \le n_0 - 3$ , and similarly dim  $D_0 \le n_0 - 3$ . Hence (8) and (9) imply

$$\dim AC_0 \le n_0 - 2,$$
$$\dim AD_0 \le n_0 - 2.$$

Therefore Lemma 4.4 applies to  $A, C_0$  and to  $A, D_0$ , possibly after puncturing one coordinate. From there we obtain that  $C_0$  and  $D_0$  (possibly punctured on a common coordinate) are Reed-Solomon codes with a common evaluation-point sequence, and Lemma 4.5 gives the desired conclusion.

An interesting consequence of Theorem 4.2 is the following characterization of Reed-Solomon codes among MDS codes. Applying Theorem 4.2 in the case C = D yields:

COROLLARY 4.6. Let  $C \subseteq F^n$  be an MDS code, with dim  $C \leq (n-1)/2$ . The code C is Reed-Solomon if and only if

$$\dim C^2 = 2\dim C - 1. \tag{10}$$

REMARK 4.7. If dim  $C \ge (n + 1)/2$ , then C being MDS we must have  $C^2 = F^n$  and the dimension of the square cannot yield any information on the structure of C. However in that case, whether C is Reed-Solomon is betrayed by the dimension of the square of the dual code  $C^{\perp}$ . The remaining case in which Corollary 4.6 does not say anything is the case dim C = n/2. One may wonder whether it still holds that C is Reed-Solomon if and only if dim  $C^2 = 2 \dim C - 1$ , and possibly Theorem 4.2 and Corollary 4.6 have not managed to capture this fact.

The answer to this question is negative, indeed there exist plenty of MDS codes of dimension n/2 satisfying (10) which are not Reed-Solomon. For instance, the codes denoted  $C_{11,8,8}$  and  $C_{13,8,21}$  in [3], of length 8 over the fields with 11 and 13 elements respectively are self-dual, therefore satisfy (10), and can be shown not to be Reed-Solomon.

## 5 Classification of PMDS pairs

We now are finally ready to focus on the paper's central result, namely Theorem 2.12.

First, we show how Randriambololona's Product Singleton Bound can be obtained as a consequence of Theorem 3.3. To be precise we obtain:

THEOREM 5.1. Let  $C_1, \ldots, C_t \subseteq F^n$  be codes. Assume that their product  $C_1 \cdots C_t$  has full support. Then

$$d_{\min}(C_1 \cdots C_t) \le \max\{t - 1, n - (\dim C_1 + \cdots + \dim C_t) + t\}.$$

REMARK 5.2. The full result of [13] is actually stronger than Theorem 5.1, as it ensures that an element of weight at most  $\max\{t-1, n-(\dim C_1 + \cdots + \dim C_t) + t\}$  can be found in the set

$$\{x_1\cdots x_t: x_1\in C_1,\ldots,x_t\in C_t\},\$$

and not only in its span. The support condition given here is also not the same as the apparently weaker hypothesis given in [13], but the two conditions are really interchangeable, as argued in [13, Remark 3(c)].

PROOF OF THEOREM 5.1. For ease of notation, set  $k_i := \dim C_i$  for all  $i = 1, \ldots, t$ ,  $P := C_1 \cdots C_t, k^* := \dim P, d^* := d_{\min}(P)$ . Assume that  $d^* \ge t$ . The classical Singleton Bound, applied to P, says that

$$k^* \le n - d^* + 1. \tag{11}$$

Repeatedly applying Kneser's Theorem 3.3 we obtain

$$k^* \ge k_1 + \dots + k_t - (t-1) \dim \operatorname{St}(P).$$
 (12)

Combining it with (11), we get

$$d^* \le n - (k_1 + \dots + k_t) + 1 + (t - 1) \dim \operatorname{St}(P), \tag{13}$$

which is apparently a weaker statement than Theorem 5.1. To improve it, we "correct" (11) to transform it into an identity, namely we define  $m := n - d^* + 1 - k^*$ . Thus, by definition, P is "*m*-far from being MDS". The combination of this identity with (12) gives an improved version of (13), namely

$$d^* = n - k^* + 1 - m \le n - (k_1 + \dots + k_t) + 1 + (t - 1)\dim \operatorname{St}(P) - m.$$
(14)

In the case of t = 2, the first claim of Lemma 2.11, rewritten as

$$\dim St(P) - (n - d^* + 1 - k^*) \le 1$$

immediately proves the theorem. In the general case, using the second claim of Lemma 2.11 instead we obtain

$$(t-1)\dim \operatorname{St}(P) - m \leq (t-1)\frac{n-k^*}{d^*-1} - m$$
$$= t-1 + (t-1)\frac{m}{d^*-1} - m$$
$$= t-1 - \frac{d^*-t}{d^*-1}m.$$
(15)

 $\triangle$ 

As  $d^* \ge t$  the conclusion follows.

From here on we focus on the case of t = 2. Recall that a pair of codes  $C, D \subseteq F^n$  is defined to be PMDS if

$$2 \le d_{\min}(CD) = n - \dim C - \dim D + 2.$$

Observe that for a PMDS pair (C, D) all inequalities in the proof of Theorem 5.1 are actually identities. From this simple observation we obtain some corollaries which relate the Product Singleton Bound with Kneser's Theorem and with the classical Singleton Bound.

COROLLARY 5.3. Let  $C, D \subseteq F^n$  be codes such that the pair (C, D) is PMDS. Then the following hold.

1. The pair (C, D) attains the bound of Kneser's Theorem, i.e.

$$\dim CD = \dim C + \dim D - \dim \operatorname{St}(CD)$$

2. Either CD is MDS or  $d_{\min}(CD) = 2$ .

PROOF. From the above observation, (12) is an identity if (C, D) is PMDS, hence the first claim is immediately proved. From (14) and (15) we obtain

$$\frac{d_{\min}(CD) - 2}{d_{\min}(CD) - 1} m = 0.$$

where  $m := n - d_{\min}(CD) + 1 - \dim CD$ , hence either m = 0 meaning CD is MDS, or  $d_{\min}(CD) = 2$ .

The two possible cases in our main Theorem 2.12 arise from the two possible situations given by the second claim of the above corollary. We distinguish the case of  $d_{\min}(CD) > 2$ , which implies that CD is MDS, and  $d_{\min}(CD) = 2$ .

PROPOSITION 5.4. Let  $C, D \subseteq F^n$  be codes such that the pair (C, D) is PMDS, and assume  $d_{\min}(CD) > 2$ . Then C, D and CD are MDS. Moreover, if  $\dim C, \dim D \ge 2$  then C, D and CD are Reed-Solomon codes with a common evaluation-point sequence.

PROOF. By the above corollary CD is MDS. Moreover the PMDS property immediately yields  $n > \dim C + \dim D$ . We now proceed to prove that C and D are also MDS through Lemma 2.2.

Set  $k := \dim C, \ell := \dim D$ . Without loss of generality, we can choose a generator matrix  $G_C$  of C that is systematic in the first k positions. Let  $G_D$  be a generator matrix of D. The matrix formed by the last n - k columns of  $G_D$  has full rank, otherwise there is a non-zero vector of D that is zero in the last n - k positions, and taking the product with a row of  $G_C$  we would obtain a vector of CD of weight 1, contradicting that CD is MDS and not the whole space  $F^n$ . So we can now assume that  $G_C$  is systematic in the first k positions and  $G_D$  is systematic in the subsequent  $\ell$  positions.

Now we focus on  $G_C$ . Assume that there is a zero entry in the *j*-th column of  $G_C$  for some  $j > k + \ell$ , say in position (i, j) of  $G_C$ . Then, since the *j*-th column of  $G_D$  is not all-zero (otherwise CD would not be full support and would not be MDS), the product of the *i*-th row of  $G_C$  with some row of  $G_D$  yields non-zero vector of CD of weight at most  $n - k - \ell + 1 = d_{\min}(CD) - 1$ , a contradiction. Therefore, all columns of  $G_C$  indexed by  $j > k + \ell$ , that exist since  $n > k + \ell$ , have no zero entries. For the same reason, this is also true of  $G_D$ , and we obtain that the product of any row of  $G_C$  with any row of  $G_D$  is non-zero.

From this last fact, we get that  $G_C$  cannot have zero entries in the columns indexed by  $\{k+1, \ldots, k+\ell\}$ , or again, by taking a product of a row of  $G_C$  with a row of  $G_D$ , we would have a non-zero vector of CD of weight at most  $n-k-\ell+1$ . Now Lemma 2.2 allows us to conclude that C is MDS. Analogously, one has that D is MDS as well.

The last statement now follows immediately by Theorem 4.2. Note that  $n > \dim C + \dim D$  is equivalent to the hypothesis  $\dim CD \leq n-2$ .

The following lemma will be useful to deal with the second case.

LEMMA 5.5. Let  $C, D \subseteq F^n$  be codes such that CD is MDS and

 $\dim CD = \dim C + \dim D - 1 = n - 1.$ 

Then there exists  $g \in (F^n)^{\times}$  such that  $C = (gD)^{\perp}$ .

PROOF. Let  $g \in F^n$  be a generator of  $(CD)^{\perp}$ , which is invertible as  $(CD)^{\perp}$  is MDS of dimension 1. For any  $x \in C, y \in D$ , we have

$$(x \mid gy) = (xy \mid g) = 0$$

so that  $C \subseteq (gD)^{\perp}$ , and equality follows by comparing dimensions.

 $\triangle$ 

PROPOSITION 5.6. Let  $C, D \subseteq F^n$  be codes such that the pair (C, D) is PMDS. Set  $h := \dim \operatorname{St}(CD)$  and let  $\{\pi_1, \ldots, \pi_h\}$  be an F-basis of  $\operatorname{St}(CD)$  of disjoint projectors with supports  $I_1, \ldots, I_h$ . Then C, D and CD decompose as

$$C = \pi_1 C \oplus \cdots \oplus \pi_h C,$$
  

$$D = \pi_1 D \oplus \cdots \oplus \pi_h D,$$
  

$$CD = \pi_1 C D \oplus \cdots \oplus \pi_h C D$$

and, for all i = 1, ..., h, we have supp  $\pi_i C = \operatorname{supp} \pi_i D = \operatorname{supp} \pi_i C D = I_i$  and

$$\dim \pi_i CD = \dim \pi_i C + \dim \pi_i D - 1. \tag{16}$$

Moreover, if  $d_{\min}(CD) = 2$  then, for all i = 1, ..., h, when  $\pi_i C$  and  $\pi_i D$  are identified with codes of  $F^{|I_i|}$  through the natural projection on their support, then  $\pi_i C = (g_i \pi_i D)^{\perp}$  for some  $g_i \in (F^{|I_i|})^{\times}$ .

**PROOF.** By Kneser's Theorem we have, for all i = 1, ..., h,

$$\dim \pi_i CD \ge \dim \pi_i C + \dim \pi_i D - 1 \tag{17}$$

since  $\pi_i CD$  has trivial stabilizer. Therefore

$$\dim CD = \sum_{i=1}^{h} \dim \pi_i CD \ge \sum_{i=1}^{h} (\dim \pi_i C + \dim \pi_i D - 1).$$
(18)

Observing that

$$C \subseteq \pi_1 C \oplus \dots \oplus \pi_h C, \quad D \subseteq \pi_1 D \oplus \dots \oplus \pi_h D$$
 (19)

we get

$$\sum_{i=1}^{h} (\dim \pi_i C + \dim \pi_i D - 1) \ge \dim C + \dim D - h,$$

but the right hand side of this inequality equals dim CD by the first claim of Corollary 5.3. From (18) we obtain therefore that all inequalities in (17) are equalities, i.e. for all  $i = 1, \ldots, h$ ,

$$\dim \pi_i CD = \dim \pi_i C + \dim \pi_i D - 1,$$

and we obtain also

$$\sum_{i=1}^{n} (\dim \pi_i C + \dim \pi_i D) = \dim C + \dim D,$$

hence both inclusions in (19) are actually identities.

h

Now assume that  $d_{\min}(CD) = 2$ . Observe that  $n = \dim C + \dim D$  by the Product Singleton Bound. From here on all codes are identified with full-support codes through the natural projection on their support. For all  $i = 1, \ldots, h$ , we have  $d_{\min}(\pi_i CD) \ge 2$ , hence  $|I_i| \ge \dim \pi_i CD + 1$  by the classical Singleton Bound applied to  $\pi_i CD$ . Therefore

$$n = \sum_{i=1}^{h} |I_i| \ge \sum_{i=1}^{h} (\dim \pi_i CD + 1)$$
$$= \sum_{i=1}^{h} (\dim \pi_i C + \dim \pi_i D)$$
$$= \dim C + \dim D = n.$$

It follows that

$$\dim \pi_i CD = \dim \pi_i C + \dim \pi_i D - 1 = |I_i| - 1$$

and  $d_{\min}(\pi_i CD) \ge 2 = |I_i| - \dim \pi_i CD + 1$  proves that  $\pi_i CD$  is MDS. Now the conclusion follows by Lemma 5.5.

Propositions 5.4 and 5.6 constitute the proof of Theorem 2.12.

## 6 Concluding comments

As mentioned in Section 4, Theorem 4.2 is arguably a coding-theoretic analogue of Vosper's Addition Theorem. The analogy with its additive counterpart is not as clear-cut however as in the case of Theorem 3.3 and Kneser's Addition Theorem. More precisely, the MDS hypothesis in Theorem 4.2 is not a very natural analogue of the prime order of the ambient group hypothesis in Vosper's original Theorem, and there may possibly be other coding-theoretic analogues to consider.

The natural question raised by Theorem 3.3 and Theorem 4.2 is whether there exists a satisfying characterization of pairs C, D such that CD is indecomposable and of codimension at least 2, and dim  $CD = \dim C + \dim D - 1$ . Beside pairs of Reed-Solomon codes, one now has Reed-Solomon codes with duplicate coordinates. Beside these, other examples turn up. In particular, one may take the amalgamated direct sum [5] of self-dual codes.

If the analogy with additive combinatorics is to be trusted, such a characterization may be tractable – though probably difficult – and would be a coding-theory equivalent of Kemperman's Structure Theorem for small sumsets [9].

Finally, it is natural to wonder whether the characterization of PMDS pairs extends to products of more than two codes. Our techniques (Corollary 5.3 and Proposition 5.4) allow to deal with the analogue of the first case of Theorem 2.12 and to prove the following: if  $(C_1, \ldots, C_t)$  is a *t*-PMDS tuple, i.e. satisfies

$$d_{\min}(C_1 \cdots C_t) = n - (\dim C_1 + \cdots + \dim C_t) + t,$$

if none of the  $C_i$ 's has dimension 1 and

$$d_{\min}(C_1 \cdots C_t) > t,$$

then all  $C_i$ 's are Reed-Solomon codes with a common evaluation-point sequence. On the other hand the arguments in the paper do not seem quite sufficient to deal with the case of

$$d_{\min}(C_1 \cdots C_t) = t$$

corresponding to the second case of our main theorem. We leave the matter open for further study.

While this paper was in review, a version of Kneser's Theorem for a family of algebras was independently posted by Beck and Lecouvey [2], and is a more general version of Theorem 3.3. The proof follows similar arguments. Moreover, [2, Section 4.3] shows how Kneser's original Theorem can be recovered from the new variant. The very same argument, which is based on the embedding of the group G into the complex group algebra  $\mathbb{C}[G]$  and on the isomorphism  $\mathbb{C}[G] \cong \mathbb{C}^{|G|}$ , allows one to recover the original theorem from our variant as well.

### References

 C. Bachoc, O. Serra, and G. Zémor. An analogue of Vosper's Theorem for extension fields. Preprint, 2015: http://arxiv.org/abs/1501.00602

- [2] V. Beck and C. Lecouvey. Additive combinatorics methods in associative algebras. Preprint, 2015: http://arxiv.org/abs/1504.02287
- [3] K. Betsumiya, S. Georgiou, T. Gulliver, M. Harada, and C. Koukouvinos. On self-dual codes over some prime fields. *Discrete Mathematics*, vol. 262(1-3), pp. 37-58, February 2003.
- [4] I. Cascudo, R. Cramer, D. Mirandola, and G. Zémor. Squares of Random Linear Codes. *IEEE Trans. Inform. Theory*, vol. 61(3), pp. 1159-1173, March 2015.
- [5] G. Cohen, I. Honkala, S. Litsyn, and A. Lobstein. Covering Codes. North-Holland, 1997.
- [6] R. Cramer, I. Damgård, and J. B. Nielsen. Secure Multiparty Computation and Secret Sharing. Cambridge University Press, 2015.
- [7] A. Dür. The automorphism groups of Reed-Solomon codes. Journal of Combinatorial Theory, Series A, vol. 44(1), pp. 69-82, January 1987.
- [8] X.-D. Hou, K. H. Leung, and Q. Xiang. A Generalization of an Addition Theorem of Kneser. *Journal of Number Theory*, vol. 97(1), pp. 1-9, November 2002.
- [9] J. H. B. Kemperman. On small sumsets in an abelian group. Acta Mathematica, vol. 103(1-2), pp. 63-88, July 1960.
- [10] M. Kneser. Abschätzung der asymptotischen Dichte von Summenmengen. Mathematische Zeitschrift, vol. 58(1), pp. 459-484, 1953.
- [11] F. J. McWilliams and N. J. A. Sloane. The Theory of Error-Correcting Codes. North-Holland, 1977.
- [12] M. Nathanson. Additive Number Theory: Inverse Problems and the Geometry of Sumsets. Springer, 1996.
- [13] H. Randriambololona. An upper bound of Singleton type for componentwise products of linear codes. *IEEE Trans. Inform. Theory*, vol. 59(12), pp. 7936-7939, December 2013.
- [14] H. Randriambololona. Linear independence of rank 1 matrices and the dimension of \*-products of codes. To appear in *Proc. IEEE Int. Symp. Inf. Theory*, Hong-Kong, June 2015.
- [15] H. Randriambololona. On products and powers of linear codes under componentwise multiplication. *Contemporary Mathematics*, vol. 637, AMS, April 2015.
- [16] T. Tao and V. H. Vu. Additive Combinatorics. Cambridge studies in advanced mathematics 105, Cambridge University Press, 2006.
- [17] A. G. Vosper. The Critical Pairs of Subsets of a Group of Prime Order. Journal of The London Mathematical Society, vol. 31, pp. 200-205, 1956.